Mechanical loading reveals an intrinsic cardiomyocyte stiffness contribution to diastolic dysfunction in murine cardiometabolic disease

Johannes V. Janssens[1,6], Antonia J.A. Raaijmakers[1], Parisa Koutsifeli[3], Kate L. Weeks[1,4], James R. Bell[1,5],

Jennifer E. Van EyK[6], Claire L. Curl[1], Kimberley M. Mellor[2,3,1] and Lea M.D. Delbridge[1]

[1] Department of Anatomy & Physiology, University of Melbourne, Australia

[2] Department of Physiology, University of Auckland, New Zealand

[3] Auckland Bioengineering Institute, University of Auckland, New Zealand

[4] Baker Department of Cardiometabolic Health (Baker), , University of Melbourne, Australia

[5] Department of Microbiology, Anatomy, Physiology & Pharmacology, La Trobe University, Australia

[6] Smidt Heart Institute, Cedars-Sinai Medical Center, Los Angeles, USA.

Running Title: *Cardiomyocyte stiffness in diastolic dysfunction*

Key words: *cardiomyocyte; diastolic dysfunction; cardiometabolic disease; stiffness; relaxation; Ca^{2+} myofilament interaction*

Corresponding author:

Prof Lea MD Delbridge

Cardiac Phenomics Laboratory

Department of Anatomy and Physiology,

University of Melbourne, Australia.

lmd@unimelb.edu.au

ABSTRACT

Increasingly, cardiometabolic syndromes including diabetes and obesity, are associated with occurrence of heart failure with diastolic dysfunction – including heart failure with preserved ejection fraction (HFpEF) and diabetic cardiomyopathy. Diagnosis of heart failure with diastolic dysfunction is independently associated with higher incidence of adverse clinical events. There is currently no specific treatment for diastolic dysfunction and therapies to manage symptoms have limited efficacy. Understanding of the cardiomyocyte origins of diastolic dysfunction is an important priority to identify new therapeutics. The goal of this investigation was to experimentally define *in vitro* stiffness properties (stress/strain) of cardiomyocytes derived from hearts of animals exhibiting diastolic dysfunction *in vivo* in response to dietary induction of cardiometabolic disease.

Mice fed a high-fat-sugar-diet (HFSD vs control) for at least 25 weeks exhibited glucose intolerance, obesity and diastolic dysfunction *in vivo* (echo E/e'). Intact isolated paced cardiomyocytes were each functionally investigated in three states: non-loaded, loaded ('afterload') and stretched ('pre-load') configurations. Mean stiffness of HFSD cardiomyocytes was 70% higher than control. The E/e' doppler ratio for the origin hearts was elevated by 35%. A significant relationship was identified between the *in vitro* stiffness of single cardiomyocytes (Youngs Modulus) and the *in vivo* dysfunction severity (Doppler E/e').

In relation to lengthening/relaxation performance, in the loaded HFSD cardiomyocytes (vs CTRL) the decrement in maximal sarcomere lengthening rate was more accentuated. With stretch, the Ca^{2+} transient decay time course was prolonged. In addition for HFSD the pacing-associated stiffness elevation was further increased and diastolic sarcomere length decreased, although diastolic Ca^{2+} level was reduced. Our findings suggest a previously undescribed alteration in Ca^{2+}- myofilament interaction contributes to stiffness in HFSD cardiomyocytes.

This study provides the first quantitative analysis of intact cardiomyocyte stiffness and performance deficits associated with acquired cardiometabolic disease. The findings demonstrate that a component of the cardiac diastolic dysfunction in cardiometabolic disease derives from intrinsic cardiomyocyte mechanical abnormality. These data offer impetus for further investigation to achieve therapeutic targeting of the specific myofilament structural modifications conferring cardiomyocyte stiffness in cardiometabolic disease.

INTRODUCTION

Increasingly, cardiometabolic syndromes, including diabetes and obesity, are associated with occurrence of heart failure with a diastolic dysfunction patho-phenotype (Sotomi *et al.*, 2021). Diastolic dysfunction is characteristic of heart failure with preserved ejection fraction (HFpEF) which has now overtaken heart failure with reduced ejection fraction (HFrEF) comprising the majority of new heart failure diagnoses (McHugh *et al.*, 2019; Jackson *et al.*, 2022). Diastolic dysfunction is also linked with progression of diabetic cardiomyopathy (Patil *et al.*, 2011; Redfield, 2016). With both HFpEF and diabetic heart disease, early development of diastolic dysfunction can be detected even when other phenotypic features/comorbidities are heterogeneous (ie hypertrophy, dilation, natriuretic factor elevation, hypertension) (Kittleson *et al.*, 2023). Common clinical features of cardiometabolic syndromes, including obesity and insulin resistance, are independent risk factors for diastolic dysfunction as a prelude to heart failure (Rozenbaum *et al.*, 2019; Kamon *et al.*, 2020). Cardiometabolic syndromes can involve the 'silent' presence and progression of diastolic dysfunction for an extended period. Symptoms can be subclinical and/or detectable only in challenge conditions (ie as reduced diastolic reserve in exercise) or when exacerbated by co-morbidity (Leung *et al.*, 2015; Lam *et al.*, 2019). Diagnosis of heart failure with diastolic dysfunction is independently associated with higher incidence of adverse clinical events (Sotomi *et al.*, 2021). Notably, there is currently no specific treatment for diastolic dysfunction and/or HFpEF, and therapies to manage symptoms have limited efficacy (Delbridge *et al.*, 2022).

In diastolic dysfunction, the relaxation and filling phases of the cardiac cycle are impaired and the condition is generally described qualitatively as an increase in left ventricular wall 'stiffness' (or by decrease in compliance, the inverse of stiffness) (Daniels *et al.*, 2022; Janssens *et al.*, 2024). Clinically and experimentally, reduced compliance can be measured invasively as an elevation in left ventricular filling pressure or an increase in the gradient of the diastolic pressure-volume relationship. Non-invasively, on doppler echocardiogram, an increase in the E/e' ratio [of early (E) mitral valve blood flow rate to early mitral annulus tissue motion (e')] is an index used to identify diastolic abnormality (Lo & Thomas, 2010). Myocardial stiffness is considered to reflect myocardial extracellular matrix abnormalities linked with increased tissue fibrosis. While a component of reduced compliance likely reflects fibrotic infiltration, diastolic dysfunction can be identified in settings where there is no evidence of increased interstitial fibrosis burden (Curl *et al.*, 2018). There is growing appreciation that alterations of myocardial compliance in diastolic dysfunction disease states represents underlying intrinsic cardiomyocyte stiffness pathology which contributes to impaired relaxation and filling (Zhou & Pu, 2016; Villalobos Lizardi *et al.*, 2022). Understanding of the cardiomyocyte origins of diastolic dysfunction is an important priority to identify new molecular targets and to support the development of new diagnostics and therapeutics to reduce morbidity and mortality in cardiometabolic disease states.

Cardiomyocyte origins of diastolic dysfunction potentially comprise structural and operational components.

Evidence of molecular changes in sarcomeric and cytoskeletal elements is extensive (Villalobos Lizardi *et al.*,

2022; Loescher *et al.*, 2023). Operationally, the cardiac cycle of contraction and relaxation is driven by the

cyclic flux of cardiomyocyte activator Ca^{2+} to initiate contraction (myocyte sarcomere shortening) and to

allow relaxation (myocyte sarcomere lengthening). Much current knowledge of cardiomyocyte failure

dysfunction derives from myocytes isolated from tissues of hearts exhibiting systolic dysfunction of various

aetiologies, now characterized as HFrEF (van Heerebeek *et al.*, 2006; Curl *et al.*, 2018). The HFrEF isolated

intact cardiomyocyte exhibits reduced shortening and decreased amplitude Ca^{2+} activation transients when

paced *in vitro* to contract isotonically (ie free shortening & non-loaded). In systolic dysfunction, a disease

mechanism of cardiomyocyte activator Ca^{2+} deficit is consistently observed in non-loaded cardiomyocyte

studies and has been a phenotypic finding for cardiomyocytes of a range of animal models. Marked

increases in interstitial and perivascular fibrosis are common features of HFrEF, conferring myocardial ECM

stiffness (Luo & Anderson, 2013). As various cardiac intervention surgeries are common for HFrEF patients,

biopsy tissues have been relatively available and clinical confirmation of cardiomyocyte experimental

findings has been extensive. HFpEF surgical interventions are much less frequent and availability of HFpEF

tissue samples limited.

In diastolic dysfunction, specifically in HFpEF animal models, evidence is accumulating that Ca^{2+} deficit and

impaired isotonic shortening are not phenotypic (Kilfoil *et al.*, 2020). Indeed non-loaded cardiomyocyte

hypercontractility and elevated Ca^{2+} levels have been reported (Curl *et al.*, 2018; Louch, 2020). Thus current

experimental enquiry focusses on cardiomyocyte performance under loaded conditions. While the term

'stiffness' is used as a general descriptor of impaired compliance in a cardiovascular setting, it is more

formally specified quantitatively in physical science disciplines (Mirsky, 1976). Stiffness is defined as the

force per cross-sectional area (stress) developed with application of a normalized stretch or deformation

(strain) to a material. The ratio of stress/strain defines stiffness of the material and is known as 'Young's

Modulus'. In the cardiac setting, a stretch or deformation state is commonly described as 'loading' – for

example 'preload' (diastolic stretch) and 'afterload' (resistance to shortening). Depending on the

experimental design, stiffness behaviour of myocardial tissues, cardiomyocytes and/or sub cellular

cytoskeleton and sarcomere structures all potentially contribute to *in vivo* functional stiffness. Little is

known about how stiffness measured at different structural levels may determine function *in vivo* and how

stiffness elements are quantitatively changed in pathophysiology – in particular in acquired cardiometabolic

disease conditions. Thus, the goal of this investigation was to experimentally characterize *in vitro* stiffness

properties (stress/strain) of cardiomyocytes derived from hearts of animals exhibiting diastolic dysfunction

in vivo in response to dietary induction of cardiometabolic disease.

Mice fed a high-fat-sugar-diet (HFSD vs control) for at least 25 weeks demonstrated glucose intolerance,

obesity and diastolic dysfunction *in vivo* (echo E/e'). Intact isolated paced cardiomyocytes were each

functionally investigated in three states: non-loaded, loaded ('afterload') and stretched ('pre-load')

configurations. Our findings suggest a previously undescribed alteration in Ca^{2+}- myofilament interaction

contributes to stiffness in HFSD cardiomyocytes, potentially by selectively reducing cross-bridge

deactivation rates and increasing the number of cycling cross-bridges during diastole.

METHODS & MATERIALS

Animal model of cardiometabolic disease

As a pre-clinical cardiometabolic disease model, the high-fat/sugar diet (HFSD) fed rodent recapitulates a common and growing human disease phenotype (obesity, insulin resistance, glucose intolerance) and has been established as a diastolic dysfunction model (Heather *et al.*, 2022). Cardiometabolic disease was induced in male C57Bl/6J mice by a high fat/high sugar (HFSD) dietary intervention commencing at 8-9 weeks of age. Animals were randomized to experimental treatment groups and were group housed (> 4 animals per cage) in a temperature-controlled environment 21-23°C with 12 hour light/dark cycles. After a 1 week transitional feeding period (alternating feed periods), animals were fed a HFSD (43% kcal from fat, 200g/kg sucrose, SF04-001, Specialty Feeds) or control reference diet (CTRL: 16% kcal from fat, 100g/kg sucrose, AIN93G, Specialty Feeds) for at least 25 weeks. All animal experiments were approved by the University of Melbourne (School of Biomedical Sciences) Animal Ethics Committee and complied with the guidelines and regulations of the Code of Practice for the Care and Use of Animals for Scientific Purposes (National Health and Medical Research Council Australia).

Glucose tolerance testing

At 23-25 weeks of feeding, glucose tolerance testing was performed in mice following 6 hours fasting as previously described (Chandramouli *et al.*, 2018). Baseline blood glucose levels were measured using an Accu-Chek glucometer with a blood sample obtained from a needle prick to the tail vein. Glucose (1.5 g/kg body weight) was injected i.p. and blood glucose was measured at 5, 15, 30, 60 and 90 minutes after the glucose injection.

Transthoracic echocardiography

At 25-26 weeks of feeding, cardiac structure, systolic and diastolic function were evaluated by transthoracic 2-dimensional B- and M-mode and blood flow and tissue doppler echocardiography (GE Vivid 9; 15mHz i13L linear array transducer) as described previously (Chandramouli *et al.*, 2018; Curl *et al.*, 2018). Briefly, mice were lightly anesthetised with Isoflurane (1.5%; Baxter healthcare, Australia). Parasternal short axis view was used to measure systolic parameters (interventricular septum IVS, left ventricular posterior wall LVPW, left ventricular internal dimension LVID, fractional shortening FS, ejection fraction EF, heart rate HR). Pulse wave Doppler and tissue Doppler scans were acquired from the apical 4 chamber view to assess LV diastolic function parameters: velocity of early (E) and late (A) mitral inflow and early (e') and late (a') diastolic velocity of mitral annulus. From these compound indices of diastolic function E/e', E/A and e'/a' were calculated. In addition, the slope of the E wave was used to calculate mitral valve deceleration time. Three consecutive cardiac cycles were sampled for each measurement.

Cardiomyocyte preparation, structural and functional assessment

A series of protocols were implemented to phenotype cardiomyocyte and sarcomere morphology, and

assess functional status in different contraction configurations and challenge settings.

Cardiomyocyte isolation, morphology & microfluorimetry. HFSD and CTRL left ventricular cardiomyocytes

were enzymatically isolated using methods previously described (Howlett *et al.*, 2006; Mellor *et al.*, 2011).

Briefly, the heart was excised, cannulated via the aorta, and subjected to retrograde perfusion on a

Langendorff apparatus with an oxygenated Ca^{2+} free buffer at 37°C for 10 minutes. The buffer was

composed of (in mM): 130 NaCl, 5 KCl, 0.33 NaH_2PO_4; 1 $MgCl_2$, 25 HEPES; 20 Glucose; 3 Na Pyruvate; 1 Na

Lactate. The Ca^{2+} free buffer was then supplemented with 0.66mg/mL collagenase (Worthington; Type II),

0.033mg/mL trypsin and 0.05mM $CaCl_2$ and perfusion continued for a further 10 minutes. The heart was

removed from the Langendorff apparatus, and the left ventricle was segmented into small pieces in a high

potassium solution composed of (in mM): 30 KCl, 90 KOH, 30 KH_2PO_4, 3 $MgSO_4$, 50 glutamic acid, 20

taurine, 0.5 EGTA, 10 glucose, and 10 HEPES. Single left ventricular cardiomyocytes were isolated via gentle

trituration. Viable cardiomyocytes were resuspended in HEPES-Krebs buffer (in mM); 146.2 NaCl, 4.69 KCl,

0.35 $NaH_2PO_4H_2O$, 1.05 $MgSO_47H_2O$, 10 HEPES, 11 glucose containing 1.8mM Ca^{2+} and were stored on a

rocking platform at room temperature. From each heart cell dimensions were collected from at least 50 rod

shaped cardiomyocytes using an inverted light microscope (Nikon) and calibrated eye piece as previously

described (Curl *et al.*, 2018). Cardiomyocyte volume was calculated by inserting experimentally determined

myocyte length and width values into the formula: volume =length x width x depth x multiplier 0.54

(elliptical area of the rectangle defined by width & length, (Satoh *et al.*, 1996)). Cardiomyocyte width: depth

ratio was assumed to be 1.44 (Natali *et al.*, 2001). Cardiomyocyte cross-sectional area (CSA) could be

calculated from measured width scaled for 2-dimensional geometry. (CSA = W [W/1.44] x 0.54 μm^2). An

aliquot of isolated cardiomyocyte suspension was pipetted into a cell chamber mounted on the stage of an

inverted fluorescence microscope (Motic AE31) and superfused with HEPES-Krebs solution with 2mM Ca^{2+},

37°C at a flow rate of 2ml/min. Cardiomyocyte sarcomere length and Ca^{2+} transients were simultaneously

recorded during isotonic contractions. Myocyte force development was evaluated in the same cells with

transducer attachment, using four protocols described below.

Characterizing cardiomyocyte diastolic function: non-loaded shortening measurements (isotonic), loaded

force measurements (auxotonic), stretch responses and pacing challenge. Cardiomyocyte performance

was evaluated in several paced conditions: designated non-loaded, loaded and stretched. The 'non-loaded'

state refers to the absence of externally applied load (noting that sarcomeric and cytoskeletal structures do

constitute an underlying internal load).

Firstly, cardiomyocyte non-loaded (isotonic) sarcomere contractions (IonOptix SarcLen, Fast Fourier

Transform, FFT) and Ca^{2+} transients (Fura2-AM, 5μM microfluorimetry F340:380nm ratio) were

simultaneously measured at 2Hz, using a microscope diaphragm to define a rectangular intra-myocyte

region of interest as depicted **Fig. S1**. During contraction cycles, cardiomyocyte sarcomere length was

monitored within an intracellular region of interest defined by an adjustable diaphragm mounted in the

microscope optical tube (**Fig S1A**)

Secondly, to obtain loaded (auxotonic) contraction data (ie 'afterload'), each cardiomyocyte was attached

to a pair of glass fibres coated with a bio-adhesive (MyoTak, IonOptix, USA). Attachment sites were

positioned to be outside the sarcomere tracking region, closer to myocyte 'ends'. One glass fibre was

connected to a force transducer (fibre stiffness 20N/m, IonOptix, USA), and the other to a piezo motor

(Nano-drive, Mad City Labs Inc. WI, USA) to adjust cardiomyocyte length. With successful attachments,

cardiomyocytes were lifted from the coverslip using motorised micromanipulators (MC1000E, Siskiyou

Corporation, USA) attached to both the force transducer and the piezo-motor. Baseline auxotonic

contractions at 2Hz were recorded. Sarcomere length changes were measured using FFT as described

above. In addition, with glass fibre attachment, tracking of mean sarcomere shortening within a myocyte

segment delineated by the glass fibre edge positions could be performed in parallel.

Thirdly, auxotonic responses during a standardized stretch protocol ('pre-load') were recorded to measure

the effect of increasing diastolic cardiomyocyte/sarcomere basal length during pacing. A quasi 'stress-

length' relation was constructed for each myocyte (where stress = force/cross-sectional area). The 'stress-

length' relation represents a cellular equivalent to the conventional muscle 'force-length' relationship. Prior

to the commencement of the myocyte stretch measurements a protocol to calibrate the stretch step was

implemented. Four consecutive 0.5V input signals were delivered to the piezo-motor to determine a

calibration constant to define the percentage cell stretch for a given voltage. For each myocyte a

Calibration Constant was calculated as: *myocyte segment length change per peizomotor mV / sarcomere*

count in segment [where *myocyte sarcomere count = myocyte segment length (μm) / average diastolic*

sarcomere length (μm)]. To track the glass fibre position a real-time contrast detection algorithm (IonOptix -

EdgeLength) was focused on the internal edge of each of the glass fibres (**Fig. S1**) (Peyronnet *et al.*, 2017).

This enabled reproducible and comparable stretch steps to be applied to different cardiomyocyte

preparations (**Fig. S1**). Prior to the 'stress-length' protocol, a cell-specific calibration constant was applied to

yield a standardised stretch for each cardiomyocyte evaluated. The stress-length protocol involved

progressive stretch steps totaling an approximate 30% cell stretch (**Fig. S1**).

Lastly, the pacing was increased to 4Hz, a rate adaptation period allowed (2mins) and the stress-length

protocol was repeated. Sarcomere length, force development and intracellular Ca^{2+} transients were

simultaneously measured. In summary, the following parameters were measured: diastolic stress

(mN/mm^2), diastolic stress-length relation [(mN/mm^2) / (%$^-$ length change), diastolic stress-sarcomere

length relations (mN/mm^2 . µm^{-1}), sarcomere shortening (% diastolic SL), maximum sarcomere shortening

and lengthening velocity (µm/s). In addition, Ca^{2+} parameters were measured: diastolic Ca^{2+}, systolic Ca^{2+},

amplitude of the Ca^{2+} transient and time constant of decay of the Ca^{2+} transient (tau).

Statistical analyses

Data are presented as mean ± SEM and statistical analysis was performed using GraphPad Prism V9.5.1.

Cardiomyocyte stress-length and stress-SL relations were modelled with simple linear regression. For

comparison between two groups an unpaired student t-test was used. For assessment between two

groups at multiple points a one-way ANOVA with Bonferroni multiple comparisons post-hoc test was used.

Two-way ANOVA with Šídák's multiple comparisons post-hoc test was used for the evaluation of data

groups with two independent variables. Pearson Correlation analyses were used to assess statistical

significance of fitted linear regression lines. A p-value of <0.05 was considered statistically significant.

Analyses were performed using a blinded protocol.

RESULTS

Systemic indicators of cardiometabolic disease and *in vivo* diastolic function in HFSD mice

High fat /sugar-fed mice exhibited obesity and glucose intolerance. Relative to CTRL, HFSD mice body

weight was significantly elevated within 5 weeks of dietary intervention increasing over the study duration

by 50% relative to CTRL (**Fig. 1A**, CTRL: 29.5±0.5 vs. HFSD: 45.7±1.2g). After 23-24 weeks diet treatment,

glucose tolerance testing of a subset of the HFSD cohort was performed. Glucose intolerance in response to

intraperitoneal glucose administration was evidenced by increased area under the curve, an indicator of

systemic insulin resistance (**Fig. 1B**).

After confirming these key cardiometabolic disease characteristics in the HFSD mice, echocardiographic

functional evaluation was undertaken. Focusing on diastolic function evaluation (**Fig. 1C-G**), there was

reduced velocity of the mitral tissue annulus displacement (CTRL: 28.5±1.3, HFSD: 25.2±0.9 mm/s) and

increased early mitral blood inflow (CTRL: 476.8±19.9, HFSD: 554.6±16.8 mm/s) in HFSD myocardium.

Deceleration time of mitral inflow was shortened (CTRL: 31.5±2.2, HFSD: 19.7±1.5 ms) and E/e' ratio

increased, all observations consistent with reduced tissue compliance and diastolic dysfunction (CTRL:

16.7±1.0, HFSD: 22.3±1.5). No differences in systolic performance parameters were observed (**Table S1**),

confirming the primary *in vivo* phenotype of diastolic dysfunction. The next step was to evaluate the

cardiomyocyte involvement in dysfunction.

Non-loaded cardiomyocyte shortening and lengthening in CTRL and HFSD treatments

Single isolated cardiomyocytes were prepared from hearts of a subset of mice which underwent

echocardiographic analyses. For intact, non-loaded cardiomyocytes sarcomere shortening and lengthening

cycles were similar for CTRL and HFSD groups (**Fig. 2A**). Diastolic sarcomere length (SL) (**Fig. 2B**, CTRL:

1.71±0.01, HFSD: 1.72±0.01 μm) and peak sarcomere shortening (**Fig. 2C**, CTRL: 3.79±0.33, HFSD: 3.94±0.40

% diastolic SL) were not different between groups. Examination of sarcomere contraction kinetics (**Fig. 2D**

exemplar trace) showed that the maximum rate of lengthening (**Fig. 2E**, CTRL: 1.08±0.14, HFSD: 1.29±0.18

μm/s) and shortening (**Fig. 2F**, CTRL: -2.32±0.21, HFSD: -2.57±0.28 μm/s) did not differ between CTRL and

HFSD groups. Collectively these findings indicate that in the non-loaded setting no differences in CTRL and

HFSD cardiomyocyte systolic and diastolic isotonic performance were detected, a finding consistent with

our previous studies in fructose-fed mice (Mellor *et al.*, 2012).

Non-loaded cardiomyocyte Ca^{2+} transient decay and sarcomere lengthening in CTRL and HFSD treatments

Next, Ca^{2+} parameters were compared with cardiomyocyte sarcomere lengthening performance during

relaxation and diastole. No significant difference was evident between dietary treatment groups in the

diastolic Ca^{2+} (**Fig. 3A**, CTRL: 0.749±0.016, HFSD: 0.761±0.017 $F_{340:380}$) or time constant (tau) of Ca^{2+} transient

decay (**Fig.3B**, CTRL: 0.160±0.011, HFSD: 0.137±0.009 sec). Thus, the significant diastolic dysfunction

measured *in vivo* for the HFSD group was not observed in an isotonic non-loaded *in vitro* setting. Treatment

differences in Ca^{2+} transient systolic parameters were also undetectable (**Fig. 3C-E**). To reconcile the

impaired echocardiographic heart performance with absence of dysfunction in isotonically contracting

single cardiomyocytes, the next strategy was to assess single cardiomyocyte response to loading. A

common feature of cardiometabolic disease is elevation in afterload – ie elevated systemic resistance

potentially impeding myocardial pump function. To simulate this *in vitro*, mechanically loaded

cardiomyocyte performance was examined.

Loaded cardiomyocyte sarcomere shortening and lengthening

Immediately after non-loaded (isotonic) cardiomyocyte recordings, glass fibres were attached to

cardiomyocytes at longitudinal positions to permit auxotonic, loaded contraction measurements. This

manoeuvre was successful for about 20% of cardiomyocytes, due to delicacy involved in manual glass fibre-

myocyte manipulations. For cells successfully attached, the mean difference in sarcomere length measured

pre- and post-conversion to auxotonic mode was <0.02% diastolic length, indicating isotonic and auxotonic

sarcomere initial lengths were well matched. For attached cardiomyocytes, a comparison of sarcomere

shortening behaviour in loaded state could be made (using FFT) to directly match with data obtained non-

loaded state.

With loading, for all myocytes (both treatment groups), the sarcomere peak shortening was reduced and

shortening and lengthening kinetics slowed when compared with non-loaded performance (**Fig. 4A**). Mean

data showed that in response to loading, sarcomere shortening was reduced by ~34% (**Fig. 4B**, non-loaded:

0.075±0.008, loaded: 0.050±0.006 μm). Maximum rate of shortening was decreased by 58% (non-loaded: -

2.85±0.28, loaded: -1.20±0.18 μm; not shown), and maximum rate of lengthening by 66% (**Fig. 4C**, non-

loaded: 1.46±0.21, loaded: 0.52±0.10 μm/s). The reductions in extent of shortening and maximum rate of

lengthening were positively correlated (**Fig. 4D**) suggesting the possibility of common underlying

mechanisms. The more marked reductions in shortening and lengthening rates (almost double reduction in

peak shortening) indicates that these kinetics reflect the influence of factors beyond those impacting

shortening alone.

To assess this, the maximum rates of lengthening and shortening were normalised by peak shortening

amplitude to derive measures of rate parameters independent of shortening extent and thus to yield a

measure of 'deformation rate' - ie rate of length change, relative to length change (units μm x sec-1)/μm =

1/sec). The effect of loading on deformation rate was accentuated in HFSD group (significant ANOVA diet x

load interaction effect, **Fig. 4E**). A selective HFSD effect was determined as the differential between

matched 'loaded' and 'non-loaded' lengthening cardiomyocytes. The decrement in maximum rate of

lengthening for CTRL and HFSD is shown in **Fig. 4F**, highlighting the more pronounced slowing of maximal

lengthening rate in HFSD (CTRL: 7.38±1.15, HFSD: 12.70±2.23 1/s) in response to loading. Treatment

differences in deformation were not evident during the shortening phase (data not shown)**.** These findings

indicated an accentuated influence of load, specific to lengthening in HFSD cardiomyocytes, and could

reflect impaired subcellular relaxation processes specific to the HFSD myocardium. Based on these data

identifying different CTRL and HFSD origin cardiomyocyte responses in loaded state, we next investigated

the effect of stretch applied during loaded myocyte pacing, as a single myocyte paradigm simulating

ventricular filling resistance ie 'preload'.

Cardiomyocyte force development: measuring strain & stress during progressive cardiomyocyte stretch

A hallmark of diastolic dysfunction is an elevated diastolic pressure-volume relationship which describes

the increased resistance of the myocardial wall to chamber filling (Aurigemma *et al.*, 2006). We sought to

design a single cardiomyocyte version of this relationship (Fowler *et al.*, 2015), by applying standardized

stretch steps to paced, loaded cardiomyocytes, defining the steps as a % of non-stretched state (ie 'strain').

Diastolic force was measured simultaneously during length stretch steps. The force measured at every

strain increment was normalized by cross sectional area (CSA) to produce a measure of 'stress' (ie

force/CSA, mN/mm^2). Thus a sarcomere 'stress/strain' relation (ie force-length) was constructed for each

myocyte.

The protocols employed to generate stress-strain are depicted in **Fig. S1** and described in detail in the

Methods. **Fig. S1** show the stretch standardisation approach used to enable reproducible application of

strain (converting voltage signals to linear displacement) as a % length change (via individual myocyte

calibration constant calculation) to allow comparisons between different cardiomyocytes. The size of each

step change in length on sarcomere length was measured (FFT) and confirmed by tracking the intra-

cardiomyocyte segment between the glass fibres (**Fig. S1**). In some cases a direct sarcomere FFT measure

was not microscopically resolvable due to myocyte contraction responses which partially positioned some

area of the myocyte outside the optimal focal plane. In these situations it was always possible to calculate

segment length change. For the example shown, stretch steps were determined to be 2.75µm, and

converted to % sarcomere length (ie strain) using sarcomere count within intracellular region of interest

area **(Fig. S1C)**. A comparison of the variability of cell stretches performed using a uniform (pre-set) stretch

constant and the individually derived stretch calibration constant demonstrates that reproducibility is

optimized with usage of the individual stretch calibration constant data (**Fig. S1D**). Force, sarcomere length

and % stretch were measured simultaneously **(Fig. S1)**.

Quantifying diastolic stiffness in CTRL vs HFSD cardiomyocytes using strain normalization methods

Superimposed exemplar stress recordings are shown in **Fig. 5A** for CTRL and HFSD cardiomyocytes,

captured during a series of progressive stretches during a 100 sec period. The broken lines are aligned with

the diastolic stress step increments for each of the 2 myocytes (using the data points collected prior to the

transition to a new step) throughout the series of progressive stretches. A robust increase in systolic force

with stretch is also evident, indicative of length-dependent activation, the cardiomyocyte basis for the

Frank-Starling mechanism.

The formal definition of stiffness is the ratio of force per cross-sectional area (stress) normalized to the

extent of stretch deformation (ie strain) (Mirsky, 1976). Thus, in this experimental setting stiffness could be

visualized as the gradient of the linear plot derived from the cardiomyocyte diastolic stress-length relation.

Our stretch protocol methodology provided two approaches for stiffness calculation (as per Methods).

A first approach used FFT measurement of sarcomere length to track cardiomyocyte stretch and was

plotted according to the standardized stretch steps (in units µm). An alternative approach used percent

length change calculated as the length of the cardiomyocyte segment between the glass attachment fibres

divided by the initial distance between the fibres prior to stretch (in units % initial length).

Applying the first approach, a linear regression was performed to fit the data from which a slope value

could be derived. **Fig. 5B** depicts the average diastolic and systolic stress-sarcomere length relations for

CTRL cardiomyocytes only. To determine cardiomyocyte diastolic stiffness using FFT-computed sarcomere

length values, the gradient of the mean slope of the diastolic stress sarcomere-length relation for each

treatment group was calculated, using mean sarcomere length change (µm) as the denominator. This

gradient was significantly higher in HFSD myocytes compared to CTRL (**Fig. 6A**). The individual slope

gradients were used to calculate the mean diastolic stress / strain gradient (**Fig. 6B,** CTRL: 2.40$\pm$0.72, HFSD:

5.50$\pm$1.51 (mN/mm^2) / µm), p = 0.07) where there was a trend for a 2.3 fold increase in stiffness of HFSD

cardiomyocytes. The statistical power of this analysis was limited due to the difficulty in achieving

resolvable diastolic sarcomere lengths by FFT for all myocytes.

Applying the second approach, it was possible to construct stress – segment length (%) relations for all

myocytes from which stress data could be obtained. In **Fig. 6C**, the stress – segment length data for HFSD

and CTRL are shown, with a robustly significant difference detected in mean gradient values. In **Fig. 6D** the

diastolic stress/strain values calculated for the myocytes from the two treatment groups using this segment

approach are directly compared. The HFSD cardiomyocyte stiffness index derived using this approach was

increased by 70% compared with CTRL (**Fig. 6D**, CTRL: 0.0075$\pm$0.0016, HFSD: 0.0128$\pm$0.0016 ((mN/mm^2)/%

length).

To examine the relationship between intrinsic *in vitro* cardiomyocyte stiffness and *in vivo* diastolic

dysfunction, diastolic stiffness (slope of the diastolic stress-length relation) averaged per animal was

plotted against the E/e' index of diastolic dysfunction. There was a significant correlation between

cardiomyocyte stiffness and E/e' in HFSD and no evidence of a relationship for the CTRL group (**Fig.6E** CTRL:

p=ns, HFSD: p=0.0096, r^2=0.9213). These data support the interpretation that intrinsic cardiomyocyte

stiffness represents a definable and inherent component of diastolic dysfunction occurring *in vivo* in

cardiometabolic disease.

Load dependence of Ca^{2+} transient decay in CTRL and HFSD cardiomyocytes

To further understand the cardiomyocyte functional elements comprising increased intrinsic stiffness in

cardiopathology, the relationship between Ca^{2+} transient features and myofilament deactivation in

response to progressive stretch was evaluated. In the intact cardiomyocyte model these electro-mechanical

coupling processes are functional for interrogation (in contrast to the disrupted cytosolic environment

which characterizes permeabilized myocyte preparations). The Ca^{2+} transient decay features were

compared for isotonic and auxotonic basal states and also in response to progressive stretch.

In CTRL cardiomyocytes there was no difference in the time constant of transient decay (tau, 1/e) between

loaded and stretched state (20% increase in length, **Fig. 7C.** loaded 1.12±0.14; stretched 1.09±0.10 fold

change vs non-loaded reference 1.00±0.12). In HFSD cardiomyocytes the time constant of Ca^{2+} transient

decay was increased in stretched cardiomyocytes relative to non-loaded (**Fig. 7C**. non-loaded, 1.00±0.12;

loaded: 1.22±0.15; stretched 1.42±0.19 fold change). A trend for an interaction effect (p=0.0549) between

diet and load was observed, suggesting potential for a differential Ca^{2+} transient response to load between

CTRL and HFSD cardiomyocytes. The effects can be visualised in **Fig. 7A-B** where the mono-exponential

decay of exemplar Ca^{2+} transients is depicted. The shallower decay of the Ca^{2+} transient in the stretched

state (**Fig 7B**, HFSD) was apparent whereas the loaded and non-loaded decay traces were virtually

overlayed for CTRL cardiomyocytes (**Fig. 7B**). These data suggest that stretch may enhance a mechanism by

which the myofilaments bind Ca^{2+} for a prolonged duration during relaxation thus reducing Ca^{2+} availability

for extrusion from the cytosol.

Pacing-induced $[Ca^{2+}]_i$ increase to examine potential for a cross-bridge derived stiffness mechanism

A further mechanistic study to probe the potential relationship between Ca^{2+} and stiffness involved

perturbation of cytosolic Ca^{2+} cycling time. Elevating stimulation frequency is a recognized physiologic

strategy to increase diastolic Ca^{2+} levels and to apply increased cardiomyocyte workload. In response to

frequency elevation (2Hz to 4Hz) cardiomyocyte diastolic Ca^{2+} levels increased in both CTRL and HFSD (**Fig.**

8A & B) groups consistent with previous studies (Varian & Janssen, 2007; Eisner *et al.*, 2020). In the HFSD

cardiomyocytes this effect was markedly diminished and about half the mean magnitude of the Ca^{2+} shift

observed for CTRL myocytes (**Fig. 8C**. CTRL 0.111±0.017; HFSD 0.054±0.017, $F_{340:380}$, 2-way ANOVA

significant interaction effect). Frequency elevation was associated with a reduction in diastolic sarcomere

length in both CTRL (3%) and HFSD (5%) loaded cardiomyocytes (**Fig. 8D**. CTRL 2Hz: 1.69±0.01, CTRL

4Hz:1.64±0.020, HFSD 2Hz: 1.69±0.014, HFSD 4Hz: 1.61±0.023 μm).The combination of normal diastolic

sarcomere length reduction, concomitant with blunted diastolic Ca^{2+} increase in HFSD cardiomyocytes, may

indicate elevated diastolic responsiveness to Ca^{2+} in the HFSD cardiomyocyte.

While frequency elevation did not change the stress/strain relation in CTRL cardiomyocytes, a significant

increase in the stress/strain gradient was observed in HFSD cardiomyocytes (**Fig. 8E&F**. CTRL 2Hz:

0.006±0.002, CTRL 4Hz:0.007±0.002; HFSD 2Hz: 0.012±0.002, HFSD 4Hz: 0.014±0.002 (mN/mm^2) / %

length). This may indicate a larger population of cross-bridges contributing to diastolic stiffness at

heightened workloads in HFSD cardiomyocytes although other potential non-cross-bridge sources cannot

be excluded.

DISCUSSION

Here we provide the first quantitative analysis of intact cardiomyocyte stiffness and performance deficits associated with acquired cardiometabolic disease. We demonstrate that a major component of the cardiac diastolic dysfunction in cardiometabolic disease derives from intrinsic cardiomyocyte mechanical abnormality.

The concept of myocardial stiffness is a frequently invoked qualitative descriptor of the pathophysiologic phenotype characterizing diastolic dysfunction. Clinically and experimentally the echocardiography parameter E/e' is used as an index of myocardial stiffness in disease diagnosis. The parameter E/e' is a compound measure of cellular and extracellular matrix stiffness contributors. Despite consistent *in vivo* findings, an understanding of the cardiomyocyte-specific contribution to tissue stiffness has been difficult to identify. Using a series of intact single cell recording manoeuvres, we have established that the stiffness characteristics of individual isolated cardiomyocytes from normal and dysfunctional hearts can be defined, normalized and compared. Further, a significant relationship was identified between the mean stiffness of single cardiomyocytes and the extent of the origin heart diastolic dysfunction (ie E/e' vs Young's Modulus).

To recapitulate briefly, cardiometabolic disease induced by dietary intervention in mice was characterized by marked body weight elevation (50%, 'obesity') and glucose intolerance. *In vivo* echocardiography diagnosed significant diastolic dysfunction with a 35% elevation of the E/e' in HFSD animals (**Figs.1 & 2**) in absence of systolic functional deficit (**Table. S1**). In paced non-loaded conditions (freely shortening 'isotonic' state), the sarcomere contraction cycles & Ca^{2+} transients (fura-2) were similar in CTRL and HFSD cardiomyocytes (**Figs. 2 & 3**). When the same cardiomyocytes were loaded, the peak shortening magnitude and maximal lengthening rates (**Fig. 4A-C**) were reduced for both diet conditions. In HFSD cardiomyocytes the reduction in lengthening rate (but not shortening) was accentuated (**Fig 4E & F**), indicating a selective negative impact of loading on relaxation of the HFSD-derived cardiomyocytes. In HFSD cardiomyocytes there was a larger increase in diastolic force per cross-sectional area (stress: mN/um^2) in response to standardized step increases in length (strain: % initial length[um]) (**Fig. 5A-B & 6A-C**). From these data a 'stress/strain' parameter was determined, defining diastolic stiffness for each myocyte ie 'Youngs Modulus' ($[mN/\mu m^2]/[\%$ initial length$[\mu m])$. Mean stiffness was 70% higher in cardiomyocytes of HFSD animals (**Fig. 6D**). Thus the increase in cardiomyocyte stiffness index was approximately double the elevation of the *in vivo* E/e' dysfunction index. Using matched *in vivo* and *in vitro* data, a significant correlation was found between diastolic dysfunction (E/e') and mean cardiomyocyte stiffness value for HFSD-diet animals, while there was no relationship between these parameters in the myocytes of CTRL-diet fed group (**Fig. 6E**). To evaluate possible influence of mechanical state on the Ca^{2+} transient, the mean transient decline parameters (tau) from non-loaded, loaded and stretched cardiomyocytes were compared. In HFSD cardiomyocytes the time constant of Ca^{2+} transient decay was increased in stretched cardiomyocytes

491 relative to non-loaded, a feature not observed in CTRL cardiomyocytes (**Fig. 7**). Finally, in response to an *in*

492 *vitro* pacing HFSD cardiomyocytes exhibited accentuated diastolic stiffness even with a smaller increase in

493 diastolic Ca^{2+} level (**Fig. 8**). Collectively, these findings may reveal a previously undescribed alteration in

494 Ca^{2+}-myofilament interactions indicative of a cross-bridge component to cardiomyocyte stiffness in

495 cardiometabolic disease.

Dietary model of acquired metabolic syndrome and cardiac diastolic disease

High-fat/sugar diet feeding (ie 'western diet') has previously been shown experimentally to induce diastolic

dysfunction in rodents with prolongation of isovolumic relaxation time (Pulinilkunnil *et al.*, 2014; Dia *et al.*,

2020; Sowers *et al.*, 2020; Wingard *et al.*, 2021, 2021). This dietary regime employed in this study has been

well characterized as a model of diastolic dysfunction, with gradual development of metabolic disturbance

(stable rate of weight gain and mild hyperglycemia), with maintained systolic function (ejection fraction and

fractional shortening). As has been reported previously, over the time span of the dietary intervention

differential fibrosis is not observed (Daniels *et al.*, 2022; Wells *et al.*, 2023; data not shown).The data

presented here are consistent with clinical observations linking cardiometabolic disease and diastolic

dysfunction.

The finding of increased stiffness here is consistent with some aspects of prior experiments involving

animal feeding of obesogenic diets, using other rodent preparation types– including permeabilized

cardiomyocytes, tissue strips/papillary muscles, and non-paced cardiomyocytes (Leopoldo *et al.*, 2010;

Gonçalves *et al.*, 2016; Sowers *et al.*, 2020). Interestingly stiffness in those studies was attributed to titin or

extracellular matrix properties. Our protocol with intact single cardiomyocytes allows a mechanistic

extension of these previous findings (see later discussion).

Loading the intact cardiomyocyte - functional deficits revealed in HFSD

Isotonically contracting cardiomyocytes of CTRL and HFSD mice exhibited identical shortening-lengthening

cycles, but a distinctly different decrement in lengthening rate when exposed to load (ie auxotonic with

progressive resistive 'afterload' increase). Our findings show unequivocally that cardiomyocyte mechanical

dysfunction cannot be detected by analysis of isotonic shortening. With loading, the peak shortening

magnitude and the maximal lengthening rates were reduced for both CTRL and HFSD. In HFSD the

lengthening rate reduction was accentuated even when shortening extent was accounted for. These

findings indicated specific influence of loading on lengthening in HFSD cardiomyocytes and could reflect

impaired subcellular relaxation processes specific to the HFSD myocardium.

Others have demonstrated a physiological effect of loading on shortening in cardiac cells/tissues in

different species (using slow pacing and sub physiologic temperature conditions), although the pronounced

effect on relaxation is not consistently reported (Lab *et al.*, 1984; White *et al.*, 1995; Shimkunas *et al.*,

2021). This is consistent with *in vivo* human clinical observations where there is a strong inverse association

between late-systolic loading and left ventricular wall displacement velocity during early diastole (Borlaug

et al., 2007). Importantly, increased dependence of relaxation on afterload is reported to be an *in vivo*

feature of heart failure (Eichhorn *et al.*, 1992; Gillebert *et al.*, 2000). Regarding the Ca^{2+} transient, in the

current study a subtle prolongation effect is observed in both CTRL and HFSD cardiomyocytes when loaded

(but not stretched) which is consistent with recently reported findings in rat LV trabeculae (Dowrick *et al.*,

2022). The variability in the sarcomere shortening recorded for both non-loaded and loaded

cardiomyocytes is notable (**Fig.4**) but consistent with other studies assessing relaxation rates in rodent

cardiomyocytes (Delbridge & Roos, 1997; Clark & Campbell, 2019).

Characterizing cardiomyocyte stretch & pacing challenge responses

In order to quantify intact single cardiomyocyte stiffness our approach was to apply a calibrated stretch

protocol, based on pre-measurement of a calibration constant for each myocyte, tracking changes in force

generation and cytosolic Ca^{2+} responses for each stretch step. This calibrated step-by-step approach

enabled the stress and strain data to be gathered from actively paced cardiomyocytes over extended

periods (ie 80-100sec) and computed to generate a definitive, comparative stiffness parameter for each

cardiomyocyte.

In the intact myocyte model, the internal cytosolic regulatory milieu is preserved and the

cytoskeletal/sarcomeric/organelle structural framework is intact. Thus contributions to diastolic 'stiffness'

reflect components of the structural passive resistance to deformation and the basal crossbridge derived

activity. Recent studies of permeabilized cardiac muscle strip bundles have explored the separate

contributions of mictrotubule, titin, actin-associated, sarcolemmal and desmin-linked components to

passive force development (Loescher *et al.*, 2023). Prior studies of 'skinned' or quiescent myocardial

preparations have also identified sources of passive stiffness, with titin a major stiffness component in

cardiometabolic disease (Leopoldo *et al.*, 2010; Hamdani *et al.*, 2013; Gonçalves *et al.*, 2016; Hopf *et al.*,

2018; Sowers *et al.*, 2020; Abdellatif *et al.*, 2021).

Our findings indicate involvement of a cytosolic Ca^{2+} component of the stretch response. Ca^{2+} fluorescence

analyses from these stretch protocols showed novel stretch-dependent prolongation of Ca^{2+} transient

decay in the HFSD cardiomyocytes. The time constant of Ca^{2+} transient decay increased with loading and

further with stretching – a feature not evident in CTRL cardiomyocytes which provides additional

mechanistic insight (discussion below).

The final pacing frequency protocol was implemented to apply a 'stress test' to the loaded cardiomyocytes

with the expectation of elevated diastolic Ca^{2+}, potentially revealing a cross-bridge stiffness component in

HFSD cardiomyocytes. With a doubling of pacing frequency, a significant increase in the stress/strain

gradient was observed in HFSD cardiomyocytes (only), and a similar reduction in diastolic sarcomere length

in both CTRL and HFSD was observed. While cardiomyocyte diastolic Ca^{2+} level increased in both CTRL and

HFSD – the HFSD increase was less marked than CTRL. This was a surprising finding, not previously

reported in any other model setting. A suggestion of Ca^{2+} hyper-sensitivity arises (Davis *et al.*, 2007)

amongst other explanations to consider.

Mechanistic insights into cross-bridge derived stiffness

In cardiometabolic disease, a cardiomyocyte stiffness contribution from intra-myocyte structural/passive

sources is undoubtedly substantial. In the intact cell, all this structural architecture is preserved, and all the

structures are vulnerable to functionally damaging post translational modifications in the free radical

myocyte soup. Titin, as the largest coded protein in the genome with hundreds of modifiable residue sites

(regulatory and non-regulatory), receives much attention. Isoform variants, and a multitude of polymorphic

genotypes likely also contribute to the stiffness signature. With this model, clearly the extracellular matrix

is not a direct player, although an effect of disrupted membrane-spanning integrins cannot be precluded.

Here we report that a cardiometabolic disease phenotype induced by obesogenic diet includes a 35%

increase in diastolic dysfunction (by echo E/e') which relates to a mean 70% increase in loaded

cardiomyocyte stiffness modulus. Several lines of evidence emerging from this study suggest that Ca^{2+} -

related mechanism/s may contribute to cardiomyocyte stiffness increase. Specifically, in the loaded HFSD

cardiomyocyte (vs CTRL):

a) load confers accentuated lengthening rate deficit

b) with stretch, the Ca^{2+} transient decay time course is prolonged

c) with more rapid pacing diastolic stiffness increases and sarcomere length decreases, although diastolic

Ca^{2+} is less elevated.

Together these cardiomyocyte phenotypes would be consistent with selective impairment of relaxation

rate, selective myofilament hyper-activation, impeded diastolic filling and reduced diastolic reserve.

This combination of observations is consistent with the notion that length-dependent cross-bridge

attachment is prolonged in HFSD cardiomyocytes (or conversely detachment is delayed). An alteration in

the 3D geometry of the low affinity Ca^{2+} binding pocket would produce this effect, by stabilizing Ca^{2+} in the

pocket for longer every activation cycle. A manifestation of this altered binding function could be apparent

Ca^{2+} hypersensitivity. The Ca^{2+} dissociation rate has been theorized as an effector of the transient decay

time-course (Robinson *et al.*, 2018). A role for the troponin complex in regulation of Ca^{2+} affinity is well

established (Chemla *et al.*, 2000). Further evidence of structural modifications of specific regions of TnI and

TnC which form and regulate this pocket is required. Our mass spectrometry studies provide proof of

concept of relevant post-translational modification types (Janssens *et al.*, 2018).

Limitations

Given the diastolic dysfunction goals of this study, investigations have primarily focused on cardiomyocyte

relaxation and lengthening behaviors. Diastolic and systolic interactions are important and additional

protocols are in development for these studies. In this regard, we defer to the extensive body of

developmental research undertaken to develop software and instrumentation to perform single

cardiomyocyte 'Pressure-Volume (PV) loops' (Helmes *et al.*, 2016). In their approach, which mimics the PV

loops in the intact myocardium, stretch is applied after relaxation has occurred by using a force clamp to

maintain a predefined diastolic force level. This approach has been used to discover that titin truncation

shifts the working sarcomere length range preserving diastolic stiffness at the expense of systolic

performance (Methawasin *et al.*, 2022) and that highly compliant titin allows cardiomyocytes to function at

high diastolic Ca^{2+} levels (Najafi *et al.*, 2019). Conceivably, this approach could also be used to evaluate the

temporal relationship between slow or incomplete relaxation and cardiomyocyte cross-bridge derived

stiffness in a semi-situ manner. Operator expertise is imperative with this system which to date has been

very productive in the hands of the developers.

In our work we have adopted a pragmatic approach, simpler to implement – although still very challenging

with about 20% isotonic-to-loaded myocyte conversion success rate. When animals have many months of

feeding history and cell isolation from older and diseased hearts is difficult, an optimized loading and

stretching approach is key. The two approaches have not been benchmarked, although it can be predicted

that the viscous components of stiffness are likely underestimated when stretches are slow and relatively

small such as in this study (Caporizzo & Prosser, 2021). The capacity to directly and reproducibly determine

Youngs Modulus (stress/strain) with a slow stretch routine has useful comparative value.

With ongoing research, our priority is to undertake longitudinal studies, including both sexes, to investigate

the relative timing of the emergence of *in vivo* and *in vitro* stiffness phenotypes as differences in Ca^{2+}

handling have previously been observed between control male and female rodents (Farrell *et al.*, 2010).

This has high translational relevance as we have previously identified a differential time course of diastolic

dysfunction development in females and males, with increased female vulnerability (Reichelt *et al.*, 2013;

Chandramouli *et al.*, 2018).

In conclusion

This is the first study to link elevated intact single cardiomyocyte stiffness with *in vivo* diastolic dysfunction

in a diet-induced cardiometabolic disease mouse model. The novel findings identify augmented load

dependence and reduced diastolic reserve as fundamental features of cardiomyocytes derived from hearts

exhibiting cardiometabolic diastolic dysfunction. Taken together, the findings emphasise the importance of

mechanical loading and pacing frequency in revealing latent diastolic abnormalities at the cardiomyocyte

level. The rationale for further studies identifying specific subcellular and molecular mechanisms of

cardiomyocyte stiffness in cardiometabolic disease prior to failure onset is compelling.

REFERENCES

Abdellatif M et al. (2021). Nicotinamide for the treatment of heart failure with preserved ejection fraction. *Sci Transl Med* **13,** eabd7064.

Aurigemma GP, Zile MR & Gaasch WH (2006). Contractile Behavior of the Left Ventricle in Diastolic Heart Failure. *Circulation* **113,** 296–304.

Borlaug BA, Melenovsky V, Redfield MM, Kessler K, Chang H-J, Abraham TP & Kass DA (2007). Impact of Arterial Load and Loading Sequence on Left Ventricular Tissue Velocities in Humans. *Journal of the American College of Cardiology* **50,** 1570–1577.

Caporizzo MA & Prosser BL (2021). Need for Speed: The Importance of Physiological Strain Rates in Determining Myocardial Stiffness. *Front Physiol* **12,** 696694.

Chandramouli C, Reichelt ME, Curl CL, Varma U, Bienvenu LA, Koutsifeli P, Raaijmakers AJA, De Blasio MJ, Qin CX, Jenkins AJ, Ritchie RH, Mellor KM & Delbridge LMD (2018). Diastolic dysfunction is more apparent in STZ-induced diabetic female mice, despite less pronounced hyperglycemia. *Sci Rep* **8,** 2346.

Chemla D, Coirault C, Hébert J-L & Lecarpentier Y (2000). Mechanics of Relaxation of the Human Heart. *Physiology* **15,** 78–83.

Clark JA & Campbell SG (2019). Diverse relaxation rates exist among rat cardiomyocytes isolated from a single myocardial region. *J Physiol* **597,** 711–722.

Curl CL, Danes VR, Bell JR, Raaijmakers AJA, Ip WTK, Chandramouli C, Harding TW, Porrello ER, Erickson JR, Charchar FJ, Kompa AR, Edgley AJ, Crossman DJ, Soeller C, Mellor KM, Kalman JM, Harrap SB & Delbridge LMD (2018). Cardiomyocyte Functional Etiology in Heart Failure With Preserved Ejection Fraction Is Distinctive-A New Preclinical Model. *J Am Heart Assoc* **7,** e007451.

Daniels LJ, Macindoe C, Koutsifeli P, Annandale M, Raaijmakers AJA, Weeks KL, Bell JR, Janssens JV, Curl CL, Delbridge LMD & Mellor KM (2022). Myocardial deformation imaging by 2D speckle tracking echocardiography for assessment of diastolic dysfunction in murine cardiopathology. 2022.08.06.503061. Available at: https://www.biorxiv.org/content/10.1101/2022.08.06.503061v1 [Accessed May 21, 2023].

Davis J, Wen H, Edwards T & Metzger JM (2007). Thin Filament Disinhibition by Restrictive Cardiomyopathy Mutant R193H Troponin I Induces Ca2+-Independent Mechanical Tone and Acute Myocyte Remodeling. *Circulation Research* **100,** 1494–1502.

Delbridge LM & Roos KP (1997). Optical methods to evaluate the contractile function of unloaded isolated cardiac myocytes. *J Mol Cell Cardiol* **29,** 11–25.

Delbridge LMD, Bell JR, Weeks KL, Raaijmakers AJA & Mellor KM (2022). HFpEF etiology - Can focus on sex-specific mechanisms deliver insights for all? *J Mol Cell Cardiol* **173,** 71–72.

Dia M, Gomez L, Thibault H, Tessier N, Leon C, Chouabe C, Ducreux S, Gallo-Bona N, Tubbs E, Bendridi N, Chanon S, Leray A, Belmudes L, Couté Y, Kurdi M, Ovize M, Rieusset J & Paillard M (2020). Reduced reticulum–mitochondria Ca2+ transfer is an early and reversible trigger of mitochondrial dysfunctions in diabetic cardiomyopathy. *Basic Res Cardiol* **115,** 74.

Dowrick JM, Tran K, Garrett AS, Anderson AJ, Nielsen PMF, Taberner AJ & Han J-C (2022). Work-loop contractions reveal that the afterload-dependent time course of cardiac Ca2+ transients is modulated by preload. *Journal of Applied Physiology* **133,** 663–675.

Eichhorn EJ, Willard JE, Alvarez L, Kim AS, Glamann DB, Risser RC & Grayburn PA (1992). Are contraction
and relaxation coupled in patients with and without congestive heart failure? *Circulation* **85,** 2132–
2139.

Eisner DA, Caldwell JL, Trafford AW & Hutchings DC (2020). The Control of Diastolic Calcium in the Heart.
Circ Res **126,** 395–412.

Farrell SR, Ross JL & Howlett SE (2010). Sex differences in mechanisms of cardiac excitation-contraction
coupling in rat ventricular myocytes. *Am J Physiol Heart Circ Physiol* **299,** H36-45.

Fowler ED, Benoist D, Drinkhill MJ, Stones R, Helmes M, Wüst RCI, Stienen GJM, Steele DS & White E (2015).
Decreased creatine kinase is linked to diastolic dysfunction in rats with right heart failure induced
by pulmonary artery hypertension. *J Mol Cell Cardiol* **86,** 1–8.

Gillebert TC, Leite-Moreira AF & De Hert SG (2000). Load dependent diastolic dysfunction in heart failure.
Heart Fail Rev **5,** 345–355.

Gonçalves N, Silva AF, Rodrigues PG, Correia E, Moura C, Eloy C, Roncon-Albuquerque R, Falcão-Pires I &
Leite-Moreira AF (2016). Early cardiac changes induced by a hypercaloric Western-type diet in
"subclinical" obesity. *American Journal of Physiology-Heart and Circulatory Physiology* **310,** H655–
H666.

Hamdani N, Franssen C, Lourenço A, Falcão-Pires I, Fontoura D, Leite S, Plettig L, López B, Ottenheijm CA,
Becher PM, González A, Tschöpe C, Díez J, Linke WA, Leite-Moreira AF & Paulus WJ (2013).
Myocardial Titin Hypophosphorylation Importantly Contributes to Heart Failure With Preserved
Ejection Fraction in a Rat Metabolic Risk Model. *Circulation: Heart Failure* **6,** 1239–1249.

Heather LC, Hafstad AD, Halade GV, Harmancey R, Mellor KM, Mishra PK, Mulvihill EE, Nabben M,
Nakamura M, Rider OJ, Ruiz M, Wende AR & Ussher JR (2022). Guidelines on models of diabetic
heart disease. *Am J Physiol Heart Circ Physiol* **323,** H176–H200.

van Heerebeek L, Borbély A, Niessen HWM, Bronzwaer JGF, van der Velden J, Stienen GJM, Linke WA,
Laarman GJ & Paulus WJ (2006). Myocardial Structure and Function Differ in Systolic and Diastolic
Heart Failure. *Circulation* **113,** 1966–1973.

Helmes M, Najafi A, Palmer BM, Breel E, Rijnveld N, Iannuzzi D & van der Velden J (2016). Mimicking the
cardiac cycle in intact cardiomyocytes using diastolic and systolic force clamps; measuring power
output. *Cardiovasc Res* **111,** 66–73.

Hopf A-E, Andresen C, Kötter S, Isić M, Ulrich K, Sahin S, Bongardt S, Röll W, Drove F, Scheerer N,
Vandekerckhove L, De Keulenaer GW, Hamdani N, Linke WA & Krüger M (2018). Diabetes-Induced
Cardiomyocyte Passive Stiffening Is Caused by Impaired Insulin-Dependent Titin Modification and
Can Be Modulated by Neuregulin-1. *Circulation Research* **123,** 342–355.

Howlett SE, Grandy SA & Ferrier GR (2006). Calcium spark properties in ventricular myocytes are altered in
aged mice. *American Journal of Physiology-Heart and Circulatory Physiology* **290,** H1566–H1574.

Jackson AM et al. (2022). Diabetes and pre-diabetes in patients with heart failure and preserved ejection
fraction. *Eur J Heart Fail* **24,** 497–509.

Janssens JV, Ma B, Brimble MA, Van Eyk JE, Delbridge LMD & Mellor KM (2018). Cardiac troponins may be
irreversibly modified by glycation: novel potential mechanisms of cardiac performance modulation.
Sci Rep **8,** 16084.

Janssens JV, Raaijmakers AJA, Weeks KL, Bell JR, Mellor KM, Curl CL & Delbridge LMD (2024). The cardiomyocyte origins of diastolic dysfunction: cellular components of myocardial stiffness. *Am J Physiol Heart Circ Physiol*; DOI: 10.1152/ajpheart.00334.2023.

Kamon T, Kaneko H, Itoh H, Kiriyama H, Koyama K, Fujiu K, Morita H, Uno K, Hayashi N & Komuro I (2020). Association between insulin resistance and left ventricular diastolic dysfunction in non-diabetic general population. *European Heart Journal* **41,** ehaa946.3057.

Kilfoil PJ, Lotteau S, Zhang R, Yue X, Aynaszyan S, Solymani RE, Cingolani E, Marbán E & Goldhaber JI (2020). Distinct features of calcium handling and β-adrenergic sensitivity in heart failure with preserved versus reduced ejection fraction. *J Physiol* **598,** 5091–5108.

Kittleson MM, Panjrath GS, Amancherla K, Davis LL, Deswal A, Dixon DL, Januzzi JL & Yancy CW (2023). 2023 ACC Expert Consensus Decision Pathway on Management of Heart Failure With Preserved Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. *Journal of the American College of Cardiology* **81,** 1835–1878.

Lab MJ, Allen DG & Orchard CH (1984). The effects of shortening on myoplasmic calcium concentration and on the action potential in mammalian ventricular muscle. *Circulation Research* **55,** 825–829.

Lam CSP, Arnott C, Beale AL, Chandramouli C, Hilfiker-Kleiner D, Kaye DM, Ky B, Santema BT, Sliwa K & Voors AA (2019). Sex differences in heart failure. *European Heart Journal* **40,** 3859–3868c.

Leopoldo AS, Sugizaki MM, Lima-Leopoldo AP, do Nascimento AF, de Azevedo Melo Luvizotto R, de Campos DHS, Okoshi K, Pai-Silva MD, Padovani CR & Cicogna AC (2010). Cardiac remodeling in a rat model of diet-induced obesity. *Can J Cardiol* **26,** 423–429.

Leung M, Phan V, Whatmough M, Heritier S, Wong VW & Leung DY (2015). Left ventricular diastolic reserve in patients with type 2 diabetes mellitus. *Open Heart* **2,** e000214.

Lo Q & Thomas L (2010). Echocardiographic evaluation of diastolic heart failure. *Australas J Ultrasound Med* **13,** 14–26.

Loescher CM, Freundt JK, Unger A, Hessel AL, Kühn M, Koser F & Linke WA (2023). Titin governs myocardial passive stiffness with major support from microtubules and actin and the extracellular matrix. *Nat Cardiovasc Res* **2,** 991–1002.

Louch WE (2020). A horse of a different colour: distinct mechanisms of HFpEF and HFrEF. *J Physiol* **598,** 5005–5006.

Luo M & Anderson ME (2013). Mechanisms of altered Ca^{2+} handling in heart failure. *Circ Res* **113,** 690–708.

McHugh K, DeVore AD, Wu J, Matsouaka RA, Fonarow GC, Heidenreich PA, Yancy CW, Green JB, Altman N & Hernandez AF (2019). Heart Failure With Preserved Ejection Fraction and Diabetes: JACC State-of-the-Art Review. *J Am Coll Cardiol* **73,** 602–611.

Mellor KM, Bell JR, Young MJ, Ritchie RH & Delbridge LMD (2011). Myocardial autophagy activation and suppressed survival signaling is associated with insulin resistance in fructose-fed mice. *Journal of Molecular and Cellular Cardiology* **50,** 1035–1043.

Mellor KM, Wendt IR, Ritchie RH & Delbridge LMD (2012). Fructose diet treatment in mice induces fundamental disturbance of cardiomyocyte Ca2+ handling and myofilament responsiveness. *Am J Physiol Heart Circ Physiol* **302,** H964-972.

Methawasin M, Farman GP, Granzier-Nakajima S, Strom J, Kiss B, Smith JE & Granzier H (2022). Shortening
the thick filament by partial deletion of titin's C-zone alters cardiac function by reducing the
operating sarcomere length range. *J Mol Cell Cardiol* **165,** 103–114.

Mirsky I (1976). Assessment of passive elastic stiffness of cardiac muscle: mathematical concepts,
physiologic and clinical considerations, directions of future research. *Prog Cardiovasc Dis* **18,** 277–
308.

Najafi A, van de Locht M, Schuldt M, Schönleitner P, van Willigenburg M, Bollen I, Goebel M, Ottenheijm
CAC, van der Velden J, Helmes M & Kuster DWD (2019). End-diastolic force pre-activates
cardiomyocytes and determines contractile force: role of titin and calcium. *J Physiol* **597,** 4521–
4531.

Natali AJ, Turner DL, Harrison SM & White E (2001). Regional effects of voluntary exercise on cell size and
contraction-frequency responses in rat cardiac myocytes. *Journal of Experimental Biology* **204,**
1191–1199.

Patil VC, Patil HV, Shah KB, Vasani JD & Shetty P (2011). Diastolic dysfunction in asymptomatic type 2
diabetes mellitus with normal systolic function. *J Cardiovasc Dis Res* **2,** 213–222.

Peyronnet R, Bollensdorff C, Capel RA, Rog-Zielinska EA, Woods CE, Charo DN, Lookin O, Fajardo G, Ho M,
Quertermous T, Ashley EA & Kohl P (2017). Load-dependent effects of apelin on murine
cardiomyocytes. *Prog Biophys Mol Biol* **130,** 333–343.

Pulinilkunnil T, Kienesberger PC, Nagendran J, Sharma N, Young ME & Dyck JRB (2014). Cardiac-specific
adipose triglyceride lipase overexpression protects from cardiac steatosis and dilated
cardiomyopathy following diet-induced obesity. *Int J Obes* **38,** 205–215.

Redfield MM (2016). Heart Failure with Preserved Ejection Fraction. *N Engl J Med* **375,** 1868–1877.

Reichelt ME, Mellor KM, Bell JR, Chandramouli C, Headrick JP & Delbridge LMD (2013). Sex, sex steroids,
and diabetic cardiomyopathy: making the case for experimental focus. *Am J Physiol Heart Circ*
Physiol **305,** H779-792.

Robinson P, Liu X, Sparrow A, Patel S, Zhang Y-H, Casadei B, Watkins H & Redwood C (2018). Hypertrophic
cardiomyopathy mutations increase myofilament Ca2+ buffering, alter intracellular Ca2+ handling,
and stimulate Ca2+-dependent signaling. *J Biol Chem* **293,** 10487–10499.

Rozenbaum Z, Topilsky Y, Khoury S, Pereg D & Laufer-Perl M (2019). Association of body mass index and
diastolic function in metabolically healthy obese with preserved ejection fraction. *Int J Cardiol* **277,**
147–152.

Satoh H, Delbridge LM, Blatter LA & Bers DM (1996). Surface:volume relationship in cardiac myocytes
studied with confocal microscopy and membrane capacitance measurements: species-dependence
and developmental effects. *Biophys J* **70,** 1494–1504.

Shimkunas R et al. (2021). Mechanical Load Regulates Excitation-Ca2+ Signaling-Contraction in
Cardiomyocyte. *Circ Res* **128,** 772–774.

Sotomi Y et al. (2021). Sex Differences in Heart Failure With Preserved Ejection Fraction. *J Am Heart Assoc*
10, e018574.

Sowers JR, Habibi J, Jia G, Bostick B, Manrique-Acevedo C, Lastra G, Yang Y, Chen D, Sun Z, Domeier TL,
Durante W, Whaley-Connell AT, Hill MA, Jaisser F, DeMarco VG & Aroor AR (2020). Endothelial

sodium channel activation promotes cardiac stiffness and diastolic dysfunction in Western diet fed
female mice. *Metabolism* **109,** 154223.

Varian KD & Janssen PML (2007). Frequency-dependent acceleration of relaxation involves decreased
myofilament calcium sensitivity. *American Journal of Physiology-Heart and Circulatory Physiology*
292, H2212–H2219.

Villalobos Lizardi JC, Baranger J, Nguyen MB, Asnacios A, Malik A, Lumens J, Mertens L, Friedberg MK,
Simmons CA, Pernot M & Villemain O (2022). A guide for assessment of myocardial stiffness in
health and disease. *Nat Cardiovasc Res* **1,** 8–22.

Wells SP, Raaijmakers AJA, Curl CL, O'Shea C, Hayes S, Mellor KM, Kalman JM, Kirchhof P, Pavlovic D,
Delbridge LMD & Bell JR (2023). Localized cardiomyocyte lipid accumulation is associated with
slowed epicardial conduction in rats. *J Gen Physiol* **155,** e202213296.

White E, Boyett MR & Orchard CH (1995). The effects of mechanical loading and changes of length on single
guinea-pig ventricular myocytes. *The Journal of Physiology* **482,** 93–107.

Wingard MC, Dalal S, Shook PL, Myers R, Connelly BA, Thewke DP, Singh M & Singh K (2021). Deficiency of
ataxia-telangiectasia mutated kinase modulates functional and biochemical parameters of the
heart in response to Western-type diet. *Am J Physiol Heart Circ Physiol* **320,** H2324–H2338.

Zhou P & Pu WT (2016). Recounting Cardiac Cellular Composition. *Circulation Research* **118,** 368–370.

 FIGURE LEGENDS

Fig.1 Systemic indicators of cardiometabolic disease and *in vivo* diastolic function in high-fat/sugar diet (HFSD) vs control (CTRL) mice.

A. Body weight elevation at 5 week intervals up to 25 weeks of dietary treatment (last timepoint full cohort intact. N = 16 – 17 mice per group).

B. Glucose tolerance testing at 23-24 weeks post dietary intervention; N = 6 mice per group).

C. Echocardiography exemplar blood flow and tissue doppler images from which diastolic indices were derived for CTRL and HFSD mice. Upper panels: HFSD mitral blood flow & annular wall motion. Lower panels: CTRL mitral blood flow & annular wall motion. (N = 9-16 mice per group)

D. Early diastolic mitral annular blood flow (E) was increased in HFSD.

E. Early diastolic mitral annular wall velocity (e') was decreased in HFSD.

F. Deceleration time of early diastolic mitral annular blood flow was decreased in HFSD.

G. Ratio of early diastolic blood flow to wall velocity (E/e') was increased in HFSD.

Time-course data analysed using two-way repeated measures ANOVA with Sidak post-hoc test ($*p < 0.05$). Comparisons between two treatment groups performed using student's t-test; $*p<0.05$, $**p<0.01$, $***p<0.001$). Data presented as mean ± standard error of the mean.

Fig.2 Comparing non-loaded sarcomere shortening and lengthening in CTRL and HFSD cardiomyocytes.

A. Mean sarcomere contraction cycles presented as a percentage of diastolic sarcomere length for CTRL and HFSD cardiomyocytes

B-C. Mean diastolic sarcomere length and peak sarcomere shortening for CTRL and HFSD cardiomyocytes.

D. Mean sarcomere shortening and lengthening rate records (time derivative of sarcomere contraction-relaxation cycle) for CTRL and HFSD cardiomyocytes. The nadir and zenith represent maximal shortening and lengthening rates respectively.

E-F. Maximal rate of lengthening and shortening rates for CTRL and HFSD cardiomyocytes.

Mean data computed from average of 20 cycles per cardiomyocyte to derive group mean (N = 7-8 mice per group, n = 38-47 cells per group). Comparisons between two treatment groups performed using student's t-test (all $p>0.05$, ns). Data presented as mean ± standard error of the mean.

Fig. 3 Ca^{2+} signals in non-loaded (isotonic) paced cardiomyocytes.

A-B. Mean diastolic Ca^{2+} levels and time constant of transient decay (*tau*) in HFSD and CTRL cardiomyocytes.

C-D. Mean systolic Ca^{2+} levels and amplitude of the Ca^{2+} transient for CTRL and HFSD.

E. Mean Ca^{2+} transient records for CTRL and HFSD cardiomyocytes.

Mean data computed from average of 20 cycles per cardiomyocyte to derive group mean (N = 6-8 mice per group, n = 32-37 cells per group). Comparisons between treatment groups were performed using Student's t-test; $*p<0.05$). Data presented as mean ± standard error of the mean.

Fig. 4 Sarcomere shortening and lengthening in non-loaded (isotonic) and loaded (auxotonic) pacing states.

A. Exemplar non-loaded (isotonic) and loaded (auxotonic) sarcomere contraction cycles for the same cardiomyocyte (CTRL).

B-C. Peak sarcomere shortening and maximum rate of lengthening, measured for each cardiomyocyte in non-loaded (isotonic) and loaded (auxotonic) state for CTRL and HFSD groups combined (N = 9 mice, n = 25 cells). Mean data for CTRL and HFSD are presented as grey points to the left (non-loaded) and right (loaded) of individual cardiomyocyte data.

D. Correlation of loading associated peak sarcomere shortening change and maximum rate of lengthening change for CTRL and HFSD combined.

E. Mean sarcomere loading effect on maximum lengthening rate (μm/s) normalized by shortening extent (μm) for CTRL vs HFSD. (CTRL: N = 5 mice, n = 13 cells; HFSD: N = 4 mice, n = 12 cells). Significant ANOVA interaction effect (diet treatment x load state).

F. Mean decrement in maximum sarcomere lengthening rate (μm/s) normalized by shortening extent (μm).

Comparison of cardiomyocyte responses in different loading conditions analysed by paired t-test. Comparison of cardiomyocytes from different treatment groups analysed by unpaired t-test. A 2-way ANOVA with repeated measures used for analysis of two independent variables Data fitted with linear regression model and calculation of Pearson's correlation coefficient. Data presented as mean ± standard error of the mean. $*p<0.05$, $**p<0.01$, $****p<0.0001$, $\#p<0.05$ ANOVA factor interaction.

Fig. 5 Quantifying diastolic stiffness in CTRL vs HFSD cardiomyocytes using strain normalization methods

A. Exemplar stress recordings of CTRL and HFSD cardiomyocytes undergoing 'stress-length' protocol (2Hz paced). Diastolic baseline stress shifts indicated by broken line.

B. Diastolic and systolic sarcomere mean stress-length relations in CTRL cardiomyocytes (N = 5 mice, n = 9 cells).

C. Diastolic sarcomere mean stress-length relation (linear regression fit) for HFSD and CTRL.

D. Diastolic stress (mN/mm^2) normalized relative to sarcomere stretch (μm) yields sarcomere stress/strain = stiffness [(mN/mm^2) /strain (μm)] for HFSD and CTRL.

E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL..

F. Diastolic stress (mN/mm^2) normalized relative to segment stretch (% length) yields cardiomyocyte segment stress/strain = stiffness [(mN/mm^2) /strain (% length)] for HFSD & CTRL.

G. Correlation (linear regression fit) of *in vitro* cardiomyocyte segment diastolic stiffness (stress/strain) with *in vivo* diastolic dysfunction E/e' (blood flow/wall motion) in CTRL and HFSD. Cardiomyocyte data averaged per heart.

Comparisons between two groups performed using student's t-test ($*p<0.05$, $**p<0.01$). Data fitted with linear regression model and calculation of Pearson's correlation coefficient, presented as mean ± standard error of the mean. For **C. & D.** (CTRL: N=5 mice, n=9 cells; HFSD: N=2 mice, n=7 cells). For **E.&F.** (CTRL: N=6 mice, n = 16 cells; HFSD: N=6 mice, n=18 cells).

Fig. 6 Load dependence of cardiomyocyte Ca^{2+} transient decay in CTRL and HFSD cardiomyocytes.

A-B. Exemplar Ca^{2+} transient decay records, with a mono-exponential decay function fit, in non-loaded
(isotonic) and loaded (auxotonic, stretched) CTRL and HFSD cardiomyocytes. Horizontal black broken lines
intersect with mono-exponential fit at 36.8% (1/e) transient amplitude. Vertical broken lines from
intersection points indicate the time constant of Ca^{2+} transient decay (tau).

C. Time constant of Ca^{2+} transient decay (tau) in cardiomyocytes with different load conditions: auxotonic
(aux) & stretched (str) shown as fold change relative to non-loaded isotonic (iso) for CTRL and HFSD.
912

For analysis of two independent variables a 2-way ANOVA with repeated measures used (CTRL: N = 5 mice,
n = 13 cells; HFSD: N = 4 mice, n = 12 cells). Data are presented as mean ± standard error of the mean.
***p<0.001 (HFSD stretched vs non-loaded isotonic). Non-significant ANOVA interaction effect (p = 0.055,
diet treatment x load state).

Fig. 7 Pacing induced changes in cardiomyocyte cytosolic Ca^{2+} levels and stiffness in CTRL & HFSD.
A. Exemplar CTRL and HFSD cardiomyocyte Ca^{2+} transients at 2 and 4 Hz pacing.
B. Diastolic cardiomyocyte Ca^{2+} levels measured before and after transition from 2 to 4 Hz pacing in HFSD &
CTRL.
C. Diastolic cardiomyocyte Ca^{2+} level mean shift with transition from 2 to 4 Hz pacing in HFSD & CTRL.
D. Diastolic sarcomere length measured before and after transition from 2 to 4 Hz pacing in HFSD & CTRL.
E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL at
2 and 4 Hz pacing.
F. Diastolic mean cardiomyocyte segment stress-length relation slope (mN/mm^2) normalized relative to
segment stretch (% length) yields cardiomyocyte mean segment stress/strain = stiffness [(mN/mm^2) /strain
(% length)] for HFSD and CTRL at 2 and 4 Hz pacing.

Comparison between two groups performed using unpaired t-test. With two independent variables a
repeated measures 2-way ANOVA Sidaks post-hoc analysis was performed (*p<0.05, **p<0.01,
****p<0.0001). Data are presented as mean ± standard error of the mean or mean with paired individual
points. For **B. & C.** (CTRL: N = 4 mice, n = 9 cells; HFSD: N = 4 mice, n = 8 cells). For **D. & E.** (CTRL: N = 4 mice,
n = 7 cells; HFSD: N = 4 mice, n = 9 cells). For **F.** (CTRL: N = 4 mice, n = 8 cells; HFSD: N = 4 mice, n = 9 cells).
936

Table S1. Echocardiographic parameters in CTRL and HFSD mice.
LVIDd left ventricular internal diameter at end diastole, *IVSd* interventricular septum diameter at end
diastole, *LVPWd* left ventricular posterior wall thickness at end diastole, *RWT* relative wall thickness, *E*
wave mitral valve blood flow velocity during the early ventricular filling phase, *A wave* mitral valve blood
flow velocity during atrial contraction, *e' wave* mitral annulus tissue motion velocity during the early
ventricular filling phase, *a' wave* mitral annulus tissue motion velocity during atrial contraction, *MV Dec T*
mitral valve deceleration time.
944
945 Comparisons between two treatment groups performed using Students' unpaired t-test (N = 9-15 mice per
946 group; *$p < 0.05$, **$p<0.01$, ***$p<0.001$). Data are presented as mean ± standard error of the mean.
947
948
Fig. S1. Cardiomyocyte standardized stretch protocol.
A. Image of a left ventricular cardiomyocyte longitudinally attached to glass rods. Edge-tracking used to
measure inter-rod distance (marked as blue bars) for cardiomyocyte segment analyses. The delineated
region of interest (marked pink) represents the area from which Fast Fourier Transform analysis of
sarcomere striations was performed to derive sarcomere-specific data.
B. Exemplar traces depict the change in glass rod internal distance in response to successive 0.5V piezo
motor inputs used for calibration of stretch protocol for each individual cardiomyocyte to allow normalized
comparisons.
C. Measuring stress and strain during progressive cardiomyocyte stretch. Exemplar force, sarcomere length
and rod edge-tracker traces during the first stretch of a stress-length protocol.
D. Comparison of variability of cell stretches performed using a uniform (preset) stretch constant and the
individually derived stretch calibration constant. Reproducibility is optimized using the individual stretch
calibration constant. Data shown as median ± interquartile range.

Fig.1 Systemic indicators of cardiometabolic disease and *in vivo* diastolic function in high-fat/sugar diet (HFSD) vs control (CTRL) mice.

A. Body weight elevation at 5 week intervals up to 25 weeks of dietary treatment (last timepoint full cohort intact. N = 16 – 17 mice per group).

B. Glucose tolerance testing at 23-24 weeks post dietary intervention; N = 6 mice per group).

C. Echocardiography exemplar blood flow and tissue doppler images from which diastolic indices were derived for CTRL and HFSD mice. Upper panels: HFSD mitral blood flow & annular wall motion. Lower panels: CTRL mitral blood flow & annular wall motion. (N = 14-16 mice per group)

D. Early diastolic mitral annular blood flow (E) was increased in HFSD.

E. Early diastolic mitral annular wall velocity (e') was decreased in HFSD.

F. Deceleration time of early diastolic mitral annular blood flow was decreased in HFSD.

G. Ratio of early diastolic blood flow to wall velocity (E/e') was increased in HFSD.

Time-course data analysed using two-way repeated measures ANOVA with Sidak post-hoc test (*p < 0.05). Comparisons between two treatment groups performed using student's t-test; *p<0.05, **p<0.01, ***p<0.001). Data presented as mean ± standard error of the mean.

Fig.2 Comparing non-loaded sarcomere shortening and lengthening in CTRL and HFSD cardiomyocytes.
A. Mean sarcomere contraction cycles presented as a percentage of diastolic sarcomere length for CTRL and HFSD cardiomyocytes
B-C. Mean diastolic sarcomere length and peak sarcomere shortening for CTRL and HFSD cardiomyocytes.
D. Mean sarcomere shortening and lengthening rate records (time derivative of sarcomere contraction-relaxation cycle) for CTRL and HFSD cardiomyocytes. The nadir and zenith represent maximal shortening and lengthening rates respectively.
E-F. Maximal rate of lengthening and shortening rates for CTRL and HFSD cardiomyocytes.

Mean data computed from average of 20 cycles per cardiomyocyte to derive group mean (N = 7-8 mice per group, n = 38-47 cells per group). Comparisons between two treatment groups performed using student's t-test (all p>0.05, ns). Data presented as mean ± standard error of the mean.

Fig. 3 Ca²⁺ signals in non-loaded (isotonic) paced cardiomyocytes.
A-B. Mean diastolic Ca²⁺ levels and time constant of transient decay (*tau*) in HFSD and CTRL cardiomyocytes.
C-D. Mean systolic Ca²⁺ levels and amplitude of the Ca²⁺ transient for CTRL and HFSD.
E. Mean Ca²⁺ transient records for CTRL and HFSD cardiomyocytes.

Mean data computed from average of 20 cycles per cardiomyocyte to derive group mean (N = 6-8 mice per group, n = 32-37 cells per group). Comparisons between treatment groups were performed using Student's t-test; *p<0.05). Data presented as mean ± standard error of the mean.

Fig. 4 Sarcomere shortening and lengthening in non-loaded (isotonic) and loaded (auxotonic) pacing states.

A. Exemplar non-loaded (isotonic) and loaded (auxotonic) sarcomere contraction cycles for the same cardiomyocyte (CTRL).

B-C. Peak sarcomere shortening and maximum rate of lengthening, measured for each cardiomyocyte in non-loaded (isotonic) and loaded (auxotonic) state for CTRL and HFSD groups combined (N = 9 mice, n = 25 cells). Mean data for CTRL and HFSD are presented as grey points to the left (non-loaded) and right (loaded) of individual cardiomyocyte data.

D. Correlation of loading associated peak sarcomere shortening change and maximum rate of lengthening change for CTRL and HFSD combined.

E. Mean sarcomere loading effect on maximum lengthening rate (µm/s) normalized by shortening extent (µm) for CTRL vs HFSD. (CTRL: N = 5 mice, n = 13 cells; HFSD: N = 4 mice, n = 12 cells). Significant ANOVA interaction effect (diet treatment x load state).

F. Mean decrement in maximum sarcomere lengthening rate (µm/s) normalized by shortening extent (µm).

Comparison of cardiomyocyte responses in different loading conditions analysed by paired t-test. Comparison of cardiomyocytes from different treatment groups analysed by unpaired t-test. A 2-way ANOVA with repeated measures used for analysis of two independent variables. Data fitted with linear regression model and calculation of Pearson's correlation coefficient. Data presented as mean ± standard error of the mean. *p<0.05, **p<0.01, ****p<0.0001, #p<0.05 ANOVA factor interaction.

A Cardiomyocyte stress during calibrated stretch steps (paced 2Hz)

B Stress-Length Relations (CTRL)

Fig. 5 Cardiomyocyte stress recordings and stress-length relations using strain normalization methods
A. Exemplar stress recordings of CTRL and HFSD cardiomyocytes undergoing 'stress-length' protocol (2Hz paced). Diastolic baseline stress shifts indicated by broken line.
B. Diastolic and systolic sarcomere mean stress-length relations in CTRL cardiomyocytes (N = 5 mice, n = 9 cells).

Data fitted with linear regression model. Data presented as mean ± standard error of the mean.

Fig. 6 Quantifying diastolic stiffness in CTRL vs HFSD cardiomyocytes using strain normalization methods (cont).

A. Diastolic sarcomere mean stress-length relation (linear regression fit) for HFSD and CTRL.

B. Diastolic stress (mN/mm^2) normalized relative to sarcomere stretch (μm) yields sarcomere stress/strain = stiffness [(mN/mm^2) /strain (μm)] for HFSD and CTRL.

C. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL..

D. Diastolic stress (mN/mm^2) normalized relative to segment stretch (% length) yields cardiomyocyte segment stress/strain = stiffness [(mN/mm^2) /strain (% length)] for HFSD & CTRL.

E. Correlation (linear regression fit) of *in vitro* cardiomyocyte segment diastolic stiffness (stress/strain) with *in vivo* diastolic dysfunction E/e' (blood flow/wall motion) in CTRL and HFSD. Cardiomyocyte data averaged per heart.

Comparisons between two groups performed using student's t-test (*p<0.05, **p<0.01). Data fitted with linear regression model and calculation of Pearson's correlation coefficient, presented as mean ± standard error of the mean. For **A. & B.** (CTRL: N=5 mice, n=9 cells; HFSD: N=2 mice, n=7 cells). For **C.&D.** (CTRL: N=6 mice, n = 16 cells; HFSD: N=6 mice, n=18 cells).

Fig. 7 Load dependence of cardiomyocyte Ca^{2+} transient decay in CTRL and HFSD cardiomyocytes.

A-B. Exemplar Ca^{2+} transient decay records, with a mono-exponential decay function fit, in non-loaded (isotonic) and loaded (auxotonic, stretched) CTRL and HFSD cardiomyocytes. Horizontal black broken lines intersect with mono-exponential fit at 36.8% (1/e) transient amplitude. Vertical broken lines from intersection points indicate the time constant of Ca^{2+} transient decay (tau).
C. Time constant of Ca^{2+} transient decay (tau) in cardiomyocytes with different load conditions: auxotonic (aux) & stretched (str) shown as fold change relative to non-loaded isotonic (iso) for CTRL and HFSD.

For analysis of two independent variables a 2-way ANOVA with repeated measures used (CTRL: N = 5 mice, n = 13 cells; HFSD: N = 4 mice, n = 12 cells). Data are presented as mean ± standard error of the mean. ***p<0.001 (HFSD stretched vs non-loaded isotonic). Non-significant ANOVA interaction effect (p = 0.055, diet treatment x load state).

Fig. 8 Pacing induced changes in cardiomyocyte cytosolic Ca²⁺ levels and stiffness in CTRL & HFSD.
A. Exemplar CTRL and HFSD cardiomyocyte Ca²⁺ transients at 2 and 4 Hz pacing.
B. Diastolic cardiomyocyte Ca²⁺ levels measured before and after transition from 2 to 4 Hz pacing in HFSD & CTRL.
C. Diastolic cardiomyocyte Ca²⁺ level mean shift with transition from 2 to 4 Hz pacing in HFSD & CTRL.
D. Diastolic sarcomere length measured before and after transition from 2 to 4 Hz pacing in HFSD & CTRL.
E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL at 2 and 4 Hz pacing.
F. Diastolic mean cardiomyocyte segment stress-length relation slope (mN/mm²) normalized relative to segment stretch (% length) yields cardiomyocyte mean segment stress/strain = stiffness
 [(mN/mm²) /strain (% length)] for HFSD and CTRL at 2 and 4 Hz pacing.

Comparison between two groups performed using unpaired t-test. With two independent variables a repeated measures 2-way ANOVA Sidaks post-hoc analysis was performed (*p<0.05, **p<0.01, ****p<0.0001). Data are presented as mean ± standard error of the mean or mean with paired individual points. For **B. & C.** (CTRL: N = 4 mice, n = 9 cells; HFSD: N = 4 mice, n = 8 cells). For **D. & E.** (CTRL: N = 4 mice, n = 7 cells; HFSD: N = 4 mice, n = 9 cells). For **F.** (CTRL: N = 4 mice, n = 8 cells; HFSD: N = 4 mice, n = 9 cells).