Mechanical loading reveals an intrinsic cardiomyocyte stiffness contribution to diastolic dysfunction in murine cardiometabolic disease

Johannes V. Janssens[1,6], Antonia J.A. Raaijmakers[1], Parisa Koutsifeli[3], Kate L. Weeks[1,4], James R. Bell[1,5], Jennifer E. Van Eyk[6], Claire L. Curl[1], Kimberley M. Mellor[2,3,1] and Lea M.D. Delbridge[1]

[1]Department of Anatomy & Physiology, University of Melbourne, Australia

[2]Department of Physiology, University of Auckland, New Zealand

[3]Auckland Bioengineering Institute, University of Auckland, New Zealand

[4]Baker Department of Cardiometabolic Health (Baker), University of Melbourne, Australia

[5]Department of Microbiology, Anatomy, Physiology & Pharmacology, La Trobe University, Australia

[6]Smidt Heart Institute, Cedars-Sinai Medical Center, Los Angeles, USA.

Running Title: *Cardiomyocyte stiffness in diastolic dysfunction*

Key words: *cardiomyocyte; diastolic dysfunction; cardiometabolic disease; stiffness; relaxation; Ca^{2+} myofilament interaction*

Corresponding author:

Prof Lea MD Delbridge

Cardiac Phenomics Laboratory

Department of Anatomy and Physiology,

University of Melbourne, Australia.

lmd@unimelb.edu.au

ABSTRACT

Cardiometabolic syndromes including diabetes and obesity are associated with occurrence of heart failure

with diastolic dysfunction. There are no specific treatments for diastolic dysfunction, and therapies to

manage symptoms have limited efficacy. Understanding of the cardiomyocyte origins of diastolic

dysfunction is an important priority to identify new therapeutics. The investigative goal was to

experimentally define *in vitro* stiffness properties of isolated cardiomyocytes derived from rodent hearts

exhibiting diastolic dysfunction *in vivo* in response to dietary induction of cardiometabolic disease.

Male mice fed a high fat/sugar diet (HFSD vs control) exhibited diastolic dysfunction (echo E/e' doppler

ratio). Intact paced cardiomyocytes were functionally investigated in three conditions: non-loaded, loaded

and stretched.

Mean stiffness of HFSD cardiomyocytes was 70% higher than control. E/e' for the origin hearts was

elevated by 35%. A significant relationship was identified between *in vitro* cardiomyocyte stiffness and *in*

vivo dysfunction severity. With conversion from non-loaded to loaded condition, the decrement in maximal

sarcomere lengthening rate was more accentuated in HFSD cardiomyocytes (vs control). With stretch, the

Ca^{2+} transient decay time course was prolonged. With increased pacing, cardiomyocyte stiffness was

elevated, yet diastolic Ca^{2+} elevation was attenuated. Our findings show unequivocally that cardiomyocyte

mechanical dysfunction cannot be detected by analysis of non-loaded shortening.

Collectively, these findings demonstrate that a component of cardiac diastolic dysfunction in

cardiometabolic disease is derived from cardiomyocyte stiffness. Differential responses to load, stretch and

pacing suggest that a previously undescribed alteration in myofilament-Ca^{2+} interaction contributes to

intrinsic cardiomyocyte stiffness in cardiometabolic disease.

KEY POINTS

• Understanding cardiomyocyte stiffness components is an important priority for identifying new

therapeutics for diastolic dysfunction, a key feature of cardiometabolic disease.

• In this study cardiac function was measured *in vivo* (echocardiography) for mice fed a high-fat/sugar

diet (HFSD, ≥25weeks). Performance of intact isolated cardiomyocytes derived from the same hearts

was measured during pacing under non-loaded, loaded and stretched conditions *in vitro.*

• Calibrated cardiomyocyte stretches demonstrated that stiffness (stress/strain) was elevated in HFSD

cardiomyocytes *in vitro* and correlated with diastolic dysfunction (E/e') *in vivo*. HFSD cardiomyocyte

Ca^{2+} transient decay was prolonged in response to stretch. Stiffness was accentuated with pacing

increase while the elevation in diastolic Ca^{2+} was attenuated.

• Data show unequivocally that cardiomyocyte mechanical dysfunction cannot be detected by analysis of

non-loaded shortening.

• These findings suggest that stretch-dependent augmentation of the myofilament-Ca^{2+} response during

diastole partially underlies elevated cardiomyocyte stiffness and diastolic dysfunction of hearts of

animals with cardiometabolic disease.

INTRODUCTION

Increasingly, cardiometabolic syndromes, including diabetes and obesity, are associated with occurrence of

heart failure with a diastolic dysfunction patho-phenotype (Sotomi *et al.*, 2021). Diastolic dysfunction is

characteristic of heart failure with preserved ejection fraction (HFpEF) which has now overtaken heart

failure with reduced ejection fraction (HFrEF) comprising the majority of new heart failure diagnoses

(McHugh *et al.*, 2019; Jackson *et al.*, 2022). Diastolic dysfunction is also linked with progression of diabetic

cardiomyopathy (Patil *et al.*, 2011; Redfield, 2016). With both HFpEF and diabetic heart disease, early

development of diastolic dysfunction can be detected even when other phenotypic features/comorbidities

are heterogeneous (i.e. hypertrophy, dilation, natriuretic factor elevation, hypertension) (Kittleson *et al.*,

2023). Common clinical features of cardiometabolic syndromes, including obesity and insulin resistance, are

independent risk factors for diastolic dysfunction as a prelude to heart failure (Rozenbaum *et al.*, 2019;

Kamon *et al.*, 2020). The experimental induction of these phenotypes using a high/fat sugar diet feeding

protocol has been shown by us and others to reliably produce diastolic dysfunction in rodents (Pulinilkunnil

et al., 2014; Dia *et al.*, 2020; Sowers *et al.*, 2020; Wingard *et al.*, 2021; Daniels *et al.*, 2022). Cardiometabolic

syndromes can involve the 'silent' presence and progression of diastolic dysfunction for an extended

period. Symptoms can be subclinical and/or detectable only in challenge conditions (i.e. as reduced

diastolic reserve in exercise) or when exacerbated by co-morbidity (Leung *et al.*, 2015; Lam *et al.*, 2019).

Diagnosis of heart failure with diastolic dysfunction is independently associated with higher incidence of

adverse clinical events (Sotomi *et al.*, 2021). Notably, there is currently no specific treatment for diastolic

dysfunction and/or HFpEF, and therapies to manage symptoms have limited efficacy (Delbridge *et al.*,

2022).

In diastolic dysfunction, the relaxation and filling phases of the cardiac cycle are impaired. The condition is

generally described qualitatively as an increase in left ventricular wall 'stiffness' (or by a decrease in

compliance, the inverse of stiffness) (Daniels *et al.*, 2022; Janssens *et al.*, 2024). Clinically and

experimentally, reduced compliance can be measured invasively as an elevation in left ventricular filling

pressure or an increase in the gradient of the diastolic pressure-volume relationship. Non-invasively, on

doppler echocardiogram, an increase in the E/e' ratio [early (E) mitral valve blood flow rate to early mitral

annulus tissue motion (e')] is an index used to identify diastolic abnormality (Lo & Thomas, 2010).

Myocardial stiffness is commonly considered to reflect myocardial extracellular matrix abnormalities linked

with increased tissue fibrosis. While a component of reduced compliance likely reflects fibrotic infiltration,

diastolic dysfunction can be identified in settings where there is no evidence of increased interstitial fibrosis

burden (Curl *et al.*, 2018). There is growing appreciation that alterations of myocardial compliance in

diastolic dysfunction disease states represents underlying intrinsic cardiomyocyte stiffness pathology which

contributes to impaired relaxation and filling (Zhou & Pu, 2016; Villalobos Lizardi *et al.*, 2022).

Understanding the cardiomyocyte origin of diastolic dysfunction is an important priority to identify new

molecular targets and to support the development of new diagnostics and therapeutics to reduce

morbidity and mortality in cardiometabolic disease states.

Cardiomyocyte origins of diastolic dysfunction potentially comprise structural and operational components.

Evidence of molecular changes in sarcomeric and cytoskeletal elements is extensive (Villalobos Lizardi *et al.*,

2022; Loescher *et al.*, 2023; Janssens *et al.*, 2024). Operationally, the cardiac cycle of contraction and

relaxation is driven by the cyclic flux of cardiomyocyte activator Ca^{2+} to initiate contraction (myocyte

sarcomere shortening) and to allow relaxation (myocyte sarcomere lengthening). The relationship between

cytosolic Ca^{2+} levels and myofilament activation is dynamic and depends on several external factors some

of which include stretch and pacing (Fabiato & Fabiato, 1978; Varian & Janssen, 2007).

Conventionally, the steady-state force-Ca^{2+} analysis (typically performed in permeabilised, quiescent

preparations) is used to summarise the myofilament-Ca^{2+} response and is the product of many complex

myofilament protein interactions. Less work has been done to investigate the dynamic changes in the

myofilament-Ca^{2+} relationship in paced intact cardiomyocytes, especially in disease contexts. In diastolic

dysfunction, specifically in HFpEF animal models, evidence is accruing that Ca^{2+} deficit and impaired non-

loaded shortening *are not* phenotypic (Kilfoil *et al.*, 2020). Indeed cardiomyocyte myofilament hyper-

activation and elevated Ca^{2+} levels have been reported (Curl *et al.*, 2018; Louch, 2020; Saad *et al.*, 2023). As

we have recently proposed, the role for elevated diastolic Ca^{2+} levels and augmented myofilament-Ca^{2+}

response comprising a cross-bridge derived component of cardiomyocyte diastolic stiffness is increasingly

plausible (Janssens *et al.*, 2024). This experimental enquiry focusses on intact cardiomyocyte mechanical

and Ca^{2+} performance in response to load, stretch and modulated pacing conditions.

While the term 'stiffness' is used as a general descriptor of impaired compliance in a cardiovascular setting,

it is more formally specified quantitatively in physical science disciplines (Mirsky, 1976). Stiffness is defined

as the extent to which an object resists deformation. Stiffness is influenced by the geometry and the

intrinsic material properties of an object. In the cardiac literature and in this study, stress/strain relations

are calculated to assess cardiomyocyte resistance to stretch. Stress is defined as force per cross-sectional

area, and strain is the application of a normalized stretch or deformation. The ratio of stress/strain defines

the cardiomyocyte resistance to stretch independent of geometry and is known as 'Young's Modulus'.

While stiffness and Young's Modulus are not equivalent, the term 'stiffness' is commonly used (even

though Young's Modulus is the correct formal term). In the cardiac setting, a stretch or deformation state is

commonly described as 'loading' – for example 'pre-load' (diastolic stretch) and 'after-load' (resistance to

shortening). Depending on the experimental design, stiffness behaviour of myocardial tissues, active

cardiomyocytes and/or myocyte sub cellular cytoskeleton and sarcomere structures all potentially

contribute to *in vivo* functional stiffness. Little is known about how stiffness measured at different

structural levels may determine function *in vivo* and how stiffness elements are quantitatively changed in

pathophysiology – in particular in acquired disease conditions. Thus, the goal of this investigation was to

experimentally characterize *in vitro* stiffness properties (stress/strain) of cardiomyocytes derived from

hearts of animals exhibiting diastolic dysfunction *in vivo* in the setting of dietary induction of

cardiometabolic disease.

METHODS & MATERIALS

Ethical approval

The authors understand the ethical principles under which *The Journal of Physiology* operates, and the

work presented in this study complies with the animal ethics checklist as outlined in Instructions to

Authors. All experiments conform to the principles and regulations as described by Grundy (2015). Animals

(male C57Bl/6J mice) were purchased from the Animal Resource Centre (Perth, WA Australia). All animal

experiments were approved by the University of Melbourne (School of Biomedical Sciences) Animal Ethics

Committee (Project Ethics Approval 10366) and complied with the guidelines and regulations of the Code of

Practice for the Care and Use of Animals for Scientific Purposes (National Health and Medical Research

Council Australia). Animals were provided with free access to food and water, were group housed in a

temperature-controlled environment 21-23°C with 12 hour light/dark cycles. Animals were euthanised

under anaesthesia (single dose of sodium pentobarbital 70 mg/kg i.p) on cardiac excision.

Animal model of cardiometabolic disease

As a pre-clinical cardiometabolic disease model, the high fat/sugar diet (HFSD) fed rodent recapitulates a

common and growing human disease phenotype (obesity, insulin resistance, glucose intolerance) and has

been established as a diastolic dysfunction model (Heather *et al.*, 2022). Cardiometabolic disease was

induced in male C57Bl/6J mice by a high fat/sugar dietary intervention commencing at 8-9 weeks of age.

Animals were randomized to experimental treatment groups and were group housed (> 4 animals per cage)

in a temperature-controlled environment 21-23°C with 12 hour light/dark cycles. After 1 week transitional

feeding period (alternating feed periods), animals were fed a HFSD (43% kcal from fat, 200 g/kg sucrose,

SF04-001, Specialty Feeds) or control reference diet (CTRL: 16% kcal from fat, 100 g/kg sucrose, AIN93G,

Specialty Feeds) for at least 25 weeks.

Glucose tolerance testing

At 23-25 weeks of feeding, glucose tolerance testing was performed in mice following 6 hours fasting as

previously described (Chandramouli *et al.*, 2018). Baseline blood glucose levels were measured using an

Accu-Chek glucometer with a blood sample obtained from a needle prick to the tail vein. Glucose (1.5 g/kg

body weight) was injected i.p. and blood glucose was measured at 5, 15, 30, 60, 90 and 120 minutes after

the glucose injection.

Transthoracic echocardiography

At 25-26 weeks of feeding, cardiac structure, systolic and diastolic function were evaluated by transthoracic

2-dimensional B- and M-mode and blood flow and tissue doppler echocardiography (GE Vivid 9; 15mHz i13L

linear array transducer) as described previously (Chandramouli *et al.*, 2018; Curl *et al.*, 2018). Briefly, mice

were lightly anesthetised with Isoflurane (1.5%; Baxter healthcare, Australia). Parasternal short axis view

was used to measure morphological and systolic parameters (interventricular septum IVS, left ventricular

posterior wall LVPW, left ventricular internal dimension LVID, fractional shortening FS, ejection fraction EF,

heart rate HR). Pulse wave Doppler and tissue Doppler scans were acquired from the apical 4 chamber view

to assess LV diastolic function parameters: velocity of early (E) and late (A) mitral inflow and early (e') and

late (a') diastolic velocity of mitral annulus. From these compound indices of diastolic function E/e', E/A and

e'/a' were calculated. In addition, the slope of the E wave was used to calculate mitral valve deceleration

time. Three consecutive cardiac cycles were sampled for each measurement.

Cardiomyocyte preparation, structural and functional assessment

A series of protocols were implemented to phenotype cardiomyocyte and sarcomere morphology, and

assess functional status in different contraction configurations and challenge settings.

Cardiomyocyte isolation, morphology & microfluorimetry. Prior to cardiomyocyte isolation animals were

aneasthetised (sodium pentobarbital 70 mg/kg i.p) and hearts excised. HFSD and CTRL left ventricular

cardiomyocytes were enzymatically isolated using methods previously described (Howlett *et al.*, 2006;

Mellor *et al.*, 2011) at 30-32 weeks feeding. Briefly, the heart was excised, cannulated via the aorta, and

perfused on a Langendorff apparatus with an oxygenated Ca^{2+} free buffer at $37^{o}C$ for 10 minutes. The

buffer was composed of (in mM): 130 NaCl, 5 KCl, 0.33 NaH_2PO_4; 1 $MgCl_2$, 25 HEPES; 20 Glucose; 3 Na

Pyruvate; 1 Na Lactate. The Ca^{2+} free buffer was then supplemented with 0.66mg/mL collagenase

(Worthington; Type II), 0.033 mg/mL trypsin and 0.05 mM $CaCl_2$ and perfusion continued for a further 10

minutes. The heart was removed from the Langendorff apparatus, and the left ventricle was segmented

into small pieces in a high potassium solution composed of (in mM): 30 KCl, 90 KOH, 30 KH_2PO_4, 3 $MgSO_4$,

50 glutamic acid, 20 taurine, 0.5 EGTA, 10 glucose, and 10 HEPES. Single left ventricular cardiomyocytes

were isolated via gentle trituration. Viable cardiomyocytes were resuspended in HEPES-Krebs buffer (in

mM); 146.2 NaCl, 4.69 KCl, 0.35 $NaH_2PO_4H_2O$, 1.05 $MgSO_47H_2O$, 10 HEPES, 11 glucose containing 1.8 mM

Ca^{2+} and were stored on a rocking platform at room temperature. From each heart cell dimensions were

collected from at least 50 rod shaped cardiomyocytes using an inverted light microscope (Nikon) and

calibrated eye piece as previously described (Curl *et al.*, 2018). Cardiomyocyte volume was calculated by

inserting experimentally determined myocyte length and width values into the formula: volume = length x

width x depth x multiplier 0.54 (elliptical area of the rectangle defined by width & length, (Satoh *et al.*,

1996)). Cardiomyocyte width: depth ratio was assumed to be 1.44 (Natali *et al.*, 2001). Cardiomyocyte

cross-sectional area (CSA) could be calculated from measured width scaled for 2-dimensional geometry

(CSA = W [W/1.44] x 0.54 μm^2). An aliquot of isolated cardiomyocyte suspension was pipetted into a cell

chamber mounted on the stage of an inverted fluorescence microscope (Motic AE31) and superfused with

HEPES-Krebs solution with 2 mM Ca^{2+}, 37°C at a flow rate of 2 mL/min. Cardiomyocyte sarcomere length

and Ca^{2+} transients were simultaneously recorded during isotonic contractions. Myocyte force development

was evaluated in the same cells with transducer attachment, using four protocols described below.

Characterizing cardiomyocyte diastolic function: non-loaded shortening measurements (isotonic), loaded

force measurements (auxotonic), stretch responses and pacing challenge. Cardiomyocyte performance

was evaluated in several paced conditions: designated 'non-loaded' (isotonic), 'loaded' (auxotonic after-

loaded), and 'stretched'(auxotonic pre-loaded). The non-loaded state refers to the absence of externally

applied load (noting that sarcomeric and cytoskeletal structures do constitute an underlying internal load).

Firstly, cardiomyocyte non-loaded (isotonic) sarcomere contractions (IonOptix SarcLen, Fast Fourier

Transform, FFT) and Ca^{2+} transients (Fura2-AM, 5 μM microfluorimetry F340:380 nm ratio) were

simultaneously measured at 2 Hz, using a microscope diaphragm to define a rectangular intra-myocyte

region of interest as depicted. During contraction cycles, cardiomyocyte sarcomere length was monitored

within an intracellular region of interest defined by an adjustable diaphragm mounted in the microscope

optical tube (**Fig 4A**).

Secondly, to obtain loaded (auxotonic) contraction data, each cardiomyocyte was attached to a pair of glass

fibres coated with a bio-adhesive (MyoTak, IonOptix, USA). Attachment sites were positioned to be outside

the sarcomere tracking region, closer to myocyte longitudinal ends. One glass fibre was connected to a

force transducer (fibre stiffness 20 N/m, IonOptix, USA), and the other to a piezo-motor (Nano-drive, Mad

City Labs Inc. WI, USA) to adjust cardiomyocyte length (**Fig. 4A**). With successful attachments,

cardiomyocytes were lifted from the coverslip using motorised micromanipulators (MC1000E, Siskiyou

Corporation, USA) attached to both the force transducer and the piezo-motor. Baseline loaded contractions

at 2 Hz were recorded. Sarcomere length changes were measured using FFT as described above. In

addition, with glass fibre attachment, tracking of mean sarcomere shortening within a myocyte segment

delineated by the glass fibre edge positions could be performed in parallel (**Fig4B-C**).

Thirdly, force was recorded during a standardized stretch protocol of loaded cardiomyocytes to measure

the effect of increasing diastolic cardiomyocyte/sarcomere basal length during pacing. A stress-length

relation was constructed for each myocyte (where stress = force/cross-sectional area). Prior to the

commencement of the myocyte stretch measurements a protocol to calibrate the stretch step was

implemented. Four consecutive 0.5 V input signals were delivered to the piezo-motor to determine a

calibration constant to define the percentage cell stretch for a given voltage (**Fig. 4B**). For each myocyte a

Calibration Constant was calculated as: *myocyte segment length change per piezo-motor mV / sarcomere*

count in segment [where *myocyte sarcomere count = myocyte segment length (μm) / average diastolic*

sarcomere length (μm)]. To track the glass fibre position a real-time contrast detection algorithm (IonOptix -

EdgeLength) was focused on the internal edge of each of the glass fibres (**Fig. 4**) (Peyronnet *et al.*, 2017).

This enabled reproducible and comparable stretch steps to be applied to different cardiomyocyte

preparations (**Fig. 4B-C**). Prior to the 'stress-length' protocol, a cell-specific calibration constant was applied

to yield a standardised stretch for each cardiomyocyte evaluated. The stress-length protocol involved

progressive stretch steps totaling an approximate 30% cell stretch (**Fig. 4B-C**).

Lastly, the pacing was increased to 4 Hz, a rate adaptation period allowed (2 min) and the stress-length

protocol was repeated. Sarcomere length, force development and intracellular Ca^{2+} transients were

simultaneously measured. An average of at least 10 twitches or transients were used to calculate

sarcomere, Ca^{2+} and force parameters at a given stretch length. In summary, the following parameters were

measured: diastolic stress (mN/mm^2), diastolic stress-length relation [(mN/mm^2) / (%$^-$ length change)],

diastolic stress-sarcomere length relations ((mN/mm^2) / μm), systolic : diastolic stress-sarcomere relation

(Frank Starling Gain Index (Bollensdorff *et al.*, 2011)), systolic : diastolic stress-length relation, systolic stress

amplitude (mN/mm^2), sarcomere shortening (% diastolic SL), maximum sarcomere shortening and

lengthening velocity (μm/s). In addition, Ca^{2+} parameters were measured: diastolic Ca^{2+}, systolic Ca^{2+} and

time constant of decay of the Ca^{2+} transient (tau).

Statistical analyses

Data are presented as mean ± standard deviation and statistical analysis was performed using GraphPad

Prism V9.5.1. Cardiomyocyte stress-length and stress-SL relations were modelled with simple linear

regression. For comparison between two groups an unpaired student t-test was used. For assessment

between two groups at multiple points a one-way ANOVA with Bonferroni multiple comparisons post-hoc

test was used. Two-way ANOVA with Šídák's multiple comparisons post-hoc test was used for the

evaluation of data groups with two independent variables. Pearson Correlation analyses were used to

assess statistical significance of fitted linear regression lines. A p-value of <0.05 was considered statistically

significant. Analyses were performed using a blinded protocol.

RESULTS

Systemic indicators of cardiometabolic disease and *in vivo* diastolic function in HFSD mice

High fat/sugar diet-fed mice exhibited obesity and glucose intolerance. Relative to CTRL, the body weight of

HFSD mice was significantly elevated within 5 weeks of dietary intervention increasing over the study

duration by 50% relative to CTRL (**Fig. 1A**, CTRL: 28.1±2.0 vs. HFSD: 44.3±1.9 g). After 23-24 weeks diet

treatment, glucose tolerance testing of a subset of the HFSD cohort was performed. Glucose intolerance in

response to intraperitoneal glucose administration was evidenced by increased area under the curve,

consistent with literature reporting insulin resistance in mice fed a similar diet (**Fig. 1B**).

After confirming these key cardiometabolic disease characteristics in the HFSD mice, echocardiographic

functional evaluation was undertaken. Focusing on diastolic function evaluation (**Fig. 1C-G**), there was

higher early mitral inflow (**Fig. 1D,** E wave, CTRL: 472±74, HFSD: 569±65 mm/s) and reduced velocity of wall

motion (**Fig. 1E,** e' wave, CTRL: 28.5±5.1, HFSD: 25.2±3.5 mm/s) in HFSD myocardium. Deceleration time of

mitral inflow was shortened (**Fig. 1F,** CTRL: 30.8±8.7, HFSD: 19.7±5.9 ms) and E/e' ratio increased (**Fig. 1G,**

CTRL: 16.7±3.9, HFSD: 22.3±6.0), all observations consistent with reduced tissue compliance and diastolic

dysfunction. No significant differences in systolic performance parameters were observed (**Table. 1**),

confirming the primary *in vivo* phenotype of diastolic dysfunction. The next step was to evaluate the

cardiomyocyte involvement in dysfunction.

Non-loaded sarcomere and Ca^{2+} dynamics in CTRL and HFSD cardiomyocytes

Single isolated cardiomyocytes were prepared from hearts of a subset of mice which underwent

echocardiographic analyses. For intact, non-loaded cardiomyocytes sarcomere shortening and lengthening

cycles were similar for CTRL and HFSD groups (**Fig. 2A**). Diastolic sarcomere length (SL) (**Fig. 2B**, CTRL:

1.71±0.04, HFSD: 1.72±0.04 μm) and extent of sarcomere shortening (**Fig. 2C**, CTRL: 3.79±2.24, HFSD:

3.94±3.15 % diastolic SL) were not significantly different between groups. Examination of sarcomere

contraction kinetics (**Fig. 2D** exemplar trace) showed that the maximum rate of lengthening (**Fig. 2E**, CTRL:

1.08±0.85, HFSD: 1.29±1.07 μm/s) and shortening (**Fig. 2F**, CTRL: -2.32±1.44, HFSD: -2.57±2.01 μm/s) did

not significantly differ between CTRL and HFSD groups. Collectively these findings indicate that in the non-

loaded setting no differences in CTRL and HFSD cardiomyocyte systolic and diastolic non-loaded

performance were detected, a finding consistent with other studies investigating non-loaded

cardiomyocyte performance in diet-induced cardiometabolic disease including our previous studies in

fructose-fed mice (Mellor *et al.*, 2012; Llano-Diez *et al.*, 2016; West *et al.*, 2019; Wang *et al.*, 2020).

Next, Ca^{2+} parameters were compared with cardiomyocyte sarcomere lengthening performance during

relaxation and diastole (**Fig. 2G**, exemplar traces). No significant difference was evident between dietary

treatment groups in the systolic Ca^{2+} (**Fig. 2H**, CTRL: 1.098±0.169, HFSD: 1.171±0.182 $F_{340:380}$), diastolic Ca^{2+}

(**Fig. 2I**, CTRL: 0.749±0.097, HFSD: 0.761±0.094 $F_{340:380}$) or time constant (tau) of Ca^{2+} transient decay (**Fig.2J**,

CTRL: 0.160±0.067, HFSD: 0.137±0.054 sec). Thus, the significant diastolic dysfunction measured *in vivo* for

the HFSD group was not observed in a non-loaded *in vitro* setting. To reconcile the impaired

echocardiographic heart performance with absence of dysfunction in non-loaded contracting single

cardiomyocytes, the next strategy was to assess single cardiomyocyte response to loading. A common

feature of cardiometabolic disease is elevation in after-load – i.e. elevated systemic resistance potentially

impeding myocardial pump function. To simulate this *in vitro*, mechanically loaded cardiomyocyte

performance was examined.

Loaded cardiomyocyte sarcomere shortening and lengthening

Immediately after non-loaded cardiomyocyte recordings, glass fibres were attached to cardiomyocyte

longitudinally separated positions to permit auxotonic, loaded contraction measurements. For cells

successfully attached, the mean difference in sarcomere length measured pre- and post-conversion to

loaded mode was <0.02% diastolic length, indicating non-loaded and loaded sarcomere initial lengths were

well matched. For attached cardiomyocytes, a comparison of sarcomere shortening behavior in the loaded

state could be made (using FFT) to directly match with data obtained non-loaded state.

In the first instance, data from all myocytes were pooled to investigate the general effects of load

regardless of treatment group. With loading, for all myocytes (both treatment groups), extent of sarcomere

shortening was reduced and shortening and lengthening kinetics slowed when compared with non-loaded

performance (**Fig. 3A**). Mean data showed that in response to loading, sarcomere shortening was reduced

by ~34% (**Fig. 3B**, non-loaded: 0.075±0.042, loaded: 0.050±0.031 µm). Maximum rate of shortening was

decreased by 58% (non-loaded: -2.85±1.40, loaded: -1.20±0.89 µm/s; not shown), and maximum rate of

lengthening by 66% (**Fig. 3C**, non-loaded: 1.46±1.03, loaded: 0.52±0.50 µm/s). The magnitude of the change

in extent of shortening and maximum rate of lengthening with loading were positively correlated (**Fig. 3D**)

suggesting the possibility of common underlying mechanisms.

To assess cardiomyocyte rate kinetics in non-loaded and loaded cardiomyocytes independent of shortening

extent, the maximum rates of lengthening and shortening were normalised by shortening amplitude to

derive 'deformation rate' - i.e. rate of length change, relative to extent of shortening (($µm$ x sec^{-1})/$µm$ =

1/sec). The effect of loading on deformation rate was accentuated in HFSD group (significant ANOVA diet x

load interaction effect, **Fig. 3E**). A selective HFSD effect was revealed in the differential between matched

'loaded' and 'non-loaded' lengthening cardiomyocytes. The decrement in deformation rate (lengthening)

for CTRL and HFSD is shown in **Fig. 3F**, and is accentuated in HFSD in response to loading (CTRL: 6.84±4.08,

HFSD: 12.35±7.15 1/s). Treatment differences in deformation rate were not evident during the shortening

phase (data not shown)**.** These findings indicated an accentuated influence of load, specific to the

lengthening (i.e. diastolic) phase in HFSD cardiomyocytes and could reflect impaired subcellular relaxation

processes specific to the HFSD myocardium. Based on these data identifying different CTRL and HFSD origin

cardiomyocyte responses in loaded state, we next investigated the effect of stretch applied to

cardiomyocytes to simulate *in vivo* 'pre-load' conditions.

Cardiomyocyte force development: measuring strain & stress during progressive cardiomyocyte stretch

A hallmark of diastolic dysfunction is an elevated diastolic pressure-volume relationship which describes

the increased resistance of the myocardial wall to chamber filling (Aurigemma *et al.*, 2006). We sought to

design a single cardiomyocyte version of this relationship (Fowler *et al.*, 2015), by applying standardized

stretch steps to cardiomyocytes, defining the steps as a % of non-stretched state (i.e. 'strain'). Diastolic

force was measured simultaneously during length stretch steps. The force measured at every strain

increment was normalized by cross sectional area (CSA) to produce a measure of 'stress' (i.e. force/CSA,

mN/mm^2). Thus a sarcomere 'stress/strain' relation (i.e. force-length) was constructed for each myocyte.

The protocols employed to generate stress-strain are depicted in **Fig. 4** and described in detail in the

Methods. **Fig. 4** shows the stretch standardization approach used to enable reproducible application of

strain (converting voltage signals to linear displacement) as a % length change (via individual myocyte

calibration constant calculation) to allow comparisons between different cardiomyocytes. The size of each

step change in length on sarcomere length was measured (FFT) and confirmed by tracking the intra-

cardiomyocyte segment between the glass fibres (**Fig. 4**). In some cases a direct sarcomere FFT measure

was not microscopically resolvable due to myocyte contraction responses which partially positioned some

area of the myocyte outside the optimal focal plane. In these situations it was always possible to calculate

segment length change. For the example shown, stretch steps were determined to be 2.75μm, and

converted to % sarcomere length (i.e. strain) using sarcomere count within intracellular region of interest

area (**Fig. 4C**). A comparison of the variability of cell stretches performed using a uniform (pre-set) stretch

constant and the individually derived stretch calibration constant demonstrates that reproducibility is

optimized with usage of the individual stretch calibration constant data (**Fig. 4D**). Force, sarcomere length

and % stretch were measured simultaneously (**Fig. 4**).

Quantifying diastolic stiffness in CTRL vs HFSD cardiomyocytes using strain normalization methods

Superimposed exemplar stress recordings are shown in **Fig. 5A** for CTRL and HFSD cardiomyocytes,

captured during a series of progressive stretches during a 100 sec period. The dashed (broken) lines are

aligned with the diastolic stress step increments for each of the 2 myocytes (using the data points collected

prior to the transition to a new step) throughout the series of progressive stretches. A robust increase in

systolic force with stretch is also evident, indicative of length-dependent activation, the cardiomyocyte

basis for the Frank-Starling mechanism.

In this experimental setting, stiffness could be visualized as the gradient of the linear plot derived from the

cardiomyocyte diastolic stress-length relation. Our stretch protocol methodology provided two approaches

for stiffness calculation (as per Methods). A first approach used FFT measurement of sarcomere length to

track cardiomyocyte stretch and was plotted according to the standardized stretch steps (in μm). An

alternative approach used percent length change calculated as the length of the cardiomyocyte segment

between the glass attachment fibres divided by the initial distance between the fibres prior to stretch (in %

initial length).

Applying the first approach, a linear regression was performed to fit the data from which a slope value

could be derived. **Fig. 5B** depicts the average diastolic and systolic stress-sarcomere length relations for

CTRL cardiomyocytes only. To determine cardiomyocyte diastolic stiffness using FFT-computed sarcomere

length values, the gradient of the mean slope of the diastolic stress sarcomere-length relation for each

treatment group was calculated, using mean sarcomere length change (μm) as the denominator. This

gradient was significantly higher in HFSD myocytes compared to CTRL (**Fig. 5C**). The individual slope

gradients were used to calculate the mean diastolic stress / strain gradient (**Fig. 5D,** CTRL: 2.58±2.25, HFSD:

5.50±4.01 (mN/mm^2) / μm, p = 0.0996) where there was a trend for a 2.3 fold increase in stiffness of HFSD

cardiomyocytes. The statistical power of this analysis was limited due to the difficulty in achieving

resolvable diastolic sarcomere lengths by FFT for all myocytes.

Applying the second approach, it was possible to construct stress-segment length (%) relations for all

myocytes from which stress data could be obtained. In **Fig. 5E**, the stress-segment length data for HFSD and

CTRL are shown, with a robustly significant difference detected in mean gradient values. In **Fig. 5F** the

diastolic stress/strain values calculated for the myocytes from the two treatment groups using this segment

approach are directly compared. The HFSD cardiomyocyte stiffness index derived using this approach was

increased by 70% compared with CTRL (**Fig. 5F**, CTRL: 0.0075±0.0064, HFSD: 0.0128±0.0068 (mN/mm^2) / %

length). There was no significant difference in 'Frank Starling Gain index' (**Table. 2,** CTRL: 2.40±1.06, HFSD:

2.77±1.45, (mN/mm^2) / μm) or the ratio of systolic to diastolic stress segment length relations (**Table. 2,**

CTRL: 1.62±0.25, HFSD: 1.73±0.50, (mN/mm^2) / % length) indicating that cardiomyocyte length-dependent

activation is preserved (Bollensdorff *et al.*, 2011; Najafi *et al.*, 2016)**.**

To examine the relationship between intrinsic *in vitro* cardiomyocyte stiffness and *in vivo* diastolic

dysfunction, diastolic stiffness (slope of the diastolic stress segment-length relation) averaged per animal

was plotted against the E/e' index of diastolic dysfunction. There was a significant correlation between

cardiomyocyte stiffness and E/e' in HFSD and no evidence of a relationship for the CTRL group (**Fig. 5G**

CTRL: p = ns, HFSD: p = 0.0096, r^2 = 0.9213). These data support the interpretation that intrinsic

cardiomyocyte stiffness represents a definable and inherent component of diastolic dysfunction occurring

in vivo in cardiometabolic disease.

Load dependence of Ca^{2+} transient decay in CTRL and HFSD cardiomyocytes

To further understand the cardiomyocyte functional elements comprising increased intrinsic stiffness in

cardiopathology, the relationship between Ca^{2+} transient features and myofilament deactivation in

response to progressive stretch was evaluated. In the intact cardiomyocyte model these electro-mechanical

coupling processes are functional for interrogation (in contrast to the disrupted cytosolic environment

which characterizes permeabilized myocyte preparations). The Ca^{2+} transient decay features were

compared for non-loaded and loaded basal states and also in response to progressive stretch.

In CTRL cardiomyocytes there was no difference in the time constant of Ca^{2+} transient decay (tau, 1/e)

between loaded and stretched state (20% increase in length, **Fig. 6A.** loaded 1.18±0.48; stretched

1.12±0.35 fold change vs non-loaded reference 1.00±0.46). In HFSD cardiomyocytes the time constant of

Ca^{2+} transient decay was increased in stretched cardiomyocytes relative to non-loaded (**Fig. 6A**. non-

loaded, 1.00±0.42; loaded 1.22±0.50; stretched 1.42±0.65 fold change). Loading prolonged time to peak

Ca^{2+} transient in both CTRL and HFSD cardiomyocytes (**Fig. 6B.** CTRL: non-loaded 1.00±0.25; loaded

1.49±0.38; stretched 1.46±0.44, HFSD: non-loaded 1.00±0.41; loaded 1.36±0.41; stretched 1.21±0.34

foldchange). The effects can be visualized in **Fig. 6C-D** where the mono-exponential decay of exemplar Ca^{2+}

transients is depicted. The shallower decay of the Ca^{2+} transient in the stretched state (**Fig 6D**, HFSD) was

apparent whereas the loaded and non-loaded decay traces were virtually overlayed for CTRL

cardiomyocytes (**Fig. 6C**). These data suggest that stretch may enhance a mechanism by which the

myofilaments bind Ca^{2+} for a prolonged duration during relaxation thus reducing Ca^{2+} availability for

extrusion from the cytosol.

Pacing-induced [Ca^{2+}]$_i$ increase to examine potential for a cross-bridge derived stiffness mechanism

A further mechanistic study to probe the potential relationship between Ca^{2+} and stiffness involved

perturbation of cytosolic Ca^{2+} cycling time. Elevating stimulation frequency is a recognized physiologic

strategy to increase diastolic Ca^{2+} levels and to apply increased cardiomyocyte workload. In response to

frequency elevation (2 Hz to 4 Hz) cardiomyocyte diastolic Ca^{2+} levels increased in both CTRL and HFSD (**Fig.**

7A & B) groups consistent with previous studies (Varian & Janssen, 2007; Eisner *et al.*, 2020). In the HFSD

cardiomyocytes this effect was markedly diminished and about half the mean magnitude of the Ca^{2+} shift

observed for CTRL myocytes (**Fig. 7B.** CTRL 2 Hz: 0.664±0.052, CTRL 4 Hz: 0.775±0.083, HFSD 2 Hz:

0.688±0.099, HFSD 4 Hz: 0.741±0.126, F$_{340:380}$, 2-way ANOVA significant interaction effect, **Fig. 7C**. CTRL

0.110±0.051; HFSD 0.054±0.049, F$_{340:380}$). No treatment or interaction effects were observed for systolic

Ca^{2+}, Ca^{2+} amplitude, or time constant of Ca^{2+} transient decay (**Table. 3**). Frequency elevation was

associated with a reduction in diastolic sarcomere length in both CTRL (3%) and HFSD (5%) loaded

cardiomyocytes (**Fig. 7D**. CTRL 2 Hz: 1.69±0.04, CTRL 4 Hz:1.64±0.05, HFSD 2 Hz: 1.69±0.04, HFSD 4 Hz:

1.61±0.07 µm). The combination of normal diastolic sarcomere length reduction, concomitant with

suppressed diastolic Ca^{2+} increase in HFSD cardiomyocytes, may indicate elevated diastolic responsiveness

to Ca^{2+} in the HFSD cardiomyocyte.

While frequency elevation did not change the stress/strain relation in CTRL cardiomyocytes, a significant

increase in the stress/strain gradient was observed with increased frequency in HFSD cardiomyocytes (**Fig.**

7E&F. CTRL 2 Hz: 0.0060±0.0056, CTRL 4 Hz:0.0068±0.0068; HFSD 2 Hz: 0.0121±0.052, HFSD 4 Hz:

0.0136±0.0052 (mN/mm^2) / % length). This may indicate a larger population of cross-bridges contributing to

diastolic stiffness at heightened workloads in HFSD cardiomyocytes although other potential non-cross-

bridge sources cannot be excluded. Since Ca^{2+} shift in response to increased pacing is modest (relative to

CTRL), increase in stiffness cannot be attributed to rise in Ca^{2+} levels *per se.* Shift in the myofilament-Ca^{2+}

response during diastole is a potential alternate explanation.

DISCUSSION

Here we provide the first quantitative analysis of intact cardiomyocyte stiffness and performance deficits

associated with acquired cardiometabolic disease. We demonstrate that a component of cardiac diastolic

dysfunction in cardiometabolic disease derives from intrinsic cardiomyocyte mechanical abnormality.

The concept of myocardial stiffness is a frequently invoked qualitative descriptor of the pathophysiologic

phenotype characterizing diastolic dysfunction. Clinically and experimentally the echocardiography

parameter E/e' is used as an index of myocardial stiffness in disease diagnosis. E/e' is a compound measure

of cellular and extracellular matrix stiffness contributors. Despite consistent *in vivo* findings, an

understanding of the cardiomyocyte-specific contribution to tissue stiffness has been difficult to identify.

Using a series of intact single cell recording maneuvers, we have established that the stiffness

characteristics of individual isolated cardiomyocytes from normal and dysfunctional hearts can be defined,

normalized and compared. Further, a significant relationship was identified between the mean stiffness of

single cardiomyocytes and the extent of the origin heart diastolic dysfunction (i.e. E/e' vs Young's Modulus

[(mN/mm2) / %length]).

To recapitulate, cardiometabolic disease induced by dietary intervention in mice was characterized by

marked body weight elevation (50%, 'obesity') and glucose intolerance. *In vivo* echocardiography

diagnosed significant diastolic dysfunction with a 35% elevation of the E/e' in HFSD animals in absence of

systolic functional deficit. In paced non-loaded conditions, the sarcomere contraction cycles & Ca^{2+}

transients (fura-2) were similar in CTRL and HFSD cardiomyocytes. When the same cardiomyocytes were

loaded, the shortening extent and maximal lengthening rate were reduced for both diet conditions. In HFSD

cardiomyocytes the reduction in deformation rate during lengthening (but not shortening) was

accentuated, indicating a selective negative impact of loading on relaxation of the HFSD-derived

cardiomyocytes. In HFSD cardiomyocytes there was a larger increase in diastolic force per cross-sectional

area (stress: mN/um^2) in response to standardized step increases in length (strain: % initial length [μm]).

From these data a 'stress/strain' parameter was determined, defining diastolic stiffness for each myocyte

i.e. 'Youngs Modulus' ([$mN/μm^2$]/[% initial length[μm]). Mean stiffness was 70% higher in cardiomyocytes

of HFSD animals. Thus the increase in cardiomyocyte stiffness index was approximately double the

elevation of the *in vivo* E/e' dysfunction index. Using matched *in vivo* and *in vitro* data, a significant

correlation was found between diastolic dysfunction (E/e') and mean cardiomyocyte stiffness value for

HFSD-diet animals, while there was no relationship between these parameters in the myocytes of CTRL-diet

fed group. To evaluate possible influence of mechanical state on the Ca^{2+} transient, the mean transient

decline parameters (tau) from non-loaded, loaded and stretched cardiomyocytes were compared. In HFSD

cardiomyocytes the time constant of Ca^{2+} transient decay was increased in stretched cardiomyocytes

relative to non-loaded, a feature not observed in CTRL cardiomyocytes. Finally, in response to an *in vitro*

pacing increase HFSD cardiomyocytes exhibited accentuated diastolic stiffness even with a smaller increase

in diastolic Ca^{2+} level. Collectively, these findings suggest a previously undescribed alteration in Ca^{2+}-

myofilament interactions indicative of a cross-bridge component to cardiomyocyte stiffness in

cardiometabolic disease.

Dietary model of acquired metabolic syndrome and cardiac diastolic disease

High fat/sugar diet feeding (i.e. 'western diet') has previously been shown experimentally to induce

diastolic dysfunction in rodents. Elevated E/e' is a remarkably consistent echocardiographic feature of high

fat/sugar diet fed mouse myocardial performance while E/A and IVRT may be elevated, reduced, or

unchanged relative to control counterparts (Pulinilkunnil *et al.*, 2014; Dia *et al.*, 2020; Sowers *et al.*, 2020;

Wingard *et al.*, 2021; Daniels *et al.*, 2022). The dietary regime employed in this study has been well

characterized as a model of diastolic dysfunction, with gradual development of metabolic disturbance

(stable rate of weight gain and mild hyperglycemia), with maintained systolic function (ejection fraction and

fractional shortening). The data presented here are consistent with clinical observations linking

cardiometabolic disease and diastolic dysfunction.

As we have reported previously, over the time span of the dietary intervention differential fibrosis is not

observed (Daniels *et al.*, 2022; Wells *et al.*, 2023). The finding of increased stiffness in cardiometabolic

disease is consistent with some aspects of prior experiments involving animal feeding of obesogenic diets,

using other rodent preparation types – including permeabilized cardiomyocytes, tissue strips/papillary

muscles, and non-paced cardiomyocytes (Leopoldo *et al.*, 2010; Gonçalves *et al.*, 2016; Sowers *et al.*, 2020).

Interestingly stiffness in those studies was attributed to titin or extracellular matrix properties. Our protocol

with intact single cardiomyocytes allows a mechanistic extension of these previous findings.

Loading the intact cardiomyocyte - functional deficits revealed in HFSD

Non-loaded contracting cardiomyocytes of CTRL and HFSD mice exhibited identical shortening-lengthening

cycles, but a distinctly different decrement in lengthening rate when exposed to load (i.e. auxotonic with

progressive 'pre-load' increase). Our findings show unequivocally that cardiomyocyte mechanical

dysfunction cannot be detected by analysis of non-loaded shortening. With loading, the shortening extent

and the maximal lengthening rates were reduced for both CTRL and HFSD. In HFSD the lengthening rate

reduction was accentuated even when shortening extent was accounted for. These findings indicated

specific influence of loading on lengthening in HFSD cardiomyocytes and could reflect impaired subcellular

relaxation processes specific to the HFSD myocardium.

Others have demonstrated a physiological effect of loading on shortening in cardiac cells/tissues in

different species (using slow pacing and sub physiologic temperature conditions), although the pronounced

effect on relaxation is not consistently reported (Lab *et al.*, 1984; White *et al.*, 1995; Shimkunas *et al.*,

2021). This is consistent with *in vivo* human clinical observations where there is a strong inverse association

between late-systolic loading and left ventricular wall displacement velocity during early diastole (Borlaug

et al., 2007). Importantly, increased dependence of relaxation on after-load is reported to be an *in vivo*

feature of heart failure with diastolic dysfunction (Eichhorn *et al.*, 1992; Gillebert *et al.*, 2000). Regarding

the Ca^{2+} transient, in the current study a subtle prolongation effect is observed in both CTRL and HFSD

cardiomyocytes when loaded (but not stretched) which is consistent with recently reported findings in rat

LV trabeculae (Dowrick *et al.*, 2022). The variability in the sarcomere shortening recorded for both non-

loaded and loaded cardiomyocytes is notable but consistent with other studies assessing relaxation of

rodent cardiomyocytes (Delbridge & Roos, 1997; Clark & Campbell, 2019). A range of non-loaded

cardiomyocyte mean diastolic sarcomere lengths are reported in the literature. When cells are evaluated *in*

vitro and not constrained by local structural elements sarcomere lengths <1.8μm are commonly observed

and consistent with our findings (Lim *et al.*, 2000; Bub *et al.*, 2010). *In vitro* experimental conditions can

also contribute to variation in diastolic sarcomere lengths (e.g. pacing frequency, temperature and buffer

tonicity) (Roos *et al.*, 1982; Chung & Campbell, 2013; Khokhlova *et al.*, 2022).

Characterizing cardiomyocyte stretch & pacing challenge responses

In order to quantify intact single cardiomyocyte stiffness our approach was to apply a calibrated stretch

protocol, based on pre-measurement of a stretch calibration constant for each myocyte, tracking changes

in force generation and cytosolic Ca^{2+} responses for each stretch step. This calibrated step-by-step

approach enabled the stress and strain data to be gathered from actively paced cardiomyocytes over

extended periods (i.e. 80-100 sec) and computed to generate a definitive, comparative stiffness parameter

for each cardiomyocyte.

In the intact myocyte model, the internal cytosolic regulatory milieu is preserved, and the

cytoskeletal/sarcomeric/organelle structural framework is integral. Thus contributions to diastolic

'stiffness' reflect components of the structural passive resistance to deformation and the basal crossbridge-

derived activity. Recent studies of permeabilized cardiac muscle strip bundles have explored the separate

contributions of mictrotubule, titin, actin-associated, sarcolemmal and desmin-linked components to

passive force development (Loescher *et al.*, 2023). Prior studies of 'skinned' or quiescent myocardial

preparations have also identified sources of passive stiffness elevation, with titin isoform switch and

modification identified as major modulators of stiffness in cardiometabolic disease (Leopoldo *et al.*, 2010;

Hamdani *et al.*, 2013; Gonçalves *et al.*, 2016; Hopf *et al.*, 2018; Sowers *et al.*, 2020; Abdellatif *et al.*, 2021).

Our findings indicate involvement of a cytosolic Ca^{2+} component of the stretch response. Ca^{2+} fluorescence

analyses from these stretch protocols showed novel stretch-dependent prolongation of Ca^{2+} transient

decay in the HFSD cardiomyocytes. The time constant of Ca^{2+} transient decay increased with loading and

further with stretching – a feature not evident in CTRL cardiomyocytes and which provides additional

mechanistic insight .

The final pacing frequency protocol was implemented as a relative workload stress to the loaded

cardiomyocyte. With a doubling of pacing frequency, a significant increase in the stress/strain gradient was

observed in HFSD cardiomyocytes (and not CTRL), and a similar reduction in diastolic sarcomere length in

both CTRL and HFSD was observed. While cardiomyocyte diastolic Ca^{2+} level increased in both CTRL and

HFSD – the HFSD increase was less marked than CTRL. This was a surprising finding, not previously

reported in any other model setting. A suggestion of Ca^{2+} hyper-sensitivity arises (Davis *et al.*, 2007)

amongst other explanations to consider.

Mechanistic insights into cross-bridge derived stiffness

Here we report that a cardiometabolic disease phenotype induced by obesogenic diet includes a 70%

increase in cardiomyocyte stiffness which contributes to a 35% increase in diastolic dysfunction *in vivo* (by

echo E/e'). Several lines of evidence emerging from this study suggest that Ca^{2+} -related mechanism/s may

contribute to cardiomyocyte stiffness increase. Specifically, in the loaded HFSD cardiomyocyte (vs CTRL):

a) load confers accentuated lengthening rate deficit

b) with stretch, the Ca^{2+} transient decay time course is prolonged

c) with more rapid pacing, relatively small increases in diastolic Ca^{2+} are associated with elevated stiffness

Together these cardiomyocyte phenotypes would be consistent with selective impairment of relaxation

rate, selective myofilament hyper-activation, impeded diastolic filling and reduced diastolic reserve.

This combination of observations is consistent with the notion that stretch-dependent cross-bridge

attachment is prolonged in HFSD cardiomyocytes (or conversely detachment is delayed). We can speculate

that an alteration in the 3-dimensional geometry of the low affinity Ca^{2+} binding pocket in troponin C (TnC)

would produce this effect, by stabilizing Ca^{2+} in the pocket for longer every activation cycle. This could

increase TnC buffering power which would explain the relatively small Ca^{2+} rise in HFSD in response to

increased pacing (Eisner *et al.*, 2023). A manifestation of this altered binding function could be apparent

Ca^{2+} hypersensitivity. The TnC-Ca^{2+} dissociation rate has been theorized as an effector of the transient

decay time-course (Robinson *et al.*, 2018). A role for the troponin complex in regulation of Ca^{2+} affinity is

well established (Chemla *et al.*, 2000). Given the load and stretch-dependent nature of these findings, a

role for stress sensor proteins such as titin is likely. Indeed, there is some evidence to suggest that titin

isoform regulates TnC low-affinity binding pocket opening and Ca^{2+} affinity in response to stretch via

regulating the number of attached cross-bridges (Li *et al.*, 2019). Further evidence of structural

modifications of specific myofilament regions which form and regulate this pocket is required. Mass

spectrometry studies by us and others provide proof of concept of relevant post-translational modification

types (Janssens *et al.*, 2018; Loescher *et al.*, 2022).

Limitations

Given the diastolic dysfunction focus of this study, investigations have primarily examined cardiomyocyte

relaxation and lengthening behaviors. Diastolic and systolic interactions are important and additional

protocols are in development for these studies. In this regard, we defer to the extensive body of

technologic research undertaken to develop software and instrumentation to perform single

cardiomyocyte 'Pressure-Volume (PV) loops' (Helmes *et al.*, 2016). Using this approach, to mimic PV cycles

in the intact myocardium, stretch is applied after relaxation has occurred by using a force clamp to

maintain a predefined diastolic force level. This approach has been used to discover that titin truncation

shifts the working sarcomere length range preserving diastolic stiffness at the expense of systolic

performance (Methawasin *et al.*, 2022) and that highly compliant titin allows cardiomyocytes to function at

high diastolic Ca^{2+} levels (Najafi *et al.*, 2019). Potentially, this approach could also be used to evaluate the

temporal relationship between slow or incomplete relaxation and cardiomyocyte cross-bridge derived

stiffness in a semi-situ manner. Operator expertise is imperative with this system which to date has been

very productive in the hands of the developers.

In our work we have adopted a pragmatic approach, simpler to implement – although still very challenging.

Our protocol involves the application of cumulative ramp steps with about 20% non-loaded-to-loaded

myocyte conversion success rate. When animals have many months of dietary treatment history, and as

cell isolation from older and diseased hearts is difficult, an optimized loading and stretching approach is

key. The two approaches have not been benchmarked, although it can be predicted that the viscous

components of stiffness are likely underestimated when stretches are slow and relatively small such as in

the current study (Caporizzo & Prosser, 2021). The capacity to directly and reproducibly determine Young's

Modulus (stress/strain) with a slow stretch routine has useful comparative value. An additional aspect of

our pragmatic approach has been to evaluate cardiomyocyte segment stretches as average sarcomere

stretch methods can be challenging to track optically.

With ongoing research, a priority is to undertake longitudinal studies, including both sexes, to investigate

the relative timing of the emergence of *in vivo* and *in vitro* stiffness phenotypes as differences in Ca^{2+}

handling have previously been observed between control male and female rodents (Farrell *et al.*, 2010).

This has high translational relevance as we have previously identified a differential time course of diastolic

dysfunction emergence in females and males, with increased female vulnerability (Reichelt *et al.*, 2013;

Chandramouli *et al.*, 2018).

In conclusion

This is the first study to link elevated intact single cardiomyocyte stiffness with *in vivo* diastolic dysfunction

in a dietary acquired cardiometabolic disease mouse model. The novel findings identify augmented load

dependent TnC-Ca^{2+} interactions as one potential feature of myofilament function which contributes to

cardiomyocyte stiffness in hearts exhibiting cardiometabolic diastolic dysfunction. Taken together, the

findings emphasize the importance of mechanical loading and pacing frequency in revealing latent diastolic

abnormalities at the cardiomyocyte level. The rationale for further studies identifying specific subcellular

and molecular mechanisms of cardiomyocyte stiffness in cardiometabolic disease prior to failure onset is

compelling.

BIOGRAPHY

Johannes Janssens completed his PhD in the Department of Anatomy and Physiology at the University of

Melbourne (Australia) where the current work was performed. After completing his PhD he received a

Fulbright Postdoctoral Scholarship to develop a novel blood biomarker of diabetic heart disease at Cedars-

Sinai Medical Center (Los Angeles, CA) where he continues to work currently. Johannes' research

aspirations involve linking large-scale proteomic features of hearts and cardiomyocytes with their

functional phenotype to develop novel therapeutic and diagnostic strategies for cardiac pathophysiology. In

particular diseases where diastolic dysfunction is a prominent characteristic.

ADDITIONAL INFORMATION

Data availability statement

The data that support the findings of this study are available from the corresponding author upon

reasonable request.

Competing interests

The authors declare that they have no competing interests.

Author contributions

All experiments were performed at the Cardiac Phenomics Laboratory at the University of Melbourne,

Australia. J.V.J., K.M.M. and L.M.D.D. contributed to the conception and research design of the work. All

authors contributed to the acquisition, analysis or interpretation of data for the work and contributed to

drafting the work or revising it critically for important intellectual content. All authors provide final

approval of the version to be published and agreement to be accountable for all aspects of the work.

Funding

This work was supported by grants from the National Health and Medical Research Council of Australia

(LMDD: NHMRCA 1157320), the Diabetes Australia Research Trust (LMDD), the National Heart Foundation

of Australia (LMDD: NHFA), the New Zealand Marsden Fund (KMM: 19-UOA-268), the Health Research

Council of New Zealand (KMM:19/190), and the National Institutes of Health USA (JVE: R01 HL155346-01 &

R01 HL144509-01).

REFERENCES

Abdellatif M et al. (2021). Nicotinamide for the treatment of heart failure with preserved ejection fraction. *Sci Transl Med* **13,** eabd7064.

Aurigemma GP, Zile MR & Gaasch WH (2006). Contractile Behavior of the Left Ventricle in Diastolic Heart Failure. *Circulation* **113,** 296–304.

Bollensdorff C, Lookin O & Kohl P (2011). Assessment of contractility in intact ventricular cardiomyocytes using the dimensionless "Frank-Starling Gain" index. *Pflugers Arch* **462,** 39–48.

Borlaug BA, Melenovsky V, Redfield MM, Kessler K, Chang H-J, Abraham TP & Kass DA (2007). Impact of Arterial Load and Loading Sequence on Left Ventricular Tissue Velocities in Humans. *Journal of the American College of Cardiology* **50,** 1570–1577.

Bub G, Camelliti P, Bollensdorff C, Stuckey DJ, Picton G, Burton RAB, Clarke K & Kohl P (2010). Measurement and analysis of sarcomere length in rat cardiomyocytes in situ and in vitro. *Am J Physiol Heart Circ Physiol* **298,** H1616–H1625.

Caporizzo MA & Prosser BL (2021). Need for Speed: The Importance of Physiological Strain Rates in Determining Myocardial Stiffness. *Front Physiol* **12,** 696694.

Chandramouli C, Reichelt ME, Curl CL, Varma U, Bienvenu LA, Koutsifeli P, Raaijmakers AJA, De Blasio MJ, Qin CX, Jenkins AJ, Ritchie RH, Mellor KM & Delbridge LMD (2018). Diastolic dysfunction is more apparent in STZ-induced diabetic female mice, despite less pronounced hyperglycemia. *Sci Rep* **8,** 2346.

Chemla D, Coirault C, Hébert J-L & Lecarpentier Y (2000). Mechanics of Relaxation of the Human Heart. *Physiology* **15,** 78–83.

Chung CS & Campbell KS (2013). Temperature and transmural region influence functional measurements in unloaded left ventricular cardiomyocytes. *Physiol Rep* **1,** e00158.

Clark JA & Campbell SG (2019). Diverse relaxation rates exist among rat cardiomyocytes isolated from a single myocardial region. *J Physiol* **597,** 711–722.

Curl CL, Danes VR, Bell JR, Raaijmakers AJA, Ip WTK, Chandramouli C, Harding TW, Porrello ER, Erickson JR, Charchar FJ, Kompa AR, Edgley AJ, Crossman DJ, Soeller C, Mellor KM, Kalman JM, Harrap SB & Delbridge LMD (2018). Cardiomyocyte Functional Etiology in Heart Failure With Preserved Ejection Fraction Is Distinctive-A New Preclinical Model. *J Am Heart Assoc* **7,** e007451.

Daniels LJ, Macindoe C, Koutsifeli P, Annandale M, Raaijmakers AJA, Weeks KL, Bell JR, Janssens JV, Curl CL, Delbridge LMD & Mellor KM (2022). Myocardial deformation imaging by 2D speckle tracking echocardiography for assessment of diastolic dysfunction in murine cardiopathology. 2022.08.06.503061. Available at: https://www.biorxiv.org/content/10.1101/2022.08.06.503061v1 [Accessed May 22, 2023].

Davis J, Wen H, Edwards T & Metzger JM (2007). Thin Filament Disinhibition by Restrictive Cardiomyopathy Mutant R193H Troponin I Induces Ca2+-Independent Mechanical Tone and Acute Myocyte Remodeling. *Circulation Research* **100,** 1494–1502.

Delbridge LM & Roos KP (1997). Optical methods to evaluate the contractile function of unloaded isolated cardiac myocytes. *J Mol Cell Cardiol* **29,** 11–25.

Delbridge LMD, Bell JR, Weeks KL, Raaijmakers AJA & Mellor KM (2022). HFpEF etiology - Can focus on sex-specific mechanisms deliver insights for all? *J Mol Cell Cardiol* **173,** 71–72.

Dia M, Gomez L, Thibault H, Tessier N, Leon C, Chouabe C, Ducreux S, Gallo-Bona N, Tubbs E, Bendridi N,
Chanon S, Leray A, Belmudes L, Couté Y, Kurdi M, Ovize M, Rieusset J & Paillard M (2020). Reduced
reticulum–mitochondria Ca2+ transfer is an early and reversible trigger of mitochondrial
dysfunctions in diabetic cardiomyopathy. *Basic Res Cardiol* **115,** 74.

Dowrick JM, Tran K, Garrett AS, Anderson AJ, Nielsen PMF, Taberner AJ & Han J-C (2022). Work-loop
contractions reveal that the afterload-dependent time course of cardiac Ca2+ transients is
modulated by preload. *Journal of Applied Physiology* **133,** 663–675.

Eichhorn EJ, Willard JE, Alvarez L, Kim AS, Glamann DB, Risser RC & Grayburn PA (1992). Are contraction
and relaxation coupled in patients with and without congestive heart failure? *Circulation* **85,** 2132–
2139.

Eisner D, Neher E, Taschenberger H & Smith G (2023). Physiology of intracellular calcium buffering.
Physiological Reviews **103,** 2767–2845.

Eisner DA, Caldwell JL, Trafford AW & Hutchings DC (2020). The Control of Diastolic Calcium in the Heart.
Circ Res **126,** 395–412.

Fabiato A & Fabiato F (1978). Myofilament-generated tension oscillations during partial calcium activation
and activation dependence of the sarcomere length-tension relation of skinned cardiac cells. *J Gen
Physiol* **72,** 667–699.

Farrell SR, Ross JL & Howlett SE (2010). Sex differences in mechanisms of cardiac excitation-contraction
coupling in rat ventricular myocytes. *Am J Physiol Heart Circ Physiol* **299,** H36-45.

Fowler ED, Benoist D, Drinkhill MJ, Stones R, Helmes M, Wüst RCI, Stienen GJM, Steele DS & White E (2015).
Decreased creatine kinase is linked to diastolic dysfunction in rats with right heart failure induced
by pulmonary artery hypertension. *J Mol Cell Cardiol* **86,** 1–8.

Gillebert TC, Leite-Moreira AF & De Hert SG (2000). Load dependent diastolic dysfunction in heart failure.
Heart Fail Rev **5,** 345–355.

Gonçalves N, Silva AF, Rodrigues PG, Correia E, Moura C, Eloy C, Roncon-Albuquerque R, Falcão-Pires I &
Leite-Moreira AF (2016). Early cardiac changes induced by a hypercaloric Western-type diet in
"subclinical" obesity. *American Journal of Physiology-Heart and Circulatory Physiology* **310,** H655–
H666.

Grundy D (2015). Principles and standards for reporting animal experiments in The Journal of Physiology
and Experimental Physiology. *J Physiol* **593,** 2547–2549.

Hamdani N, Franssen C, Lourenço A, Falcão-Pires I, Fontoura D, Leite S, Plettig L, López B, Ottenheijm CA,
Becher PM, González A, Tschöpe C, Díez J, Linke WA, Leite-Moreira AF & Paulus WJ (2013).
Myocardial Titin Hypophosphorylation Importantly Contributes to Heart Failure With Preserved
Ejection Fraction in a Rat Metabolic Risk Model. *Circulation: Heart Failure* **6,** 1239–1249.

Heather LC, Hafstad AD, Halade GV, Harmancey R, Mellor KM, Mishra PK, Mulvihill EE, Nabben M,
Nakamura M, Rider OJ, Ruiz M, Wende AR & Ussher JR (2022). Guidelines on models of diabetic
heart disease. *Am J Physiol Heart Circ Physiol* **323,** H176–H200.

Helmes M, Najafi A, Palmer BM, Breel E, Rijnveld N, Iannuzzi D & van der Velden J (2016). Mimicking the
cardiac cycle in intact cardiomyocytes using diastolic and systolic force clamps; measuring power
output. *Cardiovasc Res* **111,** 66–73.

Hopf A-E, Andresen C, Kötter S, Isić M, Ulrich K, Sahin S, Bongardt S, Röll W, Drove F, Scheerer N,
Vandekerckhove L, De Keulenaer GW, Hamdani N, Linke WA & Krüger M (2018). Diabetes-Induced
Cardiomyocyte Passive Stiffening Is Caused by Impaired Insulin-Dependent Titin Modification and
Can Be Modulated by Neuregulin-1. *Circulation Research* **123,** 342–355.

Howlett SE, Grandy SA & Ferrier GR (2006). Calcium spark properties in ventricular myocytes are altered in
aged mice. *American Journal of Physiology-Heart and Circulatory Physiology* **290,** H1566–H1574.

Jackson AM et al. (2022). Diabetes and pre-diabetes in patients with heart failure and preserved ejection
fraction. *Eur J Heart Fail* **24,** 497–509.

Janssens JV, Ma B, Brimble MA, Van Eyk JE, Delbridge LMD & Mellor KM (2018). Cardiac troponins may be
irreversibly modified by glycation: novel potential mechanisms of cardiac performance modulation.
Sci Rep **8,** 16084.

Janssens JV, Raaijmakers AJA, Weeks KL, Bell JR, Mellor KM, Curl CL & Delbridge LMD (2024). The
cardiomyocyte origins of diastolic dysfunction: cellular components of myocardial stiffness. *Am J
Physiol Heart Circ Physiol*; DOI: 10.1152/ajpheart.00334.2023.

Kamon T, Kaneko H, Itoh H, Kiriyama H, Koyama K, Fujiu K, Morita H, Uno K, Hayashi N & Komuro I (2020).
Association between insulin resistance and left ventricular diastolic dysfunction in non-diabetic
general population. *European Heart Journal* **41,** ehaa946.3057.

Khokhlova A, Myachina T, Volzhaninov D, Butova X, Kochurova A, Berg V, Gette I, Moroz G, Klinova S,
Minigalieva I, Solovyova O, Danilova I, Sokolova K, Kopylova G & Shchepkin D (2022). Type 1
Diabetes Impairs Cardiomyocyte Contractility in the Left and Right Ventricular Free Walls but
Preserves It in the Interventricular Septum. *Int J Mol Sci* **23,** 1719.

Kilfoil PJ, Lotteau S, Zhang R, Yue X, Aynaszyan S, Solymani RE, Cingolani E, Marbán E & Goldhaber JI (2020).
Distinct features of calcium handling and β-adrenergic sensitivity in heart failure with preserved
versus reduced ejection fraction. *J Physiol* **598,** 5091–5108.

Kittleson MM, Panjrath GS, Amancherla K, Davis LL, Deswal A, Dixon DL, Januzzi JL & Yancy CW (2023). 2023
ACC Expert Consensus Decision Pathway on Management of Heart Failure With Preserved Ejection
Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. *Journal
of the American College of Cardiology* **81,** 1835–1878.

Lab MJ, Allen DG & Orchard CH (1984). The effects of shortening on myoplasmic calcium concentration and
on the action potential in mammalian ventricular muscle. *Circulation Research* **55,** 825–829.

Lam CSP, Arnott C, Beale AL, Chandramouli C, Hilfiker-Kleiner D, Kaye DM, Ky B, Santema BT, Sliwa K &
Voors AA (2019). Sex differences in heart failure. *European Heart Journal* **40,** 3859–3868c.

Leopoldo AS, Sugizaki MM, Lima-Leopoldo AP, do Nascimento AF, de Azevedo Melo Luvizotto R, de Campos
DHS, Okoshi K, Pai-Silva MD, Padovani CR & Cicogna AC (2010). Cardiac remodeling in a rat model
of diet-induced obesity. *Can J Cardiol* **26,** 423–429.

Leung M, Phan V, Whatmough M, Heritier S, Wong VW & Leung DY (2015). Left ventricular diastolic reserve
in patients with type 2 diabetes mellitus. *Open Heart* **2,** e000214.

Li K-L, Methawasin M, Tanner BCW, Granzier HL, Solaro RJ & Dong W-J (2019). Sarcomere length–
dependent effects on Ca2+-troponin regulation in myocardium expressing compliant titin. *J Gen
Physiol* **151,** 30–41.

Lim CC, Apstein CS, Colucci WS & Liao R (2000). Impaired Cell Shortening and Relengthening with Increased
Pacing Frequency are Intrinsic to the Senescent Mouse Cardiomyocyte. *Journal of Molecular and*
Cellular Cardiology **32,** 2075–2082.

Llano-Diez M, Sinclair J, Yamada T, Zong M, Fauconnier J, Zhang S-J, Katz A, Jardemark K, Westerblad H,
Andersson DC & Lanner JT (2016). The Role of Reactive Oxygen Species in β-Adrenergic Signaling in
Cardiomyocytes from Mice with the Metabolic Syndrome. *PLoS One* **11,** e0167090.

Lo Q & Thomas L (2010). Echocardiographic evaluation of diastolic heart failure. *Australas J Ultrasound Med*
13, 14–26.

Loescher CM, Freundt JK, Unger A, Hessel AL, Kühn M, Koser F & Linke WA (2023). Titin governs myocardial
passive stiffness with major support from microtubules and actin and the extracellular matrix. *Nat*
Cardiovasc Res **2,** 991–1002.

Loescher CM, Hobbach AJ & Linke WA (2022). Titin (TTN): from molecule to modifications, mechanics, and
medical significance. *Cardiovascular Research* **118,** 2903–2918.

Louch WE (2020). A horse of a different colour: distinct mechanisms of HFpEF and HFrEF. *J Physiol* **598,**
5005–5006.

McHugh K, DeVore AD, Wu J, Matsouaka RA, Fonarow GC, Heidenreich PA, Yancy CW, Green JB, Altman N
& Hernandez AF (2019). Heart Failure With Preserved Ejection Fraction and Diabetes: JACC State-
of-the-Art Review. *J Am Coll Cardiol* **73,** 602–611.

Mellor KM, Bell JR, Young MJ, Ritchie RH & Delbridge LMD (2011). Myocardial autophagy activation and
suppressed survival signaling is associated with insulin resistance in fructose-fed mice. *Journal of*
Molecular and Cellular Cardiology **50,** 1035–1043.

Mellor KM, Wendt IR, Ritchie RH & Delbridge LMD (2012). Fructose diet treatment in mice induces
fundamental disturbance of cardiomyocyte Ca2+ handling and myofilament responsiveness. *Am J*
Physiol Heart Circ Physiol **302,** H964-972.

Methawasin M, Farman GP, Granzier-Nakajima S, Strom J, Kiss B, Smith JE & Granzier H (2022). Shortening
the thick filament by partial deletion of titin's C-zone alters cardiac function by reducing the
operating sarcomere length range. *J Mol Cell Cardiol* **165,** 103–114.

Mirsky I (1976). Assessment of passive elastic stiffness of cardiac muscle: mathematical concepts,
physiologic and clinical considerations, directions of future research. *Prog Cardiovasc Dis* **18,** 277–
308.

Najafi A, van de Locht M, Schuldt M, Schönleitner P, van Willigenburg M, Bollen I, Goebel M, Ottenheijm
CAC, van der Velden J, Helmes M & Kuster DWD (2019). End-diastolic force pre-activates
cardiomyocytes and determines contractile force: role of titin and calcium. *J Physiol* **597,** 4521–
4531.

Najafi A, Sequeira V, Helmes M, Bollen IAE, Goebel M, Regan JA, Carrier L, Kuster DWD & Van Der Velden J
(2016). Selective phosphorylation of PKA targets after β-adrenergic receptor stimulation impairs
myofilament function in Mybpc3-targeted HCM mouse model. *Cardiovasc Res* **110,** 200–214.

Natali AJ, Turner DL, Harrison SM & White E (2001). Regional effects of voluntary exercise on cell size and
contraction-frequency responses in rat cardiac myocytes. *Journal of Experimental Biology* **204,**
1191–1199.

Patil VC, Patil HV, Shah KB, Vasani JD & Shetty P (2011). Diastolic dysfunction in asymptomatic type 2
diabetes mellitus with normal systolic function. *J Cardiovasc Dis Res* **2,** 213–222.

Peyronnet R, Bollensdorff C, Capel RA, Rog-Zielinska EA, Woods CE, Charo DN, Lookin O, Fajardo G, Ho M,
Quertermous T, Ashley EA & Kohl P (2017). Load-dependent effects of apelin on murine
cardiomyocytes. *Prog Biophys Mol Biol* **130,** 333–343.

Pulinilkunnil T, Kienesberger PC, Nagendran J, Sharma N, Young ME & Dyck JRB (2014). Cardiac-specific
adipose triglyceride lipase overexpression protects from cardiac steatosis and dilated
cardiomyopathy following diet-induced obesity. *Int J Obes* **38,** 205–215.

Redfield MM (2016). Heart Failure with Preserved Ejection Fraction. *N Engl J Med* **375,** 1868–1877.

Reichelt ME, Mellor KM, Bell JR, Chandramouli C, Headrick JP & Delbridge LMD (2013). Sex, sex steroids,
and diabetic cardiomyopathy: making the case for experimental focus. *Am J Physiol Heart Circ
Physiol* **305,** H779-792.

Robinson P, Liu X, Sparrow A, Patel S, Zhang Y-H, Casadei B, Watkins H & Redwood C (2018). Hypertrophic
cardiomyopathy mutations increase myofilament Ca2+ buffering, alter intracellular Ca2+ handling,
and stimulate Ca2+-dependent signaling. *J Biol Chem* **293,** 10487–10499.

Roos KP, Brady AJ & Tan ST (1982). Direct measurement of sarcomere length from isolated cardiac cells.
American Journal of Physiology-Heart and Circulatory Physiology **242,** H68–H78.

Rozenbaum Z, Topilsky Y, Khoury S, Pereg D & Laufer-Perl M (2019). Association of body mass index and
diastolic function in metabolically healthy obese with preserved ejection fraction. *Int J Cardiol* **277,**
147–152.

Saad NS, Mashali MA, Repas SJ & Janssen PML (2023). Altering Calcium Sensitivity in Heart Failure: A
Crossroads of Disease Etiology and Therapeutic Innovation. *Int J Mol Sci* **24,** 17577.

Satoh H, Delbridge LM, Blatter LA & Bers DM (1996). Surface:volume relationship in cardiac myocytes
studied with confocal microscopy and membrane capacitance measurements: species-dependence
and developmental effects. *Biophys J* **70,** 1494–1504.

Shimkunas R et al. (2021). Mechanical Load Regulates Excitation-Ca2+ Signaling-Contraction in
Cardiomyocyte. *Circ Res* **128,** 772–774.

Sotomi Y et al. (2021). Sex Differences in Heart Failure With Preserved Ejection Fraction. *J Am Heart Assoc*
10, e018574.

Sowers JR, Habibi J, Jia G, Bostick B, Manrique-Acevedo C, Lastra G, Yang Y, Chen D, Sun Z, Domeier TL,
Durante W, Whaley-Connell AT, Hill MA, Jaisser F, DeMarco VG & Aroor AR (2020). Endothelial
sodium channel activation promotes cardiac stiffness and diastolic dysfunction in Western diet fed
female mice. *Metabolism* **109,** 154223.

Varian KD & Janssen PML (2007). Frequency-dependent acceleration of relaxation involves decreased
myofilament calcium sensitivity. *American Journal of Physiology-Heart and Circulatory Physiology*
292, H2212–H2219.

Villalobos Lizardi JC, Baranger J, Nguyen MB, Asnacios A, Malik A, Lumens J, Mertens L, Friedberg MK,
Simmons CA, Pernot M & Villemain O (2022). A guide for assessment of myocardial stiffness in
health and disease. *Nat Cardiovasc Res* **1,** 8–22.

Wang Q, Wang Y, West TM, Liu Y, Reddy GR, Barbagallo F, Xu B, Shi Q, Deng B, Wei W & Xiang YK (2020).
Carvedilol induces biased β1 adrenergic receptor-Nitric oxide synthase 3-cyclic guanylyl
monophosphate signaling to promote cardiac contractility. *Cardiovasc Res*cvaa266.

Wells SP, Raaijmakers AJA, Curl CL, O'Shea C, Hayes S, Mellor KM, Kalman JM, Kirchhof P, Pavlovic D,
Delbridge LMD & Bell JR (2023). Localized cardiomyocyte lipid accumulation is associated with
slowed epicardial conduction in rats. *J Gen Physiol* **155,** e202213296.

West TM, Wang Q, Deng B, Zhang Y, Barbagallo F, Reddy GR, Chen D, Phan KS, Xu B, Isidori A & Xiang YK
(2019). Phosphodiesterase 5 Associates With β2 Adrenergic Receptor to Modulate Cardiac Function
in Type 2 Diabetic Hearts. *JAHA*; DOI: 10.1161/JAHA.119.012273.

White E, Boyett MR & Orchard CH (1995). The effects of mechanical loading and changes of length on single
guinea-pig ventricular myocytes. *The Journal of Physiology* **482,** 93–107.

Wingard MC, Dalal S, Shook PL, Myers R, Connelly BA, Thewke DP, Singh M & Singh K (2021). Deficiency of
ataxia-telangiectasia mutated kinase modulates functional and biochemical parameters of the
heart in response to Western-type diet. *Am J Physiol Heart Circ Physiol* **320,** H2324–H2338.

Zhou P & Pu WT (2016). Recounting Cardiac Cellular Composition. *Circulation Research* **118,** 368–370.

FIGURE LEGENDS

Fig.1 Systemic indicators of cardiometabolic disease and *in vivo* diastolic function in high-fat/sugar diet
(HFSD) vs control (CTRL) mice.
A. Body weight elevation at 5 week intervals up to 30 weeks of dietary treatment (last timepoint full cohort
intact. N = 16 – 17 mice per group).
B. Glucose tolerance testing at 23-24 weeks post dietary intervention; N = 6 mice per group).
C. Echocardiography exemplar blood flow and tissue doppler images from which diastolic indices were
derived for CTRL and HFSD mice. Panels with orange outline: HFSD mitral blood flow & annular wall motion.
Panels with blue outline: CTRL mitral blood flow & annular wall motion. (N = 14-16 mice per group)
D. Early diastolic mitral annular blood flow (E) was increased in HFSD.
E. Early diastolic mitral annular wall velocity (e') was decreased in HFSD.
F. Deceleration time of early diastolic mitral annular blood flow was decreased in HFSD.
G. Ratio of early diastolic blood flow to wall velocity (E/e') was increased in HFSD.
944
945 Time-course data analysed using two-way repeated measures ANOVA with Sidak post-hoc test with p-value
946 for 'time x diet' effect presented. Comparisons between two treatment groups performed using student's
947 t-test. Data presented as mean ± standard deviation.
948
949
Fig.2 Non-loaded sarcomere and Ca^{2+} dynamics in CTRL and HFSD cardiomyocytes*.*
A. Mean sarcomere contraction cycles presented as a percentage of diastolic sarcomere length for CTRL
and HFSD cardiomyocytes
B-C. Mean diastolic sarcomere length and extent of sarcomere shortening for CTRL and HFSD
cardiomyocytes.
D. Mean sarcomere shortening and lengthening rate records (time derivative of sarcomere contraction-
relaxation cycle) for CTRL and HFSD cardiomyocytes. The nadir and zenith represent maximal shortening
and lengthening rates respectively.
E-F. Maximal rate of lengthening and shortening rates for CTRL and HFSD cardiomyocytes.
G. Mean Ca^{2+} transient records for CTRL and HFSD cardiomyocytes.
H-J. Mean Ca^{2+} levels (systolic and diastolic) and time constant of transient decay (*tau*) in HFSD and CTRL
cardiomyocytes.
962
963 Mean data computed from average of 20 cycles per cardiomyocyte to derive group mean (N = 6-8 mice per
964 group, n = 32-49 cells per group). Comparisons between two treatment groups performed using student's
965 t-test (all p>0.05, ns). Data presented as mean ± standard deviation.
966

Fig. 3 Sarcomere shortening and lengthening in non-loaded and loaded pacing states.
A. Exemplar non-loaded and loaded sarcomere contraction cycles for the same cardiomyocyte (CTRL).
B-C. Extent of sarcomere shortening and maximum rate of lengthening, measured for each cardiomyocyte
in non-loaded and loaded state for CTRL and HFSD groups combined (N = 9 mice, n = 25 cells). Mean data
for CTRL and HFSD are presented as grey points to the left (non-loaded) and right (loaded) of individual
cardiomyocyte data.
D. Correlation of loading associated sarcomere shortening change and maximum rate of lengthening
change for CTRL and HFSD combined.
E. Mean loading effect on maximum deformation rate during lengthening ((μm/s) / μm; maximum rate of
lengthening (μm/s) normalized by shortening extent (μm)) for CTRL vs HFSD. (CTRL: N = 5 mice, n = 14 cells;
HFSD: N = 4 mice, n = 12 cells). Significant ANOVA interaction effect (diet treatment x load state).
F. Mean decrement in deformation rate during lengthening ((μm/s) / μm).
979
980 Comparison of cardiomyocyte responses in different loading conditions analysed by paired t-test.
981 Comparison of cardiomyocytes from different treatment groups analysed by unpaired t-test. A 2-way
982 ANOVA with repeated measures used for analysis of two independent variables. Data fitted with linear
983 regression model and calculation of Pearson's correlation coefficient. Data presented as mean ± standard
984 deviation.
985
986
Fig. 4. Cardiomyocyte standardized stretch protocol.
A. Image of a left ventricular cardiomyocyte longitudinally attached to glass rods. Edge-tracking used to
measure inter-rod distance (marked as blue bars) for cardiomyocyte segment analyses. The delineated
region of interest (marked pink) represents the area from which Fast Fourier Transform analysis of
sarcomere striations was performed to derive sarcomere-specific data.
B. Exemplar traces depict the change in glass rod internal distance in response to successive 0.5V piezo
motor inputs used for calibration of stretch protocol for each individual cardiomyocyte to allow normalized
comparisons.
C. Measuring stress and strain during progressive cardiomyocyte stretch. Exemplar force, sarcomere length
and rod edge-tracker traces during the first stretch of a stress-length protocol.
D. Comparison of variability of cell stretches performed using a uniform (preset) stretch constant and the
individually derived stretch calibration constant. Strain was measured as the %change in segment length
relative to the initial segment length. Reproducibility is optimized using the individual stretch calibration
constant. Data shown as median ± interquartile range.
1001

Fig. 5 Quantifying diastolic stiffness in CTRL vs HFSD cardiomyocytes using strain normalization methods

A. Exemplar stress recordings of CTRL and HFSD cardiomyocytes undergoing 'stress-length' protocol (2Hz

paced). Diastolic baseline stress shifts indicated by broken line.

B. Diastolic (pair of lower lines) and systolic (pair of upper lines) sarcomere mean stress- sarcomere length

(SL) relations in CTRL cardiomyocytes (N = 5 mice, n = 9 cells).

C. Diastolic mean stress-sarcomere length relation (linear regression fit) for HFSD and CTRL.

D. Diastolic stress (mN/mm^2) normalized relative to sarcomere stretch (µm) yields sarcomere stress/strain =

stiffness [(mN/mm^2) /strain (µm)] for HFSD and CTRL.

E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL.

F. Diastolic stress (mN/mm^2) normalized relative to segment stretch (% length) yields cardiomyocyte

segment stress/strain = stiffness [(mN/mm^2) /strain (% length)] for HFSD & CTRL.

G. Correlation (linear regression fit) of *in vitro* cardiomyocyte segment diastolic stiffness (stress/strain) with

in vivo diastolic dysfunction E/e' (blood flow/wall motion) in CTRL and HFSD. Cardiomyocyte data averaged

per heart.

1016

1017 Comparisons between two groups performed using student's t-test. Data fitted with linear regression

1018 model and calculation of Pearson's correlation coefficient, presented as mean ± standard deviation (D, F)

1019 and mean ± standard error of the mean (B, C, E) . For C. & D. (CTRL: N=4 mice, n=8 cells; HFSD: N=2 mice,

1020 n=7 cells). For E.&F. (CTRL: N=6 mice, n = 16 cells; HFSD: N=6 mice, n=18 cells).

1021

1022

Fig. 6 Load dependence of cardiomyocyte Ca^{2+} transient decay in CTRL and HFSD cardiomyocytes.

A-B. Time constant of Ca^{2+} transient decay (tau) and time to peak Ca^{2+} transient in cardiomyocytes with

different load conditions: loaded (L) & stretched (S) shown as fold change relative to non-loaded (N) for

CTRL and HFSD.

C-D. Exemplar Ca^{2+} transient decay records, with a mono-exponential decay function fit, in non-loaded and

stretched CTRL and HFSD cardiomyocytes. Horizontal black broken lines intersect with mono-exponential fit

at 36.8% (1/e) transient amplitude. Vertical broken lines from intersection points indicate the time constant

of Ca^{2+} transient decay (tau).

1031

1032 For analysis of two independent variables a 2-way ANOVA with repeated measures used (CTRL: N = 5 mice,

1033 n = 13 cells; HFSD: N = 4 mice, n = 12 cells). Data are presented as mean ± standard deviation.

1034

Fig. 7 Pacing induced changes in cardiomyocyte cytosolic Ca^{2+} levels and stiffness in CTRL & HFSD.

A. Exemplar CTRL and HFSD cardiomyocyte Ca^{2+} transients at 2 and 4 Hz pacing.

B. Diastolic cardiomyocyte Ca^{2+} levels measured before and after transition from 2 to 4 Hz pacing in HFSD &
CTRL.

C. Diastolic cardiomyocyte Ca^{2+} level mean shift with transition from 2 to 4 Hz pacing in HFSD & CTRL.

D. Diastolic sarcomere length measured before and after transition from 2 to 4 Hz pacing in HFSD & CTRL.

E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL at
2 and 4 Hz pacing.

F. Diastolic mean cardiomyocyte segment stress-length relation slope (mN/mm^2) normalized relative to
segment stretch (% length) yields cardiomyocyte mean segment stress/strain = stiffness
[(mN/mm^2) /strain (% length)] for HFSD and CTRL at 2 and 4 Hz pacing.

1046

1047 Comparison between two groups performed using unpaired t-test. With two independent variables a
1048 repeated measures 2-way ANOVA Sidaks post-hoc analysis was performed. Data are presented as mean ±
1049 standard deviation (C), mean ± standard error of the mean (E) or mean with paired individual points (B, D,
1050 F). For B. & C. (CTRL: N = 4 mice, n = 9 cells; HFSD: N = 4 mice, n = 8 cells). For D. & E. (CTRL: N = 4 mice, n =
1051 7 cells; HFSD: N = 4 mice, n = 9 cells). For F. (CTRL: N = 4 mice, n = 8 cells; HFSD: N = 4 mice, n = 9 cells).

Stiff hearts are characterized by stiff cardiomyocytes in metabolic disease

Loaded cardiomyocytes

Stiffness: heart vs cardiomyocytes

Abstract Figure Legend

Understanding cardiomyocyte stiffness components is an important priority for identifying new therapeutics for diastolic dysfunction, a key feature of cardiometabolic disease. In this study cardiac function was measured *in vivo* (echocardiography) for mice fed a high-fat/sugar diet (HFSD, ≥25weeks). Performance of intact isolated cardiomyocytes derived from the same hearts was measured during pacing under non-loaded, loaded and stretched conditions *in vitro.* Calibrated cardiomyocyte stretches demonstrated that stiffness was elevated in HFSD cardiomyocytes *in vitro* and correlated with diastolic dysfunction (E/e') *in vivo*. These findings show that stiff hearts are characterized by stiff cardiomyocytes in metabolic disease.

Table 1.	Echocardiographic parameters for CTRL and HFSD mice		
	CTRL	**HFSD**	**P-value**
Heart rate (bpm)	515±74	551±29	0.174
LV mass (mg)	125±27	131±21	0.314
LVIDd (mm)	3.60±0.17	4.16±0.47	0.004
IVSd (mm)	0.84±0.06	0.80±0.10	0.349
LVPWd (mm)	0.94±0.03	0.94±0.10	0.830
RWT (ratio)	0.49±0.06	0.43±0.09	0.079
Ejection Fraction (%)	67.8±3.7	64.9±3.2	0.093
Fractional Shortening (%)	31.8±3.6	30.6±2.1	0.400
E wave (mm/s)	472±74	569±65	0.003
A wave (mm/s)	348±99	400±116	0.190
e' wave (mm/s)	28.5±5.1	25.2±3.5	0.046
a' wave (mm/s)	23.5±7.8	23.5±6.4	0.995
E/A	1.51±0.41	1.47±0.37	0.095
e'/a'	1.33±0.44	1.12±0.39	0.171
E/e'	16.7±3.9	22.3±6.0	0.005
MV Deceleration Time (ms)	30.8±8.7	19.7±5.9	<0.001

Table 1. Echocardiographic parameters in CTRL and HFSD mice.
LVIDd left ventricular internal diameter at end diastole, *IVSd* interventricular septum diameter at end diastole, *LVPWd* left ventricular posterior wall thickness at end diastole, *RWT* relative wall thickness, *E wave* mitral valve blood flow velocity during the early ventricular filling phase, *A wave* mitral valve blood flow velocity during atrial contraction, *e' wave* mitral annulus tissue motion velocity during the early ventricular filling phase, *a' wave* mitral annulus tissue motion velocity during atrial contraction, *MV* mitral valve.

Comparisons between two treatment groups performed using Students' unpaired t-test (N = 9-15 mice per group). Data are presented as mean ± standard deviation.

Table 2. Stress-length relation mean values for CTRL and HFSD mouse cardiomyocytes			
	CTRL 2 Hz	**HFSD 2 Hz**	**p-value**
Systolic Stress-SL Relation slope ($(mN/mm^2) / \mu m$)	6.54±4.97	16.35±15.09	0.087
Systolic Stress-Length Relation slope ($(mN/mm^2) / \%length$)	0.013±0.011	0.022±0.011	0.022
Systolic : Diastolic Stress-SL relation 'Frank-Starling Gain Index' (ratio)	2.40±1.06	2.77±1.45	0.562
Systolic : Diastolic Stress-Length relation (ratio)	1.62±0.25	1.73±0.50	0.471

Table 2. Stress-length relation mean values for CTRL and HFSD mouse cardiomyocytes.
Comparisons between two treatment groups performed using Students' unpaired t-test (CTRL: N = 4-6 mice, n = 9-17 cells cells; HFSD: N = 2-6 mice, n = 7-19 cells). Data are presented as mean ± standard deviation.

Table 3. Ca^{2+} transient parameters for loaded CTRL and HFSD mouse cardiomyocytes							
	CTRL 2 Hz	**CTRL 4 Hz**	**HFSD 2 Hz**	**HFSD 4 Hz**	**Frequency p-value**	**Diet p-value**	**Interaction p-value**
Systolic Ca^{2+} (F$_{340:380}$)	0.93±0.08	1.05±0.10$^{\#}$	0.99±0.15	1.08±0.20	p = 0.0004	p = 0.571	p = 0.100
Amplitude Ca^{2+} (F$_{340:380}$)	0.26±0.06	0.28±0.06	0.32±0.10	0.33±0.11	p = 0.569	p = 0.0672	p = 0.723
Time constant of Ca^{2+} decay (tau, sec)	0.21±0.07	0.09±0.02$^{\#}$	0.17±0.07	0.09±0.03$^{\#}$	p < 0.0001	p = 0.207	p = 0.350

Table 3. Ca^{2+} transient parameters for loaded CTRL and HFSD mouse cardiomyocytes.
Comparisons in data with two independent variable were performed using a mixed analysis 2-way ANOVA with Sidak's post-hoc analysis (CTRL, 2 Hz: N = 5 mice, n = 13 cells; CTRL, 4 Hz: N = 4 mice, n = 9 cells; HFSD, 2 Hz: N = 5 mice, n = 14 cells; HFSD, 4 Hz: N = 4 mice, n = 8 cells). Data are presented as mean ± standard deviation. P-values for ANOVA effects (Frequency, Diet, Interaction) are listed within the table. #<0.05: Significant pairwise post-hoc frequency differences vs CTRL 2 Hz or HFSD 2 Hz. No pairwise post-hoc treatment differences were observed.

Fig.1 Systemic indicators of cardiometabolic disease and *in vivo* diastolic function in high-fat/sugar diet (HFSD) vs control (CTRL) mice.

A. Body weight elevation at 5 week intervals up to 30 weeks of dietary treatment (last timepoint full cohort intact. N = 16 – 17 mice per group).

B. Glucose tolerance testing at 23-24 weeks post dietary intervention; N = 6 mice per group).

C. Echocardiography exemplar blood flow and tissue doppler images from which diastolic indices were derived for CTRL and HFSD mice. Panels with orange outline: HFSD mitral blood flow & annular wall motion. Panels with blue outline: CTRL mitral blood flow & annular wall motion. (N = 14-16 mice per group)

D. Early diastolic mitral annular blood flow (E) was increased in HFSD.

E. Early diastolic mitral annular wall velocity (e') was decreased in HFSD.

F. Deceleration time of early diastolic mitral annular blood flow was decreased in HFSD.

G. Ratio of early diastolic blood flow to wall velocity (E/e') was increased in HFSD.

Time-course data analysed using two-way repeated measures ANOVA with Sidak post-hoc test with p-value for 'time x diet' effect presented. Comparisons between two treatment groups performed using student's t-test. Data presented as mean ± standard deviation.

Fig.2 Non-loaded sarcomere and Ca²⁺ dynamics in CTRL and HFSD cardiomyocytes.
A. Mean sarcomere contraction cycles presented as a percentage of diastolic sarcomere length for CTRL and HFSD cardiomyocytes
B-C. Mean diastolic sarcomere length and extent of sarcomere shortening for CTRL and HFSD cardiomyocytes.
D. Mean sarcomere shortening and lengthening rate records (time derivative of sarcomere contraction-relaxation cycle) for CTRL and HFSD cardiomyocytes. The nadir and zenith represent maximal shortening and lengthening rates respectively.
E-F. Maximal rate of lengthening and shortening rates for CTRL and HFSD cardiomyocytes.
G. Mean Ca²⁺ transient records for CTRL and HFSD cardiomyocytes.
H-J. Mean Ca²⁺ levels (systolic and diastolic) and time constant of transient decay (*tau*) in HFSD and CTRL cardiomyocytes.

Mean data computed from average of 20 cycles per cardiomyocyte to derive group mean (N = 6-8 mice per group, n = 32-49 cells per group). Comparisons between two treatment groups performed using student's t-test (all p>0.05, ns). Data presented as mean ± standard deviation.

Fig. 3 Sarcomere shortening and lengthening in non-loaded and loaded pacing states.
A. Exemplar non-loaded and loaded sarcomere contraction cycles for the same cardiomyocyte (CTRL).
B-C. Extent of sarcomere shortening and maximum rate of lengthening, measured for each cardiomyocyte in non-loaded and loaded state for CTRL and HFSD groups combined (N = 9 mice, n = 25 cells). Mean data for CTRL and HFSD are presented as grey points to the left (non-loaded) and right (loaded) of individual cardiomyocyte data.
D. Correlation of loading associated sarcomere shortening change and maximum rate of lengthening change for CTRL and HFSD combined.
E. Mean loading effect on maximum deformation rate during lengthening ((μm/s) / μm; maximum rate of lengthening (μm/s) normalized by shortening extent (μm)) for CTRL vs HFSD. (CTRL: N = 5 mice, n = 14 cells; HFSD: N = 4 mice, n = 12 cells). Significant ANOVA interaction effect (diet treatment x load state).
F. Mean decrement in deformation rate during lengthening ((μm/s) / μm).

Comparison of cardiomyocyte responses in different loading conditions analysed by paired t-test. Comparison of cardiomyocytes from different treatment groups analysed by unpaired t-test. A 2-way ANOVA with repeated measures used for analysis of two independent variables. Data fitted with linear regression model and calculation of Pearson's correlation coefficient. Data presented as mean ± standard deviation.

Fig. 4. Cardiomyocyte standardized stretch protocol.
A. Image of a left ventricular cardiomyocyte longitudinally attached to glass rods. Edge-tracking used to measure inter-rod distance (marked as blue bars) for cardiomyocyte segment analyses. The delineated region of interest (marked pink) represents the area from which Fast Fourier Transform analysis of sarcomere striations was performed to derive sarcomere-specific data.
B. Exemplar traces depict the change in glass rod internal distance in response to successive 0.5V piezo motor inputs used for calibration of stretch protocol for each individual cardiomyocyte to allow normalized comparisons.
C. Measuring stress and strain during progressive cardiomyocyte stretch. Exemplar force, sarcomere length and rod edge-tracker traces during the first stretch of a stress-length protocol.
D. Comparison of variability of cell stretches performed using a uniform (preset) stretch constant and the individually derived stretch calibration constant. Strain was measured as the %change in segment length relative to the initial segment length. Reproducibility is optimized using the individual stretch calibration constant. Data shown as median ± interquartile range.

Fig. 5 Quantifying diastolic stiffness in CTRL vs HFSD cardiomyocytes using strain normalization methods

A. Exemplar stress recordings of CTRL and HFSD cardiomyocytes undergoing 'stress-length' protocol (2Hz paced). Diastolic baseline stress shifts indicated by broken line.

B. Diastolic (pair of lower lines) and systolic (pair of upper lines) sarcomere mean stress- sarcomere length (SL) relations in CTRL cardiomyocytes (N = 5 mice, n = 9 cells).

C. Diastolic mean stress-sarcomere length relation (linear regression fit) for HFSD and CTRL.

D. Diastolic stress (mN/mm^2) normalized relative to sarcomere stretch (μm) yields sarcomere stress/strain = stiffness [(mN/mm^2) /strain (μm)] for HFSD and CTRL.

E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL..

F. Diastolic stress (mN/mm^2) normalized relative to segment stretch (% length) yields cardiomyocyte segment stress/strain = stiffness [(mN/mm^2) /strain (% length)] for HFSD & CTRL.

G. Correlation (linear regression fit) of *in vitro* cardiomyocyte segment diastolic stiffness (stress/strain) with *in vivo* diastolic dysfunction E/e' (blood flow/wall motion) in CTRL and HFSD. Cardiomyocyte data averaged per heart.

Comparisons between two groups performed using student's t-test. Data fitted with linear regression model and calculation of Pearson's correlation coefficient, presented as mean ± standard deviation (D, F) and mean ± standard error of the mean (B, C, E) . For **C. & D.** (CTRL: N=4 mice, n=8 cells; HFSD: N=2 mice, n=7 cells). For **E.&F.** (CTRL: N=6 mice, n = 16 cells; HFSD: N=6 mice, n=18 cells).

(N: Non-loaded, L: Loaded, S: Stretched)

Fig. 6 Load dependence of cardiomyocyte Ca²⁺ transient decay in CTRL and HFSD cardiomyocytes.

A-B. Time constant of Ca²⁺ transient decay (tau) and time to peak Ca²⁺ transient in cardiomyocytes with different load conditions: loaded (L) & stretched (S) shown as fold change relative to non-loaded (N) for CTRL and HFSD.

C-D. Exemplar Ca²⁺ transient decay records, with a mono-exponential decay function fit, in non-loaded and stretched CTRL and HFSD cardiomyocytes. Horizontal black broken lines intersect with mono-exponential fit at 36.8% (1/e) transient amplitude. Vertical broken lines from intersection points indicate the time constant of Ca²⁺ transient decay (tau).

For analysis of two independent variables a 2-way ANOVA with repeated measures used (CTRL: N = 5 mice, n = 13 cells; HFSD: N = 4 mice, n = 12 cells). Data are presented as mean ± standard deviation.

Fig. 7 Pacing induced changes in cardiomyocyte cytosolic Ca²⁺ levels and stiffness in CTRL & HFSD.
A. Exemplar CTRL and HFSD cardiomyocyte Ca²⁺ transients at 2 and 4 Hz pacing.
B. Diastolic cardiomyocyte Ca²⁺ levels measured before and after transition from 2 to 4 Hz pacing in HFSD & CTRL.
C. Diastolic cardiomyocyte Ca²⁺ level mean shift with transition from 2 to 4 Hz pacing in HFSD & CTRL.
D. Diastolic sarcomere length measured before and after transition from 2 to 4 Hz pacing in HFSD & CTRL.
E. Diastolic mean cardiomyocyte segment stress-length relation (linear regression fit) for HFSD and CTRL at 2 and 4 Hz pacing.
F. Diastolic mean cardiomyocyte segment stress-length relation slope (mN/mm²) normalized relative to segment stretch (% length) yields cardiomyocyte mean segment stress/strain = stiffness
 [(mN/mm²) /strain (% length)] for HFSD and CTRL at 2 and 4 Hz pacing.

Comparison between two groups performed using unpaired t-test. With two independent variables a repeated measures 2-way ANOVA Sidaks post-hoc analysis was performed. Data are presented as mean ± standard deviation (C), mean ± standard error of the mean (E) or mean with paired individual points (B, D, F). For **B. & C.** (CTRL: N = 4 mice, n = 9 cells; HFSD: N = 4 mice, n = 8 cells). For **D. & E.** (CTRL: N = 4 mice, n = 7 cells; HFSD: N = 4 mice, n = 9 cells). For **F.** (CTRL: N = 4 mice, n = 8 cells; HFSD: N = 4 mice, n = 9 cells).

Table 1.	Echocardiographic parameters for CTRL and HFSD mice		
	CTRL	**HFSD**	**P-value**
Heart rate (bpm)	515±74	551±29	0.174
LV mass (mg)	125±27	131±21	0.314
LVIDd (mm)	3.60±0.17	4.16±0.47	0.004
IVSd (mm)	0.84±0.06	0.80±0.10	0.349
LVPWd (mm)	0.94±0.03	0.94±0.10	0.830
RWT (ratio)	0.49±0.06	0.43±0.09	0.079
Ejection Fraction (%)	67.8±3.7	64.9±3.2	0.093
Fractional Shortening (%)	31.8±3.6	30.6±2.1	0.400
E wave (mm/s)	472±74	569±65	0.003
A wave (mm/s)	348±99	400±116	0.190
e' wave (mm/s)	28.5±5.1	25.2±3.5	0.046
a' wave (mm/s)	23.5±7.8	23.5±6.4	0.995
E/A	1.51±0.41	1.47±0.37	0.095
e'/a'	1.33±0.44	1.12±0.39	0.171
E/e'	16.7±3.9	22.3±6.0	0.005
MV Deceleration Time (ms)	30.8±8.7	19.7±5.9	<0.001

Table 1. Echocardiographic parameters in CTRL and HFSD mice.
LVIDd left ventricular internal diameter at end diastole, *IVSd* interventricular septum diameter at end diastole, *LVPWd* left ventricular posterior wall thickness at end diastole, *RWT* relative wall thickness, *E wave* mitral valve blood flow velocity during the early ventricular filling phase, *A wave* mitral valve blood flow velocity during atrial contraction, *e' wave* mitral annulus tissue motion velocity during the early ventricular filling phase, *a' wave* mitral annulus tissue motion velocity during atrial contraction, *MV* mitral valve.

Comparisons between two treatment groups performed using Students' unpaired t-test (N = 9-15 mice per group). Data are presented as mean ± standard deviation.

Table 2. Stress-length relation mean values for CTRL and HFSD mouse cardiomyocytes

	CTRL 2 Hz	HFSD 2 Hz	p-value
Systolic Stress-SL Relation slope ((mN/mm^2) / µm)	6.54±4.97	16.35±15.09	0.087
Systolic Stress-Length Relation slope ((mN/mm^2) / %length)	0.013±0.011	0.022±0.011	0.022
Systolic : Diastolic Stress-SL relation 'Frank-Starling Gain Index' (ratio)	2.40±1.06	2.77±1.45	0.562
Systolic : Diastolic Stress-Length relation (ratio)	1.62±0.25	1.73±0.50	0.471

Table 2. Stress-length relation mean values for CTRL and HFSD mouse cardiomyocytes.
Comparisons between two treatment groups performed using Students' unpaired t-test (CTRL: N = 4-6 mice, n = 9-17 cells cells; HFSD: N = 2-6 mice, n = 7-19 cells). Data are presented as mean ± standard deviation.

Table 3. Ca^{2+} transient parameters for loaded CTRL and HFSD mouse cardiomyocytes

	CTRL 2 Hz	CTRL 4 Hz	HFSD 2 Hz	HFSD 4 Hz	Frequency p-value	Diet p-value	Interaction p-value
Systolic Ca^{2+} (F$_{340:380}$)	0.93±0.08	1.05±0.10#	0.99±0.15	1.08±0.20	p = 0.0004	p = 0.571	p = 0.100
Amplitude Ca^{2+} (F$_{340:380}$)	0.26±0.06	0.28±0.06	0.32±0.10	0.33±0.11	p = 0.569	p = 0.0672	p = 0.723
Time constant of Ca^{2+} decay (tau, sec)	0.21±0.07	0.09±0.02#	0.17±0.07	0.09±0.03#	p < 0.0001	p = 0.207	p = 0.350

Table 3. Ca^{2+} transient parameters for loaded CTRL and HFSD mouse cardiomyocytes.
Comparisons in data with two independent variable were performed using a mixed analysis 2-way ANOVA with Sidak's post-hoc analysis (CTRL, 2 Hz: N = 5 mice, n = 13 cells; CTRL, 4 Hz: N = 4 mice, n = 9 cells; HFSD, 2 Hz: N = 5 mice, n = 14 cells; HFSD, 4 Hz: N = 4 mice, n = 8 cells). Data are presented as mean ± standard deviation. P-values for ANOVA effects (Frequency, Diet, Interaction) are listed within the table. #<0.05: Significant pairwise post-hoc frequency differences vs CTRL 2 Hz or HFSD 2 Hz. No pairwise post-hoc treatment differences were observed.