Multi-tissue transition of A-to-I RNA editing pattern and its regulatory relevance in

transcription, splicing, and translation during development

Jia-Qi Pan,[a,b,c,1] Xu-Bin Pan,[a,b,c,d,1] Yan-Shan Liu,[a,b,c,e] Yun-Yun Jin,[a,b,c] Jian-Huan Chen[a,b,c,□]

[a]Laboratory of Genomic and Precision Medicine, Wuxi School of Medicine, Jiangnan

University, Wuxi 214000, Jiangsu, China

[b]Joint Primate Research Center for Chronic Diseases, Institute of Zoology of Guangdong

Academy of Science, Jiangnan University, Wuxi 214000, Jiangsu, China

[c]Jiangnan University Brain Institute, Wuxi 214000, Jiangsu, China

[d]Department of Ophthalmology, Affiliated Hospital of Jiangnan University, Wuxi 214000,

Jiangsu, China

[e]Department of Pediatric Laboratory, Affiliated Children's Hospital of Jiangnan University

(Wuxi Children's Hospital), Wuxi 214000, Jiangsu, China

[□]Corresponding author at: Laboratory of Genomic and Precision Medicine, Wuxi School of

Medicine, Jiangnan University, Wuxi, Jiangsu, China. cjh_bio@hotmail.com

[1]These authors contributed equally to this work.

Abstract

Previous studies have shown that A-to-I RNA editing can occur in various organs and tissues of normal physiological conditions. However, the dynamics of RNA editing and its functional relevance in multiple tissues and organs during the embryo-to-adult transition in mammals remains to be elucidated. Herein, we performed a comprehensive analysis of RNA sequencing and ribosome profiling of six mouse tissues at embryonic and adult stages, to elucidate the tissue- and stage-specific landscape of A-to-I RNA editing. Our result identified transcriptome-wide A-to-I RNA editing in six tissue types. Furthermore, differential expression was concurrently observed in a set of distinct differential RNA editing genes at both mRNA and protein levels across different tissues. Gene function and pathway enrichment analysis indicated that these genes with both differential editing and expression were involved in not only tissue-specific biological functions, but also common fundamental processes of post-transcriptional and post-translational modification. Further analysis showed a dynamic interaction between A-to-I RNA editing and alternative splicing in cell survival, death, signal transduction, and cell-cell interactions during development. Overall, our study demonstrates the potential role played by A-to-I editing during development, providing new insight into the effects of RNA editing within both transcriptional and translational landscape on it.

Keywords: A-to-I editing, transcription, translation, alternative splicing, development

1. Introduction

RNA editing is a post-transcriptional mechanism that alters the transcript sequence and can increase transcript and protein diversities [1]. Adenosine-to-inosine (A-to-I) RNA editing, catalyzed by the adenosine deaminases acting on RNA (*ADAR*s) family, is a canonical type of RNA editing in mammals [2, 3]. A-to-I RNA editing is involved in diverse biological processes, depending on the editing location. RNA editing profiles may vary among tissues and can also dynamically regulate critical biological processes, such as brain development, cancers, and metabolic disorders [4-7].

Mammals have almost identical genome sequences but exhibit great functional diversity in different tissues [8-11]. Correspondingly, A-to-I RNA editing in different tissues and developmental stages may also present complex landscapes and functions [12]. Therefore, the study of A-to-I RNA editing during mammalian development could not only reveal the dynamic regulation of RNA editing but also help unravel the complex mechanisms of this process. Previous studies have shown that A-to-I editing plays a crucial role in brain and neurological development, especially in the context of ion channels, neurotransmitter receptors, and synaptic plasticity, and contributes to the complexity and adaptability of the nervous system [4, 13]. Recent genome-wide characterization of RNA editing highlighted roles of high editing events of glutamatergic synapse during retinal development [14]. However, most of these studies have focused on the analysis of RNA editing in single tissues or organs, while further investigation is needed to compare RNA editing and their functional relevance to expression processes such as transcription, splicing, and translation across different tissues or organs.

In the current study, we performed comprehensive transcriptome and translatome analysis of six types of mouse tissues at both embryonic and adult stages. Our research findings suggested the substantial regulatory role of A-to-I RNA editing in transcription,

splicing and translation, providing new insight into the potential effects of RNA editing on

both transcriptional and translational landscape during development.

2. Materials and methods

2.1. Dataset download

Reads of RNA sequencing (RNA-seq) and ribosome profiling (Ribo-seq) from a dataset

(GSE94982) were downloaded from the NCBI Gene Expression Omnibus (GEO,

http://www.ncbi.nlm.nih.gov/geo) database. The dataset contains six types of tissue samples

from wild-type C57BL/6 mice at embryonic day (E) 15.5 and postnatal day (P) 42 (N = 2

each tissue group per time point), including ectoderm-derived brain and retina,

mesoderm-derived heart and kidney, as well as endoderm-derived liver and lung. The tissue

collection, RNA extraction, and library construction and sequencing were performed as

described in the original study [15].

2.2. Data processing

The raw sequencing reads of RNA-Seq and Ribo-seq were processed using FASTP (version

0.23.4) to remove adapters and low-quality reads [16]. The obtained Clean data were mapped to

the mouse reference genome (UCSC mm10) using RNA STAR (Version 2.7.0e) [17], generating

alignment files in BAM format. SAMtools (Version 1.9) was used to eliminate multiple

mappings and optical duplications of reads, retaining reads uniquely mapped to the mouse

genome [18]. GATK (Version 4.1.3) was used to recalibrate the base quality scores of the BAM

files by following the best practices suggested by the software [19].

2.3. Identification and annotation of A-to-I editing sites

VarScan (version 2.4.3) was utilized to call variants to identify single nucleotide variations

(SNVs) [20]. Given that inosine (I) is interpreted as guanosine (G) during reverse transcription

and translation, A-to-I RNA editing is also referred to as A-to-G RNA editing [21, 22]. Filtering

was performed using a standard bioinformatic pipeline described in our previous study [23, 24].

Annotation was carried out using the Ensembl Variation Effect Predictor (VEP) [25]. Briefly,

only SNVs that met the following criteria were kept: base quality $\geq$ 25, total sequencing

depth $\geq$ 10, alternative allele depth $\geq$ 2, and alternative allele frequency (AAF) $\geq$ 1%. SNVs

that met the following criteria were filtered and removed unless annotated in the REDIportal

V2.0 database [26, 27]: (1) located within the mitochondria DNA, simple repeat sequences, or

homopolymer runs $\geq$ 5 nucleotides (nt); (2) located within six nt of splice junctions or one nt

of insertions-deletions (INDELs); (3) annotated in the dbSNP database as known variant

types; (4) with an AAF equal to 100% or ranging between 40% and 60% in over 90% of

samples; (5) covered by high-quality reads < 10, or edited reads < 2, or editing levels < 1%.

2.4. Identification of differential RNA editing sites

Generalized Linear Models (GLM) and Likelihood Ratio Tests (LRT) were used to compare

RNA editing levels between embryonic and adult stages. Differential RNA editing (DRE)

sites were identified using the following criteria: (1) differential editing levels $\geq$ 1%; (2) false

discovery rate (FDR) < 0.05 [28].

2.5. Quantification of gene expression

FeatureCounts (Version 2.0.1) was utilized to extract pseudo-counts of gene expression from

BAM files [29]. Subsequently, EdgeR (Version 3.7) was used to calculate transcripts per

kilobase million (TPM) values for each gene, and differential gene expression analysis was

carried out by the Generalized Linear Model (GLM) [30]. *P*-values < 0.05 were determined to

be statistically significant. The correlation coefficients (*r*) between RNA editing level and

gene expression were calculated using Pearson correlation analysis.

2.6. Enrichment analysis of gene function and pathways

Enrichr (https://maayanlab.cloud/Enrichr/) [31] and DAVID online prediction tools

(https://david.ncifcrf.gov/tools.jsp) [32] was used to perform Gene Ontology (GO) biological

process enrichment analysis. Items with *P*-value < 0.05 were considered to be statistically

significant.

2.7. Prediction and mapping of RNA binding proteins binding sites

114 RBPmap website (http://rbpmap.technion.ac.il) was used to predict the binding patterns of

115 RNA binding proteins (RBPs) to differential A-to-I RNA editing sites [33]. The results of RBPs

116 binding prediction were visualized with WordCloud package (Version 2.6).

2.8. Analysis of differential alternative splicing

The replicate multivariate analysis of transcript splicing (rMATS) (Version 4.0.2) was used to

detect differential alternative splicing (DAS) events across different tissues [34]. The obtained

alternative splicing (AS) events were classified into five categories: skipped exon (SE),

retained intron (RI), alternative 5′ splice sites (A5SS), alternative 3′ splice site (A3SS) and

mutually exclusive exons (MXE). For each AS event, inclusion and exclusion junction reads

were averaged from replicates and used to calculate the Inclusion Level Difference (ΔPSI).

DAS events were identified using the following criteria: (1) $|\Delta\text{PSI}| > 10\%$; (2) P-values $<$

0.05.

3. Results

3.1. Tissue-specific A-to-I RNA Editing across mouse development

From the RNA-seq data of six tissues at E15.5 and P42, we identified a total of 618

high-confidence A-to-I RNA editing sites in 317 genes in brain, 501 high-confidence A-to-I

RNA editing sites in 265 genes in the heart, 503 high-confidence A-to-I RNA editing sites in

254 genes in the kidney, 479 high-confidence A-to-I RNA editing sites in 235 genes in the

liver, 660 high-confidence A-to-I RNA editing sites in 323 genes in the lung, 853

high-confidence A-to-I RNA editing sites in 404 genes in the retina **(Figure 1A;**

Supplementary Table S1). Analysis of the functional categories showed that the majority

(68.5-86.4%) of them were 3′-untranslated region (3′UTR) RNA editing. Missense RNA

editing constituted to only a small proportion (2.7 -7.3%) of the total variants **(Figure 1B,**

Supplementary Figure S1A). Sorting intolerant from tolerant (SIFT) analysis predicted 15.4

- 33.3% of the missense variants could potentially impact function of the encoded protein

(Figure 1C, Supplementary Figure S1B). Motif analysis showed that G was suppressed at

one base upstream but preferred at one base downstream of the editing sites **(Figure 1D)**,

consistent with the reported sequence preference of *ADAR* editing.

To investigate the dynamic regulation of A-to-I RNA editing, we compared the existence

of editing sites and edited genes between E15.5 and P42 in different tissues. As shown in

Figure 2A and Supplementary Figure S2, generally more editing sites and edited genes at

P42 than E15.5 were observed across different tissues, except for the heart.

3.2. Differential A-to-I RNA Editing between embryonic and adult stages

To systematically analyze RNA editing events during development, we then analyzed

differential RNA editing (DRE) between embryonic and adult stages. We identified 186, 110,

117, 112, 208, 294 DRE sites in these six tissues, respectively **(Figure 2B; Supplementary**

Table S2). The top 50 DRE sites in different tissues ranked by p-values are shown in **Figure**

3. In addition, we found that most of the DRE sites were up-regulated at P42 compared to

E15.5. Consistent with the overall RNA editing landscape, most of the DRE sites were also

found in 3UTR **(Figure 2C)**, and showed similar sequence motifs across all tissues

(Supplementary Figure S3A).

We then evaluated the expression of *Adar*s across different tissues at both stages. Our

findings revealed a dramatic increase in Adarb1 expression during the development of the

brain and retina at both transcriptional and translational levels **(Figure 2E)**. *Adar* was

significantly up-regulated in the brain at the translational level but not at the transcriptional

level and significantly up-regulated in the retina at the transcriptional level but not the

translational level **(Figure 2D)**. In addition, we observed a significant positive correlation

between the mean editing level and *Adarb1* mRNA expression both in the brain and retina

(*Pearson*'s $r = 0.99$, $P = 0.006$ and *Pearson*'s $r = 0.97$, $P = 0.029$, respectively) (**Figure 2F,**

G). We then identified *Adarb1*-correlated DRE sites by evaluating the correlation of their

editing levels with *Adarb1* mRNA expression levels in the brain and retina. 11

Adarb1-correlated DRE sites in 11 genes. 18 ones in 13 genes were identified in the brain and

retina, respectively, all of which sites were significantly up-regulated and positively

correlated with *Adarb1* expression (**Supplementary Figure S4, S5**). These observations

indicated that *Adarb1* may be the primary editing enzyme of A-to-I editing sites in the brain

and retina. However, we failed to observe a significant correlation between editing levels and

ADAR expression in the other four tissues (**Supplementary Figure S3B, C**).

3.3. Functional enrichment analysis of differential A-to-I RNA Editing genes

To explore the biological function of tissue-specific A-to-I RNA editing in development, we

performed enrichment analysis of DRE genes in each tissue, respectively. GO analysis

indicated that these DRE genes were not only enriched for certain tissue-specific biological

functions but also various routine biological processes related to development, such as

'positive regulation of growth hormone secretion', 'regulation of mitotic cell cycle',

'RNA-dependent DNA biosynthetic process' and 'phosphate-containing compound metabolic

process' (**Figure 2H; Supplementary Table S3**).

3.4. Potential effects of A-to-I RNA editing on transcription and translation

To explore the landscape function of A-to-I RNA editing at transcriptional and translational

levels, we compared the DRE genes and differentially expressed genes (DEGs) identified by

mRNA-Seq or Ribo-seq in different tissues (**Figure 4A; Supplementary Table S4**). Our

results revealed that 46, 18, 40, 37, 59, and 58 DRE genes were differentially expressed both

in mRNA-Seq and Ribo-Seq in the brain, heart, kidney, liver, lung, and retina, respectively

(**Figure 4B**). We then evaluated these DRE genes with differential expression for significant

functional enrichment in at least two tissue types. DRE genes with up-regulated expression

were enriched in routine biological processes of protein modification and metabolic

regulation, such as 'protein-containing complex localization', 'dephosphorylation',

'regulation of lipid metabolic process', and 'glycerolipid metabolic process', whereas DRE

genes with down-regulated expression were enriched in biological processes related to DNA

and RNA dynamics, such as 'DNA conformation change', 'DNA replication', 'mRNA export

from nucleus' and 'RNA localization' **(Figures 4C, D; Supplementary Table S5)**.

In addition, it was also noticed that 59, 40, 20, 14, 37, and 51 DRE genes were

differentially expressed at the transcription level only but not at the translation level in the

brain, heart, kidney, liver, lung, and retina, respectively **(Supplementary Figure S6A)**. We

then evaluated these DRE genes with differential RNA expression for significant functional

enrichment in at least two tissue types. Likewise, our enrichment analysis indicated that these

DRE genes, with transcriptional differential expression only, were enriched for biological

processes involved in the regulation of cellular processes and metabolic processes, such as

'cellular component disassembly', 'negative regulation of cell motility', 'thioester metabolic

process' and 'nucleoside bisphosphate metabolic process', while these DRE genes with

down-regulated expression were mainly enriched for biological processes involved in DNA

regulation and gene silencing, such as 'positive regulation of DNA repair', 'positive

regulation of DNA metabolic process', 'regulation of gene silencing' and 'positive regulation

of posttranscriptional gene silencing' **(Supplementary Figure S6B, C; Supplementary**

Table S4).

3.5. RNA binding prediction of differential A-to-I RNA editing sites

To gain insights into the potential functional impact of RNA editing, we utilized the RBPmap

database to predict the binding patterns of RBP overlapping with differential A-to-I RNA

editing sites across different tissues **(Supplementary Figure S7)**. **Figure 5** illustrates the top

10 RBPs with the highest binding frequencies to differential A-to-I RNA editing sites across

different tissues. Notably, RNA binding motif protein 45 (*RBM45*), Heterogeneous nuclear

ribonucleoprotein A0 (*HNRNPA0*), Serine and arginine-rich splicing factor 10 (*SRSF10*), Unk

Zinc Finger (*UNK*), CCR4-NOT transcription complex subunit 4 (*CNOT4*) were identified

across all six tissues. This suggests that these RBPs may play a crucial and ubiquitous role in

regulating RNA editing, indicating potential functional significance across gene expression

and cellular processes.

3.6. Interaction between A-to-I RNA editing and alternative splicing

To gain insights into the interplay between A-to-I RNA editing and alternative splicing, we

used rMATS to identify alternative splicing (AS) events and differential alternative splicing

(DAS) events in different tissues. We found that 39-54% of genes with A-to-I RNA editing

also exhibited alternative splicing in these six tissues, suggesting a potential interaction

between A-to-I RNA editing and alternative splicing **(Figure 6A)**. Then, we identified DAS

events and genes between embryonic and adult stages **(Figure 6B; Supplementary Table**

S6). Amongst the six tissues, the brain and retina exhibit the largest numbers of DAS events

and genes. Comparing DRE genes and DAS genes, we found that 4-40 DAS genes were also

differentially edited in these six tissues during development **(Figure 6C)**. Enrichment

analysis indicated that these genes were involved in various routine biological processes

related to development, such as 'negative regulation of programmed necrotic cell death',

'positive regulation of cytoplasmic transport', 'fibroblast growth factor receptor signaling

pathway', and 'neuron cell-cell adhesion.' This suggests that RNA editing and alternative

splicing were closely intertwined in cellular processes such as survival, death, signal

transduction, and cell-cell interactions during development **(Figure 6D)**.

4. Discussion

ADAR-mediated A-to-I RNA editing could be crucial for the mammalian development of

various organs and tissues [35]. However, the dynamic regulation of RNA editing during the

mammalian embryo-to-adult transition remains to be further elucidated. In the current study,

we performed a transcriptome-wide RNA editing analysis of six mouse tissues to study the

tissue- and stage-specific landscape of A-to-I RNA editing during development. Our analysis

revealed dramatic RNA editing changes in the development from embryonic into adult stages

across different tissues, most of which exhibited significantly increased editing levels and

were involved in potential biological functions related to development.

Among these DRE sites, 61 were found to be missense RNA editing. Notably,

significant up-regulation of two missense editing c.284A>G (p.K95R) and c.232A>G

(p.R78G) in the insulin-like growth factor binding protein 7 (*Igfbp7*) gene were observed in

the mouse brain, heart and kidney at the adult stage compared to the embryonic stage. These

two conserved missense sites were also found in other mammals, such as humans and pigs [36].

Igfbp7 encodes a member of the soluble proteins that bind insulin growth factor (IGF) and

suppress downstream signaling, thereby inhibiting protein synthesis, cell growth, and survival

[36]. Recent studies have observed that K95R and R78G editing in *IGFBP7* showed a high level

in the adult brain and cardiovascular tissues, and are reported to be up-regulated in

atherosclerosis and cardiomyopathies [37, 38]. R/G substitutions in the IGF binding domain may

disrupt the interaction between the protein and its IGF ligands [39]. In addition, the R78G and

K95R recordings have been reported to possibly modulate its susceptibility to proteolysis [40].

Our results also found another missense editing c.14A>G (p.Q5R) in the bladder

cancer-associated protein (*Blcap*) gene was observed in the mouse heart, kidney, and retina.

The editing level of this site decreased in the heart and kidney yet increased in the retina

during the development from the embryo to adult. *Blcap* encodes a tumor suppressor that

prevents tumorigenesis by promoting apoptosis and inhibiting cell proliferation. RNA

over-editing of *BLCAP* can activate the Akt/mTOR signaling pathway by regulating the

phosphorylation level, thus promoting the progression of hepatocellular carcinoma [41] and

cervical carcinogenesis [42]. In addition, missense editing in Neuroligin 2 (*Nlgn2*) c.406A>G

(p.N136D) was also observed to be up-regulated in the brain, lung, and retina in adult mice

compared to the embryo, which probably abolishes an N-glycosylation target site in Nlgn2 [43]

and thus might affect neurotransmission. *Nlgn2* encodes a transmembrane scaffolding protein

and promotes synapse development by forming trans-synaptic bridges spanning the synaptic

cleft [44]. Moreover, *Nlgn2* has been reported to be associated with neurological and psychiatric

disorders, such as schizophrenia [45], autism [46], and anxiety [47]. The functional relevance of

most of these missense DRE sites in development, especially those in *Igfbp7* and *Nlgn2,*

remains poorly understood.

Interestingly, our results showed that up-regulated RNA editing in a subset of long

intergenic non-coding RNA (lincRNA), such as small nucleolar RNA host gene 11 (*Snhg11*),

H19 imprinted maternally expressed transcript (*H19*), and maternally expressed 3 (*Meg3*)

during development. One intronic editing variant in *Snhg11* and two exonic editing variants

in *Meg3* were significantly up-regulated in the brain in adult mice. Seven intronic editing

variants in *Snhg11* and four exonic editing variants in *Meg3* were significantly up-regulated

in the retina in adult mice (**Supplementary Table S2**). *Snhg11* has been reported to promote

cell proliferation, migration, apoptosis, and autophagy during cancer development [48, 49].

Recent studies implicated that *Snhg11* also plays a neuroprotective role in neuronal injury [50].

Previous studies have suggested *Meg3* not only contributed to cancer development but is also

associated with the pathological stages of cancer and the prognosis of cancer patients [51].

Recent studies have shown that *Meg3* regulates AMPA receptor expression on the surface of

primary cortical neurons and plays a vital role in the nervous system [52]. The hyper editing in

Snhg11 and *Meg3* and differential RNA editing in both genes suggest a potential role of these

two lincRNA genes during brain and retina development.

In mammals, A-to-I editing is catalyzed by *ADAR*s, including *ADAR1*, *ADARB1* and

ADARB2 [53]. Our findings revealed that *Adarb1* could be primarily responsible for the

dynamics of A-to-I editing during the development of the brain and retina, compared to *Adar*

and *Adarb2*. However, the expression of *Adar*, *Adarb1*, and *Adarb2* have no significant

difference in the other four tissues. These results suggest that the *ADAR* family exhibits

tissue-specific mediation in A-to-I RNA editing, highlighting the vital role of *Adarb1* in

regulating A-to-I RNA editing activity during central nervous system (CNS) development.

Dynamic regulation of transcription and translation plays a key role in embryonic

development and maintenance of adult tissues [54, 55]. This regulation involves multiple

dimensions, with RNA modification standing out as a crucial facet in this intricate process [56,

57]. Previous studies have shown that A-to-I editing could play an important role in regulating

transcriptome diversity [58]. However, the impact of RNA editing on translation during

mammalian development remains largely unexplored. Ribo-seq can reveal the intricacies of

translation dynamics, enabling exploration of the specific regions of mRNA that are actively

undergoing translation by ribosomes [59]. Our results revealed certain differentially edited

genes with differential expression at both mRNA and protein levels, which were primarily

involved in both tissue-specific and routine biological processes related to development.

Previous studies have shown that A-to-I editing is crucial in neurological development [13].

Ectoderm-derived brain and retina are important components of the CNS [60]. Our results

consistently showed that DRE might contribute to brain and retina development by regulating

the expression of genes related to synaptic vesicles and neurotransmitters, highlighting its

crucial role in CNS development.

Overlapping of differentially edited sites and RBP binding sites provides a potential link

between RNA editing and RBP functions in development, such as the regulation of

pre-mRNA splicing and high-order RNA structures [61]. Our study identified a set of 5 core

RBPs with the highest binding frequencies to DRE sites across different tissues, including

RBM45, *HNRNPA0*, *SRSF10*, *UNK*, and *CNOT4*. *RBM45* encodes an RBP involved in neural development, and its pathological aggregation is closely related to neurodegenerative diseases, such as amyotrophic lateral sclerosis (ALS) and frontotemporal lobar dementia (FTLD) [62]. *SRSF10* encodes an atypical member of serine/arginine-rich (SR) proteins [63], and has been reported to influence glucose, fat, and cholesterol metabolism, embryonic heart development, and neural processes by controlling alternative splicing of specific transcripts [64]. *CNOT4* encodes an evolutionarily conserved E3 ubiquitin ligase [65]. *CNOT4* has been demonstrated to play synergistic biochemical functions during development and is required for post-implantation embryo development and meiosis progression during spermatogenesis [66]. Our study suggested the involvement of these RBPs in the downstream effects of differential RNA editing during development. Further experimental and functional studies will be needed to further understand the interplay between these RBPs and RNA editing during development.

Previous studies have shown that A-to-I editing of pre-mRNA before RNA splicing could be closely related to alternative pre-mRNA splicing [61, 67]. However, a global understanding of the interplay between RNA editing and alternative splicing during development is still missing. Our study systematically compared differential A-to-I RNA editing genes and differential alternative splicing genes in six tissues, revealing their closely intertwined in cell survival, death, signal transduction, and cell-cell interactions during development, providing a new insight into the relationship between RNA editing and alternative splicing.

In conclusion, our study suggested a transition of A-to-I RNA editing pattern across multiple tissues and its regulatory relevance in transcription, splicing, and translation during development. Further investigation will be needed to evaluate the role and mechanisms of RNA editing within both transcriptional and translational landscapes.

Funding

This study was supported in part by grants from the National Natural Science Foundation of China (no. 31671311), the "Six Talent Peak" Plan of Jiangsu Province (no. SWYY-127), the Program for High-Level Entrepreneurial and Innovative Talents of Jiangsu Province, Natural Science Foundation of Guangdong Province/Guangdong Basic and Applied Basic Research Foundation (2019A1515012062), Taihu Lake Talent Plan, and Fundamental Research Funds for the Central Universities (JUSRP51712B and JUSRP1901XNC).

CRediT authorship contribution statement

Jia-Qi Pan: Conceptualization, Data curation, Formal analysis, Writing - original draft. **Xu-Bin Pan:** Conceptualization, Methodology, Software. **Yan-Shan Liu**: Methodology, Software, Supervision. **Yun-Yun Jin:** Methodology, Project administration, Supervision. **Jian-Huan Chen:** Conceptualization, Supervision, Funding acquisition, Writing – review & editing.

Declaration of Competing Interest

All authors declare that they have no known financial or personal relationships that might influence the publication of this paper.

References

1. Tan MH, Li Q, Shanmugam R, et al. Dynamic landscape and regulation of RNA editing in mammals. *Nature* 2017;550:249-254.

2. Bass BL. RNA editing by adenosine deaminases that act on RNA. *Annu Rev Biochem* 2002;71:817-846.

3. Nishikura K. Functions and regulation of RNA editing by ADAR deaminases. *Annu Rev*

Biochem 2010;79:321-349.

4. Hwang T, Park CK, Leung AK, et al. Dynamic regulation of RNA editing in human brain

development and disease. *Nat Neurosci* 2016;19:1093-1099.

5. Kung CP, Maggi LB, Jr., Weber JD. The Role of RNA Editing in Cancer Development

and Metabolic Disorders. *Front Endocrinol (Lausanne)* 2018;9:762.

6. Xu X, Wang Y, Liang H. The role of A-to-I RNA editing in cancer development. *Curr*

Opin Genet Dev 2018;48:51-56.

7. Buchumenski I, Holler K, Appelbaum L, Eisenberg E, Junker JP, Levanon EY.

Systematic identification of A-to-I RNA editing in zebrafish development and adult organs.

Nucleic Acids Res 2021;49:4325-4337.

8. Yanai I, Korbel JO, Boue S, McWeeney SK, Bork P, Lercher MJ. Similar gene

expression profiles do not imply similar tissue functions. *Trends Genet* 2006;22:132-138.

9. Ong CT, Corces VG. Enhancer function: new insights into the regulation of

tissue-specific gene expression. *Nat Rev Genet* 2011;12:283-293.

10. Battle A, Brown CD, Engelhardt BE, Montgomery SB. Genetic effects on gene

expression across human tissues. *Nature* 2017;550:204-213.

11. Sonawane AR, Platig J, Fagny M, et al. Understanding Tissue-Specific Gene Regulation.

Cell Rep 2017;21:1077-1088.

12. Yao L, Wang H, Song Y, et al. Large-scale prediction of ADAR-mediated effective

human A-to-I RNA editing. *Brief Bioinform* 2019;20:102-109.

13. Yang Y, Okada S, Sakurai M. Adenosine-to-inosine RNA editing in neurological

development and disease. *RNA Biol* 2021;18:999-1013.

14. Li C, Shi X, Yang J, et al. Genome-wide characterization of RNA editing highlights roles

of high editing events of glutamatergic synapse during mouse retinal development. *Comput*

Struct Biotechnol J 2022;20:2648-2656.

15. Wang H, Wang Y, Yang J, et al. Tissue- and stage-specific landscape of the mouse

translatome. *Nucleic Acids Res* 2021;49:6165-6180.

16. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics 2018;34:i884-i890.

17. Dobin A, Gingeras TR. Mapping RNA-seq Reads with STAR. *Curr Protoc*

Bioinformatics 2015;51:11 14 11-11 14 19.

18. Li H, Handsaker B, Wysoker A, et al. The Sequence Alignment/Map format and

SAMtools. *Bioinformatics* 2009;25:2078-2079.

19. Van der Auwera GA, Carneiro MO, Hartl C, et al. From FastQ data to high confidence

variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr Protoc*

Bioinformatics 2013;43:11 10 11-11 10 33.

20. Koboldt DC, Zhang Q, Larson DE, et al. VarScan 2: somatic mutation and copy number

alteration discovery in cancer by exome sequencing. *Genome Res* 2012;22:568-576.

21. Goldstein B, Agranat-Tamir L, Light D, Ben-Naim Zgayer O, Fishman A, Lamm AT.

A-to-I RNA editing promotes developmental stage-specific gene and lncRNA expression.

Genome Res 2017;27:462-470.

22. Xiang JF, Yang Q, Liu CX, Wu M, Chen LL, Yang L. N(6)-Methyladenosines Modulate

A-to-I RNA Editing. *Mol Cell* 2018;69:126-135 e126.

23. Tao J, Ren CY, Wei ZY, Zhang F, Xu J, Chen JH. Transcriptome-Wide Identification of

G-to-A RNA Editing in Chronic Social Defeat Stress Mouse Models. *Front Genet*

2021;12:680548.

24. Zhang JQ, Pan JQ, Wei ZY, et al. Brain Epitranscriptomic Analysis Revealed Altered

A-to-I RNA Editing in Septic Patients. *Front Genet* 2022;13:887001.

25. McLaren W, Gil L, Hunt SE, et al. The Ensembl Variant Effect Predictor. *Genome Biol*

2016;17:122.

26. Picardi E, D'Erchia AM, Lo Giudice C, Pesole G. REDIportal: a comprehensive database of A-to-I RNA editing events in humans. *Nucleic Acids Res* 2017;45:D750-D757.

27. Mansi L, Tangaro MA, Lo Giudice C, et al. REDIportal: millions of novel A-to-I RNA editing events from thousands of RNAseq experiments. *Nucleic Acids Res* 2021;49:D1012-D1019.

28. Srivastava PK, Bagnati M, Delahaye-Duriez A, et al. Genome-wide analysis of differential RNA editing in epilepsy. *Genome Res* 2017;27:440-450.

29. Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 2014;30:923-930.

30. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2010;26:139-140.

31. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res* 2016;44:W90-97.

32. Huang da W, Sherman BT, Lempicki RA. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* 2009;37:1-13.

33. Paz I, Kosti I, Ares M, Jr., Cline M, Mandel-Gutfreund Y. RBPmap: a web server for mapping binding sites of RNA-binding proteins. *Nucleic Acids Res* 2014;42:W361-367.

34. Shen S, Park JW, Lu ZX, et al. rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. *Proc Natl Acad Sci U S A* 2014;111:E5593-5601.

35. Slotkin W, Nishikura K. Adenosine-to-inosine RNA editing and human disease. *Genome Med* 2013;5:105.

36. Levitsky LI, Ivanov MV, Goncharov AO, et al. Massive Proteogenomic Reanalysis of Publicly Available Proteomic Datasets of Human Tissues in Search for Protein Recoding via Adenosine-to-Inosine RNA Editing. *J Proteome Res* 2023;22:1695-1711.

37. Huang J, Lin L, Dong Z, et al. A porcine brain-wide RNA editing landscape. *Commun*

Biol 2021;4:717.

38. Mann TD, Kopel E, Eisenberg E, Levanon EY. Increased A-to-I RNA editing in

atherosclerosis and cardiomyopathies. *PLoS Comput Biol* 2023;19:e1010923.

39. Levanon EY, Hallegger M, Kinar Y, et al. Evolutionarily conserved human targets of

adenosine to inosine RNA editing. *Nucleic Acids Res* 2005;33:1162-1168.

40. Godfried Sie C, Hesler S, Maas S, Kuchka M. IGFBP7's susceptibility to proteolysis is

altered by A-to-I RNA editing of its transcript. *FEBS Lett* 2012;586:2313-2317.

41. Hu X, Wan S, Ou Y, et al. RNA over-editing of BLCAP contributes to

hepatocarcinogenesis identified by whole-genome and transcriptome sequencing. *Cancer Lett*

2015;357:510-519.

42. Chen W, He W, Cai H, et al. A-to-I RNA editing of BLCAP lost the inhibition to STAT3

activation in cervical cancer. *Oncotarget* 2017;8:39417-39429.

43. Scott H, Panin VM. The role of protein N-glycosylation in neural transmission.

Glycobiology 2014;24:407-417.

44. Ali H, Marth L, Krueger-Burg D. Neuroligin-2 as a central organizer of inhibitory

synapses in health and disease. *Sci Signal* 2020;13.

45. Sun C, Cheng MC, Qin R, et al. Identification and functional characterization of rare

mutations of the neuroligin-2 gene (NLGN2) associated with schizophrenia. *Hum Mol Genet*

2011;20:3042-3051.

46. Parente DJ, Garriga C, Baskin B, et al. Neuroligin 2 nonsense variant associated with

anxiety, autism, intellectual disability, hyperphagia, and obesity. *Am J Med Genet A*

2017;173:213-216.

47. Babaev O, Botta P, Meyer E, et al. Neuroligin 2 deletion alters inhibitory synapse

function and anxiety-associated neuronal activation in the amygdala. *Neuropharmacology*

2016;100:56-65.

48. Liu S, Yang N, Wang L, Wei B, Chen J, Gao Y. lncRNA SNHG11 promotes lung cancer

cell proliferation and migration via activation of Wnt/β-catenin signaling pathway. *J Cell*

Physiol 2020;235:7541-7553.

49. Huang W, Huang F, Lei Z, Luo H. LncRNA SNHG11 Promotes Proliferation, Migration,

Apoptosis, and Autophagy by Regulating hsa-miR-184/AGO2 in HCC. *Onco Targets Ther*

2020;13:413-421.

50. Chen Y, Fan Z, Wu Q. Dexmedetomidine improves oxygen-glucose

deprivation/reoxygenation (OGD/R) -induced neurological injury through regulating

SNHG11/miR-324-3p/VEGFA axis. *Bioengineered* 2021;12:4794-4804.

51. He Y, Luo Y, Liang B, Ye L, Lu G, He W. Potential applications of MEG3 in cancer

diagnosis and prognosis. *Oncotarget* 2017;8:73282-73295.

52. Tan MC, Widagdo J, Chau YQ, et al. The Activity-Induced Long Non-Coding RNA

Meg3 Modulates AMPA Receptor Surface Expression in Primary Cortical Neurons. *Front*

Cell Neurosci 2017;11:124.

53. Fritzell K, Xu LD, Lagergren J, Öhman M. ADARs and editing: The role of A-to-I RNA

modification in cancer progression. *Semin Cell Dev Biol* 2018;79:123-130.

54. Matkovich SJ, Edwards JR, Grossenheider TC, de Guzman Strong C, Dorn GW, 2nd.

Epigenetic coordination of embryonic heart transcription by dynamically regulated long

noncoding RNAs. *Proc Natl Acad Sci U S A* 2014;111:12264-12269.

55. Falcon KT, Watt KEN, Dash S, et al. Dynamic regulation and requirement for ribosomal

RNA transcription during mammalian development. *Proc Natl Acad Sci U S A*

2022;119:e2116974119.

56. Roundtree IA, Evans ME, Pan T, He C. Dynamic RNA Modifications in Gene

Expression Regulation. *Cell* 2017;169:1187-1200.

57. Cui L, Ma R, Cai J, et al. RNA modifications: importance in immune cell biology and

related diseases. *Signal Transduct Target Ther* 2022;7:334.

58. Bahn JH, Lee JH, Li G, Greer C, Peng G, Xiao X. Accurate identification of A-to-I RNA

editing in human by transcriptome sequencing. *Genome Res* 2012;22:142-150.

59. Ingolia NT, Hussmann JA, Weissman JS. Ribosome Profiling: Global Views of

Translation. *Cold Spring Harb Perspect Biol* 2019;11.

60. Fernando M, Lee S, Wark JR, et al. Differentiation of brain and retinal organoids from

confluent cultures of pluripotent stem cells connected by nerve-like axonal projections of

optic origin. *Stem Cell Reports* 2022;17:1476-1492.

61. Hu X, Zou Q, Yao L, Yang X. Survey of the binding preferences of RNA-binding

proteins to RNA editing events. *Genome Biol* 2022;23:169.

62. Chen X, Yang Z, Wang W, et al. Structural basis for RNA recognition by the N-terminal

tandem RRM domains of human RBM45. *Nucleic Acids Res* 2021;49:2946-2958.

63. Zhou X, Wu W, Li H, et al. Transcriptome analysis of alternative splicing events

regulated by SRSF10 reveals position-dependent splicing modulation. *Nucleic Acids Res*

2014;42:4019-4030.

64. Shkreta L, Delannoy A, Salvetti A, Chabot B. SRSF10: an atypical splicing regulator

with critical roles in stress response, organ development, and viral replication. *Rna*

2021;27:1302-1317.

65. Chalabi Hagkarim N, Grand RJ. The Regulatory Properties of the Ccr4-Not Complex.

Cells 2020;9.

66. Dai XX, Jiang Y, Gu JH, et al. The CNOT4 Subunit of the CCR4-NOT Complex is

Involved in mRNA Degradation, Efficient DNA Damage Repair, and XY Chromosome

Crossover during Male Germ Cell Meiosis. *Adv Sci (Weinh)* 2021;8:2003636.

67. Hsiao YE, Bahn JH, Yang Y, et al. RNA editing in nascent RNA affects pre-mRNA

splicing. *Genome Res* 2018;28:812-823.

Figure Legends

Figure. 1. Transcriptome-wide analysis of A-to-I RNA editing in different tissues from embryonic (E15.5) and adult (P42) mice. (A) Circos plot of gene expression and A-to-I RNA editing sites in all six tissues. The outer circle denotes the mean expression level of genes, and the inner circle denotes the A-to-I RNA editing sites. **(B)** Functional categories of A-to-I RNA editing variants in all six tissues. **(C)** SIFT prediction of missense A-to-I RNA editing variants. **(D)** Motif context around A-to-I RNA editing sites.

Figure. 2. Differential A-to-I RNA editing in different tissues between embryonic (E15.5)

and adult (P42) mice. (A) Venn plots of genes and sites with A-to-I RNA editing at

embryonic and adult stages. **(B)** Bar plot showing the number of DRE sites between

embryonic and adult stages. **(C)** Functional categories of DRE sites. **(D)** Normalized

expression of *Adar* at the transcriptional (left) and translational level (right). **(E)** Normalized

expression of *Adarb1* at the transcriptional (left) and translational level (right). **: $P < 0.01$;

*: $P < 0.05$. *Pearson*'s correlation analysis between editing levels and normalized expression

of *Adars* at the transcriptional level in the brain **(F)** and the retina **(G)**. **(H)** Heatmap showing

terms from GO items enriched by differential edited genes in ≥ three tissues.

Figure. 3. A-to-I DRE sites in different tissues during development. Heatmap showing the

top 50 editing levels of DRE sites in different tissues during development.

Figure. 4. Possible functional impact of A-to-I DRE on transcription and translation during development. (A) Venn plots comparing genes with DRE and DEGS identified by mRNA-Seq (DEG_mRNA) or Ribo-seq (DEG_Ribo) across different tissues. **(B)** Bar plot showing the number of DRE genes differentially expressed both in mRNA-Seq and Ribo-Seq. Heatmap showing the enriched GO terms for DRE genes with up-regulated **(C)** and down-regulated expression **(D)** in at least two tissue types.

Figure. 5. The top 10 frequent RNA binding proteins (RBPs) ranked by their binding frequencies to DRE sites in different tissues.

Figure. 6. Developmental interrelation between A-to-I RNA editing and alternative splicing. (A) Proportion and number of edited genes with and without alternative splicing (AS) in different tissues. **(B)** Bar plot comparing the number of differential alternative splicing (DAS) events between embryonic and adult stages. **(C)** Venn plots of DAS genes and DRE genes in different tissues. **(D)** Heatmap showing the enriched GO terms for DAS genes with differential editing in at least two tissue types. RI, retained intron; MXE, mutually exclusive exons; A3SS, alternative 3′ splice sites; A5SS, alternative 5′ splice sites; SE, skipped exon.