Multi-tissue transition of A-to-I RNA editing pattern and its regulatory relevance in

transcription, splicing, and translation during development

Jia-Qi Pan, [1,2,3,4,#] Xu-Bin Pan, [1,2,3,4,5,#] Yan-Shan Liu,[1,2,3,4,6] Yun-Yun Jin,[1,2,3,4] Jian-Huan Chen[1,2,3,4,*]

[1]Laboratory of Genomic and Precision Medicine, Wuxi School of Medicine, Jiangnan University, Wuxi 214000, Jiangsu, China

[2]Joint Primate Research Center for Chronic Diseases, Institute of Zoology of Guangdong Academy of Science, Jiangnan University, Wuxi 214000, Jiangsu, China

[3]Jiangnan University Brain Institute, Wuxi 214000, Jiangsu, China

[4]MOE Medical Basic Research Innovation Center for Gut Microbiota and Chronic Diseases, Wuxi School of Medicine, Jiangnan University, Wuxi 214000, Jiangsu, China

[5]Department of Ophthalmology, Affiliated Hospital of Jiangnan University, Wuxi 214000, Jiangsu, China

[6]Department of Pediatric Laboratory, Affiliated Children's Hospital of Jiangnan University (Wuxi Children's Hospital), Wuxi 214000, Jiangsu, China

[□]Corresponding author:

Prof. Jian-Huan Chen

Email: cjh_bio@hotmail.com

Laboratory of Genomic and Precision Medicine, Wuxi School of Medicine, Jiangnan University, Wuxi, Jiangsu, China.

[#]These authors contributed equally to this work.

Abstract

Previous studies have shown that A-to-I RNA editing can occur in various organs and tissues under normal physiological conditions. However, the dynamics of RNA editing and its functional relevance in multiple tissues and organs during the embryo-to-adult transition in mammals remain to be elucidated. Here, we performed a comprehensive analysis of RNA-Seq and Ribo-Seq data from six mouse tissues at embryonic and adult stages to elucidate the A-to-I RNA editing transition during development and to validate it in additional mouse brain datasets spanning multiple developmental stages. Our results revealed a general transition of up-regulated A-to-I RNA editing activity across numerous tissue types during embryonic-adult development, indicated by significantly increased average RNA editing levels. Consistently, differential RNA editing (DRE) analysis showed more up-regulated than down-regulated RNA editing sites in all six tissue types. Furthermore, such RNA editing transitions during development could contribute to differential gene expression (DEG) at both transcriptional and translational levels, as well as differential alternative splicing (DAS) across multiple tissues. Differentially edited genes with DEG or DAS could be involved not only in tissue-specific biological functions but also in common routine biological processes during development. Notably, *Adarb1* was found to be a more important player than *Adar* in the CNS (the brain and retina), with its expression levels increasing gradually and correlating with the A-to-I RNA editing activity. Our study demonstrates the potential role of A-to-I editing during development across multiple tissues, providing new insights into its regulatory relevance in both transcriptional and translational landscapes.

Keywords: A-to-I editing, transcription, translation, alternative splicing, development

Introduction

RNA editing is a post-transcriptional mechanism that alters the transcript sequence and can increase transcript and protein diversities (Tan et al. 2017b). Adenosine-to-inosine (A-to-I) RNA editing, catalyzed by the adenosine deaminases acting on RNA (ADARs) family, is a canonical type of RNA editing in mammals (Bass 2002; Nishikura 2010). A-to-I RNA editing is involved in diverse biological processes, depending on the editing location. RNA editing profiles may vary among tissues and can also dynamically regulate critical biological processes, such as brain development, cancers, and metabolic disorders (Hwang et al. 2016; Kung et al. 2018; Xu et al. 2018; Buchumenski et al. 2021).

Mammals have almost identical genome sequences but exhibit substantial functional diversity across tissues (Yanai et al. 2006; Ong and Corces 2011; Battle et al. 2017; Sonawane et al. 2017). Correspondingly, A-to-I RNA editing in different tissues and developmental stages may also present complex landscapes and functions (Yao et al. 2019). Therefore, the study of A-to-I RNA editing during mammalian development could not only reveal the dynamic regulation of RNA editing but also help unravel the complex mechanisms of this process. Previous studies have shown that A-to-I editing plays a crucial role in brain and neurological development, particularly in ion channels, neurotransmitter receptors, and synaptic plasticity, and contributes to the complexity and adaptability of the nervous system (Hwang et al. 2016; Yang et al. 2021). Recent genome-wide characterization of RNA editing highlighted roles of editing events of glutamatergic synapse during retinal development (Li et al. 2022). However, most of these studies have focused on RNA editing in a single tissue or organ (Yang et al. 2024), and further investigation is needed to compare RNA editing and its functional relevance across different tissues or organs, including transcription, splicing, and translation.

In the current study, we performed comprehensive transcriptome and translatome

analyses of six mouse tissues at both embryonic and adult stages. Our research findings

suggest a substantial regulatory role for A-to-I RNA editing in transcription, splicing, and

translation, providing new insights into its potential effects on both transcriptional and

translational landscapes during development.

Results

Tissue-specific A-to-I RNA Editing across mouse development

From the RNA-seq data of six tissues at E15.5 and P42, we identified a total of 618

high-confidence A-to-I RNA editing sites in 317 genes in brain, 501 high-confidence A-to-I

RNA editing sites in 265 genes in the heart, 503 high-confidence A-to-I RNA editing sites in

254 genes in the kidney, 479 high-confidence A-to-I RNA editing sites in 235 genes in the

liver, 660 high-confidence A-to-I RNA editing sites in 323 genes in the lung, 853

high-confidence A-to-I RNA editing sites in 404 genes in the retina **(Figure 1A;**

Supplementary Table S1). Analysis of the functional categories showed that the majority

(68.5-86.4%) of them were 3′-untranslated region (3′UTR) RNA editing. Missense RNA

editing accounted for only a small proportion (2.7 -7.3%) of the total variants **(Figure 1B,**

Supplementary Figure S1A). Sorting intolerant from tolerant (SIFT) analysis predicted that

15.4-33.3% of missense variants could potentially impact the function of the encoded protein

(Figure 1C, Supplementary Figure S1B). Motif analysis showed that G was suppressed at

one base upstream but preferred at one base downstream of the editing sites **(Figure 1D)**,

consistent with the reported sequence preference of *ADAR* editing.

To investigate the dynamic regulation of A-to-I RNA editing, we compared the existence

of editing sites and edited genes between E15.5 and P42 in different tissues. As shown in

Figure 2A and Supplementary Figure S2, more editing sites and edited genes were

observed at P42 than at E15.5 across different tissues, except in the heart.

Differential A-to-I RNA Editing between embryonic and adult stages

To systematically analyze RNA editing events during development, we then analyzed

differential RNA editing (DRE) between embryonic and adult stages. We identified 186, 110,

117, 112, 208, 294 DRE sites in these six tissues, respectively **(Figure 2B; Supplementary**

Table S2). The top 50 DRE sites across different tissues, ranked by p-values, are shown in

Figure 3. In addition, we found that most DRE sites were up-regulated at P42 compared with

E15.5. Consistent with the overall RNA editing landscape, most of the DRE sites were also

found in 3UTR **(Figure 2C)** and showed similar sequence motifs across all tissues

(Supplementary Figure S3A).

We then evaluated the expression of Adars across different tissues at both stages. Our

findings revealed a dramatic increase in Adarb1 expression during brain and retinal

development at both transcriptional and translational levels **(Figure 2E)**. *Adar* was

significantly up-regulated in the brain at the translational level but not at the transcriptional

level, and in the retina at the transcriptional level but not at the translational level **(Figure**

2D). In addition, we observed a significant positive correlation between the mean editing

level and *Adarb1* mRNA expression both in the brain and retina (*Pearson*'s $r = 0.99$, $P =$

0.006, and *Pearson*'s $r = 0.97$, $P = 0.029$, respectively) **(Figure 2F, G)**. We then identified

Adarb1-correlated DRE sites by evaluating the correlation of their editing levels with *Adarb1*

mRNA expression levels in the brain and retina. 11 *Adarb1*-correlated DRE sites in 11 genes.

18 ones in 13 genes were identified in the brain and retina, respectively, all of which sites

were significantly up-regulated and positively correlated with *Adarb1* expression

(Supplementary Figure S4, S5). These observations indicated that *Adarb1* may be the

primary editing enzyme of A-to-I editing sites in the brain and retina. However, we failed to

observe a significant correlation between editing levels and *ADAR* expression in the other

four tissues **(Supplementary Figure S3B and C)**.

Functional enrichment analysis of differentially edited genes

To explore the biological function of tissue-specific A-to-I RNA editing in development, we

performed enrichment analysis of DRE genes in each tissue. GO analysis indicated that these

DRE genes were not only enriched for certain tissue-specific biological functions but also

various routine biological processes related to development, such as 'positive regulation of

growth hormone secretion', 'regulation of mitotic cell cycle', 'RNA-dependent DNA

biosynthetic process', and 'phosphate-containing compound metabolic process' **(Figure 2H;**

Supplementary Table S3).

Potential effects of A-to-I RNA editing on transcription and translation

To explore the landscape function of A-to-I RNA editing at transcriptional and translational

levels, we compared the DRE genes and differentially expressed genes (DEGs) identified by

mRNA-Seq or Ribo-seq in different tissues **(Figure 4A; Supplementary Table S4)**. Our

results revealed that 46, 18, 40, 37, 59, and 58 DRE genes were differentially expressed both

in mRNA-Seq and Ribo-Seq in the brain, heart, kidney, liver, lung, and retina, respectively

(Figure 4B). We then evaluated these DRE genes with differential expression for significant

functional enrichment in at least two tissue types. DRE genes with up-regulated expression

were enriched in routine biological processes of protein modification and metabolic

regulation, such as 'protein-containing complex localization', 'dephosphorylation',

'regulation of lipid metabolic process', and 'glycerolipid metabolic process'. In contrast,

DRE genes with down-regulated expression were enriched in biological processes related to

DNA and RNA dynamics, such as 'DNA conformation change', 'DNA replication', 'mRNA

export from nucleus', and 'RNA localization' **(Figures 4C, D; Supplementary Table S5)**.

In addition, it was also noticed that 59, 40, 20, 14, 37, and 51 DRE genes were

differentially expressed at the transcription level only but not at the translation level in the

brain, heart, kidney, liver, lung, and retina, respectively **(Supplementary Figure S6A)**. We

then evaluated these DRE genes with differential RNA expression for significant functional

enrichment in at least two tissue types. Likewise, our enrichment analysis indicated that these

DRE genes, with transcriptional differential expression only, were enriched for biological

processes involved in the regulation of cellular processes and metabolic processes, such as

'cellular component disassembly', 'negative regulation of cell motility', 'thioester metabolic

process', and 'nucleoside bisphosphate metabolic process'. In contrast, these DRE genes with

down-regulated expression were mainly enriched for biological processes involved in DNA

regulation and gene silencing, such as 'positive regulation of DNA repair', 'positive

regulation of DNA metabolic process', 'regulation of gene silencing', and 'positive regulation

of posttranscriptional gene silencing' **(Supplementary Figure S6B, C; Supplementary**

Table S4).

RNA binding prediction of differential A-to-I RNA editing sites

To gain insights into the potential functional impact of RNA editing, we utilized the RBPmap

database to predict the binding patterns of RBP overlapping with differential A-to-I RNA

editing sites across different tissues **(Supplementary Figure S7)**. **Figure 5** illustrates the top

10 RBPs with the highest binding frequencies to differential A-to-I RNA editing sites across

various tissues. Notably, RNA binding motif protein 45 (*RBM45*), Heterogeneous nuclear

ribonucleoprotein A0 (*HNRNPA0*), Serine and arginine-rich splicing factor 10 (*SRSF10*), Unk

Zinc Finger (*UNK*), and CCR4-NOT transcription complex subunit 4 (*CNOT4*) were

identified across all six tissues. This suggests that these RBPs may play a crucial and

ubiquitous role in regulating RNA editing, indicating potential functional significance across

gene expression and cellular processes.

Interaction between A-to-I RNA editing and alternative splicing

To gain insights into the interplay between A-to-I RNA editing and alternative splicing, we

used rMATS to identify alternative splicing (AS) and differential alternative splicing (DAS)

events across tissues. We found that 39-54% of genes with A-to-I RNA editing also exhibited

alternative splicing in these six tissues, suggesting a potential interaction between A-to-I

RNA editing and alternative splicing **(Figure 6A)**. Then, we identified DAS events and genes

that differed between the embryonic and adult stages **(Figure 6B; Supplementary Table S6)**.

Among the six tissues, the brain and retina exhibit the highest numbers of DAS events and

genes. Comparing DRE genes and DAS genes, we found that 4-40 DAS genes were also

differentially edited in these six tissues during development **(Figure 6C)**. Enrichment

analysis indicated that these genes were involved in various routine biological processes

related to development, such as 'negative regulation of programmed necrotic cell death',

'positive regulation of cytoplasmic transport', 'fibroblast growth factor receptor signaling

pathway', and 'neuron cell-cell adhesion.' This suggests that RNA editing and alternative

splicing were closely intertwined in cellular processes such as survival, death, signal

transduction, and cell-cell interactions during development **(Figure 6D)**.

Validation of A-to-I RNA editing pattern during brain development

To validate the common characteristics of A-to-I RNA editing pattern during development,

we conducted a comparative analysis using additional RNA-Seq and Ribo-Seq datasets

(GSE157425 and GSE169457) of brain neocortex at embryonic (E12.5–E17) and postnatal

(P0) stages from an independent study. Using the same analysis procedure, 557 to 723 A-to-I

RNA editing sites were identified across different developmental stages, with a significant

increase in the number of editing sites over time. Consistent with the findings in the brain

RNA editing data described above, RNA editing was also enriched in 3UTR **(Figure 7A)**. A

large proportion (283, 30.01%) of the editing sites were shared across all stages, with a

considerable number also shared by adjacent time points, whereas only a small proportion

(9.23%, 87) were exclusively found in a single stage **(Figure 7B)**. Analysis of editing levels

between different stages revealed an increasing trend of the average editing level during brain

development **(Figure 7C)**. To systematically explore the temporal dynamics of RNA editing,

DRE across these time points was analyzed. Our results showed that 453 sites underwent

differential editing during brain development **(Figure 7D and Supplementary Figure S8A)**.

The top 50 DRE sites over time, ranked by p-values, are shown in **Figure 7E**. Most DRE

sites were up-regulated during development, with the most rapid up-regulation occurring after

birth. By comparing the discovery and validation datasets, 52 differentially edited genes and

42 DRE sites were shared between the two **(Supplementary Figure S8B, C)**. Gene function

enrichment analysis indicated that the DRE in the validation dataset was mainly involved in

chromatin and gene expression regulation, such as 'negative regulation of organelle

organization', 'covalent chromatin modification', and 'regulation of chromosome

organization' **(Supplementary Figure S8D)**.

Comparison of *Adarb1* and Adarb2 expression among different stages showed a

dramatic increase in *Adarb1* and *Adarb2* expression during brain development at both

transcriptional and translational levels **(Figure 7F)**. In addition, we observed significant

positive correlations of *Adarb1* and *Adarb2* expressions with the number of RNA editing sites

and edited genes, as well as with the average editing level (all *Pearson r* range from 0.96 to

0.98 and all $P < 0.001$) **(Figure 7G, H)**.

Then the potential effects of A-to-I RNA editing on transcription and translation were

further validated during brain development. We observed that 134 differentially edited genes

were differentially expressed both in RNA-Seq and Ribo-Seq over time **(Figure 8A)**. Gene

enrichment analysis indicated that differentially edited genes with up-regulated expression

were primarily involved in functions related to synapses and neurotransmitters, such as

'regulation of postsynaptic membrane potential', 'receptor diffusion trapping', and 'synaptic

vesicle cycle'. In contrast, those with down-regulated expression were mainly involved in

biological processes related to RNA dynamics, such as 'positive regulation of mRNA

processing' and 'regulation of transcription of nucleolar large rRNA by RNA polymerase I'

(Figure 8B). Comparing splicing events between adjacent time points, we revealed

1622~2741 DAS events over time **(Figure 8D)**. We then found that 85 genes with both DAS

and differentially edited during brain development **(Figure 8C),** which were mainly enriched

in functions related to synaptic regulation and brain development, such as 'regulation of

neuronal synaptic plasticity', 'excitatory postsynaptic potential', and 'forebrain development'

(Figure 8E).

Discussion

ADAR-mediated A-to-I RNA editing could be crucial for the mammalian development of

various organs and tissues (Slotkin and Nishikura 2013). Yet, its regulatory relevance remains

to be further elucidated. In the current study, our analysis revealed a dramatic RNA editing

transition characterized by increased RNA editing activity across different tissues during

embryonic-to-adult development, with potential biological impacts on RNA splicing,

transcription, and translation.

Differential missense RNA editing was observed in multiple tissues during the

development from E15.5 to P42. Notably, significant up-regulation of two missense editing

c.284A>G (p.K95R) and c.232A>G (p.R78G) in the insulin-like growth factor binding

protein 7 (*Igfbp7*) gene was observed in the mouse brain, heart, and kidney at the adult stage

(P42) compared to the embryonic (E15.5) stage. These two conserved missense editing sites

were also found in other mammals, such as humans and pigs (Levitsky et al. 2023). *Igfbp7*

encodes a soluble protein that binds insulin growth factor (IGF) and suppresses downstream

signaling, thereby inhibiting protein synthesis, cell growth, and survival (Levitsky et al. 2023).

Recent studies have reported that K95R and R78G editing in *IGFBP7* are elevated in the

adult brain and cardiovascular tissues, as well as in atherosclerosis and cardiomyopathies

(Huang et al. 2021; Mann et al. 2023). R/G substitutions in the IGF binding domain may

disrupt the interaction between the protein and its IGF ligands (Levanon et al. 2005). In

addition, the R78G and K95R recordings have been reported to possibly modulate its

susceptibility to proteolysis (Godfried Sie et al. 2012). Our results also showed that another

differential missense editing c.14A>G (p.Q5R) in the bladder cancer-associated protein

(*Blcap*) gene was observed in the mouse heart, kidney, and retina. The editing level of this

site decreased in the heart and kidney, yet increased in the retina, during development from

embryo to adult. *Blcap* encodes a tumor suppressor that prevents tumorigenesis by promoting

apoptosis and inhibiting cell proliferation. RNA over-editing of *BLCAP* can activate the

Akt/mTOR signaling pathway by regulating phosphorylation levels, thereby promoting

hepatocellular carcinoma progression (Hu et al. 2015) and cervical carcinogenesis (Chen et al.

2017). In addition, missense editing in Neuroligin 2 (*Nlgn2*) c.406A>G (p.N136D) was

up-regulated in the brain, lung, and retina during the embryo-to-adult development, which

probably abolishes an N-glycosylation target site in *Nlgn2* (Scott and Panin 2014) and thus

might affect neurotransmission. *Nlgn2* encodes a transmembrane scaffolding protein and

promotes synapse development by forming trans-synaptic bridges spanning the synaptic cleft

(Ali et al. 2020). Moreover, *Nlgn2* has been reported to be associated with neurological and

psychiatric disorders, such as schizophrenia (Sun et al. 2011), autism (Parente et al. 2017),

and anxiety (Babaev et al. 2016). The functional relevance of most of these missense DRE

sites in development, especially those in *Igfbp7* and *Nlgn2,* remains poorly understood. Our

findings thus warranted functional investigations in future studies.

Interestingly, our results showed that RNA editing was up-regulated in a subset of long

intergenic non-coding RNAs (lincRNAs), such as small nucleolar RNA host gene 11

(*Snhg11*), H19 imprinted maternally expressed transcript (*H19*), and maternally expressed

gene 3 (*Meg3*), during development. One intronic editing variant in *Snhg11* and two exonic

editing variants in *Meg3* were significantly up-regulated in the brain in adult mice. Seven

intronic editing variants in *Snhg11* and four exonic editing variants in *Meg3* were

significantly up-regulated in the retina in adult mice (**Supplementary Table S2**). *Snhg11* has

been reported to promote cell proliferation, migration, apoptosis, and autophagy during

cancer development (Liu et al. 2020; Huang et al. 2020). Recent studies implicated that

Snhg11 also plays a neuroprotective role in neuronal injury (Chen et al. 2021b). Previous

studies have suggested *that Meg3* not only contributes to cancer development but is also

associated with the pathological stages of cancer and the prognosis of cancer patients (He et

al. 2017). Recent studies have shown that *Meg3* regulates surface AMPA receptor expression

in primary cortical neurons and plays a vital role in the nervous system (Tan et al. 2017a).

The hyper-editing in *Snhg11* and *Meg3,* and the differential RNA editing in both genes,

suggest a potential role for these two lincRNA genes during brain and retina development.

In mammals, A-to-I editing is catalyzed by *ADAR*s, including *ADAR1*, *ADARB1,* and

ADARB2 (Fritzell et al. 2018). Our findings revealed that *Adarb1* was primarily responsible

for the dynamics of A-to-I editing during brain and retinal development, compared to *Adar*

and *Adarb2*. In particular, *Adarb1* increased gradually and showed a strong correlation with

the up-regulated A-to-I RNA editing activity during development, which was further

validated in multi-stage brain development datasets. In contrast, the expression of *Adar*,

Adarb1, and *Adarb2* did not differ significantly across the other four tissues. These results

suggest a tissue-specific role for the *ADAR* family in mediating A-to-I RNA editing,

highlighting the vital role of *Adarb1* in regulating A-to-I RNA editing activity during CNS

development.

Dynamic regulation of transcription and translation plays a key role in embryonic

development and maintenance of adult tissues (Matkovich et al. 2014; Falcon et al. 2022).

This regulation involves multiple dimensions, with RNA modification standing out as a

crucial facet in this intricate process (Roundtree et al. 2017; Cui et al. 2022). Previous studies

have shown that A-to-I editing could play an essential role in regulating transcriptome

diversity (Bahn et al. 2012). However, the impact of RNA editing on translation during

mammalian development remains largely unexplored. Ribo-seq can reveal the intricacies of

translation dynamics, enabling exploration of the specific regions of mRNA that are actively

undergoing translation by ribosomes (Ingolia et al. 2019). Our results revealed differentially

edited genes with differential expression at both mRNA and protein levels, primarily

involved in both tissue-specific and routine biological processes related to development.

Previous studies have shown that A-to-I editing is crucial in neurological development (Yang

et al. 2021). Ectoderm-derived brain and retina are essential components of the CNS

(Fernando et al. 2022). Our results consistently showed that DRE may contribute to brain and

retinal development by regulating the expression of genes involved in synaptic vesicle and

neurotransmitter function, highlighting its crucial role in CNS development.

Previous studies have shown that A-to-I editing of pre-mRNA before RNA splicing

could be closely related to alternative pre-mRNA splicing (Hu et al. 2022; Hsiao et al. 2018).

However, a global understanding of the interplay between RNA editing and alternative

splicing during development remains to be further elucidated. Our study systematically

compared differential A-to-I RNA editing genes and differential alternative splicing genes

across six tissues, revealing their close intertwining in cell survival, death, signal transduction,

and cell-cell interactions during development, providing new insights into the relationship

between RNA editing and alternative splicing.

The overlap between differentially edited RNA sites and RBP binding sites provides a

potential link between RNA editing and RBP functions in development, such as the

regulation of pre-mRNA splicing and higher-order RNA structures (Hu et al. 2022). Our

study identified five core RBPs with the highest binding frequencies to DRE sites across

different tissues, including *RBM45*, *HNRNPA0*, *SRSF10*, *UNK*, and *CNOT4*. *RBM45* encodes

an RBP involved in neural development, and its pathological aggregation is closely related to

neurodegenerative diseases, such as amyotrophic lateral sclerosis (ALS) and frontotemporal

lobar dementia (FTLD) (Chen et al. 2021a). *SRSF10* encodes an atypical member of

serine/arginine-rich (SR) proteins (Zhou et al. 2014), and has been reported to influence

glucose, fat, and cholesterol metabolism, embryonic heart development, and neural processes

by controlling alternative splicing of specific transcripts (Shkreta et al. 2021). *CNOT4*

encodes an evolutionarily conserved E3 ubiquitin ligase (Chalabi Hagkarim and Grand 2020).

CNOT4 has been demonstrated to play synergistic biochemical functions during development

and is required for post-implantation embryo development and meiosis progression during

spermatogenesis (Dai et al. 2021). Our study suggested that these RBPs are involved in the

downstream effects of differential RNA editing during development. Further experimental

and functional studies will be needed to further understand the interplay between these RBPs

and RNA editing during development.

In conclusion, our study suggested a transition in the A-to-I RNA editing pattern across

multiple tissues and its regulatory relevance for transcription, splicing, and translation during

development. Further investigation will be needed to evaluate the role and mechanisms of

RNA editing within both transcriptional and translational landscapes.

Materials and methods

Dataset download

Reads of RNA sequencing (RNA-seq) and ribosome profiling (Ribo-seq) from a dataset

(GSE94982) were downloaded from the NCBI Gene Expression Omnibus (GEO,

http://www.ncbi.nlm.nih.gov/geo) database. The dataset contains six types of tissue samples

from wild-type C57BL/6 mice at embryonic day (E) 15.5 and postnatal day (P) 42 (N = 2

each tissue group per time point), including ectoderm-derived brain and retina,

mesoderm-derived heart and kidney, as well as endoderm-derived liver and lung (Wang et al.

2021). The validation datasets (GSE157425 and GSE169457) were downloaded from the

NCBI Gene Expression Omnibus (GEO, http://www.ncbi.nlm.nih.gov/geo). The dataset

contains neocortex tissue from wild-type CD-1 mice at embryonic (E12.5–E17) and early

postnatal (P0) period (N = 2 per time point) (Harnett et al. 2022). The tissue collection, RNA

extraction, and library construction and sequencing were performed as described in the

original studies.

Data processing

The raw RNA-Seq and Ribo-Seq sequencing reads were processed using FASTP (version

0.23.4) to remove adapters and low-quality reads (Chen et al. 2018). The obtained Clean data

were mapped to the mouse reference genome (UCSC mm10) using RNA STAR (Version

2.7.0e) (Dobin and Gingeras 2015), generating BAM alignment files. SAMtools (Version 1.9)

was used to eliminate multiple mappings and optical duplications of reads, retaining reads

uniquely mapped to the mouse genome (Li et al. 2009). GATK (Version 4.1.3) was used to

recalibrate the base quality scores of the BAM files by following the best practices suggested

by the software (Van der Auwera et al. 2013).

Identification and annotation of A-to-I editing sites

VarScan (version 2.4.3) was utilized to call variants to identify single-nucleotide variations

(SNVs) (Koboldt et al. 2012). Given that inosine (I) is interpreted as guanosine (G) during

reverse transcription and translation, A-to-I RNA editing is also referred to as A-to-G RNA

editing (Goldstein et al. 2017; Xiang et al. 2018). Filtering was performed using a standard

bioinformatic pipeline described in our previous study (Tao et al. 2021; Zhang et al. 2022).

Annotation was carried out using the Ensembl Variation Effect Predictor (VEP) (McLaren et

al. 2016). Briefly, only SNVs that met the following criteria were kept: base quality ≥ 25,

total sequencing depth ≥ 10, alternative allele depth ≥ 2, and alternative allele frequency

(AAF) $\geq 1\%$. SNVs that met the following criteria were filtered and removed unless

annotated in the REDIportal V2.0 database (Picardi et al. 2017; Mansi et al. 2021): (1)

located within the mitochondria DNA, simple repeat sequences, or homopolymer runs ≥ 5

nucleotides (nt); (2) located within six nt of splice junctions or one nt of insertions-deletions

(INDELs); (3) annotated in the dbSNP database as known variant types; (4) with an AAF

equal to 100% or ranging between 40% and 60% in over 90% of samples; (5) covered by

high-quality reads < 10, or edited reads < 2, or editing levels < 1%.

Identification of differential RNA editing sites

Generalized Linear Models (GLM) and Likelihood Ratio Tests (LRT) were used to compare

RNA editing levels between embryonic and adult stages. Differential RNA editing (DRE)

sites were identified using the following criteria: (1) differential editing levels ≥ 1%; (2) false

discovery rate (FDR) < 0.05 (Srivastava et al. 2017).

Quantification of gene expression

FeatureCounts (Version 2.0.1) was utilized to extract pseudo-counts of gene expression from

BAM files (Liao et al. 2014). Subsequently, EdgeR (Version 3.7) was used to calculate

transcripts per kilobase million (TPM) values for each gene, and differential gene expression

analysis was performed using the Generalized Linear Model (GLM) (Robinson et al. 2010).

P-values < 0.05 were determined to be statistically significant. The correlation coefficients (*r*)

between RNA editing level and gene expression were calculated using Pearson correlation

analysis.

Enrichment analysis of gene function and pathways

Enrichr (https://maayanlab.cloud/Enrichr/) (Kuleshov et al. 2016) and DAVID online

prediction tools (https://david.ncifcrf.gov/tools.jsp) (Huang da et al. 2009) were used to

perform Gene Ontology (GO) biological process enrichment analysis. Items with a *P*-value <

0.05 were considered to be statistically significant.

Prediction and mapping of RNA-binding protein binding sites

The RBPmap website (http://rbpmap.technion.ac.il) was used to predict the binding patterns

of RNA-binding proteins (RBPs) to differential A-to-I RNA editing sites (Paz et al. 2014).

The results of the RBPs binding prediction were visualized with the WordCloud package

(Version 2.6).

Analysis of differential alternative splicing

The replicate multivariate analysis of transcript splicing (rMATS) (Version 4.0.2) was used to

detect differential alternative splicing (DAS) events across different tissues (Shen et al. 2014).

The obtained alternative splicing (AS) events were classified into five categories: skipped

exon (SE), retained intron (RI), alternative 5′ splice sites (A5SS), alternative 3′ splice sites

(A3SS), and mutually exclusive exons (MXE). For each AS event, junction reads from

inclusion and exclusion groups were averaged across replicates and used to calculate the

Inclusion Level Difference (ΔPSI). DAS events were identified using the following criteria:

(1) $|\Delta PSI| > 10\%$; (2) P-values < 0.05.

Author contributions

410 **Jia-Qi Pan:** Conceptualization, Data curation, Formal analysis, Writing - original draft.

411 **Xu-Bin Pan:** Conceptualization, Methodology, Software. **Yan-Shan Liu**: Methodology,

412 Software, Supervision. **Yun-Yun Jin:** Methodology, Project administration, Supervision.

413 **Jian-Huan Chen:** Conceptualization, Supervision, Funding acquisition, Writing – review &

414 editing.

Funding

This study was supported in part by grants from the National Natural Science Foundation of

China (no. 31671311), the "Six Talent Peak" Plan of Jiangsu Province (no. SWYY-127), the

Program for High-Level Entrepreneurial and Innovative Talents of Jiangsu Province, Natural

Science Foundation of Guangdong Province/Guangdong Basic and Applied Basic Research

Declarations

All authors declare that they have no known financial or personal relationships that might

influence the publication of this paper.

References

Ali H, Marth L, Krueger-Burg D. 2020. Neuroligin-2 as a central organizer of inhibitory
synapses in health and disease. *Sci Signal* **13**.

Babaev O, Botta P, Meyer E, Müller C, Ehrenreich H, Brose N, Lüthi A, Krueger-Burg D. 2016.
Neuroligin 2 deletion alters inhibitory synapse function and anxiety-associated
neuronal activation in the amygdala. *Neuropharmacology* **100**: 56–65.

Bahn JH, Lee JH, Li G, Greer C, Peng G, Xiao X. 2012. Accurate identification of A-to-I RNA
editing in human by transcriptome sequencing. *Genome Res* **22**: 142–50.

Bass BL. 2002. RNA editing by adenosine deaminases that act on RNA. *Annu Rev Biochem* **71**:
817–46.

Battle A, Brown CD, Engelhardt BE, Montgomery SB. 2017. Genetic effects on gene
expression across human tissues. *Nature* **550**: 204–213.

Buchumenski I, Holler K, Appelbaum L, Eisenberg E, Junker JP, Levanon EY. 2021.
Systematic identification of A-to-I RNA editing in zebrafish development and adult
organs. *Nucleic Acids Res* **49**: 4325–4337.

Chalabi Hagkarim N, Grand RJ. 2020. The Regulatory Properties of the Ccr4-Not Complex.
Cells **9**.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**: i884–i890.

Chen W, He W, Cai H, Hu B, Zheng C, Ke X, Xie L, Zheng Z, Wu X, Wang H. 2017. A-to-I
RNA editing of BLCAP lost the inhibition to STAT3 activation in cervical cancer.
Oncotarget **8**: 39417–39429.

Chen X, Yang Z, Wang W, Qian K, Liu M, Wang J, Wang M. 2021a. Structural basis for RNA
recognition by the N-terminal tandem RRM domains of human RBM45. *Nucleic Acids
Res* **49**: 2946–2958.

Chen Y, Fan Z, Wu Q. 2021b. Dexmedetomidine improves oxygen-glucose
deprivation/reoxygenation (OGD/R) -induced neurological injury through regulating
SNHG11/miR-324-3p/VEGFA axis. *Bioengineered* **12**: 4794–4804.

Cui L, Ma R, Cai J, Guo C, Chen Z, Yao L, Wang Y, Fan R, Wang X, Shi Y. 2022. RNA
modifications: importance in immune cell biology and related diseases. *Signal
Transduct Target Ther* **7**: 334.

Dai XX, Jiang Y, Gu JH, Jiang ZY, Wu YW, Yu C, Yin H, Zhang J, Shi QH, Shen L, et al. 2021.
The CNOT4 Subunit of the CCR4-NOT Complex is Involved in mRNA Degradation,
Efficient DNA Damage Repair, and XY Chromosome Crossover during Male Germ
Cell Meiosis. *Adv Sci (Weinh)* **8**: 2003636.

Dobin A, Gingeras TR. 2015. Mapping RNA-seq Reads with STAR. *Curr Protoc
Bioinformatics* **51**: 11 14 1-11 14 19.

Falcon KT, Watt KEN, Dash S, Zhao R, Sakai D, Moore EL, Fitriasari S, Childers M, Sardiu
ME, Swanson S, et al. 2022. Dynamic regulation and requirement for ribosomal RNA
transcription during mammalian development. *Proc Natl Acad Sci U S A* **119**:
e2116974119.

Fernando M, Lee S, Wark JR, Xiao D, Lim BY, O'Hara-Wright M, Kim HJ, Smith GC, Wong T, Teber ET, et al. 2022. Differentiation of brain and retinal organoids from confluent cultures of pluripotent stem cells connected by nerve-like axonal projections of optic origin. *Stem Cell Reports* **17**: 1476–1492.

Fritzell K, Xu LD, Lagergren J, Öhman M. 2018. ADARs and editing: The role of A-to-I RNA modification in cancer progression. *Semin Cell Dev Biol* **79**: 123–130.

Godfried Sie C, Hesler S, Maas S, Kuchka M. 2012. IGFBP7's susceptibility to proteolysis is altered by A-to-I RNA editing of its transcript. *FEBS Lett* **586**: 2313–7.

Goldstein B, Agranat-Tamir L, Light D, Ben-Naim Zgayer O, Fishman A, Lamm AT. 2017. A-to-I RNA editing promotes developmental stage-specific gene and lncRNA expression. *Genome Res* **27**: 462–470.

Harnett D, Ambrozkiewicz MC, Zinnall U, Rusanova A, Borisova E, Drescher AN, Couce-Iglesias M, Villamil G, Dannenberg R, Imami K, et al. 2022. A critical period of translational control during brain development at codon resolution. *Nat Struct Mol Biol* **29**: 1277–1290.

He Y, Luo Y, Liang B, Ye L, Lu G, He W. 2017. Potential applications of MEG3 in cancer diagnosis and prognosis. *Oncotarget* **8**: 73282–73295.

Hsiao YE, Bahn JH, Yang Y, Lin X, Tran S, Yang EW, Quinones-Valdez G, Xiao X. 2018. RNA editing in nascent RNA affects pre-mRNA splicing. *Genome Res* **28**: 812–823.

Hu X, Wan S, Ou Y, Zhou B, Zhu J, Yi X, Guan Y, Jia W, Liu X, Wang Q, et al. 2015. RNA over-editing of BLCAP contributes to hepatocarcinogenesis identified by whole-genome and transcriptome sequencing. *Cancer Lett* **357**: 510–9.

Hu X, Zou Q, Yao L, Yang X. 2022. Survey of the binding preferences of RNA-binding proteins to RNA editing events. *Genome Biol* **23**: 169.

Huang da W, Sherman BT, Lempicki RA. 2009. Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**: 1–13.

Huang J, Lin L, Dong Z, Yang L, Zheng T, Gu W, Zhang Y, Yin T, Sjöstedt E, Mulder J, et al. 2021. A porcine brain-wide RNA editing landscape. *Commun Biol* **4**: 717.

Huang W, Huang F, Lei Z, Luo H. 2020. LncRNA SNHG11 Promotes Proliferation, Migration, Apoptosis, and Autophagy by Regulating hsa-miR-184/AGO2 in HCC. *Onco Targets Ther* **13**: 413–421.

Hwang T, Park CK, Leung AK, Gao Y, Hyde TM, Kleinman JE, Rajpurohit A, Tao R, Shin JH, Weinberger DR. 2016. Dynamic regulation of RNA editing in human brain development and disease. *Nat Neurosci* **19**: 1093–9.

Ingolia NT, Hussmann JA, Weissman JS. 2019. Ribosome Profiling: Global Views of Translation. *Cold Spring Harb Perspect Biol* **11**.

Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, Miller CA, Mardis ER, Ding L, Wilson RK. 2012. VarScan 2: somatic mutation and copy number alteration discovery in cancer by exome sequencing. *Genome Res* **22**: 568–76.

Kuleshov MV, Jones MR, Rouillard AD, Fernandez NF, Duan Q, Wang Z, Koplev S, Jenkins SL, Jagodnik KM, Lachmann A, et al. 2016. Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res* **44**: W90-7.

Kung CP, Maggi LB, Weber JD. 2018. The Role of RNA Editing in Cancer Development and
Metabolic Disorders. *Front Endocrinol (Lausanne)* **9**: 762.

Levanon EY, Hallegger M, Kinar Y, Shemesh R, Djinovic-Carugo K, Rechavi G, Jantsch MF,
Eisenberg E. 2005. Evolutionarily conserved human targets of adenosine to inosine
RNA editing. *Nucleic Acids Res* **33**: 1162–8.

Levitsky LI, Ivanov MV, Goncharov AO, Kliuchnikova AA, Bubis JA, Lobas AA, Solovyeva
EM, Pyatnitskiy MA, Ovchinnikov RK, Kukharsky MS, et al. 2023. Massive
Proteogenomic Reanalysis of Publicly Available Proteomic Datasets of Human Tissues
in Search for Protein Recoding via Adenosine-to-Inosine RNA Editing. *J Proteome Res*
22: 1695–1711.

Li C, Shi X, Yang J, Li K, Dai L, Zhang Y, Zhou M, Su J. 2022. Genome-wide characterization
of RNA editing highlights roles of high editing events of glutamatergic synapse during
mouse retinal development. *Comput Struct Biotechnol J* **20**: 2648–2656.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and
SAMtools. *Bioinformatics* **25**: 2078–9.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**: 923–30.

Liu S, Yang N, Wang L, Wei B, Chen J, Gao Y. 2020. lncRNA SNHG11 promotes lung cancer
cell proliferation and migration via activation of Wnt/β-catenin signaling pathway. *J*
Cell Physiol **235**: 7541–7553.

Mann TD, Kopel E, Eisenberg E, Levanon EY. 2023. Increased A-to-I RNA editing in
atherosclerosis and cardiomyopathies. *PLoS Comput Biol* **19**: e1010923.

Mansi L, Tangaro MA, Lo Giudice C, Flati T, Kopel E, Schaffer AA, Castrignano T, Chillemi G,
Pesole G, Picardi E. 2021. REDIportal: millions of novel A-to-I RNA editing events
from thousands of RNAseq experiments. *Nucleic Acids Res* **49**: D1012–D1019.

Matkovich SJ, Edwards JR, Grossenheider TC, de Guzman Strong C, Dorn GW. 2014.
Epigenetic coordination of embryonic heart transcription by dynamically regulated
long noncoding RNAs. *Proc Natl Acad Sci U S A* **111**: 12264–9.

McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, Flicek P, Cunningham F. 2016.
The Ensembl Variant Effect Predictor. *Genome Biol* **17**: 122.

Nishikura K. 2010. Functions and regulation of RNA editing by ADAR deaminases. *Annu Rev*
Biochem **79**: 321–49.

Ong CT, Corces VG. 2011. Enhancer function: new insights into the regulation of
tissue-specific gene expression. *Nat Rev Genet* **12**: 283–93.

Parente DJ, Garriga C, Baskin B, Douglas G, Cho MT, Araujo GC, Shinawi M. 2017.
Neuroligin 2 nonsense variant associated with anxiety, autism, intellectual disability,
hyperphagia, and obesity. *Am J Med Genet A* **173**: 213–216.

Paz I, Kosti I, Ares M, Cline M, Mandel-Gutfreund Y. 2014. RBPmap: a web server for
mapping binding sites of RNA-binding proteins. *Nucleic Acids Res* **42**: W361-7.

Picardi E, D'Erchia AM, Lo Giudice C, Pesole G. 2017. REDIportal: a comprehensive
database of A-to-I RNA editing events in humans. *Nucleic Acids Res* **45**: D750–D757.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**:
139–40.

Roundtree IA, Evans ME, Pan T, He C. 2017. Dynamic RNA Modifications in Gene
Expression Regulation. *Cell* **169**: 1187–1200.

Scott H, Panin VM. 2014. The role of protein N-glycosylation in neural transmission.
Glycobiology **24**: 407–17.

Shen S, Park JW, Lu ZX, Lin L, Henry MD, Wu YN, Zhou Q, Xing Y. 2014. rMATS: robust
and flexible detection of differential alternative splicing from replicate RNA-Seq data.
Proc Natl Acad Sci U S A **111**: E5593-601.

Shkreta L, Delannoy A, Salvetti A, Chabot B. 2021. SRSF10: an atypical splicing regulator
with critical roles in stress response, organ development, and viral replication. *Rna* **27**:
1302–1317.

Slotkin W, Nishikura K. 2013. Adenosine-to-inosine RNA editing and human disease. *Genome*
Med **5**: 105.

Sonawane AR, Platig J, Fagny M, Chen CY, Paulson JN, Lopes-Ramos CM, DeMeo DL,
Quackenbush J, Glass K, Kuijjer ML. 2017. Understanding Tissue-Specific Gene
Regulation. *Cell Rep* **21**: 1077–1088.

Srivastava PK, Bagnati M, Delahaye-Duriez A, Ko JH, Rotival M, Langley SR, Shkura K,
Mazzuferi M, Danis B, van Eyll J, et al. 2017. Genome-wide analysis of differential
RNA editing in epilepsy. *Genome Res* **27**: 440–450.

Sun C, Cheng MC, Qin R, Liao DL, Chen TT, Koong FJ, Chen G, Chen CH. 2011.
Identification and functional characterization of rare mutations of the neuroligin-2 gene
(NLGN2) associated with schizophrenia. *Hum Mol Genet* **20**: 3042–51.

Tan MC, Widagdo J, Chau YQ, Zhu T, Wong JJ, Cheung A, Anggono V. 2017a. The
Activity-Induced Long Non-Coding RNA Meg3 Modulates AMPA Receptor Surface
Expression in Primary Cortical Neurons. *Front Cell Neurosci* **11**: 124.

Tan MH, Li Q, Shanmugam R, Piskol R, Kohler J, Young AN, Liu KI, Zhang R, Ramaswami G,
Ariyoshi K, et al. 2017b. Dynamic landscape and regulation of RNA editing in
mammals. *Nature* **550**: 249–254.

Tao J, Ren CY, Wei ZY, Zhang F, Xu J, Chen JH. 2021. Transcriptome-Wide Identification of
G-to-A RNA Editing in Chronic Social Defeat Stress Mouse Models. *Front Genet* **12**:
680548.

Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A,
Jordan T, Shakir K, Roazen D, Thibault J, et al. 2013. From FastQ data to high
confidence variant calls: the Genome Analysis Toolkit best practices pipeline. *Curr*
Protoc Bioinformatics **43**: 11 10 1-11 10 33.

Wang H, Wang Y, Yang J, Zhao Q, Tang N, Chen C, Li H, Cheng C, Xie M, Yang Y, et al. 2021.
Tissue- and stage-specific landscape of the mouse translatome. *Nucleic Acids Res* **49**:
6165–6180.

Xiang JF, Yang Q, Liu CX, Wu M, Chen LL, Yang L. 2018. N(6)-Methyladenosines Modulate
A-to-I RNA Editing. *Mol Cell* **69**: 126-135 e6.

Xu X, Wang Y, Liang H. 2018. The role of A-to-I RNA editing in cancer development. *Curr*

Opin Genet Dev **48**: 51–56.

Yanai I, Korbel JO, Boue S, McWeeney SK, Bork P, Lercher MJ. 2006. Similar gene
expression profiles do not imply similar tissue functions. *Trends Genet* **22**: 132–8.

Yang L, Yi L, Yang J, Zhang R, Xie Z, Wang H. 2024. Temporal landscape and translational
regulation of a-to-I RNA editing in mouse retina development. *BMC Biol* **22**: 106.

Yang Y, Okada S, Sakurai M. 2021. Adenosine-to-inosine RNA editing in neurological
development and disease. *RNA Biol* **18**: 999–1013.

Yao L, Wang H, Song Y, Dai Z, Yu H, Yin M, Wang D, Yang X, Wang J, Wang T, et al. 2019.
Large-scale prediction of ADAR-mediated effective human A-to-I RNA editing. *Brief*
Bioinform **20**: 102–109.

Zhang JQ, Pan JQ, Wei ZY, Ren CY, Ru FX, Xia SY, He YS, Lin K, Chen JH. 2022. Brain
Epitranscriptomic Analysis Revealed Altered A-to-I RNA Editing in Septic Patients.
Front Genet **13**: 887001.

Zhou X, Wu W, Li H, Cheng Y, Wei N, Zong J, Feng X, Xie Z, Chen D, Manley JL, et al. 2014.
Transcriptome analysis of alternative splicing events regulated by SRSF10 reveals
position-dependent splicing modulation. *Nucleic Acids Res* **42**: 4019–30.

Figure 1. Transcriptome-wide analysis of A-to-I RNA editing in different tissues from embryonic (E15.5) and adult (P42) mice. (A) Circos plot of gene expression and A-to-I RNA editing sites in all six tissues. The outer circle denotes the mean expression level of genes, and the inner circle denotes the A-to-I RNA editing sites. **(B)** Functional categories of A-to-I RNA editing variants in all six tissues. **(C)** SIFT prediction of missense A-to-I RNA editing variants. **(D)** Motif context around A-to-I RNA editing sites.

Figure 2. Differential A-to-I RNA editing in different tissues between embryonic (E15.5)
and adult (P42) mice. (A) Venn plots of genes and sites with A-to-I RNA editing at
embryonic and adult stages. **(B)** Bar plot showing the number of DRE sites between
embryonic and adult stages. **(C)** Functional categories of DRE sites. **(D)** Normalized
expression of *Adar* at the transcriptional (left) and translational level (right). **(E)** Normalized
expression of *Adarb1* at the transcriptional (left) and translational level (right). **:** $P < 0.01$;
*: $P < 0.05$. *Pearson*'s correlation analysis between editing levels and normalized expression
of *Adars* at the transcriptional level in the brain **(F)** and the retina **(G)**. **(H)** Heatmap showing
terms from GO items enriched by differential edited genes in ≥ three tissues.

Figure 3. A-to-I DRE sites in different tissues during development. Heatmap showing the top 50 editing levels of DRE sites in different tissues during development.

Figure 4. Possible functional impact of A-to-I DRE on transcription and translation during development. (A) Venn plots comparing genes with DRE and DEGs identified by mRNA-seq (DEG_mRNA) or Ribo-seq (DEG_Ribo) across different tissues. **(B)** Bar plot showing the number of DRE genes differentially expressed both in mRNA-Seq and Ribo-Seq. Heatmap showing the enriched GO terms for DRE genes with up-regulated **(C)** and down-regulated expression **(D)** in at least two tissue types.

Figure 5. The top 10 frequent RNA binding proteins (RBPs) ranked by their binding frequencies to DRE sites in different tissues.

Figure 6. Developmental interrelation between A-to-I RNA editing and alternative splicing. (A) Proportion and number of edited genes with and without alternative splicing (AS) in different tissues. **(B)** Bar plot comparing the number of differential alternative splicing (DAS) events between embryonic and adult stages. **(C)** Venn plots of DAS genes and DRE genes in different tissues. **(D)** Heatmap showing the enriched GO terms for DAS genes with differential editing in at least two tissue types. RI, retained intron; MXE, mutually exclusive exons; A3SS, alternative 3′ splice sites; A5SS, alternative 5′ splice sites; SE, skipped exon.

Figure 7. Validation of RNA editing pattern during brain development in dataset GSE157425 and GSE169457. (A) Functional categories of A-to-I RNA editing variants in the brain's neocortex at embryonic (E12.5–E17) and postnatal (P0) stages. **(B)** Intersection of RNA editing sites in each stage. **(C)** Editing levels in each stage. ***: $P < 0.001$; **: $P < 0.01$. **(D)** Bar plot showing the number of differential editing sites between adjacent developmental time points. **(E)** Heatmap showing the top 50 editing levels of DRE sites during brain development. **(F)** Normalized expression of three *Adars* at the transcriptional (left) and translational level (right). **(G)** *Pearson*'s correlation analysis between number of editing sites and normalized expression of *Adars* at the transcriptional level. **(H)** *Pearson*'s correlation

669   analysis between editing levels and normalized expression of *Adars* at the transcriptional

670   level.

Figure 8. Potential functional impact of A-to-I DRE in transcription, translation, and splicing during brain development. (A) Venn plots comparing genes with DRE and DEGs identified by mRNA-seq (DEG_mRNA) or Ribo-seq (DEG_Ribo). **(B)** Heatmap showing the enriched GO terms for DRE genes with up-regulated and down-regulated expression. **(C)** Venn plots comparing DAS genes and DRE genes. **(D)** Bar plot showing the number of DAS events between adjacent developmental time points. **(E)** Heatmap showing the enriched GO terms for DAS genes with differential editing. RI, retained intron; MXE, mutually exclusive exons; A3SS, alternative 3′ splice sites; A5SS, alternative 5′ splice sites; SE, skipped exon.