Words in manuscript: 3541

Tables: 3 (Supplemental)

Figures: 21 (16 supplemental)

A multi-symptomatic model of heroin use disorder in rats reveals distinct behavioral profiles and

neuronal correlates of heroin vulnerability versus resiliency

Brittany N. Kuhn[1], Ph.D., Nazzareno Cannella[2], Ph.D., Ayteria D. Crow[1], B.S., Veronica Lunerti[2], Ph.D.,

Arkobrato Gupta[3-5], M.S., Stephen J. Walterhouse[1], M.S, Carter Allen[4-5], Ph.D., Reda M. Chalhoub[1], B.S.,

Eric Dereschewitz[1], B.S., Analyse T. Roberts[1], B.S., Mackenzie Cockerham[6], H.S., Angela Beeson[6], H.S.,

Rusty W. Nall[7], Ph.D., Abraham A. Palmer[8], Ph.D., Gary Hardiman[9], Ph.D., Leah C. Solberg Woods[6], Ph.D.,

Dongjun Chung[4-5], Ph.D., Roberto Ciccocioppo[2], Ph.D., and Peter W. Kalivas[1], Ph.D.

[1]Department of Neuroscience, Medical University of South Carolina, Charleston, SC, USA

[2]School of Pharmacy, Center for Neuroscience, Pharmacology Unit, University of Camerino, Camerino, IT

[3]The Interdisciplinary Ph.D. Program in Biostatistics, The Ohio State University, OH, USA

[4]Department of Biomedical Informatics, The Ohio State University, Columbus, OH, USA

[5]Pelotonia Institute for Immuno-Oncology, The James Comprehensive Cancer Center, The Ohio State

university, OH, USA

[6]Department of Internal Medicine, Wake Forest University, Winston-Salem, NC, USA

[7]Department of Psychology, Jacksonville State University, Jacksonville, AL, USA

[8]Institute for Genomic Medicine, University of California San Diego, La Jolla, CA, USA

[9]School of Biological Sciences, Queen's University Belfast, Belfast, Northern Ireland, UK

Corresponding author:

Brittany Kuhn
Medical University of South Carolina
173 Ashley Avenue
BSB 403- MSC 510
Charleston, SC 29425
e-mail: kuhnb@musc.edu
Fax: +1-843-792-4423
29
30

Location of work

Forced swim testing occurred at Wake Forest University (Winston-Salem, NC) in the Department of Internal Medicine. All tests for heroin taking, extinction and seeking measures, along with open field, elevated plus maze and tail flick testing, occurred at the Medical University of South Carolina (Department of Neuroscience, Charleston, SC) and the University of Camerino (School of Pharmacy, Camerino, IT). Neurobiological measures along with estrous phase, ultrasonic vocalizations and punished heroin-taking procedures occurred at the Medical University of South Carolina (Department of Neuroscience, Charleston, SC). Nonlinear clustering was performed at Ohio State University.

Funding and Disclosure

The authors declare no conflict of interest. Funding for these studies was provided by the National Institute on Drug Abuse branch of the National Institutes of Health U01DA45300 (awarded to PWK) and T32DA007288 (BNK).

Acknowledgements

The experiments were designed by BNK, NC, AAP, GH, DC, RC and PWK, and conducted by BNK, NC, ADC, VL, SJW, ED, ATR, MC, AB and RWN. Statistical modeling was performed by CA, AG, DC, and RMC. LCSW provided all heterogeneous stock rats used in these experiments. Data was analyzed by BNK, CA, AG, DC and RMC. BNK and PWK wrote the manuscript. We would like to thank Dr. Carmela Reichel for providing the equipment necessary to perform the punished heroin taking test, and the Department of Neuroscience Behavioral Core at Medical University of South Carolina for the equipment needed to record ultrasonic vocalizations.

Abstract (249 words)

Objective: The behavioral and diagnostic heterogeneity within human opioid use disorder (OUD) diagnosis is not readily captured in current animal models, limiting translational relevance of the mechanistic research that is conducted in experimental animals. We hypothesize that a non-linear clustering of OUD-like behavioral traits will capture population heterogeneity and yield subpopulations of OUD vulnerable rats with distinct behavioral and neurocircuit profiles.

Methods: Over 900 male and female heterogeneous stock rats, a line capturing genetic and behavioral heterogeneity present in humans, were assessed for several measures of heroin use and rewarded and non-rewarded seeking behaviors. Using a non-linear stochastic block model clustering analysis, rats were assigned to OUD vulnerable, intermediate and resilient clusters. Additional behavioral tests and circuit analyses using c-fos activation were conducted on the vulnerable and resilient subpopulations.

Results: OUD vulnerable rats exhibited greater heroin taking and seeking behaviors relative to those in the intermediate and resilient clusters. Akin to human OUD diagnosis, further vulnerable rat sub-clustering revealed subpopulations with different combinations of behavioral traits, including sex differences. Lastly, heroin cue-induced neuronal patterns of circuit activation differed between resilient and vulnerable phenotypes. Behavioral sex differences were recapitulated in patterns of circuitry activation, including males preferentially engaging extended amygdala stress circuitry, and females cortico-striatal drug cue-seeking circuitry.

Conclusion: Using a non-linear clustering approach in rats, we captured behavioral diagnostic heterogeneity reflective of human OUD diagnosis. OUD vulnerability and resiliency were associated with distinct neuronal activation patterns, posing this approach as a translational tool in assessing neurobiological mechanisms underpinning OUD.

Introduction (3523/3500 words)

Opioid use disorder (OUD) inflicts worldwide social and personal harm, with over 16 million individuals affected (1) and approximately 1 million a year seeking treatment (2). Greater understanding of the neurobiological mechanisms underlying OUD is necessary to advance treatment options. A consequential impediment towards attaining this goal is the behavioral heterogeneity inherent in a diagnosis of OUD based on DSM-V criteria. Different behavioral traits, or diagnostic criteria, interact and combine in divergent ways to form an OUD diagnosis, resulting in individuals having the same diagnosis with distinct symptom profiles. Animal models necessary for mechanistic neurobiological experimentation fail to recapitulate the multi-dimensional diagnosis of OUD and largely fail to capitalize the preclinical biological discoveries into useful treatments. We hypothesized that a non-linear clustering protocol applied to multiple behaviors commonly used in mechanistic animal studies of substance use disorder (SUD) would better recapitulate the multidimensional diagnosis of SUD and behavioral diversity conferring vulnerability. We further hypothesized this approach would yield heroin-conditioned cue circuit activation differences between OUD vulnerable and resilient subpopulations.

Typically preclinical studies use individual behavioral traits or summated linear relationships between a few traits to predict and explore underpinning neurobiological causes of SUD (3, 4). These models either do not capture a multi-symptomatic diagnosis, or when doing so rely on linear interactions that do not capture the complex multidimensional interactions between symptoms that creates subpopulations of SUDs patients. Here we describe a rat model and analysis workflow that clusters behaviors using non-linear modeling (5) on data from over 900 male and female outbred heterogeneous stock rats, a line that emulates the complex within species genetic and behavioral heterogeneity found in outbred mammalian populations, including humans (6). Rats were clustered as OUD vulnerable, resilient or intermediate. We further assessed variability within the OUD vulnerable subpopulation and identified distinct subclusters that, akin to human OUD, varied in combinations of traits conferring vulnerability. Also, by correlating Fos

1   protein synthesis (indicator of neuronal activity) stimulated by heroin-conditioned cues across different

2   brain nuclei we identified distinct patterns of connectivity between vulnerable and resilient rats.

Methods

The Medical University of South Carolina Institutional Animal Care and Use Committee, and the Italian

Ministry of Health, approved of all experimental procedures. All procedures complied with the National

Institute of Health Guide for the Care and Use of Laboratory Animals and the Assessment and

Accreditation of Laboratory Animals Care, as well as the European Community Council Directive for Care

and Use of Laboratory Animals.

Subjects

Heterogeneous stock rats (NMcwiWFsm:HS) were bred and shipped from Wake Forest University (WFU,

USA) to either Medical University of South Carolina (MUSC, USA) or the University of Camerino (UCAM,

Italy). Litters were equally represented at both sites in an effort to create genetically comparable testing

cohorts. Final analyses were comprised of 917 rats (male, n=465 (MUSC, n=271; UCAM, n=194); females,

n= 452 (MUSC, n=258; UCAM, n=194)). Yoked saline control rats underwent identical testing procedures

as their counterparts.

Drugs

Heroin hydrochloride supplied by the National Institute on Drug Abuse (Bethesda, MD) was dissolved in

0.9% sterile saline.

Behavioral testing

The behavioral testing procedure is shown in Figure 1a. Rats underwent long-access heroin self-

administration to quantify consumption and escalation of use. Extinction assessed perseverance of

seeking in the absence of reward. Estimates of clinical laboratory-induced craving included both heroin-

prime and cued reinstatement. Motivation for reward was assessed using progressive ratio test.

Additional behavioral traits associated with SUD were also examined prior to and following the heroin

self-administration protocol, including the elevated plus maze (EPM) and open field test (OFT) to assess

stress- and anxiety-like behaviors and analgesic threshold was evaluated using the tail-flick (TF) test. TF

consisted of two phases: baseline (1 mg/kg saline injection, s.c.) and test (0.75 mg/kg heroin or saline

given to control rats). Additionally, a forced swim task (FST) was administered at WFU prior to shipment

to assess stress-coping strategy.

Heroin self-administration, extinction and reinstatement procedures

Rats were outfitted with an indwelling jugular catheter prior to training. During self-administration

presses on the active lever resulted in an infusion of heroin (20 µg/kg/100 µl infusion over 3 seconds) and

presentation of a tone and light cue (Fig. 1b). Saline-control animals received a non-contingent infusion

of saline (100 µl) every 20 min, accompanied with tone/cue-light presentation. There were four 12-h

sessions/week for a total of 12 training sessions. A progressive ratio test occurred on day 36 (Fig. 1a), then

self-administration was re-established. A 6-h extinction/prime session occurred (Fig. 1b) and rats were

administered a heroin priming injection (0.25 mg/kg, s.c.; saline to control rats) after 4-h of extinction.

Daily extinction training sessions then commenced prior to a test for cue-induced reinstatement (Fig. 1b).

Estrous cycle phase was identified in a subset of MUSC rats (n=36) following the tests for heroin-prime

and cued reinstatement.

Additional behavioral testing

Punished heroin-taking behavior: We hypothesized that vulnerable rats would continue seeking heroin

more than the resilient rats in the presence of an adverse stimuli. Three weeks following all testing, a

subset of MUSC rats (males, n=15; females, n=14) re-established heroin self-administration training for 3

days prior to a test session where there was a 50% chance of foot shock delivery (0.40 mA, 0.5 seconds)

with each infusion earned.

<u>Ultrasonic vocalizations (USVs):</u> We examined the potential involvement that different withdrawal-

induced affective states were associated with vulnerable and resilient rats. Four hours prior to extinction

on day 41 (Fig. 1a), a subset of MUSC rats (males, n=22; Females, n=21; Saline, n= 8) were assessed for

acute heroin withdrawal-induced USVs, a preclinical test for assessing affective state (7, 8).

<u>Fos protein quantification:</u> A subset of MUSC vulnerable and resilient rats (Females, n=13; Males, n=18;

Saline, n= 11) underwent 9 additional extinction training sessions and a second cued reinstatement test

~3 weeks after testing was completed. Fos protein expression in regions of interest were manually

quantified.

Statistical analyses

To capture behaviors across OUD phases (heroin use, extinction and seeking), seven behaviors were

selected for analyses as the dependent variables: Heroin use included escalation of intake (µg/kg; average

consumption days 1-3 subtracted from days 10-12) and total consumption (µg/kg) across the first 12

training sessions; Extinction behaviors include active lever presses made during the first 2 hours of the

extinction-prime test (extinction burst) and during the last day of extinction training (extinction day 6);

Seeking measures include break point (maximum active lever presses expended to receive an infusion)

from the progressive ratio test, active lever presses during heroin-prime reinstatement and cued

reinstatement.

Normality was evaluated using a Kolmogorov-Smirnov test. Raw data were assessed for group differences

using a 2-way ANOVA with sex and site as the independent variables and Bonferroni post-hoc test, with

several differences found (ST1). Accordingly, data were z-score transformed within site and assessed for

sex (independent variable) differences using the Mann-Whitney test (SF1). Sex differences were present

for every behavior, and data were standardized within sex and site prior to being recombined for

clustering analyses. Sexes were then analyzed as independent groups for all subsequent analyses.

We previously described a nonlinear network-based clustering analysis (9) and provided an R package

'mlsbm' (5). A rat-rat similarity network was constructed from the standardized data to assess behavioral

similarities between rats, and then a Bayesian stochastic block model (SBM) was applied to define clusters

within the data set. Cluster number was set to 3 (vulnerable, intermediate, resilient) in order to identify

subpopulations of rats potentially most susceptible and least susceptible to developing OUD and in

alignment with a previous study where cluster number was determined according to Bayesian Information

Criterion (SF2) (9). Behavioral differences between clusters (independent variable) were analyzed using

either an ANOVA and Bonferroni post-hoc test or Kruskal-Wallis test and Dunn's post-hoc test.

Repeated-measures ANOVAs (cluster and session as independent variables) and Bonferroni post-hoc test

were used to assess behavior during heroin-reacquisition. Unpaired Student's-t or Mann-Whitney tests

were used for punishment test, USV, and Fos protein analyses (cluster as independent variable for all) and

estrous phase (phase as independent variable). Fos protein was standardized to saline control rats prior

to analyses. Spearman correlation coefficient was used for behavioral correlations with Bonferroni test to

correct for multiple comparisons and Fos protein correlations which employed a false discovery rate of

$q=0.01$ to correct for multiple comparisons. Differences in group composition were evaluated with a Chi-

squared test. Sub-clustering analyses were performed using an agglomerative hierarchical clustering

strategy. The Euclidian distance between subjects with a threshold of 0.7*max(linkage) was used to create

sub-clusters and a minimum of 15 rats/subcluster was necessary to be included in analyses. Multiple

comparisons across different analyses were not corrected for. Unless stated otherwise, analyses were

performed using GraphPad Prism version 9.5.1 (San Diego, CA), with $p < 0.05$ for statistical significance.

Results

Raw data

Substantial site and sex differences existed in the raw data for the OUD-like traits and the behaviors

quantified prior to and following heroin experience (ST1; SF3-7). Female rats exhibited an overall more

OUD vulnerable phenotype than male rats (SF5) (10). Despite standardization of experimental procedures

across sites, MUSC rats showed augmented levels of heroin-related behaviors compared to UCAM rats

(SF5). These source of these site differences presumably results from variables we did not control, such

as shipping, aspects of animal housing protocols and handling, etc (see Supplemental Methods for

detailed description). Because these uncontrolled environmental factors were not quantified and the

study was designed to examine environmental factors that were experimenter controlled and more

related to opioid use and relapse, we z-scored the data within site and sex for further analysis (SF8-9).

However, in future studies it would be of interest to determine which if any environmental differences

between sites contributed to behavioral differences. Following standardizations, linear regression

revealed ubiquitous co-variance across variables for both sites, necessitating an alternative analysis

strategy to better define OUD vulnerability (SF10).

OUD-like trait clustering analysis

SBM clustering was used to model the multidimensional nature of the diagnostic criteria for OUD. Three

statistically distinct (SF2) clusters were established as vulnerable, resilient and intermediate (Male, Fig.

1c; Female, Fig. 1e), that were distributed equivalently between sexes (SF11). The mean value of

behavioral traits was higher in the vulnerable compared with resilient rats in both sexes (Figs. 1d,f),

although the distribution of individuals showed substantial overlap between the clusters (Male, SF8;

Female, SF9). Vulnerable, but not resilient rats in estrus showed potentiated cue-reactivity compared to

non-estrus phase rats. Estrous phase had no effect on behavior during heroin-primed reinstatement (Fig.

1g).

Non-OUD traits

Stress-coping strategy or anxiety-like behavior in basal conditions were not associated with OUD

vulnerability in either sex, as measured by the FS test (SF12) and EPM (Males, SF13c,f; Females, SF14c,f),

respectively. Novelty-induced locomotion was augmented in male but not female vulnerable compared

to resilient rats (SF13a,b), consistent with our previous finding (10) that exploratory behavior can predict

OUD vulnerability in male (r^2=0.18, p<0.001), but not female rats (r^2=0.09, p=0.08; SF14a,b). Analgesic

threshold was not associated with OUD vulnerability in either sex (Males, SF15; Females, SF16). However,

following heroin use, female vulnerable rats had a lower analgesic threshold compared to resilient rats,

implying heroin-induced alterations in pain processing.

Additional trait testing after completing the protocol in F1a

<u>Punishment training:</u> Following ~3 weeks of a forced abstinence vulnerable and resilient rats readily

reacquired heroin self-administration, though vulnerable animals to a greater extent (Males: Fig. 2a, SF

17a; Females: Fig. 2d, SF 17c). Furthermore, both female subpopulations exhibited potentiated heroin

taking on the first day of reacquisition relative to the last day of heroin self-administration training,

indicative of a deprivation effect described for alcohol (11) but not previously shown for opioids (Fig. 2d).

During the punished heroin-taking test, vulnerable male rats endured ~3x the number of infusion-shock

pairings, and females ~2x, prior to attenuating heroin-taking behavior to levels of resilient rats (Males,

Fig. 2b, SF17b; Females, Fig. 2e, SF17d).

<u>Ultrasonic vocalizations:</u> The effect of acute heroin withdrawal on emotional state was quantified using

USVs (8). USVs indicative of a positive affective state (50-kHz) did not differ between clusters in male (Fig.

2c) or female (Fig. 2f) rats. Male vulnerable rats showed potentiated USVs associated with a negative

affective state (22-kHz) compared to resilient rats (Fig. 2c). Female clusters did not differ in negative affect

(Fig. 2f).

Behavioral phenotype sub-clustering

Clusters were probed for the presence of subpopulations to assess behavioral heterogeneity conferring

OUD vulnerability and resiliency. Distinct subclusters existed only in the vulnerable phenotype for both

male (Fig. 3a; n=3 subclusters) and female (Fig. 3b; n=4 subclusters) rats. For each sex, subclusters

corresponded to augmented responding when behavior was: 1) reinforced with heroin (break point,

consumption, escalation, primed reinstatement); 2) not reinforced by heroin (break point, extinction

burst, extinction day 6, cued reinstatement); or 3) in both behavioral categories (Fig. 3c-d). Female rats in

the non-reinforced subcluster exhibited further heterogeneity, with one group having potentiated

responses during acute heroin withdrawal (break point, extinction burst) and the other after more

protracted withdrawal (extinction day 6 and cued reinstatement; Fig. 3d). Overall, males were biased

toward the heroin reinforced subcluster, whereas females toward the heroin non-reinforced subclusters

(Fig. 3e). These data emulate behavioral heterogeneity in OUD diagnosis and emphasize sex differences

in conferring vulnerability.

Cue-induced Fos protein expression

The number of neurons expressing cue-induced Fos protein relative to saline rats differed between

vulnerable and resilient rats in the prelimbic (PrL) and infralimbic cortex (IL), and the anterior

paraventricular nucleus of the thalamus (aPVT; SF18-20). We were surprised to find differences in brain

nuclei with opposite functions. Much of the animal and human literature show activation of PrL and IL

during cued seeking (12) and of the aPVT in suppressing seeking (13, 14). We further evaluated this

apparent contradiction by examining correlated Fos protein between nuclei, which revealed both shared

and distinct patterns between the vulnerable and resilient phenotypes. We subdivided ROI connectivity

into subcircuits that in animal and human imaging studies (15-18) are generally associated with cue-

reactivity (Fig. 4b, f), stress (Fig. 4c, g) and behavioral inhibition (Fig. 4d, h). Commensurate with vulnerable

rats reinstating more that resilient rats (Mann-Whitney U= 18.50, p<0.0001; Mean ± SD: Vulnerable, 42.18

± 25.96; Resilient, 6.07 ± 12.23; SF15m), vulnerable rats showed greater levels of correlated neuronal

engagement (Fig. 4a,e), most notably in ROIs in the extended amygdala associated with stress responses

(19) (Fig. 4c,g). It was also notable that the ventral pallidum (VP), which contains subpopulations of

neurons contributing to all three subcircuits was highly interconnected within all subcircuits in vulnerable

but not resilient rats. To gain insight into possible sex differences contributing toward vulnerability,

vulnerable males and females were assessed separately. Strikingly, sexes shared almost no common

subcircuitry (Fig 5a, e). Female rats exhibited engagement of top-down processes, including the classic

cue-reactivity pathways (20) with correlated neuronal activation between the PrL, nucleus accumbens

core (NAcc) and ventral pallidum (VP; Fig. 5b). Males engaged subcortical cue-reactivity regions,

specifically the NAcc and VP (Fig. 5g, h). Strikingly, only males showed engagement of the extended

amygdala stress circuit (Fig. 5g). Since the sexes exhibited equivalent cued reinstatement (t(15)=0.27,

p=0.79; Mean ± SD: Male, 44.26 ± 17.59; Female, 40.7 ± 31.40), these data suggest sexually dimorphic

circuits contributing to cue reactivity.

Discussion

OUD has increased in all demographics over the past two decades (21), regardless of age, sex, or socio-

economic status (22). To capture the diversity of OUD subpopulations and different combinations of DSM-

V diagnostic symptoms we used genetically heterogeneous rats and a non-linear network-based clustering

of 7 rat behavioral traits. Using this approach, we succeeded in separating vulnerable from resilient rats

and further sub-clustering vulnerable into subpopulations of rats possessing different combinations of

OUD-like behaviors. We validated this approach by showing distinct neurocircuit activation by heroin cues

between vulnerable and resilient subpopulations. We propose that this statistical workflow and

behavioral findings are akin to the multi-dimensional diagnosis of OUD patients and are ideal for

identifying subpopulation heterogeneity in brain mechanisms underpinning OUD vulnerability and

resilience.

OUD vulnerability clustering

The nonlinear clustering approach identified a vulnerable subpopulation of rats exhibiting greater levels

of heroin use, resistance to extinction and seeking compared to rats in the intermediate and resilient

clusters for both sexes. Akin to diagnostic criteria for OUD, behavioral heterogeneity was present in the

traits that confer OUD-like vulnerability in both sexes, with distinct subpopulations exhibiting behavioral

vulnerability in heroin reinforced drug-seeking, non-reinforced drug-seeking or a combination of both.

Male vulnerable sub-clusters were biased toward heroin reinforced seeking, and females toward heroin

non-reinforced seeking. Female rats in the non-reinforced subpopulation were further subdivided into a

subpopulation more motivated to respond during early (i.e., progressive ratio and extinction burst) versus

late extinction conditions (i.e., extinction day 6), with the latter also exhibiting high cue reactivity during

cued reinstatement, suggesting differential sensitivity to contextual and discrete drug-associated cues.

Other measures associated with vulnerability

In addition to the seven traits used for SBM clustering, hallmark features of OUD were identified in

subsequent testing done in a portion of rats. Vulnerable rats re-acquired heroin self-administration and

required more infusion-shock pairings to attenuate heroin-taking behavior. Male vulnerable rats exhibited

more distress than their resilient counterparts during acute heroin withdrawal, with no differences

observed in females. However, male rats are more prone to emit stress-induced USVs (23), suggesting

alternative withdrawal assays may be more applicable for female rats. Circulating ovarian hormones

affected behavioral heterogeneity in female rats. Behavioral responding is potentiated during the estrus

phase of the cycle due to estradiol regulation of mesolimbic dopamine (24), and vulnerable estrus phase

rats showed potentiated cued seeking relative to non-estrus phase rats. No differences were observed

during prime reinstatement, suggesting estradiol involvement in modulating discrete environmental cues,

but not heroin interoceptive cues.

Correlated Fos activity patterns in OUD vulnerable versus resilient rats

Little is known about the neurobiological mechanisms mediating SUD resilience. Decline in prefrontal

cortex grey matter volume (25) and engagement of top-down cortical processes within the dorsal striatum

(26) confers SUD resiliency. Resilient OUD rats exhibit hypo-function of neuronal activation in medial

prefrontal cortex (SF19), suggesting that resiliency may be mediated by prefrontal cortex neuroplasticity.

Resilient rats appear to be recruiting circuitry and regions engaged in negative affective states and

avoidance, such as the PrL-BLA pathway that mediates anxiety-like behavior and fear memory acquisition

(27) and the IL, a region contributing toward suppressing drug seeking (28). This profile of correlated

activation suggests that OUD resiliency is in part governed by circuits that inhibit behavioral responding

to cued seeking.

Commensurate with greater cue-induced heroin seeking, vulnerable rats engaged more complex circuitry,

similar to what is observed in human OUD individuals (16). Key features of connectivity distinctions

between clusters include a majority of connectivity in vulnerable rats centered around the aPVT, a region

involved in mediating anxiety and stress response (29) and suppression of reward seeking (13), the VP

which exhibits cell specific regulation over drug-seeking and drug-refraining (i.e., withholding of seeking)

(30), and the CeA, part of the extended amygdala that mediates stress response and stress-induced

reinstatement (31). In an effort to disentangle the differences in circuit activation we isolated nuclei

generally found activated in the animal and human imaging literature during a cue-reactivity test, stress

responding with a focus on the extended amygdala, and behavioral inhibition (16-19). Most striking was

the involvement of the extended amygdala stress circuit in vulnerable, but not resilient rats. Importantly,

stress facilitates relapse in animal models and humans (32), which can account for the robust cued relapse

in vulnerable, but not resilient rats. Surprisingly, the vulnerable rats also showed greater overall

connectivity in behavioral inhibition circuitry, notably in the interconnectivity between the lHb, PVT and

VP, all structures experimentally implicated in a cell, region or circuit-specific manner to promote or

induce avoidance behaviors (14, 30, 33). It is possible that following chronic heroin self-administration in

vulnerable rats, these brain nuclei have undergone adaptations rendering them less capable of inhibiting

cued seeking (34), or a shift in cell-specific opposing regulation of drug seeking versus avoidance may have

occurred (35).

Sex Differences

In parallel with behavioral sex differences frequently observed in this study, we found minimal overlap in

vulnerability circuitry in male or female rats. Approximately 40% of female rats were in the non-heroin

reinforced sub-cluster, and females heavily engaged the PrL→NAcc→VP series circuit that mediates cue

reactivity in animals and humans (20). In contrast, male rats predominately engaged extended amygdala

circuitry involved in stress responding (31). Vulnerable males also exhibited withdrawal-induced distress

posing the possibility that male rats were experiencing more distress following prolonged heroin

abstinence, resulting in stress circuitry contributing to cued reinstatement. Females, however, showed

lower levels of distress across training (SF 12). Together, these data suggest sexual dimorphism in neuroadaptations governing vulnerability in male and female rats, with male vulnerability driven more by stress circuitry and female vulnerability by cue responsivity. Importantly, the literature supports nuanced sex differences translating to human OUD studies (24), though a more focused approach is necessary in preclinical and clinical research to sufficiently disentangle SUD sex differences (37). Also, we found that female rats show greater inability to withhold from seeking during early extinction, which can result from greater impulsivity, a trait that is more pronounced in human females with OUD (38).

Conclusions and Future Directions

Medication-assisted treatment is the current standard of OUD treatment (39). However, diagnostic heterogeneity is an important consideration for treatment, as individual symptom clusters are likely driven by distinct neurobiological mechanisms and genetics. By using a non-linear clustering model that allowed us to capture aspects of the behavioral heterogeneity and diagnostic diversity in human OUD, we identified distinct behavioral profiles and sexually dimorphic neurobiological circuitry associated with OUD resiliency versus vulnerability in rats. Further sub-clustering revealed variability in traits conferring vulnerability, with notable differences in sub-cluster composition between sexes. This study constitutes a first step in applying non-linear clustering of SUD vulnerability traits and provides a preclinical database for continued studies into the neurobiological mechanisms and genetic vulnerabilities contributing to OUD phenotypes. Additionally, neurobiological underpinnings of sex differences in human SUDs is not well characterized, and despite rates of OUD steadily increasing in females (40), females remain underrepresented in clinical OUD research (<15% of participants) (16). The marked behavioral and circuit sex differences we found highlights using non-linear clustering in animal models to discern mechanisms of sex differences in behavioral phenotypes contributing to human OUD.

References

1. Chang HY, Kharrazi H, Bodycombe D, Weiner JP, Alexander GC: Healthcare costs and utilization associated with high-risk prescription opioid use: a retrospective cohort study. BMC Med 2018;16(1):69

2. National Survey of Substance Abuse Treatment Services (N-SSATS): 2020 [Internet]. Substabce Abuse and Mental Health Services Administration. 2021. Available from: https://www.samhsa.gov/data/sites/default/files/reports/rpt35313/2020_NSSATS_FINAL.pdf

3. Kuhn BN, Kalivas PW, Bobadilla AC: Understanding Addiction Using Animal Models. Front Behav Neurosci 2019;13:262

4. Spanagel R: Animal models of addiction. Dialogues Clin Neurosci 2017;19(3):247-58

5. Allen C, Chung D. mlsbm: Efficient Estimation of Bayesian SBMs & MLSBMs. R package version 0.99.2 ed2021

6. Consortium S, Saar K, Beck A, Bihoreau MT, Birney E, Brocklebank D, et al.: SNP and haplotype mapping for genetic analysis in the rat. Nat Genet 2008;40(5):560-6

7. Covington HE, 3rd, Miczek KA: Vocalizations during withdrawal from opiates and cocaine: possible expressions of affective distress. Eur J Pharmacol 2003;467(1-3):1-13

8. Barker DJ, Simmons SJ, West MO: Ultrasonic Vocalizations as a Measure of Affect in Preclinical Models of Drug Abuse: A Review of Current Findings. Curr Neuropharmacol 2015;13(2):193-210

9. Allen C, Kuhn BN, Cannella N, Crow AD, Roberts AT, Lunerti V, et al.: Network-Based Discovery of Opioid Use Vulnerability in Rats Using the Bayesian Stochastic Block Model. Front Psychiatry 2021;12:745468

10. Kuhn BN, Cannella N, Crow AD, Roberts AT, Lunerti V, Allen C, et al.: Novelty-induced locomotor behavior predicts heroin addiction vulnerability in male, but not female, rats. Psychopharmacology (Berl) 2022;239(11):3605-20

11. Vengeliene V, Bilbao A, Spanagel R: The alcohol deprivation effect model for studying relapse behavior: a comparison between rats and mice. Alcohol 2014;48(3):313-20

12. Hearing M: Prefrontal-accumbens opioid plasticity: Implications for relapse and dependence. Pharmacol Res 2019;139:158-65

13. Engelke DS, Zhang XO, O'Malley JJ, Fernandez-Leon JA, Li S, Kirouac GJ, et al.: A hypothalamic-thalamostriatal circuit that controls approach-avoidance conflict in rats. Nat Commun 2021;12(1):2517

14. McGinty JF, Otis JM: Heterogeneity in the Paraventricular Thalamus: The Traffic Light of Motivated Behaviors. Front Behav Neurosci 2020;14:590528

15. Shin LM, Liberzon I: The neurocircuitry of fear, stress, and anxiety disorders. Neuropsychopharmacology 2010;35(1):169-91

16. Moningka H, Lichenstein S, Worhunsky PD, DeVito EE, Scheinost D, Yip SW: Can neuroimaging help combat the opioid epidemic? A systematic review of clinical and pharmacological challenge fMRI studies with recommendations for future research. Neuropsychopharmacology 2019;44(2):259-73

17. Ray S, Haney M, Hanson C, Biswal B, Hanson SJ: Modeling Causal Relationship Between Brain Regions Within the Drug-Cue Processing Network in Chronic Cocaine Smokers. Neuropsychopharmacology 2015;40(13):2960-8

18. Reiner DJ, Fredriksson I, Lofaro OM, Bossert JM, Shaham Y: Relapse to opioid seeking in rat models: behavior, pharmacology and circuits. Neuropsychopharmacology 2019;44(3):465-77

19. Koob GF, Volkow ND: Neurobiology of addiction: a neurocircuitry analysis. Lancet Psychiatry 2016;3(8):760-73

20. Kalivas PW, Volkow ND: The neural basis of addiction: a pathology of motivation and choice. Am J Psychiatry 2005;162(8):1403-13

21. Wide-ranging online data for epidemiologic research (WONDER) [Internet]. 2020. Available from: http://wonder.cdc.gov/

22. Jones CM, Logan J, Gladden RM, Bohm MK: Vital Signs: Demographic and Substance Use Trends Among Heroin Users - United States, 2002-2013. MMWR Morb Mortal Wkly Rep 2015;64(26):719-25

23. Laine MA, Mitchell JR, Rhyner J, Clark R, Kannan A, Keith J, et al.: Sounding the Alarm: Sex Differences in Rat Ultrasonic Vocalizations during Pavlovian Fear Conditioning and Extinction. eNeuro 2022;9(6)

24. Becker JB, Chartoff E: Sex differences in neural mechanisms mediating reward and addiction. Neuropsychopharmacology 2019;44(1):166-83

25. Ersche KD, Williams GB, Robbins TW, Bullmore ET: Meta-analysis of structural brain abnormalities associated with stimulant drug dependence and neuroimaging of addiction vulnerability and resilience. Curr Opin Neurobiol 2013;23(4):615-24

26. Ersche KD, Meng C, Ziauddeen H, Stochl J, Williams GB, Bullmore ET, et al.: Brain networks underlying vulnerability and resilience to drug addiction. Proc Natl Acad Sci U S A 2020;117(26):15253-61

27. Grossman YS, Fillinger C, Manganaro A, Voren G, Waldman R, Zou T, et al.: Structure and function differences in the prelimbic cortex to basolateral amygdala circuit mediate trait vulnerability in a novel model of acute social defeat stress in male mice. Neuropsychopharmacology 2022;47(3):788-99

28. Peters J, LaLumiere RT, Kalivas PW: Infralimbic prefrontal cortex is responsible for inhibiting cocaine seeking in extinguished rats. J Neurosci 2008;28(23):6046-53

29. Kirouac GJ: The Paraventricular Nucleus of the Thalamus as an Integrating and Relay Node in the Brain Anxiety Network. Front Behav Neurosci 2021;15:627633

30. Kupchik YM, Prasad AA: Ventral pallidum cellular and pathway specificity in drug seeking. Neurosci Biobehav Rev 2021;131:373-86

31. Nall RW, Heinsbroek JA, Nentwig TB, Kalivas PW, Bobadilla AC: Circuit selectivity in drug versus natural reward seeking behaviors. J Neurochem 2021;157(5):1450-72

32. Mantsch JR, Baker DA, Funk D, Le AD, Shaham Y: Stress-Induced Reinstatement of Drug Seeking: 20 Years of Progress. Neuropsychopharmacology 2016;41(1):335-56

33. Velasquez KM, Molfese DL, Salas R: The role of the habenula in drug addiction. Front Hum Neurosci 2014;8:174

34. Paniccia JE, Vollmer KM, Green LM, Grant RI, Winston KT, Buchmaier S, et al.: Restoration of a paraventricular thalamo-accumbal behavioral suppression circuit prevents reinstatement of heroin seeking. Neuron 2023

35. Heinsbroek JA, Bobadilla AC, Dereschewitz E, Assali A, Chalhoub RM, Cowan CW, et al.: Opposing Regulation of Cocaine Seeking by Glutamate and GABA Neurons in the Ventral Pallidum. Cell Rep 2020;30(6):2018-27 e3

36. Russo SJ, Murrough JW, Han MH, Charney DS, Nestler EJ: Neurobiology of resilience. Nat Neurosci 2012;15(11):1475-84

37. Nicolas C, Zlebnik NE, Farokhnia M, Leggio L, Ikemoto S, Shaham Y: Sex Differences in Opioid and Psychostimulant Craving and Relapse: A Critical Review. Pharmacol Rev 2022;74(1):119-40

38. Fattore L, Melis M: Sex differences in impulsive and compulsive behaviors: a focus on drug addiction. Addict Biol 2016;21(5):1043-51

39. Meyer B, Utter GL, Hillman C: A Personalized, Interactive, Cognitive Behavioral Therapy-Based Digital Therapeutic (MODIA) for Adjunctive Treatment of Opioid Use Disorder: Development Study. JMIR Ment Health 2021;8(10):e31173

40. Marsh JC, Park K, Lin YA, Bersamira C: Gender differences in trends for heroin use and nonmedical prescription opioid use, 2007-2014. J Subst Abuse Treat 2018;87:79-85

Figure 1 Experimental timeline and animal OUD phenotype behavioral data. **a)** Rats underwent behavioral

testing (OFT: open field test; EPM: elevated-plus maze; TF: tail flick; FS: forced swim test) prior to jugular

catheterization surgery, and again at the end of heroin experience (excluding FS test). Heroin self-

administration then commenced for 3 weeks (12 hr/session, 4 days a week), followed by a progressive

ratio test and 3 additional days of self-administration training. A within-session extinction-prime

reinstatement test then occurred, followed by extinction training and a test for cued reinstatement. **b)** Illustrations of behavioral paradigm and testing chamber configuration (created with BioRender.com). Active lever presses during self-administration training resulted in tone/cue-light presentation for 5-seconds and a heroin infusion. Active lever presses had no consequence during the within-session extinction-prime test or during extinction training. During cued reinstatement, active lever presses resulted in tone/cue-light presentation, but no heroin infusion. **c)** UMAP representation of male rats separated into distinct OUD vulnerability clusters using stochastic block model network-based clustering analysis. **d)** Mean ± SD for heroin taking, extinction and seeking behavioral measures in male rats. OUD vulnerable rats showing augmented behavioral responding across selected behaviors relative to OUD intermediate and resilient rats (repeated-measure ANOVA for comparison of behaviors across the cluster, cluster x behavior interaction $F_{(12, 2772)}=9.37$, $p<0.0001$). Clusters differed from one another on all behaviors ($p<0.001$) with exception of the intermediate and resilient cluster for escalation of heroin intake. **e)** UMAP presentation of the distinct cluster formation of OUD vulnerability in female rats following stochastic-block model network-based clustering analysis. **f)** Mean ± SD for heroin taking, extinction and seeking behaviors between clusters in female rats. Vulnerable rats showed potentiated behavioral responding relative to intermediate and resilient rats (repeated-measure ANOVA for comparison of behaviors across the cluster, cluster x behavior interaction $F_{(12, 2694)}=4.84$, $p<0.0001$) in all behaviors ($p<0.001$) with the exception of escalation of intake between intermediate and resilient rats. **g)** Mean ± SD for behavior during the reinstatement tests in female rats. Rats (vulnerable, n=31; resilient, n=27) in the estrus phase of the estrous cycle exhibited potentiated responding during the cued reinstatement test relative to non-estrus rats ($t_{(28)}= 0.91$, $p=0.01$). Estrous phase cycle did not affect behavior in vulnerable rats during heroin-primed reinstatement ($t_{(29)}= 0.08$, $p=0.94$), or in resilient rats (heroin-primed reinstatement: $t_{(25)}= 0.1.88$, $p=0.07$; cued reinstatement: $t_{(25)}= 0.57$, $p=0.57$). *$p<0.05$

1   (Males: **c-d**, Vulnerable, n= 182; Intermediate, n=168; Resilient, n=115; Females **e-f**: Vulnerable, n= 204;

2   Intermediate, n=132; Resilient, n=116)

Figure 2 Post-hoc behavioral testing. Mean + SD **a)** for infusions earned the last day of heroin self-administration training (LgA12) and heroin reacquisition in a subset of male rats (n=8/cluster). Vulnerable male rats maintained higher levels of heroin taking during reacquisition (cluster x session interaction, $F_{(3,42)}=5.29$, p=0.004; all post-hoc tests p<0.01). **b)** There was a 50% chance of foot shock delivery with each heroin infusion earned during the punishment task. Foot shock were never delivered during the 20-sec time out period following an infusion. Vulnerable male rats endured more shocks while maintaining heroin-taking behavior compared to resilient rats (inset, Mann-Whitney U= 0, p=0.0002), with differences centralized to the first hour of testing (cluster x hour interaction, $F_{(11,154)}=7.29$, p<0.001; significant post-hoc test p=0.02). **c)** Mean ± SD for acute withdrawal-induced ultrasonic distress vocalization. Calls were higher in vulnerable male rats compared to resilient male rats (22 khz; Mann-Whitney U= 14.5, p=0.02), with no difference in appetitive state calls (50 kHz; $t_{(17)}=0.92$, p=0.37; Vulnerable, n=8; Resilient, n=10). Mean + SD **d)** for infusions earned the last day of heroin self-administration training (LgA12) and heroin

reacquisition in a subset of female rats (vulnerable, n=9; resilient, n=7). A main effect of cluster, but not

cluster x session interaction, was present, indicating vulnerable female rats in general maintained higher

levels of heroin taking during reacquisition (cluster, $F_{(1, 14)}=36.02$, $p<0.0001$; cluster x session interaction,

$F_{(3,42)}=1.43$, $p=0.25$), and **e)** endured more shock-heroin pairing across the test session relative to

resilient rats (inset, Mann-Whitney U= 11, $p=0.03$). However, phenotypes only differed in cumulative

shocks endured across the test session, not across the independent hours of testing (cluster x hour

interaction, $F_{(11,154)}=1.56$; $p=0.12$). **g)** Mean ± SD for acute withdrawal-induced ultrasonic vocalization

calls that did not differ between female phenotypes (22 kHz, $t_{(16)}= 0.17$, $p=0.87$; 50 kHz, Mann-Whitney

U= 30, $p=0.48$; Vulnerable, n=11; Resilient, n=7). *$p<0.05$

Figure 3 Behavioral heterogeneity present within OUD vulnerable clusters. Hierarchical analysis showed that the vulnerable group only in **a)** male and **b)** female rats was composed of distinct behavioral sub-clusters. Clusters with less than 15 animals were excluded from further analysis. Spatial representation of vulnerable sub-cluster behavioral heterogeneity presented as spider plots, where the value at each node represents the average z-score for the indicated behavior for each sub-cluster. **c-d)** Both sexes contained sub-clusters comprised of rats that showed augmented behavior when reinforced by heroin (purple), not reinforced by heroin (red/yellow) and by both circumstances (blue). **e)** Composition of sub-clusters differed between sexes, with vulnerable male rats biased toward heroin reinforced behaviors and female rats toward heroin non-reinforced behaviors (combined yellow and red subclusters in females; $x^2(1,366)=39.93$, $p<0.001$). Number within bar is the number of rats within each sub-cluster. *$p<0.05$ (**a-b**, Males: vulnerable, n= 182; intermediate, n=168; resilient, n=115; Females: vulnerable, n= 204; intermediate, n=132; resilient, n=116)

Figure 4 Correlated neuronal activation patterns between heroin cue-induced c-Fos ROIs in OUD vulnerable and resilient rats. Light purple line illustrates connectivity shared between vulnerable and resilient subclusters; black and blue lines indicate connectivity distinct for vulnerable and resilient subclusters, respectively. **a, e)** Rats in the vulnerable cluster exhibited more correlated neuronal activation compared to resilient rats. Connectivity was further broken down into three different functional circuitry categories: ROIs associated with **b, f)** cue-reactivity, **c, g)** extended amygdala stress response, and **d, h)** behavioral inhibition. Both phenotypes engaged ROIs associated with cue-reactivity, however, the vulnerable group recruited several ROIs that mediate stress response and behavioral inhibition. PrL (prelimbic cortex), IL (infralimbic cortex), NAcc (nucleus accumbens core), NAcs (nucleus accumbens shell), VP (ventral pallidum), lHb (lateral habenula), aPVT (anterior paraventricular nucleus of the thalamus), pPVT (posterior paraventricular nucleus of the thalamus), BLA (basolateral amygdala), CeA (central amygdala), LH (lateral hypothalamus, AP), BNST (bed nucleus of the stria terminalis). Data for

1   each ROI was standardized to saline control rats and correlations shown are p<0.05 with q=0.1

2   (Vulnerable, n= 17; Resilient, n=14; Saline, n=11)

Figure 5 Sexual dimorphic correlated neuronal activation patterns between ROIs in female and male OUD vulnerable rats. All correlated activity patterns for **a)** male and **e)** female vulnerable rats. Connectivity patterns were further broken down into functional circuit categories: ROIs associated with **b, f)** cue-reactivity, **c, g)** extended amygdala stress response, and **d, h)** behavioral inhibition. Male rats show pronounced subcortical activity patterns, specifically engaging the extended amygdala stress circuitry. In contrast, female rats engage top-down cortical processing, with minimal engagement of stress circuitry. PrL (prelimbic cortex), IL (infralimbic cortex), NAcc (nucleus accumbens core), NAcs (nucleus accumbens shell), VP (ventral pallidum), lHb (lateral habenula), aPVT (anterior paraventricular nucleus of the thalamus), pPVT (posterior paraventricular nucleus of the thalamus), BLA (basolateral amygdala), CeA (central amygdala), LH (lateral hypothalamus, AP), BNST (bed nucleus of the stria terminalis). Data for each ROI was standardized to saline control rats and correlations shown are p<0.05 with q=0.1 (Females, n= 10; Males, n= 7; Saline, n=11)

1   **Supplemental methods**

Subjects

Heterogeneous stock rats from NIH (N/NIH-HS)[1] bred at Wake Forest University since 2017 (Rat Genome

Database number 13673907) were used in the experiments. Rats were shipped from Wake Forest

University (n=1200) to MUSC or UCAM at 4-6 weeks of age in batches of 40 (20 females and 20 males/site).

Animals arrived at MUSC via van transport the same day at shipping from Wake Forest. Rats were shipped

via air travel same day from the United States to Rome, Italy, and required 48 of processing and van transit

before arriving at UCAM. During transit, all animals had access to food and hydrogel, and were kept in a

climate controlled environment. Upon arrival at both sites, rats were left undisturbed for 3 weeks in a

climate-controlled vivarium (standard 12-hr light:dark cycle) prior to testing. Rats were pair-housed and

had ad libitum access to standard chow (MUSC: Lab Diet 5V75; UCAM: Mucedola 4RF18) and water

throughout testing. Housing chambers at MUSC contained 1204.5 cm^2 floor space and 22,885.5 cm^3 total

volume (36.5 x 33 cm base by 19 cm tall) and contained corncob bedding and were attached to a Lixit

system for water delivery and ventilation. At UCAM, housing chambers were of similar dimensions at

1126.3 cm^2 floor space and 21,398.8 cm^3 total volume (26.5 x 42.5 cm base by 19 cm tall), contained

wooden dust free bedding and were kept on a free standing rack with 2 water bottles per cage. The

following animals were excluded from analyses: 107 due to death (surgery, n= 23; illness, n= 84), 29 due

to technical issues in data collection, and 147 were used for control self-administration procedures (saline

self-administration, n= 111; sucrose, n=36).

Rats at MUSC had to be transported via freight elevator daily from the floor containing the housing facility

to the testing suite. At UCAM, the housing and testing facilities were located near-adjacent on the same

floor. All testing occurred during the dark phase with the exception of the elevated-plus maze, open field

test and the tail flick test (Figure 1). These behavioral assays occurred in the following sequence starting

at 07:00, with an hour between each test whereupon rats were left undisturbed in their home cage:

elevated-plus maze, open field test then tail flick test. Animals were handled daily the 3 days prior to

beginning the behavioral protocol. All testing procedures, including equipment used and general housing

conditions, were standardized between sites.

Behavioral testing

Forced swim (FS) test: Sessions lasted for 6 minutes and the last 5 minutes of each session was manually

scored in 5 second intervals for measures of immobility (e.g., floating) and mobility (e.g., swimming and

climbing). Testing apparatus was a container (diameter 28 cm, height 36 cm) with a water depth 27 cm

and temperature of 25°C ± 2. Subsequent behavioral testing occurred 3 weeks after the rats' arrival to the

study site location, immediately preceding heroin self-administration training.

Elevated-plus maze (EPM): The EPM test lasted for 5 minutes. The apparatus (San Diego Instruments; 4

arms at 43.5" long and 4" wide) consisted of two enclosed arms (12" high walls; "closed" arms) and two

"open" arms (i.e. not enclosed). Testing occurred under dark conditions, with infrared lighting overhead

of the apparatus to allow for accurate scoring. Behavior was recorded and automatically scored using

ANY-maze behavioral tracking software (Stoelting, Wood Dale, IL; version 6.17) with a threshold of 85%

of animal's body within an arm to be counted in it.

Open field test (OFT): Sessions lasted for 60 minutes in a Plexiglas chamber sitting within a metal frame

(Omnitech Electronics, Columbus, OH; 16" L x 16" W x 12" H) housed within a dark room. Photocells within

the chambers tracked lateral and vertical movements, and data was recorded and analyzed using

Versamax (Omnitech Electronics, Columbus, OH; version 1.80-0142).

Tail flick test: Testing occurred on a platform outfitted with a motion sensor that turns the beam off once

the animal removed its tail, or after 10 seconds has passed (Ugo Basile S.R.L., Gemonio, Italy). Injections

(i.e., saline or heroin) were administered 15 minutes prior to session start. The two phases were separated

by 1 hour and consisted of 4 trials that were averaged to determine the mean latency to remove the tail

from the sensor. To prevent tissue damage, the location of the rat's tail over the beam was adjust by 1 cm

each trial.

Heroin self-administration, extinction and reinstatement procedures

Prior to training, rats were outfitted with an indwelling jugular catheter and administered an analgesic

(Ketorolac, 2 mg/kg, s.c.; or Meloxicam, 0.5 mg/rat, s.c.) and an antibiotic (Cefazolin, 0.2 mg/kg, s.c.; or

enrofloxacin, 1 mg/kg, i.v.) post-operatively. Catheter patency was verified prior to training starting via

blood collection for additional analyses (data not presented in this manuscript). If a rat showed a sudden

decrease in heroin-taking behavior, or the catheter port was compromised (e.g., leaking), patency was

checked using methohexital sodium diluted in sterile saline (10 mg/ml, i.v.). Rats were re-catheterized

(MUSC, n= 15; UCAM, n= 30) if ataxia was not observed within 5 seconds. Following a 3-day recovering,

rats were reintroduced back into experimentation. All training occurred in standard behavioral testing

chambers (Med Associates, St. Albans, VT) comprised of two levers with a light above each, and a speaker

and house light on the opposite wall (Figure 1b). A fixed-ratio 1 schedule of reinforcement was used for

self-administration. Once an infusion was earned a 20-second time out period was signaled by the house

light turning off whereupon additional active lever presses were recorded, but did not result in a heroin

infusion. Sessions terminated after 12 hours or 300 earned infusions, with four training sessions/week

Monday-Friday, one randomized day off per week. A progressive ratio test was used to assess motivation

to take heroin, during which the number of lever presses needed to receive an infusion exponentially

increased according to the following formula: $(5*e^{0.2n})-5$[2]. Testing terminated after 1 hour of no earned

infusion or 12 total hours of testing. During extinction training, active lever presses were without

consequence (i.e., no infusion or cue presentation). Active lever presses during cued reinstatement

resulted in tone/cue-light presentation, but no infusion.

Estrous cycle phase identification

Vaginal lumen samples were collected and processed with a hematology stain from a subset of female

MUSC rats (n=36) following the tests for heroin-prime and cued reinstatement to determine estrous cycle

phase at time of testing. Phase was determined via cell morphology using a Leica Thunder DM6B scope.

Ultrasonic vocalizations

Rats were placed into a self-administration testing chamber housed within a sound attenuating cabinet in

a quiet room, and ultrasonic vocalizations (USVs) were recorded for 5-minutes. Testing chambers were

off (i.e., no levers or cues were present). The recording microphone was placed approximately 12.25 cm

from the top of the chamber, beyond physical reach from the rat. Vocalizations were recorded using

Avisoft UltraSoundGate equipment (Avisoft Bioacoustics, Glienicke/Nordbahn, Germany), and the

following parameters were used: FFT was set to 256, bandwidth of 3672 Hz and resolution of 977 Hz.

Recordings were sorted using Avisoft SAS Lab Pro. Both the audio recording and the spectrogram were

used to identify USVs and associated frequency range, as well as to exclude ambient noise (e.g., animal

locomotive behaviors, other environmental noises, etc.). USV duration had to exceed 25 ms to be included

in analysis. USVs between 18-33 kHz (i.e., 22-kHz range) and 38-80 kHz (i.e., 50-kHz range) frequency range

were quantified in 60-second intervals (i.e., 5 60-sec bins).

Behavioral phenotype subclusters

Sub-clustering analyses were performed using an agglomerative hierarchical clustering strategy on

individual subclusters, using custom MATLAB scripts. Similarity between subjects was measured using

Euclidian distance (pdist function) along the variables by which the animals were originally clustered:

escalation of intake, total consumption, break points, and active lever presses during extinction burst,

extinction day 6, heroin-prime reinstatement and cued reinstatement. A hierarchal clustering tree was

subsequently created by the *ward* linkage method on the calculated distances between subjects. The tree

was then divided into subsequent subclusters using a threshold of 0.7*max(linkage). A minimum of 15

rats per subcluster was used as a cut-off to be included in post-hoc analyses on the generated subclusters.

Fos protein quantification

Immediately following the second reinstatement test, animals underwent transcardial perfusion and

brains were extracted and post-fixed prior to being sliced (40 μm) for Fos protein immunohistochemistry.

Brains were first washed in 0.1 M phosphate-buffered saline (PBS; 3x5 min/wash) and blocked in 1%

hydrogen perozide in PBS for 10 min prior to undergoing 4 additional PBS washes. Slices were blocked in

PBS + 0.4% Triton-X + 2.5% normal donkey serum (Jackson ImmunoResearch, Product #017-000-121) for

1 hour at room temperature, then incubated overnight in rabbit anti-cFos primary antibody (1:2000

dilution; Cell Signaling, Product #5348S). Brain slices were rinsed in PBS and incubated in biotinylated

donkey anti-rabbit secondary antibody (1:500 dilution; Jackson ImmunoResearch, Catology #711-065-

152) at room temperature for 1 hour and then rinsed in PBS. Incubation in ABC-Elite followed (1:1000

dilution; Vector Labs, Product #PK-6100) for 1 hour and tissue was rinsed in PBS. Last, sliced were

incubated for 8 minutes in diaminobenzidine tetrahydrochloride (DAB; Signma-Aldrich, Product #D5905)

and 0.012% hydrogen eproxide prior to additional rinses in PBS. Slices were mounted onto slides,

underwent graduated ethanol dilution washes and a Xylene wash prior to being cover slipped with

Permount mounting medium.

Images of regions of interest (ROI) were acquired as a 20x brightfield tilescan using a Leica Thunder DM6B

scope outfitted with a Flexacam C1 camera. ImageJ was used for manual c-Fos protein cell quantification,

and ROI parameters were identical for a region across animals to standardize data. Fos protein number

for each animal and ROI was standardized to the average Fos protein number in saline control rats for the

respected ROI (i.e., Fos protein for ROI/average Fos protein level in saline control rats for ROI). Fos protein

levels across the ROIs were correlated with one another [3-5] to assess potential neural circuitry involved

in OUD vulnerability versus resiliency.

References

1. Hansen, C. and K. Spuhler, *Development of the National Institutes of Health genetically heterogeneous rat stock.* Alcohol Clin Exp Res, 1984. **8**(5): p. 477-9. https://www.ncbi.nlm.nih.gov/pubmed/6391259

2. Richardson, N.R. and D.C. Roberts, *Progressive ratio schedules in drug self-administration studies in rats: a method to evaluate reinforcing efficacy.* J Neurosci Methods, 1996. **66**(1): p. 1-11. https://www.ncbi.nlm.nih.gov/pubmed/8794935

3. Flagel, S.B., et al., *A food predictive cue must be attributed with incentive salience for it to induce c-fos mRNA expression in cortico-striatal-thalamic brain regions.* Neuroscience, 2011. **196**: p. 80-96. PMC3206316

4. Haight, J.L. and S.B. Flagel, *A potential role for the paraventricular nucleus of the thalamus in mediating individual variation in Pavlovian conditioned responses.* Front Behav Neurosci, 2014. **8**: p. 79. PMC3953953

5. Ruiz, C.M., et al., *Pharmacokinetic, behavioral, and brain activity effects of Delta(9)-tetrahydrocannabinol in adolescent male and female rats.* Neuropsychopharmacology, 2021. **46**(5): p. 959-969. PMC8115040

a   Raw data: Behavioral tests

	Site			Sex			Site * Sex		
	DF	F	P	DF	F	P	DF	F	P
Tail flick baseline 1	1,878	46.84	**<0.0001**	1,878	2.56	0.11	1,878	5.01	**0.03**
Tail flick test 1	1,878	533.3	**<0.0001**	1,878	80.50	**<0.0001**	1,878	20.74	**<0.0001**
Tail flick difference 1	1,878	320.5	**<0.0001**	1,878	58.17	**<0.0001**	1,878	27.63	**<0.0001**
Time travelled 1	1,874	2257	**<0.0001**	1,874	2.37	0.12	1,874	26.40	**<0.0001**
Distance travelled 1	1,874	61.76	**<0.0001**	1,874	119.0	**<0.0001**	1,874	56.95	**<0.0001**
Percent time open arm 1	1,877	126.0	**<0.0001**	1,877	83.14	**<0.0001**	1,877	1.90	0.17
Number of climbs	1,843	0.16	0.69	1,843	26.66	**<0.0001**	1,843	0.10	0.75
Time swimming	1,843	4.56	**0.043**	1,843	1.43	0.23	1,843	0.05	0.82
Time floating	1,843	3.81	0.05	1,843	0.69	0.41	1,843	0.04	0.84
Tail flick baseline 2	1,878	103.6	**<0.0001**	1,878	5.74	**0.02**	1,878	14.36	**0.0002**
Tail flick test 2	1,878	287.4	**<0.0001**	1,878	21.26	**<0.0001**	1,878	9.29	**0.002**
Tail flick difference 2	1,878	136.7	**<0.0001**	1,878	11.18	**0.0009**	1,878	21.02	**<0.0001**
Time travelled 2	1,872	1496	**<0.0001**	1,872	0.13	0.72	1,872	16.79	**<0.0001**
Distance travelled 2	1,872	125.6	**<0.0001**	1,872	47.29	**<0.0001**	1,872	15.67	**<0.0001**
Percent time open arm 2	1,877	4.03	**0.04**	1,877	52.68	**<0.0001**	1,877	0.31	0.58

b   Raw data: Heroin taking, extinction and seeking

	Site			Sex			Site * Sex		
	DF	F	P	DF	F	P	DF	F	P
Escalation	1,913	52.17	**<0.0001**	1,913	2.98	0.08	1,913	0.46	0.50
Consumption	1,913	29.35	**<0.0001**	1,913	66.53	**<0.0001**	1,913	0.003	0.96
Break point	1,913	118.4	**<0.0001**	1,913	1.31	0.25	1,913	0.11	0.74
Extinction burst	1,913	114.4	**<0.0001**	1,913	5.46	**0.02**	1,913	0.0006	0.98
Prime reinstatement	1,913	0.73	0.39	1,913	15.69	**<0.0001**	1,913	0.68	0.41
Extinction day 6	1,913	17.13	**<0.0001**	1,913	6.09	**0.01**	1,913	1.14	0.29
Cued reinstatement	1,913	21.18	**<0.0001**	1,913	15.89	**<0.0001**	1,913	3.09	0.08

Supplemental Table 1 Raw data statistics for all behavioral testing and heroin taking, extinction (extinction behaviors) and seeking measures. **a)** Behavioral differences were highly prevalent between sites (MUSC vs UCAM) and sex (male vs female) both prior to (test 1) and following (test 2) heroin self-administration training. Forced swim testing only occurred prior to heroin experience. **b)** Differences were present between sites for every measure except active lever pressing during heroin-prime reinstatement. Behavior between males and females were also present, with sexes not differing to escalation of heroin taking and break point achieved during the progressive ration test only. No interactions were present. Abbreviations: DF, degrees of freedom; F, F-test value; P, probability value

a

<div align="center">

OUD: Vulnerable

</div>

ROI 1	ROI 2	Spearman r	p-value	n
PrL	IL	0.8	0.00019	17
	VP	0.74	0.001	17
IL	VP	0.58	0.015	17
	aPVT	0.5	0.049	16
NAcc	VP	0.6	0.013	17
NAcs	aPVT	0.61	0.013	16
	BLA	0.89	<0.0001	14
	CeA	0.81	0.0007	14
	BNST	0.6	0.013	17
	VP	0.67	0.0043	17
VP	lHb	0.82	0.001	13
	aPVT	0.68	0.0046	16
	pPVT	0.75	0.0046	13
	BLA	0.6	0.025	14
	CeA	0.61	0.022	14
	LH	0.76	0.0022	14
LH	lHb	0.83	0.0027	14
	CeA	0.55	0.044	14
	BNST	0.72	0.0047	14
BNST	BLA	0.79	0.0014	14
	CeA	0.75	0.0029	14
CeA	BLA	0.91	<0.0001	14
	aPVT	0.55	0.044	14
BLA	aPVT	0.71	0.0062	14
pPVT	aPVT	0.62	0.027	13
	lHb	0.69	0.011	13

b

<div align="center">

OUD Resilient

</div>

ROI 1	ROI 2	Spearman r	p-value	n
PrL	pPVT	0.69	0.016	12
	BLA	0.66	0.013	14
	VP	0.75	0.0027	14
	IL	0.8	0.001	14
IL	LH	0.58	0.033	14
	VP	0.76	0.0024	14
NAcc	pPVT	0.64	0.028	12
	BLA	0.75	0.022	14
	NAcs	0.9	<0.0001	14
NAcs	BLA	0.75	0.0029	14
VP	LH	0.67	0.011	14
LH	lHb	0.7	0.019	11
	BLA	0.59	0.03	14
pPVT	aPVT	0.87	0.0005	12
	lHb	0.66	0.03	11

Supplemental Table 2 Data statistics for significant correlated neuronal activation in OUD vulnerable and resilient male and female rats. **a)** Vulnerable rats showed overall augmented levels of activation patterns relative to **b)** resilient rats. All data was standardized to saline control animals (n=11) prior to comparison. All correlations were positive, with a p<0.05 and a q=0.01 to correct for multiple comparisons. Abbreviations: ROI, region of interest; Spearman r, correlation coefficient; n, number of rats/comparison; PrL (prelimbic cortex), IL (infralimbic cortex), NAcc (nucleus accumbens core), NAcs (nucleus accumbens shell), VP (ventral pallidum), lHb (lateral habenula), aPVT (anterior paraventricular nucleus of the thalamus), pPVT (posterior paraventricular nucleus of the thalamus), BLA (basolateral amygdala), CeA (central amygdala), LH (lateral hypothalamus), BNST (bed nucleus of the stria terminalis)

a

		OUD Vulnerable: Males		
ROI 1	ROI 2	Spearman r	p-value	n
PrL	IL	0.86	0.0009	7
NAcs	lHb	1	0.018	5
	BLA	0.75	0.018	7
	CeA	0.86	0.021	7
	VP	0.96	0.0009	7
VP	lHb	1	0.013	5
	CeA	0.82	0.026	7
	LH	0.87	0.034	7
LH	lHb	1	0.0033	5
CeA	BLA	0.96	0.0005	7
BLA	aPVT	0.96	0.023	7

b

		OUD Vulnerable: Females		
ROI 1	ROI 2	Spearman r	p-value	n
PrL	IL	0.78	0.011	10
	VP	0.90	0.00016	10
	aPVT	0.80	0.014	9
	NAcc	0.73	0.02	10
IL	aPVT	0.82	0.011	9
NAcc	VP	0.81	0.0072	10
aPVT	VP	0.78	0.017	9
pPVT	lHb	0.89	0.0058	8
BNST	LH	0.93	0.0067	7

Supplemental Table 3 Data statistics for significant correlated neuronal activation in OUD vulnerable male and female rats. Vulnerable **a)** male and **b)** female rats showed distinct neuronal activation patterns. All data was standardized to saline control animals (n=11) prior to comparison. All correlations were positive, with a p<0.05 and a q=0.01 to correct for multiple comparisons. Abbreviations: ROI, region of interest; Spearman r, correlation coefficient; n, number of rats/comparison; PrL (prelimbic cortex), IL (infralimbic cortex), NAcs (nucleus accumbens shell), VP (ventral pallidum), lHb (lateral habenula), aPVT (anterior paraventricular nucleus of the thalamus), BLA (basolateral amygdala), CeA (central amygdala), LH (lateral hypothalamus)

Supplemental Figure 1 Data for heroin-taking, extinction and seeking behaviors following z-score

transformation to standardize data between sites (MUSC vs UCAM). The black bar within each violin plot

indicates the median value of the data set, and all data represented as z-scores. Females exhibited

potentiated behavioral responding relative to males for **a)** escalation (Mann-Whitney U= 97160, p<0.05)

and **b)** consumption (Mann-Whitney U= 71799, p<0.0001) of heroin, **c)** break point achieved during the

progressive ratio test (Mann-Whitney U= 92258, p<0.001), extinction behavior during the **d)** first (Mann-

Whitney U= 86294, p<0.0001) and **f)** last (Mann-Whitney U= 92872, p<0.002) extinction training, as well

as seeking behavior during the **e)** heroin-primed (Mann-Whitney U= 79385, p<0.0001) and **g)** cue-

induced (Mann-Whitney U= 85034, p<0.0001) reinstatement. *p<0.05 (Females, n= 452; Males, n=465)

Supplemental Figure 2 Stochastic block model (SBM) and behavioral clustering parameters. **a)** Bayesian

information criterion (BIC) for SBM fits with a range of cluster (K) number between 2 to 10. **b)** Adjacency

matrix of inferred clusters from the SBM using K=3 clusters. **c)** Cluster uncertainty score for each rat on

UMAP space. Numbers represent rat ID for any rats with less than a 90% certainty of belonging to the

assigned cluster (n=7; 0.008% of total population). **d)** Cluster connectivity (Θ) parameters for within ($\Theta_{1,1}$,

$\Theta_{1,2}$, and $\Theta_{1,3}$) and between ($\Theta_{1,2}$, $\Theta_{1,3}$, and $\Theta_{2,3}$) assigned clusters. Higher Θ value indicates greater

connectivity and 95% credible interval reported below Θ. Greater connectivity was observed within a

cluster relative to between clusters, indicating strong behavioral cohesiveness within SBM cluster analysis

assignments. (Vulnerable, n=386; Intermediate, n=300; Resilient, n=231)

Supplemental Figure 3 Raw data for tail flick (TF) test behavior for site (MUSC and UCAM) and sex (female and male) in rats. The black bar within each violin plot indicates the median value of the data set. Significant interactions between site and sex were present for both baseline and test sessions prior to (Baseline, a, $F_{1,878}$=5.01, p=0.03; Test, **b**, $F_{1,878}$=20.74, p<0.0001; Difference, **c**, $F_{1,878}$=27.63, p<0.0001) and following (Baseline, a, $F_{1,878}$=14.36, p=0.0002; Test, **b**, $F_{1,878}$=9.29, p=0.002; Difference, **c**, $F_{1,878}$=21.02, p<0.0001) heroin self-administration training. *p<0.05 (MUSC: Male, n= 253; Female, n= 241; UCAM: Male, n= 194; Female, n=194)

Supplemental Figure 4 Raw data for open field test (OFT), elevated pluz maze (EPM) and forced swim test (FST) behaviors for site (MUSC and UCAM) and sex (female and male) in rats. Median value for data sets represented as the black bar within each violin plot. A significant testing location and sex interaction was present for time spent moving and distance travelled during the OFT both prior to (Time, **a**, $F_{1,874}$=26.40, p<0.0001; Distance, **b**, $F_{1,874}$=56.95, p<0.0001) and following (Time, **d**, $F_{1,872}$=16.79, p<0.0001; Distance, **e**,

$F_{1,872}=15.67$, p<0.0001) heroin self-administration experience. Effects of site and sex, but no interactions, were present for time spent in the open arm of the EPM **c)** before (Site, $F_{1,877}=126$, p<0.0001; Sex, $F_{1,877}=83.14$, p<0.0001) and **f)** after (Site, $F_{1,877}=4.03$, p=0.04; Sex, $F_{1,877}=52.68$, p<0.0001) heroin experience. During the FST, females exhibited more **g)** climbing behavior relative to males ($F_{1,843}=26.66$, p<0.0001), and rats that were trained at UCAM showed more **h)** swimming behavior ($F_{1,843}=4.56$, p=0.04) compared to those at MUSC. No differences were present for **i)** time spent floating, or interactions for any FST measures. *p<0.05 (n for each test designated as 'time point 1/time point 2'; OFT: MUSC: Male, n= 253/253; Female, n= 240/241; UCAM: Male, n= 193/189; Female, n=192/193; EPM: MUSC: Male, n= 252/252; Female, n= 241/241; UCAM: Male, n= 194/194; Female, n=194/194; FST: Male, n= 245; Female, n= 236; UCAM: Male, n= 183; Female, n= 182)

Supplemental Figure 5 Data for heroin-taking, extinction and seeking behavior for site (MUSC and UCAM) and sex (female and male) rats. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. Rats undergoing training at MUSC versus UCAM differed in **a)** escalation ($F_{1,913}$=52.17, p<0.0001) and **b)** consumption of heroin ($F_{1,913}$=29.35, p<0.0001) and **c)** break point achieved during the progressive ratio test ($F_{1,913}$=118.40, p<0.0001). Active lever presses made during the **d)** first ($F_{1,913}$=114.40, p<0.0001) and **f)** last day of extinction training ($F_{1,913}$=17.13, p<0.0001), as well as during **g)** cued reinstatement ($F_{1,913}$=21.18, p<0.0001), between sites were also present. Relative to males, females exhibited potentiated levels of **b)** heroin consumption ($F_{1,913}$=66.53, p<0.0001), and greater levels of extinction (first day, **d**, $F_{1,913}$=5.46, p=0.02; last day, f, $F_{1,913}$=6.09, p=0.01) and seeking behavior (prime reinstatement, **e**, $F_{1,913}$=15.69, p<0.0001; cued reinstatement, **g**, $F_{1,913}$=15.89, p<0.0001. No significant interactions were present. *p<0.05 (MUSC: Male, n= 271; Female, n= 258; UCAM: Male, n= 194; Female, n=194)

Supplemental Figure 6 Raw data for heroin-taking, extinction and seeking behavior for male and female rats at MUSC. All data shown with mean ± SD. **a)** All rats escalated intake across heroin self-administration training session (session, $F_{(5.41,2829)}=71.06$, $p<0.0001$), with females maintaining higher lever of consumption relative to males (sex, $F_{(1,525)}=39.83$, $p<0.0001$) infusions earned across heroin self-administration training. **b)** Female rats exhibited higher break points during the progressive ratio test compared to males (Mann-Whitney U= 31196, $p<0.03$). **c)** All rats showed attenuated responding on levers as training continued (session x lever, $F_{(5,5245)}=7.58$, $p<0.0001$), however, female rats exhibited higher levels of responding on both levers (sex x lever, $F_{(1,1050)}=4.03$, $p<0.05$), and across training session (session x sex, $F_{(5,5245)}=2.96$, $p=0.01$) compares to males. All rats showed greater responding during the reinstatement test relative to the preceding extinction training session, and differentiated between the levers during both sessions (**d**, lever x session, $F_{(1,1054)}=32.95$, $p<0.0001$, all post-hocs $p<0.05$; **e**, lever x session, $F_{(1,1054)}=152.6$, $p<0.0001$, all post-hocs $p<0.001$). Females showed higher levels of responding

during both extinction and prime reinstatement (sex, $F(1,1054)=6.76$, $p=0.01$). *$p<0.05$ (Male, n= 271; Female, n= 258)

Supplemental Figure 7 Raw data for heroin-taking, extinction and seeking behavior for male and female rats at UCAM. All data shown with mean ± SD. **a)** Females exhibited higher levels of heroin consumption across all training sessions compared to males (session x sex, $F_{(11,4282)}=2.53$, $p=0.04$, all post-hoc $p<0.02$). **b)** Females achieved high break points during the progressive ratio test compared to male rats (Mann-Whitney U= 15796, $p<0.05$). **c)** Rats continued to differentiate levers across training (session x lever, $F_{(5,3965)}=7.58$, $p=0.03$), with females showing higher levels of responding on both levers (sex x lever, $F_{(1,802)}=8.01$, $p=0.005$). A lever x session x sex interaction was present for behavior during the extinction session preceding reinstatement and reinstatement for both **d)** prime ($F_{(1,772)}=9.54$, $p<0.02$, all post-hocs $p<0.05$) and **e)** cue ($F_{(1,772)}=9.60$, $p=0.002$, all post-hocs $p<0.005$). *$p<0.05$ (Male, n= 194; Female, n= 194)

Supplemental Figure 8 Data for heroin-taking, extinction and seeking behavior for OUD vulnerability clusters in male rats. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. Behavior differed across clusters for **a)** escalation (Kruskal-Wallis, H(2)=206.9, p<0.0001) and **b)** consumption (Kruskal-Wallis, H(2)=278.2, p<0.0001) of heroin, **c)** break point achieved during the progressive ratio test (Kruskal-Wallis, H(2)=240.5 p<0.0001), extinction behavior during the **d)** first (Kruskal-Wallis, H(2)=140.7, p<0.0001) and **f)** last (Kruskal-Wallis, H(2)=177.7, p<0.0001) extinction training session and seeking behavior during the **e)** heroin-primed (Kruskal-Wallis, H(2)=115.7, p<0.0001), **g)** cued reinstatement test (Kruskal-Wallis, H(2)=212.6, p<0.0001) and **h)** summation of z-scores for the behaviors is a-g (Kruskal-Wallis, H(2)=388.6, p<0.0001). All significant post-hoc tests were p<0.0001. *p<0.05 (Vulnerable, n= 182; Intermediate, n= 168; Resilient, n=115)

Supplemental Figure 9 Data for heroin-taking, extinction and seeking behavior for OUD vulnerability clusters in female rats. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. Behavioral differences are present in clusters for **a)** escalation (Kruskal-Wallis, H(2)=133.9, p<0.0001) and **b)** consumption (Kruskal-Wallis, H(2)=274.8, p<0.0001) of heroin, **c)** break point achieved during the progressive ratio test (Kruskal-Wallis, H(2)=230.4 p<0.0001), extinction behavior during the **d)** first (Kruskal-Wallis, H(2)=135.5, p<0.0001) and **f)** last (Kruskal-Wallis, H(2)=171.9, p<0.0001) extinction training session and seeking behavior during the **e)** heroin-primed (Kruskal-Wallis, H(2)=150.0, p<0.0001), **g)** cued reinstatement test (Kruskal-Wallis, H(2)=191.8, p<0.0001) and **h)** summation of z-scores for the behaviors is a-g (Kruskal-Wallis, H(2)=372, p<0.0001). All significant post-hoc tests were p<0.0001. *p<0.05 (Vulnerable, n= 204; Intermediate, n= 132; Resilient, n=116)

Supplemental Figure 10 Correlation matrixes for behaviors used in non-linear clustering analysis for male and female rats. The value within each box is the Spearman's correlation coefficient. Significant covariance was present for all behavioral interactions for both **a)** male and **b)** female rats. p<0.05 (Male, n= 465; Female, n= 452)

Supplemental Figure 11 OUD phenotypic composition for male and female vulnerable (V), intermediate (I) and resilient (R) rats. Cluster composition within sex did not differ between male and female rats (x^2(1,917)=5.4, p=0.07). Number inside of bar is the number of animals for that cluster.

Supplemental Figure 12 Forced-swim test behavior for OUD vulnerability clusters in female and male rats. Rats underwent a forced-swim test prior to heroin experience. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. Clusters did not differ from one another for either female or male rats in **a, d)** number of climbs (Females: Kruskal-Wallis, H(2)=4.1, p=0.13; Males: Kruskal-Wallis, H(2)=3.3, p=0.19), **b, e)** time spent swimming (Females: Kruskal-Wallis, H(2)=3.5, p=0.18; Males: Kruskal-Wallis, H(2)=2.1, p=0.35), **c, f)** or time spent floating (Females: Kruskal-Wallis, H(2)=1.7, p=0.43; Males: Kruskal-Wallis, H(2)=2.1, p=0.36) for either females or males. (Females, n= 418; Males, n=429)

Supplemental Figure 13 Open field test (OFT) and elevated plus maze (EPM) behavior for OUD vulnerability clusters in male rats. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. Vulnerable rats spent more **a)** time moving (ANOVA, $F_{(2,442)}=4.19$, $p=0.02$; post-hoc $p=0.01$) and **b)** travelled a greater distance (ANOVA, $F_{(2,442)}=5.34$, $p=0.001$; post-hoc $p=0.01$) than resilient rats during the open field task prior to heroin experience. Following heroin experience, rats did not differ in **d)** time spent moving (ANOVA, $F_{(2,439)}=2.70$, $p=0.07$), but vulnerable rats maintained **e)** higher levels of distance travelled relative to resilient rats (Kruskal-Wallis, $H_{(2)}=7.5$, $p=0.02$; post-hoc $p=0.02$). Phenotypes overall differed in percent time spent in the open

arm of the elevated plus maze **c)** prior to (Kruskal-Wallis, H(2)=6.6, p=0.04, no significant post-hocs) and

f) following heroin experience (Kruskal-Wallis, H(2)=14.2, p=0.0008; vulnerable vs intermediate post-hoc

p=0.0005). *p<0.05 (OFT: Vulnerable, n=175 (174 OFT 2); Intermediate, n=161 (159 OFT 2); Resilient,

n=109; EPM: Vulnerable, n= 175 (174 OFT 2); Intermediate, n= 163 (161 OFT 2); Resilient, n=109)

Open field and elevated plus maze: Females

Supplemental Figure 14 Open field test (OFT) and elevated plus maze (EPM) behavior for OUD vulnerability clusters in female rats. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. Rats did not differ in time spent moving or total distance travelled prior to (**a**) Time: Kruskal-Wallis, H(2)=0.21, p=0.90; **b)** Distance: Kruskal-Wallis, H(2)=0.22, p=0.89) and following (**d**) Time: F(2,432)=2.88, p=0.06; **e)** Distance: Kruskal-Wallis, H(2)=3.1, p=0.22) heroin experience. **c)** A difference in behavior between the phenotypes were present in EPM 1 (Kruskal-Wallis, H(2)=11.8, p=0.003), with intermediate rats spending less percent time in the open arm compared to vulnerable rats (p=0.002). **f)** No phenotypic differences were observed in the second EPM

(F(2,432)=2.06, p=0.13). *p<0.05 (OFT: Vulnerable, n= 192 (198 OFT 2); Intermediate, n= 130; Resilient,

n=107; EPM: Vulnerable, n= 198; Intermediate, n= 130; Resilient, n=107)

Supplemental Figure 15 Tail flick test (TF) behavior for OUD vulnerability clusters in male rats. The black

bar within each violin plot indicates the median value of the data set, and all data represented as z-scores.

No phenotypics differences were observed in the **a)** baseline session (Kruskal-Wallis, H(2)=2.8, p=0.24),

and though a difference in phenotype was present for the **b)** test session (Kruskal-Wallis, H(2)=6.6,

p=0.04), there were no significant post hoc. **c)** A difference in latency between the two sessions were not

observed (F(2,444)=1.49, p=0.23). Phenotypes did not differ in behavior in the **d)** baseline session

following heroin experience (Kruskal-Wallis, H(2)=5.9, p=0.05), but did in the **e)** test session (Kruskal-

Wallis, H(2)=13.5, p=0.001) with vulnerable rats having lower latency times than intermediate rats

(p=0.001). **f)** This relationship was present in the latency difference between the two sessions (Kruskal-

Wallis, H(2)=11.9, p=0.003), with intermediate rats having a higher latency than vulnerable rats (p=0.002).

*p<0.05 (Vulnerable, n= 175; Intermediate, n= 163; Resilient, n=109)

Tail flick test: Females

☐ Vulnerable (V)   ☐ Intermediate (I)   ☐ Resilient (R)

Tail flick: Time point 1

Supplemental Figure 16 Tail flick test (TF) behavior for OUD vulnerability clusters in female rats. The black bar within each violin plot indicates the median value of the data set, and all data represented as z-scores. No phenotypics differences were observed in the **a)** baseline session (Kruskal-Wallis, H(2)=1.6, p=0.46) and **b)** test session (Kruskal-Wallis, H(2)=3.7, p=0.16), or the **c)** latency difference between the two sessions (F(2,432)=2.02, p=0.13). Following heroin experience, clusters did not differ during the **d)** baseline session (Kruskal-Wallis, H(2)=5.3, p=0.07), however, vulnerabe rats showed lower latency responses compared to the resilient rats during the **e)** test session (Kruskal-Wallis, H(2)=9.5, p=0.009;

vulnerable vs intermdiate p=0.006) and **f)** the latency difference between the two sessions (F(2,432)=7.76,

p=0.0005). *p<0.05 (Vulnerable, n= 198; Intermediate, n= 130; Resilient, n=107)

Heroin reacquisition and punishment testing: Infusions

a   Heroin reacquisition: Males

b   Punishment testing: Males

c   Heroin reacquisition: Females

d   Punishment testing: Females

Supplemental Figure 17 Infusions earned across the last day of heroin reacquisition and the punishment test. All data represented as mean + SD. Vulnerable animals (males, n=8; females, n=7) maintained higher levels of infusions earned across the last day of heroin reacquisition training (session 3) relative to resilient animals (males, n=8; females, n=7), with **a)** effects localized to the first hour of the session in males (hour x cluster, $F_{(11,154)}=2.64$, p=0.004) and **d)** across training for females (cluster, $F_{(1,14)}=12.65$, p=0.003). These effects were maintained during the punishement test, with **b)** vulnerable males earning more infusions in the first two hours of the test (hour x cluster, $F_{(11,154)}=7.05$, p<0.001, and female vulnerable

rats in the first hour of the test (hour x cluster, $F_{(11,132)}=2.59$, p=0.005). *p<0.05 (Male, n= 194; Female,

n= 194)

Vulnerable   # Resilient

Supplemental Figure 18 Representative micrographs of c-Fos protein density in vulnerable and resilient rats. Images are from regions that showed phenotypic differences in Fos protein quantification. PrL (prelimbic cortex), IL (infralimbic cortex); aPVT (anterior paraventricular nucleus of the thalamus

Supplemental Figure 19 Fos protein quantification comparisons between OUD vulnerable and resilient

rats. All data was standardized to saline control rats prior to comparison. Relative to rats in the vulnerable

cluster, resilient rats exhibited hypofunctioning in the **a)** PrL, **b)** IL and **g)** aPVT. Phenotypes did not differ

in Fos quantity in other ROIs. **m)** Vulnerable animals maintained higher levels of cued seeking behavior

during the second cued reinstatement test compared to resilient animals (Mann-Whitney U= 18.50,

p<0.0001). The anterior-posterior (AP) coordinate from the skull relative to Bregma (mm) where

quantification of select ROIs occurred are within the parantheses. PrL (prelimbic cortex, AP 3), IL

(infralimbic cortex, AP 3), NAcc (nucleus accumbens core, AP 1.56), NAcs (nucleus accumbens shell, AP

1.56), VP (ventral pallidum, AP 0), lHb (lateral habenula, AP -3), aPVT (anterior paraventricular nucleus of

the thalamus, AP -2), pPVT (posterior paraventricular nucleus of the thalamus, AP -3), BLA (basolateral

amygdala, AP -2.4), CeA (central amygdala, AP -2.4), LH (lateral hypothalamus, AP -2.4), BNST (bed nucleus

of the stria terminalis, AP 0). *p<0.05 (Vulnerable, n= 17; Resilient, n= 14; Saline, n=11)

Supplemental Figure 20 Raw Fos protein cell quantification comparisons between saline control, OUD

resilient and vulnerable rats. Relative to rats in the vulnerable cluster, resilient rats exhibited

hypofunctioning in the **g)** aPVT ($F(2,36)= 4.18$, $p=0.02$, post-hoc $p=0.03$). Groups did not differ in Fos

quantity in other ROIs. **m)** Vulnerable animals maintained higher levels of cued seeking behavior during

the second cued reinstatement test compared to saline control and resilient animals (Kruskal-Wallis,

$H(2)= 21.96$, $p<0.0001$, post-hocs $p=0.0002$). The anterior-posterior (AP) coordinate from the skull

relative to Bregma (mm) where quantification of select ROIs occurred are within the parantheses. PrL

(prelimbic cortex, AP 3), IL (infralimbic cortex, AP 3), NAcc (nucleus accumbens core, AP 1.56), NAcs

(nucleus accumbens shell, AP 1.56), VP (ventral pallidum, AP 0), lHb (lateral habenula, AP -3), aPVT

(anterior paraventricular nucleus of the thalamus, AP -2), pPVT (posterior paraventricular nucleus of the

thalamus, AP -3), BLA (basolateral amygdala, AP -2.4), CeA (central amygdala, AP -2.4), LH (lateral

hypothalamus, AP -2.4), BNST (bed nucleus of the stria terminalis, AP 0). *p<0.05 (Vulnerable, n= 17;

Resilient, n= 14; Saline, n=11)