Another fly diuretic hormone: tachykinins increase fluid and ion transport by adult *Drosophila melanogaster* Malpighian 'renal' tubules

Marishia A. Agard,[a] **Meet Zandawala,**[b] **and Jean-Paul V. Paluzzi**[a,*]
[a]Department of Biology, York University, 4700 Keele Street, Toronto, Ontario, M3J 1P3, Canada
[b]Department of Biochemistry and Molecular Biology, University of Nevada Reno, Reno, 89557, NV, USA
*Correspondence should be addressed to J.P.P. (paluzzi@yorku.ca)

Due to dynamic environmental conditions including temperature, water and food availability, insects like the model organism Drosophila melanogaster, must modulate their internal physiology to withstand these challenges. Regulation of the excretory system by peptidergic hormones is one mechanism by which insects maintain their internal homeostasis. Tachykinins are a family of neuropeptides that have been shown to stimulate fluid secretion from the Malpighian 'renal' tubules (MTs) in some insect species, but it is unclear if that is the case in the fruit fly, D. melanogaster. A central objective of the current study was to examine the physiological role of tachykinin signaling in the MTs of adult D. melanogaster. Using the genetic toolbox available in this model organism along with in vitro and whole animal bioassays, our results indicate that Drosophila tachykinins (DTKs) function as diuretic hormones by binding to the DTK receptor (DTKR) expressed in stellate cells of the MTs. Specifically, DTK activates cation and anion transport across the stimulated MTs, which negatively impacts their survival in response to desiccation due to their inability to conserve water. Thus, besides their previously described roles in neuromodulation of pathways controlling locomotion and food search, olfactory processing, aggression, lipid metabolism and metabolic stress, processing of noxious stimuli and hormone release, DTKs also appear to function as bona fide endocrine factors regulating the excretory system and appear essential for the maintenance of hydromineral balance.

Introduction

Insects like the fruit fly, *Drosophila melanogaster*, are routinely faced with challenges in their natural environment including changes to temperature, humidity, food availability and access to water, which necessitate feedback mechanisms and other physiological processes to maintain homeostasis [1,2]. Insects living in arid habitats have evolved to increase their tolerance to desiccation by reducing the rate of water loss and conserving body water content [1]. One strategy that insects utilize to enhance survival in response to environmental stresses involves the regulation of the excretory organs, composed of the Malpighian tubules (MTs) and hindgut [3,4]. The MTs in *D. melanogaster* are composed of two main epithelial cell types: (1) the principal cells, which are the most abundant cells making up the MTs that mediate transepithelial secretion of Na+ and K+ ions using the ionomotive enzyme V-type H+-ATPase; and (2) the relatively less abundant stellate cells that are responsible for Cl- and water transport [4,5]. Hormones that act upon the MTs can be released from neurosecretory cells or endocrine cells localized in the central nervous system or peripheral tissues (e.g. intestines), respectively [6]. These hormones target distinct epithelial cell types of the MTs to function as either diuretic or anti-diuretic factors by mainly interacting with specific G protein-coupled receptors (GPCRs) that can stimulate or inhibit transepithelial ion transport and, consequently, osmotically obliged water [6–8]. Therefore, to overcome excessive or limited amounts of water and solutes associated with their particular diet or environment, *Drosophila* modulate their MTs and other components of the excretory system by peptidergic hormones and other neurochemicals to regulate primary urine production and reabsorptive processes [9] .

Tachykinins are a group of neuropeptides that are expressed in the central nervous system and intestine in both vertebrates and invertebrates and are widely considered as pleiotropic factors [10]. Insect tachykinin-related peptides have been shown to function as myostimulatory peptides on peripheral tissues including the foregut, midgut, hindgut, and oviduct [11,12]. Studies have shown that insect tachykinins, such as those found in locusts and cockroaches, display myotropic effects as determined via muscle contraction assays [13,14]. One prototypical insect tachykinin-related peptide, locustatachykinin, has also been shown to stimulate tubule writhing activity, which may assist with urine flow [15]. In addition to their myotropic effects, exogenous tachykinins stimulate secretion by the MTs in the tobacco hornworm, *Manduca sexta* [16]. Similarly, locustatachykinin has been shown to elicit dose-dependent diuretic effects in isolated tubules from two locust species, *Schistocerca gregaria* and *Locusta migratoria* [17]. These effects of tachykinins and tachykinin-related peptides can be mediated via both cAMP and PLC-IP$_3$-calcium signal transduction pathways [18,19].

We ask here whether *Drosophila* tachykinins (DTKs), act as diuretic (neuro)peptide hormones with a potential role in regulating excretory organs that contribute towards ion-water balance and homeostasis. The *Drosophila tachykinin* (*Dtk*) gene is expressed in interneurons of the central nervous system including in the brain and subesophageal

ganglion [20]. DTKs are localized in about 150 neurons in the adult fly central nervous system aiding in olfactory and locomotor behaviour in search of food during times of hunger [21,22]. DTKs are also co-expressed with ion-transport peptide (ITP), and short neuropeptide F (sNPF) in neurosecretory cells in the adult *Drosophila* brain, where they may function as neuromodulators [23]. Studies have also revealed that DTKs are expressed in enteroendocrine cells in the midgut where they regulate physiological functions including intestinal lipid metabolism [20,24]. *Dtk* encodes for six structurally related peptide isoforms (DTK-1-6) that bind and activate two GPCRs, *Drosophila* tachykinin receptor (DTKR or TkR99D) and neurokinin K receptor (NKD or TkR86C) [20,25,26]. While NKD has been detected in the central nervous system and midgut [27,28], some evidence indicated that NKD is not a tachykinin receptor. Instead, it has more recently been recognized as a *bona fide* receptor of natalisin, which is a tachykinin-related neuropeptide family in arthropods that influences reproductive physiology in insects [28]. Comparatively, DTKR is expressed in the central nervous system, midgut, and MTs and is activated by DTK-1-5 [19]. However, recent studies have revealed that DTKs from a certain subset of brain neurons target neurons that express both receptors, DTKR and NKD, to function as neuromodulators of aggressive behaviour and visual aversive responses [29,30]. In the current study, we focused on the potential function of DTKs in regulating transepithelial secretion by the adult *D. melanogaster* MTs via DTKR.

Here, we confirm the cell-specific expression of DTKR in the fly MTs and provide insight into the physiological role of DTK/DTKR signaling in relation to ion-water balance. Based on earlier reports examining tachykinin activity on insect MTs [16,17,31,32], we predicted that fly DTKs function as diuretic hormones by activating DTKR expressed in the stellate cells of the MTs to stimulate fluid transport, which may be important for the maintenance of hydromineral balance, especially during times of stress. To address these hypotheses, we first conducted a single-nucleus RNA sequencing (snRNA-seq) analysis of *Drosophila* MTs to determine the expression of DTKR (TkR99D). We then used two independent cell-specific driver lines to knockdown *DTKR* expression in the MTs and measured the transcript levels of *DTKR* in this organ of the adult fruit fly. To further establish the physiological role of DTK/DTKR signaling in the MTs, various *in vitro* assays were conducted on isolated MTs along with whole animal stress assays where DTKR was specifically knocked down in MTs. Our results strongly support the notion that DTKR is indeed expressed in the stellate cells, and its knockdown in the MTs affects renal function in the adult fruit fly that ultimately disrupts ion transport capacity. Finally, knockdown of DTKR exclusively in the MTs leads to increased lifespan during desiccation stress, which corroborates the diuretic activity of DTKs on the MTs and their influence on hydromineral homeostasis. Thus, these findings indicate that DTK/DTKR signaling contributes towards ionic and osmotic balance as DTKR knockdown flies displayed lower ion secretion and enhanced survival during desiccation stress. Overall, these findings establish the role of DTKs in hormonal control of the renal organs and build upon a

growing list of diuretic and anti-diuretic factors in *Drosophila* that are fundamental for hydromineral homeostasis.

Results

Single-nucleus RNA sequencing analysis of MTs confirms DTKR expression in the stellate cells

snRNA-seq analysis was conducted to confirm the cell-type specific expression of DTKR (TkR99D) in the MTs. The MTs are composed of 10 distinct clusters, which can express a variety of diuretic (or anti-diuretic) hormone receptors. Based on the snRNA-seq analysis, a number of diuretic hormone receptors [33–36] including *TkR99D*, *leucokinin receptor* (*Lkr*), and *diuretic hormone 44 receptor 2* (*DH44-R2*) are highly expressed in the cluster comprised of the stellate cells of the main segments (**Fig 1A-C**). Specifically, roughly 75% of the stellate cells of the main segment express *TkR99D*, *Lkr* and *DH44-R2*. In contrast, distinct peptidergic regulators of the MTs have receptors that are not expressed in stellate cells including those for diuretic hormone 31 (DH_{31}) and Capa peptides, but instead display high expression in clusters representing the principal cells (**Fig 1A-C**) in strong agreement with earlier findings [37,38]. Notably, the *tyramine receptor* (*TyrR*) was strongly expressed in both the principal cells of the lower segment and stellate cells of the main segments (**Fig 1A-C**). Intriguingly, *TkR99D* is only expressed in the star-shaped (main segment) stellate cells while the *Lkr* is expressed in both bar (initial segment) and star-shaped (main segment) stellate cells (**Fig 1C**). Interestingly, similarity clustering of stellate cells based on expression of diuretic hormone receptors shows that cells expressing *Tkr99D* also express *Lkr* (**Fig 1D**). Independently, TkR99D-GAL4 drives strong GFP expression in the star-shaped stellate cells, thus validating our snRNA-seq analysis (**Fig 1E**). Further examination of *TkR99D*-positive stellate cells shows that they express a variety of diuretic hormone receptors and water channels (*Drip and Prip*), the latter of which might be central for primary urine production when tubules are stimulated by DTKs (**Fig 1F**).

Examining the physiological role of DTK on isolated MTs

Secretion rates were measured from isolated MTs in an *ex vivo* assay to determine the full dose-response characteristics of DTK-1, which was used as a representative of all *Dtk* derived peptides since all peptide isoforms have similar efficacy in DTKR activation [19]. MTs were isolated from wild-type (w^{1118}) flies and treated with different concentrations of DTK-1 (ranging from 10^{-4} to 10^{-13} M) to determine an intermediate concentration that could then be used to test out secretion rates from MTs isolated from DTKR-RNAi flies since high doses might not be as sensitive to reduced receptor expression. The dose-response analysis demonstrates the diuretic activity of DTK-1, with an EC_{50} of 1.126 x 10^{-7} M (**Fig 2A**), which is similar to other insect tachykinins such as locustatachykinin I that was tested on *Locusta migratoria* and *Schistocerca gregaria* MTs [17]. Indeed 10^{-7} M DTK-1 was shown to be an intermediate dose as it elicited significantly higher fluid secretion rates compared to unstimulated and 10^{-9} M DTK-1 treated MTs while displaying significantly lower rates of secretion when compared to 10^{-5} M DTK-1 treated tubules (**Fig 2B**). In addition, the MTs were treated separately with

DTK-1, DTK-2, and DTK-3 at 10^{-7} M, which revealed similar secretion rates and diuretic activity (**Fig 2C**), indicating that multiple DTK peptide isoforms have similar effects in driving secretion by isolated tubules.

Knockdown of DTKR expression in adult fly MTs

To functionally validate the expression of DTKR in the adult fruit fly MTs, we utilized the bipartite GAL4-UAS system to drive *DTKR* RNAi using two independent cell-specific GAL4 drivers: (1) *uro*-GAL4 and (2) *c724*-GAL4, which drive GFP expression in principal and stellate cells, respectively (**Fig 3A and 3B**). Previous studies have established the expression patterns of these two lines with *uro*-GAL4 driving expression in the more abundant principal cells found along the length of both the anterior and posterior MTs, including the initial and the main segments [39,40]. Comparatively, *c724*-GAL4 drives expression in the star-shaped stellate cells from the initial to the main segment of both anterior and posterior MTs [39]. First, the principal cell *uro*-GAL4 driver line was crossed with two different RNAi lines (*uro*-GAL4>UAS-*DTKR* RNAi 1 and *uro*-GAL4>UAS *DTKR* RNAi 2) in an attempt to knockdown *DTKR* in the principal cells of the MTs. Tubules from the experimental flies displayed a similar *DTKR* abundance to the controls (**Fig 3C**) with no evidence of a reduction in *DTKR* transcript levels. Next, the stellate cell driver was crossed with the two DTKR-RNAi lines (*c724*-GAL4>UAS-*DTKR* RNAi 1 & *c724*-GAL4>UAS-*DTKR* RNAi 2) to knockdown *DTKR* in stellate cells of the MTs (**Fig 3D**). In contrast to the principal cell manipulation, a significant reduction in *DTKR* abundance was observed in MTs isolated from stellate cell driven DTKR-RNAi flies relative to MTs from controls that displayed normal DTKR expression (**Fig 3D**). These results support the notion that DTKR is expressed in the stellate cells, but not the principal cells of the MTs, and hence identify the former cell type in MTs as the cellular target of circulating DTK. These results are consistent with the snRNA-seq data from the adult fruit fly that detected high expression of *DTKR* in the main segment stellate cells (**Fig. 1A, 1C, 1E**) of the MTs [41]. This evidence also indicates that *DTKR* is co-expressed in stellate cells with other diuretic hormone receptors (**Fig. 1F**), including *Lkr* and *TyrR* [42–44].

Secretion rates from tubules isolated from knockdown flies

To examine the functional impact of *DTKR* knockdown in the renal tubules, the MT fluid secretion assay was completed on DTKR-RNAi flies in response to DTK-1 using the classic *in vitro* assay described by Ramsay [45]. Fluid secretion rates by MTs isolated from flies with DTKR-RNAi driven by the principal cell driver (*uro*-GAL4>UAS-*DTKR* RNAi 1 & *uro*-GAL4>UAS *DTKR* RNAi 2) and the genetic control flies displayed a significant increase (~4-fold) in fluid secretion rates when treated with 10^{-7} M DTK-1 (**Fig 3E and 3F**). The MTs from *uro*-GAL4 driven RNAi lines displayed similar secretion rates to the parental control lines under both basal (unstimulated) and DTK-1 stimulated conditions. These *ex vivo* bioassay results support the RT-qPCR analysis indicating that DTKR is not expressed in the principal cells of the MTs.

Fluid secretion rates were then measured in MTs isolated from flies with DTKR-RNAi driven by the stellate cell driver (*c724*-GAL4>UAS-*DTKR* RNAi 1 & *c724*-GAL4>UAS-*DTKR* RNAi 2) and the genetic control flies (**Fig 3G and 3H**). Basal unstimulated fluid secretion rates by MTs from *c724*-GAL4>UAS-*DTKR* RNAi 1 & *c724*-GAL4>UAS-*DTKR* RNAi 2 flies were similar to parental control lines. However, the MTs with *DTKR* knockdown in stellate cells displayed a significantly reduced response to DTK-1 with an 81-84% lower stimulated secretion rate compared to parental control lines. Thus, the knockdown of DTKR in stellate cells decreases stimulated secretion rates in response to DTK-1, but rates were marginally (and significantly) elevated compared to the unstimulated basal rates of secretion. Altogether, the results confirm that reduced expression of DTKR specifically in stellate cells leads to lower fluid secretion rates in response to DTK-1, which complements the earlier data demonstrating strong GFP expression in the star-shaped stellate cells driven by TkR99D-GAL4 (**Fig 1E**). Similarly, this functional data for DTK-1 on stellate cells substantiates the cell-type specific expression results observed for *DTKR* in the snRNA-seq (**Fig 1A, 1C and 1F**) and RT-qPCR analysis (**Fig 3C and 3D**).

DTK-1 alters transepithelial ion transport by adult Malpighian tubules

Given that DTK elicits stimulatory effects on fluid secretion rates of MTs isolated *ex vivo* from wild-type flies, we predicted that ion concentrations within the secreted fluid and/or transport rate may be altered since passive water movement is coupled to ion transport. Before investigating the effects on transepithelial transport across MTs isolated from DTKR knockdown flies, the activity of DTK-1 was examined in MTs from wild-type flies. Using ion-selective microelectrodes (ISME) to measure select ion concentrations in the primary urine droplets from MTs [46], we first quantified Na^+ concentration in the secreted fluid and its transport rate following treatment with DTK-1, which revealed a significant decrease in Na^+ concentration in the secreted fluid (**Fig 4A**) but a significant increase in Na^+ transport rates (**Fig 4B**) compared to unstimulated MTs. Similarly, there was a significant decrease in the K^+ concentration (**Fig 4C**) in the secreted fluid and a significant increase in K^+ transport rate (**Fig 4D**) when wild-type MTs were treated with DTK-1. Notably, wild-type MTs treated with DTK-1 demonstrated a significant increase in Cl^- concentration (**Fig 4E**) in the secreted fluid and an increase in the transport rate of this anion (**Fig 4F**) compared to unstimulated tubules. Therefore, these results suggest that DTK drives the movement of anions, specifically Cl^-, similar to the ***Drosophila*** leucokinin that displays effects on overall chloride transport [47]. Altogether, these data indicate that, along with increased fluid secretion rates, DTK stimulates an increase in cation and anion transport by targeting DTKR on stellate cells and thereby increasing primarily chloride permeability.

We next examined how stellate cell-specific knockdown of DTKR in the MTs might impact transepithelial ion transport. In light of the significantly diminished DTK-1 stimulated fluid secretion in flies with *DTKR* knockdown in the stellate cells of the MTs using two different RNAi lines, we proceeded to use one of these (UAS-***DTKR*** RNAi 1) for further analysis using ISME to ex-

amine the effect on tubule transepithelial ion transport. We observed a significant decrease in Na^+ concentration in the *DTKR* knockdown flies when exposed to DTK-1 compared to the unstimulated treatment (**Fig 5A**). However, there was no significant change in Na^+ concentration between the *DTKR* knockdown flies and control flies in the DTK-1 treatment (**Fig 5A**). The K^+ concentration in tubule secreted fluid by the DTKR-RNAi flies displayed no significant change following DTK-1 treatment when compared to the unstimulated treatment (**Fig 5B).** Notably, there was no change in transport rate of either cation between unstimulated and DTK-1 stimulated MTs in *DTKR* knockdown flies (**Fig 5C and 5D**) whereas their transport rates significantly increased after DTK-1 stimulation in control flies consistent with earlier data on wildtype flies (**Fig. 4B and 4D**). Comparatively, the Cl^- concentration in fluid secreted by MTs and the transport rate of this anion in *DTKR*-RNAi flies was significantly reduced in comparison to the control flies (**Fig 5E and 5F**). In summary, while there was an increase in the Cl^- concentration in the secreted fluid and transport rate of this anion in control flies, *DTKR* knockdown abolished these activated epithelial transport characteristics. Together, these results demonstrate that *DTKR* knockdown in stellate cells of the MTs primarily reduces chloride conductance, which in turn decreases counterion (cation) transport that collectively leads to diminished diuretic activity.

Renal-specific DTKR knockdown improves desiccation survival

Considering our data revealing that *DTKR* knockdown in stellate cells reduces DTK-1 stimulated transepithelial transport, we predicted that these flies may also exhibit an increase in lifespan during stress conditions challenging fluid and ion homeostasis, such as desiccation. Stress assays were completed on DTKR-RNAi 1 flies driven by the stellate cell driver as it was the most effective in knocking down *DTKR* levels (**Fig 1D**). Flies were deprived of water and food to induce desiccation stress, which revealed flies exhibiting DTKR knockdown in stellate cells of the MTs (*c724*-GAL4>UAS-***DTKR*** RNAi 1) exhibited a significant increase in lifespan with a maximal survival of 90 hours compared to the parental control flies (**Fig 6A**). Comparably, in the starvation stress assay where flies were deprived only of food, but not water, there was no significant difference in survival between the DTKR knockdown and parental control flies (**Fig 6B**), suggesting that DTK/DTKR renal signaling may not be involved in feedback mechanisms related to nutritional stress. Altogether, the results indicate that *DTKR* knockdown in the MTs may contribute towards greater water conservation in the animal due to decreased diuretic activity, which in turn enhances their dehydration tolerance and survival during desiccation stress.

Discussion

In this study, we combined cell-type specific RNAi screening and various *ex vivo* and *in vivo* biological assays to provide evidence that DTKs are peptidergic hormones that interact with DTKR expressed in the MTs to contribute towards ionic and osmotic balance in the adult fruit fly, *D. melanogaster*. Previous studies in other insects have indicated that tachykinins are not only myostimulatory on

peripheral tissues but may also function as a diuretic factors increasing fluid secretion rates of isolated MTs [11,12,20]. However, since the MTs are not innervated, tachykinins elicit their activity on MTs by functioning as circulating hormones [32]. There have been attempts to localize the expression of the DTKR in the MTs, but these proved challenging due to apparent contradictory evidence [19,48]. Therefore, we attempted to functionally localize DTKR in specific cells of the MTs by utilizing the snRNA-seq analysis and GAL4/UAS bipartite system to knockdown expression in the renal tubules using cell-specific drivers and characterize deficiencies in DTK-DTKR signaling by complementary *in vivo* and *ex vivo* bioassays.

Based on the snRNA-seq analysis, the expression of *TKR99D* was localized to the star-shaped stellate cells of the main segments of the MTs. In addition, we were able to confirm the localization to other diuretic hormone receptors that are either expressed in the principal cells or stellate cells. These include the receptors for leucokinin, DH_{44}, and tyramine displaying high expression in the stellate cells of the MTs, which for the most part is consistent with previous studies [36,41,44] Contrastingly, earlier evidence on the DH_{44} receptor suggested that it is expressed in principal cells where DH_{44} was reported to increase cAMP levels [49]. Interestingly, however, principal cell driven RNAi of the DH_{44} receptor failed to abolish the diuretic activity of DH_{44} on isolated MTs [36]. The high expression of *TkR99D* in the stellate cells of the main tubule segment suggests that DTK/DTKR signaling may be an essential diuretic hormone for maintaining hydromineral balance by mediating transcellular transport of primarily chloride ions and water due to the high expression of chloride and water channels localized to the stellate cells [5,47,50]. Intriguingly, the fact that TkR99D-positive stellate cells display high expression of other diuretic hormone receptors, including the overlap in expression with the *Lkr*, suggest that these diuretic hormones may be functioning together to aid in transepithelial transport to maintain ion-water balance [5,36].

Tachykinins in both vertebrates and invertebrates are potent stimulators inducing contractions of peripheral tissues, including the hindgut, oviduct, and the MTs in insects [11,12,14,15,20]. Since previous studies demonstrated that tachykinins in other insect species act as diuretic hormones [16,17], we aimed to determine whether this peptidergic signaling system plays a similar role in the fruit fly. Indeed, we observed that DTKs are also potent stimulators of fluid secretion by isolated MTs whereby a dose-dependent increase in fluid secretion rate is observed, with an EC_{50} in the nanomolar range. This physiological dose of DTK is consistent with previous studies that have investigated other diuretic factors like tyramine and leucokinin that target stellate cells in the MTs inducing a rapid increase in fluid transport [7,44]. Relatedly, previous studies have highlighted the diuretic role of different isoforms of insect tachykinin-related peptides at various concentrations, including in the pharate adult *Manduca sexta*, where concentrations in the range of 0.01-1 μM elicit diuretic activity on isolated MTs [16].

The next question we addressed in this study was where are DTKs specifically acting in the MTs to elicit their diu-

retic activity? Our findings indicate that DTKR is expressed in the main segment stellate cells and not in principal cells of the MTs. This conclusion is based on multiple lines of evidence including *DTKR* knockdown driven by the stellate cell driver which had significantly lower *DTKR* transcript abundance whereas no change in transcript level was observed when knockdown was attempted using the principal cell driver. These findings are consistent with our snRNA-seq analysis using a recent dataset that characterized the cell types of the adult fly MTs [41]. Specifically, the star-shaped main segment stellate cells were found to express *DTKR* in support of the notion that these cells in the MTs are the target of circulating DTKs [41]. The results also corroborate the initial study that immunolocalized DTKR in the bar-shaped stellate cells in the larval fruit fly MTs [19]. Not only was DTKR transcript abundance affected, but stellate cell driven knockdown also significantly reduced fluid secretion rates stimulated by DTK-1 by more than 80%. Comparatively, MTs with principal cell driven knockdown of DTKR had normal DTK-1 stimulated secretion rates. These results mirror data from earlier studies that found knockdown of receptors for diuretic factors in the stellate cells resulted in reduced stimulated secretion rates compared to control fly lines [36,44,51].

Altogether the data implies that DTK is a diuretic hormone that targets DTKR in the tubule stellate cells, which like leucokinin and tyramine [7,36,44,51], contribute towards the regulation of hydromineral balance in the insect. In light of the diuretic activity of DTK on the stellate cells of MTs and considering DTKR downstream signaling involves increases in Ca^{2+} [19,52], it may trigger a second messenger pathway similar to the other diuretics targeting the stellate cells to stimulate fluid secretion. The importance of calcium and confirmation of its involvement in eliciting DTK-DTKR signaling in the tubule stellate cells awaits further investigation, but such signaling has been deemed important for modulating nociceptive sensitization in *Drosophila* [52].

Besides establishing its role as a diuretic hormone leading to increased fluid secretion, we were particularly interested in discerning how this diuretic activity is achieved given water cannot be moved across the tubule epithelia without a favourable osmotic gradient, provided largely through passage of inorganic ions. To further characterize the mechanism by which DTK elicits diuretic action on the MTs, we measured Na^+, K^+, and Cl^- conductance by isolated MTs. These *ex vivo* bioassay results revealed that DTK stimulates primarily Cl^- transport, which in turn increases transport of counter-ions (the cations Na^+ and K^+) and osmotically-obliged water leading to fluid secretion from the tubules. Our findings are consistent with previous studies in the mosquito *Ae. aegypti*, that found kinin-related neuropeptide hormones increase Cl^- conductance, which drives transepithelial transport of Na^+ and K^+ [53,54]. Similarly, the endocrine control of chloride conductance by MTs was shown in *Drosophila*, where leucokinin was found to stimulate chloride flux transcellularly through the chloride channel encoded by CG31116 [47]. Similarly, the biogenic amine tyramine, which is produced endogenously by MTs and thus elicits its control in a paracrine fashion, acts identically to

leucokinin leading to increased chloride permeability via stellate cells [43,44]. Therefore, DTK-1 stimulated fluid secretion likely involves increased chloride flux, which ultimately leads to depolarization (or collapsing) of the transepithelial membrane potential across the tubules [50]. The transport of chloride in the stellate cells is coupled with the active transport of cations in the principal cells, which in turn increases transepithelial transport across the tubules upon DTK stimulation with the help of transporters localized in the principal cells, including the Na^+-K^+-$2Cl^-$ (NKCC) and Na^+-K^+-ATPase (NKA) [43,44,53,55,56]. This movement of cations and anions may drive the movement of water via aquaporins localized in the stellate cells or the entomoglyceroporins in principal cells [5]. Therefore, this may explain the decrease in both Na^+ and K^+ concentration in response to DTK-1 stimulation due to the movement of water that may dilute cation concentration in the lumen of the tubules in the absence of co-activated cation-transporting principal cells. Future investigations should look at the activity and involvement of specific channels and transporters localized in the stellate cells when MTs are stimulated by DTK peptides. For example, previous studies have shown that stellate cell targeting diuretic factors (leucokinin and tyramine) utilize different chloride channels (*ClC-a* and *secCl*) either on the apical or basolateral membranes of the stellate cells in the MTs of the fruit fly [47,51]. The current results involving DTK-1 stimulated diuretic activity appears to demonstrate that chloride is the main driver for changes in transepithelial transport. In particular, chloride concentration and transport increased most significantly in response to DTK-1 in wildtype flies, corroborating that this peptidergic hormone is directly targeting the stellate cells having abundant chloride channels necessary for stimulated fluid secretion, and osmotically drawn water transport in the same (stellate) and adjacent (principal) cells [5]. Moreover, knockdown of DTKR specifically in the tubule stellate cells mainly affects the transport rate and not the concentration of Na^+ and K^+ while affecting both the transport rate and concentration of Cl^- ions. Therefore, the dramatic reduction in DTK-1 stimulated fluid secretion rates in *DTKR* knockdown flies is likely due to a drop in transepithelial ion transport resulting in reduced osmotic pressure that lowers the driving force for water transport.

Considering the functional consequences of DTKR knockdown on isolated MTs *ex vivo*, we questioned whether this reduced DTK-DTKR signaling involving the Malpighian 'renal' tubules alone might impact adult fly survival during starvation or desiccation stress, raising the notion that such signaling plays an important role in whole organismal physiology. Indeed, our data indicates that *DTKR* knockdown in the MTs prolonged survival of flies during desiccation stress, with no impact to survival under starvation stress. Despite converging on the same renal cell type and driving chloride conductance, one important difference between leucokinin and DTK signaling is that knockdown of LKR in stellate cells improves starvation tolerance but not desiccation tolerance [36], which represents the opposite phenotype following *DTKR* knockdown. DTKs are extensively distributed and synthesized in the central nervous system including over 150 neurons in the adult fly

brain, although it has been argued that this neural origin likely does not result in hormonal release since most DTK producing cells in the brain are interneurons [10]. A subset of ten DTK producing neurons in adult *D. melanogaster* brain (ITPn cells) are neurosecretory cells, that co-express ITP and sNPF [57]. Since ITP from ITPn functions as an anti-diuretic hormone, it seems counter-intuitive for diuretic DTK to be co-released into the circulation [58]. In support of this hypothesis, it has been argued that it is unlikely that DTK (or sNPF) derived from ITPn cells are released as circulating hormones but instead might act locally as neuromodulators in other brain regions or presynaptically on ITPn axon terminals [10,23]. Interestingly, most recent evidence suggests that this speculation of local release might be incorrect since ITP producing neurosecretory cells do not form synapses in the brain [58]. On the other hand, strong evidence supports that the midgut is responsible for endocrine action of DTKs in circulation as intestinal release has been demonstrated in other insects, including locusts and cockroaches [32]. Circulating DTKs acting as hormones target, as confirmed herein, DTKR expressed in stellate cells of the MTs triggering ion and fluid transport, which may lead to greater water loss from the organism. Relatedly, *DTKR* knockdown solely in the stellate cells of the MTs was sufficient to reduce water loss, providing these animals with greater tolerance of desiccation stress. These results are consistent with an earlier study where knockdown of diuretic hormones in the fruit fly, including the leucokinin hormone, resulted in increased survival during desiccation stress [59]. Interestingly, internal stimuli may be integrated differently in relation to DTK signaling and it may depend on the source of this neuropeptide since desiccation stress led to increased water loss when *Dtk* was diminished in a subset of DTK producing neurons in the brain, resulting in increased desiccation sensitivity [57], the opposite phenotype to what we observed when DTKR was reduced exclusively in stellate cells. This could be explained by considering brain-derived DTKs in ITPn [57,58] might affect other hormones acting on the MTs and other organs controlling hydromineral homeostasis, whereas DTKR knockdown exclusively in stellate cells reveals a more direct phenotype of altered DTK/DTKR signaling.

Conclusion

In summary, our data confirm that DTKs regulate fluid and ion transport by acting on the stellate cells of the MTs in the adult fruit fly, *D. melanogaster*. Although the source of circulating DTKs acting on the MTs remains unconfirmed in flies, they are likely released from enteroendocrine cells in the midgut as established in other insects, including cockroach and locust [20,32]. DTKs are localized in enteroendocrine cells in the midgut and may regulate lipid metabolism, especially during times of starvation [24]. Indeed, this earlier study established that tachykinins derived from the midgut regulate feeding-related physiological processes, while tachykinins derived from the nervous system govern behaviour [24]. Overall, this suggests that tachykinins could be released in response to physiological stress in the fly and future studies should confirm the source of hormonal DTKs along with stimuli that control their release. To conclude, DTKs are essential for osmo-

regulation in *D. melanogaster* given our findings reveal these peptide hormones contribute towards organismal ion-water balance by targeting DTKR expressed in stellate cells of the MTs to trigger diuresis. DTK/DTKR diuretic signaling may contribute towards hydromineral homeostasis, but considering its earlier documented involvement in increasing resistance to starvation stress [48], these two pathways may converge to coordinate metabolic and osmoregulatory demands of the fly, particularly in response to nutritional and desiccation stress.

References

1. Chown SL, Sørensen JG, Terblanche JS. Water loss in insects: An environmental change perspective. Journal of Insect Physiology. 2011. doi:10.1016/j.jinsphys.2011.05.004
2. Leopold P, Perrimon N. *Drosophila* and the genetics of the internal milieu. Nature. 2007. doi:10.1038/nature06286
3. Benoit JB. Water management by dormant insects: comparisons between dehydration resistance during summer aestivation and winter diapause. Progress in molecular and subcellular biology. 2010. pp. 209–229. doi:10.1007/978-3-642-02421-4_10
4. Beyenbach KW, Skaer H, Dow JAT. The developmental, molecular, and transport biology of Malpighian tubules. Annual Review of Entomology. 2010. pp. 351–374. doi:10.1146/annurev-ento-112408-085512
5. Cabrero P, Terhzaz S, Dornan AJ, Ghimire S, Holmes HL, Turin DR, et al. Specialized stellate cells offer a privileged route for rapid water flux in *Drosophila* renal tubule. Proceedings of the National Academy of Sciences. 2020;117: 1779–1787. doi:10.1073/pnas.1915943117
6. Hartenstein V. The neuroendocrine system of invertebrates: A developmental and evolutionary perspective. Journal of Endocrinology. 2006. doi:10.1677/joe.1.06964
7. Terhzaz S, O'Connell FC, Pollock VP, Kean L, Davies SA, Veenstra JA, et al. Isolation and characterization of a leucokinin-like peptide of *Drosophila melanogaster*. Journal of Experimental Biology. 1999;202. doi:10.1242/jeb.202.24.3667
8. Audsley N, McIntosh C, Phillips JE. Isolation of a neuropeptide from locust corpus cardiacum which influences ileal transport. J Exp Biol. 1992;173. doi:10.1242/jeb.173.1.261
9. Coast GM, Orchard I, Phillips JE, Schooley DA. Insect diuretic and antidiuretic hormones. Advances in Insect Physiology. 2002. doi:10.1016/s0065-2806(02)29004-9
10. Nässel DR, Zandawala M, Kawada T, Satake H. Tachykinins: Neuropeptides That Are Ancient, Diverse, Widespread and Functionally Pleiotropic. Frontiers in Neuroscience. 2019. doi:10.3389/fnins.2019.01262
11. Schoofs L, Holman GM, Hayes TK, Nachman RJ, De Loof A. Locustatachykinin I and II, two novel insect neuropeptides with homology to peptides of the vertebrate tachykinin family. FEBS Lett. 1990;261: 397–401. doi:10.1016/0014-5793(90)80601-E
12. Schoofs L, Holman GM, Hayes TK, Kochansky JP, Nachman RJ, De Loof A. Locustatachykinin III and IV: two additional insect neuropeptides with homology to peptides of the vertebrate tachykinin family. Regul Pept. 1990;31: 199–212. doi:10.1016/0167-0115(90)90006-I
13. Muren JE, Nässel DR. Isolation of five tachykinin-related peptides from the midgut of the cockroach *Leucaphaea maderae*: Existence of N-terminally extended isoforms. Regul Pept. 1996;65. doi:10.1016/0167-0115(96)00092-4
14. Winther ÅME, Muren JE, Lundquist CT, Osborne RH, Nässel DR. Characterization of actions of *Leucophaea* tachykinin-related peptides (LemTRPs) and proctolin on cockroach hindgut contractions. Peptides (NY). 1998;19. doi:10.1016/S0196-9781(97)00463-4
15. Coast GM. The influence of neuropeptides on Malpighian tubule writhing and its significance for excretion. Peptides (NY). 1998;19. doi:10.1016/S0196-9781(97)00461-0
16. Skaer NJV, Nässel DR, Maddrell SHP, Tublitz NJ. Neurochemical fine tuning of a peripheral tissue: Peptidergic and aminergic regulation of fluid secretion by Malpighian tubules in the tobacco hawkmoth *M. sexta*. Journal of Experimental Biology. 2002;205. doi:10.1242/jeb.205.13.1869
17. Johard HAD, Coast GM, Mordue W, Nässel DR. Diuretic action of the peptide locustatachykinin I: Cellular localisation and effects on fluid

secretion in Malpighian tubules of locusts. Peptides. 2003. doi:10.1016/j.peptides.2003.08.012

18. Torfs H, Shariatmadari R, Guerrero F, Parmentier M, Poels J, Van Poyer W, et al. Characterization of a receptor for insect tachykinin-like peptide agonists by functional expression in a stable *Drosophila* Schneider 2 cell line. J Neurochem. 2000;74: 2182–2189. doi:10.1046/j.1471-4159.2000.0742182.x

19. Birse RT, Johnson EC, Taghert PH, Nässel DR. Widely distributed Drosophila G-protein-coupled receptor (CG7887) is activated by endogenous tachykinin-related peptides. J Neurobiol. 2006;66: 33–46. doi:10.1002/neu.20189

20. Siviter RJ, Coast GM, Winther ÅME, Nachman RJ, Taylor CAM, Shirras AD, et al. Expression and functional characterization of a *Drosophila* neuropeptide precursor with homology to mammalian preprotachykinin A. Journal of Biological Chemistry. 2000;275: 23273–23280. doi:10.1074/jbc.M002875200

21. Winther ÅME, Siviter RJ, Isaac RE, Predel R, Nässel DR. Neuronal expression of tachykinin-related peptides and gene transcript during postembryonic development of *Drosophila*. Journal of Comparative Neurology. 2003;464: 180–196. doi:10.1002/cne.10790

22. Winther ÅME, Acebes A, Ferrús A. Tachykinin-related peptides modulate odor perception and locomotor activity in *Drosophila*. Molecular and Cellular Neuroscience. 2006;31: 399–406. doi:10.1016/j.mcn.2005.10.010

23. Kahsai L, Kapan N, Dircksen H, Winther ÅME, Nässel DR. Metabolic stress responses in *Drosophila* are modulated by brain neurosecretory cells that produce multiple neuropeptides. PLoS One. 2010;5. doi:10.1371/journal.pone.0011480

24. Song W, Veenstra JA, Perrimon N. Control of lipid metabolism by tachykinin in *Drosophila*. Cell Rep. 2014;9. doi:10.1016/j.celrep.2014.08.060

25. Monnier D, Colas JF, Rosay P, Hen R, Borrelli E, Maroteaux L. NKD, a developmentally regulated tachykinin receptor in *Drosophila*. Journal of Biological Chemistry. 1992;267: 1298–1302. doi:10.1016/s0021-9258(18)48429-3

26. Li XJ, Wolfgang W, Wu YN, North RA, Forte M. Cloning, heterologous expression and developmental regulation of a *Drosophila* receptor for tachykinin-like peptides. EMBO Journal. 1991;10: 3221–3229. doi:10.1002/j.1460-2075.1991.tb04885.x

27. Poels J, Birse RT, Nachman RJ, Fichna J, Janecka A, Vanden Broeck J, et al. Characterization and distribution of NKD, a receptor for *Drosophila* tachykinin-related peptide 6. Peptides (NY). 2009;30: 545–556. doi:10.1016/j.peptides.2008.10.012

28. Jiang H, Lkhagva A, Daubnerová I, Chae HS, Šimo L, Jung SH, et al. Natalisin, a tachykinin-like signaling system, regulates sexual activity and fecundity in insects. Proc Natl Acad Sci U S A. 2013;110. doi:10.1073/pnas.1310676110

29. Tsuji M, Nishizuka Y, Emoto K. Threat gates visual aversion via theta activity in Tachykinergic neurons. Nat Commun. 2023;14. doi:10.1038/s41467-023-39667-z

30. Wohl MP, Liu J, Asahina K. *Drosophila* Tachykininergic Neurons Modulate the Activity of Two Groups of Receptor-Expressing Neurons to Regulate Aggressive Tone. Journal of Neuroscience. 2023;43. doi:10.1523/JNEUROSCI.1734-22.2023

31. Vanden Broeck J, Torfs H, Poels J, Van Poyer W, Swinnen E, Ferket K, et al. Tachykinin-like peptides and their receptors. A review. Annals of the New York Academy of Sciences. 1999. doi:10.1111/j.1749-6632.1999.tb07907.x

32. Winther ÅME, Nässel DR. Intestinal peptides as circulating hormones: Release of tachykinin-related peptide from the locust and cockroach midgut. Journal of Experimental Biology. 2001;204: 1269–1280. doi:10.1242/jeb.204.7.1269

33. Coast GM, Orchard I, Phillips JE, Schooley DA. Insect diuretic and antidiuretic hormones. Adv In Insect Phys. 2002;29: 279–409. doi:10.1016/S0065-2806(02)29004-9

34. Radford JC, Davies SA, Dow JA. Systematic G-protein-coupled receptor analysis in *Drosophila melanogaster* identifies a leucokinin receptor with novel roles. J Biol Chem. 2002/08/07. 2002;277: 38810–38817. doi:10.1074/jbc.M203694200 M203694200 [pii]

35. Johnson EC, Bohn LM, Taghert PH. *Drosophila* CG8422 encodes a functional diuretic hormone receptor. Journal of Experimental Biology. 2004;207: 743–748. doi:10.1242/JEB.00818

36. Cannell E, Dornan AJ, Halberg KA, Terhzaz S, Dow JAT, Davies SA. The corticotropin-releasing factor-like diuretic hormone 44 (DH44) and kinin neuropeptides modulate desiccation and starvation toler-

ance in *Drosophila melanogaster*. Peptides (NY). 2016;80: 96. doi:10.1016/J.PEPTIDES.2016.02.004

37. Johnson EC, Shafer OT, Trigg JS, Park J, Schooley DA, Dow JA, et al. A novel diuretic hormone receptor in *Drosophila*: evidence for conservation of CGRP signaling. Journal of Experimental Biology. 2005;208: 1239–1246. doi:10.1242/JEB.01529

38. Terhzaz S, Cabrero P, Robben JH, Radford JC, Hudson BD, Milligan G, et al. Mechanism and function of *Drosophila* capa GPCR: a desiccation stress-responsive receptor with functional homology to human neuromedinU receptor. PLoS One. 2012;7: e29897. doi:10.1371/journal.pone.0029897

39. Sözen MA, Armstrong JD, Yang M, Kaiser K, Dow JAT. Functional domains are specified to single-cell resolution in a *Drosophila* epithelium. Proc Natl Acad Sci U S A. 1997;94. doi:10.1073/pnas.94.10.5207

40. Weaver LN, Ma T, Drummond-Barbosa D. Analysis of Gal4 expression patterns in adult *Drosophila* females. G3: Genes, Genomes, Genetics. 2020;10. doi:10.1534/g3.120.401676

41. Xu J, Liu Y, Li H, Tarashansky AJ, Kalicki CH, Hung RJ, et al. Transcriptional and functional motifs defining renal function revealed by single-nucleus RNA sequencing. Proc Natl Acad Sci U S A. 2022;119. doi:10.1073/pnas.2203179119

42. Radford JC, Davies SA, Dow JAT. Systematic G-protein-coupled receptor analysis in *Drosophila melanogaster* identifies a leucokinin receptor with novel roles. Journal of Biological Chemistry. 2002;277. doi:10.1074/jbc.M203694200

43. Blumenthal EM. Regulation of chloride permeability by endogenously produced tyramine in the *Drosophila* Malpighian tubule. Am J Physiol Cell Physiol. 2003;284. doi:10.1152/ajpcell.00359.2002

44. Cabrero P, Richmond L, Nitabach M, Davies SA, Dow JAT. A biogenic amine and a neuropeptide act identically: Tyramine signals through calcium in *Drosophila* tubule stellate cells. Proceedings of the Royal Society B: Biological Sciences. 2013;280. doi:10.1098/rspb.2012.2943

45. Ramsay JA. Active transport of water by the Malpighian tubules of the stick insect, *Dixippus morosus* (Orthoptera, Phasmidae). J Exp Biol. 1954;31: 104–113.

46. Lajevardi A, Sajadi F, Donini A, Paluzzi JP V. Studying the Activity of Neuropeptides and Other Regulators of the Excretory System in the Adult Mosquito. JoVE (Journal of Visualized Experiments). 2021;2021: e61849. doi:10.3791/61849

47. Cabrero P, Terhzaz S, Romero MF, Davies SA, Blumenthal EM, Dow JAT. Chloride channels in stellate cells are essential for uniquely high secretion rates in neuropeptide stimulated *Drosophila* diuresis. Proc Natl Acad Sci U S A. 2014;111: 14301–14306. doi:10.1073/pnas.1412706111

48. Söderberg JAE, Birse RT, Nässel DR. Insulin production and signaling in renal tubules of *Drosophila* is under control of tachykinin-related peptide and regulates stress resistance. PLoS One. 2011;6. doi:10.1371/journal.pone.0019866

49. Cabrero P, Radford JC, Broderick KE, Costes L, Veenstra JA, Spana EP, et al. The Dh gene of *Drosophila melanogaster* encodes a diuretic peptide that acts through cyclic AMP. J Exp Biol. 2002/11/15. 2002;205: 3799–3807. doi:10.1242/jeb.205.24.3799

50. O'Donnell MJ, Rheault MR, Davies SA, Rosay P, Harvey BJ, Maddrell SHP, et al. Hormonally controlled chloride movement across *Drosophila* tubules is via ion channels in stellate cells. Am J Physiol Regul Integr Comp Physiol. 1998;274. doi:10.1152/ajpregu.1998.274.4.r1039

51. Feingold D, Knogler L, Starc T, Drapeau P, O'Donnell MJ, Nilson LA, et al. secCl is a cys-loop ion channel necessary for the chloride conductance that mediates hormone-induced fluid secretion in *Drosophila*. Sci Rep. 2019;9. doi:10.1038/s41598-019-42849-9

52. Im SH, Takle K, Jo J, Babcock DT, Ma Z, Xiang Y, et al. Tachykinin acts upstream of autocrine Hedgehog signaling during nociceptive sensitization in *Drosophila*. Elife. 2015;4. doi:10.7554/eLife.10735

53. Pannabecker TL, Hayest TK, Beyenbach KW. Regulation of epithelial shunt conductance by the peptide leucokinin. J Membr Biol. 1993;132: 63–76. doi:10.1007/BF00233052

54. Hayes TK, Mark Holman G, Pannabecker TL, Wright MS, Strey AA, Nachman RJ, et al. Culekinin depolarizing peptide: a mosquito leucokinin-like peptide that influences insect Malpighian tubule ion transport. Regul Pept. 1994;52. doi:10.1016/0167-0115(94)90058-2

55. Rodan AR, Baum M, Huang CL. The *Drosophila* NKCC Ncc69 is required for normal renal tubule function. Am J Physiol Cell Physiol. 2012;303. doi:10.1152/ajpcell.00201.2012

56. Linton SM, O'Donnell MJ. Contributions of K+:Cl- cotransport and Na+/K+-ATPase to basolateral ion transport in Malpighian tubules of *Drosophila melanogaster*. Journal of Experimental Biology. 1999;202. doi:10.1242/jeb.202.11.1561

57. Kahsai L, Kapan N, Dircksen H, Winther ÅME, Nässel DR. Metabolic Stress Responses in *Drosophila* Are Modulated by Brain Neurosecretory Cells That Produce Multiple Neuropeptides. PLoS One. 2010;5: e11480. doi:10.1371/JOURNAL.PONE.0011480

58. Gera J, Agard M, Nave H, Sajadi F, Thorat L, Kondo S, et al. Anti-diuretic hormone ITP signals via a guanylate cyclase receptor to modulate systemic homeostasis in *Drosophila*. bioRxiv. 2024; 2024.02.07.579245. doi:10.1101/2024.02.07.579245

59. Zandawala M, Marley R, Davies SA, Nässel DR. Characterization of a set of abdominal neuroendocrine cells that regulate stress physiology using colocalized diuretic peptides in *Drosophila*. Cellular and Molecular Life Sciences. 2018;75. doi:10.1007/s00018-017-2682-y

60. Halberg KA, Rainey SM, Veland IR, Neuert H, Dornan AJ, Klämbt C, et al. The cell adhesion molecule Fasciclin2 regulates brush border length and organization in *Drosophila* renal tubules. Nat Commun. 2016;7. doi:10.1038/ncomms11266

61. Zandawala M, Nguyen T, Segura MB, Johard HAD, Amcoff M, Wegener C, et al. A neuroendocrine pathway modulating osmotic stress in *Drosophila*. PLoS Genet. 2021;17. doi:10.1371/journal.pgen.1009425

62. Li H, Janssens J, de Waegeneer M, Kolluru SS, Davie K, Gardeux V, et al. Fly Cell Atlas: A single-nucleus transcriptomic atlas of the adult fruit fly. Science (1979). 2022;375. doi:10.1126/science.abk2432

63. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis of multimodal single-cell data. Cell. 2021;184. doi:10.1016/j.cell.2021.04.048

64. Ignell R, Root CM, Birse RT, Wang JW, Nässel DR, Winther ÅME. Presynaptic peptidergic modulation of olfactory receptor neurons in *Drosophila*. Proc Natl Acad Sci U S A. 2009;106. doi:10.1073/pnas.0813004106

65. Birse RT, Söderberg JAE, Luo J, Winther ÅME, Nässel DR. Regulation of insulin-producing cells in the adult *Drosophila* brain via the tachykinin peptide receptor DTKR. Journal of Experimental Biology. 2011;214: 4201–4208. doi:10.1242/jeb.062091

66. Ponton F, Chapuis MP, Pernice M, Sword GA, Simpson SJ. Evaluation of potential reference genes for reverse transcription-qPCR studies of physiological responses in *Drosophila melanogaster*. J Insect Physiol. 2011;57. doi:10.1016/j.jinsphys.2011.03.014

67. Sajadi F, Uyuklu A, Paputsis C, Lajevardi A, Wahedi A, Ber LT, et al. CAPA neuropeptides and their receptor form an anti-diuretic hormone signaling system in the human disease vector, *Aedes aegypti*. Sci Rep. 2020;10. doi:10.1038/s41598-020-58731-y

68. Ramsay JA. Active Transport of Water by the Malpighian Tubules of the Stick Insect, *Dixippus morosus* (Orthoptera, Phasmidae) . Journal of Experimental Biology. 1954;31. doi:10.1242/jeb.31.1.104

69. Vanderveken M, O'Donnell MJ. Effects of diuretic hormone 31, drosokinin, and allatostatin a on transepithelial K+ transport and contraction frequency in the midgut and hindgut of larval *Drosophila melanogaster*. Arch Insect Biochem Physiol. 2014;85. doi:10.1002/arch.21144

70. Rheault MR, O'Donnell MJ. Organic cation transport by Malpighian tubules of *Drosophila melanogaster*: Application of two novel electrophysiological methods. Journal of Experimental Biology. 2004;207. doi:10.1242/jeb.01003

71. Donini A, O'Donnell MJ, Orchard I. Differential actions of diuretic factors on the Malpighian tubules of *Rhodnius prolixus*. Journal of Experimental Biology. 2008;211. doi:10.1242/jeb.011882

72. Paluzzi JP V., Naikkhwah W, O'Donnell MJ. Natriuresis and diuretic hormone synergism in *R. prolixus* upper Malpighian tubules is inhibited by the anti-diuretic hormone, RhoprCAPA-α2. J Insect Physiol. 2012;58. doi:10.1016/j.jinsphys.2011.11.020

73. Liao S, Post S, Lehmann P, Veenstra JA, Tatar M, Nässel DR. Regulatory Roles of *Drosophila* Insulin-Like Peptide 1 (DILP1) in Metabolism Differ in Pupal and Adult Stages. Front Endocrinol (Lausanne). 2020;11. doi:10.3389/fendo.2020.00180

Acknowledgements

The authors would like to thank Dr. Kenneth Halberg (University of Copenhagen) and Dr. Laura Nilson (McGill University) for providing the following fly lines: *uro*-GAL4 and *c724*-GAL4, respectively. This work was supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant and an Ontario Ministry of Research Innovation Early Researcher Award to JPP.

Author contributions

M.A. and J.P.P. conceived the study. J.P.P. supervised the project. M.A. performed the experimental work and analyzed the data. M.Z. performed computational analyses. M.A. wrote the first draft of the manuscript while M.Z. and J.P.P. provided feedback and edited the manuscript. All authors read, provided feedback, and approved the final manuscript.

Competing interest statement

The authors declare we have no competing interests.

Materials and Methods

Fly lines and husbandry

Fly lines were maintained at room temperature with a 12hr:12hr light:dark cycle. The wild-type (w^{1118}) flies, GAL4 and UAS fly lines used in this study are listed in Table 1. All fly lines were reared in vials containing food prepared as follows: 100 g/L sucrose, 50 g/L yeast, 12 g/L agar, 3 ml/L propionic acid, and 1.5 g/L nipagin. All experiments utilized 5-7 day-old (post-eclosion) adult females.

Table 1. Fly lines used in the study.

Fly lines	Inserted on chromosome	Role	Source or reference	Stock number
w^{1118} (Wild-type)	-----	Wild-type (control)	BDSC	5905
TkR99D (DTKR) GAL4	III	Expresses GAL4 in TkR99D pattern.	BDSC	76208
uro-GAL4 (PC GAL4)	II	Expresses GAL4 in the pattern of the urate oxidase gene in principal cells of the Malpighian tubules	Dr. Kenneth Halberg, [60]	--------
c724-GAL4 (SC GAL4)	II	Expresses GAL4 in the stellate cells of the Malpighian tubules	Dr. Laura Nilson, [51]	--------
UAS-*DTKR*-RNAi (RNAi 1)	III	UAS-DTKR-RNAi 1 line that encodes for the DTKR-RNAi construct	BDSC	27513
UAS-*DTKR*-RNAi (RNAi 2)	II	UAS-DTKR-RNAi 2 line that encodes for the DTKR-RNAi construct	BDSC	55732
JFRC81-10xUAS-IVS-Syn21-GFP-p10	-----	GFP is strongly expressed in cells where GAL4 has been expressed.	[61]	---------

Single-nucleus RNA-sequencing Analysis

Single-nucleus transcriptomes of Malpighian tubules were mined using the FlyCellAtlas dataset generated earlier [62]. The parameters used to identify Tkr99D positive stellate cells were: TkR99D > 0 & annotation == "adult Malpighian tubule stellate cell of main segment". All analyses were performed in R-Studio (v2022.02.0) using the Seurat package (v4.1.1 [63]). Heatmaps were constructed using the pheatmap package. Code available upon request.

Fluorescence microscopy

Fluorescence microscopy was performed to visualize GFP expression in the MTs. The JFRC81-10xUAS-IVS-Syn21-GFP-p10 (GFP) fly line was crossed with the following GAL4 lines: *TkR99D* (*DTKR*)-GAL4, *uro*-GAL4, and *c724*-GAL4. The MTs were dissected and isolated from the transgenic flies in phosphate-buffered saline (PBS). The guts were then fixed with 4% paraformaldehyde for 1 hour on the rocker at room temperature followed by three washes each for 10-15 minutes with PBS. The guts were mounted in 50% glycerol and visualized with the EVOS FL Auto live-cell imaging system (Life Technologies, Burlington, ON).

RT-qPCR

RT-qPCR was used to measure knockdown efficiency of DTKR by RNAi in fly MTs. Total RNA was extracted from 30-35 sets of MTs isolated from 5-7-day old female flies in five independent biological replicates using the Monarch Total RNA Miniprep kit (New England BioLabs, Whitby, ON, Canada). Total RNA was quantified using a UV spectrophotometer (Synergy 2 microplate reader, BioTek, Winooski, VT) and cDNA synthesis was performed using the iScript Reverse Transcription Supermix (Bio-Rad, Mississauga, ON, Canada). RT-qPCR was set up with quadruplicate technical replicates using PowerUP™ SYBR® Green Master Mix (Applied Biosystems, Carlsbad, CA, USA). The template cDNA was diluted 5-fold and used for RT-qPCR using a StepOnePlus Real-Time PCR System (Applied Biosystems, Carlsbad, CA, USA) that ran for 40 cycles: denaturation (95 °C), annealing (60 °C), and extension (72 °C). Forward (GAGTAAGCGAAGGGTGGTGAAG) and reverse (GAACGGCAGCCAGCAGAT) gene-specific primers described previously [64,65] were used to quantify DTKR expression in the MTs. Ribosomal protein L32 (RpL32) and alpha-tubulin were used as the reference genes for the RT-qPCR, as previously described [66]. Standard curves were obtained using serial dilutions of the MTs cDNA template from female w^{1118} yielding primer efficiencies ranging from 90-103%. For each replicate, the ΔΔCt method was used to determine the relative quantification (RQ) value, which was normalized to the geometric mean of RpL32 and alpha-tubulin reference genes [67]. Statistical analysis was performed on the log-transformed data.

Malpighian tubules secretion assay

The fluid secretion assay was adapted from Ramsay [68]. Female adult fruit flies were isolated and dissected in *Drosophila* saline (mmol l⁻¹): 6.8 glutamine, 5.6 glucose, 4.8 MOPS, 8.3 NaH_2PO_4, 11.9 $NaHCO_3$, 4.9 $MgCl_2$ (hexahydrate), 6.9 $CaCl_2$ (dihydrate), 13.4 KCl, 17.1 NaCl that was titrated to pH 7 [69]. The MTs were placed into small wells in a Sylgard-lined Petri dish containing 20µL of the DTK treatment in a 1:1 ratio of *Drosophila* saline: Schneider's Insect Medium (Sigma-Aldrich, Oakville, ON). The MTs were treated with 10^{-4}-10^{-13} M DTK-1 (APTSSFIGMR-amide) to determine the full dose-response curve and 10^{-7} M DTK-1 (~EC_{50}) was the selected dose to test out secretion rates from knockdown flies as identical secretion rates were observed when MTs were treated with the same concentration of other DTK isoforms, including DTK-2 and DTK-3. The proximal end of the tubule was wrapped around a minutien pin to measure the urine droplet from the MTs after a 60 min incubation. After the treatment period, the urine droplet was measured using a calibrated eyepiece micrometer from the dissecting microscope to calculate the fluid secretion rate (nL min⁻¹) with the following equation: secretion rate = $\frac{V}{\text{secretion time}}$, where secretion time refers to the 60-minute incubation, and V refers to the volume of the secreted droplet.

Use of Ion-selective microelectrodes (ISME) to measure ion concentration from isolated tubules

Ion-selective microelectrodes (ISME) were used to measure Na⁺, K⁺, and Cl⁻ concentrations from MTs isolated from wild-type (w^{1118}) and transgenic flies treated with DTK-1 (treatment) and 0.5% DMSO (control as DTK stocks were made up in DMSO). Glass capillaries for microelec-

trodes (TW150-4, World Precision Instruments, Sarasota, FL, USA) and reference electrodes (1B200F-4, World Precision Instruments, Sarasota, FL, USA) were prepared using a Sutter P-97 Flaming Brown pipette puller (Sutter Instruments, San Raffael, CA, USA). The ion-selective microelectrodes were silanized using N, N-dimethyltrimethylsilylamine (Sigma-Aldrich, Oakville, ON) as previously described [46]. The reference electrode was backfilled with 500 mmol l⁻¹ KCl. The Na⁺-selective microelectrode was backfilled with 100 mmol l⁻¹ NaCl, while the K⁺-selective microelectrode was backfilled with 100 mmol l⁻¹ KCl, and the Cl⁻-selective microelectrode was backfilled with 150 mmol l⁻¹ KCl. The tip of the ion-selective microelectrodes were forward-filled with an ion-specific ionophore: Na⁺ ionophore (sodium ionophore II cocktail A, Fluka), K⁺ ionophore (potassium ionophore I cocktail B, Fluka), and Cl⁻ ionophore (Chloride ionophore I - Cocktail A, Sigma-Aldrich). The tips of the electrodes were coated with polyvinyl chloride dissolved in tetrahydrofuran (Sigma-Aldrich) to prevent the movement of the ionophore when submerged in the mineral oil as described previously [46].

The silver chloride wires from the electrometer were inserted into the microelectrodes so that the differential voltage could be measured and recorded with the electrometer and a data acquisition system (Picolog for Windows, version 5.25.3). The ISME was calibrated in the following standard solutions: Na⁺ microelectrode, 200 mmol l⁻¹ NaCl and 20 mmol l⁻¹ NaCl +180 mmol l⁻¹ LiCl; K⁺ microelectrode, 150 mmol l⁻¹ KCl and 15 mmol l⁻¹ KCl + 135 mmol l⁻¹ LiCl; Cl⁻ microelectrode, 150 mmol l⁻¹ KCl and 15 mmol l⁻¹ KCl. The ISME and reference electrodes were positioned into the secreted droplet using the micromanipulators. The ion concentration was calculated using the equation previously described [70–72]: $[\text{ion}]_{sf} = [C] \times 10^{\Delta V/m}$, where $[C]$ is the concentration of the calibration solution in mmol l⁻¹, ΔV is the difference between the voltage recorded in the secreted fluid droplet and the voltage recorded in the calibration solution, and m is the voltage difference between the two standard calibrations (slope). Finally, the ion transport rates were calculated by multiplying the fluid secretion rates and the $[\text{ion}]_{sf}$.

Stress resistance assays

The stress assays were performed on 5-7 day-old female flies. For each of the four biological replicates, 15-20 flies were selected, and the number of surviving flies was recorded every 8 hours until all flies were dead. The desiccation assay consisted of placing 5-7 day-old female DTKR-RNAi flies and control flies into empty vials to deprive flies of food and water to induce desiccation [48,73]. The starvation assay consisted of placing 5-7 day-old female DTKR-RNAi flies and control flies into vials containing 1% aqueous agar (Fisher Scientific, USA) [48,73].

Statistical Analyses

All experimental data was graphed as mean±SEM. The data were analyzed using GraphPad Prism Version 8.0.0 (San Diego, California USA) to complete the appropriate statistical test such as the unpaired t-test, one-way ANOVA & two-way ANOVA with the addition of the post-hoc test (Tukeys, and Šídák's), and the Log-rank (Mantel-Cox) test.

Fig 1. Single-nucleus RNA sequencing analysis distinguishing the Malpighian tubules cell types and expression of hormone receptors and channels. (A) Expression of hormone receptors in single cell transcriptomes of Malpighian tubules. (B) Clustering of different cell types in Malpighian tubules. (C) Dotplot showing expression of hormone receptors in different cell types of Malpighian tubules. (D) Heatmap of all stellate cells showing expression of hormone receptors. Leucokinin receptor (Lkr) and Tkr99D show overlapping expression pattern. (E) Tubules isolated from TkR99D>JFRC81-10xUAS-IVS-Syn21-GFP-p10 (GFP) flies displayed high GFP expression in the star-shaped stellate cells (scale bar: 50 m). (F) Dot plot showing expression of diuretic hormone receptors and aquaporins in Tkr99D positive cells.

Fig 2. Secretion rates by in vitro isolated Malpighian tubules in response to different concentrations of Drosophila tachykinin-1 (DTK-1). The MTs isolated from wild-type (WT) flies (A) displayed a dose-dependent activation of fluid secretion as shown by the activity of various concentrations of DTK-1, where the EC_{50} was 1.126 x 10^{-7} M DTK-1 (assessed by log(agonist) vs. response (three parameters) non-linear regression). (B) Tubules treated with 10^{-7} M DTK-1 displayed significant stimulation of secretion compared to unstimulated and 10^{-9} M DTK-1 treatment, while displaying significantly lower secretion rates compared to 10^{-5} M DTK-1 ($p<0.05$, assessed by One-way ANOVA). (C) Tubules displayed identical significant increases in fluid secretion rates when treated with DTK-1, DTK-2, and DTK-3 at 10^{-7} M compared to unstimulated MTs ($p<0.05$, assessed by One-way ANOVA).

Fig 3. DTKR transcript levels in Malpighian tubules and fluid secretion rates isolated from different crosses to assess DTKR knockdown using either a principal or stellate cell GAL4 driver. (A) Tubules isolated from *uro*-GAL4>JFRC81-10xUAS-IVS-Syn21-GFP-p10 (GFP) flies displayed high GFP expression in the principal cells (scale bar: 400μm). (B) High GFP expression in the stellate cells of the tubules isolated from *c724*-GAL4>GFP flies (scale bar: 400μm). (C) Malpighian tubules isolated from flies where knockdown was driven by the principal cell *uro*-GAL4 displayed no changes in DTKR transcript levels. (D) Tubules isolated from flies where knockdown was driven by the stellate cell *c724*-GAL4 displayed a significant reduction in DTKR abundance. Similar DTKR knockdown in stellate cells was achieved using two different DTKR RNAi lines ($p < 0.05$, assessed by two-way ANOVA). (E) & (F) Tubules isolated from knockdown flies driven by the *uro*-GAL4 displayed a significant increase in fluid secretion rates in response to DTK-1 ($p < 0.05$, assessed by two-way ANOVA). (G) & (H) Tubules isolated from *c724*-GAL4 and two UAS-DTKR RNAi crosses displayed significantly reduced stimulated secretion rates in response to DTK-1 treatment ($p < 0.05$, assessed by two-way ANOVA).

Fig 4. Ion concentration in the secreted fluid and transport rates from wild-type fly (WT) Malpighian tubules stimulated by *Drosophila* tachykinin-1. (A) Reduced Na$^+$ concentration in the secreted fluid and (B) increased Na$^+$ transport rate from tubules treated with DTK-1 ($p<0.05$, assessed by unpaired t-test). Similarly, (C) lower K$^+$ concentration in the secreted fluid and (D) elevated K$^+$ transport rates by tubules treated with the DTK-1 ($p<0.05$, assessed by unpaired t-test). (E) Cl$^-$ concentration in the secreted fluid and (F) Cl$^-$ transport rate significantly increased in MTs treated with DTK-1 ($p<0.05$, assessed by unpaired t-test).

Fig 5. Effects of *Drosophila* tachykinin-1 on the ion concentration in secreted fluid and their transport rates following knockdown of DTKR in stellate cells. (A) Na$^+$ concentration in secreted fluid of the DTKR knockdown MTs increased after DTK-1 treatment when compared to the unstimulated treatment whereas compared to the control flies there was no significance. (B) No changes in K$^+$ concentration in DTKR knockdown MTs after DTK-1 treatment while displaying significance with one of the control fly lines. (C) Na$^+$ transport rates significantly decreased compared to control flies. (D) A significant decrease in K$^+$ transport was observed in comparison to control flies. (E) Cl$^-$ concentration in the secreted fluid significantly decreased in DTKR knockdown MTs following DTK-1 treatment compared to control flies. Similarly, (F) Cl$^-$ transport rates also significantly decreased following DTK-1 treatment compared to control flies. All statistical comparisons were assessed by two-way ANOVA and Šídák's post-hoc test with p<0.05 considered significant.

Fig 6. DTKR knockdown DTKR knockdown enhances desiccation stress survival but does not influence survival in response to starvation stress conditions. (A) DTKR-RNAi flies driven by the stellate cell specific *c724*-GAL4 display significantly longer survival compared to the control flies under conditions of desiccation stress. (B) DTKR-RNAi did not influence survival of flies compared to the control flies in response to starvation stress. (A) & (B) data are presented as survival curves (assessed by Log-rank (Mantel-Cox) test).

Legend:
- $c724$-GAL4>UAS-$DTKR$ RNAi 1
- WT>UAS-$DTKR$ RNAi 1
- $c724$-GAL4>WT

A — Na⁺ concentration (mmol l⁻¹) — Na^+ concentration (mmol l^{-1}); x-axis: Unstimulated / 10^{-7} M DTK-1 for each genotype; labels a, bc, ab, bc, bc, c

B — K^+ concentration (mmol l^{-1}); labels a, ac, bc, b, ab, bc

C — Na^+ transport rate (pmol min^{-1}); labels a, a, a, b, a, b

D — K^+ transport rate (pmol min^{-1}); labels a, a, a, b, a, b

E — Cl^- concentration (mmol l^{-1}); labels a, a, a, b, a, b

F — Cl^- transport rate (pmol min^{-1}); labels a, a, a, b, a, b