ASTROGLIOSIS AND NEUROINFLAMMATION UNDERLIE SCOLIOSIS UPON CILIA DYSFUNCTION

Morgane Djebar[1], Isabelle Anselme[1], Guillaume Pezeron[2], Pierre-Luc Bardet[1], Yasmine Cantaut-Belarif[3], Alexis Eschstruth[1], Diego López Santos[1], Hélène Le Ribeuz[1], Arnim Jenett[4], Hanane Khoury[1], Joëlle Véziers[5], Caroline Parmentier[6], Aurélie Hirschler[7], Christine Carapito[7], Ruxandra Bachmann-Gagescu[8], Sylvie Schneider-Maunoury[1]* and Christine Vesque[1]*

Sorbonne Université, CNRS UMR7622, INSERM U1156, Institut de Biologie Paris Seine (IBPS) - Developmental Biology Unit, 75005, Paris, France
Molecular Physiology and Adaptation (PhyMA - UMR7221), Muséum National d'Histoire Naturelle, CNRS, Paris, France
Institut du Cerveau et de la Moelle épinière (ICM), Sorbonne Université, Inserm U 1127, CNRS UMR 7225, F-75013, Paris, France
TEFOR Paris-Saclay, CNRS UMS2010 / INRA UMS1451, Université Paris-Saclay.
Inserm UMR 1229, CHU Nantes PHU4 OTONN, SC3M facility, Inserm UMS 016, CNRS 3556, Université de Nantes, Nantes, France.
Sorbonne Université, CNRS UMR8246, INSERM U1130, Institut de Biologie Paris Seine (IBPS) – Neurosciences Paris Seine (NPS), 75005, Paris, France
Laboratoire de Spectrométrie de Masse Bio-Organique, IPHC, UMR 7178, Université de Strasbourg, CNRS, Infrastructure Nationale de Protéomique ProFI - FR2048, Strasbourg, France.
Institute of Medical Genetics, University of Zurich, Wagistrasse 12, 8952 Schlieren, Zurich, Switzerland; Institute of Molecular Life Sciences, University of Zurich, Winterthurerstrassse 190, 8057 Zurich, Switzerland
* Co-corresponding authors, co-last authors.
sylvie.schneider-maunoury@sorbonne-universite.fr
christine.vesque@sorbonne-universite.fr

ABSTRACT

Cilia defects lead to scoliosis in zebrafish, but the underlying pathogenic mechanisms are poorly understood and may diverge depending on the mutated gene. Here, we dissected the mechanisms of scoliosis onset in a zebrafish mutant for the *rpgrip1l* gene encoding a ciliary transition zone protein. *rpgrip1l* mutant fish developed scoliosis with near-total penetrance but asynchronous onset in juveniles. Taking advantage of this asynchrony, we found that curvature onset was preceded by ventricle dilations and was concomitant to the perturbation of Reissner fiber polymerization and to the loss of multiciliated tufts around the subcommissural organ. Rescue experiments showed that Rpgrip1l was exclusively required in *foxj1a*-expressing cells to prevent axis curvature. Genetic interactions investigations ruled out Urp1/2 levels as a main driver of scoliosis in *rpgrip1* mutants. Transcriptomic and proteomic studies identified neuroinflammation associated with increased Annexin levels as a potential mechanism of scoliosis development in *rpgrip1l* juveniles. Investigating the cell types associated with *annexin2* over-expression, we uncovered astrogliosis, arising in glial cells surrounding the diencephalic and rhombencephalic ventricles just before scoliosis onset and increasing with time in severity. Anti-inflammatory drug treatment reduced scoliosis penetrance and severity and this correlated with reduced astrogliosis and macrophage/microglia enrichment around the diencephalic ventricle. Mutation of the *cep290* gene encoding another transition zone protein also associated astrogliosis with scoliosis. Thus, we propose astrogliosis induced by perturbed ventricular homeostasis and associated with immune cell activation as a novel pathogenic mechanism of zebrafish scoliosis caused by cilia dysfunction.

INTRODUCTION

Idiopathic scoliosis (IS) is a 3D rotation of the spine without vertebral anomalies that affects about 3% of adolescents worldwide. Its etiology remained mysterious, mainly because of the lack of appropriate animal models and the complexity of its inheritance profile in large families presenting a severe scoliosis trait (Patten et al., 2015). Since the description of the first IS model in the zebrafish *ptk7* gene mutant (Hayes et al., 2014), a number of mutants for genes encoding ciliary proteins were shown to develop scoliosis at late larval and juvenile stage (4 to 12 weeks post-fertilization, wpf) without any vertebral fusion or fracture, highlighting the link between cilia function and straight axis maintenance in that species (Grimes et al., 2016; Wang et al., 2022).

Cilia are microtubular organelles with sensory and/or motile functions. Zebrafish mutants in genes involved in cilia motility or in intraflagellar transport display severe embryonic axis curvature and are usually not viable beyond larval stages (Grimes et al., 2016; Rose et al., 2020). In contrast, mutants in genes implicated in ciliary gating or ciliary trafficking encoding respectively components of the transition zone (TZ) or of the BBsome complex survive to adult stage and develop scoliosis with variable penetrance (Bentley-Ford et al., 2022; Masek et al., 2022; Song et al., 2020; Wang et al., 2022). The molecular basis for these differences is still poorly understood, even if progress has been made in our understanding of zebrafish scoliosis. Cilia-driven movements of the cerebrospinal fluid (CSF) are involved in zebrafish axis straightness, both in embryos and juveniles (Boswell and Ciruna, 2017; Grimes et al., 2016) and are tightly linked to the assembly and maintenance of the Reissner fiber (RF), a SCO-spondin polymer secreted by the subcommissural organ (SCO) and running down the brain and spinal cord CSF-filled cavities (Cantaut-Belarif et al., 2018). RF loss at embryonic stage in null *sspo* mutants generates ventral curvatures of the posterior axis of the body and is lethal while its loss at juvenile stage in hypomorphic *sspo* mutants triggers scoliosis with full penetrance (Cantaut-Belarif et al., 2018; Rose et al., 2020; Troutwine et al., 2020). Signaling downstream of the RF in embryos implicates Urp1 and Urp2, two neuropeptides of the Urotensin 2 family expressed in ventral CSF-contacting neurons (CSF-cNs), which act on dorsal muscles in embryos and larvae (Cantaut-Belarif et al., 2020; Gaillard et al., 2023; Lu

et al., 2020; Zhang et al., 2018). Their combined mutations or the mutation of their receptor gene *uts2r3* lead to scoliosis at adult stages (Bearce et al., 2022; Gaillard et al., 2023; Zhang et al., 2018). However, whether RF maintenance and Urp1/2 signaling are perturbed in juvenile scoliotic mutants of genes encoding TZ proteins ("TZ mutants") and how these perturbations are linked to scoliosis is unknown, especially because TZ mutants do not display any sign of embryonic or larval cilia motility defects (Wang et al., 2022). Moreover, two scoliotic models develop axial curvature while maintaining a RF showing that their curvature onset occurs independently of RF loss (Meyer-Miner et al., 2022; Xie et al., 2023). Finally, neuro-inflammation has been described in a small subset of zebrafish IS models for which anti-inflammatory/anti-oxidant treatments (with NAC or NACET) partially rescue scoliosis penetrance and severity (Rose et al., 2020; Van Gennip et al., 2018).

In this paper we dissect the mechanisms of scoliosis appearance in a novel deletion allele of the zebrafish TZ *rpgrip1l* gene. *RPGRIP1L* mutations have been found in neurodevelopmental ciliopathies, Meckel syndrome (MKS) and Joubert syndrome (JS) (Delous et al., 2007)(Arts et al., 2007) characterized by severe brain malformations. Other organs such as the kidney and liver are variably affected. Some JS patients also present with severe scoliosis, but this phenotypic trait is of variable penetrance (Delous et al., 2007)(Brancati et al., 2008). Functional studies performed in mouse and zebrafish have shown that *Rpgrip1l* is required for proper central nervous system patterning due to complex perturbations of Sonic Hedgehog signaling (in mice) and cilia planar positioning (in mice and zebrafish) (Andreu-Cervera et al., 2021; Mahuzier et al., 2012; Vierkotten et al., 2007). Zebrafish *rpgrip1l* mutants are viable and develop an axis curvature phenotype at juvenile stages with nearly full penetrance. Using a variety of approaches, we show that i) Rpgrip1l is required in ciliated cells surrounding ventricular cavities for maintaining a straight axis; ii) at juvenile stages, *rpgrip1l* mutants develop cilia defects associated with ventricular dilations and loss of the RF; iii) increased *urp1/2* expression in *rpgrip1l* mutants does not account for spine curvature defects; and iv) a pervasive astrogliosis process associated with neuroinflammation participates in scoliosis onset and progression. Finally, we show that astrogliosis is conserved in another zebrafish ciliary TZ mutant, *cep290,* identifying a novel mechanism involved in scoliosis upon ciliary dysfunction.

RESULTS

Rpgrip1l $^{-/-}$ fish develop scoliosis asynchronously at juvenile stage

To study the mechanisms of scoliosis appearance upon ciliary dysfunction in zebrafish, we produced a viable zebrafish deletion mutant in the *rpgrip1l* gene encoding a ciliary transition zone protein (*rpgrip1l$^\Delta$*, identified in ZFIN as *rpgrip1l^{bps1}*) Supplementary Figure S1A). We made a 26 kb deletion by CRISPR-Cas9 double-cut, which removes 22 out of the 26 exons of the protein (Supplementary Figure S1A) and triggers protein truncation at amino-acid (aa) position 88 out of 1256. 90-100% of *rpgrip1l$^{\Delta/\Delta}$* embryos and larvae were straight (0-10%, depending on the clutch, displayed a sigmoid curvature). They did not display any additional defects found in many ciliary mutant embryos or larvae such as randomized left-right asymmetry, kidney cysts or retinal anomalies (Bachmann-Gagescu et al., 2011; Kramer-Zucker et al., 2005; Tsujikawa and Malicki, 2004)(Figure 1A and Supplementary Figure S1B, D, E). Another allele with a frameshift mutation in exon 4 (*rpgrip1l^{ex4}*, identified in ZFIN as *rpgrip1l^{bps2}*) showed a similar phenotype (Supplementary Figure S1A, B). Considering the large size of the deletion, we considered the *rpgrip1l$^\Delta$* allele as a null and called it *rpgrip1l$^{-/-}$* thereafter. *rpgrip1l$^{-/-}$* animals developed scoliosis during juvenile stages. Scoliosis appearance was asynchronous between clutches, from 4 weeks post fertilization (wpf) (around 1 cm length) to 11 wpf (1.8- 2.3 cm), and also within a clutch. It started with slight upward bending of the tail (the *tail-up* phenotype) and progressed toward severe curvature (Figure 1A, B) with 90% penetrance in adults (100% in females and 80% in males) (Figure 1C). Micro-computed tomography (µCT) at two different stages (5 wpf and 23 wpf) confirmed that spine curvature was three-dimensional and showed no evidence of vertebral fusion, malformation or fracture (Figure 1D-G and Supplementary Movie 1).

Thus, the ciliary *rpgrip1l$^{-/-}$* mutant shows almost totally penetrant, late-onset juvenile scoliosis without vertebral anomalies, making it a valuable model for studying the etiology of human idiopathic scoliosis. Furthermore, the asynchronous onset of curvature is an asset for studying the chronology of early defects leading to scoliosis.

***Reintroducing* RPGRIP1L *in foxj1a lineages suppresses scoliosis in rpgrip1l*$^{-/-}$**

fish

Since *rpgrip1l* is ubiquitously expressed in zebrafish (Mahuzier et al., 2012), scoliosis could have different tissue origins, including a neurological origin as shown for the *ptk7* and *ktnb1* scoliotic fish (Meyer-Miner et al., 2022; Van Gennip et al., 2018). Indeed, no tissue-specific rescue has been performed in mutants for zebrafish ciliary TZ or BBS genes. We tested if introducing by transgenesis a tagged human RPGRIP1L protein under the control of the *col2a1a (Dale and Topczewski, 2011)* or *foxj1a* (Van Gennip et al., 2018) enhancers (Figure 1H) would reduce scoliosis penetrance and severity. The *foxj1a* enhancer drives expression in motile ciliated cells, among which ependymal cells lining CNS cavities (Van Gennip et al., 2018), while the *col2a1a* regulatory region drives expression in cartilage cells including intervertebral discs and semi-circular canals, tissues that have been proposed to be defective in mammalian scoliosis models (Hitier et al., 2015; Karner et al., 2015). After selecting F1 transgenic fish that expressed the Myc-tagged RPGRIP1L protein in the targeted tissue at 2.5 dpf, we introduced one copy of each transgene into the *rpgrip1l*$^{-/-}$ background. The Foxj1a:5XMycRPGRIP1L transgene was sufficient to fully suppress scoliosis in *rpgrip1l*$^{-/-}$ fish: none of the transgenic *rpgrip1l*$^{-/-}$ fish became scoliotic while virtually all the non-transgenic *rpgrip1l*$^{-/-}$ siblings did (Figure 1I, J). In contrast, RPGRIP1L expression triggered by the *col2a1a* promoter did not have any beneficial effect on scoliosis penetrance (Figure 1I, J). In conclusion, *rpgrip1l* expression in *foxj1a*-expressing cells is sufficient to maintain a straight axis.

Rpgrip1l*$^{-/-}$ *adult fish display cilia defects along CNS cavities

We observed normal cilia in the neural tube of *rpgrip1l*$^{-/-}$ embryos (Supplementary Figure S1C). In 3 month-adults, scanning electron microscopy (SEM) at the level of the rhombencephalic ventricle (RhV) showed cilia tufts of multiciliated ependymal cells (MCC) lining the ventricle in controls and straight *rpgrip1l*$^{-/-}$ fish (Supplementary Figure S1F-G, F'-G'). In scoliotic *rpgrip1l*$^{-/-}$ fish, cilia tufts were sparse and disorganized (Supplementary Figure S1H, H'). Cilia of mono-ciliated ependymal cells

were present in *rpgrip1l*[-/-] fish, some of them with irregular shape (Supplementary

Figure S1, F", G", H" and H"').

MCC defects at SCO level correlate with scoliosis appearance at juvenile

stages

We then took advantage of the asynchrony in scoliosis onset to correlate, at a given

stage, cilia defects at different levels of the CNS cavities with the spine curvature

status (straight or curved) of *rpgrip1l*[-/-] juveniles. We analyzed cilia defects in three

CNS regions implicated in scoliosis appearance: the spinal cord central canal (since

scoliosis develops in the spine), the forebrain choroid plexus (fChP) whose cilia

defects have been associated with scoliosis in *katnb1*[-/-] fish (Meyer-Miner et al.,

2022) and the subcommissural organ (SCO), which secretes SCO-spondin whose

absence leads to scoliosis (Rose et al., 2020; Troutwine et al., 2020).

In the spinal cord central canal of 8 wpf juvenile fish, cilia density was reduced to the

same extent in *rpgrip1l*[-/-] fish irrespective of the fish curvature status (Figure 2 A-C,

E). In addition, cilia length increased in straight mutants and was more heterogenous

than controls in scoliotic mutants (Figure 2 A'-C', D). Arl13b content appeared

severely reduced in mutants (Figure 2 A"-C"). In the fChP, mono- and multi-ciliated

cell populations were observed with a specific distribution along the anterior-posterior

axis, as previously described (D'Gama et al., 2021). At the level of the habenula

nuclei, cells of dorsal and ventral midline territories were monociliated, while lateral

cells presented multicilia tufts (Supplementary Figure S2 K-K"). *rpgrip1l*[-/-] juveniles

presented an incomplete penetrance of ciliary defects in the fChP, with no correlation

between cilia defects and the curvature status (Supplementary Figure S2, L-O").

The SCO lies at the dorsal midline of the diencephalic ventricle (DiV), under the

posterior commissure (PC) (Figure 2 G, H, Supplementary Figure 2A, C). Cilia in and

around the SCO have not been described previously in zebrafish. We showed that, in

control animals, Sco-spondin (Sspo) secreting cells varied in number along SCO

length, from 24 cells anteriorly to 6-8 cells posteriorly (Supplementary Figure 2 A, A',

C) and were monociliated at all anteroposterior levels (Figure 2G-H, Supplementary

Figure 2 A'). The Sspo protein produced from one copy of the transgene Scospondin-

GFP[ut24] (Troutwine et al., 2020) formed cytoplasmic granules around SCO cell nuclei,

under emerging primary cilia (Supplementary Figure 2A'). Conversely, MCCs lateral

to the SCO did not secrete Sspo and were prominent towards posterior (Figure 2H,
Supplementary Figure 2C), as observed in the rat brain (Collins and Woollam, 1979).
Monocilia of Sspo secreting cells were still present but appeared longer in straight
rpgrip1l[-/-] juveniles and could not be individualized anymore at anterior SCO level in
scoliotic fish: we observed a weaker and denser acetylated and glutamylated tubulin
staining that likely corresponded to longer intertwined cilia (Figure 2 I, I', K, K').
Lateral multicilia tufts were preserved in straight *rpgrip1l*[-/-] juveniles (n=5) (Figure 2 J)
but were missing at posterior SCO level in tail-up or scoliotic *rpgrip1l*[-/-] fish (n=7)
(Figure 2L).
Thus, *rpgrip1l* mutants develop both monocilia and multicilia defects at juvenile
stages, but cilia defects in the spinal cord and in the fChP do not correlate with axis
curvature status of the fish. By contrast, the loss of MCC cilia at the SCO strictly
correlates with scoliosis onset, suggesting a causal relationship.
178

Rpgrip1l[-/-] juveniles show ventricular dilations and loss of the Reissner fiber at scoliosis onset

Ciliary beating is an essential actor of CSF flow and of ventricular development in
zebrafish larvae (Kramer-Zucker et al., 2005) (Olstad et al., 2019). Thus, ciliary
defects in the brain and spinal cord of *rpgrip1l*[-/-] fish could lead to abnormal
ventricular and central canal volume and content. In the spinal cord, the lumen of the
central canal was enlarged at thoracic and lumbar levels in all *rpgrip1l*[-/-] mutant
juveniles, with no correlation to the axis curvature status (Figure 2 A-C; F). To
determine the timing of brain ventricular dilations relative to scoliosis, we analyzed
ventricular volume at the onset of spine curvature (Figure 3A-G). Ventricular
reconstruction was performed on cleared brains of 4 control and 4 *rpgrip1l*[-/-] (3 tail-up
and 1 straight) fish at 5 wpf, stained with ZO1 to highlight the ventricular border and
with DiI to outline brain shape, focusing on the RhV at the level of the posterior
midbrain and hindbrain (Figure 3A-D). We identified a significant increase in ventricle
volume in *rpgrip1l*[-/-] fish compared to controls, that was restricted to the ventral
regions of the RhV in regions ROI4.4 (green sphere in Figure 3C) and ROI6 (green
sphere in Figure 3D), as confirmed by numerical sections (Figure 3E, G). Ventricle

dilations were also present, although to a lesser extent, in the RhV of the unique
straight *rpgrip1l*[-/-] fish of this study (green squares and triangle in Figure 3F, G).
Defective ependymal ciliogenesis has been associated with abnormal CSF flow and
defective Reissner Fiber (RF) maintenance in several scoliotic zebrafish models
(Grimes et al., 2016)(Rose et al., 2020). The RF runs along the length of the body,
within brain ventricles and central canal (Figure 3H). It is mainly composed of Sspo
secreted by the SCO and floor plate (FP) and its loss at juvenile stages triggers
scoliosis in zebrafish (Rose et al., 2020)(Troutwine et al., 2020). To visualize the RF,
we used an antiserum that labels the RF in zebrafish embryos (Didier et al.,
1995)(Cantaut-Belarif et al., 2018) and we confirmed the results by introducing one
copy of the *sspo-GFP* knock-in allele, sco-spondin-GFP[ut24,] (Troutwine et al., 2020)
by genetic cross into *rpgrip1l*[-/-] animals. The RF formed normally in *rpgrip1l*[-/-]
embryos, as expected given the absence of embryonic curvature (Supplementary
Figure S1 C). To study its maintenance in *rpgrip1l*[-/-] juveniles, we immunostained
spinal cord longitudinal sections of 7-8 wpf curved or straight fish of the same clutch.
The RF was visualized as a 1 µm diameter rod in the central canal of the neural tube
and few dots were present in the apical cytoplasm of FP cells (Figure 3I) as well as in
the SCO of all controls (n=8, Figure 3L, O) and in ventricles caudal to the SCO, but
not rostral to it as expected (Supplementary Figure S2E, H). The RF was also
present in all *rpgrip1l*[-/-] straight fish (n=9, Figure 3J, Supplementary Figure S2I). In
contrast, in *rpgrip1l*[-/-] tail-up and scoliotic fish, the RF was absent at all axis levels and
a few SCO-spondin-positive debris were present in the central canal (n=8, Figure
3K). Transverse sections at the level of the SCO and fChP as well as caudal to the
SCO in severely scoliotic fish revealed abnormally packed material in the ventricles
(Figure 3N, Q, Supplementary Figure S2G, J). In juveniles with a slight tail-up
phenotype, SCO-secreting cells produced Sspo-GFP cytoplasmic granules as in
controls (Supplementary Figure S2B, B', n=7/7), and polymers of Sspo-GFP
accumulate in the adjacent DiV, which suggests that Sspo-GFP aggregates are
produced by the SCO.
Thus, our study uncovers a strict temporal correlation between the loss of MCC at
posterior SCO level, impaired RF polymerization and scoliosis onset in *rpgrip1l*[-/-]
juvenile fish.
229

Urp1/2 upregulation does not contribute to scoliosis penetrance or severity in rpgrip1l$^{-/-}$ fish

In scoliotic fish where the RF is not maintained at juvenile stage, *urp1* and *urp2* levels are downregulated (Rose et al., 2020). *urp1* and *urp2* encode peptides of the Urotensin 2 family, expressed in ventral spinal CSF-cNs (Parmentier et al., 2011; Quan et al., 2015). Their expression levels have to be finely tuned to keep a straight axis since their combined loss-of-function induces larval kyphosis that evolves into adult scoliosis (Bearce et al., 2022; Gaillard et al., 2023), while an increased dose of Urp1/2 peptides induces an upward curvature in embryos (Zhang et al., 2018) and juveniles (Gaillard et al., 2023; Lu et al., 2020). We therefore monitored via qRT-PCR *urp1 and urp2* expression levels on individual straight or scoliotic *rpgrip1l$^{-/-}$* juvenile and/or adult fish. *urp2* was the most highly expressed gene among all *urotensin 2* members at juvenile stage (Figure 4A, D, E and Supplementary Figure S3A-E. *urp2* expression was upregulated in 5 weeks juveniles, both in straight and scoliotic fish and its upregulation was maintained in adult scoliotic fish (Figure 4A).

Given the early and long-lasting upregulation of *urp1/2* in *rpgrip1l$^{-/-}$* fish, we attempted to rescue axis curvature by decreasing global Urp1/2 production via genetic means. To that end, we generated double (*rpgrip1l$^{+/-}$; urp2$^{+/-}$*) heterozygous fish using the recently generated *urp2* deletion allele (Gaillard et al., 2023) and crossed them to produce double mutants. Using a curvature quantification method on whole fish (Supplementary Figure S3I and methods), we observed that the removal of one or two *urp2* functional allele(s) in *rpgrip1l$^{-/-}$* fish did not lower their curvature index (Figure 4B, C), neither at two mpf (Supplementary Figure S3 F) nor at four mpf (Supplementary Figure S3 G-H). We observed that *urp2* mRNA amounts were reduced by removing one or two functional copies of *urp2*, probably as a consequence of mRNA decay (Gaillard et al., 2023) (Figure 4D). We also showed that none of the other *urotensin 2* family members was upregulated by a compensation mechanism upon *urp2* loss (Figure 4E for *urp1* and Supplementary Figure S3C-E for *urp, uts2a* and *uts2b*).

Thus, lowering global Urp1/2 expression level in *rpgrip1l* mutants did not have any beneficial effect on the axis curvature phenotype, ruling out altered *urp1/2* signaling as a main driver of scoliosis in *rpgrip1l* mutants.

Regulators of motile ciliogenesis, inflammation genes and annexins are upregulated in rpgrip1l$^{-/-}$ juveniles

To get further insight into the early mechanisms driving spine curvature in *rpgrip1l* mutants, we obtained the transcriptome of the brain and dorsal trunk of 7 *rpgrip1l*$^{-/-}$ juveniles at scoliosis onset (6 tail-up and 1 straight 5 wpf juveniles) and 5 control siblings (2 *rpgrip1l*$^{+/+}$ and 3 *rgprip1l*$^{+/-}$). Differential gene expression (DGE) analysis was performed with the DSEQ2 software within the Galaxy environment. A complete annotated gene list is displayed in Supplementary Table 1 (Trunk) and Table 2 (Brain). Filtering the gene list to select for P-adj < 0.05 and Log2 FC > 0.75 or < -0.75 recovered a vast majority of upregulated genes in the dorsal trunk (75 downregulated vs 614 upregulated genes) and, to a lesser extent, in the brain (1 downregulated, *rpgrip1l,* vs 60 upregulated genes) as illustrated on the volcano plots from DGE analysis of the brain (Figure 5A) and trunk (Figure 5B). 54 out of 60 genes upregulated in the brain were also upregulated in the trunk, suggesting that, in the trunk, these genes were specific to the CNS (Figure 5D). In both the brain (Figure 5C) and the trunk (Supplementary Figure S4A), the most upregulated biological processes in GO term analysis included cilium movement and cilium assembly. Among the 20 most significant GO terms found for upregulated genes in the trunk, 8 were related to ciliary movement/assembly (Supplementary Figure S4A). The upregulation in both tissues of *foxj1a,* encoding a master transcriptional activator of genes involved in cilia motility (Yu et al., 2008) (Figure 5E) likely contributed to this enrichment. To confirm this hypothesis, we compared our dataset with those of targets of Foxj1 from (Choksi et al., 2014; Mukherjee et al., 2019; Quigley and Kintner, 2017). We found that 106 out of 614 upregulated genes were targets of Foxj1, among which 77 were direct targets (Supplementary Table 3). Several genes encoding regulators of embryonic axis curvature were also upregulated in *rpgrip1l*$^{-/-}$ fish, among which *pkd1b* (Mangos et al., 2010), *urp1* and *urp2* (Bearce et al., 2022; Gaillard et al., 2023; Zhang et al., 2018) and *sspo* (Cantaut-Belarif et al., 2018; Rose et al., 2020; Troutwine et al., 2020) (Figure 5E). Finally, GO terms enrichment analysis also highlighted processes associated with inflammation and innate and adaptive immune responses, as described for the scoliotic *ptk7* model (Van Gennip

et al., 2018) sharing for example *irg1l* and complement genes up-regulation (Figure 5

C, E, and Supplementary Figure S4A).

In order to confirm these results and to identify processes affected at the post-

transcriptional level in *rpgrip1l* mutants, we performed a quantitative proteomic

analysis by mass spectrometry of brains from 5 *rpgrip1l$^{-/-}$* adult fish and 5 control

siblings. This analysis detected 5706 proteins present in at least 4 over the 5

replicates, of which 172 are associated with a P value < 0.05 and a Log2FC > +0.75

and < -0.75. 127 proteins were present in higher and 45 in lower quantity, as

illustrated on the volcano plot (Figure 5F and Supplementary Table 4). Pathways

enriched in *rpgrip1l$^{-/-}$* brains included regulation of vesicular trafficking and membrane

repair processes, as well as proteolysis and catabolic processes, suggesting

profound perturbation of cellular homeostasis (Supplementary Figure S4B). Among

the proteins that were present in highest quantity in mutants, several members of the

Annexin family were identified: Anxa2a, Anxa1a, Anxa5b (Figure 5F and

Supplementary Figure S4C. Annexins are associated with membranes from endo

and exovesicles as well as plasma membranes and involved in a wide array of

intracellular trafficking processes such as cholesterol endocytosis, plasma membrane

repair as well as modulation of immune system functions (Boye and Nylandsted,

2016; Grewal et al., 2016; Rentero et al., 2018). Of note, only four proteins of that list,

Anxa2a, C4, C7a and Stat3, were the products of genes up-regulated in the

transcriptomic study (indicated in red in Figure 5E). In agreement with the common

enriched GO terms in both studies (Figure 5C and Supplementary Figure S4B), these

four proteins are associated with immune response. Overall, *rpgrip1l$^{-/-}$* mutants at

scoliosis onset present deregulated pathways in common with other scoliotic models

such as activation of immune response genes, but also specific deregulated

pathways such a marked activation of the Foxj1a ciliary motility program and an

increase of Annexin levels.

Astrogliosis precedes scoliosis onset of rpgrip1l$^{-/-}$ juvenile fish

Both transcriptomic and proteomic analysis highlighted the occurrence of innate and

adaptive immune responses in *rpgrip1l$^{-/-}$* fish. Several upregulated genes, such as

irg1l, mmp9 and anxa2, are reported to be expressed in macrophages and/or

microglia but also in epithelia or neurons (Hall et al., 2014) (Kuhlmann et al., 2014)
(Shan et al., 2015). We therefore investigated in which cell type(s) Anxa2, the most
upregulated protein in the *rpgrip1l*$^{-/-}$ brain proteome, was expressed. Immunostaining
of brain sections revealed a strong Anxa2 staining on cells lining brain ventricles,
such as SCO cells and ventral ependymal cells of the rhombencephalic ventricle
(RhV) (Fig 6B', D') in *rpgrip1l*$^{-/-}$ adult fish, which was barely detectable in controls
(Figure 6A', C'), and to a lesser extent on other brain regions (Supplementary Fig S5
A'-F'), suggesting a deregulation of ependymal cell homeostasis. In mutants, Anxa2
staining spread over long cellular extensions reaching the pial surface of the
rhombencephalon (Figure 6D' and Supplementary Figure S5F'). This cell shape is
characteristic of radial glial cells, a cell type that behaves as neural and glial
progenitors and also plays astrocytic functions in zebrafish, as anamniote brains are
devoid of stellate astrocytes (Jurisch☐Yaksi et al., 2020; Lyons and Talbot, 2015). To
confirm the glial nature of Anxa2-positive cells around brain ventricles in *rpgrip1l*$^{-/-}$
fish, we double-labeled brain sections with GFAP, a glial cell marker, and Anxa2. In
adult *rpgrip1l*$^{-/-}$ fish, SCO cells were strongly double-positive for Anxa2 and GFAP
(Fig 6B, B'), while controls displayed much weaker labeling for GFAP in the SCO and
no labeling for Anxa2 (Fig 6 A, A'). We made similar observations at
rhombencephalic ventricle (RhV) and other brain levels: ependymal cells were
positive for Anxa2 and strongly positive for GFAP, and GFAP labeling was much
higher in mutants than in controls (Fig 6 C-D'; supplementary Figure S5A-F'). GFAP
overexpression is a hallmark of astrogliosis, a process in which astroglial cells react
to the perturbed environment, as shown during brain infection, injury, ischemia or
neurodegeneration (Escartin et al., 2021; Matusova et al., 2023; Zamanian et al.,
2012). Thus, the strong increase of GFAP staining combined with Anxa2
overexpression in brain ependymal cells of *rpgrip1l*$^{-/-}$ adult fish strongly suggests that
these cells undergo astrogliosis as observed in other animal models following CNS
injury (Chen et al., 2017; Zamanian et al., 2012).
To assay whether an astrogliosis-like phenotype was already present at scoliotic
onset, we immunostained GFAP on juvenile brain sections of straight and tail-up
mutants and controls. Tail-up juveniles (n=3/3) presented a strong GFAP staining in
almost all cells of the SCO and along the ventral rhombencephalic ventricle, higher
than control levels (Figure 6 E, I, H, L, M) while the phenotype of straight fish was

heterogeneous: two out of four juveniles presented GFAP-positive cells (Figure 6G, K

and red dots on Figure 6M graph), while the two other did not (Figure 6F, J and

green dots on Figure 6M graph) as confirmed by cell quantification (Figure 6M). As

scoliosis onset is asynchronous in *rpgrip1l*[-/-], heterogenous GFAP staining among

straight mutant fish may support an early role of astrogliosis at SCO and RhV levels

in scoliosis onset.

Immune cell enrichment around the SCO and within tectum parenchyma is

detected prior to scoliosis onset.

Astrogliosis may be reinforced or induced by an inflammatory environment containing

cytokines released by microglia or macrophages or the presence of high levels of

ROS and NO (Sofroniew, 2015). We thus performed immunostaining for the LCP1/L-

Plastin marker, which labels both macrophages and microglia (Figure 6N-Q'), in

adjacent sections to those used for GFAP (Figure 6E-L). We found that the scoliotic

fish presented a higher number of LCP1-positive cells of ameboid shape within SCO

cells (Figure 6Q, Q', R). Strikingly, the straight mutants (2/4) that presented a high

GFAP staining within SCO cells (Figure 6G, K) also showed a higher number of

LCP1-positive cells around the SCO (Figure 6P, P'; red dots in Figure 6R) than in

controls (Figure 6N, N'), while the other straight mutants did not ((Figure 6F, J, O, O',

green dots in Figure 6M) suggesting positive regulation between astrogliosis and

macrophages/microglia recruitment. LCP1+ cells were also detected in other regions

of the brain (Supplementary Figure S5 G-K). In the optic tectum, where resident

LCP1-positive macrophages/microglia were present in controls, a two-fold increase in

the number of ramified LCP1[+] cells was observed in straight *rpgrip1l*[-/-] juveniles

compared to controls (Supplementary Figure S5H-J). Surprisingly, scoliotic *rpgrip1l*[-/-]

fish showed similar LCP1[+] cell number as control fish in the tectum (Supplementary

Figure S5G), suggesting a transient increase of macrophage/microglia activation

preceding scoliosis.

Our data thus uncovers an early astrogliosis process initiated asynchronously at the

SCO and RhV levels of straight *rpgrip1l*[-/-] fish, favored by a transient inflammatory

state, and preceding the loss of multiciliated tufts around the SCO and that of the RF,

that we only detected in curved *rpgrip1l*[-/-] fish. Astrogliosis then spreads along the

brain ventricles and becomes prominent in brain of scoliotic fish. In addition, we
found among the upregulated genes in dorsal trunk at scoliosis onset, a number of
activated astrocyte markers (Matusova et al., 2023) such as *s100b*, *c4b*, *gfap*,
glast/slc1a3b, *viml*, *ctsb* (Supplementary Figure S4D), showing that astrogliosis
spread over the whole central nervous system.

Brain neuroinflammation and astrogliosis contribute to scoliosis penetrance
and severity in rpgrip1l*^{-/-}* ***fish

Our data suggest that astrogliosis and neuro-inflammation could play a role in
triggering scoliosis in *rpgrip1l* mutants. We thus aimed to counteract inflammation
before curvature initiation by performing drug treatment with the hope of reducing
scoliosis penetrance and severity. We used the NACET anti-oxidant and anti-
inflammatory drug that successfully reduced scoliosis of *sspo* hypomorphic mutant
fish (Rose et al., 2020). We first assayed the biological activity of our NACET batch
on the ciliary motility mutant *dnaaf1*^{-/-} (Sullivan-Brown et al., 2008). Indeed, NACET
was shown to suppress the embryonic tail-down phenotype of the ciliary motility
mutant *ccdc151*^{ts} (Rose et al., 2020). We found that 3mM NACET treatment
suppressed the *dnaaf1*^{-/-} tail-down phenotype (Supplementary Figure S6A-F). To
suppress scoliosis, we treated the progeny of *rpgrip1l*^{+/-} incrosses from 4 to 12 wpf
with 1.5 mM NACET as in (Rose et al., 2020).This long-term treatment reduced
scoliosis penetrance from 92% to 58% and markedly reduced spinal curvature index
(Figure 7A-C), measured and illustrated as shown in Supplementary Figure S3I.
We then asked whether NACET treatment suppressed astrogliosis and
neuroinflammation in *rpgrip1l*^{-/-} treated fish. Indeed, the intensity and number of
GFAP+ cells within the SCO were reduced in slightly curved fish and alleviated in
straight fish (Figure 7E, F). Furthermore, the quantification of Lcp1-positive cells in
the SCO region showed a drastic reduction of these cells in NACET-treated fish,
back to levels found in *rpgrip1l*^{+/+} fish (Figure 7D, E).
Our data thus show that *rpgrip1l*^{-/-} juvenile fish display brain astrogliosis and
inflammation at scoliosis onset and that an anti-inflammatory/anti-oxidant treatment
of *rpgrip1l*^{-/-} juveniles is able to maintain a straight axis in approximately one third of
the mutants and to slow down curve progression in the other two thirds. The

beneficial effect of NACET on axis curvature correlates with the combined reduction

of astrogliosis and immune cell density.

Brain astrogliosis is present in zebrafish *cep290 $^{fh297/fh297}$* scoliotic fish

To investigate whether abnormal SCO cilia function and brain astrogliosis may be a

general mechanism of scoliosis development in TZ mutants, we studied the

cep290$^{fh297/fh297}$ zebrafish mutant. Zebrafish *cep290* mutants display variable axis

curvature defects at embryonic stages as well as slow retinal degeneration, and

develop juvenile scoliosis (Lessieur et al., 2019; Wang et al., 2022). Like *RPGRIP1L,*

the human *CEP290* gene encodes a TZ protein and its mutation leads to severe

neurodevelopmental ciliopathies (Baala et al., 2007). All *cep290$^{fh297/fh297}$* juveniles

developed a fully penetrant scoliosis by 4 wpf in our fish facility (n: 30/30). We

analyzed cilia presence within the SCO by immunostaining on sections (Figure 8A-

F'). At 4 wpf, the SCO had not reached its final adult width (14 versus 24 cells wide in

the anterior SCO), and each of its cells harbored a short monocilium labeled by

glutamylated tubulin (Figure 8A-C). Multicilia tufts on SCO lateral walls were not as

clearly visible as in adults (Figure 2F) but patches of glutamylated tubulin staining

were detected on both sides of the posterior SCO (Figure 8B-C'). In *cep290$^{-/-}$*,

monocilia of the SCO appeared longer than in controls (N=3/3) and pointed towards

an abnormally elongated diencephalic ventricle (Figure 8D-F). Lateral patches of

glutamylated tubulin staining were present in *cep290$^{-/-}$* posterior SCO as in controls

(Figure 8E-F'). Thus, cilia morphology is perturbed within *cep290$^{-/-}$* SCO monociliated

cells, as well as the overall morphology of the diencephalic ventricle at this level.

To test for the presence of astrogliosis, we quantified the number of SCO cells

presenting high GFAP labeling on several sections along the A/P extent of the SCO

in 3 control and 3 *cep290$^{-/-}$* brains (Figure 8G-K). 50% of SCO cells presented strong

GFAP staining in fully scoliotic juveniles, which was not detectable in controls (Figure

8H, K, N=2/3), while a lower proportion (22%) of SCO cells was GFAP positive in the

juvenile with a milder curvature (Figure 8M, N=1/3). Strong GFAP staining was also

detected in some diencephalic radial glial cells (Figure 8K). On the same sections,

immune cells were quantified by LCP1 staining. A mean of 4 LCP1 positive cells per

SCO field was found in *cep290$^{-/-}$,* versus 1 in controls (Figure 8I, L and N). As in

rpgrip1l mutants, abnormal monocilia, astrogliosis and immune cell enrichment were

evidenced at SCO levels in *Cep290* [-/-](3/3). Furthermore, astrogliosis was observed

within radial glial cells lining the RhV at cerebellar level (Figure 8O-R) and at anterior

hindbrain level (Figure 8S-V), as shown in *rpgrip1l*[-/-] scoliotic brain (Figure 6D, K, L).

This indicates that astrogliosis is shared by these two zebrafish scoliotic models,

even though they develop axis curvature at different stages (3-4 wpf for *cep290*[-/-]

versus 5-12 wpf for *rpgrip1l*[-/-]).

DISCUSSION

In this paper we dissected the mechanisms of scoliosis appearance in a zebrafish

mutant for the TZ gene *rpgrip1l*, taking advantage of its asynchronous onset to

decipher the chronology of events leading to spine torsion. Transcriptome analysis of

the brain and trunk at scoliosis onset in *rpgrip1l*[-/-] fish compared to control siblings

revealed increased expression of *urp1/2*, *foxj1a* and its target genes, as well as

genes linked to immune response. Genetic experiments ruled out an involvement of

increased Urp1/2 signaling in scoliosis development. Cilia defects appeared

asynchronously in different CNS regions, while RF loss strictly coincided with

scoliosis onset. Strikingly, we observed a strong increase of GFAP staining along the

diencephalic and rhombencephalic ventricles, indicative of astrogliosis. This

astrogliosis preceded scoliosis onset and then spread posteriorly and increased in

severity. In the region of the SCO, the appearance of astrogliosis and

neuroinflammation coincided temporally, preceding the onset of RF depolymerization

and spine torsion. NACET treatment showed that increased astrogliosis and

inflammation participated in the penetrance and severity of scoliosis in *rpgrip1l*[-/-] fish.

Finally, we showed that another TZ gene mutant, *cep290*, also displayed astrogliosis

at the time of scoliosis onset. Thus, the cascade of events uncovered in the *rpgrip1l*

mutant sheds a new light on the origin of zebrafish scoliosis caused by ciliary TZ

defects.

Cilia motility defects lead to RF loss and to scoliosis in several zebrafish mutants

(Grimes et al., 2016; Rose et al., 2020)). *Rpgrip1l* encodes a TZ protein important for

cilia integrity and function in many eukaryotic organisms (Andreu-Cervera et al.,

2019; Gogendeau et al., 2020; Jensen et al., 2015; Wiegering et al., 2018). Our data

show that *rpgrip1l* deficiency in zebrafish affects cilia maintenance differentially

depending on the CNS territory and the developmental stage, raising the question of

which ciliated cells are responsible for RF loss and scoliosis initiation in this mutant.

The full rescue observed after re-introducing RPGRIP1L in *foxj1a*-expressing cells

points towards a crucial function of the protein in cells lining CNS cavities. Foxj1a-

positive cell populations encompass both monociliated and multiciliated ependymal

cells, radial glial progenitors, glial cells of the SCO and the ChP, as well as CSF-

contacting sensory neurons in the spinal cord (D'Gama et al., 2021; Prendergast et

al., 2023; Ribeiro et al., 2017; Van Gennip et al., 2018). Recent work suggested that

impairment of ciliogenesis in the fChP of the *katnb1* mutant could play a role in

inducing axis curvature (Meyer-Miner et al., 2022). In the case of the *rpgrip1l* mutant,

we temporally uncorrelated fChP cilia defects from scoliosis onset. This, combined

with the fact that the total loss of multicilia in the fChP of [*foxj1b$^{-/-}$, gmnc$^{-/-}$*] double

mutants does not trigger axis curvature (D'Gama et al., 2021) suggests that another

foxj1a-expressing population is crucial for axis straightness. Our results strongly

suggest that cilia defects in the SCO trigger axis curvature in *rpgrip1l* mutants. Two

cilia populations were affected in the SCO of *rpgrip1l* mutant juveniles at the onset of

scoliosis: cilia of monociliated glial cells that secrete SCO-spondin, which appeared

longer in mutants than in controls, and multiciliated tufts at SCO exit, which were lost

in scoliotic mutants. Both defects could be responsible for the phenotype, by

perturbing CSF flow and content (presence of SCO-spondin aggregates in ventricles)

(left side on the Graphical Abstract) and leading to ventricular dilations or by

perturbing primary cilia signaling within SCO cells (right side of the Graphical

Abstract). The observation that multiciliated cells were not yet fully differentiated in

cep290 mutants at scoliosis onset is in favor of a prominent role of monocilia. This is

consistent with the literature showing that most ciliary mutant models develop axis

curvature at 3-4 weeks, a stage when brain multiciliated cells are not yet

differentiated (D'Gama et al., 2021). Still, as scoliosis onset in *rpgrip1l* mutant is

asynchronous from 5 to 12 weeks, MCC loss at SCO exit could contribute to

defective RF polymerization.

Our model for scoliosis appearance in the *rpgrip1l* mutant is presented in the
graphical abstract. Since we have observed several straight *rpgrip1l*[-/-] fish with
astrogliosis and increased immune cells presence around the SCO, we propose that
rpgrip1l[-/-] SCO and astroglial cells experience altered signaling that activates immune
cell recruitment, producing an inflammatory environment. This would promote
multicilia loss as a secondary event (Graphical Abstract, orange arrow) (Lattke et al.,
2012) leading to RF depolymerization and spine curvature. However, as *sspo* [-/-]
hypomorphic mutants which present aggregates within ventricles have been shown
to present an inflammatory response (Rose et al., 2020), it is equally possible that
SCO astrogliosis is a downstream consequence of the presence of these
proteoglycans aggregates in contact with primary cilia or ventricular cell surfaces
(Graphical Abstract , green arrow). These two scenarios may also act in parallel to
reinforce scoliosis onset and ependymal cilia loss. Future experiments will help to
discriminate between these two scenarios.
540
Our transcriptome analysis and qRT-PCR data challenges a model proposed for axis
straightness of zebrafish embryos in which the loss of RF leads to a down-regulation
of *urp2* and *urp1* expression (Zhang et al., 2018) since we observed an upregulation
of both genes before and after RF loss. As forced expression of *urp1/2* induces a tail-
up phenotype in embryos and juveniles (Zhang et al., 2018) (Gaillard et al., 2023; Lu
et al., 2020), we attempted to rescue *rpgrip1l*[-/-] axis curvature by down-regulating
urp1/2 expression. No beneficial effect on scoliosis penetrance or severity was
observed, indicating that increased *urp1/2* expression level does not significantly
contribute to axis curvature in *rpgrip1l*[-/-] fish.
550
In this study, we found an unexpected and strong defect characterized by enhanced
GFAP and Anxa2 expression, which we identified as astrogliosis (also called reactive
astrogliosis). Astrogliosis is a reaction of astroglial cells to perturbed homeostasis of
the CNS, characterized by molecular and phenotypic changes in these cells (Escartin
et al., 2021; Matusova et al., 2023). In zebrafish, radial glial cells are endowed with
both neurogenic and astrocytic functions, and are thus also called astroglial cells
(Jurisch☐Yaksi et al., 2020). Astrogliosis in *rpgrip1l* mutants arose in a subdomain of
Foxj1a-positive cells, within the DiV and ventral ependymal cells lining the RhV. The

observation of astrogliosis in some straight juvenile mutants suggests an early role of
this defect in scoliosis onset.

Astrogliosis along the ventricular cavities could arise in response to local mechanical
or chemical perturbations caused by cilia motility defects, abnormal CSF flow and
content and ventricular dilations. In two murine models of primary ciliary dyskinesia,
decreased CSF flow was associated with gliosis at juvenile stage (Finn et al., 2014).
A response to CSF pertubation in *rpgrip1l* mutants is also suggested by the strong
upregulation of Foxj1a and its target genes. *foxj1a* expression is also reported to be
upregulated in the CNS of three other scoliotic models in addition to *rpgrip1l*, namely
in *ptk7, sspo* and *katnb1* mutants (Escartin et al., 2021; Meyer-Miner et al., 2022;
Van Gennip et al., 2018), suggesting that it constitutes a common response to similar
ventricular and CSF defects. Interestingly, *foxj1a* expression is also upregulated
upon zebrafish embryonic (Cavone et al., 2021; Hellman et al., 2010) or adult
(Ribeiro et al., 2017) spinal cord injury. The up-regulation in the *rpgrip1l* mutant
proteome of the membrane repair module Anxa2-S100a10-Ahnak (Bharadwaj et al.,
2021) (Figure 4C) together with the Foxj1-induced motility module detected in the
transcriptomic analysis may indicate ongoing reparation attempts of CNS ventricles
and associated neural tissues.

Moreover, the presence of few Lcp1+ cells around the mutant SCO suggests that
microglia/macrophages could participate in astrogliosis establishment and
reinforcement. This dialogue between immune Lcp1+ cells and astroglial cells is
evidenced during early stages of regeneration after acute injury or cellular damages
(Cavone et al., 2021). In our bulk RNAseq analysis performed at scoliosis onset, we
have indications of the upregulation of microglia and macrophage markers (*p2ry12*
and *mpeg1*) at trunk levels and of numerous markers of activated astrocytes
characterized in several mammalian astrogliosis models (Matusova et al., 2023). A
similar dialogue is detected in polycystic kidney disease models where compromised
primary cilia signalling leads to uncontrolled cytokines secretion by epithelial cells, a
situation that favors local immune cells recruitment and proliferation (Viau et al.,
2018).

What are the intracellular mechanisms involved in astrogliosis induction in ciliated
ventricular cells? Candidate pathways emerge from our multi-omics studies.

Proteomic data showed a 2.4-fold increase in GSK3bb amount in *rpgrip1l*[−/−] mutant
brains (P value < 0.001, Supplementary Table 3). In several neurodegenerative
murine animal models, GSK3b enzymatic activity was shown to promote
inflammation and gliosis (Jorge-Torres et al., 2018; Medina and Avila, 2010; Mines et
al., 2011). Another potential trigger of astrogliosis may be a defective mitochondrial
activity as the amounts of four proteins involved in electron transport chain activity
(*gpd1b, mt-nd6, pdia5 and dmgdh*) were reduced by 40% to 86% in the analysis of
mutant brain proteome (Supplementary Table 4). We think this is of particular interest
in light of a recent report demonstrating that impaired mitochondrial activity within
ciliated astrocytes leads to the induction of the Foxj1 ciliary motility program, the
elongation and distortion of astrocyte primary cilia as well as reactive astrogliosis
(Ignatenko et al., 2023), three phenotypes also observed in *rpgrip1l*[−/−] brains.
603
Finally, our observation of widespread astrogliosis in two TZ gene mutants, *rpgrip1l*
and *cep290*, suggests that it could represent a general mechanism involved in
scoliosis downstream of cilia dysfunction. Of note, astrogliosis markers such as
glast/slc1a3b, *vim*, *s100b*, *c4*, *timp2b*, *gfap* and *ctssb.2*, are also upregulated in the
transcriptome of *ptk7 mutants* (Van Gennip et al., 2018). We propose that sustained
astrogliosis might impair neuronal survival and activity, crucial for proper
interoception and locomotor control, thus leading to axis curvature in the context of a
rapidly growing axial skeleton. A similar context of astrocytosis associated with spinal
cord dilation has been observed in human patients with syringomyelia. This
population with a high incidence of scoliosis can present hyper-signals on T2 MRI
scans, which were proposed to reflect local astrocytosis and could be validated by
biopsies in rare cases (Sherman et al., 1987). Further imaging studies will need to be
performed to validate a potential link between developing astrogliosis and the onset
and progression of idiopathic scoliosis in humans.
618

619 **ACKNOWLEDGEMENTS**
620
We are grateful to the IBPS aquatic animal, imaging and bioinformatics facilities and
to the ICM sequencing facility for their technical assistance. We thank Michaël
Trichet from the IBPS electron microscopy facility for participating in the MEB

experiments, the TACGENE facility for providing the Cas9 protein; the TEFOR Paris-

Saclay facility for the brain clearing experiment; Thierry Jaffredo and Pierre Charbord

for their precious help in transcriptome analysis; Nicolas Baylé for initial

characterization of the *rpgrip1l* mutant; Claudia Hoffman for providing *Cep290* fixed

samples; Brian Ciruna for sharing the *foxj1a* expressing constructs and detailed

NACET treatment protocol; Ryan Gray for sharing the *scospondin-GFP^{ut24}* zebrafish

line. This work was supported by funding to SSM from the Fondation pour la

Recherche Médicale (Equipe FRM EQU201903007943) and the Fondation Yves

Cotrel.

REFERENCES

Affaticati P, Simion M, De Job E, Rivière L, Hermel J-M, Machado E, Joly J-S, Jenett A. 2017. zPACT: Tissue Clearing and Immunohistochemistry on Juvenile Zebrafish Brain. *BIO-PROTOCOL* **7**. doi:10.21769/BioProtoc.2636

Andreu-Cervera A, Anselme I, Karam A, Laclef C, Catala M, Schneider-Maunoury S. 2019. The ciliopathy gene *ftm/rpgrip1l* controls mouse forebrain patterning via region-specific modulation of hedgehog/gli signaling. *J Neurosci* 2199–18. doi:10.1523/JNEUROSCI.2199-18.2019

Andreu-Cervera A, Catala M, Schneider-Maunoury S. 2021. Cilia, ciliopathies and hedgehog-related forebrain developmental disorders. *Neurobiology of Disease* **150**:105236. doi:10.1016/j.nbd.2020.105236

Arts HH, Doherty D, Van Beersum SEC, Parisi MA, Letteboer SJF, Gorden NT, Peters TA, Märker T, Voesenek K, Kartono A, Ozyurek H, Farin FM, Kroes HY, Wolfrum U, Brunner HG, Cremers FPM, Glass IA, Knoers NVAM, Roepman R. 2007. Mutations in the gene encoding the basal body protein RPGRIP1L, a nephrocystin-4 interactor, cause Joubert syndrome. *Nat Genet* **39**:882–888. doi:10.1038/ng2069

Baala L, Audollent S, Martinovic J, Ozilou C, Babron M-C, Sivanandamoorthy S, Saunier S, Salomon R, Gonzales M, Rattenberry E, Esculpavit C, Toutain A, Moraine C, Parent P, Marcorelles P, Dauge M-C, Roume J, Merrer ML, Meiner V, Meir K, Menez F, Beaufrère A-M, Francannet C, Tantau J, Sinico M, Dumez Y, MacDonald F, Munnich A, Lyonnet S, Gubler M-C, Génin E, Johnson CA, Vekemans M, Encha-Razavi F, Attié-Bitach T. 2007. Pleiotropic Effects of CEP290 (NPHP6) Mutations Extend to Meckel Syndrome. *The American Journal of Human Genetics* **81**:170–179. doi:10.1086/519494

Bachmann-Gagescu R, Phelps IG, Stearns G, Link BA, Brockerhoff SE, Moens CB, Doherty D. 2011. The ciliopathy gene cc2d2a controls zebrafish photoreceptor outer segment development through a role in Rab8-dependent vesicle trafficking. *Human Molecular Genetics* **20**:4041–4055. doi:10.1093/hmg/ddr332

Bearce EA, Irons ZH, O'Hara-Smith JR, Kuhns CJ, Fisher SI, Crow WE, Grimes DT. 2022. Urotensin II-related peptides, Urp1 and Urp2, control zebrafish spine morphology. *eLife* **11**:e83883. doi:10.7554/eLife.83883

Bentley-Ford MR, LaBonty M, Thomas HR, Haycraft CJ, Scott M, LaFayette C, Croyle MJ, Andersen RS, Parant JM, Yoder BK. 2022. Evolutionarily conserved genetic interactions between *nphp-4* and *bbs-5* mutations exacerbate ciliopathy phenotypes. *Genetics*

220:iyab209. doi:10.1093/genetics/iyab209

Bharadwaj A, Kempster E, Waisman DM. 2021. The Annexin A2/S100A10 Complex: The

Mutualistic Symbiosis of Two Distinct Proteins. *Biomolecules* **11**:1849.

doi:10.3390/biom11121849

Boswell CW, Ciruna B. 2017. Understanding Idiopathic Scoliosis: A New Zebrafish School

of Thought. *Trends in Genetics* **33**:183–196. doi:10.1016/j.tig.2017.01.001

Bouyssié D, Hesse A-M, Mouton-Barbosa E, Rompais M, Macron C, Carapito C, Gonzalez

De Peredo A, Couté Y, Dupierris V, Burel A, Menetrey J-P, Kalaitzakis A, Poisat J,

Romdhani A, Burlet-Schiltz O, Cianférani S, Garin J, Bruley C. 2020. Proline: an efficient

and user-friendly software suite for large-scale proteomics. *Bioinformatics* **36**:3148–3155.

doi:10.1093/bioinformatics/btaa118

Boye TL, Nylandsted J. 2016. Annexins in plasma membrane repair. *Biological Chemistry*

397:961–969. doi:10.1515/hsz-2016-0171

Brancati F, Travaglini L, Zablocka D, Boltshauser E, Accorsi P, Montagna G, Silhavy JL,

Barrano G, Bertini E, Emma F, Rigoli L, International JSRD Study Group, Dallapiccola B,

Gleeson JG, Valente EM. 2008. RPGRIP1L mutations are mainly associated with the

cerebello-renal phenotype of Joubert syndrome-related disorders. *Clin Genet* **74**:164–170.

doi:10.1111/j.1399-0004.2008.01047.x

Cantaut-Belarif Y, Orts Del'Immagine A, Penru M, Pézeron G, Wyart C, Bardet P-L. 2020.

Adrenergic activation modulates the signal from the Reissner fiber to cerebrospinal fluid-

contacting neurons during development. *eLife* **9**:e59469. doi:10.7554/eLife.59469

Cantaut-Belarif Y, Sternberg JR, Thouvenin O, Wyart C, Bardet P-L. 2018. The Reissner

Fiber in the Cerebrospinal Fluid Controls Morphogenesis of the Body Axis. *Current Biology*

28:2479-2486.e4. doi:10.1016/j.cub.2018.05.079

Cavone L, McCann T, Drake LK, Aguzzi EA, Oprişoreanu A-M, Pedersen E, Sandi S,

Selvarajah J, Tsarouchas TM, Wehner D, Keatinge M, Mysiak KS, Henderson BEP, Dobie R,

Henderson NC, Becker T, Becker CG. 2021. A unique macrophage subpopulation signals

directly to progenitor cells to promote regenerative neurogenesis in the zebrafish spinal cord.

Developmental Cell **56**:1617-1630.e6. doi:10.1016/j.devcel.2021.04.031

Chen J, Cui Z, Yang S, Wu C, Li W, Bao G, Xu G, Sun Y, Wang L, Zhang J. 2017. The

upregulation of annexin A2 after spinal cord injury in rats may have implication for astrocyte

proliferation. *Neuropeptides* **61**:67–76. doi:10.1016/j.npep.2016.10.007

Choksi SP, Lauter G, Swoboda P, Roy S. 2014. Switching on cilia: transcriptional networks

regulating ciliogenesis. *Development* **141**:1427–1441. doi:10.1242/dev.074666

Collins P, Woollam DH. 1979. The ventricular surface of the subcommissural organ: a

scanning and transmission electron microscopic study. *J Anat* **129**:623–631.

Dale RM, Topczewski J. 2011. Identification of an evolutionarily conserved regulatory

element of the zebrafish col2a1a gene. *Developmental Biology* **357**:518–531.

doi:10.1016/j.ydbio.2011.06.020

Delous M, Baala L, Salomon R, Laclef C, Vierkotten J, Tory K, Golzio C, Lacoste T, Besse

L, Ozilou C, Moutkine I, Hellman NE, Anselme I, Silbermann F, Vesque C, Gerhardt C,

Rattenberry E, Wolf MTF, Gubler MC, Martinovic J, Encha-Razavi F, Boddaert N, Gonzales

M, Macher MA, Nivet H, Champion G, Berthélémé JP, Niaudet P, McDonald F, Hildebrandt

F, Johnson CA, Vekemans M, Antignac C, Rüther U, Schneider-Maunoury S, Attié-Bitach T,

Saunier S. 2007. The ciliary gene RPGRIP1L is mutated in cerebello-oculo-renal syndrome

(Joubert syndrome type B) and Meckel syndrome. *Nat Genet* **39**:875–881.

doi:10.1038/ng2039

D'Gama PP, Qiu T, Cosacak MI, Rayamajhi D, Konac A, Hansen JN, Ringers C, Acuña-

Hinrichsen F, Hui SP, Olstad EW, Chong YL, Lim CKA, Gupta A, Ng CP, Nilges BS,

Kashikar ND, Wachten D, Liebl D, Kikuchi K, Kizil C, Yaksi E, Roy S, Jurisch-Yaksi N.

2021. Diversity and function of motile ciliated cell types within ependymal lineages of the

zebrafish brain. *Cell Reports* **37**:109775. doi:10.1016/j.celrep.2021.109775

Didier R, Dastugue B, Meiniel A. 1995. The secretory material of the subcommissural organ

of the chick embryo. Characterization of a specific polypeptide by two-dimensional

electrophoresis. *Int J Dev Biol* **39**:493–499.

Escartin C, Galea E, Lakatos A, O'Callaghan JP, Petzold GC, Serrano-Pozo A, Steinhäuser C,

Volterra A, Carmignoto G, Agarwal A, Allen NJ, Araque A, Barbeito L, Barzilai A, Bergles

DE, Bonvento G, Butt AM, Chen W-T, Cohen-Salmon M, Cunningham C, Deneen B, De

Strooper B, Díaz-Castro B, Farina C, Freeman M, Gallo V, Goldman JE, Goldman SA, Götz

M, Gutiérrez A, Haydon PG, Heiland DH, Hol EM, Holt MG, Iino M, Kastanenka KV,

Kettenmann H, Khakh BS, Koizumi S, Lee CJ, Liddelow SA, MacVicar BA, Magistretti P,

Messing A, Mishra A, Molofsky AV, Murai KK, Norris CM, Okada S, Oliet SHR, Oliveira

JF, Panatier A, Parpura V, Pekna M, Pekny M, Pellerin L, Perea G, Pérez-Nievas BG,

Pfrieger FW, Poskanzer KE, Quintana FJ, Ransohoff RM, Riquelme-Perez M, Robel S, Rose

CR, Rothstein JD, Rouach N, Rowitch DH, Semyanov A, Sirko S, Sontheimer H, Swanson

RA, Vitorica J, Wanner I-B, Wood LB, Wu J, Zheng B, Zimmer ER, Zorec R, Sofroniew

MV, Verkhratsky A. 2021. Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci* **24**:312–325. doi:10.1038/s41593-020-00783-4

Finn R, Evans CC, Lee L. 2014. Strain-dependent brain defects in mouse models of primary ciliary dyskinesia with mutations in Pcdp1 and Spef2. *Neuroscience* **277**:552–567. doi:10.1016/j.neuroscience.2014.07.029

Gaillard A-L, Mohamad T, Quan FB, De Cian A, Mosimann C, Tostivint H, Pézeron G. 2023. Urp1 and Urp2 act redundantly to maintain spine shape in zebrafish larvae. *Developmental Biology* **496**:36–51. doi:10.1016/j.ydbio.2023.01.010

Gogendeau D, Lemullois M, Le Borgne P, Castelli M, Aubusson-Fleury A, Arnaiz O, Cohen J, Vesque C, Schneider-Maunoury S, Bouhouche K, Koll F, Tassin A-M. 2020. MKS-NPHP module proteins control ciliary shedding at the transition zone. *PLoS Biol* **18**:e3000640. doi:10.1371/journal.pbio.3000640

Grewal T, Wason SJ, Enrich C, Rentero C. 2016. Annexins – insights from knockout mice. *Biological Chemistry* **397**:1031–1053. doi:10.1515/hsz-2016-0168

Grimes DT, Boswell CW, Morante NFC, Henkelman RM, Burdine RD, Ciruna B. 2016. Zebrafish models of idiopathic scoliosis link cerebrospinal fluid flow defects to spine curvature. *Science* **352**:1341–1344. doi:10.1126/science.aaf6419

Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud J-B, Schneider-Maunoury S, Shkumatava A, Teboul L, Kent J, Joly J-S, Concordet J-P. 2016. Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol* **17**:148. doi:10.1186/s13059-016-1012-2

Hall CJ, Boyle RH, Sun X, Wicker SM, Misa JP, Krissansen GW, Print CG, Crosier KE, Crosier PS. 2014. Epidermal cells help coordinate leukocyte migration during inflammation through fatty acid-fuelled matrix metalloproteinase production. *Nat Commun* **5**:3880. doi:10.1038/ncomms4880

Hayes M, Gao X, Yu LX, Paria N, Henkelman RM, Wise CA, Ciruna B. 2014. ptk7 mutant zebrafish models of congenital and idiopathic scoliosis implicate dysregulated Wnt signalling in disease. *Nat Commun* **5**:4777. doi:10.1038/ncomms5777

Hellman NE, Liu Y, Merkel E, Austin C, Le Corre S, Beier DR, Sun Z, Sharma N, Yoder BK, Drummond IA. 2010. The zebrafish *foxj1a* transcription factor regulates cilia function in response to injury and epithelial stretch. *Proc Natl Acad Sci USA* **107**:18499–18504. doi:10.1073/pnas.1005998107

Hitier M, Hamon M, Denise P, Lacoudre J, Thenint M-A, Mallet J-F, Moreau S, Quarck G.

2015. Lateral Semicircular Canal Asymmetry in Idiopathic Scoliosis: An Early Link between
Biomechanical, Hormonal and Neurosensory Theories? *PLoS ONE* **10**:e0131120.
doi:10.1371/journal.pone.0131120

Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, Krijgsveld J. 2014.
Ultrasensitive proteome analysis using paramagnetic bead technology. *Molecular Systems*
Biology **10**:757. doi:10.15252/msb.20145625

Ignatenko O, Malinen S, Rybas S, Vihinen H, Nikkanen J, Kononov A, Jokitalo ES, Ince-
Dunn G, Suomalainen A. 2023. Mitochondrial dysfunction compromises ciliary homeostasis
in astrocytes. *Journal of Cell Biology* **222**:e202203019. doi:10.1083/jcb.202203019

Jensen VL, Li C, Bowie RV, Clarke L, Mohan S, Blacque OE, Leroux MR. 2015. Formation
of the transition zone by Mks5/Rpgrip1L establishes a ciliary zone of exclusion (CIZE) that
compartmentalises ciliary signalling proteins and controls PIP $_2$ ciliary abundance. *The EMBO*
Journal **34**:2537–2556. doi:10.15252/embj.201488044

Jorge-Torres OC, Szczesna K, Roa L, Casal C, Gonzalez-Somermeyer L, Soler M, Velasco
CD, Martínez-San Segundo P, Petazzi P, Sáez MA, Delgado-Morales R, Fourcade S, Pujol A,
Huertas D, Llobet A, Guil S, Esteller M. 2018. Inhibition of Gsk3b Reduces Nfkb1 Signaling
and Rescues Synaptic Activity to Improve the Rett Syndrome Phenotype in Mecp2-Knockout
Mice. *Cell Reports* **23**:1665–1677. doi:10.1016/j.celrep.2018.04.010

Jurisch☐Yaksi N, Yaksi E, Kizil C. 2020. Radial glia in the zebrafish brain: Functional,
structural, and physiological comparison with the mammalian glia. *Glia* **68**:2451–2470.
doi:10.1002/glia.23849

Karner CM, Long F, Solnica-Krezel L, Monk KR, Gray RS. 2015. *Gpr126/Adgrg6* deletion in
cartilage models idiopathic scoliosis and pectus excavatum in mice. *Hum Mol Genet*
24:4365–4373. doi:10.1093/hmg/ddv170

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of embryonic
development of the zebrafish. *Developmental Dynamics* **203**:253–310.
doi:10.1002/aja.1002030302

Kramer-Zucker AG, Olale F, Haycraft CJ, Yoder BK, Schier AF, Drummond IA. 2005. Cilia-
driven fluid flow in the zebrafish pronephros, brain and Kupffer's vesicle is required for
normal organogenesis. *Development* **132**:1907–1921. doi:10.1242/dev.01772

Kuhlmann K, Tschapek A, Wiese H, Eisenacher M, Meyer HE, Hatt HH, Oeljeklaus S,
Warscheid B. 2014. The Membrane Proteome of Sensory Cilia to the Depth of Olfactory
Receptors. *Molecular & Cellular Proteomics* **13**:1828–1843. doi:10.1074/mcp.M113.035378

Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, Parant JM, Yost

HJ, Kanki JP, Chien C. 2007. The Tol2kit: A multisite gateway-based construction kit for

Tol2 transposon transgenesis constructs. *Developmental Dynamics* **236**:3088–3099.

doi:10.1002/dvdy.21343

Lattke M, Magnutzki A, Walther P, Baumann B. 2012. Nuclear Factor κB Activation Impairs

Ependymal Ciliogenesis and Links Neuroinflammation to Hydrocephalus Formation. *J*

Neurosci **32**:11511–11523. doi:10.1523/JNEUROSCI.0182-12.2012

Lessieur EM, Song P, Nivar GC, Piccillo EM, Fogerty J, Rozic R, Perkins BD. 2019. Ciliary

genes arl13b, ahi1 and cc2d2a differentially modify expression of visual acuity phenotypes

but do not enhance retinal degeneration due to mutation of cep290 in zebrafish. *PLoS One*

14:e0213960. doi:10.1371/journal.pone.0213960

Lu H, Shagirova A, Goggi JL, Yeo HL, Roy S. 2020. Reissner fibre-induced urotensin

signalling from cerebrospinal fluid-contacting neurons prevents scoliosis of the vertebrate

spine. *Biology Open* **9**:bio052027. doi:10.1242/bio.052027

Lyons DA, Talbot WS. 2015. Glial Cell Development and Function in Zebrafish. *Cold Spring*

Harb Perspect Biol **7**:a020586. doi:10.1101/cshperspect.a020586

Mahuzier A, Gaudé H-M, Grampa V, Anselme I, Silbermann F, Leroux-Berger M, Delacour

D, Ezan J, Montcouquiol M, Saunier S, Schneider-Maunoury S, Vesque C. 2012. Dishevelled

stabilization by the ciliopathy protein Rpgrip1l is essential for planar cell polarity. *Journal of*

Cell Biology **198**:927–940. doi:10.1083/jcb.201111009

Mangos S, Lam P, Zhao A, Liu Y, Mudumana S, Vasilyev A, Liu A, Drummond IA. 2010.

The ADPKD genes *pkd1a/b* and *pkd2* regulate extracellular matrix formation. *Disease Models*

& Mechanisms **3**:354–365. doi:10.1242/dmm.003194

Masek M, Etard C, Hofmann C, Hülsmeier AJ, Zang J, Takamiya M, Gesemann M, Neuhauss

SCF, Hornemann T, Strähle U, Bachmann-Gagescu R. 2022. Loss of the Bardet-Biedl protein

Bbs1 alters photoreceptor outer segment protein and lipid composition. *Nat Commun*

13:1282. doi:10.1038/s41467-022-28982-6

Matusova Z, Hol EM, Pekny M, Kubista M, Valihrach L. 2023. Reactive astrogliosis in the

era of single-cell transcriptomics. *Front Cell Neurosci* **17**:1173200.

doi:10.3389/fncel.2023.1173200

Medina M, Avila J. 2010. Glycogen Synthase Kinase-3 (GSK-3) Inhibitors for the Treatment

of Alzheimers Disease. *CPD* **16**:2790–2798. doi:10.2174/138161210793176581

Meyer-Miner A, Van Gennip JLM, Henke K, Harris MP, Ciruna B. 2022. Resolving primary

pathomechanisms driving idiopathic-like spinal curvature using a new katnb1 scoliosis model.

iScience **25**:105028. doi:10.1016/j.isci.2022.105028

Mines MA, Beurel E, Jope RS. 2011. Regulation of Cell Survival Mechanisms in Alzheimer's

Disease by Glycogen Synthase Kinase-3. *International Journal of Alzheimer's Disease*

2011:1–11. doi:10.4061/2011/861072

Mukherjee I, Roy S, Chakrabarti S. 2019. Identification of Important Effector Proteins in the

FOXJ1 Transcriptional Network Associated With Ciliogenesis and Ciliary Function. *Front*

Genet **10**:23. doi:10.3389/fgene.2019.00023

Olstad EW, Ringers C, Hansen JN, Wens A, Brandt C, Wachten D, Yaksi E, Jurisch-Yaksi N.

2019. Ciliary Beating Compartmentalizes Cerebrospinal Fluid Flow in the Brain and

Regulates Ventricular Development. *Current Biology* **29**:229-241.e6.

doi:10.1016/j.cub.2018.11.059

Parmentier C, Hameury E, Dubessy C, Quan FB, Habert D, Calas A, Vaudry H, Lihrmann I,

Tostivint H. 2011. Occurrence of Two Distinct Urotensin II-Related Peptides in Zebrafish

Provides New Insight into the Evolutionary History of the Urotensin II Gene Family.

Endocrinology **152**:2330–2341. doi:10.1210/en.2010-1500

Patten SA, Margaritte-Jeannin P, Bernard J-C, Alix E, Labalme A, Besson A, Girard SL,

Fendri K, Fraisse N, Biot B, Poizat C, Campan-Fournier A, Abelin-Genevois K, Cunin V,

Zaouter C, Liao M, Lamy R, Lesca G, Menassa R, Marcaillou C, Letexier M, Sanlaville D,

Berard J, Rouleau GA, Clerget-Darpoux F, Drapeau P, Moldovan F, Edery P. 2015.

Functional variants of POC5 identified in patients with idiopathic scoliosis. *J Clin Invest*

125:1124–1128. doi:10.1172/JCI77262

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S,

Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A,

Vizcaíno JA. 2022. The PRIDE database resources in 2022: a hub for mass spectrometry-

based proteomics evidences. *Nucleic Acids Research* **50**:D543–D552.

doi:10.1093/nar/gkab1038

Prendergast AE, Jim KK, Marnas H, Desban L, Quan FB, Djenoune L, Laghi V,

Hocquemiller A, Lunsford ET, Roussel J, Keiser L, Lejeune F-X, Dhanasekar M, Bardet P-L,

Levraud J-P, Van De Beek D, Vandenbroucke-Grauls CMJE, Wyart C. 2023. CSF-contacting

neurons respond to Streptococcus pneumoniae and promote host survival during central

nervous system infection. *Current Biology* **33**:940-956.e10. doi:10.1016/j.cub.2023.01.039

Quan FB, Dubessy C, Galant S, Kenigfest NB, Djenoune L, Leprince J, Wyart C, Lihrmann I,

Tostivint H. 2015. Comparative Distribution and In Vitro Activities of the Urotensin II-
Related Peptides URP1 and URP2 in Zebrafish: Evidence for Their Colocalization in Spinal
Cerebrospinal Fluid-Contacting Neurons. *PLoS ONE* **10**:e0119290.
doi:10.1371/journal.pone.0119290

Quigley IK, Kintner C. 2017. Rfx2 Stabilizes Foxj1 Binding at Chromatin Loops to Enable
Multiciliated Cell Gene Expression. *PLoS Genet* **13**:e1006538.
doi:10.1371/journal.pgen.1006538

Rentero C, Blanco-Muñoz P, Meneses-Salas E, Grewal T, Enrich C. 2018. Annexins—
Coordinators of Cholesterol Homeostasis in Endocytic Pathways. *IJMS* **19**:1444.
doi:10.3390/ijms19051444

Ribeiro A, Monteiro JF, Certal AC, Cristovão AM, Saúde L. 2017. Foxj1a is expressed in
ependymal precursors, controls central canal position and is activated in new ependymal cells
during regeneration in zebrafish. *Open Biol* **7**:170139. doi:10.1098/rsob.170139

Rose CD, Pompili D, Henke K, Van Gennip JLM, Meyer-Miner A, Rana R, Gobron S, Harris
MP, Nitz M, Ciruna B. 2020. SCO-Spondin Defects and Neuroinflammation Are Conserved
Mechanisms Driving Spinal Deformity across Genetic Models of Idiopathic Scoliosis.
Current Biology **30**:2363-2373.e6. doi:10.1016/j.cub.2020.04.020

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K,
Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis.
Nat Methods **9**:676–682. doi:10.1038/nmeth.2019

Shan Y, Fang C, Cheng C, Wang Y, Peng J, Fang W. 2015. Immersion infection of germ-free
zebrafish with Listeria monocytogenes induces transient expression of innate immune
response genes. *Front Microbiol* **6**. doi:10.3389/fmicb.2015.00373

Sherman J, Barkovich A, Citrin C. 1987. The MR appearance of syringomyelia: new
observations. *American Journal of Roentgenology* **148**:381–391. doi:10.2214/ajr.148.2.381

Sofroniew MV. 2015. Astrogliosis. *Cold Spring Harb Perspect Biol* **7**:a020420.
doi:10.1101/cshperspect.a020420

Song P, Fogerty J, Cianciolo LT, Stupay R, Perkins BD. 2020. Cone Photoreceptor
Degeneration and Neuroinflammation in the Zebrafish Bardet-Biedl Syndrome 2 (bbs2)
Mutant Does Not Lead to Retinal Regeneration. *Front Cell Dev Biol* **8**:578528.
doi:10.3389/fcell.2020.578528

Sullivan-Brown J, Schottenfeld J, Okabe N, Hostetter CL, Serluca FC, Thiberge SY, Burdine

RD. 2008. Zebrafish mutations affecting cilia motility share similar cystic phenotypes and

suggest a mechanism of cyst formation that differs from pkd2 morphants. *Developmental*

Biology **314**:261–275. doi:10.1016/j.ydbio.2007.11.025

Troutwine BR, Gontarz P, Konjikusic MJ, Minowa R, Monstad-Rios A, Sepich DS, Kwon

RY, Solnica-Krezel L, Gray RS. 2020. The Reissner Fiber Is Highly Dynamic In Vivo and

Controls Morphogenesis of the Spine. *Current Biology* **30**:2353-2362.e3.

doi:10.1016/j.cub.2020.04.015

Tsujikawa M, Malicki J. 2004. Intraflagellar Transport Genes Are Essential for

Differentiation and Survival of Vertebrate Sensory Neurons. *Neuron* **42**:703–716.

doi:10.1016/S0896-6273(04)00268-5

Van Gennip JLM, Boswell CW, Ciruna B. 2018. Neuroinflammatory signals drive spinal

curve formation in zebrafish models of idiopathic scoliosis. *Sci Adv* **4**:eaav1781.

doi:10.1126/sciadv.aav1781

van Rooijen E, Giles RH, Voest EE, van Rooijen C, Schulte-Merker S, van Eeden FJ. 2008.

LRRC50, a conserved ciliary protein implicated in polycystic kidney disease. *J Am Soc*

Nephrol **19**:1128–1138. doi:10.1681/ASN.2007080917

Viau A, Bienaimé F, Lukas K, Todkar AP, Knoll M, Yakulov TA, Hofherr A, Kretz O,

Helmstädter M, Reichardt W, Braeg S, Aschman T, Merkle A, Pfeifer D, Dumit VI, Gubler

M, Nitschke R, Huber TB, Terzi F, Dengjel J, Grahammer F, Köttgen M, Busch H, Boerries

M, Walz G, Triantafyllopoulou A, Kuehn EW. 2018. Cilia‐localized LKB 1 regulates

chemokine signaling, macrophage recruitment, and tissue homeostasis in the kidney. *The*

EMBO Journal **37**:e98615. doi:10.15252/embj.201798615

Vierkotten J, Dildrop R, Peters T, Wang B, Rüther U. 2007. Ftm is a novel basal body protein

of cilia involved in Shh signalling. *Development* **134**:2569–2577. doi:10.1242/dev.003715

Wang J, Thomas HR, Thompson RG, Waldrep SC, Fogerty J, Song P, Li Z, Ma Y, Santra P,

Hoover JD, Yeo NC, Drummond IA, Yoder BK, Amack JD, Perkins B, Parant JM. 2022.

Variable phenotypes and penetrance between and within different zebrafish ciliary transition

zone mutants. *Disease Models & Mechanisms* **15**:dmm049568. doi:10.1242/dmm.049568

Wieczorek S, Combes F, Lazar C, Giai Gianetto Q, Gatto L, Dorffer A, Hesse A-M, Couté Y,

Ferro M, Bruley C, Burger T. 2017. DAPAR & ProStaR: software to perform statistical

analyses in quantitative discovery proteomics. *Bioinformatics* **33**:135–136.

doi:10.1093/bioinformatics/btw580

Wiegering A, Rüther U, Gerhardt C. 2018. The ciliary protein Rpgrip1l in development and

disease. *Developmental Biology* **442**:60–68. doi:10.1016/j.ydbio.2018.07.024

Xie H, Kang Y, Liu J, Huang M, Dai Z, Shi J, Wang S, Li L, Li Y, Zheng P, Sun Y, Han Q,

Zhang J, Zhu Z, Xu L, Yelick PC, Cao M, Zhao C. 2023. Ependymal polarity defects coupled

with disorganized ciliary beating drive abnormal cerebrospinal fluid flow and spine curvature

in zebrafish. *PLoS Biol* **21**:e3002008. doi:10.1371/journal.pbio.3002008

Yu X, Ng CP, Habacher H, Roy S. 2008. Foxj1 transcription factors are master regulators of

the motile ciliogenic program. *Nat Genet* **40**:1445–1453. doi:10.1038/ng.263

Zamanian JL, Xu L, Foo LC, Nouri N, Zhou L, Giffard RG, Barres BA. 2012. Genomic

Analysis of Reactive Astrogliosis. *J Neurosci* **32**:6391–6410.

doi:10.1523/JNEUROSCI.6221-11.2012

Zhang X, Jia S, Chen Z, Chong YL, Xie H, Feng D, Wu X, Song DZ, Roy S, Zhao C. 2018.

Cilia-driven cerebrospinal fluid flow directs expression of urotensin neuropeptides to

straighten the vertebrate body axis. *Nat Genet* **50**:1666–1673. doi:10.1038/s41588-018-0260-

3

Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda

SK. 2019. Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. *Nat Commun* **10**:1523. doi:10.1038/s41467-019-09234-6

MATERIAL AND METHODS

Zebrafish

Wild-type, *rpgrip1l^ex4* and *rpgrip1l^Δ* zebrafish embryos and adults were raised, staged and maintained as previously described (Kimmel et al., 1995). All our experiments were made in agreement with the european Directive 210/63/EU on the protection of animals used for scientific purposes, and the French application decree 'Décret 2013-118'. The projects of our group have been approved by our local ethical committee 'Comité d'éthique Charles Darwin'. The authorization number is APAFIS #31540-2021051809258003 v4. The fish facility has been approved by the French 'Service for animal protection and health' with approval number A750525. All experiments were performed on Danio rerio embryos of mixed AB/TL background. Animals were raised at 28.5°C under a 14/10 light/dark cycle.

Rpgrip1l mutant generation and genotyping

<u>Guide RNA preparation and microinjection</u>
CRISPR target sites were selected for their high predicted specificity and efficiency using the CRISPOR online tool (Haeussler et al., 2016). Real efficiency was assessed on zebrafish embryos by T7E1 test. The two most efficient guides (Rpgrip1l_x4_G1: GCTTACGGTCCTTCACCAGACGG and Rpgrip1l_x25_G3: CCTCAGTTGACAGGTTTCAGCGG) respectively situated 24 nt from the beginning of exon 4 and 82 nt downstream of exon 25 were kept for further experiments.
sgRNA transcription templates were obtained by PCR using T7_Rpgrip1l-x4_G1_Fw primer (5′-GAAATTAATACGACTCACTATAGGCTTACGGTCCTTCACCAGAGTTTTAGAGCTA GAAATAGC- 3′) or T7_Rpgrip1l-x25_G3_Fw (5′-GAAATTAATACGACTCACTATAGGCCTCAGTTGACAGGTTTCAGGTTTTAGAGCT AGAAATAG C-3′) as forward primer and sgRNA_R universal primer (5′-AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTT AACTTGCTA TTTCTAGCTCTAAAAC-3′) as reverse primer. sgRNAs were transcribed using Megascript T7 Transcription Kit (Thermo Fisher Scientific, Waltham, MA) and purified using NucleoSpin® RNA Clean up XS kit (Macherey Nagel, Du☐ren, Germany). sgRNA:Cas9 RNP complex was obtain by incubating

Cas9 protein (gift of TACGENE, Paris, France) (7.5 µM) with sgRNA (10 µM) in 20
mM Hepes-NaOH pH 7.5, 150 mM KCl for 10 min at 28 °C. 1–2 nl was injected per
embryo. For deletion, Rpgrip1l_x4_G1 and Rpgrip1l_x25_G3 RNP complexes were
mixed half and half.
We kept two F1 fish, one with a 15 Kb deletion between exon 4 and 25 of *rpgrip1l*,
that we called *rpgrip1l*$^{\Delta}$ and that we submitted to ZFIN under the name *rpgrip1l*
^{bps1} and a second one with a stop codon within exon 4 called *rpgrip1l* ^{ex4}, that is
submitted to ZFIN under the name *rpgrip1l* ^{bps2}.
991
<u>Screening and genotyping</u>
Injected (F0) fish were screened for germline transmission by crossing with wild type
fish and extracting genomic DNA from resulting embryos. For genomic DNA
extraction, caudal fin (juveniles/adults) or whole embryo DNA were used. Genomic
DNA was isolated with Proteinase K (PK) digestion in 40 of lysis buffer (100 mM Tris-
HCl pH 7.5, 1 mM EDTA, 250 mM NaCl, 0.2% SDS, 0.1 µg/µl Proteinase K (PK)) for
embryos (300 µl for adult fin) overnight at 37°C with agitation. PK was inactivated for
10 min at 90°C and a five-fold dilution was used as template for PCR amplification.
Combined genotyping of wild type and mutant alleles (Supplementary Figure S1A')
was performed by PCR using 3 primers, a common reverse primer for both alleles :
Rpgrip1l_ex25_R3 (GTTGTGTCTCTGCCATATATTG), a specific forward primer for
the deleted allele: Rpgrip1l-ex4-del (CCCACACTGCATACGCACTC) and a specific
forward   primer   for   the   wild   type   allele   :   Rpgrip1l_ex25F
(AGTGTGCGGTACATCTCCAA) at an annealing temperature of 60°C for 35 cycles.
Expected size of the amplified fragment is 364 nt for the wild type allele and 544nt for
the mutant allele. The deleted allele presents a frameshift at position N77 leading to
the generation of a STOP codon after 11 aas, thus leading to a N-terminally
truncated protein at position 88/1256 aas.
For genotyping the *urp2* mutant line, caudal fins from adult fish were extracted in
300µl of lysis solution and PCR was performed as above to detect exon 5 deletion
using the following primers: urp2-F1 (TGATTACTAGCCCTGTCCCAAC) and urp2-R1
(AGGTACAGTACACACGTCACAG).
1014
µCT scans

The samples were scanned on a Bruker micro scanner (Skyscan 1272) with a
resolution of 8.5 µm, a rotation step of 0.55° and a total rotation of 180°. For the
acquisition of adult fish (2.5 cm), a 0.25 mm aluminium filter was used, for a voltage
of 50 kV and an intensity of 180 mA, for juvenile fish (0.9 to 1.2 cm) the filter was
omitted, and a voltage of 60 kV was used with an intensity of 166 mA. Each image
contained 1008 x 672 pixels and was based on the average of 3 images. The 3D
reconstruction by backprojection was carried out by the NRecon software and the 2D
image overlays were then cleaned by the CT Analyser software. The Dataviewer
software allowed all fish skeletons to be oriented in the same way taking the otoliths
as landmarks. The CTvox software then allowed 3D visualizing of the samples.
Morphometric analysis was performed with the CT Analyser software.
1027
Scanning electron microscopy
3 month fish (3 controls, 5 mutants) were euthanized using lethal concentration of
MS222 (0.028 mg/mL). The brains were quickly dissected in 1.22 X PBS (pH 7.4),
0.1 M sodium cacodylate and fixed overnight with 2% glutaraldehyde in 0.61 X PBS
(pH 7.4), 0.1 M sodium cacodylate at 4°C. They were sectioned along the dorsal
midline with a razor blade to expose their ventricular surfaces. Both halves were
washed four times in 1.22 X PBS and post-fixed for 15 minutes in 1.22 X PBS
containing 1% OsO_4. Fixed samples were washed four times in ultrapure water,
dehydrated with a graded series of ethanol and critical point dried (CPD 300, Leica)
at 79 bar and 38 °C with liquid CO_2 as the transition fluid and then depressurized
slowly (0,025 bar/s). They were then mounted on aluminum mounts with conductive
silver cement. Sample surfaces were coated with a 5 nm platinum layer using a
sputtering device (ACE 600, Leica). Samples were observed under high vacuum
conditions using a Field Emission Scanning Electron Microscope (Gemini 500, Zeiss)
operating at 5 kV, with a 20 µm objective aperture diameter and a working distance
around 3 mm. Secondary electrons were collected with an in-lens detector. Scan
speed and line compensation integrations were adjusted during observation.
1045
Histological sections of juvenile and immunofluorescence on sections
Juvenile and adult zebrafish were euthanized using lethal concentration of MS222
(0.028 mg/mL). Pictures and size measurements were systematically taken before

fixation. Fish were fixed in Zamboni fixative [35 ml PFA 4 %, 7.5 ml saturated picric
acid (1.2 %), 7.5 ml 0.2M Phosphate Buffer (PB)] [55] overnight at 4°C under
agitation. Fish were washed with Ethanol 70% and processed for dehydration by
successive 1 h incubation in Ethanol (3 x 70% and 2 x 100%) at room temperature
under agitation, then for paraffin inclusion. 14 µm sagittal paraffin sections were
obtained using a Leica RM2125RT microtome. Sections were deparaffinized and
antigen retrieval was performed by incubation for 7 min in boiling citrate buffer (10
mM, pH 6). Immunofluorescence staining was performed as described previously
(Andreu-Cervera et al., 2019). The following primary antibodies were used: anti-RF,
anti-Acetylated Tubulin, anti-Glutamylated Tubulin, anti-Arl13b, anti-LCP1, anti-
AnnexinA2. Corresponding primary and secondary antibodies are described and
referenced in the Resources Table.
1061

Immunofluorescence on whole embryos
Embryos from 24 to 40 hpf were fixed 4 hr to overnight in 4% paraformaldehyde
(PFA) at 4°C. For Reissner fiber staining, larvae at 48 and 72 hpf were fixed 2 hr in
4% PFA and 3% sucrose at 4°C, skin from the rostral trunk and yolk were removed.
Samples from 24 to 40 hpf embryos were blocked overnight in a solution containing
0.5% Triton, 1% DMSO, 10% normal goat serum and 2 mg/mL BSA. For older
samples (48 to 72 hpf) triton concentration was increased to 0.7%. Primary
antibodies were incubated one to two nights at 4°C in a buffer containing 0.5% Triton,
1% DMSO, 1% NGS and 1 mg/mL BSA. All secondary antibodies were from
Molecular Probes, used at 1:400 in blocking buffer, and incubated a minimum of 2.5
hr at room temperature. The primary antibodies chosen for in toto immuno-labeling
against Reissner fiber, Acetylated-tubulin, Myc, Arl13b and Gamma-tubulin as well as
the corresponding secondary antibodies are referenced in the Key Resources Table.
Whole mount zebrafish embryos (dorsal or lateral mounting in Vectashield Mounting
Medium) were imaged on a Leica SP5 confocal microscope and Zeiss LSM910
confocal microscope, both equipped with a 63X immersion objective. Images were
then processed using Fiji (Schindelin et al., 2012).
1079

Whole-mount brain clearing

Brains were dissected from 4-5 wpf size-matched (0.9-1.2 mm length) zebrafish after
in toto fixation with formaldehyde. Whole–mount tissue clearing was performed
following the zPACT protocol (Affaticati et al., 2017). In brief, brains were infused for
2 days in hydrogel monomer solution (4% acrylamide, 0.25% VA- 044, 1%
formaldehyde and 5% DMSO in 1X PBS) at 4°C. Polymerization was carried out for
2h30 at 37°C in a desiccation chamber filled with pure nitrogen. Brains were
transferred into histology cassettes and incubated in clearing solution (8% SDS and
200 mM boric acid in dH2O) at 37°C with gentle agitation for 8 days. Cleared brains
were washed in 1X PBS with 0.1% Tween-20 (PBT) for 3 days at room temperature
and kept in 0.5% formaldehyde, 0.05% sodium azide in PBT at 4°C until further
processing. Brains were subsequently placed for 1h in depigmentation pre-incubation
solution (0.5X SSC, 0.1% Tween-20 in dH2O) at room temperature. The solution was
replaced by depigmentation solution (0.5X SSC, Triton X-100 0.5%, formamide
0.05% and H2O2 0.03% in dH2O) for 45 minutes at room temperature. Depigmented
brains were washed for 4h in PBT and post-fixed (2% formaldehyde and 2% DMSO
in PBT) overnight at 4°C.
1097

Cleared brain immunostaining of whole adult brains
Whole-mount immunolabeling of cleared brains was performed as described in the
zPACT protocol with slight modifications. Briefly, brains were washed in PBT for one
day at room temperature and blocked for 10h in 10% normal goat serum, 10%
DMSO, 5% PBS-glycine 1M and 0.5% Triton X-100 in PBT at room temperature.
Brains were washed again in PBT for 1h prior to incubation with anti- ZO-1 antibody
(ZO1-1A12, Thermofisher, 1:150) in staining solution (2% normal goat serum, 10%
DMSO, 0.1% Tween-20, 0.1% Triton X-100 and 0.05% sodium azide in PBT) for 12
days at room temperature under gentle agitation. Primary antibody was renewed
once after 6 days of incubation. Samples were washed three times in PBT and
thereafter incubated with Alexa Fluor 488-conjugated secondary antibody (A11001,
Invitrogen, 1:200) for 10 days in the staining solution at room temperature under
gentle agitation. Secondary antibody was renewed once after 5 days of incubation.
Samples were washed three times in PBT prior to a counterstaining with DiIC18
(D282, Invitrogen, 1µM) in the staining solution for three days at room temperature

with gentle agitation. Samples were washed in PBT for 3h before mounting
procedure.

1115

Mounting and confocal imaging

Samples were placed in 50% fructose-based high-refractive index solution (fbHRI,
see (Affaticati et al., 2017) /50% PBT for 1h, then in 100% fbHRI. Brains were
mounted in agarose-coated (1% in standard embryo medium) 60 mm Petri dish with
custom imprinted niches to help orientation. Niche-fitted brains were embedded in
1% phytagel and the Petri dish filled with 100% fbHRI. fbHRI was changed three
times before imaging until its refractive index matched 1.457. Images were acquired
with a Leica TCS SP8 laser scanning confocal microscope equipped with a Leica HC
FLUOTAR L 25x/1.00 IMM motCorr objective. Brains were scanned at a resolution of
1.74x1.74x1.74 µm (xyz) and tiled into 45 to 70 individual image stacks, depending
on brain dimensions, subsequently stitched, using LAS X software.

1127

Volumetric analysis of the posterior ventricles

The volumes of the posterior ventricles were segmented, reconstructed and analyzed
using Amira for Life & Biomedical Sciences software (Thermo Fisher Scientific). In
essence the ventricles volumes were manually segmented in Amira's segmentation
editor and subsequently refined by local thresholding and simplification of the
corresponding surfaces. Volumes, which were open to the environment were
artificially closed with minimal surfaces by connecting the distal-most points of their
surface to the contralaterally corresponding points using straight edges. Due to the
biologic variability of the sample population the overall size of the specimens needed
to be normalized to keep the measured volumes comparable. For this spatial
normalization one of the specimens was randomly selected from the wildtype
population as template (1664, grey) and the 'Registration' module in Amira was used
to compute region-specific rigid registrations for the other specimens, allowing for
isotropic scaling only [details in supp. mat.]. For excluding the influence of the
ventricular volumes, the registration was computed on the basis of the independent
reference stain (DiIC18) (details in supp. mat.). Region specific volume differences
between the mutant and wildtype population were evaluated on seven subvolumes of
the posterior ventricles (details in supp. mat.).

1146

RNA extraction for transcriptome analysis and quantitative RT-PCR

1147

Juvenile and adult zebrafish were euthanized using lethal concentration of MS222

(0.28 mg/mL). Their length was measured and their fin cut-off for genotyping. For

transcriptomic analysis, brain and dorsal trunk from 1 month juvenile (0.9 to 1.0 cm)

zebrafish were dissected in cold PBS with forceps and lysed in QIAzol (QIAGEN)

after homogenization with plunger pistons and 1ml syringes. Samples were either

stored in QIAzol at -80°C or immediately processed. Extracts containing RNA were

loaded onto QIAGEN-mini columns, DNAse digested and purified in the miRNAeasy

QIAGEN kit (Cat 217004) protocol. Samples were stored at -80°C until use. RNA

concentration and size profile were obtained on the Tapestation of ICM platform. All

preparations had a RIN above 9.2. For Q-PCR from juveniles, whole fish minus

internal organs were lysed in Trizol (Life technologies) using the Manufacturer

protocol.

1160

Quantitative PCR from individual juvenile or adult fish

1161

The cDNA from the isolated RNAs was obtained following GoScript Reverse

Transcription System protocol (Promega), using 3 to 4 µg of total RNA for each

sample. The 20 µL RT-reaction was diluted 4-fold and 4µl was used for each

amplification performed in duplicates. The primers used for qPCR were the following:

urp2 (F: ACCAGAGGAAACAGCAATGGAC; R: TGAGGTTTCCATCCGTCACTAC),

urp1 (F: ACATTCTGGCTGTGGTTTGTTC; R: CTCTTTTGCACCTCTCTGAAGC),

urp (F: GCGGAAAAATGTCATGCCTCTTC; R: TTGAGCTCCTTTTGAAGCTCCTG),

uts2a (F: CACTGCTCAACAGAGACAGTATCA; R:

CCAAAAGACCACTGGGAGGAAC), *uts2b* (F: TACCCGTCTCTCATCAGTGGAG; R:

TTTTCCAGCAAGGCCTCTTTTAC) where all of them are from Gaillard et al. 2023,

rpgrip1l (F: CAGACACCTGCTGGAGTTACA; R: TCCTGACTCACATCAAACGCA),

irg1l (F: TCCCTAAGAGGTTCCATCCTCC; R: AAGATGCAGCGATGGCCAAA),

stat3 (F: CAGGACTGGGCGTATGCG; R: GAAGCGGCTGTACTGCTGAT), *c4b* (F:

GGGTGTTCTTATGGCGGTGG and R: GCGCACAACAAGCTTTTCTCATC) *lsm12b*

(F: GAGACTCCTCCTCCTCTAGCAT and R: GATTGCATAGGCTTGGGACAAC).

The Q-PCR were performed using the StepOne real-time PCR system and following

its standard amplification protocol (Thermo Scientific). Relative gene expression

levels were quantified using the comparative CT method ($2^{-\Delta\Delta CT}$ method or $2^{-\Delta CT}$) on
the basis of CT values for target genes and *lsm12b* as internal control.

1181

RNA-sequencing and analysis

RNA-seq libraries were constructed by the ICM Platform (PARIS) from 100 ng of total
RNA using the KAPA mRNA Hyperprep kit (Roche) that allows to obtain stranded
polyA libraries, and their quality controlled on Agilent Tape station. The sequencing
was performed on a NovaSeq 6000 Illumina for both strands. Sequences were
aligned against the GRCz11 version of zebrafish genome using **RNA Star function**
in "Galaxy" environment. The 12 Brain libraries reached between 22 to 26 Millions of
assigned reads, and the trunk libraries between 26 to 35 Millions of assigned reads.
Expression level for each gene was determined using **Feature counts function** from
RNA Star BAM files. **DESeq2 function** was used to calculate the log2 Fold change
and adjusted P values from the Wald test for each gene comparing the 7 mutants
raw count collection to the 5 controls raw counts collection. Volcano plots were
generated using the Log2 fold change and P adj. value of each Gene ID from the
DESeq2 table. A threshold of 5.10 E-2 was chosen for the Padj. values and a log2
fold change > to +0.75 or < to -0.75 to select significantly up or down-regulated
regulated genes. Both gene lists were used as input to search for enriched GO terms
and KEGGs pathways using Metascape software (Zhou et al., 2019) as well as to
compare up-regulated genes in the dorsal trunk versus the brain using Galaxy.

1200

Quantitative proteomics

Sample preparation: Adult zebrafish were euthanized using lethal concentration of
MS222 (0.28 mg/mL). For proteomic analysis, brain from 3 months zebrafish were
dissected in cold PBS with forceps and immediately flash frozen in liquid nitrogen
and stored at -80°C. Zebrafish brain samples were lysed in RIPA buffer (Sigma) with
1/100 antiprotease and sonicated for 5 minutes. After a 660 nm protein assay
(Thermo), samples were digested using a Single Pot Solid Phase enhanced Sample
Preparation (SP3) protocol (Hughes et al., 2014). In brief, 40 μg of each protein
extract was reduced with 12mM dithiothreitol (DTT) and alkylated using 40 mM
iodoacetamic acid (IAM). A mixture of hydrophilic and hydrophobic magnetic beads
was used to clean-up the proteins at a ratio of 20:1 beads:proteins (Sera-Mag Speed

beads, Fisher Scientific). After addition of ACN to a final concentration of 50%, the

beads were allowed to bind to the proteins for 18 minutes. Protein-bead mixtures

were washed twice with 80% EtOH and once with 100% ACN. The protein-bead

complexes were digested with a mixture of trypsin:LysC (Promega) at a 1:20 ratio

overnight at 37°C. Extracted peptides were cleaned-up using automated C18 solid

phase extraction on the Bravo AssayMAP platform (Agilent Technologies).

1218

NanoLC-MS-MS analysis

The peptide extracts were analysed on a nanoLC-TimsTof Pro coupling (Bruker

Daltonics). The peptides (200ng) were separated on an IonOpticks column (25cm X

75µm 1.6µm C18 resin) using a gradient of 2-37% B (2%ACN, 0.1%FA) in 100

minutes at a flow rate of 0.3 µl/min. The dual TIMS had a ramp time and

accumulation time of 166 ms resulting in a total cycle time of 1.89s. Data was

acquired in Data Dependent Acquisition-Parallel Accumulation Serial Fragmentation

(DDA-PASEF) mode with 10 PASEF scans in a mass range from 100 m/z to 1700

m/z. The ion mobility scan range was fixed from 0.7-1.25 Vs/cm^2. All samples were

injected using a randomized injection sequence. To minimize carry-over, one solvent

blank injection was performed after each sample.

1230

Data interpretation

Raw files were converted to .mgf peaklists using DataAnalysis (version 5.3, Bruker

Daltonics) and were submitted to Mascot database searches (version 2.5.1,

MatrixScience, London, UK) against a *Danio rerio* protein sequence database

downloaded from UniProtKB-SwissProt (2022_05_18, 61 732 entries, taxonomy ID:

7955), to which common contaminants and decoy sequences were added. Spectra

were searched with a mass tolerance of 15 ppm in MS mode and 0.05 Da in MS/MS

mode. One trypsin missed cleavage was tolerated. Carbamidomethylation of cysteine

residues was set as a fixed modification. Oxidation of methionine residues and

acetylation of proteins' n-termini were set as variable modifications. Identification

results were imported into the Proline software (version 2.1.2,

http://proline.profiproteomics.fr) (Bouyssié et al., 2020) for validation. Peptide

Spectrum Matches (PSM) with pretty ranks equal to one were retained. False

Discovery Rate (FDR) was then optimized to be below 1% at PSM level using

Mascot Adjusted E-value and below 1% at protein level using Mascot Mudpit score. For label free quantification, peptide abundances were extracted without cross assignment between the samples. Protein abundances were computed using the sum of the unique peptide abundances normalized at the peptide level using the median.

To be considered, proteins must be identified in at least four out of the five replicates in at least one condition. Imputation of the missing values and differential data analysis were performed using the open-source ProStaR software (Wieczorek et al., 2017). Imputation of missing values was done using the approximation of the lower limit of quantification by the 2.5% lower quantile of each replicate intensity distribution ("det quantile"). A Limma moderated t-test was applied on the dataset to perform differential analysis. The adaptive Benjamini-Hochberg procedure was applied to adjust the p-values and FDR. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol et al., 2022), with the dataset identifier PXD042283.

Embryos drug treatment

27 hpf embryos from *dnaaf1*[tm317b/+] incross were dechorionated manually. N-acetyl cysteine ethyl ester (NACET) (BIOLLA Chemicals #59587-09-6) was prepared extemporaneously at the final concentration of 3 mM in pure water (pH: 7.2; conductivity 650 µS). Embryos were treated between 27 hpf to 60 hpf and embryos were anesthetized before being imaged laterally.

Juvenile drug treatment and preparation of treated samples

Two hundred larvae were housed off-system in 1.8 liter tanks with 20 fish per tank, being fed twice a day throughout the experiment. 100 were treated with NACET (BIOLLA Chemicals #59587-09-6), which was prepared extemporaneously at 1.5 mM (286.5mg/L) in fish water (pH 7.2, Conductivity 650 µS) which was changed once per day. Fish were treated from 30 dpf to 84 dpf and monitored for curvature onset once per week. At 85 dpf, fish were euthanized, imaged to measure the curvature index. Half of the fish were fixed for histology analysis in zamboni fixative [35 ml PFA 4 %, 15 ml saturated picric acid (1.2 %)] overnight at 4°C under agitation, while the other half was processed for RT-qPCR analysis. To prepare dorsal trunk RNA, internal

1278 organs were removed in cold PBS 1X, and the remaining tissue was cut in small
1279 pieces before lysis in QIAzol (QIAGEN). Homogenization was achieved using
1280 plunger pistons and passages through 1ml syringes.
1281
Transgenesis
The *Tol2 -5.2foxj1a:5xmyc-RPGRIP1L* plasmid was produced using the Gateway
system by combining four plasmids. P5E-foxj1a enhancer was kindly donated by [3].
5xmyc-RPGRIP1L cDNA was amplified from the plasmid pCS2-5xmyc-RPGRIP1L
(Mahuzier et al., 2012) using CloneAmp HiFi PCR Premix (Takara # 639298) using
primers : forward 5'-
GGGGACAAGTTTGTACAAAAAGCAGGCTGCAGGATCCCATCGATTTAAAGCT-3'
and reverse 5'
GGGGACCACTTTGTACAAGAAAGCTGGGTCTCCGAGCTCGGTTCAAGCCTCCA
AGTCATCTCTGT-3. The PCR product was gel purified using QIAEX II gel extraction
kit (QIAGEN, #20021) . BP recombination (Invitrogren "Gateway BP Clonase II
Enzyme Mix" #11789-020) was performed into pDONR221 #218 to generate pMe-
5xmyc-RPGRIP1L. The 3' entry polyadenylation signal plasmid was the one
described in [61]. The 3 plasmids were shuttled into the backbone containing the
cmcl2:GFP selection cassette (Kwan et al., 2007) using the LR recombination
(Invitrogen "Gateway LR Clonase II Plus Enzyme Mix" #12538-120). *rpgrip1l*$^{+/-}$ X AB
embryos outcrosses were injected at the one cell stage with 1nl of a mix containing
20 ng/µl plasmid and 25 ng/µl Tol2 transposase RNA and screened at 48 hpf for GFP
expression in the heart. Some F0 founders produced F1 zebrafish carrying one copy
of the transgene that were further checked for RPGRIP1L specific expression in
foxj1a territory, using Myc labelling at the base of FP cilia in 2,5 dpf embryos.
The Tol2-1.7kb col2a1a:5xMyc-RPGRIP1L plasmid was produced using the Gateway
System by combining four plasmids. Col2a1a enhancer was amplified from the
plasmid -1.7kbcol2a1a:EGFP-CAAX (Dale and Topczewski, 2011) using CloneAmp
HiFi premix (Takara #639298) with the primers: forward 5'-GGG GAC AAC TTT GTA
TAG AAA AGT TGG CCC TCT GAC ACC TGA TGC CAA TTG C-3' and reverse 5'-
GGG GAC TGC TTT TTT GTA CAA ACT TGC TTG CAG GTC CTA AGG GGT GAA
AGT CG-. The PCR product was gel purified and BP recombination was performed
into pDONR_P4-P1R-#219. The middle entry plasmid and the 3' plasmid were the

same as those used to generate the Tol-5.2foxj1a: 5xmyc-RPGRIP1L plasmid. F1
transgenic were first selected for GFP expression in the heart and then on Myc
labelling within notochordal cells at 2,5 dpf within their embryonic progeny.

Skeletal preparations, Alizarin staining and imaging

Animals were euthanized using a lethal concentration of MS222 (0.28 mg/mL), skin
was removed and fixation was performed with 4% paraformaldehyde overnight at
4°C. After evisceration, samples were incubated in borax 5%, rinsed several times in
1% KOH and incubated in solution composed with Alizarin 0.01% (Sigma, A5533)
and KOH 1% during 2 days. Fish were rinsed several times in KOH 1% and
incubated in trypsin 1% and borax 2% for 2 days until cleared. Samples were rinsed
several times in distilled water and transferred in glycerol 80% (in KOH 1%) using
progressive dilutions. Samples were kept in glycerol 80% until imaging.

Cobb angles measurements

To quantitatively evaluate the severity of spine curvature in *rpgrip1l$^{+/-}$; urp2$^{+/-}$*
incrosses, we used the zebrafish skeletal preparations stained with Alizarin. We drew
parallel lines to the top and bottom most displaced vertebrae on lateral and dorsal
views. The Cobb angle was then measured as the angle of intersection between
lines drawn perpendicular to the original 2 lines. This was conducted using FIJI
software. For each fish, the total Cobb angle was calculated by summing all Cobb
angles.

Curvature index measurements

We implemented a novel procedure to quantify curvatures on both axes by drawing a
line along the body of the fish as shown in Figure supplementary 4I and curvature
was calculated in MATLAB (code available on demand). We decomposed the line in
a series of equidistant points and we measured the curvature at each of these points
using the LineCurvature2D function. We then added the absolute values of these
measures, which are expressed as an angle (in radian) per unit of length. The higher
the sum is for a given line, the more this line is curvated. We verified the accuracy
of this metric by comparing it with the visual assessment of curvatures by 3
independent observers. For *rpgrip1l$^{+/-}$; urp2$^{+/-}$* incross and NACET experiment, a line

1344 following the body deformation was drawn from the mouth to the base of the tail
1345 following the midline of the fish in lateral position and another one along the dorsal
1346 axis. Curvature index of both curves were summed for each fish. Curvature analysis
1347 was performed blinded to fish genotype. *rpgri1l*$^{+/+}$ or *rpgrip1l*$^{+/-}$ siblings were used as
1348 controls.

1349

Data acquisition for body-curvature analysis at embryonic stage
Zebrafish embryos were anesthetized and imaged laterally with the head pointing to
the left. The angle between the line connecting the center of the eye to the center of
the yolk and the line connecting the center of the yolk to the tip of the tail was
measured to evaluate the body curvature of the embryos. For quantitative analysis,
the angles of each embryo were put in the same concentric circles represented with
a Roseplot, with 0° pointing to the right and 90° pointing to the top. Each triangle
represents a 30° quadrant, and its height indicates the number of embryos within the
same quadrant.

1359

Quantification, statistical analysis and figure preparation
For all experiments the number of samples analyzed is indicated in the text and/or
visible in the figures. Statistical analysis was performed using the Prism software.
****: P value < 0.0001; ***: P value < 0.001; **: P value < 0.01; *: P value < 0.05.
Graph were made using Prism and Matlab and figures were assembled using
Photoshop software.

1366

1367 **REAGENT AND RESOURCE TABLE**

Reagent type or resource	Designation	Source or reference	Identifiers	Additional Infos
Antibody (antiserum)	Rabbit polyclonal IgG anti-bovine	Didier et al 1995	Courtesy of Dr. S.Gobron	1:200

	Reissner Fiber			
Antibody	Mouse monoclonal IgG1k anti ZO1 -1A12	Thermofisher	#33-9100	1:150
Antibody	Mouse monoclonal IgG2b anti-acetylated-tubulin (clone 6-11B-1)	Sigma-Aldrich	#T 6793 RRID: AB_477585	1:400
Antibody	Mouse monoclonal IgG1 anti glutamylated Tubulin	Adipogen	#AG-20B-0020-C100	1:500
Antibody	Rabbit polyclonal anti-Arl13b	Proteintech	#17711-1-AP	1:200
Antibody	Mouse monoclonal IgG1 anti-GFAP	Sigma	#G3893	1:400
Antibody	Chicken monoclonal IgY anti-GFP	AVES lab	#GFP-1020	1:200
Antibody	Mouse monoclonal	Cell signaling	#2276	1:200

	IgG2a anti-myc (clone 9b11)			
Antibody	Rabbit polyclonal anti-LCP1	GeneTex	#GTX124420	1:200
Antibody	Mouse monoclonal IgG1 anti-gamma-tubulin	Sigma	#T6557	1:500
Antibody	Mouse polyclonal anti-annexin A2	Proteintech	#11256-1-AP	1:200
Antibody	Goat anti-mouse IgG1 Alexa633	Molecular probes	# A-21126 RRID:AB_2535768	1:400
Antibody	Goat anti-mouse IgG1 Alexa568	Molecular Probes	# A-21124 RRID: AB_2535766	1:400
Antibody	Goat anti-mouse IgG2a Alexa568	Molecular probes	# A-21134 RRID:AB_2535773	1:400
Antibody	Goat anti-mouse IgG2a Alexa488	Molecular probes	# A-21131 RRID:AB_141618	1:400

Antibody	Goat anti-mouse IgG2b Alexa633	Molecular probes	# A-21146 RRID:AB_2535782	1:400
Antibody	Goat anti-mouse IgG2b Alexa568	Molecular probes	# A-21144 RRID: AB_2535780	1:400
Antibody	Goat anti-rabbit IgG Alexa568	Molecular probes	# A-11011 RRID:AB_143157	1:400
Antibody	Goat anti-rabbit IgG Alexa488	Molecular probes	# A-11008 RRID: AB_143165	1:400
Antibody	FITC donkey Anti-Chicken IgY	Jackson ImmunoResearch	# 703-096-155	1:200
Chemical	Vectashield	Vector Laboratories	H-1000	
Strain (Danio rerio)	zebrafish wild-type AB or (TL x AB) hybrid strains	N/A	N/A	
Strain (Danio rerio)	Zebrafish *rpgrip1l$^{\Delta}$* mutant strain	This manuscript, Zfin:*rpgrip1l^{bps1}*		
Strain (Danio rerio)	Zebrafish *rpgrip1l^{ex4}*	This manuscript,		

	mutant strain	Zfin:*rpgrip1l* [bps2]		
Strain (Danio rerio)	Zebrafish *urp2*[+/-]	(Gaillard et al., 2023)	ZDB-ALT-230207-5	
Strain (Danio rerio)	Zebrafish *dnaaf1*[tm317b/+]	(van Rooijen et al., 2008)		
Strain (Danio rerio)	Zebrafish Scospondin-GFP[ut24]	(Troutwine et al., 2020)		
Recomb. DNA	Foxj1 promoter-enhancer sequence	(Dale and Topczewski, 2011)		
Software	LASX	Leica		
Software	Amira for Life & Biomedical Sciences	Thermo Fisher Scientific		
Software	CRISPOR	(Haeussler et al., 2016)	http://crispor.tefor.net/	
Software	Fiji/ImageJ	(Schindelin et al., 2012)	https://imagej.net/Fiji/Downloads	
Software	PRISM	GraphPad	https://www.graphpad.com/	
Software	Metascape		https://metascape.org/gp/index.html#/main/step1	
Software	Matlab	The Mathworks, Inc.	https://fr.mathworks.com/products/matlab.	

			html	
Software	NRecon reconstruction software	Micro Photonics Inc.		
Software	CTvox	Bruker		
Software	CT Analysis	Bruker		
Other				
Refractometer	Roth Sochiel EURL	DR 301-95		
Confocal microscope	Leica	SP5		
Confocal microscope	Zeiss	LSM 980 Inverted		
Confocal microscope	Zeiss	LSM 710		
Macroscope	Zeiss	AXIOZOOM V16		
Micro-scanner	Bruker	Skyscan 1272		
Microtome	Leica	RM2125RT		
Q-PCR apparatus	Applied Biosystems	Step One plus		

1368

1369

1370

1371

1372

1373

1374

1375

FIGURE LEGENDS

1377

Figure 1: *rpgrip1l^{-/-}* juvenile fish develop scoliosis, which is rescued by RPGRIP1L expression in *foxj1a*-positive cells.

(A) Representative *rpgrip1l^{-/-}* fish at 5 days post-fertilization (dpf) (4 mm body length, larvae), 5 weeks pf (wpf) (1.2 cm body length, juveniles) and 12 wpf (2.5 cm body length, adults), showing absence of morphological defects in embryos, onset of spine curvature (tail-up) in juveniles and scoliosis in adults. **(B-C)** Graph showing the time course of scoliosis appearance **(B)** in *rpgrip1l^{-/+}* incrosses (total 4 clutches, 252 fish) and scoliosis penetrance **(C)** in adults. **(D-E)** Micro-computed tomography (µCT) scans of control siblings (D) and *rpgrip1l^{-/-}* (E) adult fish. 4 fish were analyzed for each condition. **(F-G)** Dorsal and lateral superimposed µCT views of the spines of one control (yellow) and two *rpgrip1l^{-/-}* (pink, cyan) adult fish illustrating the 3D spine curvatures in mutants. **(H)** Schematic representation of the two rescue constructs. **(I)** Table presenting the number and phenotype of *rpgrip1l^{-/-}* fish generated for both rescue experiments by stable transgenesis **(J)** Graph representing scoliosis penetrance in transgenic and non-transgenic *rpgrip1l^{-/-}* siblings.

1393

Figure 2 Altered ciliogenesis at trunk and subcommissural organ (SCO) levels.

(**A-C'''**) Immunostaining of trunk sagittal sections at the level of the spinal cord central canal in *rpgrip1l^{+/+}* (A-A'''; n=5), straight *rpgrip1l^{-/-}* (B-B'''; n=4) and scoliotic *rpgrip1l^{-/-}* fish (C-C'''; n=6)) at 8 weeks (N=2 independent experiments). Cilia were immunostained with Arl13b (cyan, A'', B'', C'') and glutamylated-tubulin (magenta, A''', B''', C'''). Panels A-C' are merged images with DAPI. Panels A', B' and C' show higher magnification of the squared regions in A, B and C, respectively. Scale bars: 10 µm in A, 5 µm in A'. **(D)** Violin plot showing the distribution of cilia length along the central canal of *rpgrip1l^{+/+}* (n=300 cilia from 5 fish), straight *rpgrip1l^{-/-}* (n=267 from 4

fish) and scoliotic *rpgrip1l^{/-}* (n=414 from 6 fish) juvenile fish (N=2). **(E)** Violin plot showing the distribution of cilia density along the central canal of juvenile *rpgrip1l^{+/+}* (n=5), straight *rpgrip1l^{/-}* (n=4) and scoliotic *rpgrip1l^{/-}* fish (n=6, N=2). Density is the average number of ventral and dorsal cilia lining the CC over a length of 100 µm from 3 to 5 sections for each fish. Each dot represents the mean cilia length (D) or cilia density (E) of 4-8 sections analyzed at different antero-posterior levels. Statistical analysis was performed using Tukey's multiple comparisons test where * means P-value < 0.05, and ** means P-value < 0.01. **(F)** Violin plot showing the distribution of width of the central canal (nuclear-free region) (in µm) of juvenile *rpgrip1l^{+/+}* (n=5), straight *rpgrip1l^{/-}* (n=4) and scoliotic *rpgrip1l^{/-}* fish (n=6, N=2). Each dot represents the mean of 3 measurements made at different AP levels. Statistical analysis was performed using Tukey's multiple comparisons test where ns means non-significant, * means P-value < 0.05. **(G-L)** Immunostaining of cilia on transverse sections of the brain at SCO level using glutamylated-tubulin and acetylated-tubulin antibodies (both in white), in *rpgrip1l^{+/+}* (n=6) (G, G', H), straight *rpgrip1^{/-}* (n=3) (I, I', J) and scoliotic *rpgrip1^{/-}* (n=7) (K, K', L) 8 wpf fish (N=2). Nuclei are stained with DAPI (blue). G', I' and K' are higher magnifications of squared regions in G, I and K, respectively. In H, J and L the bottom-right inset represents a higher magnification of the squared region. Scale bars in G and G': 10 µm.

Figure 3: *rpgrip1^{-/-}* juveniles show ventricular dilations and loss of the Reissner Fiber at scoliosis onset.

(A) Reconstruction of the RhV at the level of posterior midbrain and hindbrain in a transparised 5 wpf control zebrafish brain, stained with ZO1 antibody (ventricular surface) and DiI (global brain shape). Several brain regions are annotated on the reconstruction where Tel means telencephalon, ChP forebrain choroid plexus, SCO subcommissural organ, TeO optic tectum, CCe corpus cerebelli, CC crista cerebellaris and Rh rhombencephalon. **(B-D)** Cyan circles represent the ROIs measured in F. The orange line in D indicates the level of optical sections in E. **(E)** Optical transverse sections showing the caudal part of reconstructed ventricles of a control and two *rpgrip1l^{/-}* fish, one straight and two tail-up. **(F)** Graphs of the ventricle volume at the onset of scoliosis in 3 control and 4 *rpgrip1l^{/-}* (one straight and three tail-up) fish. The green dot corresponds to the straight mutant fish. Each dot

represents a fish. Statistical analysis was performed using unpaired t-test where ns
means non-significant, and * means P-value < 0.05. **(G)** Graph of the surface of the
optical sections as illustrated in E. The green dot corresponds to a straight mutant
fish. Each dot represents a fish. Statistical analysis was performed using unpaired t-
test where ns means non-significant, and * means P-value < 0.05. **(H)** Schematic
representation of a portion of the spinal cord central canal in a lateral view of the fish
trunk, showing the Reissner Fiber (RF), and of the SCO in a transverse view of the
brain at the level of diencephalic Ventricle (DiV). **(I-Q)** Visualization of the GFP
fluorescence in the *sspo-GFP*$^{uts24/+}$ transgenic line (fire LUT) in sagittal sections of the
trunk (I-K) (N=3) and transverse sections of the brain at the level of the SCO (N=2)
(L-Q) in juvenile fish. The corresponding fish are 8 wpf *rpgrip1l*$^{+/+}$ (I, L, O) (n=10);
straight *rpgrip1l*$^{/-}$ (J, M, P) (n=9) and scoliotic *rpgrip1l*$^{/-}$ (K, N, Q) (n=8). L-N and O-Q
show sections at anterior and posterior SCO levels, respectively. Scale bars: 5 µm in
I-K, 10 µm in L-Q.

1450

Figure 4: Upregulation of *urp1/2* expression does not contribute to axis
curvature of *rpgrip1l*$^{/-}$ fish.

(A) Quantitative reverse transcription polymerase chain reaction (qRT-PCR)
assaying *urp2* expression level in *rpgrip1l*$^{/-}$ and *rpgrip1l*$^{+/+}$ (WT) siblings. Each dot
represents the value obtained for one fish. Green and purple dots represent straight
and scoliotic *rpgrip1l*$^{/-}$ fish, respectively. qRT-PCR analysis was performed at 5 wpf
(juvenile stage, 5 controls and 7 *rpgrip1l*$^{/-}$) and 12 wpf (adult stage, 5 controls and 9
rpgrip1l$^{/-}$). *urp2* expression levels are increased by approximately 3.5-fold in *rpgrip1l*
$^{/-}$ at both stages. Statistical analysis was performed using unpaired t-test where **P <
0.01. Error bars represent s.d. **(B)** Quantification of body axis curvature from dorsal
and lateral views in 4 mpf fish issued from [*rpgrip1l*$^{+/-}$, *urp2*$^{+/-}$] incrosses. Each point
represents the value measured for one fish. Statistical analysis was performed using
Tukey's multiple comparisons test where **** P < 0,0001 with error bars represent
s.d. n=9 [*rpgrip1l*$^{+/+}$; *urp2*$^{+/+}$], 7 [*rpgrip1l*$^{+/+}$; *urp2*$^{/-}$], 27 [*rpgrip1l*$^{/-}$; *urp2*$^{+/+}$], 48 [*rpgrip1l*
$^{/-}$; *urp2*$^{+/-}$] and 27 [*rpgrip1l*$^{/-}$; *urp2*$^{/-}$] fish. **(C)** Representative [*rpgrip1l*$^{/-}$; *urp2*$^{+/+}$] and
sibling [*rpgrip1l*$^{/-}$; *urp2*$^{/-}$] fish on lateral and dorsal views at 12 wpf. Scale bar 5mm.
(D, E) Expression levels of *urp2* (D) and *urp1* (E) in adult fish (12 wpf). Each dot
represents the value for one fish. Green and purple dots represent straight and

scoliotic *rpgrip1l^{-/-}* fish, respectively (n=5 [*rpgrip1l^{+/+}; urp2^{+/+}*], 9 [*rpgrip1l^{+/+}; urp2^{-/-}*], 11

[*rpgrip1l^{-/-}; urp2^{+/-}*] and 10 [*rpgrip1l^{-/-}; urp2^{-/-}*]). Each gene expression level is

compared to the *lsm12b* housekeeping gene. Statistical analysis was performed

using Tukey's multiple comparisons test where ns means non-significant and ** P <

0.01, **** P < 0.0001. Error bars represent s.d.

1474

Figure 5: Comparative transcriptome and proteome analysis of control and

rpgrip1l^{-/-} fish.

(A-B) Volcano-plots showing differentially regulated genes in the transcriptomes of

the brain (A) and dorsal trunk (B) comparing 5 controls (2 wt, 3 *rpgrip1l^{/+}*) and 7

rpgrip1l^{/-} (6 tail-up and 1 straight) fish at 5 wpf (0.9 cm body length). Each dot

corresponds to one gene. A gene was considered deregulated if its associated P-adj.

value was inferior to 0,05. Red and blue dots represent up-regulated (Log2FC >

0.75), and down-regulated (Log2FC < -0.75) genes, respectively. The names of the

top 25 most deregulated genes are boxed. **(C)** GO term analysis of biological

processes in the brain samples using Metascape. **(D)** Venn diagram of the

upregulated genes in brain and trunk samples, showing many common genes

between the two regions. **(E)** Selected genes of interest upregulated in the trunk

and/or brain of *rpgrip1l^{/-}* fish. The four genes in red were also found upregulated in

the proteome of adult *rpgrip1l^{/-}* fish. **(F)** Volcano-plot showing differentially expressed

proteins in the brain of 5 *rpgrip1l^{+/+}* fish versus 5 *rpgrip1l^{/-}* adult scoliotic fish (3

months pf). Each dot corresponds to one protein. A protein was considered

differentially expressed if its associated P-adj. value was inferior to 0.05. Red and

blue dots represent proteins enriched (Log2FC > 0.75) or depleted (Log2FC < -0.75)

in mutant brains, respectively. The names of the top 25 most differentially expressed

proteins are boxed.

1495

Figure 6: Anxa2 and GFAP upregulation and increased number of LCP1-

positive cells around CNS ventricles of *rpgrip1l^{/-}* juvenile fish at scoliosis

onset.

(A-D') Immunostaining for GFAP (A, B, C, D) and Anxa2 (A', B', C', D') on adult brain

transverse sections at DiV (A-B') and RhV (cerebellar) (C-D') levels in controls (A, A',

C, C') (n=3) and *rpgrip1l^{/-}* scoliotic (B, B', D, D') (n=4) fish. In A-D, nuclei are stained

with DAPI (blue). White stars indicate the SCO, white triangles point to the ventral RhV region. Open arrows point to a long cellular process co-labelled by Anxa2 and GFAP. **(E-L)** Immunostaining for GFAP (magenta) on transverse sections of the brain at SCO (E-H) and RhV (I-L) levels in controls (E, I) (n=4), straight *rpgrip1l$^{-/-}$* (F, G, J, K) (n=4) and scoliotic *rpgrip1l$^{-/-}$* (H, L) (n=3) juvenile (8 wpf) fish. Nuclei are stained with DAPI (white). Scale bar: 125µM for A-D', 20 µm for E-H and 100 µm for I-L. **(M)** Graph representing the percentage of GFAP-positive cells among the total number of SCO cells in control (n=4), straight *rpgrip1l$^{-/-}$* (n=4), and scoliotic *rpgrip1l$^{-/-}$* (n=3) fish at scoliosis onset (N=1). Each dot represents the mean ratio of GFAP-positive over negative cells in 3-6 sections per fish. Green dots correspond to fish with no or very low percentage of GFAP+ cells, red dots to fish with a high percentage of GFAP+ cells. Statistical analysis was performed with Tukey's multiple comparisons test where ns means non-significant, ** means P-value < 0.01 and *** P-value < 0.001. Error bars represent s.d. "str": straight; "sco": scoliotic. **(N-Q')** Immunostaining for LCP1+ microglia/macrophages (magenta) on brain transverse sections at SCO levels in a rectangle of 100 µm x 60 µm located under the posterior commissure in controls (n=4) (N, N'), straight *rpgrip1l$^{-/-}$* (n=4) (O-P') and *rpgrip1l$^{-/-}$* at scoliosis onset (n=3) (Q, Q') 8 wpf fish (N=1). Nuclei are stained with DAPI (white). Scale bar: 20 µm. **(R)** Quantification of LCP1+ cell density around the SCO in controls (n=4), straight *rpgrip1l$^{-/-}$* (n=4) and scoliotic *rpgrip1l$^{-/-}$* (n=3) at 8 wpf. 23 sections were counted for control, 21 sections for *rpgrip1l$^{-/-}$* straight and 18 sections for *rpgrip1l$^{-/-}$* scoliotic fish. Each dot represents the mean value for one fish. Green dots correspond to the fish without any GFAP+ SCO cells, red dots correspond to the fish with GFAP+ SCO cells (SCO cells are defined as the dorsal cells located underneath the posterior commissure and above the diencephalic ventricle). Note that straight *rpgrip1l$^{-/-}$* fish with GFAP+ SCO cells present a higher number of LCP1+ cells. Statistical analysis was performed with Tukey's multiple comparisons test where ns means non-significant. Error bars represent s.e.m. "str": straight "sco": scoliotic.

Figure 7: NACET treatment reduces scoliosis penetrance and severity in *rpgrip1l$^{-/-}$* and decreases astrogliosis and LCP1+ cell number at SCO level.

(A) Kinetics of scoliosis development in *rpgrip1l$^{+/-}$* incrosses in the presence (n=150 fish) or absence (n=174 fish) of NACET treatment (1.5 mM) from 4 to 12 weeks

(N=1). "str": straight; "sco": scoliotic. **(B)** Quantification of body axis curvature from

dorsal and lateral views at 13 wpf, at the end of the NACET treatment. Each dot

corresponds to one fish. Among the 174 fish raised in untreated condition, we

measured 18 controls (*rpgrip1l*$^{+/-}$) and 33 *rpgrip1l*$^{-/-}$, and upon NACET treatment, 16

controls (*rpgrip1l*$^{+/+}$) and 23 *rpgrip1l*$^{-/-}$. *rpgrip1l*$^{+/-}$ represents a pool of *rpgrip1l*$^{+/+}$ and

rpgrip1l$^{+/-}$. Error bars represent s.d. **(C)** Representative pictures of *rpgrip1l*$^{-/-}$ fish with

their curvature index without or after NACET treatment, at 13 wpf. Scale bar is 0,5

cm. Left panels correspond to an untreated *rpgrip1l*$^{-/-}$ scoliotic fish on lateral (upper)

and dorsal (lower) views. Middle panels represent treated *rpgrip1l*$^{-/-,}$ with a lower

curvature index than scoliotic fish on lateral (upper) and dorsal (lower) views and

right panels are pictures of rescued straight *rpgrip1l*$^{-/-}$ fish on lateral (upper) and

dorsal (lower) views. **(D)** Graph representing the number of LCP1+ cells in wt (n=3),

untreated *rpgrip1l*$^{-/-}$ (n=3) and NACET-treated *rgprip1l*$^{-/-}$ (n=3) fish in the SCO region

(N=1). **(E)** Immunostaining for GFAP (green), Acetylated tubulin (white) and LCP1

(red) on adult brain transverse sections at SCO level in untreated (upper panels) and

NACET-treated (lower panels) fish. Nuclei are stained with DAPI (blue). No GFAP

staining was present in controls (n=3) (upper left), while untreated mutants displayed

strong GFAP labeling (upper right) (n=3). NACET treatment totally suppressed gliosis

in a straight *rpgrip1l*$^{-/-}$ fish (lower right panel) (n=1) and diminished the intensity of

GFAP labeling in slightly tail-up *rpgrip1l*$^{-/-}$ (n=2). Scale bar: 12,5 µm. **(F)** Graph

presenting the percentage of GFAP-positive cells among the total number of SCO

secretory cells compared between controls (n=3), untreated *rpgrip1l*$^{-/-}$ (n=3), and

NACET treated *rpgrip1l*$^{-/-}$ adult fish (n=3) (N=1). Each dot represents the mean of %

of GFAP-positive cells present in 3-6 sections per fish. Statistical analysis was

performed using Tukey's multiple comparisons test where * means P-value <0.05

and **** means P-value < 0.0001.

1561

Figure 8: Altered ciliogenesis, GFAP upregulation and increased number of

LCP1-positive cells around ventricles of *cep290*$^{-/-}$ juvenile fish at 4 wpf.

(A-F') Immunostaining of the cilia marker glutamylated tubulin (white) on 4 wpf brain

transverse sections at 3 antero-posterior SCO levels in 4 wpf control (A-C', n=3) and

cep290$^{-/-}$ scoliotic (D-F', n=3) fish. Nuclei are stained with DAPI (blue). White stars

indicate the diencephalic ventricle (DiV), above which lies the SCO. White rectangles

in C and F delineate the close-up on cilia (C', F'). Scale bar: 10 µm for A-F, 2,5 µm

for C'-F'. **(G-L)** Immunostaining for Acetylated Tubulin (G, J), GFAP (H, K) and LCP1

(I, L) on transverse sections of the brain at SCO level in controls (G-I) (n=3), and

cep290$^{-/-}$ (J-L) (n=3) juvenile (4 wpf) fish. In G and J, nuclei are stained with DAPI

(blue). Scale bar: identical to A-F. **(M)** Graph representing the percentage of GFAP-

positive cells among the total number of SCO secretory cells in control (n=3) and

scoliotic *cep290*$^{-/-}$ (n=3) fish at scoliosis onset (N=1). Each dot represents the mean

ratio of GFAP-positive over total SCO secretory cells in 5-7 sections per fish. **(N)**

Quantification of LCP1+ cell density around the SCO in controls (n=3) and curved

cep290$^{-/-}$ (n=3) fish at 4 wpf. 16 sections were counted for control, 18 sections for

cep290$^{-/-}$ scoliotic fish. Each dot represents the mean value for one fish. Statistical

analysis was performed with unpaired T test where * means P-value < 0.05. Error

bars represent s.d. **(O-V)** Immunostaining for GFAP (white) on brain transverse

sections at cerebellum levels (P, R) and anterior hindbrain level (T, V) in control (P,

T) (n=3), and *cep290*$^{-/-}$ (R, V) (n=3) fish. Corresponding nuclear staining (DAPI in

white) is shown above (O, Q, S, U). Scale bar: 125 µm.

1584

 SUPPLEMENTARY FIGURE LEGENDS

1586

Supplementary figure S1: Production and characterization of the *rpgrip1l$^{\Delta}$* and

***rpgrip1l^{ex4}* fish lines.**

(A) Schematic exon-intron structure of the *Danio rerio rpgrip1l* gene. Two alleles

were produced by CRISPR/cas9-mediated genome engineering,

rpgrip1l^{ex4}(rpgrip1l^{bps2} in ZFIN) with a nonsense mutation in exon 4 and *rpgrip1l$^{\Delta4-25}$*

(*(rpgrip1l^{bps1} in ZFIN)* also called *rpgrip1l$^{\Delta}$* or *rpgrip1l*) with a large deletion between

exons 4 and 25, encompassing most of the protein coding sequence. Both alleles

had identical embryonic phenotype (panel B and data not shown). (A')

Electrophoresis gel showing the migration profile of amplicons for each genotype,

rpgrip1l$^{\Delta/\Delta}$, *rpgrip1l$^{\Delta/+}$*, *rpgrip1l$^{+/+}$* produced by triplex PCR (B) Absence of laterality

defects and kidney cysts in *rpgrip1l$^{\Delta/\Delta}$* and *rpgrip1l$^{ex4/ex4}$* embryos. **(C)**

Immunostaining on whole 30 hpf control (left panels) and *rpgrip1l$^{/-}$* (right panels)

embryos to visualize the RF (anti-RF antibody, middle panels) and cilia (acetylated

tubulin antibody, lower panels) in the caudal region. Nuclei (upper panels) are

stained with DAPI. The RF is present in the central canal of *rpgrip1l$^{/-}$* embryos, as

well as cilia in the whole neural tube width. Scale bar: 6 µm for upper panels; 15 µm

for lower panels. **(D-E)** Absence of retinal morphological defects in 5 dpf *rpgrip1l$^{/-}$*

larvae. Retinal sections of control (D) and *rpgrip1l$^{/-}$* (E) larvae were immunostained

with Zpr3 (red) to label the external segment of photoreceptors and Acetylated

Tubulin (Ac Tub) (green) to label axonemes. D' and E' are insets in D and E focusing

on the photoreceptor layer. Scale bar: 50 µm. (**F-H'''**) Scanning electron microscopy

of brain ventricles in control (F-F''), straight (str.) *rpgrip1l$^{/-}$* (G-G'') and scoliotic (sco.)

rpgrip1l$^{/-}$ (H-H''') adult (3 months) fish. Ependymal multiciliated cells of the

rhombencephalic ventricle (RhV) are almost totally absent in *rpgrip1l$^{/-}$* scoliotic fish

(I-I'), while they are present in straight r*pgrip1l$^{/-}$*(H-H') and controls (G-G''). Cilia of

hindbrain mono-ciliated cells in *rpgrip1l$^{/-}$* (F'', G'') and controls (G''). Scale bars: 10

µm for F-I, 2 µm for F'-H', 1 µm for F'', G'', H'''' and H'''.

1614

1615

1616

Supplementary figure S2: Presence of scospondin aggregates and heterogenous ciliary defects in the fChP of *rpgrip1l$^{-/-}$* juvenile fish.

(A-D) Immunostaining on transverse sections of the SCO, at anterior level (A, A', B, B') or posterior level (C, D) in *rpgrip1l$^{+/+}$* (n=3) and tail-up *rpgrip1l$^{-/-}$* (n=3), transgenic for Sspo-GFP (N=2). Sspo-GFP forms small cytoplasmic aggregates within SCO cells of both genotypes (* in A' and B'), under the primary cilia extending towards the DiV, and accumulates at the apical surface of tail-up *rpgrip1l$^{-/-}$*, creating a double positive area for cilia and Sspo-GFP staining (yellow in B'). Multiciliated cells lateral to the posterior SCO (mcc) are negative for Sspo-GFP staining in controls (C) and absent in tail-up *rpgrip1l$^{-/-}$* (D). **(E-J)** GFP immunostaining on transverse section of the fChP (E-G) or at the level of the optic tectum (H-J) in *rpgrip1l$^{+/+}$* (E, H, n=7), straight *rpgrip1l$^{-/-}$* (F, I, n=4) and scoliotic *rpgrip1l$^{-/-}$* (G, J, n=5) 8 wpf fish transgenic for *sspo-GFP* (N=2). Scoliotic *rpgrip1l$^{-/-}$* fish present with filamentous material and aggregates within the ventricles (G, J) while a small RF fragment is present within the ventricle of controls (H) and straight mutants (I) at optic tectum level. * indicates aggregates. **(K-O'')** Representative maximum intensity Z-stack projections of confocal transverse brain section at the level of the fChP in *rpgrip1l$^{+/+}$* (K-K'') (n=7), straight *rpgrip1l$^{-/-}$* (L-M'') (n=4) and scoliotic *rpgrip1l$^{-/-}$* (N-O'') (n=5) 8 wpf fish (N=2). Cilia are stained with glutamylated-tubulin and acetylated-tubulin antibodies (both in white) and nuclei are stained with DAPI (blue). Sections shown are at the level of the habenular nuclei. At this level, monocilia in the dorsal midline region (K') and multicilia on lateral sides (K'') (n=7/7) are present in controls. Straight *rpgrip1l$^{-/-}$* exhibited either normal (L', L'') (n=2/4) or abnormal and sparser (M', M'') (n=2/4) monocilia and multicilia. Similarly, scoliotic *rpgrip1l$^{-/-}$* presented either normal monocilia and multicilia (N', N'') (n=3/5) or long monocilia (O') and decreased cilia density of multi-ciliated tufts (O'') (n=2/5). Scale bars: 20 μm.

Supplementary figure S3: Genetic invalidation of *urp2* in *rpgrip1l$^{-/-}$* fish does not have any beneficial effect on scoliosis penetrance or severity.

(A-E) Absence of compensatory upregulation of *urp* family members in *urp2$^{-/-}$* fish. Expression levels of *urp2* (A), *urp1* (B), *urp* (C), *uts2a* (D) and *uts2b* (E) in zebrafish in 13 wpf adults. Each dot represents the value measured for one fish; green and purple dots represent straight and curved fish, respectively. n = 5 [*rpgrip1l$^{+/+}$; urp2$^{+/+}$*],

9 [*rpgrip1l$^{+/+}$; urp2$^{-/}$*], 11 [*rpgrip1l$^{-/-}$; urp2$^{+/}$*] and 10 [*rpgrip1l$^{-/-}$; urp2$^{-/}$*]. Expression levels for each gene are compared to the *lsm12b* housekeeping gene. Statistical analysis was performed using the Tukey test where ns means not significant, ** P < 0.01, **** P < 0.0001. Error bars represent s.d. None of the *uts2* family members is significantly upregulated in [*rpgrip1l$^{-/-}$*; urp2$^{-/-}$] fish compared to *rpgrip1l$^{-/-}$*, suggesting an absence of compensation. **(F-G)** Quantification of body axis curvature on dorsal and lateral positions at 2 (F) and 4 (G) mpf. Each dot represents the curvature index of one fish. Statistical analysis was done using Tukey's multiple comparisons test where ****P < 0.0001, with error bars representing s.d. At 2 mpf, n=30 [*rpgrip1l$^{+/+}$; urp2$^{+/+}$*], 26 [*rpgrip1l$^{+/+}$; urp2$^{-/}$*], 27 [*rpgrip1l$^{-/-}$; urp2$^{+/+}$*], 55 [*rpgrip1l$^{-/-}$; urp2$^{+/}$*] and 32 [*rpgrip1l$^{-/-}$; urp2$^{-/}$*]. At 4 mpf, n=9 [*rpgrip1l$^{+/+}$; urp2$^{+/+}$*], 7 [*rpgrip1l$^{+/+}$; urp2$^{-/}$*], 27 [*rpgrip1l$^{-/-}$; urp2$^{+/+}$*], 48 [*rpgrip1l$^{-/-}$; urp2$^{+/}$*] and 27 [*rpgrip1l$^{-/-}$; urp2$^{-/}$*]. **(H)** Rose plots representing combined Cobb angle measurements at 4 mpf of [*rpgrip1l$^{+/+}$; urp2$^{+/+}$*] (n=7) in green, [*rpgrip1l$^{+/+}$; urp2$^{-/}$*] (n=6) in blue, [*rpgrip1l$^{-/-}$; urp2$^{+/+}$*] (n=10) in pink and [*rpgrip1l$^{-/-}$; urp2$^{-/}$*] (n=7) in yellow. Cobb angle measurements were performed on Alizarin red fish skeletons. One example of Alizarin red skeletons for each genotype is presented below its corresponding rose-plot. **(I)** Illustration of combined curvature index measurement from traced lines on lateral (left) and dorsal (right) views of fish. The curvature index of both traced lines was measured using a modified MATLAB program (LineCurvature2D function) which provided a measurement in radian/length in pixels. Scale bar: 5 mm.

Supplementary Figure S4: Complementary comparative transcriptome and proteomic analysis of *rpgrip1l$^{-/-}$* versus control fish.

(A, B) GO term analysis of biological processes enriched in *rpgrip1l$^{-/-}$* for genes upregulated in the trunk transcriptome data (A) or enriched in *rpgrip1l$^{-/-}$* for proteins in the brain adult proteome data (B) using Metascape. **(C)** List of the top 25 most enriched proteins in the *rpgrip1l$^{-/-}$* vs control brain proteome. **(D)** List of selected genes upregulated in the zebrafish *rpgrip1l* trunk transcriptome and found in different reactive astrogliosis models in single cell transcriptome studies (Matusova et al, 2023).

Supplementary figure S5: Increased GFAP and Annexin 2 staining and LCP1+ cell density in different brain regions.

(A-F') Immunostaining of Annexin A2 (Anxa2) (A-F) and GFAP (A'- F') on transverse brain sections at habenula (A, A', D, D'), tectum (B, B', E, E') and hindbrain (C, C', F, F') levels in controls (A-C') and scoliotic *rpgrip1l⁻/⁻* (D-F') fish. Nuclei are stained with DAPI (blue). Scoliotic *rpgrip1l⁻/⁻* fish upregulate Annexin A2 and GFAP in overlapping territories. Scale bar: 20 µm. **(G)** Quantification of LCP1+ cell density in the optic tectum region of *rpgrip1l⁺/⁺* (n=4), straight *rpgrip1l⁻/⁻* (n=4) and scoliotic *rpgrip1l⁻/⁻* (n=3) 8 wpf fish. Each dot represents a fish. 15-20 sections were counted for each brain level. Straight *rpgrip1l⁻/⁻* fish sections displayed around twice more Lcp1+ cells at tectum level than controls, while scoliotic *rpgrip1l⁻/⁻* fish had a slightly increased LCP1+ cell density compared to controls. Statistical analysis was performed using Tukey's multiple comparisons test where ns means non significative and * Pvalue < 0.05, ** Pvalue < 0.01 with error bars representing s.d. **(H-J')** Immunostaining of LCP1+ immune cells on transverse sections of the optic tectum region of *rpgrip1l⁺/⁺* (H, H') (n=4), straight *rpgrip1l⁻/⁻* (I, I') (n=4) and scoliotic *rpgrip1l⁻/⁻* (J, J') (n=3) 8 wpf fish. Immune cells are labelled with the LCP1 antibody (magenta) and nuclei with DAPI (blue). H', I' and J' are higher magnifications of the squared regions in H, I and J, respectively. Scale bars: 100 µm for H, I, J; 30 µm for H', I', J'. str: straight, sco: scoliotic. **(K)** Quantification of LCP1+ cell density in the cerebellum region of 8 wpf fish (N=1). The analysis compared straight *rpgrip1l⁻/⁻* (n=4) and scoliotic *rpgrip1l⁻/⁻* (n=3) fish to *rpgrip1l⁺/⁺* siblings (n=4). No significant difference was found between genotypes. Statistical analysis was performed using Tukey's multiple comparisons test where ns means not significant. Error bars represent s.d. "str" means straight and "sco" means scoliotic.

Supplementary figure S6: Biological assay of NACET activity on *dnaaf1⁻/⁻* embryos and expression level of three immune markers at adult stage. (A) Theta angle (θ) measurement between the eye, the caudal end of the yolk extension and the tail extremity, corresponding to the embryo curvature index. **(B)** Lateral views of 60 hpf representative embryos showing axial curvature phenotypes. In untreated conditions, *dnaaf1⁻/⁻* embryos show a characteristic curly tail-down (CTD) phenotype, whereas *dnaaf1⁺/⁺* siblings have a straight axis. Treatment at 27 hpf with

3 mM NACET suppresses the CTD phenotype of *dnaaf1*[-/-] embryos and does not

affect axis straightness of *dnaaf1*[+/+] siblings. Embryos were measured around 2.5 to

3 mm. **(C-F)** Rose plots representing the $(360 - \theta)$ angle of *dnaaf1*[+/+] and *dnaaf1*[-/-] in

untreated and 3 mM NACET-treated conditions. Embryos were treated from 27hpf to

2.5 dpf and θ angles were measured at the end of the treatment. Untreated *dnaaf1*[+/+]

are represented in (C – dark green), and treated ones are in (D – light green).

Untreated *dnaaf1*[-/-] are in (E – blue) and treated ones are in (F – red). NACET

treatment is able to rescue curly tail down phenotype in *dnaaf1*[-/-] embryos without

affecting the controls.

1724

1725   **Supplementary Movie 1: Micro-CT scans of *rpgrip1l*[Δ/Δ] juvenile vertebral axis**

1726   **compared to controls**

1727   Superposition of 3D-reconstructed spines of 5 wpf juvenile fish, 1 control (white), 3

1728   rpgrip1lΔ/Δ of different severities: 2 tail-up (yellow, blue) and 1 scoliotic (pink)

1729

1730

A

rpgrip1l$^{-/-}$

5 dpf

5 wpf

12 wpf

B

Scoliosis appareance in clutches from
rpgrip1l$^{-/+}$ incrosses

C

Scoliosis penetrance in adults

D Ctrl **E** *rpgrip1l$^{-/-}$*

F Dorsal view

G Lateral view

ctrl *rpgrip1l$^{-/-}$* *rpgrip1l$^{-/-}$*

H

| Tol2 | Foxj1a enh. | 5XMyc-RPGRIP1L | polyA. | polyA | GFP. | cmlc2 enh. | Tol2 |
| Tol2 | Col2a1 enh. | 5XMyc-RPGRIP1L | polyA. | polyA | GFP. | cmlc2 enh. | Tol2 |

I

Genotype	*rpgrip1l$^{-/-}$*	*rpgrip1l$^{-/-}$; Tg Foxj1a :RPGRIP1L*	*rpgrip1l$^{-/-}$*	*rpgrip1l$^{-/-}$; Tg Col2a1 :RPGRIP1L*
Adult Scoliotic fish (n)	72/74	0/63	28/29	28/29
Clutch (N)	N=2	N=2	N=1	N=1

J

% of scoliotic *rpgrip1l$^{-/-}$*

A

Stage: Juvenile — WT, *rpgrip1l⁻ᐟ⁻* | Adult — WT, *rpgrip1l⁻ᐟ⁻*

Fold change (*Urp2* expression)

Straight (green)
Scoliotic (magenta)

C

rpgrip1l⁻ᐟ⁻ ;urp2⁺ᐟ⁺

rpgrip1l⁻ᐟ⁻ ;urp2⁻ᐟ⁻

B

Curvature index

Radian/length in pixels

ns
****

rpgrip1l⁺ᐟ⁻;urp2⁺ᐟ⁺
rpgrip1l⁺ᐟ⁺;urp2⁻ᐟ⁻
rpgrip1l⁻ᐟ⁻ ; urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻ ; urp2⁺ᐟ⁻
rpgrip1l⁻ᐟ⁻ ; urp2⁻ᐟ⁻

D

urp2

2^{-dct}

***
**
****

rpgrip1l⁺ᐟ⁺;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁻
rpgrip1l⁻ᐟ⁻;urp2⁻ᐟ⁻

E

urp1

2^{-dct}

Straight (green)
Scoliotic (magenta)

**
ns

rpgrip1l⁺ᐟ⁺;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁻
rpgrip1l⁻ᐟ⁻;urp2⁻ᐟ⁻

A **Brain**

B **Dorsal trunk**

C

GO:0035082: axoneme assembly
GO:0001539: cilium or flagellum-dependent cell motility
GO:0002460: adaptive immune response
GO:0003351: epithelial cilium movement involved in extracellular fluid movement
GO:0007276: gamete generation
R-HSA-6785807: Interleukin-4 and Interleukin-13 signaling
GO:0045921: positive regulation of exocytosis

D **Up-regulated genes**

Trunk   Brain

560   54   6

E

Process/Pathway	Gene	Trunk.LogFC	Trunk P adj.	Brain.LogFC	Brain P adj.
Foxj1a and	*tbata*	2,25	8,82E-22	0,84	1,80E-07
Its targets	*foxj1a*	1,90	3,77E-23	1,36	9,67E-21
	enkur	1,88	8,98E-24	0,76	2,13E-06
	stpg1	1,80	1,10E-09	0,69	3,00E-05
	ccdc65	1,73	8,65E-11	0,74	1,25E-05
Regulators of	*pkd1b*	3,20	2,21E-47	1,90	1,08E-45
axis curvature	*urp1*	1,92	1,32E-16	-	-
	urp2	0,96	1,22E-03	0,87	5,44E-09
	sspo	1,18	4,16E-05	-	-
Immune	*irg1l*	2,45	1,20E-16	-	-
Response	*il11a*	2,23	3,09E-16	0,67	1,21E-05
	c4	1,97	1,50E-12	1,54	5,63E-29
	anxa2a	1,52	8,15E-17	1,53	7,04E-26
	mmp9	1,00	3,94E-04	0,46	4,02E-02
	c7a	1,86	8,82E-22	0,80	9,38E-07
	stat3	0,76	3,63E-07	0,80	5,23E-10

F **Brain proteome**

A **Scoliosis penetrance in *rpgrip1l*⁺/⁻ incross**

B **Curvature index**

C **Scoliotic phenotype and associated curvature index of *rpgrip1l*⁻/⁻ adults**

D **Lcp1 cells in SCO region**

E

F **% of GFAP(+) cells within the SCO**

DAPI Arl13b Glu-tub

Arl13b

Glu-tub

rpgrip1l$^{+/+}$

A A' A'' A'''

rpgrip1l$^{-/-}$ str.

B B' B'' B'''

rpgrip1l$^{-/-}$ sco.

C C' C'' C'''

D

Cilia length mean along central canal (μm)

P value 0,09

** *

rpgrip1l +/+ -/- str. -/- sco.

E

Cilia density mean along central canal

**** *** ns

rpgrip1l +/+ -/- str. -/- sco.

F

Central canal width (μm)

P Value 0,07

* ns

rpgrip1l +/+ -/- str. -/- sco.

rpgrip1l$^{+/+}$ *rpgrip1l*$^{-/-}$ straight *rpgrip1l*$^{-/-}$ scoliotic

Anterior SCO

G I K

G' I' K'

Post SCO

H J L

Ac.tub/Glu.Tub DAPI

Adult rpgrip1l⁺/⁺

Scoliotic adult rpgrip1l⁻/⁻

DiV

A DAPI GFAP
A' ANXA2
B DAPI GFAP
B' ANXA2

RhV

C
C'
D
D'

rpgrip1l⁺/⁺

straight *rpgrip1l⁻/⁻*

rpgrip1l⁻/⁻ scoliosis onset

DiV

E
F
G
H

RhV (cerebellar lev.)

I
J
K
L DAPI GFAP

M

% of GFAP(+) cells among SCO cells

ns   ***
      **

rpgrip1l⁺/⁺   *rpgrip1l⁻/⁻* str.   *rpgrip1l⁻/⁻*sco.

rpgrip1l⁺/⁺

straight *rpgrip1l⁻/⁻*

rpgrip1l⁻/⁻ scoliosis onset

N
O
P
Q' DAPI LCP1

N'
O'
P'
Q'

R

LCP1(+) cells density

P value 0,07

ns   ns

WT   *rpgrip1l⁻/⁻* str.   *rpgrip1l⁻/⁻* sco.

A

Exon 4   Exon 25

Deletion

Start Codon   CRISPR site   CRISPR site

A'

544 bp : ex4F/ex25R for MUT*
364 bp : ex25F/ex25R for WT

B

	Heart looping	Kidney cysts	Total nb embryos
wt	1	0	900
rpgrip1l$^{ex4/ex4}$	0	0	73
rpgrip1l$^{\Delta/\Delta}$	3	0	694

C

Control   *rpgrip1l*-/-

DAPI

RF

Ac-tub

Zpr3 Ac-tub DAPI

Control   *rpgrip1l*-/-

D   **E**

D'   **E'**

rpgrip1l+/+   **F**   **F'**   **F''**

           I'

rpgrip1l-/- str.   **G**   **G'**   **G''**

rpgrip1l-/- sco.   **H**   **H'**   **H''**   **H'''**

Supplementary Figure S1

Supplementary Figure S2

Supplementary Fig S3

A: Up-regulated genes in trunk

GO:0003341: cilium movement
GO:0070286: axonemal dynein complex assembly
GO:0003352: regulation of cilium movement
GO:0090660: cerebrospinal fluid circulation
GO:1904158: axonemal central apparatus assembly
GO:0002252: immune effector process
GO:0071345: cellular response to cytokine stimulus
GO:0007420: brain development
hsa04814: Motor proteins
GO:0002521: leukocyte differentiation
GO:0070663: regulation of leukocyte proliferation
WP2855: Dopaminergic neurogenesis
GO:0032735: positive regulation of interleukin-12 production
GO:0120229: protein localization to motile cilium
GO:0150063: visual system development
GO:0040017: positive regulation of locomotion
GO:0035239: tube morphogenesis
GO:0048598: embryonic morphogenesis
GO:0044703: multi-organism reproductive process
GO:0060541: respiratory system development

B: Enriched proteins in adult brain

GO:0051336: regulation of hydrolase activity
GO:0010638: positive regulation of organelle organization
GO:0006897: endocytosis
GO:1905686: positive regulation of plasma membrane repair
GO:0006886: intracellular protein transport
GO:0030162: regulation of proteolysis
R-HSA-2454202: Fc epsilon receptor (FCERI) signaling
WP5323: Fatty Acids and Lipoproteins Transport in Hepatocytes
GO:0060627: regulation of vesicle-mediated transport
GO:2000644: regulation of receptor catabolic process
GO:0006954: inflammatory response
GO:0071363: cellular response to growth factor stimulus
GO:0016310: phosphorylation
GO:0031329: regulation of cellular catabolic process
GO:0006887: exocytosis
GO:0032944: regulation of mononuclear cell proliferation
GO:0030036: actin cytoskeleton organization
GO:0150105: protein localization to cell-cell junction
GO:0032386: regulation of intracellular transport
GO:0050778: positive regulation of immune response

C: 25 highest P Value

Protein	FC	P value
anxa2a	7,28	4,82E-08
echdc2	4,97	1,99E-06
anxa1a	6,06	9,6856E-06
anxa5b	2,37	3,3303E-05
capsla	3,72	0,00011272
tagln2	2,53	0,00011315
psph	1,72	0,00011655
c7a	3,07	0,0002431
mapk9	2,49	0,00029106
vav3b	2,31	0,00036912
serpina7	2,46	0,0003851
gps1	1,92	0,00057247
synrg	2,08	0,00066171
c3 like	3,77	0,00068862
tut7	2,01	0,00102053
gsk3bb	2,40	0,00109597
tmed3	3,44	0,00148983
mrc1a	2,10	0,00152123
wdr12	2,78	0,00186884
wdfy2	1,97	0,00235225
plcd4b	2,24	0,00249521
itchb	1,81	0,00250875
ehd1b	1,94	0,00253596
exoc6	1,77	0,00298918
stat3	5,31	0,00319733

D: Activated astrocytes markers

Hum. Gene	Zf Gene	FC (trunk)	P.adj. Value
S100B	s100b	2,98	7,35711E-27
C4b	c4b	5,15	1,48224E-21
GFAP	gfap	2,49	1,99844E-09
CTSB	ctslb	3,60	1,29502E-08
CAPS	capsla	2,67	2,31406E-06
GLAST/SLC1A3	slc1a3b	2,68	1,46788E-05
PLCXD3	plcxd3	1,74	1,6557E-05
VIM	viml	2,14	8,30568E-05

Supplementary Figure S4

Supplementary Figure S5

Supplementary Figure S6

Subcommissural Organ (SCO)

SCO

RF

Defective CSF flow
Motility/Coordination defects

Posterior
Commissure

DiV

mcc

Defective Cilia signalling
Astrogliosis/
immune cells attraction

• Sco-spondin

GFAP+

Immune cells

SCO