ASTROGLIOSIS AND NEUROINFLAMMATION UNDERLIE SCOLIOSIS UPON CILIA DYSFUNCTION

Morgane Djebar[1], Isabelle Anselme[1], Guillaume Pezeron[2], Pierre-Luc Bardet[1], Yasmine Cantaut-Belarif[3], Alexis Eschstruth[1], Diego López Santos[1], Hélène Le Ribeuz[1], Arnim Jenett[4], Hanane Khoury[1], Joëlle Véziers[5], Caroline Parmentier[6], Aurélie Hirschler[7], Christine Carapito[7], Ruxandra Bachmann-Gagescu[8], Sylvie Schneider-Maunoury[1]* and Christine Vesque[1]*

Sorbonne Université, CNRS UMR7622, INSERM U1156, Institut de Biologie Paris Seine (IBPS) - Developmental Biology Unit, 75005, Paris, France
Molecular Physiology and Adaptation (PhyMA - UMR7221), Muséum National d'Histoire Naturelle, CNRS, Paris, France
Institut du Cerveau et de la Moelle épinière (ICM), Sorbonne Université, Inserm U 1127, CNRS UMR 7225, F-75013, Paris, France
TEFOR Paris-Saclay, CNRS UMS2010 / INRA UMS1451, Université Paris-Saclay.
Inserm UMR 1229, CHU Nantes PHU4 OTONN, SC3M facility, Inserm UMS 016, CNRS 3556, Université de Nantes, Nantes, France.
Sorbonne Université, CNRS UMR8246, INSERM U1130, Institut de Biologie Paris Seine (IBPS) – Neurosciences Paris Seine (NPS), 75005, Paris, France
Laboratoire de Spectrométrie de Masse Bio-Organique, IPHC, UMR 7178, Université de Strasbourg, CNRS, Infrastructure Nationale de Protéomique ProFI - FR2048, Strasbourg, France.
Institute of Medical Genetics, University of Zurich, Wagistrasse 12, 8952 Schlieren, Zurich, Switzerland; Institute of Molecular Life Sciences, University of Zurich, Winterthurerstrassse 190, 8057 Zurich, Switzerland
* Co-corresponding authors, co-last authors.
sylvie.schneider-maunoury@sorbonne-universite.fr
christine.vesque@sorbonne-universite.fr

ABSTRACT

Cilia defects lead to scoliosis in zebrafish, but the underlying pathogenic mechanisms are poorly understood and may diverge depending on the mutated gene. Here, we dissected the mechanisms of scoliosis onset in a zebrafish mutant for the *rpgrip1l* gene encoding a ciliary transition zone protein. *rpgrip1l* mutant fish developed scoliosis with near-total penetrance but asynchronous onset in juveniles. Taking advantage of this asynchrony, we found that curvature onset was preceded by ventricle dilations and was concomitant to the perturbation of Reissner fiber polymerization and to the loss of multiciliated tufts around the subcommissural organ. Rescue experiments showed that Rpgrip1l was exclusively required in *foxj1a*-expressing cells to prevent axis curvature. Genetic interactions investigations ruled out Urp1/2 levels as a main driver of scoliosis in *rpgrip1* mutants. Transcriptomic and proteomic studies identified neuroinflammation associated with increased Annexin levels as a potential mechanism of scoliosis development in *rpgrip1l* juveniles. Investigating the cell types associated with *annexin2* over-expression, we uncovered astrogliosis, arising in glial cells surrounding the diencephalic and rhombencephalic ventricles just before scoliosis onset and increasing with time in severity. Anti-inflammatory drug treatment reduced scoliosis penetrance and severity and this correlated with reduced astrogliosis and macrophage/microglia enrichment around the diencephalic ventricle. Mutation of the *cep290* gene encoding another transition zone protein also associated astrogliosis with scoliosis. Thus, we propose astrogliosis induced by perturbed ventricular homeostasis and associated with immune cell activation as a novel pathogenic mechanism of zebrafish scoliosis caused by cilia dysfunction.

INTRODUCTION

Idiopathic scoliosis (IS) is a 3D rotation of the spine without vertebral anomalies that affects about 3% of adolescents worldwide. Its etiology remained mysterious, mainly because of the lack of appropriate animal models and the complexity of its inheritance profile in large families presenting a severe scoliosis trait (Patten et al., 2015). Since the description of the first IS model in the zebrafish *ptk7* gene mutant (Hayes et al., 2014), a number of mutants for genes encoding ciliary proteins were shown to develop scoliosis at late larval and juvenile stage (4 to 12 weeks post-fertilization, wpf) without any vertebral fusion or fracture, highlighting the link between cilia function and straight axis maintenance in that species (Grimes et al., 2016; Wang et al., 2022).

Cilia are microtubular organelles with sensory and/or motile functions. Zebrafish mutants in genes involved in cilia motility or in intraflagellar transport display severe embryonic axis curvature and are usually not viable beyond larval stages (Grimes et al., 2016; Rose et al., 2020). In contrast, mutants in genes implicated in ciliary gating or ciliary trafficking encoding respectively components of the transition zone (TZ) or of the BBsome complex survive to adult stage and develop scoliosis with variable penetrance (Bentley-Ford et al., 2022; Masek et al., 2022; Song et al., 2020; Wang et al., 2022). The molecular basis for these differences is still poorly understood, even if progress has been made in our understanding of zebrafish scoliosis. Cilia-driven movements of the cerebrospinal fluid (CSF) are involved in zebrafish axis straightness, both in embryos and juveniles (Boswell and Ciruna, 2017; Grimes et al., 2016) and are tightly linked to the assembly and maintenance of the Reissner fiber (RF), a SCO-spondin polymer secreted by the subcommissural organ (SCO) and running down the brain and spinal cord CSF-filled cavities (Cantaut-Belarif et al., 2018). RF loss at embryonic stage in null *sspo* mutants generates ventral curvatures of the posterior axis of the body and is lethal while its loss at juvenile stage in hypomorphic *sspo* mutants triggers scoliosis with full penetrance (Cantaut-Belarif et al., 2018; Rose et al., 2020; Troutwine et al., 2020). Signaling downstream of the RF in embryos implicates Urp1 and Urp2, two neuropeptides of the Urotensin 2 family expressed in ventral CSF-contacting neurons (CSF-cNs), which act on dorsal muscles in embryos and larvae (Cantaut-Belarif et al., 2020; Gaillard et al., 2023; Lu et al., 2020; Zhang et al., 2018). Their combined mutations or the mutation of their receptor gene *uts2r3* lead to scoliosis

at adult stages (Bearce et al., 2022; Gaillard et al., 2023; Zhang et al., 2018). However,

whether RF maintenance and Urp1/2 signaling are perturbed in juvenile scoliotic

mutants of genes encoding TZ proteins ("TZ mutants") and how these perturbations

are linked to scoliosis is unknown, especially because TZ mutants do not display any

sign of embryonic or larval cilia motility defects (Wang et al., 2022). Moreover, two

scoliotic models develop axial curvature while maintaining a RF showing that their

curvature onset occurs independently of RF loss (Meyer-Miner et al., 2022; Xie et al.,

2023). Finally, neuro-inflammation has been described in a small subset of zebrafish

IS models for which anti-inflammatory/anti-oxidant treatments (with NAC or NACET)

partially rescue scoliosis penetrance and severity (Rose et al., 2020; Van Gennip et

al., 2018).

In this paper we dissect the mechanisms of scoliosis appearance in a novel deletion

allele of the zebrafish TZ *rpgrip1l* gene. *RPGRIP1L* mutations have been found in

neurodevelopmental ciliopathies, Meckel syndrome (MKS) and Joubert syndrome (JS)

(Delous et al., 2007)(Arts et al., 2007) characterized by severe brain malformations.

Other organs such as the kidney and liver are variably affected. Some JS patients also

present with severe scoliosis, but this phenotypic trait is of variable penetrance (Delous

et al., 2007)(Brancati et al., 2008). Functional studies performed in mouse and

zebrafish have shown that *Rpgrip1l* is required for proper central nervous system

patterning due to complex perturbations of Sonic Hedgehog signaling (in mice) and

cilia planar positioning (in mice and zebrafish) (Andreu-Cervera et al., 2021; Mahuzier

et al., 2012; Vierkotten et al., 2007). Zebrafish *rpgrip1l* mutants are viable and develop

an axis curvature phenotype at juvenile stages with nearly full penetrance. Using a

variety of approaches, we show that i) Rpgrip1l is required in ciliated cells surrounding

ventricular cavities for maintaining a straight axis; ii) at juvenile stages, *rpgrip1l*

mutants develop cilia defects associated with ventricular dilations and loss of the RF;

iii) increased *urp1/2* expression in *rpgrip1l* mutants does not account for spine

curvature defects; and iv) a pervasive astrogliosis process associated with

neuroinflammation participates in scoliosis onset and progression. Finally, we show

that astrogliosis is conserved in another zebrafish ciliary TZ mutant, *cep290*, identifying

a novel mechanism involved in scoliosis upon ciliary dysfunction.

RESULTS

Rpgrip1l ^{-/-} fish develop scoliosis asynchronously at juvenile stage

To study the mechanisms of scoliosis appearance upon ciliary dysfunction in zebrafish, we produced a viable zebrafish deletion mutant in the *rpgrip1l* gene encoding a ciliary transition zone protein (*rpgrip1l*$^\Delta$, identified in ZFIN as *rpgrip1l*^{bps1}) Supplementary Figure S1A). We made a 26 kb deletion by CRISPR-Cas9 double-cut, which removes 22 out of the 26 exons of the protein (Supplementary Figure S1A) and triggers protein truncation at amino-acid (aa) position 88 out of 1256. 90-100% of *rpgrip1l*$^{\Delta/\Delta}$ embryos and larvae were straight (0-10%, depending on the clutch, displayed a sigmoid curvature). They did not display any additional defects found in many ciliary mutant embryos or larvae such as randomized left-right asymmetry, kidney cysts or retinal anomalies (Bachmann-Gagescu et al., 2011; Kramer-Zucker et al., 2005; Tsujikawa and Malicki, 2004)(Figure 1A and Supplementary Figure S1B, D, E). Another allele with a frameshift mutation in exon 4 (*rpgrip1l*^{ex4}, identified in ZFIN as *rpgrip1l*^{bps2}) showed a similar phenotype (Supplementary Figure S1A, B). Considering the large size of the deletion, we considered the *rpgrip1l*$^\Delta$ allele as a null and called it *rpgrip1l*$^{-/-}$ thereafter. *rpgrip1l*$^{-/-}$ animals developed scoliosis during juvenile stages. Scoliosis appearance was asynchronous between clutches, from 4 weeks post fertilization (wpf) (around 1 cm length) to 11 wpf (1.8- 2.3 cm), and also within a clutch. It started with slight upward bending of the tail (the *tail-up* phenotype) and progressed toward severe curvature (Figure 1A, B) with 90% penetrance in adults (100% in females and 80% in males) (Figure 1C). Micro-computed tomography (µCT) at two different stages (5 wpf and 23 wpf) confirmed that spine curvature was three-dimensional and showed no evidence of vertebral fusion, malformation or fracture (Figure 1D-G and Supplementary Movie 1).

Thus, the ciliary *rpgrip1l*$^{-/-}$ mutant shows almost totally penetrant, late-onset juvenile scoliosis without vertebral anomalies, making it a valuable model for studying the etiology of human idiopathic scoliosis. Furthermore, the asynchronous onset of curvature is an asset for studying the chronology of early defects leading to scoliosis.

Reintroducing RPGRIP1L *in foxj1a lineages suppresses scoliosis in rpgrip1l⁻ᐟ⁻ fish*

Since *rpgrip1l* is ubiquitously expressed in zebrafish (Mahuzier et al., 2012), scoliosis could have different tissue origins, including a neurological origin as shown for the *ptk7* and *ktnb1* scoliotic fish (Meyer-Miner et al., 2022; Van Gennip et al., 2018). Indeed, no tissue-specific rescue has been performed in mutants for zebrafish ciliary TZ or BBS genes. We tested if introducing by transgenesis a tagged human RPGRIP1L protein under the control of the *col2a1a (Dale and Topczewski, 2011)* or *foxj1a* (Van Gennip et al., 2018) enhancers (Figure 1H) would reduce scoliosis penetrance and severity. The *foxj1a* enhancer drives expression in motile ciliated cells, among which ependymal cells lining CNS cavities (Van Gennip et al., 2018), while the *col2a1a* regulatory region drives expression in cartilage cells including intervertebral discs and semi-circular canals, tissues that have been proposed to be defective in mammalian scoliosis models (Hitier et al., 2015; Karner et al., 2015). After selecting F1 transgenic fish that expressed the Myc-tagged RPGRIP1L protein in the targeted tissue at 2.5 dpf, we introduced one copy of each transgene into the *rpgrip1l⁻ᐟ⁻* background. The Foxj1a:5XMycRPGRIP1L transgene was sufficient to fully suppress scoliosis in *rpgrip1l⁻ᐟ⁻* fish: none of the transgenic *rpgrip1l⁻ᐟ⁻* fish became scoliotic while virtually all the non-transgenic *rpgrip1l⁻ᐟ⁻* siblings did (Figure 1I, J). In contrast, RPGRIP1L expression triggered by the *col2a1a* promoter did not have any beneficial effect on scoliosis penetrance (Figure 1I, J). In conclusion, *rpgrip1l* expression in *foxj1a*-expressing cells is sufficient to maintain a straight axis.

Rpgrip1l⁻ᐟ⁻ adult fish display cilia defects along CNS cavities

We observed normal cilia in the neural tube of *rpgrip1l⁻ᐟ⁻* embryos (Supplementary Figure S1C). In 3 month-adults, scanning electron microscopy (SEM) at the level of the rhombencephalic ventricle (RhV) showed cilia tufts of multiciliated ependymal cells (MCC) lining the ventricle in controls and straight *rpgrip1l⁻ᐟ⁻* fish (Supplementary Figure S1F-G, F'-G'). In scoliotic *rpgrip1l⁻ᐟ⁻* fish, cilia tufts were sparse and disorganized (Supplementary Figure S1H, H'). Cilia of mono-ciliated ependymal cells were present in *rpgrip1l⁻ᐟ⁻* fish, some of them with irregular shape (Supplementary Figure S1, F", G", H" and H"').

MCC defects at SCO level correlate with scoliosis appearance at juvenile stages

We then took advantage of the asynchrony in scoliosis onset to correlate, at a given stage, cilia defects at different levels of the CNS cavities with the spine curvature status (straight or curved) of *rpgrip1l^-/-* juveniles. We analyzed cilia defects in three CNS regions implicated in scoliosis appearance: the spinal cord central canal (since scoliosis develops in the spine), the forebrain choroid plexus (fChP) whose cilia defects have been associated with scoliosis in *katnb1^-/-* fish (Meyer-Miner et al., 2022) and the subcommissural organ (SCO), which secretes SCO-spondin whose absence leads to scoliosis (Rose et al., 2020; Troutwine et al., 2020).

In the spinal cord central canal of 8 wpf juvenile fish, cilia density was reduced to the same extent in *rpgrip1l^-/-* fish irrespective of the fish curvature status (Figure 2 A-C, E). In addition, cilia length increased in straight mutants and was more heterogenous than controls in scoliotic mutants (Figure 2 A'-C', D). Arl13b content appeared severely reduced in mutants (Figure 2 A''-C''). In the fChP, mono- and multi-ciliated cell populations were observed with a specific distribution along the anterior-posterior axis, as previously described (D'Gama et al., 2021). At the level of the habenula nuclei, cells of dorsal and ventral midline territories were monociliated, while lateral cells presented multicilia tufts (Supplementary Figure S2 K-K''). *rpgrip1l^-/-* juveniles presented an incomplete penetrance of ciliary defects in the fChP, with no correlation between cilia defects and the curvature status (Supplementary Figure S2, L-O'').

The SCO lies at the dorsal midline of the diencephalic ventricle (DiV), under the posterior commissure (PC) (Figure 2 G, H, Supplementary Figure 2A, C). Cilia in and around the SCO have not been described previously in zebrafish. We showed that, in control animals, Sco-spondin (Sspo) secreting cells varied in number along SCO length, from 24 cells anteriorly to 6-8 cells posteriorly (Supplementary Figure 2 A, A', C) and were monociliated at all anteroposterior levels (Figure 2G-H, Supplementary Figure 2 A'). The Sspo protein produced from one copy of the transgene Scospondin-GFP^ut24 (Troutwine et al., 2020) formed cytoplasmic granules around SCO cell nuclei, under emerging primary cilia (Supplementary Figure 2A'). Conversely, MCCs lateral to the SCO did not secrete Sspo and were prominent towards posterior (Figure 2H, Supplementary Figure 2C), as observed in the rat brain (Collins and Woollam, 1979). Monocilia of Sspo secreting cells were still present but appeared longer in straight *rpgrip1l^-/-* juveniles and could not be individualized anymore at anterior SCO level in

scoliotic fish: we observed a weaker and denser acetylated and glutamylated tubulin
staining that likely corresponded to longer intertwined cilia (Figure 2 I, I', K, K'). Lateral
multicilia tufts were preserved in straight *rpgrip1l^{-/-}* juveniles (n=5) (Figure 2 J) but were
missing at posterior SCO level in tail-up or scoliotic *rpgrip1l^{-/-}* fish (n=7) (Figure 2L).
Thus, *rpgrip1l* mutants develop both monocilia and multicilia defects at juvenile stages,
but cilia defects in the spinal cord and in the fChP do not correlate with axis curvature
status of the fish. By contrast, the loss of MCC cilia at the SCO strictly correlates with
scoliosis onset, suggesting a causal relationship.
173
174 ***Rpgrip1l^{-/-} juveniles show ventricular dilations and loss of the Reissner fiber at***
175 ***scoliosis onset***
176
Ciliary beating is an essential actor of CSF flow and of ventricular development in
zebrafish larvae (Kramer-Zucker et al., 2005) (Olstad et al., 2019). Thus, ciliary defects
in the brain and spinal cord of *rpgrip1l^{-/-}* fish could lead to abnormal ventricular and
central canal volume and content. In the spinal cord, the lumen of the central canal
was enlarged at thoracic and lumbar levels in all *rpgrip1l^{-/-}* mutant juveniles, with no
correlation to the axis curvature status (Figure 2 A-C; F). To determine the timing of
brain ventricular dilations relative to scoliosis, we analyzed ventricular volume at the
onset of spine curvature (Figure 3A-G). Ventricular reconstruction was performed on
cleared brains of 4 control and 4 *rpgrip1l^{-/-}* (3 tail-up and 1 straight) fish at 5 wpf, stained
with ZO1 to highlight the ventricular border and with DiI to outline brain shape, focusing
on the RhV at the level of the posterior midbrain and hindbrain (Figure 3A-D). We
identified a significant increase in ventricle volume in *rpgrip1l^{-/-}* fish compared to
controls, that was restricted to the ventral regions of the RhV in regions ROI4.4 (green
sphere in Figure 3C) and ROI6 (green sphere in Figure 3D), as confirmed by numerical
sections (Figure 3E, G). Ventricle dilations were also present, although to a lesser
extent, in the RhV of the unique straight *rpgrip1l^{-/-}* fish of this study (green squares and
triangle in Figure 3F, G).
Defective ependymal ciliogenesis has been associated with abnormal CSF flow and
defective Reissner Fiber (RF) maintenance in several scoliotic zebrafish models
(Grimes et al., 2016)(Rose et al., 2020). The RF runs along the length of the body,
within brain ventricles and central canal (Figure 3H). It is mainly composed of Sspo

secreted by the SCO and floor plate (FP) and its loss at juvenile stages triggers
scoliosis in zebrafish (Rose et al., 2020)(Troutwine et al., 2020). To visualize the RF,
we used an antiserum that labels the RF in zebrafish embryos (Didier et al.,
1995)(Cantaut-Belarif et al., 2018) and we confirmed the results by introducing one
copy of the *sspo-GFP* knock-in allele, sco-spondin-GFP[ut24,] (Troutwine et al., 2020) by
genetic cross into *rpgrip1l^-/-* animals. The RF formed normally in *rpgrip1l^-/-* embryos, as
expected given the absence of embryonic curvature (Supplementary Figure S1 C). To
study its maintenance in *rpgrip1l^-/-* juveniles, we immunostained spinal cord longitudinal
sections of 7-8 wpf curved or straight fish of the same clutch. The RF was visualized
as a 1 µm diameter rod in the central canal of the neural tube and few dots were
present in the apical cytoplasm of FP cells (Figure 3I) as well as in the SCO of all
controls (n=8, Figure 3L, O) and in ventricles caudal to the SCO, but not rostral to it as
expected (Supplementary Figure S2E, H). The RF was also present in all *rpgrip1l^-/-*
straight fish (n=9, Figure 3J, Supplementary Figure S2I). In contrast, in *rpgrip1l^-/-* tail-
up and scoliotic fish, the RF was absent at all axis levels and a few SCO-spondin-
positive debris were present in the central canal (n=8, Figure 3K). Transverse sections
at the level of the SCO and fChP as well as caudal to the SCO in severely scoliotic fish
revealed abnormally packed material in the ventricles (Figure 3N, Q, Supplementary
Figure S2G, J). In juveniles with a slight tail-up phenotype, SCO-secreting cells
produced Sspo-GFP cytoplasmic granules as in controls (Supplementary Figure S2B,
B', n=7/7), and polymers of Sspo-GFP accumulate in the adjacent DiV, which suggests
that Sspo-GFP aggregates are produced by the SCO.
Thus, our study uncovers a strict temporal correlation between the loss of MCC at
posterior SCO level, impaired RF polymerization and scoliosis onset in *rpgrip1l^-/-*
juvenile fish.
223
224 ***Urp1/2 upregulation does not contribute to scoliosis penetrance or severity in***
225 ***rpgrip1l^-/- fish***
226
In scoliotic fish where the RF is not maintained at juvenile stage, *urp1* and *urp2* levels
are downregulated (Rose et al., 2020). *urp1* and *urp2* encode peptides of the Urotensin
2 family, expressed in ventral spinal CSF-cNs (Parmentier et al., 2011; Quan et al.,
2015). Their expression levels have to be finely tuned to keep a straight axis since their

combined loss-of-function induces larval kyphosis that evolves into adult scoliosis
(Bearce et al., 2022; Gaillard et al., 2023), while an increased dose of Urp1/2 peptides
induces an upward curvature in embryos (Zhang et al., 2018) and juveniles (Gaillard
et al., 2023; Lu et al., 2020). We therefore monitored via qRT-PCR *urp1 and urp2*
expression levels on individual straight or scoliotic *rpgrip1l^-/-* juvenile and/or adult fish.
urp2 was the most highly expressed gene among all *urotensin 2* members at juvenile
stage (Figure 4A, D, E and Supplementary Figure S3A-E. *urp2* expression was
upregulated in 5 weeks juveniles, both in straight and scoliotic fish and its upregulation
was maintained in adult scoliotic fish (Figure 4A).
Given the early and long-lasting upregulation of *urp1/2* in *rpgrip1l^-/-* fish, we attempted
to rescue axis curvature by decreasing global Urp1/2 production via genetic means.
To that end, we generated double (*rpgrip1l^+/-; urp2^+/-*) heterozygous fish using the
recently generated *urp2* deletion allele (Gaillard et al., 2023) and crossed them to
produce double mutants. Using a curvature quantification method on whole fish
(Supplementary Figure S3I and methods), we observed that the removal of one or two
urp2 functional allele(s) in *rpgrip1l^-/-* fish did not lower their curvature index (Figure 4B,
C), neither at two mpf (Supplementary Figure S3 F) nor at four mpf (Supplementary
Figure S3 G-H). We observed that *urp2* mRNA amounts were reduced by removing
one or two functional copies of *urp2*, probably as a consequence of mRNA decay
(Gaillard et al., 2023) (Figure 4D). We also showed that none of the other *urotensin 2*
family members was upregulated by a compensation mechanism upon *urp2* loss
(Figure 4E for *urp1* and Supplementary Figure S3C-E for *urp, uts2a* and *uts2b*).
Thus, lowering global Urp1/2 expression level in *rpgrip1l* mutants did not have any
beneficial effect on the axis curvature phenotype, ruling out altered *urp1/2* signaling as
a main driver of scoliosis in *rpgrip1l* mutants.
256

257 ***Regulators of motile ciliogenesis, inflammation genes and annexins are***
258 ***upregulated in rpgrip1l^-/- juveniles***
259

260 To get further insight into the early mechanisms driving spine curvature in *rpgrip1l*
261 mutants, we obtained the transcriptome of the brain and dorsal trunk of 7 *rpgrip1l^-/-*
262 juveniles at scoliosis onset (6 tail-up and 1 straight 5 wpf juveniles) and 5 control
263 siblings (2 *rgprip1l^+/+* and 3 *rpgrip1l^+/-*). Differential gene expression (DGE) analysis

was performed with the DSEQ2 software within the Galaxy environment. A complete annotated gene list is displayed in Supplementary Table 1 (Trunk) and Table 2 (Brain). Filtering the gene list to select for P-adj < 0.05 and Log2 FC > 0.75 or < -0.75 recovered a vast majority of upregulated genes in the dorsal trunk (75 downregulated vs 614 upregulated genes) and, to a lesser extent, in the brain (1 downregulated, *rpgrip1l,* vs 60 upregulated genes) as illustrated on the volcano plots from DGE analysis of the brain (Figure 5A) and trunk (Figure 5B). 54 out of 60 genes upregulated in the brain were also upregulated in the trunk, suggesting that, in the trunk, these genes were specific to the CNS (Figure 5D). In both the brain (Figure 5C) and the trunk (Supplementary Figure S4A), the most upregulated biological processes in GO term analysis included cilium movement and cilium assembly. Among the 20 most significant GO terms found for upregulated genes in the trunk, 8 were related to ciliary movement/assembly (Supplementary Figure S4A). The upregulation in both tissues of *foxj1a,* encoding a master transcriptional activator of genes involved in cilia motility (Yu et al., 2008) (Figure 5E) likely contributed to this enrichment. To confirm this hypothesis, we compared our dataset with those of targets of Foxj1 from (Choksi et al., 2014; Mukherjee et al., 2019; Quigley and Kintner, 2017). We found that 106 out of 614 upregulated genes were targets of Foxj1, among which 77 were direct targets (Supplementary Table 3). Several genes encoding regulators of embryonic axis curvature were also upregulated in *rpgrip1l*$^{-/-}$ fish, among which *pkd1b* (Mangos et al., 2010), *urp1* and *urp2* (Bearce et al., 2022; Gaillard et al., 2023; Zhang et al., 2018) and *sspo* (Cantaut-Belarif et al., 2018; Rose et al., 2020; Troutwine et al., 2020) (Figure 5E). Finally, GO terms enrichment analysis also highlighted processes associated with inflammation and innate and adaptive immune responses, as described for the scoliotic *ptk7* model (Van Gennip et al., 2018) sharing for example *irg1l* and complement genes up-regulation (Figure 5 C, E, and Supplementary Figure S4A).

In order to confirm these results and to identify processes affected at the post-transcriptional level in *rpgrip1l* mutants, we performed a quantitative proteomic analysis by mass spectrometry of brains from 5 *rpgrip1l*$^{-/-}$ adult fish and 5 control siblings. This analysis detected 5706 proteins present in at least 4 over the 5 replicates, of which 172 are associated with a P value < 0.05 and a Log2FC > +0.75 and < -0.75. 127 proteins were present in higher and 45 in lower quantity, as illustrated on the volcano plot (Figure 5F and Supplementary Table 4). Pathways enriched in *rpgrip1l*$^{-/-}$

brains included regulation of vesicular trafficking and membrane repair processes, as
well as proteolysis and catabolic processes, suggesting profound perturbation of
cellular homeostasis (Supplementary Figure S4B). Among the proteins that were
present in highest quantity in mutants, several members of the Annexin family were
identified: Anxa2a, Anxa1a, Anxa5b (Figure 5F and Supplementary Figure S4C.
Annexins are associated with membranes from endo and exovesicles as well as
plasma membranes and involved in a wide array of intracellular trafficking processes
such as cholesterol endocytosis, plasma membrane repair as well as modulation of
immune system functions (Boye and Nylandsted, 2016; Grewal et al., 2016; Rentero
et al., 2018). Of note, only four proteins of that list, Anxa2a, C4, C7a and Stat3, were
the products of genes up-regulated in the transcriptomic study (indicated in red in
Figure 5E). In agreement with the common enriched GO terms in both studies (Figure
5C and Supplementary Figure S4B), these four proteins are associated with immune
response. Overall, *rpgrip1l-/-* mutants at scoliosis onset present deregulated pathways
in common with other scoliotic models such as activation of immune response genes,
but also specific deregulated pathways such a marked activation of the Foxj1a ciliary
motility program and an increase of Annexin levels.

Astrogliosis precedes scoliosis onset of rpgrip1l-/- juvenile fish

Both transcriptomic and proteomic analysis highlighted the occurrence of innate and
adaptive immune responses in *rpgrip1l-/-* fish. Several upregulated genes, such as
irg1l, mmp9 and anxa2, are reported to be expressed in macrophages and/or microglia
but also in epithelia or neurons (Hall et al., 2014) (Kuhlmann et al., 2014) (Shan et al.,
2015). We therefore investigated in which cell type(s) Anxa2, the most upregulated
protein in the *rpgrip1l-/-* brain proteome, was expressed. Immunostaining of brain
sections revealed a strong Anxa2 staining on cells lining brain ventricles, such as SCO
cells and ventral ependymal cells of the rhombencephalic ventricle (RhV) (Fig 6B', D')
in *rpgrip1l-/-* adult fish, which was barely detectable in controls (Figure 6A', C'), and to
a lesser extent on other brain regions (Supplementary Fig S5 A'-F'), suggesting a
deregulation of ependymal cell homeostasis. In mutants, Anxa2 staining spread over
long cellular extensions reaching the pial surface of the rhombencephalon (Figure 6D'
and Supplementary Figure S5F'). This cell shape is characteristic of radial glial cells,

a cell type that behaves as neural and glial progenitors and also plays astrocytic

functions in zebrafish, as anamniote brains are devoid of stellate astrocytes (Jurisch-

Yaksi et al., 2020; Lyons and Talbot, 2015). To confirm the glial nature of Anxa2-

positive cells around brain ventricles in *rpgrip1l*$^{-/-}$ fish, we double-labeled brain sections

with GFAP, a glial cell marker, and Anxa2. In adult *rpgrip1l*$^{-/-}$ fish, SCO cells were

strongly double-positive for Anxa2 and GFAP (Fig 6B, B'), while controls displayed

much weaker labeling for GFAP in the SCO and no labeling for Anxa2 (Fig 6 A, A').

We made similar observations at rhombencephalic ventricle (RhV) and other brain

levels: ependymal cells were positive for Anxa2 and strongly positive for GFAP, and

GFAP labeling was much higher in mutants than in controls (Fig 6 C-D'; supplementary

Figure S5A-F'). GFAP overexpression is a hallmark of astrogliosis, a process in which

astroglial cells react to the perturbed environment, as shown during brain infection,

injury, ischemia or neurodegeneration (Escartin et al., 2021; Matusova et al., 2023;

Zamanian et al., 2012). Thus, the strong increase of GFAP staining combined with

Anxa2 overexpression in brain ependymal cells of *rpgrip1l*$^{-/-}$ adult fish strongly suggests

that these cells undergo astrogliosis as observed in other animal models following CNS

injury (Chen et al., 2017; Zamanian et al., 2012).

To assay whether an astrogliosis-like phenotype was already present at scoliotic onset,

we immunostained GFAP on juvenile brain sections of straight and tail-up mutants and

controls. Tail-up juveniles (n=3/3) presented a strong GFAP staining in almost all cells

of the SCO and along the ventral rhombencephalic ventricle, higher than control levels

(Figure 6 E, I, H, L, M) while the phenotype of straight fish was heterogeneous: two out

of four juveniles presented GFAP-positive cells (Figure 6G, K and red dots on Figure

6M graph), while the two other did not (Figure 6F, J and green dots on Figure 6M

graph) as confirmed by cell quantification (Figure 6M). As scoliosis onset is

asynchronous in *rpgrip1l*$^{-/-}$, heterogenous GFAP staining among straight mutant fish

may support an early role of astrogliosis at SCO and RhV levels in scoliosis onset.

Immune cell enrichment around the SCO and within tectum parenchyma is

detected prior to scoliosis onset.

Astrogliosis may be reinforced or induced by an inflammatory environment containing

cytokines released by microglia or macrophages or the presence of high levels of ROS

and NO (Sofroniew, 2015). We thus performed immunostaining for the LCP1/L-Plastin
marker, which labels both macrophages and microglia (Figure 6N-Q'), in adjacent
sections to those used for GFAP (Figure 6E-L). We found that the scoliotic fish
presented a higher number of LCP1-positive cells of ameboid shape within SCO cells
(Figure 6Q, Q', R). Strikingly, the straight mutants (2/4) that presented a high GFAP
staining within SCO cells (Figure 6G, K) also showed a higher number of LCP1-positive
cells around the SCO (Figure 6P, P'; red dots in Figure 6R) than in controls (Figure
6N, N'), while the other straight mutants did not ((Figure 6F, J, O, O', green dots in
Figure 6M) suggesting positive regulation between astrogliosis and
macrophages/microglia recruitment. LCP1+ cells were also detected in other regions
of the brain (Supplementary Figure S5 G-K). In the optic tectum, where resident LCP1-
positive macrophages/microglia were present in controls, a two-fold increase in the
number of ramified LCP1$^+$ cells was observed in straight $rpgrip1l^{-/-}$ juveniles compared
to controls (Supplementary Figure S5H-J). Surprisingly, scoliotic $rpgrip1l^{-/-}$ fish showed
similar LCP1$^+$ cell number as control fish in the tectum (Supplementary Figure S5G),
suggesting a transient increase of macrophage/microglia activation preceding
scoliosis.
Our data thus uncovers an early astrogliosis process initiated asynchronously at the
SCO and RhV levels of straight $rpgrip1l^{-/-}$ fish, favored by a transient inflammatory
state, and preceding the loss of multiciliated tufts around the SCO and that of the RF,
that we only detected in curved $rpgrip1l^{-/-}$ fish. Astrogliosis then spreads along the brain
ventricles and becomes prominent in brain of scoliotic fish. In addition, we found among
the upregulated genes in dorsal trunk at scoliosis onset, a number of activated
astrocyte markers (Matusova et al., 2023) such as $s100b$, $c4b$, $gfap$, $glast/slc1a3b$,
$viml$, $ctsb$ (Supplementary Figure S4D), showing that astrogliosis spread over the
whole central nervous system.
389
390 ***Brain neuroinflammation and astrogliosis contribute to scoliosis penetrance***
391 ***and severity in rpgrip1l$^{-/-}$ fish***
392
393 Our data suggest that astrogliosis and neuro-inflammation could play a role in
394 triggering scoliosis in $rpgrip1l$ mutants. We thus aimed to counteract inflammation
395 before curvature initiation by performing drug treatment with the hope of reducing

scoliosis penetrance and severity. We used the NACET anti-oxidant and anti-inflammatory drug that successfully reduced scoliosis of *sspo* hypomorphic mutant fish (Rose et al., 2020). We first assayed the biological activity of our NACET batch on the ciliary motility mutant *dnaaf1⁻/⁻* (Sullivan-Brown et al., 2008). Indeed, NACET was shown to suppress the embryonic tail-down phenotype of the ciliary motility mutant *ccdc151ᵗˢ* (Rose et al., 2020). We found that 3mM NACET treatment suppressed the *dnaaf1⁻/⁻* tail-down phenotype (Supplementary Figure S6A-F). To suppress scoliosis, we treated the progeny of *rpgrip1l⁺/⁻* incrosses from 4 to 12 wpf with 1.5 mM NACET as in (Rose et al., 2020).This long-term treatment reduced scoliosis penetrance from 92% to 58% and markedly reduced spinal curvature index (Figure 7A-C), measured and illustrated as shown in Supplementary Figure S3I.

We then asked whether NACET treatment suppressed astrogliosis and neuroinflammation in *rpgrip1l⁻/⁻* treated fish. Indeed, the intensity and number of GFAP+ cells within the SCO were reduced in slightly curved fish and alleviated in straight fish (Figure 7E, F). Furthermore, the quantification of Lcp1-positive cells in the SCO region showed a drastic reduction of these cells in NACET-treated fish, back to levels found in *rpgrip1l⁺/⁺* fish (Figure 7D, E).

Our data thus show that *rpgrip1l⁻/⁻* juvenile fish display brain astrogliosis and inflammation at scoliosis onset and that an anti-inflammatory/anti-oxidant treatment of *rpgrip1l⁻/⁻* juveniles is able to maintain a straight axis in approximately one third of the mutants and to slow down curve progression in the other two thirds. The beneficial effect of NACET on axis curvature correlates with the combined reduction of astrogliosis and immune cell density.

Brain astrogliosis is present in zebrafish *cep290 ᶠʰ²⁹⁷/ᶠʰ²⁹⁷* scoliotic fish

To investigate whether abnormal SCO cilia function and brain astrogliosis may be a general mechanism of scoliosis development in TZ mutants, we studied the *cep290ᶠʰ²⁹⁷/ᶠʰ²⁹⁷* zebrafish mutant. Zebrafish *cep290* mutants display variable axis curvature defects at embryonic stages as well as slow retinal degeneration, and develop juvenile scoliosis (Lessieur et al., 2019; Wang et al., 2022). Like *RPGRIP1L*, the human *CEP290* gene encodes a TZ protein and its mutation leads to severe neurodevelopmental ciliopathies (Baala et al., 2007). All *cep290ᶠʰ²⁹⁷/ᶠʰ²⁹⁷* juveniles

developed a fully penetrant scoliosis by 4 wpf in our fish facility (n: 30/30). We analyzed

cilia presence within the SCO by immunostaining on sections (Figure 8A-F'). At 4 wpf,

the SCO had not reached its final adult width (14 versus 24 cells wide in the anterior

SCO), and each of its cells harbored a short monocilium labeled by glutamylated

tubulin (Figure 8A-C). Multicilia tufts on SCO lateral walls were not as clearly visible as

in adults (Figure 2F) but patches of glutamylated tubulin staining were detected on

both sides of the posterior SCO (Figure 8B-C'). In $cep290^{-/-}$, monocilia of the SCO

appeared longer than in controls (N=3/3) and pointed towards an abnormally elongated

diencephalic ventricle (Figure 8D-F). Lateral patches of glutamylated tubulin staining

were present in $cep290^{-/-}$ posterior SCO as in controls (Figure 8E-F'). Thus, cilia

morphology is perturbed within $cep290^{-/-}$ SCO monociliated cells, as well as the overall

morphology of the diencephalic ventricle at this level.

To test for the presence of astrogliosis, we quantified the number of SCO cells

presenting high GFAP labeling on several sections along the A/P extent of the SCO in

3 control and 3 $cep290^{-/-}$ brains (Figure 8G-K). 50% of SCO cells presented strong

GFAP staining in fully scoliotic juveniles, which was not detectable in controls (Figure

8H, K, N=2/3), while a lower proportion (22%) of SCO cells was GFAP positive in the

juvenile with a milder curvature (Figure 8M, N=1/3). Strong GFAP staining was also

detected in some diencephalic radial glial cells (Figure 8K). On the same sections,

immune cells were quantified by LCP1 staining. A mean of 4 LCP1 positive cells per

SCO field was found in $cep290^{-/-}$, versus 1 in controls (Figure 8I, L and N). As in $rpgrip1l$

mutants, abnormal monocilia, astrogliosis and immune cell enrichment were evidenced

at SCO levels in $Cep290^{-/-}$(3/3). Furthermore, astrogliosis was observed within radial

glial cells lining the RhV at cerebellar level (Figure 8O-R) and at anterior hindbrain level

(Figure 8S-V), as shown in $rpgrip1l^{-/-}$ scoliotic brain (Figure 6D, K, L). This indicates

that astrogliosis is shared by these two zebrafish scoliotic models, even though they

develop axis curvature at different stages (3-4 wpf for $cep290^{-/-}$ versus 5-12 wpf for

$rpgrip1l^{-/-}$).

DISCUSSION

In this paper we dissected the mechanisms of scoliosis appearance in a zebrafish

mutant for the TZ gene $rpgrip1l$, taking advantage of its asynchronous onset to

decipher the chronology of events leading to spine torsion. Transcriptome analysis of
the brain and trunk at scoliosis onset in *rpgrip1l^-/-* fish compared to control siblings
revealed increased expression of *urp1/2*, *foxj1a* and its target genes, as well as genes
linked to immune response. Genetic experiments ruled out an involvement of
increased Urp1/2 signaling in scoliosis development. Cilia defects appeared
asynchronously in different CNS regions, while RF loss strictly coincided with scoliosis
onset. Strikingly, we observed a strong increase of GFAP staining along the
diencephalic and rhombencephalic ventricles, indicative of astrogliosis. This
astrogliosis preceded scoliosis onset and then spread posteriorly and increased in
severity. In the region of the SCO, the appearance of astrogliosis and
neuroinflammation coincided temporally, preceding the onset of RF depolymerization
and spine torsion. NACET treatment showed that increased astrogliosis and
inflammation participated in the penetrance and severity of scoliosis in *rpgrip1l^-/-* fish.
Finally, we showed that another TZ gene mutant, *cep290*, also displayed astrogliosis
at the time of scoliosis onset. Thus, the cascade of events uncovered in the *rpgrip1l*
mutant sheds a new light on the origin of zebrafish scoliosis caused by ciliary TZ
defects.
479
Cilia motility defects lead to RF loss and to scoliosis in several zebrafish mutants
(Grimes et al., 2016; Rose et al., 2020)). *Rpgrip1l* encodes a TZ protein important for
cilia integrity and function in many eukaryotic organisms (Andreu-Cervera et al., 2019;
Gogendeau et al., 2020; Jensen et al., 2015; Wiegering et al., 2018). Our data show
that *rpgrip1l* deficiency in zebrafish affects cilia maintenance differentially depending
on the CNS territory and the developmental stage, raising the question of which ciliated
cells are responsible for RF loss and scoliosis initiation in this mutant. The full rescue
observed after re-introducing RPGRIP1L in *foxj1a*-expressing cells points towards a
crucial function of the protein in cells lining CNS cavities. Foxj1a-positive cell
populations encompass both monociliated and multiciliated ependymal cells, radial
glial progenitors, glial cells of the SCO and the ChP, as well as CSF-contacting sensory
neurons in the spinal cord (D'Gama et al., 2021; Prendergast et al., 2023; Ribeiro et
al., 2017; Van Gennip et al., 2018). Recent work suggested that impairment of
ciliogenesis in the fChP of the *katnb1* mutant could play a role in inducing axis
curvature (Meyer-Miner et al., 2022). In the case of the *rpgrip1l* mutant, we temporally

uncorrelated fChP cilia defects from scoliosis onset. This, combined with the fact that

the total loss of multicilia in the fChP of [*foxj1b^-/-, gmnc^-/-*] double mutants does not

trigger axis curvature (D'Gama et al., 2021) suggests that another *foxj1a*-expressing

population is crucial for axis straightness. Our results strongly suggest that cilia defects

in the SCO trigger axis curvature in *rpgrip1l* mutants. Two cilia populations were

affected in the SCO of *rpgrip1l* mutant juveniles at the onset of scoliosis: cilia of

monociliated glial cells that secrete SCO-spondin, which appeared longer in mutants

than in controls, and multiciliated tufts at SCO exit, which were lost in scoliotic mutants.

Both defects could be responsible for the phenotype, by perturbing CSF flow and

content (presence of SCO-spondin aggregates in ventricles) (left side on the Graphical

Abstract) and leading to ventricular dilations or by perturbing primary cilia signaling

within SCO cells (right side of the Graphical Abstract). The observation that

multiciliated cells were not yet fully differentiated in *cep290* mutants at scoliosis onset

is in favor of a prominent role of monocilia. This is consistent with the literature showing

that most ciliary mutant models develop axis curvature at 3-4 weeks, a stage when

brain multiciliated cells are not yet differentiated (D'Gama et al., 2021). Still, as

scoliosis onset in *rpgrip1l* mutant is asynchronous from 5 to 12 weeks, MCC loss at

SCO exit could contribute to defective RF polymerization.

Our model for scoliosis appearance in the *rpgrip1l* mutant is presented in the graphical

abstract. Since we have observed several straight *rpgrip1l^-/-* fish with astrogliosis and

increased immune cells presence around the SCO, we propose that *rpgrip1l^-/-* SCO

and astroglial cells experience altered signaling that activates immune cell recruitment,

producing an inflammatory environment. This would promote multicilia loss as a

secondary event (Graphical Abstract, orange arrow) (Lattke et al., 2012) leading to RF

depolymerization and spine curvature. However, as *sspo^-/-* hypomorphic mutants

which present aggregates within ventricles have been shown to present an

inflammatory response (Rose et al., 2020), it is equally possible that SCO astrogliosis

is a downstream consequence of the presence of these proteoglycans aggregates in

contact with primary cilia or ventricular cell surfaces (Graphical Abstract , green arrow).

These two scenarios may also act in parallel to reinforce scoliosis onset and

ependymal cilia loss. Future experiments will help to discriminate between these two

scenarios.

Our transcriptome analysis and qRT-PCR data challenges a model proposed for axis straightness of zebrafish embryos in which the loss of RF leads to a down-regulation of *urp2* and *urp1* expression (Zhang et al., 2018) since we observed an upregulation of both genes before and after RF loss. As forced expression of *urp1/2* induces a tail-up phenotype in embryos and juveniles (Zhang et al., 2018) (Gaillard et al., 2023; Lu et al., 2020), we attempted to rescue *rpgrip1l*[-/-] axis curvature by down-regulating *urp1/2* expression. No beneficial effect on scoliosis penetrance or severity was observed, indicating that increased *urp1/2* expression level does not significantly contribute to axis curvature in *rpgrip1l*[-/-] fish.

In this study, we found an unexpected and strong defect characterized by enhanced GFAP and Anxa2 expression, which we identified as astrogliosis (also called reactive astrogliosis). Astrogliosis is a reaction of astroglial cells to perturbed homeostasis of the CNS, characterized by molecular and phenotypic changes in these cells (Escartin et al., 2021; Matusova et al., 2023). In zebrafish, radial glial cells are endowed with both neurogenic and astrocytic functions, and are thus also called astroglial cells (Jurisch-Yaksi et al., 2020). Astrogliosis in *rpgrip1l* mutants arose in a subdomain of Foxj1a-positive cells, within the DiV and ventral ependymal cells lining the RhV. The observation of astrogliosis in some straight juvenile mutants suggests an early role of this defect in scoliosis onset.

Astrogliosis along the ventricular cavities could arise in response to local mechanical or chemical perturbations caused by cilia motility defects, abnormal CSF flow and content and ventricular dilations. In two murine models of primary ciliary dyskinesia, decreased CSF flow was associated with gliosis at juvenile stage (Finn et al., 2014). A response to CSF pertubation in *rpgrip1l* mutants is also suggested by the strong upregulation of Foxj1a and its target genes. *foxj1a* expression is also reported to be upregulated in the CNS of three other scoliotic models in addition to *rpgrip1l*, namely in *ptk7, sspo* and *katnb1* mutants (Escartin et al., 2021; Meyer-Miner et al., 2022; Van Gennip et al., 2018), suggesting that it constitutes a common response to similar ventricular and CSF defects. Interestingly, *foxj1a* expression is also upregulated upon zebrafish embryonic (Cavone et al., 2021; Hellman et al., 2010) or adult (Ribeiro et al., 2017) spinal cord injury. The up-regulation in the *rpgrip1l* mutant proteome of the

membrane repair module Anxa2-S100a10-Ahnak (Bharadwaj et al., 2021) (Figure 4C)

together with the Foxj1-induced motility module detected in the transcriptomic analysis

may indicate ongoing reparation attempts of CNS ventricles and associated neural

tissues.

Moreover, the presence of few Lcp1+ cells around the mutant SCO suggests that

microglia/macrophages could participate in astrogliosis establishment and

reinforcement. This dialogue between immune Lcp1+ cells and astroglial cells is

evidenced during early stages of regeneration after acute injury or cellular damages

(Cavone et al., 2021). In our bulk RNAseq analysis performed at scoliosis onset, we

have indications of the upregulation of microglia and macrophage markers (*p2ry12* and

mpeg1) at trunk levels and of numerous markers of activated astrocytes characterized

in several mammalian astrogliosis models (Matusova et al., 2023). A similar dialogue

is detected in polycystic kidney disease models where compromised primary cilia

signalling leads to uncontrolled cytokines secretion by epithelial cells, a situation that

favors local immune cells recruitment and proliferation (Viau et al., 2018).

What are the intracellular mechanisms involved in astrogliosis induction in ciliated

ventricular cells? Candidate pathways emerge from our multi-omics studies. Proteomic

data showed a 2.4-fold increase in GSK3bb amount in *rpgrip1l*[-/-] mutant brains (P value

< 0.001, Supplementary Table 3). In several neurodegenerative murine animal models,

GSK3b enzymatic activity was shown to promote inflammation and gliosis (Jorge-

Torres et al., 2018; Medina and Avila, 2010; Mines et al., 2011). Another potential

trigger of astrogliosis may be a defective mitochondrial activity as the amounts of four

proteins involved in electron transport chain activity (*gpd1b, mt-nd6, pdia5 and dmgdh*)

were reduced by 40% to 86% in the analysis of mutant brain proteome (Supplementary

Table 4). We think this is of particular interest in light of a recent report demonstrating

that impaired mitochondrial activity within ciliated astrocytes leads to the induction of

the Foxj1 ciliary motility program, the elongation and distortion of astrocyte primary

cilia as well as reactive astrogliosis (Ignatenko et al., 2023), three phenotypes also

observed in *rpgrip1l*[-/-] brains.

Finally, our observation of widespread astrogliosis in two TZ gene mutants, *rpgrip1l*

and *cep290*, suggests that it could represent a general mechanism involved in scoliosis

downstream of cilia dysfunction. Of note, astrogliosis markers such as *glast/slc1a3b*,

vim, *s100b*, *c4*, *timp2b*, *gfap* and *ctssb.2*, are also upregulated in the transcriptome of
ptk7 mutants (Van Gennip et al., 2018). We propose that sustained astrogliosis might
impair neuronal survival and activity, crucial for proper interoception and locomotor
control, thus leading to axis curvature in the context of a rapidly growing axial skeleton.
A similar context of astrocytosis associated with spinal cord dilation has been observed
in human patients with syringomyelia. This population with a high incidence of scoliosis
can present hyper-signals on T2 MRI scans, which were proposed to reflect local
astrocytosis and could be validated by biopsies in rare cases (Sherman et al., 1987).
Further imaging studies will need to be performed to validate a potential link between
developing astrogliosis and the onset and progression of idiopathic scoliosis in
humans.
605

606 **ACKNOWLEDGEMENTS**
607

We are grateful to the IBPS aquatic animal, imaging and bioinformatics facilities and
to the ICM sequencing facility for their technical assistance. We thank Michaël Trichet
from the IBPS electron microscopy facility for participating in the MEB experiments,
the TACGENE facility for providing the Cas9 protein; the TEFOR Paris-Saclay facility
for the brain clearing experiment; Thierry Jaffredo and Pierre Charbord for their
precious help in transcriptome analysis; Nicolas Baylé for initial characterization of the
rpgrip1l mutant; Claudia Hoffman for providing *Cep290* fixed samples; Brian Ciruna for
sharing the *foxj1a* expressing constructs and detailed NACET treatment protocol; Ryan
Gray for sharing the *scospondin-GFP*[ut24] zebrafish line. This work was supported by
funding to SSM from the Fondation pour la Recherche Médicale (Equipe FRM
EQU201903007943) and the Fondation Yves Cotrel.
619

REFERENCES

Affaticati P, Simion M, De Job E, Rivière L, Hermel J-M, Machado E, Joly J-S, Jenett A. 2017. zPACT: Tissue Clearing and Immunohistochemistry on Juvenile Zebrafish Brain. *BIO-PROTOCOL* **7**. doi:10.21769/BioProtoc.2636

Andreu-Cervera A, Anselme I, Karam A, Laclef C, Catala M, Schneider-Maunoury S. 2019. The ciliopathy gene *ftm/rpgrip1l* controls mouse forebrain patterning via region-specific modulation of hedgehog/gli signaling. *J Neurosci* 2199–18. doi:10.1523/JNEUROSCI.2199-18.2019

Andreu-Cervera A, Catala M, Schneider-Maunoury S. 2021. Cilia, ciliopathies and hedgehog-related forebrain developmental disorders. *Neurobiology of Disease* **150**:105236. doi:10.1016/j.nbd.2020.105236

Arts HH, Doherty D, Van Beersum SEC, Parisi MA, Letteboer SJF, Gorden NT, Peters TA, Märker T, Voesenek K, Kartono A, Ozyurek H, Farin FM, Kroes HY, Wolfrum U, Brunner HG, Cremers FPM, Glass IA, Knoers NVAM, Roepman R. 2007. Mutations in the gene encoding the basal body protein RPGRIP1L, a nephrocystin-4 interactor, cause Joubert syndrome. *Nat Genet* **39**:882–888. doi:10.1038/ng2069

Baala L, Audollent S, Martinovic J, Ozilou C, Babron M-C, Sivanandamoorthy S, Saunier S, Salomon R, Gonzales M, Rattenberry E, Esculpavit C, Toutain A, Moraine C, Parent P, Marcorelles P, Dauge M-C, Roume J, Merrer ML, Meiner V, Meir K, Menez F, Beaufrère A-M, Francannet C, Tantau J, Sinico M, Dumez Y, MacDonald F, Munnich A, Lyonnet S, Gubler M-C, Génin E, Johnson CA, Vekemans M, Encha-Razavi F, Attié-Bitach T. 2007. Pleiotropic Effects of CEP290 (NPHP6) Mutations Extend to Meckel Syndrome. *The American Journal of Human Genetics* **81**:170–179. doi:10.1086/519494

Bachmann-Gagescu R, Phelps IG, Stearns G, Link BA, Brockerhoff SE, Moens CB, Doherty D. 2011. The ciliopathy gene cc2d2a controls zebrafish photoreceptor outer segment development through a role in Rab8-dependent vesicle trafficking. *Human Molecular Genetics* **20**:4041–4055. doi:10.1093/hmg/ddr332

Bearce EA, Irons ZH, O'Hara-Smith JR, Kuhns CJ, Fisher SI, Crow WE, Grimes DT. 2022. Urotensin II-related peptides, Urp1 and Urp2, control zebrafish spine morphology. *eLife* **11**:e83883. doi:10.7554/eLife.83883

Bentley-Ford MR, LaBonty M, Thomas HR, Haycraft CJ, Scott M, LaFayette C, Croyle MJ, Andersen RS, Parant JM, Yoder BK. 2022. Evolutionarily conserved genetic interactions between *nphp-4* and *bbs-5* mutations exacerbate ciliopathy phenotypes. *Genetics* **220**:iyab209.

doi:10.1093/genetics/iyab209

Bharadwaj A, Kempster E, Waisman DM. 2021. The Annexin A2/S100A10 Complex: The Mutualistic Symbiosis of Two Distinct Proteins. *Biomolecules* **11**:1849. doi:10.3390/biom11121849

Boswell CW, Ciruna B. 2017. Understanding Idiopathic Scoliosis: A New Zebrafish School of Thought. *Trends in Genetics* **33**:183–196. doi:10.1016/j.tig.2017.01.001

Bouyssié D, Hesse A-M, Mouton-Barbosa E, Rompais M, Macron C, Carapito C, Gonzalez De Peredo A, Couté Y, Dupierris V, Burel A, Menetrey J-P, Kalaitzakis A, Poisat J, Romdhani A, Burlet-Schiltz O, Cianférani S, Garin J, Bruley C. 2020. Proline: an efficient and user-friendly software suite for large-scale proteomics. *Bioinformatics* **36**:3148–3155. doi:10.1093/bioinformatics/btaa118

Boye TL, Nylandsted J. 2016. Annexins in plasma membrane repair. *Biological Chemistry* **397**:961–969. doi:10.1515/hsz-2016-0171

Brancati F, Travaglini L, Zablocka D, Boltshauser E, Accorsi P, Montagna G, Silhavy JL, Barrano G, Bertini E, Emma F, Rigoli L, International JSRD Study Group, Dallapiccola B, Gleeson JG, Valente EM. 2008. RPGRIP1L mutations are mainly associated with the cerebello-renal phenotype of Joubert syndrome-related disorders. *Clin Genet* **74**:164–170. doi:10.1111/j.1399-0004.2008.01047.x

Cantaut-Belarif Y, Orts Del'Immagine A, Penru M, Pézeron G, Wyart C, Bardet P-L. 2020. Adrenergic activation modulates the signal from the Reissner fiber to cerebrospinal fluid-contacting neurons during development. *eLife* **9**:e59469. doi:10.7554/eLife.59469

Cantaut-Belarif Y, Sternberg JR, Thouvenin O, Wyart C, Bardet P-L. 2018. The Reissner Fiber in the Cerebrospinal Fluid Controls Morphogenesis of the Body Axis. *Current Biology* **28**:2479-2486.e4. doi:10.1016/j.cub.2018.05.079

Cavone L, McCann T, Drake LK, Aguzzi EA, Oprişoreanu A-M, Pedersen E, Sandi S, Selvarajah J, Tsarouchas TM, Wehner D, Keatinge M, Mysiak KS, Henderson BEP, Dobie R, Henderson NC, Becker T, Becker CG. 2021. A unique macrophage subpopulation signals directly to progenitor cells to promote regenerative neurogenesis in the zebrafish spinal cord. *Developmental Cell* **56**:1617-1630.e6. doi:10.1016/j.devcel.2021.04.031

Chen J, Cui Z, Yang S, Wu C, Li W, Bao G, Xu G, Sun Y, Wang L, Zhang J. 2017. The upregulation of annexin A2 after spinal cord injury in rats may have implication for astrocyte proliferation. *Neuropeptides* **61**:67–76. doi:10.1016/j.npep.2016.10.007

Choksi SP, Lauter G, Swoboda P, Roy S. 2014. Switching on cilia: transcriptional networks

regulating ciliogenesis. *Development* **141**:1427–1441. doi:10.1242/dev.074666

Collins P, Woollam DH. 1979. The ventricular surface of the subcommissural organ: a scanning

and transmission electron microscopic study. *J Anat* **129**:623–631.

Dale RM, Topczewski J. 2011. Identification of an evolutionarily conserved regulatory element

of the zebrafish col2a1a gene. *Developmental Biology* **357**:518–531.

doi:10.1016/j.ydbio.2011.06.020

Delous M, Baala L, Salomon R, Laclef C, Vierkotten J, Tory K, Golzio C, Lacoste T, Besse L,

Ozilou C, Moutkine I, Hellman NE, Anselme I, Silbermann F, Vesque C, Gerhardt C,

Rattenberry E, Wolf MTF, Gubler MC, Martinovic J, Encha-Razavi F, Boddaert N, Gonzales

M, Macher MA, Nivet H, Champion G, Berthélémé JP, Niaudet P, McDonald F, Hildebrandt

F, Johnson CA, Vekemans M, Antignac C, Rüther U, Schneider-Maunoury S, Attié-Bitach T,

Saunier S. 2007. The ciliary gene RPGRIP1L is mutated in cerebello-oculo-renal syndrome

(Joubert syndrome type B) and Meckel syndrome. *Nat Genet* **39**:875–881. doi:10.1038/ng2039

D'Gama PP, Qiu T, Cosacak MI, Rayamajhi D, Konac A, Hansen JN, Ringers C, Acuña-

Hinrichsen F, Hui SP, Olstad EW, Chong YL, Lim CKA, Gupta A, Ng CP, Nilges BS, Kashikar

ND, Wachten D, Liebl D, Kikuchi K, Kizil C, Yaksi E, Roy S, Jurisch-Yaksi N. 2021. Diversity

and function of motile ciliated cell types within ependymal lineages of the zebrafish brain. *Cell*

Reports **37**:109775. doi:10.1016/j.celrep.2021.109775

Didier R, Dastugue B, Meiniel A. 1995. The secretory material of the subcommissural organ of

the chick embryo. Characterization of a specific polypeptide by two-dimensional

electrophoresis. *Int J Dev Biol* **39**:493–499.

Escartin C, Galea E, Lakatos A, O'Callaghan JP, Petzold GC, Serrano-Pozo A, Steinhäuser C,

Volterra A, Carmignoto G, Agarwal A, Allen NJ, Araque A, Barbeito L, Barzilai A, Bergles

DE, Bonvento G, Butt AM, Chen W-T, Cohen-Salmon M, Cunningham C, Deneen B, De

Strooper B, Díaz-Castro B, Farina C, Freeman M, Gallo V, Goldman JE, Goldman SA, Götz

M, Gutiérrez A, Haydon PG, Heiland DH, Hol EM, Holt MG, Iino M, Kastanenka KV,

Kettenmann H, Khakh BS, Koizumi S, Lee CJ, Liddelow SA, MacVicar BA, Magistretti P,

Messing A, Mishra A, Molofsky AV, Murai KK, Norris CM, Okada S, Oliet SHR, Oliveira JF,

Panatier A, Parpura V, Pekna M, Pekny M, Pellerin L, Perea G, Pérez-Nievas BG, Pfrieger FW,

Poskanzer KE, Quintana FJ, Ransohoff RM, Riquelme-Perez M, Robel S, Rose CR, Rothstein

JD, Rouach N, Rowitch DH, Semyanov A, Sirko S, Sontheimer H, Swanson RA, Vitorica J,

Wanner I-B, Wood LB, Wu J, Zheng B, Zimmer ER, Zorec R, Sofroniew MV, Verkhratsky A.

2021. Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci*

24:312–325. doi:10.1038/s41593-020-00783-4

Finn R, Evans CC, Lee L. 2014. Strain-dependent brain defects in mouse models of primary ciliary dyskinesia with mutations in Pcdp1 and Spef2. *Neuroscience* **277**:552–567. doi:10.1016/j.neuroscience.2014.07.029

Gaillard A-L, Mohamad T, Quan FB, De Cian A, Mosimann C, Tostivint H, Pézeron G. 2023. Urp1 and Urp2 act redundantly to maintain spine shape in zebrafish larvae. *Developmental Biology* **496**:36–51. doi:10.1016/j.ydbio.2023.01.010

Gogendeau D, Lemullois M, Le Borgne P, Castelli M, Aubusson-Fleury A, Arnaiz O, Cohen J, Vesque C, Schneider-Maunoury S, Bouhouche K, Koll F, Tassin A-M. 2020. MKS-NPHP module proteins control ciliary shedding at the transition zone. *PLoS Biol* **18**:e3000640. doi:10.1371/journal.pbio.3000640

Grewal T, Wason SJ, Enrich C, Rentero C. 2016. Annexins – insights from knockout mice. *Biological Chemistry* **397**:1031–1053. doi:10.1515/hsz-2016-0168

Grimes DT, Boswell CW, Morante NFC, Henkelman RM, Burdine RD, Ciruna B. 2016. Zebrafish models of idiopathic scoliosis link cerebrospinal fluid flow defects to spine curvature. *Science* **352**:1341–1344. doi:10.1126/science.aaf6419

Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud J-B, Schneider-Maunoury S, Shkumatava A, Teboul L, Kent J, Joly J-S, Concordet J-P. 2016. Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol* **17**:148. doi:10.1186/s13059-016-1012-2

Hall CJ, Boyle RH, Sun X, Wicker SM, Misa JP, Krissansen GW, Print CG, Crosier KE, Crosier PS. 2014. Epidermal cells help coordinate leukocyte migration during inflammation through fatty acid-fuelled matrix metalloproteinase production. *Nat Commun* **5**:3880. doi:10.1038/ncomms4880

Hayes M, Gao X, Yu LX, Paria N, Henkelman RM, Wise CA, Ciruna B. 2014. ptk7 mutant zebrafish models of congenital and idiopathic scoliosis implicate dysregulated Wnt signalling in disease. *Nat Commun* **5**:4777. doi:10.1038/ncomms5777

Hellman NE, Liu Y, Merkel E, Austin C, Le Corre S, Beier DR, Sun Z, Sharma N, Yoder BK, Drummond IA. 2010. The zebrafish *foxj1a* transcription factor regulates cilia function in response to injury and epithelial stretch. *Proc Natl Acad Sci USA* **107**:18499–18504. doi:10.1073/pnas.1005998107

Hitier M, Hamon M, Denise P, Lacoudre J, Thenint M-A, Mallet J-F, Moreau S, Quarck G. 2015. Lateral Semicircular Canal Asymmetry in Idiopathic Scoliosis: An Early Link between

Biomechanical, Hormonal and Neurosensory Theories? *PLoS ONE* **10**:e0131120.

doi:10.1371/journal.pone.0131120

Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, Krijgsveld J. 2014.

Ultrasensitive proteome analysis using paramagnetic bead technology. *Molecular Systems*

Biology **10**:757. doi:10.15252/msb.20145625

Ignatenko O, Malinen S, Rybas S, Vihinen H, Nikkanen J, Kononov A, Jokitalo ES, Ince-Dunn

G, Suomalainen A. 2023. Mitochondrial dysfunction compromises ciliary homeostasis in

astrocytes. *Journal of Cell Biology* **222**:e202203019. doi:10.1083/jcb.202203019

Jensen VL, Li C, Bowie RV, Clarke L, Mohan S, Blacque OE, Leroux MR. 2015. Formation

of the transition zone by Mks5/Rpgrip1L establishes a ciliary zone of exclusion (CIZE) that

compartmentalises ciliary signalling proteins and controls PIP $_2$ ciliary abundance. *The EMBO*

Journal **34**:2537–2556. doi:10.15252/embj.201488044

Jorge-Torres OC, Szczesna K, Roa L, Casal C, Gonzalez-Somermeyer L, Soler M, Velasco CD,

Martínez-San Segundo P, Petazzi P, Sáez MA, Delgado-Morales R, Fourcade S, Pujol A,

Huertas D, Llobet A, Guil S, Esteller M. 2018. Inhibition of Gsk3b Reduces Nfkb1 Signaling

and Rescues Synaptic Activity to Improve the Rett Syndrome Phenotype in Mecp2-Knockout

Mice. *Cell Reports* **23**:1665–1677. doi:10.1016/j.celrep.2018.04.010

Jurisch-Yaksi N, Yaksi E, Kizil C. 2020. Radial glia in the zebrafish brain: Functional,

structural, and physiological comparison with the mammalian glia. *Glia* **68**:2451–2470.

doi:10.1002/glia.23849

Karner CM, Long F, Solnica-Krezel L, Monk KR, Gray RS. 2015. *Gpr126/Adgrg6* deletion in

cartilage models idiopathic scoliosis and pectus excavatum in mice. *Hum Mol Genet* **24**:4365–

4373. doi:10.1093/hmg/ddv170

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of embryonic

development of the zebrafish. *Developmental Dynamics* **203**:253–310.

doi:10.1002/aja.1002030302

Kramer-Zucker AG, Olale F, Haycraft CJ, Yoder BK, Schier AF, Drummond IA. 2005. Cilia-

driven fluid flow in the zebrafish pronephros, brain and Kupffer's vesicle is required for normal

organogenesis. *Development* **132**:1907–1921. doi:10.1242/dev.01772

Kuhlmann K, Tschapek A, Wiese H, Eisenacher M, Meyer HE, Hatt HH, Oeljeklaus S,

Warscheid B. 2014. The Membrane Proteome of Sensory Cilia to the Depth of Olfactory

Receptors. *Molecular & Cellular Proteomics* **13**:1828–1843. doi:10.1074/mcp.M113.035378

Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, Parant JM, Yost

HJ, Kanki JP, Chien C. 2007. The Tol2kit: A multisite gateway-based construction kit for *Tol2*
transposon transgenesis constructs. *Developmental Dynamics* **236**:3088–3099.
doi:10.1002/dvdy.21343

Lattke M, Magnutzki A, Walther P, Baumann B. 2012. Nuclear Factor κB Activation Impairs
Ependymal Ciliogenesis and Links Neuroinflammation to Hydrocephalus Formation. *J*
Neurosci **32**:11511–11523. doi:10.1523/JNEUROSCI.0182-12.2012

Lessieur EM, Song P, Nivar GC, Piccillo EM, Fogerty J, Rozic R, Perkins BD. 2019. Ciliary
genes arl13b, ahi1 and cc2d2a differentially modify expression of visual acuity phenotypes but
do not enhance retinal degeneration due to mutation of cep290 in zebrafish. *PLoS One*
14:e0213960. doi:10.1371/journal.pone.0213960

Lu H, Shagirova A, Goggi JL, Yeo HL, Roy S. 2020. Reissner fibre-induced urotensin
signalling from cerebrospinal fluid-contacting neurons prevents scoliosis of the vertebrate
spine. *Biology Open* **9**:bio052027. doi:10.1242/bio.052027

Lyons DA, Talbot WS. 2015. Glial Cell Development and Function in Zebrafish. *Cold Spring*
Harb Perspect Biol **7**:a020586. doi:10.1101/cshperspect.a020586

Mahuzier A, Gaudé H-M, Grampa V, Anselme I, Silbermann F, Leroux-Berger M, Delacour
D, Ezan J, Montcouquiol M, Saunier S, Schneider-Maunoury S, Vesque C. 2012. Dishevelled
stabilization by the ciliopathy protein Rpgrip1l is essential for planar cell polarity. *Journal of*
Cell Biology **198**:927–940. doi:10.1083/jcb.201111009

Mangos S, Lam P, Zhao A, Liu Y, Mudumana S, Vasilyev A, Liu A, Drummond IA. 2010. The
ADPKD genes *pkd1a/b* and *pkd2* regulate extracellular matrix formation. *Disease Models &*
Mechanisms **3**:354–365. doi:10.1242/dmm.003194

Masek M, Etard C, Hofmann C, Hülsmeier AJ, Zang J, Takamiya M, Gesemann M, Neuhauss
SCF, Hornemann T, Strähle U, Bachmann-Gagescu R. 2022. Loss of the Bardet-Biedl protein
Bbs1 alters photoreceptor outer segment protein and lipid composition. *Nat Commun* **13**:1282.
doi:10.1038/s41467-022-28982-6

Matusova Z, Hol EM, Pekny M, Kubista M, Valihrach L. 2023. Reactive astrogliosis in the era
of single-cell transcriptomics. *Front Cell Neurosci* **17**:1173200.
doi:10.3389/fncel.2023.1173200

Medina M, Avila J. 2010. Glycogen Synthase Kinase-3 (GSK-3) Inhibitors for the Treatment
of Alzheimers Disease. *CPD* **16**:2790–2798. doi:10.2174/138161210793176581

Meyer-Miner A, Van Gennip JLM, Henke K, Harris MP, Ciruna B. 2022. Resolving primary
pathomechanisms driving idiopathic-like spinal curvature using a new katnb1 scoliosis model.

iScience **25**:105028. doi:10.1016/j.isci.2022.105028

Mines MA, Beurel E, Jope RS. 2011. Regulation of Cell Survival Mechanisms in Alzheimer's Disease by Glycogen Synthase Kinase-3. *International Journal of Alzheimer's Disease* **2011**:1–11. doi:10.4061/2011/861072

Mukherjee I, Roy S, Chakrabarti S. 2019. Identification of Important Effector Proteins in the FOXJ1 Transcriptional Network Associated With Ciliogenesis and Ciliary Function. *Front Genet* **10**:23. doi:10.3389/fgene.2019.00023

Olstad EW, Ringers C, Hansen JN, Wens A, Brandt C, Wachten D, Yaksi E, Jurisch-Yaksi N. 2019. Ciliary Beating Compartmentalizes Cerebrospinal Fluid Flow in the Brain and Regulates Ventricular Development. *Current Biology* **29**:229-241.e6. doi:10.1016/j.cub.2018.11.059

Parmentier C, Hameury E, Dubessy C, Quan FB, Habert D, Calas A, Vaudry H, Lihrmann I, Tostivint H. 2011. Occurrence of Two Distinct Urotensin II-Related Peptides in Zebrafish Provides New Insight into the Evolutionary History of the Urotensin II Gene Family. *Endocrinology* **152**:2330–2341. doi:10.1210/en.2010-1500

Patten SA, Margaritte-Jeannin P, Bernard J-C, Alix E, Labalme A, Besson A, Girard SL, Fendri K, Fraisse N, Biot B, Poizat C, Campan-Fournier A, Abelin-Genevois K, Cunin V, Zaouter C, Liao M, Lamy R, Lesca G, Menassa R, Marcaillou C, Letexier M, Sanlaville D, Berard J, Rouleau GA, Clerget-Darpoux F, Drapeau P, Moldovan F, Edery P. 2015. Functional variants of POC5 identified in patients with idiopathic scoliosis. *J Clin Invest* **125**:1124–1128. doi:10.1172/JCI77262

Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S, Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A, Vizcaíno JA. 2022. The PRIDE database resources in 2022: a hub for mass spectrometry-based proteomics evidences. *Nucleic Acids Research* **50**:D543–D552. doi:10.1093/nar/gkab1038

Prendergast AE, Jim KK, Marnas H, Desban L, Quan FB, Djenoune L, Laghi V, Hocquemiller A, Lunsford ET, Roussel J, Keiser L, Lejeune F-X, Dhanasekar M, Bardet P-L, Levraud J-P, Van De Beek D, Vandenbroucke-Grauls CMJE, Wyart C. 2023. CSF-contacting neurons respond to Streptococcus pneumoniae and promote host survival during central nervous system infection. *Current Biology* **33**:940-956.e10. doi:10.1016/j.cub.2023.01.039

Quan FB, Dubessy C, Galant S, Kenigfest NB, Djenoune L, Leprince J, Wyart C, Lihrmann I, Tostivint H. 2015. Comparative Distribution and In Vitro Activities of the Urotensin II-Related Peptides URP1 and URP2 in Zebrafish: Evidence for Their Colocalization in Spinal Cerebrospinal Fluid-Contacting Neurons. *PLoS ONE* **10**:e0119290.

doi:10.1371/journal.pone.0119290

Quigley IK, Kintner C. 2017. Rfx2 Stabilizes Foxj1 Binding at Chromatin Loops to Enable
Multiciliated Cell Gene Expression. *PLoS Genet* **13**:e1006538.
doi:10.1371/journal.pgen.1006538

Rentero C, Blanco-Muñoz P, Meneses-Salas E, Grewal T, Enrich C. 2018. Annexins—
Coordinators of Cholesterol Homeostasis in Endocytic Pathways. *IJMS* **19**:1444.
doi:10.3390/ijms19051444

Ribeiro A, Monteiro JF, Certal AC, Cristóvão AM, Saúde L. 2017. Foxj1a is expressed in
ependymal precursors, controls central canal position and is activated in new ependymal cells
during regeneration in zebrafish. *Open Biol* **7**:170139. doi:10.1098/rsob.170139

Rose CD, Pompili D, Henke K, Van Gennip JLM, Meyer-Miner A, Rana R, Gobron S, Harris
MP, Nitz M, Ciruna B. 2020. SCO-Spondin Defects and Neuroinflammation Are Conserved
Mechanisms Driving Spinal Deformity across Genetic Models of Idiopathic Scoliosis. *Current*
Biology **30**:2363-2373.e6. doi:10.1016/j.cub.2020.04.020

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak
P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*
9:676–682. doi:10.1038/nmeth.2019

Shan Y, Fang C, Cheng C, Wang Y, Peng J, Fang W. 2015. Immersion infection of germ-free
zebrafish with Listeria monocytogenes induces transient expression of innate immune response
genes. *Front Microbiol* **6**. doi:10.3389/fmicb.2015.00373

Sherman J, Barkovich A, Citrin C. 1987. The MR appearance of syringomyelia: new
observations. *American Journal of Roentgenology* **148**:381–391. doi:10.2214/ajr.148.2.381

Sofroniew MV. 2015. Astrogliosis. *Cold Spring Harb Perspect Biol* **7**:a020420.
doi:10.1101/cshperspect.a020420

Song P, Fogerty J, Cianciolo LT, Stupay R, Perkins BD. 2020. Cone Photoreceptor
Degeneration and Neuroinflammation in the Zebrafish Bardet-Biedl Syndrome 2 (bbs2) Mutant
Does Not Lead to Retinal Regeneration. *Front Cell Dev Biol* **8**:578528.
doi:10.3389/fcell.2020.578528

Sullivan-Brown J, Schottenfeld J, Okabe N, Hostetter CL, Serluca FC, Thiberge SY, Burdine
RD. 2008. Zebrafish mutations affecting cilia motility share similar cystic phenotypes and
suggest a mechanism of cyst formation that differs from pkd2 morphants. *Developmental*
Biology **314**:261–275. doi:10.1016/j.ydbio.2007.11.025

Troutwine BR, Gontarz P, Konjikusic MJ, Minowa R, Monstad-Rios A, Sepich DS, Kwon RY, Solnica-Krezel L, Gray RS. 2020. The Reissner Fiber Is Highly Dynamic In Vivo and Controls Morphogenesis of the Spine. *Current Biology* **30**:2353-2362.e3. doi:10.1016/j.cub.2020.04.015

Tsujikawa M, Malicki J. 2004. Intraflagellar Transport Genes Are Essential for Differentiation and Survival of Vertebrate Sensory Neurons. *Neuron* **42**:703–716. doi:10.1016/S0896-6273(04)00268-5

Van Gennip JLM, Boswell CW, Ciruna B. 2018. Neuroinflammatory signals drive spinal curve formation in zebrafish models of idiopathic scoliosis. *Sci Adv* **4**:eaav1781. doi:10.1126/sciadv.aav1781

van Rooijen E, Giles RH, Voest EE, van Rooijen C, Schulte-Merker S, van Eeden FJ. 2008. LRRC50, a conserved ciliary protein implicated in polycystic kidney disease. *J Am Soc Nephrol* **19**:1128–1138. doi:10.1681/ASN.2007080917

Viau A, Bienaimé F, Lukas K, Todkar AP, Knoll M, Yakulov TA, Hofherr A, Kretz O, Helmstädter M, Reichardt W, Braeg S, Aschman T, Merkle A, Pfeifer D, Dumit VI, Gubler M, Nitschke R, Huber TB, Terzi F, Dengjel J, Grahammer F, Köttgen M, Busch H, Boerries M, Walz G, Triantafyllopoulou A, Kuehn EW. 2018. Cilia-localized LKB 1 regulates chemokine signaling, macrophage recruitment, and tissue homeostasis in the kidney. *The EMBO Journal* **37**:e98615. doi:10.15252/embj.201798615

Vierkotten J, Dildrop R, Peters T, Wang B, Rüther U. 2007. Ftm is a novel basal body protein of cilia involved in Shh signalling. *Development* **134**:2569–2577. doi:10.1242/dev.003715

Wang J, Thomas HR, Thompson RG, Waldrep SC, Fogerty J, Song P, Li Z, Ma Y, Santra P, Hoover JD, Yeo NC, Drummond IA, Yoder BK, Amack JD, Perkins B, Parant JM. 2022. Variable phenotypes and penetrance between and within different zebrafish ciliary transition zone mutants. *Disease Models & Mechanisms* **15**:dmm049568. doi:10.1242/dmm.049568

Wieczorek S, Combes F, Lazar C, Giai Gianetto Q, Gatto L, Dorffer A, Hesse A-M, Couté Y, Ferro M, Bruley C, Burger T. 2017. DAPAR & ProStaR: software to perform statistical analyses in quantitative discovery proteomics. *Bioinformatics* **33**:135–136. doi:10.1093/bioinformatics/btw580

Wiegering A, Rüther U, Gerhardt C. 2018. The ciliary protein Rpgrip1l in development and disease. *Developmental Biology* **442**:60–68. doi:10.1016/j.ydbio.2018.07.024

Xie H, Kang Y, Liu J, Huang M, Dai Z, Shi J, Wang S, Li L, Li Y, Zheng P, Sun Y, Han Q, Zhang J, Zhu Z, Xu L, Yelick PC, Cao M, Zhao C. 2023. Ependymal polarity defects coupled with disorganized ciliary beating drive abnormal cerebrospinal fluid flow and spine curvature

in zebrafish. *PLoS Biol* **21**:e3002008. doi:10.1371/journal.pbio.3002008

Yu X, Ng CP, Habacher H, Roy S. 2008. Foxj1 transcription factors are master regulators of

the motile ciliogenic program. *Nat Genet* **40**:1445–1453. doi:10.1038/ng.263

Zamanian JL, Xu L, Foo LC, Nouri N, Zhou L, Giffard RG, Barres BA. 2012. Genomic

Analysis of Reactive Astrogliosis. *J Neurosci* **32**:6391–6410. doi:10.1523/JNEUROSCI.6221-

11.2012

Zhang X, Jia S, Chen Z, Chong YL, Xie H, Feng D, Wu X, Song DZ, Roy S, Zhao C. 2018.

Cilia-driven cerebrospinal fluid flow directs expression of urotensin neuropeptides to straighten

the vertebrate body axis. *Nat Genet* **50**:1666–1673. doi:10.1038/s41588-018-0260-3

Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda

SK. 2019. Metascape provides a biologist-oriented resource for the analysis of systems-level

datasets. *Nat Commun* **10**:1523. doi:10.1038/s41467-019-09234-6

 MATERIAL AND METHODS

Zebrafish

Wild-type, *rpgrip1l^ex4* and *rpgrip1l^Δ* zebrafish embryos and adults were raised, staged

and maintained as previously described (Kimmel et al., 1995). All our experiments

were made in agreement with the european Directive 210/63/EU on the protection of

animals used for scientific purposes, and the French application decree 'Décret 2013-

118'. The projects of our group have been approved by our local ethical committee

'Comité d'éthique Charles Darwin'. The authorization number is APAFIS #31540-

2021051809258003 v4. The fish facility has been approved by the French 'Service for

animal protection and health' with approval number A750525. All experiments were

performed on Danio rerio embryos of mixed AB/TL background. Animals were raised

at 28.5°C under a 14/10 light/dark cycle.

Rpgrip1l mutant generation and genotyping
945
<u>Guide RNA preparation and microinjection</u>

CRISPR target sites were selected for their high predicted specificity and efficiency

using the CRISPOR online tool (Haeussler et al., 2016). Real efficiency was assessed

on zebrafish embryos by T7E1 test. The two most efficient guides (Rpgrip1l_x4_G1:

GCTTACGGTCCTTCACCAGACGG   and   Rpgrip1l_x25_G3:

CCTCAGTTGACAGGTTTCAGCGG) respectively situated 24 nt from the beginning of

exon 4 and 82 nt downstream of exon 25 were kept for further experiments.

sgRNA transcription templates were obtained by PCR using T7_Rpgrip1l-x4_G1_Fw

primer   (5′-

GAAATTAATACGACTCACTATAGGCTTACGGTCCTTCACCAGAGTTTTAGAGCTA

GAAATAGC-   3′)   or   T7_Rpgrip1l-x25_G3_Fw   (5′-

GAAATTAATACGACTCACTATAGGCCTCAGTTGACAGGTTTCAGGTTTTAGAGCT

AGAAATAG C-3′) as forward primer and sgRNA_R universal primer (5′-

AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTT

AACTTGCTA TTTCTAGCTCTAAAAC-3′) as reverse primer. sgRNAs were

transcribed using Megascript T7 Transcription Kit (Thermo Fisher Scientific, Waltham,

MA) and purified using NucleoSpin® RNA Clean up XS kit (Macherey Nagel, Düren,

Germany). sgRNA:Cas9 RNP complex was obtain by incubating Cas9 protein (gift of

TACGENE, Paris, France) (7.5 µM) with sgRNA (10 µM) in 20 mM Hepes-NaOH pH

7.5, 150 mM KCl for 10 min at 28 °C. 1–2 nl was injected per embryo. For deletion,

Rpgrip1l_x4_G1 and Rpgrip1l_x25_G3 RNP complexes were mixed half and half.

We kept two F1 fish, one with a 15 Kb deletion between exon 4 and 25 of *rpgrip1l*, that

we called *rpgrip1l*$^\Delta$ and that we submitted to ZFIN under the name *rpgrip1l* ^{bps1}and a

second one with a stop codon within exon 4 called *rpgrip1l* ^{ex4}, that is submitted to ZFIN

under the name *rpgrip1l* ^{bps2}.

<u>Screening and genotyping</u>

Injected (F0) fish were screened for germline transmission by crossing with wild type

fish and extracting genomic DNA from resulting embryos. For genomic DNA extraction,

caudal fin (juveniles/adults) or whole embryo DNA were used. Genomic DNA was

isolated with Proteinase K (PK) digestion in 40 of lysis buffer (100 mM Tris-HCl pH 7.5,

1 mM EDTA, 250 mM NaCl, 0.2% SDS, 0.1 µg/µl Proteinase K (PK)) for embryos (300

µl for adult fin) overnight at 37°C with agitation. PK was inactivated for 10 min at 90°C

and a five-fold dilution was used as template for PCR amplification. Combined

genotyping of wild type and mutant alleles (Supplementary Figure S1A') was

performed by PCR using 3 primers, a common reverse primer for both alleles :

Rpgrip1l_ex25_R3 (GTTGTGTCTCTGCCATATATTG), a specific forward primer for

the deleted allele: Rpgrip1l-ex4-del (CCCACACTGCATACGCACTC) and a specific

forward primer for the wild type allele : Rpgrip1l_ex25F

(AGTGTGCGGTACATCTCCAA) at an annealing temperature of 60°C for 35 cycles.

Expected size of the amplified fragment is 364 nt for the wild type allele and 544nt for

the mutant allele. The deleted allele presents a frameshift at position N77 leading to

the generation of a STOP codon after 11 aas, thus leading to a N-terminally truncated

protein at position 88/1256 aas.

For genotyping the *urp2* mutant line, caudal fins from adult fish were extracted in 300µl

of lysis solution and PCR was performed as above to detect exon 5 deletion using the

following primers: urp2-F1 (TGATTACTAGCCCTGTCCCAAC) and urp2-R1

(AGGTACAGTACACACGTCACAG).

µCT scans

The samples were scanned on a Bruker micro scanner (Skyscan 1272) with a
resolution of 8.5 μm, a rotation step of 0.55° and a total rotation of 180°. For the
acquisition of adult fish (2.5 cm), a 0.25 mm aluminium filter was used, for a voltage of
50 kV and an intensity of 180 mA, for juvenile fish (0.9 to 1.2 cm) the filter was omitted,
and a voltage of 60 kV was used with an intensity of 166 mA. Each image contained
1008 x 672 pixels and was based on the average of 3 images. The 3D reconstruction
by backprojection was carried out by the NRecon software and the 2D image overlays
were then cleaned by the CT Analyser software. The Dataviewer software allowed all
fish skeletons to be oriented in the same way taking the otoliths as landmarks. The
CTvox software then allowed 3D visualizing of the samples. Morphometric analysis
was performed with the CT Analyser software.
1007
Scanning electron microscopy
3 month fish (3 controls, 5 mutants) were euthanized using lethal concentration of
MS222 (0.028 mg/mL). The brains were quickly dissected in 1.22 X PBS (pH 7.4), 0.1
M sodium cacodylate and fixed overnight with 2% glutaraldehyde in 0.61 X PBS (pH
7.4), 0.1 M sodium cacodylate at 4°C. They were sectioned along the dorsal midline
with a razor blade to expose their ventricular surfaces. Both halves were washed four
times in 1.22 X PBS and post-fixed for 15 minutes in 1.22 X PBS containing 1% OsO_4.
Fixed samples were washed four times in ultrapure water, dehydrated with a graded
series of ethanol and critical point dried (CPD 300, Leica) at 79 bar and 38 °C with
liquid CO_2 as the transition fluid and then depressurized slowly (0,025 bar/s). They
were then mounted on aluminum mounts with conductive silver cement. Sample
surfaces were coated with a 5 nm platinum layer using a sputtering device (ACE 600,
Leica). Samples were observed under high vacuum conditions using a Field Emission
Scanning Electron Microscope (Gemini 500, Zeiss) operating at 5 kV, with a 20 μm
objective aperture diameter and a working distance around 3 mm. Secondary electrons
were collected with an in-lens detector. Scan speed and line compensation integrations
were adjusted during observation.
1025
Histological sections of juvenile and immunofluorescence on sections
Juvenile and adult zebrafish were euthanized using lethal concentration of MS222
(0.028 mg/mL). Pictures and size measurements were systematically taken before

fixation. Fish were fixed in Zamboni fixative [35 ml PFA 4 %, 7.5 ml saturated picric

acid (1.2 %), 7.5 ml 0.2M Phosphate Buffer (PB)] [55] overnight at 4°C under agitation.

Fish were washed with Ethanol 70% and processed for dehydration by successive 1 h

incubation in Ethanol (3 x 70% and 2 x 100%) at room temperature under agitation,

then for paraffin inclusion. 14 µm sagittal paraffin sections were obtained using a Leica

RM2125RT microtome. Sections were deparaffinized and antigen retrieval was

performed by incubation for 7 min in boiling citrate buffer (10 mM, pH 6).

Immunofluorescence staining was performed as described previously (Andreu-

Cervera et al., 2019). The following primary antibodies were used: anti-RF, anti-

Acetylated Tubulin, anti-Glutamylated Tubulin, anti-Arl13b, anti-LCP1, anti-AnnexinA2.

Corresponding primary and secondary antibodies are described and referenced in the

Resources Table.

1041

Immunofluorescence on whole embryos

Embryos from 24 to 40 hpf were fixed 4 hr to overnight in 4% paraformaldehyde (PFA)

at 4°C. For Reissner fiber staining, larvae at 48 and 72 hpf were fixed 2 hr in 4% PFA

and 3% sucrose at 4°C, skin from the rostral trunk and yolk were removed. Samples

from 24 to 40 hpf embryos were blocked overnight in a solution containing 0.5% Triton,

1% DMSO, 10% normal goat serum and 2 mg/mL BSA. For older samples (48 to 72

hpf) triton concentration was increased to 0.7%. Primary antibodies were incubated

one to two nights at 4°C in a buffer containing 0.5% Triton, 1% DMSO, 1% NGS and

1 mg/mL BSA. All secondary antibodies were from Molecular Probes, used at 1:400 in

blocking buffer, and incubated a minimum of 2.5 hr at room temperature. The primary

antibodies chosen for in toto immuno-labeling against Reissner fiber, Acetylated-

tubulin, Myc, Arl13b and Gamma-tubulin as well as the corresponding secondary

antibodies are referenced in the Key Resources Table. Whole mount zebrafish

embryos (dorsal or lateral mounting in Vectashield Mounting Medium) were imaged on

a Leica SP5 confocal microscope and Zeiss LSM910 confocal microscope, both

equipped with a 63X immersion objective. Images were then processed using Fiji

(Schindelin et al., 2012).

1059

Whole-mount brain clearing

Brains were dissected from 4-5 wpf size-matched (0.9-1.2 mm length) zebrafish after

in toto fixation with formaldehyde. Whole–mount tissue clearing was performed

following the zPACT protocol (Affaticati et al., 2017). In brief, brains were infused for 2

days in hydrogel monomer solution (4% acrylamide, 0.25% VA- 044, 1% formaldehyde

and 5% DMSO in 1X PBS) at 4°C. Polymerization was carried out for 2h30 at 37°C in

a desiccation chamber filled with pure nitrogen. Brains were transferred into histology

cassettes and incubated in clearing solution (8% SDS and 200 mM boric acid in dH2O)

at 37°C with gentle agitation for 8 days. Cleared brains were washed in 1X PBS with

0.1% Tween-20 (PBT) for 3 days at room temperature and kept in 0.5% formaldehyde,

0.05% sodium azide in PBT at 4°C until further processing. Brains were subsequently

placed for 1h in depigmentation pre-incubation solution (0.5X SSC, 0.1% Tween-20 in

dH2O) at room temperature. The solution was replaced by depigmentation solution

(0.5X SSC, Triton X-100 0.5%, formamide 0.05% and H2O2 0.03% in dH2O) for 45

minutes at room temperature. Depigmented brains were washed for 4h in PBT and

post-fixed (2% formaldehyde and 2% DMSO in PBT) overnight at 4°C.

1076

1077 *Cleared brain immunostaining of whole adult brains*

Whole-mount immunolabeling of cleared brains was performed as described in the

zPACT protocol with slight modifications. Briefly, brains were washed in PBT for one

day at room temperature and blocked for 10h in 10% normal goat serum, 10% DMSO,

5% PBS-glycine 1M and 0.5% Triton X-100 in PBT at room temperature. Brains were

washed again in PBT for 1h prior to incubation with anti- ZO-1 antibody (ZO1-1A12,

Thermofisher, 1:150) in staining solution (2% normal goat serum, 10% DMSO, 0.1%

Tween-20, 0.1% Triton X-100 and 0.05% sodium azide in PBT) for 12 days at room

temperature under gentle agitation. Primary antibody was renewed once after 6 days

of incubation. Samples were washed three times in PBT and thereafter incubated with

Alexa Fluor 488-conjugated secondary antibody (A11001, Invitrogen, 1:200) for 10

days in the staining solution at room temperature under gentle agitation. Secondary

antibody was renewed once after 5 days of incubation. Samples were washed three

times in PBT prior to a counterstaining with DiIC18 (D282, Invitrogen, 1µM) in the

staining solution for three days at room temperature with gentle agitation. Samples

were washed in PBT for 3h before mounting procedure.

1093

Mounting and confocal imaging

Samples were placed in 50% fructose-based high-refractive index solution (fbHRI, see (Affaticati et al., 2017) /50% PBT for 1h, then in 100% fbHRI. Brains were mounted in agarose-coated (1% in standard embryo medium) 60 mm Petri dish with custom imprinted niches to help orientation. Niche-fitted brains were embedded in 1% phytagel and the Petri dish filled with 100% fbHRI. fbHRI was changed three times before imaging until its refractive index matched 1.457. Images were acquired with a Leica TCS SP8 laser scanning confocal microscope equipped with a Leica HC FLUOTAR L 25x/1.00 IMM motCorr objective. Brains were scanned at a resolution of 1.74x1.74x1.74 µm (xyz) and tiled into 45 to 70 individual image stacks, depending on brain dimensions, subsequently stitched, using LAS X software.

Volumetric analysis of the posterior ventricles

The volumes of the posterior ventricles were segmented, reconstructed and analyzed using Amira for Life & Biomedical Sciences software (Thermo Fisher Scientific). In essence the ventricles volumes were manually segmented in Amira's segmentation editor and subsequently refined by local thresholding and simplification of the corresponding surfaces. Volumes, which were open to the environment were artificially closed with minimal surfaces by connecting the distal-most points of their surface to the contralaterally corresponding points using straight edges. Due to the biologic variability of the sample population the overall size of the specimens needed to be normalized to keep the measured volumes comparable. For this spatial normalization one of the specimens was randomly selected from the wildtype population as template (1664, grey) and the 'Registration' module in Amira was used to compute region-specific rigid registrations for the other specimens, allowing for isotropic scaling only [details in supp. mat.]. For excluding the influence of the ventricular volumes, the registration was computed on the basis of the independent reference stain (DiIC18) (details in supp. mat.). Region specific volume differences between the mutant and wildtype population were evaluated on seven subvolumes of the posterior ventricles (details in supp. mat.).

RNA extraction for transcriptome analysis and quantitative RT-PCR

Juvenile and adult zebrafish were euthanized using lethal concentration of MS222 (0.28 mg/mL). Their length was measured and their fin cut-off for genotyping. For transcriptomic analysis, brain and dorsal trunk from 1 month juvenile (0.9 to 1.0 cm) zebrafish were dissected in cold PBS with forceps and lysed in QIAzol (QIAGEN) after homogenization with plunger pistons and 1ml syringes. Samples were either stored in QIAzol at -80°C or immediately processed. Extracts containing RNA were loaded onto QIAGEN-mini columns, DNAse digested and purified in the miRNAeasy QIAGEN kit (Cat 217004) protocol. Samples were stored at -80°C until use. RNA concentration and size profile were obtained on the Tapestation of ICM platform. All preparations had a RIN above 9.2. For Q-PCR from juveniles, whole fish minus internal organs were lysed in Trizol (Life technologies) using the Manufacturer protocol.

Quantitative PCR from individual juvenile or adult fish

The cDNA from the isolated RNAs was obtained following GoScript Reverse Transcription System protocol (Promega), using 3 to 4 µg of total RNA for each sample. The 20 µL RT-reaction was diluted 4-fold and 4µl was used for each amplification performed in duplicates. The primers used for qPCR were the following: urp2 (F: ACCAGAGGAAACAGCAATGGAC; R: TGAGGTTTCCATCCGTCACTAC), *urp1* (F: ACATTCTGGCTGTGGTTTGTTC; R: CTCTTTTGCACCTCTCTGAAGC), *urp* (F: GCGGAAAAATGTCATGCCTCTTC; R: TTGAGCTCCTTTTGAAGCTCCTG), *uts2a* (F: CACTGCTCAACAGAGACAGTATCA; R: CCAAAAGACCACTGGGAGGAAC), *uts2b* (F: TACCCGTCTCTCATCAGTGGAG; R: TTTTCCAGCAAGGCCTCTTTTAC) where all of them are from Gaillard et al. 2023, *rpgrip1l* (F: CAGACACCTGCTGGAGTTACA; R: TCCTGACTCACATCAAACGCA), *irg1l* (F: TCCCTAAGAGGTTCCATCCTCC; R: AAGATGCAGCGATGGCCAAA), *stat3* (F: CAGGACTGGGCGTATGCG; R: GAAGCGGCTGTACTGCTGAT), *c4b* (F: GGGTGTTCTTATGGCGGTGG and R: GCGCACAACAAGCTTTTCTCATC) *lsm12b* (F: GAGACTCCTCCTCCTCTAGCAT and R: GATTGCATAGGCTTGGGACAAC). The Q-PCR were performed using the StepOne real-time PCR system and following its standard amplification protocol (Thermo Scientific). Relative gene expression levels were quantified using the comparative CT method ($2^{-\Delta\Delta CT}$ method or $2^{-\Delta CT}$) on the basis of CT values for target genes and *lsm12b* as internal control.

 RNA-sequencing and analysis

RNA-seq libraries were constructed by the ICM Platform (PARIS) from 100 ng of total

RNA using the KAPA mRNA Hyperprep kit (Roche) that allows to obtain stranded

polyA libraries, and their quality controlled on Agilent Tape station. The sequencing

was performed on a NovaSeq 6000 Illumina for both strands. Sequences were aligned

against the GRCz11 version of zebrafish genome using **RNA Star function** in "Galaxy"

environment. The 12 Brain libraries reached between 22 to 26 Millions of assigned

reads, and the trunk libraries between 26 to 35 Millions of assigned reads. Expression

level for each gene was determined using **Feature counts function** from RNA Star

BAM files. **DESeq2 function** was used to calculate the log2 Fold change and adjusted

P values from the Wald test for each gene comparing the 7 mutants raw count

collection to the 5 controls raw counts collection. Volcano plots were generated using

the Log2 fold change and P adj. value of each Gene ID from the DESeq2 table. A

threshold of 5.10 E-2 was chosen for the Padj. values and a log2 fold change > to

+0.75 or < to -0.75 to select significantly up or down-regulated regulated genes. Both

gene lists were used as input to search for enriched GO terms and KEGGs pathways

using Metascape software (Zhou et al., 2019) as well as to compare up-regulated

genes in the dorsal trunk versus the brain using Galaxy.

1177

Quantitative proteomics

Sample preparation: Adult zebrafish were euthanized using lethal concentration of

MS222 (0.28 mg/mL). For proteomic analysis, brain from 3 months zebrafish were

dissected in cold PBS with forceps and immediately flash frozen in liquid nitrogen and

stored at -80°C. Zebrafish brain samples were lysed in RIPA buffer (Sigma) with 1/100

antiprotease and sonicated for 5 minutes. After a 660 nm protein assay (Thermo),

samples were digested using a Single Pot Solid Phase enhanced Sample Preparation

(SP3) protocol (Hughes et al., 2014). In brief, 40 μg of each protein extract was reduced

with 12mM dithiothreitol (DTT) and alkylated using 40 mM iodoacetamic acid (IAM). A

mixture of hydrophilic and hydrophobic magnetic beads was used to clean-up the

proteins at a ratio of 20:1 beads:proteins (Sera-Mag Speed beads, Fisher Scientific).

After addition of ACN to a final concentration of 50%, the beads were allowed to bind

to the proteins for 18 minutes. Protein-bead mixtures were washed twice with 80%

EtOH and once with 100% ACN. The protein-bead complexes were digested with a

mixture of trypsin:LysC (Promega) at a 1:20 ratio overnight at 37°C. Extracted peptides

were cleaned-up using automated C18 solid phase extraction on the Bravo AssayMAP

platform (Agilent Technologies).

1195

NanoLC-MS-MS analysis

The peptide extracts were analysed on a nanoLC-TimsTof Pro coupling (Bruker

Daltonics). The peptides (200ng) were separated on an IonOpticks column (25cm X

75µm 1.6µm C18 resin) using a gradient of 2-37% B (2%ACN, 0.1%FA) in 100 minutes

at a flow rate of 0.3 µl/min. The dual TIMS had a ramp time and accumulation time of

166 ms resulting in a total cycle time of 1.89s. Data was acquired in Data Dependent

Acquisition-Parallel Accumulation Serial Fragmentation (DDA-PASEF) mode with 10

PASEF scans in a mass range from 100 m/z to 1700 m/z. The ion mobility scan range

was fixed from 0.7-1.25 Vs/cm^2. All samples were injected using a randomized injection

sequence. To minimize carry-over, one solvent blank injection was performed after

each sample.

1207

Data interpretation

Raw files were converted to .mgf peaklists using DataAnalysis (version 5.3, Bruker

Daltonics) and were submitted to Mascot database searches (version 2.5.1,

MatrixScience, London, UK) against a *Danio rerio* protein sequence database

downloaded from UniProtKB-SwissProt (2022_05_18, 61 732 entries, taxonomy ID:

7955), to which common contaminants and decoy sequences were added. Spectra

were searched with a mass tolerance of 15 ppm in MS mode and 0.05 Da in MS/MS

mode. One trypsin missed cleavage was tolerated. Carbamidomethylation of cysteine

residues was set as a fixed modification. Oxidation of methionine residues and

acetylation of proteins' n-termini were set as variable modifications. Identification

results were imported into the Proline software (version 2.1.2,

http://proline.profiproteomics.fr) (Bouyssié et al., 2020) for validation. Peptide

Spectrum Matches (PSM) with pretty ranks equal to one were retained. False

Discovery Rate (FDR) was then optimized to be below 1% at PSM level using Mascot

Adjusted E-value and below 1% at protein level using Mascot Mudpit score. For label

free quantification, peptide abundances were extracted without cross assignment

between the samples. Protein abundances were computed using the sum of the unique peptide abundances normalized at the peptide level using the median.

To be considered, proteins must be identified in at least four out of the five replicates in at least one condition. Imputation of the missing values and differential data analysis were performed using the open-source ProStaR software (Wieczorek et al., 2017). Imputation of missing values was done using the approximation of the lower limit of quantification by the 2.5% lower quantile of each replicate intensity distribution ("det quantile"). A Limma moderated *t*-test was applied on the dataset to perform differential analysis. The adaptive Benjamini-Hochberg procedure was applied to adjust the p-values and FDR. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol et al., 2022), with the dataset identifier PXD042283.

Embryos drug treatment

27 hpf embryos from *dnaaf1*[tm317b/+] incross were dechorionated manually. N-acetyl cysteine ethyl ester (NACET) (BIOLLA Chemicals #59587-09-6) was prepared extemporaneously at the final concentration of 3 mM in pure water (pH: 7.2; conductivity 650 µS). Embryos were treated between 27 hpf to 60 hpf and embryos were anesthetized before being imaged laterally.

Juvenile drug treatment and preparation of treated samples

Two hundred larvae were housed off-system in 1.8 liter tanks with 20 fish per tank, being fed twice a day throughout the experiment. 100 were treated with NACET (BIOLLA Chemicals #59587-09-6), which was prepared extemporaneously at 1.5 mM (286.5mg/L) in fish water (pH 7.2, Conductivity 650 µS) which was changed once per day. Fish were treated from 30 dpf to 84 dpf and monitored for curvature onset once per week. At 85 dpf, fish were euthanized, imaged to measure the curvature index. Half of the fish were fixed for histology analysis in zamboni fixative [35 ml PFA 4 %, 15 ml saturated picric acid (1.2 %)] overnight at 4°C under agitation, while the other half was processed for RT-qPCR analysis. To prepare dorsal trunk RNA, internal organs were removed in cold PBS 1X, and the remaining tissue was cut in small pieces before lysis in QIAzol (QIAGEN). Homogenization was achieved using plunger pistons and passages through 1ml syringes.

1257

Transgenesis

The *Tol2 -5.2foxj1a:5xmyc-RPGRIP1L* plasmid was produced using the Gateway
system by combining four plasmids. P5E-foxj1a enhancer was kindly donated by [3].
5xmyc-RPGRIP1L cDNA was amplified from the plasmid pCS2-5xmyc-RPGRIP1L
(Mahuzier et al., 2012) using CloneAmp HiFi PCR Premix (Takara # 639298) using
primers : forward 5'-
GGGGACAAGTTTGTACAAAAAGCAGGCTGCAGGATCCCATCGATTTAAAGCT-3'
and reverse 5'
GGGGACCACTTTGTACAAGAAAGCTGGGTCTCCGAGCTCGGTTCAAGCCTCCA
AGTCATCTCTGT-3. The PCR product was gel purified using QIAEX II gel extraction
kit (QIAGEN, #20021) . BP recombination (Invitrogren "Gateway BP Clonase II
Enzyme Mix" #11789-020) was performed into pDONR221 #218 to generate pMe-
5xmyc-RPGRIP1L. The 3' entry polyadenylation signal plasmid was the one described
in [61]. The 3 plasmids were shuttled into the backbone containing the cmcl2:GFP
selection cassette (Kwan et al., 2007) using the LR recombination (Invitrogen
"Gateway LR Clonase II Plus Enzyme Mix" #12538-120). *rpgrip1l*[+/-] X AB embryos
outcrosses were injected at the one cell stage with 1nl of a mix containing 20 ng/µl
plasmid and 25 ng/µl Tol2 transposase RNA and screened at 48 hpf for GFP
expression in the heart. Some F0 founders produced F1 zebrafish carrying one copy
of the transgene that were further checked for RPGRIP1L specific expression in *foxj1a*
territory, using Myc labelling at the base of FP cilia in 2,5 dpf embryos.

The Tol2-1.7kb col2a1a:5xMyc-RPGRIP1L plasmid was produced using the Gateway
System by combining four plasmids. Col2a1a enhancer was amplified from the plasmid
-1.7kbcol2a1a:EGFP-CAAX (Dale and Topczewski, 2011) using CloneAmp HiFi
premix (Takara #639298) with the primers: forward 5'-GGG GAC AAC TTT GTA TAG
AAA AGT TGG CCC TCT GAC ACC TGA TGC CAA TTG C-3' and reverse 5'-GGG
GAC TGC TTT TTT GTA CAA ACT TGC TTG CAG GTC CTA AGG GGT GAA AGT
CG-. The PCR product was gel purified and BP recombination was performed into
pDONR_P4-P1R-#219. The middle entry plasmid and the 3' plasmid were the same
as those used to generate the Tol-5.2foxj1a: 5xmyc-RPGRIP1L plasmid. F1 transgenic
were first selected for GFP expression in the heart and then on Myc labelling within
notochordal cells at 2,5 dpf within their embryonic progeny.

1290

Skeletal preparations, Alizarin staining and imaging

Animals were euthanized using a lethal concentration of MS222 (0.28 mg/mL), skin was removed and fixation was performed with 4% paraformaldehyde overnight at 4°C. After evisceration, samples were incubated in borax 5%, rinsed several times in 1% KOH and incubated in solution composed with Alizarin 0.01% (Sigma, A5533) and KOH 1% during 2 days. Fish were rinsed several times in KOH 1% and incubated in trypsin 1% and borax 2% for 2 days until cleared. Samples were rinsed several times in distilled water and transferred in glycerol 80% (in KOH 1%) using progressive dilutions. Samples were kept in glycerol 80% until imaging.

Cobb angles measurements

To quantitatively evaluate the severity of spine curvature in *rpgrip1l*$^{+/-}$*; urp2*$^{+/-}$ incrosses, we used the zebrafish skeletal preparations stained with Alizarin. We drew parallel lines to the top and bottom most displaced vertebrae on lateral and dorsal views. The Cobb angle was then measured as the angle of intersection between lines drawn perpendicular to the original 2 lines. This was conducted using FIJI software. For each fish, the total Cobb angle was calculated by summing all Cobb angles.

Curvature index measurements

We implemented a novel procedure to quantify curvatures on both axes by drawing a line along the body of the fish as shown in Figure supplementary 4I and curvature was calculated in MATLAB (code available on demand). We decomposed the line in a series of equidistant points and we measured the curvature at each of these points using the LineCurvature2D function. We then added the absolute values of these measures, which are expressed as an angle (in radian) per unit of length. The higher the sum is for a given line, the more this line is curvated. We verified the accurancy of this metric by comparing it with the visual assessment of curvatures by 3 independent observers. For *rpgrip1l*$^{+/-}$*; urp2*$^{+/-}$ incross and NACET experiment, a line following the body deformation was drawn from the mouth to the base of the tail following the midline of the fish in lateral position and another one along the dorsal axis. Curvature index of both curves were summed for each fish. Curvature analysis was performed blinded to fish genotype. *rpgri1l*$^{+/+}$ or *rpgrip1l*$^{+/-}$ siblings were used as controls.

1323

Data acquisition for body-curvature analysis at embryonic stage

1324

Zebrafish embryos were anesthetized and imaged laterally with the head pointing to

the left. The angle between the line connecting the center of the eye to the center of

the yolk and the line connecting the center of the yolk to the tip of the tail was measured

to evaluate the body curvature of the embryos. For quantitative analysis, the angles of

each embryo were put in the same concentric circles represented with a Roseplot, with

0° pointing to the right and 90° pointing to the top. Each triangle represents a 30°

quadrant, and its height indicates the number of embryos within the same quadrant.

1332

Quantification, statistical analysis and figure preparation

1333

1334 For all experiments the number of samples analyzed is indicated in the text and/or

1335 visible in the figures. Statistical analysis was performed using the Prism software. ****:

1336 P value < 0.0001; ***: P value < 0.001; **: P value < 0.01; *: P value < 0.05. Graph

1337 were made using Prism and Matlab and figures were assembled using Photoshop

1338 software.

1339

1340 **REAGENT AND RESOURCE TABLE**

Reagent type or resource	Designation	Source or reference	Identifiers	Additional Infos
Antibody (antiserum)	Rabbit polyclonal IgG anti-bovine Reissner Fiber	Didier et al 1995	Courtesy of Dr. S.Gobron	1:200
Antibody	Mouse monoclonal	Thermofisher	#33-9100	1:150

	IgG1k anti ZO1 -1A12			
Antibody	Mouse monoclonal IgG2b anti-acetylated-tubulin (clone 6-11B-1)	Sigma-Aldrich	#T 6793 RRID: AB_477585	1:400
Antibody	Mouse monoclonal IgG1 anti glutamylated Tubulin	Adipogen	#AG-20B-0020-C100	1:500
Antibody	Rabbit polyclonal anti-Arl13b	Proteintech	#17711-1-AP	1:200
Antibody	Mouse monoclonal IgG1 anti-GFAP	Sigma	#G3893	1:400
Antibody	Chicken monoclonal IgY anti-GFP	AVES lab	#GFP-1020	1:200
Antibody	Mouse monoclonal IgG2a anti-myc (clone 9b11)	Cell signaling	#2276	1:200

Antibody	Rabbit polyclonal anti-LCP1	GeneTex	#GTX124420	1:200
Antibody	Mouse monoclonal IgG1 anti-gamma-tubulin	Sigma	#T6557	1:500
Antibody	Mouse polyclonal anti-annexin A2	Proteintech	#11256-1-AP	1:200
Antibody	Goat anti-mouse IgG1 Alexa633	Molecular probes	# A-21126 RRID:AB_2535768	1:400
Antibody	Goat anti-mouse IgG1 Alexa568	Molecular Probes	# A-21124 RRID: AB_2535766	1:400
Antibody	Goat anti-mouse IgG2a Alexa568	Molecular probes	# A-21134 RRID:AB_2535773	1:400
Antibody	Goat anti-mouse IgG2a Alexa488	Molecular probes	# A-21131 RRID:AB_141618	1:400
Antibody	Goat anti-mouse IgG2b Alexa633	Molecular probes	# A-21146 RRID:AB_2535782	1:400

Antibody	Goat anti-mouse IgG2b Alexa568	Molecular probes	# A-21144 RRID: AB_2535780	1:400
Antibody	Goat anti-rabbit IgG Alexa568	Molecular probes	# A-11011 RRID:AB_143157	1:400
Antibody	Goat anti-rabbit IgG Alexa488	Molecular probes	# A-11008 RRID: AB_143165	1:400
Antibody	FITC donkey Anti-Chicken IgY	Jackson ImmunoResearch	# 703-096-155	1:200
Chemical	Vectashield	Vector Laboratories	H-1000	
Strain (Danio rerio)	zebrafish wild-type AB or (TL x AB) hybrid strains	N/A	N/A	
Strain (Danio rerio)	Zebrafish $rpgrip1l^{\Delta}$ mutant strain	This manuscript, Zfin:$rpgrip1l^{bps1}$		
Strain (Danio rerio)	Zebrafish $rpgrip1l^{ex4}$ mutant strain	This manuscript, Zfin:$rpgrip1l^{bps2}$		
Strain (Danio rerio)	Zebrafish $urp2^{+/-}$	(Gaillard et al., 2023)	ZDB-ALT-230207-5	

Strain (Danio rerio)	Zebrafish *dnaaf1[tm317b/+]*	(van Rooijen et al., 2008)		
Strain (Danio rerio)	Zebrafish Scospondin-GFP[ut24]	(Troutwine et al., 2020)		
Recomb. DNA	Foxj1 promoter-enhancer sequence	(Dale and Topczewski, 2011)		
Software	LASX	Leica		
Software	Amira for Life & Biomedical Sciences	Thermo Fisher Scientific		
Software	CRISPOR	(Haeussler et al., 2016)	http://crispor.tefor.net/	
Software	Fiji/ImageJ	(Schindelin et al., 2012)	https://imagej.net/Fiji/Downloads	
Software	PRISM	GraphPad	https://www.graphpad.com/	
Software	Metascape		https://metascape.org/gp/index.html#/main/step1	
Software	Matlab	The Mathworks, Inc.	https://fr.mathworks.com/products/matlab.html	
Software	NRecon reconstruction software	Micro Photonics Inc.		

Software	CTvox	Bruker		
Software	CT Analysis	Bruker		
Other				
Refractometer	Roth Sochiel EURL	DR 301-95		
Confocal microscope	Leica	SP5		
Confocal microscope	Zeiss	LSM 980 Inverted		
Confocal microscope	Zeiss	LSM 710		
Macroscope	Zeiss	AXIOZOOM V16		
Micro-scanner	Bruker	Skyscan 1272		
Microtome	Leica	RM2125RT		
Q-PCR apparatus	Applied Biosystems	Step One plus		

1341

1342

1343

1344

1345

1346

1347

1348

    FIGURE LEGENDS

Figure 1: *rpgrip1l^(-/-)* juvenile fish develop scoliosis, which is rescued by
RPGRIP1L expression in *foxj1a*-positive cells.

(A) Representative *rpgrip1l^(-/-)* fish at 5 days post-fertilization (dpf) (4 mm body length,
larvae), 5 weeks pf (wpf) (1.2 cm body length, juveniles) and 12 wpf (2.5 cm body
length, adults), showing absence of morphological defects in embryos, onset of spine
curvature (tail-up) in juveniles and scoliosis in adults. **(B-C)** Graph showing the time
course of scoliosis appearance **(B)** in *rpgrip1l^(-/+)* incrosses (total 4 clutches, 252 fish)
and scoliosis penetrance **(C)** in adults. **(D-E)** Micro-computed tomography (μCT) scans
of control siblings (D) and *rpgrip1l^(-/-)* (E) adult fish. 4 fish were analyzed for each
condition. **(F-G)** Dorsal and lateral superimposed μCT views of the spines of one
control (yellow) and two *rpgrip1l^(-/-)* (pink, cyan) adult fish illustrating the 3D spine
curvatures in mutants. **(H)** Schematic representation of the two rescue constructs. **(I)**
Table presenting the number and phenotype of *rpgrip1l^(-/-)* fish generated for both rescue
experiments by stable transgenesis *(J)* Graph representing scoliosis penetrance in
transgenic and non-transgenic *rpgrip1l^(-/-)* siblings.

1366

Figure 2 Altered ciliogenesis at trunk and subcommissural organ (SCO) levels.

(**A-C'''**) Immunostaining of trunk sagittal sections at the level of the spinal cord central
canal in *rpgrip1l^(+/+)* (A-A'''; n=5), straight *rpgrip1l^(-/-)* (B-B'''; n=4) and scoliotic *rpgrip1l^(-/-)*
fish (C-C'''; n=6)) at 8 weeks (N=2 independent experiments). Cilia were
immunostained with Arl13b (cyan, A'', B'', C'') and glutamylated-tubulin (magenta, A''',
B''', C'''). Panels A-C' are merged images with DAPI. Panels A', B' and C' show higher
magnification of the squared regions in A, B and C, respectively. Scale bars: 10 μm in
A, 5 μm in A'. **(D)** Violin plot showing the distribution of cilia length along the central
canal of *rpgrip1l^(+/+)* (n=300 cilia from 5 fish), straight *rpgrip1l^(-/-)* (n=267 from 4 fish) and
scoliotic *rpgrip1l^(-/-)* (n=414 from 6 fish) juvenile fish (N=2). **(E)** Violin plot showing the
distribution of cilia density along the central canal of juvenile *rpgrip1l^(+/+)* (n=5), straight
rpgrip1l^(-/-) (n=4) and scoliotic *rpgrip1l^(-/-)* fish (n=6, N=2). Density is the average number
of ventral and dorsal cilia lining the CC over a length of 100 μm from 3 to 5 sections for
each fish. Each dot represents the mean cilia length (D) or cilia density (E) of 4-8

sections analyzed at different antero-posterior levels. Statistical analysis was performed using Tukey's multiple comparisons test where * means P-value < 0.05, and ** means P-value < 0.01. **(F)** Violin plot showing the distribution of width of the central canal (nuclear-free region) (in μm) of juvenile *rpgrip1l*[+/+] (n=5), straight *rpgrip1l*[-/-] (n=4) and scoliotic *rpgrip1l*[-/-] fish (n=6, N=2). Each dot represents the mean of 3 measurements made at different AP levels. Statistical analysis was performed using Tukey's multiple comparisons test where ns means non-significant, * means P-value < 0.05. **(G-L)** Immunostaining of cilia on transverse sections of the brain at SCO level using glutamylated-tubulin and acetylated-tubulin antibodies (both in white), in *rpgrip1l*[+/+] (n=6) (G, G', H), straight *rpgrip1l*[-/-] (n=3) (I, I', J) and scoliotic *rpgrip1l*[-/-] (n=7) (K, K', L) 8 wpf fish (N=2). Nuclei are stained with DAPI (blue). G', I' and K' are higher magnifications of squared regions in G, I and K, respectively. In H, J and L the bottom-right inset represents a higher magnification of the squared region. Scale bars in G and G': 10 μm.

Figure 3: *rpgrip1* [-/-] juveniles show ventricular dilations and loss of the Reissner Fiber at scoliosis onset.

(A) Reconstruction of the RhV at the level of posterior midbrain and hindbrain in a transparised 5 wpf control zebrafish brain, stained with ZO1 antibody (ventricular surface) and DiI (global brain shape). Several brain regions are annotated on the reconstruction where Tel means telencephalon, ChP forebrain choroid plexus, SCO subcommissural organ, TeO optic tectum, CCe corpus cerebelli, CC crista cerebellaris and Rh rhombencephalon. **(B-D)** Cyan circles represent the ROIs measured in F. The orange line in D indicates the level of optical sections in E. **(E)** Optical transverse sections showing the caudal part of reconstructed ventricles of a control and two *rpgrip1l*[-/-] fish, one straight and two tail-up. **(F)** Graphs of the ventricle volume at the onset of scoliosis in 3 control and 4 *rpgrip1l*[-/-] (one straight and three tail-up) fish. The green dot corresponds to the straight mutant fish. Each dot represents a fish. Statistical analysis was performed using unpaired t-test where ns means non-significant, and * means P-value < 0.05. **(G)** Graph of the surface of the optical sections as illustrated in E. The green dot corresponds to a straight mutant fish. Each dot represents a fish. Statistical analysis was performed using unpaired t-test where ns means non-significant, and * means P-value < 0.05. **(H)** Schematic representation of a portion of

the spinal cord central canal in a lateral view of the fish trunk, showing the Reissner Fiber (RF), and of the SCO in a transverse view of the brain at the level of diencephalic Ventricle (DiV). **(I-Q)** Visualization of the GFP fluorescence in the *sspo-GFP^{uts24/+}* transgenic line (fire LUT) in sagittal sections of the trunk (I-K) (N=3) and transverse sections of the brain at the level of the SCO (N=2) (L-Q) in juvenile fish. The corresponding fish are 8 wpf *rpgrip1l^{+/+}* (I, L, O) (n=10); straight *rpgrip1l^{-/-}* (J, M, P) (n=9) and scoliotic *rpgrip1l^{-/-}* (K, N, Q) (n=8). L-N and O-Q show sections at anterior and posterior SCO levels, respectively. Scale bars: 5 µm in I-K, 10 µm in L-Q.

Figure 4: Upregulation of *urp1/2* expression does not contribute to axis curvature of *rpgrip1l^{-/-}* fish.

(A) Quantitative reverse transcription polymerase chain reaction (qRT-PCR) assaying *urp2* expression level in *rpgrip1l^{-/-}* and *rpgrip1l^{+/+}* (WT) siblings. Each dot represents the value obtained for one fish. Green and purple dots represent straight and scoliotic *rpgrip1l^{-/-}* fish, respectively. qRT-PCR analysis was performed at 5 wpf (juvenile stage, 5 controls and 7 *rpgrip1l^{-/-}*) and 12 wpf (adult stage, 5 controls and 9 *rpgrip1l^{-/-}*). *urp2* expression levels are increased by approximately 3.5-fold in *rpgrip1l^{-/-}* at both stages. Statistical analysis was performed using unpaired t-test where **P < 0.01. Error bars represent s.d. **(B)** Quantification of body axis curvature from dorsal and lateral views in 4 mpf fish issued from [*rpgrip1l^{+/-}*, *urp2^{+/-}*] incrosses. Each point represents the value measured for one fish. Statistical analysis was performed using Tukey's multiple comparisons test where **** P < 0,0001 with error bars represent s.d. n=9 [*rpgrip1l^{+/+}; urp2^{+/+}*]*, 7 [*rpgrip1l^{+/+}; urp2^{-/-}*], 27 [*rpgrip1l^{-/-}; urp2^{+/+}*], 48 [*rpgrip1l^{-/-}; urp2^{+/-}*] and 27 [*rpgrip1l^{-/-}; urp2^{-/-}*] fish. **(C)** Representative [*rpgrip1l^{-/-}; urp2^{+/+}*] and sibling [*rpgrip1l^{-/-}; urp2^{-/-}*] fish on lateral and dorsal views at 12 wpf. Scale bar 5mm. **(D, E)** Expression levels of *urp2* (D) and *urp1* (E) in adult fish (12 wpf). Each dot represents the value for one fish. Green and purple dots represent straight and scoliotic *rpgrip1l^{-/-}* fish, respectively (n=5 [*rpgrip1l^{+/+}; urp2^{+/+}*], 9 [*rpgrip1l^{+/+}; urp2^{-/-}*], 11 [*rpgrip1l^{-/-}; urp2^{+/-}*] and 10 [*rpgrip1l^{-/-}; urp2^{-/-}*]). Each gene expression level is compared to the *lsm12b* housekeeping gene. Statistical analysis was performed using Tukey's multiple comparisons test where ns means non-significant and ** P < 0.01, **** P < 0.0001. Error bars represent s.d.

Figure 5: Comparative transcriptome and proteome analysis of control and *rpgrip1l^-/-* fish.

(A-B) Volcano-plots showing differentially regulated genes in the transcriptomes of the brain (A) and dorsal trunk (B) comparing 5 controls (2 wt, 3 *rpgrip1l^-/+*) and 7 *rpgrip1l^-/-* (6 tail-up and 1 straight) fish at 5 wpf (0.9 cm body length). Each dot corresponds to one gene. A gene was considered deregulated if its associated P-adj. value was inferior to 0,05. Red and blue dots represent up-regulated (Log2FC > 0.75), and down-regulated (Log2FC < -0.75) genes, respectively. The names of the top 25 most deregulated genes are boxed. **(C)** GO term analysis of biological processes in the brain samples using Metascape. **(D)** Venn diagram of the upregulated genes in brain and trunk samples, showing many common genes between the two regions. **(E)** Selected genes of interest upregulated in the trunk and/or brain of *rpgrip1l^-/-* fish. The four genes in red were also found upregulated in the proteome of adult *rpgrip1l^-/-* fish. **(F)** Volcano-plot showing differentially expressed proteins in the brain of 5 *rpgrip1l^+/+* fish versus 5 *rpgrip1l^-/-* adult scoliotic fish (3 months pf). Each dot corresponds to one protein. A protein was considered differentially expressed if its associated P-adj. value was inferior to 0.05. Red and blue dots represent proteins enriched (Log2FC > 0.75) or depleted (Log2FC < -0.75) in mutant brains, respectively. The names of the top 25 most differentially expressed proteins are boxed.

Figure 6: Anxa2 and GFAP upregulation and increased number of LCP1-positive cells around CNS ventricles of *rpgrip1l^-/-* juvenile fish at scoliosis onset.

(A-D') Immunostaining for GFAP (A, B, C, D) and Anxa2 (A', B', C', D') on adult brain transverse sections at DiV (A-B') and RhV (cerebellar) (C-D') levels in controls (A, A', C, C') (n=3) and *rpgrip1l^-/-* scoliotic (B, B', D, D') (n=4) fish. In A-D, nuclei are stained with DAPI (blue). White stars indicate the SCO, white triangles point to the ventral RhV region. Open arrows point to a long cellular process co-labelled by Anxa2 and GFAP. **(E-L)** Immunostaining for GFAP (magenta) on transverse sections of the brain at SCO (E-H) and RhV (I-L) levels in controls (E, I) (n=4), straight *rpgrip1l^-/-* (F, G, J, K) (n=4) and scoliotic *rpgrip1l^-/-* (H, L) (n=3) juvenile (8 wpf) fish. Nuclei are stained with DAPI (white). Scale bar: 125µM for A-D', 20 µm for E-H and 100 µm for I-L. **(M)** Graph representing the percentage of GFAP-positive cells among the total number of SCO cells in control (n=4), straight *rpgrip1l^-/-* (n=4), and scoliotic *rpgrip1l^-/-* (n=3) fish at

scoliosis onset (N=1). Each dot represents the mean ratio of GFAP-positive over negative cells in 3-6 sections per fish. Green dots correspond to fish with no or very low percentage of GFAP+ cells, red dots to fish with a high percentage of GFAP+ cells. Statistical analysis was performed with Tukey's multiple comparisons test where ns means non-significant, ** means P-value < 0.01 and *** P-value < 0.001. Error bars represent s.d. "str": straight; "sco": scoliotic. **(N-Q')** Immunostaining for LCP1+ microglia/macrophages (magenta) on brain transverse sections at SCO levels in a rectangle of 100 µm x 60 µm located under the posterior commissure in controls (n=4) (N, N'), straight *rpgrip1l^{-/-}* (n=4) (O-P') and *rpgrip1l^{-/-}* at scoliosis onset (n=3) (Q, Q') 8 wpf fish (N=1). Nuclei are stained with DAPI (white). Scale bar: 20 µm. **(R)** Quantification of LCP1+ cell density around the SCO in controls (n=4), straight *rpgrip1l^{-/-}* (n=4) and scoliotic *rpgrip1l^{-/-}* (n=3) at 8 wpf. 23 sections were counted for control, 21 sections for *rpgrip1l^{-/-}* straight and 18 sections for *rpgrip1l^{-/-}* scoliotic fish. Each dot represents the mean value for one fish. Green dots correspond to the fish without any GFAP+ SCO cells, red dots correspond to the fish with GFAP+ SCO cells (SCO cells are defined as the dorsal cells located underneath the posterior commissure and above the diencephalic ventricle). Note that straight *rpgrip1l^{-/-}* fish with GFAP+ SCO cells present a higher number of LCP1+ cells. Statistical analysis was performed with Tukey's multiple comparisons test where ns means non-significant. Error bars represent s.e.m. "str": straight "sco": scoliotic.

Figure 7: NACET treatment reduces scoliosis penetrance and severity in *rpgrip1l^{-/-}* and decreases astrogliosis and LCP1+ cell number at SCO level.

(A) Kinetics of scoliosis development in *rpgrip1l^{+/-}* incrosses in the presence (n=150 fish) or absence (n=174 fish) of NACET treatment (1.5 mM) from 4 to 12 weeks (N=1). "str": straight; "sco": scoliotic. **(B)** Quantification of body axis curvature from dorsal and lateral views at 13 wpf, at the end of the NACET treatment. Each dot corresponds to one fish. Among the 174 fish raised in untreated condition, we measured 18 controls (*rpgrip1l^{+/-}*) and 33 *rpgrip1l^{-/-}*, and upon NACET treatment, 16 controls (*rpgrip1l^{+/+}*) and 23 *rpgrip1l^{-/-}*. *rpgrip1l^{+/-}* represents a pool of *rpgrip1l^{+/+}* and *rpgrip1l^{+/-}*. Error bars represent s.d. **(C)** Representative pictures of *rpgrip1l^{-/-}* fish with their curvature index without or after NACET treatment, at 13 wpf. Scale bar is 0,5 cm. Left panels correspond to an untreated *rpgrip1l^{-/-}* scoliotic fish on lateral (upper) and dorsal (lower)

views. Middle panels represent treated *rpgrip1f*[-/-,] with a lower curvature index than

scoliotic fish on lateral (upper) and dorsal (lower) views and right panels are pictures

of rescued straight *rpgrip1f*[-/-] fish on lateral (upper) and dorsal (lower) views. **(D)** Graph

representing the number of LCP1+ cells in wt (n=3), untreated *rpgrip1f*[-/-] (n=3) and

NACET-treated *rgprip1f*[-/-] (n=3) fish in the SCO region (N=1). **(E)** Immunostaining for

GFAP (green), Acetylated tubulin (white) and LCP1 (red) on adult brain transverse

sections at SCO level in untreated (upper panels) and NACET-treated (lower panels)

fish. Nuclei are stained with DAPI (blue). No GFAP staining was present in controls

(n=3) (upper left), while untreated mutants displayed strong GFAP labeling (upper

right) (n=3). NACET treatment totally suppressed gliosis in a straight *rpgrip1f*[-/-] fish

(lower right panel) (n=1) and diminished the intensity of GFAP labeling in slightly tail-

up *rpgrip1f*[-/-] (n=2). Scale bar: 12,5 µm. **(F)** Graph presenting the percentage of GFAP-

positive cells among the total number of SCO secretory cells compared between

controls (n=3), untreated *rpgrip1f*[-/-] (n=3), and NACET treated *rpgrip1f*[-/-] adult fish (n=3)

(N=1). Each dot represents the mean of % of GFAP-positive cells present in 3-6

sections per fish. Statistical analysis was performed using Tukey's multiple

comparisons test where * means P-value <0.05 and **** means P-value < 0.0001.

1530

Figure 8: Altered ciliogenesis, GFAP upregulation and increased number of

LCP1-positive cells around ventricles of *cep290*[-/-] juvenile fish at 4 wpf.

(A-F') Immunostaining of the cilia marker glutamylated tubulin (white) on 4 wpf brain

transverse sections at 3 antero-posterior SCO levels in 4 wpf control (A-C', n=3) and

cep290[-/-] scoliotic (D-F', n=3) fish. Nuclei are stained with DAPI (blue). White stars

indicate the diencephalic ventricle (DiV), above which lies the SCO. White rectangles

in C and F delineate the close-up on cilia (C', F'). Scale bar: 10 µm for A-F, 2,5 µm for

C'-F'. **(G-L)** Immunostaining for Acetylated Tubulin (G, J), GFAP (H, K) and LCP1 (I,

L) on transverse sections of the brain at SCO level in controls (G-I) (n=3), and *cep290*[-]

[/-] (J-L) (n=3) juvenile (4 wpf) fish. In G and J, nuclei are stained with DAPI (blue). Scale

bar: identical to A-F. **(M)** Graph representing the percentage of GFAP-positive cells

among the total number of SCO secretory cells in control (n=3) and scoliotic *cep290*[-/-]

(n=3) fish at scoliosis onset (N=1). Each dot represents the mean ratio of GFAP-

positive over total SCO secretory cells in 5-7 sections per fish. **(N)** Quantification of

LCP1+ cell density around the SCO in controls (n=3) and curved *cep290*[-/-] (n=3) fish

at 4 wpf. 16 sections were counted for control, 18 sections for *cep290⁻ᐟ⁻* scoliotic fish.

Each dot represents the mean value for one fish. Statistical analysis was performed

with unpaired T test where * means P-value < 0.05. Error bars represent s.d. **(O-V)**

Immunostaining for GFAP (white) on brain transverse sections at cerebellum levels (P,

R) and anterior hindbrain level (T, V) in control (P, T) (n=3), and *cep290⁻ᐟ⁻* (R, V) (n=3)

fish. Corresponding nuclear staining (DAPI in white) is shown above (O, Q, S, U). Scale

bar: 125 µm.

1553

SUPPLEMENTARY FIGURE LEGENDS

1555

Supplementary figure S1: Production and characterization of the *rpgrip1$^\Delta$* and

***rpgrip1l^{ex4}* fish lines.**

(A) Schematic exon-intron structure of the *Danio rerio rpgrip1l* gene. Two alleles were

produced by CRISPR/cas9-mediated genome engineering, *rpgrip1l^{ex4}(rpgrip1l^{bps2} in*

ZFIN) with a nonsense mutation in exon 4 and *rpgrip1l$^{\Delta4\text{-}25}$ ((rpgrip1l^{bps1} in ZFIN)* also

called *rpgrip1l$^\Delta$* or *rpgrip1l$^-$*) with a large deletion between exons 4 and 25,

encompassing most of the protein coding sequence. Both alleles had identical

embryonic phenotype (panel B and data not shown). (A') Electrophoresis gel showing

the migration profile of amplicons for each genotype, *rpgrip1l$^{\Delta/\Delta}$*, *rpgrip1l$^{\Delta/+}$*, *rpgrip1$^{+l/+}$*

produced by triplex PCR (B) Absence of laterality defects and kidney cysts in *rpgrip1l$^{\Delta/\Delta}$*

and *rpgrip1l$^{ex4/ex4}$* embryos. **(C)** Immunostaining on whole 30 hpf control (left panels)

and *rpgrip1l$^{-/-}$* (right panels) embryos to visualize the RF (anti-RF antibody, middle

panels) and cilia (acetylated tubulin antibody, lower panels) in the caudal region. Nuclei

(upper panels) are stained with DAPI. The RF is present in the central canal of *rpgrip1l$^-$*

$^{/-}$ embryos, as well as cilia in the whole neural tube width. Scale bar: 6 μm for upper

panels; 15 μm for lower panels. **(D-E)** Absence of retinal morphological defects in 5

dpf *rpgrip1l$^{-/-}$* larvae. Retinal sections of control (D) and *rpgrip1l$^{-/-}$* (E) larvae were

immunostained with Zpr3 (red) to label the external segment of photoreceptors and

Acetylated Tubulin (Ac Tub) (green) to label axonemes. D' and E' are insets in D and

E focusing on the photoreceptor layer. Scale bar: 50 μm. (**F-H'''**) Scanning electron

microscopy of brain ventricles in control (F-F''), straight (str.) *rpgrip1l$^{-/-}$* (G-G'') and

scoliotic (sco.) *rpgrip1l$^{-/-}$* (H-H''') adult (3 months) fish. Ependymal multiciliated cells of

the rhombencephalic ventricle (RhV) are almost totally absent in *rpgrip1l$^{-/-}$* scoliotic fish

(I-I'), while they are present in straight r*pgrip1l$^{-/-}$*(H-H') and controls (G-G''). Cilia of

hindbrain mono-ciliated cells in *rpgrip1l$^{-/-}$* (F'', G'') and controls (G''). Scale bars: 10 μm

for F-I, 2 μm for F'-H', 1 μm for F'', G'', H'''' and H'''.

1582

1583

1584

 Supplementary figure S2: Presence of scospondin aggregates and
heterogenous ciliary defects in the fChP of _rpgrip1l_^{-/-} juvenile fish.

(A-D) Immunostaining on transverse sections of the SCO, at anterior level (A, A', B,
B') or posterior level (C, D) in _rpgrip1l_^{+/+} (n=3) and tail-up _rpgrip1l_^{-/-} (n=3), transgenic
for Sspo-GFP (N=2). Sspo-GFP forms small cytoplasmic aggregates within SCO cells
of both genotypes (* in A' and B'), under the primary cilia extending towards the DiV,
and accumulates at the apical surface of tail-up _rpgrip1l_^{-/-}, creating a double positive
area for cilia and Sspo-GFP staining (yellow in B'). Multiciliated cells lateral to the
posterior SCO (mcc) are negative for Sspo-GFP staining in controls (C) and absent in
tail-up _rpgrip1l_^{-/-} (D). **(E-J)** GFP immunostaining on transverse section of the fChP (E-
G) or at the level of the optic tectum (H-J) in _rpgrip1l_^{+/+} (E, H, n=7), straight _rpgrip1l_^{-/-}
(F, I, n=4) and scoliotic _rpgrip1l_^{-/-} (G, J, n=5) 8 wpf fish transgenic for _sspo-GFP_ (N=2).
Scoliotic _rpgrip1l_^{-/-} fish present with filamentous material and aggregates within the
ventricles (G, J) while a small RF fragment is present within the ventricle of controls
(H) and straight mutants (I) at optic tectum level. * indicates aggregates. **(K-O'')**
Representative maximum intensity Z-stack projections of confocal transverse brain
section at the level of the fChP in _rpgrip1l_^{+/+} (K-K'') (n=7), straight _rpgrip1l_^{-/-} (L-M'')
(n=4) and scoliotic _rpgrip1l_^{-/-} (N-O'') (n=5) 8 wpf fish (N=2). Cilia are stained with
glutamylated-tubulin and acetylated-tubulin antibodies (both in white) and nuclei are
stained with DAPI (blue). Sections shown are at the level of the habenular nuclei. At
this level, monocilia in the dorsal midline region (K') and multicilia on lateral sides (K'')
(n=7/7) are present in controls. Straight _rpgrip1l_^{-/-} exhibited either normal (L', L'')
(n=2/4) or abnormal and sparser (M', M'') (n=2/4) monocilia and multicilia. Similarly,
scoliotic _rpgrip1l_^{-/-} presented either normal monocilia and multicilia (N', N'') (n=3/5) or
long monocilia (O') and decreased cilia density of multi-ciliated tufts (O'') (n=2/5). Scale
bars: 20 μm.

Supplementary figure S3: Genetic invalidation of _urp2_ in _rpgrip1l_^{-/-} fish does not
have any beneficial effect on scoliosis penetrance or severity.

(A-E) Absence of compensatory upregulation of _urp_ family members in _urp2_^{-/-} fish.
Expression levels of _urp2_ (A), _urp1_ (B), _urp_ (C), _uts2a_ (D) and _uts2b_ (E) in zebrafish in
13 wpf adults. Each dot represents the value measured for one fish; green and purple
dots represent straight and curved fish, respectively. n = 5 [_rpgrip1l_^{+/+}_; urp2_^{+/+}], 9

[*rpgrip1l^{+/+}; urp2^{-/-}*], 11 [*rpgrip1l^{-/-}; urp2^{+/-}*] and 10 [*rpgrip1l^{-/-}; urp2^{-/-}*]. Expression levels for each gene are compared to the *lsm12b* housekeeping gene. Statistical analysis was performed using the Tukey test where ns means not significant, ** $P < 0.01$, **** $P < 0.0001$. Error bars represent s.d. None of the *uts2* family members is significantly upregulated in [*rpgrip1l^{-/-}; urp2^{-/-}*] fish compared to *rpgrip1l^{-/-}*, suggesting an absence of compensation. **(F-G)** Quantification of body axis curvature on dorsal and lateral positions at 2 (F) and 4 (G) mpf. Each dot represents the curvature index of one fish. Statistical analysis was done using Tukey's multiple comparisons test where ****$P < 0.0001$, with error bars representing s.d. At 2 mpf, n=30 [*rpgrip1l^{+/+}; urp2^{+/+}*], 26 [*rpgrip1l^{+/+}; urp2^{-/-}*], 27 [*rpgrip1l^{-/-}; urp2^{+/+}*], 55 [*rpgrip1l^{-/-}; urp2^{+/-}*] and 32 [*rpgrip1l^{-/-}; urp2^{-/-}*]. At 4 mpf, n=9 [*rpgrip1l^{+/+}; urp2^{+/+}*], 7 [*rpgrip1l^{+/+}; urp2^{-/-}*], 27 [*rpgrip1l^{-/-}; urp2^{+/+}*], 48 [*rpgrip1l^{-/-}; urp2^{+/-}*] and 27 [*rpgrip1l^{-/-}; urp2^{-/-}*]. **(H)** Rose plots representing combined Cobb angle measurements at 4 mpf of [*rpgrip1l^{+/+}; urp2^{+/+}*] (n=7) in green, [*rpgrip1l^{+/+}; urp2^{-/-}*] (n=6) in blue, [*rpgrip1l^{-/-}; urp2^{+/+}*] (n=10) in pink and [*rpgrip1l^{-/-}; urp2^{-/-}*] (n=7) in yellow. Cobb angle measurements were performed on Alizarin red fish skeletons. One example of Alizarin red skeletons for each genotype is presented below its corresponding rose-plot. **(I)** Illustration of combined curvature index measurement from traced lines on lateral (left) and dorsal (right) views of fish. The curvature index of both traced lines was measured using a modified MATLAB program (LineCurvature2D function) which provided a measurement in radian/length in pixels. Scale bar: 5 mm.

Supplementary Figure S4: Complementary comparative transcriptome and proteomic analysis of *rpgrip1l^{-/-}* versus control fish.

(A, B) GO term analysis of biological processes enriched in *rpgrip1l^{-/-}* for genes upregulated in the trunk transcriptome data (A) or enriched in *rpgrip1l^{-/-}* for proteins in the brain adult proteome data (B) using Metascape. **(C)** List of the top 25 most enriched proteins in the *rpgrip1l^{-/-}* vs control brain proteome. **(D)** List of selected genes upregulated in the zebrafish *rpgrip1l* trunk transcriptome and found in different reactive astrogliosis models in single cell transcriptome studies (Matusova et al, 2023).

Supplementary figure S5: Increased GFAP and Annexin 2 staining and LCP1+ cell density in different brain regions.

(A-F') Immunostaining of Annexin A2 (Anxa2) (A-F) and GFAP (A'- F') on transverse brain sections at habenula (A, A', D, D'), tectum (B, B', E, E') and hindbrain (C, C', F, F') levels in controls (A-C') and scoliotic *rpgrip1l*$^{-/-}$ (D-F') fish. Nuclei are stained with DAPI (blue). Scoliotic *rpgrip1l*$^{-/-}$ fish upregulate Annexin A2 and GFAP in overlapping territories. Scale bar: 20 μm. **(G)** Quantification of LCP1+ cell density in the optic tectum region of *rpgrip1l*$^{+/+}$ (n=4), straight *rpgrip1l*$^{-/-}$ (n=4) and scoliotic *rpgrip1l*$^{-/-}$ (n=3) 8 wpf fish. Each dot represents a fish. 15-20 sections were counted for each brain level. Straight *rpgrip1l*$^{-/-}$ fish sections displayed around twice more Lcp1+ cells at tectum level than controls, while scoliotic *rpgrip1l*$^{-/-}$ fish had a slightly increased LCP1+ cell density compared to controls. Statistical analysis was performed using Tukey's multiple comparisons test where ns means non significative and * Pvalue < 0.05, ** Pvalue < 0.01 with error bars representing s.d. **(H-J')** Immunostaining of LCP1+ immune cells on transverse sections of the optic tectum region of *rpgrip1l*$^{+/+}$ (H, H') (n=4), straight *rpgrip1l*$^{-/-}$ (I, I') (n=4) and scoliotic *rpgrip1l*$^{-/-}$ (J, J') (n=3) 8 wpf fish. Immune cells are labelled with the LCP1 antibody (magenta) and nuclei with DAPI (blue). H', I' and J' are higher magnifications of the squared regions in H, I and J, respectively. Scale bars: 100 μm for H, I, J; 30 μm for H', I', J'. str: straight, sco: scoliotic. **(K)** Quantification of LCP1+ cell density in the cerebellum region of 8 wpf fish (N=1). The analysis compared straight *rpgrip1l*$^{-/-}$ (n=4) and scoliotic *rpgrip1l*$^{-/-}$ (n=3) fish to *rpgrip1l*$^{+/+}$ siblings (n=4). No significant difference was found between genotypes. Statistical analysis was performed using Tukey's multiple comparisons test where ns means not significant. Error bars represent s.d. "str" means straight and "sco" means scoliotic.

Supplementary figure S6: Biological assay of NACET activity on *dnaaf1*$^{-/-}$ embryos and expression level of three immune markers at adult stage. (A) Theta angle (θ) measurement between the eye, the caudal end of the yolk extension and the tail extremity, corresponding to the embryo curvature index. **(B)** Lateral views of 60 hpf representative embryos showing axial curvature phenotypes. In untreated conditions, *dnaaf1*$^{-/-}$ embryos show a characteristic curly tail-down (CTD) phenotype, whereas *dnaaf1*$^{+/+}$ siblings have a straight axis. Treatment at 27 hpf with 3 mM NACET suppresses the CTD phenotype of *dnaaf1*$^{-/-}$ embryos and does not affect axis straightness of *dnaaf1*$^{+/+}$ siblings. Embryos were measured around 2.5 to 3 mm. **(C-F)**

Rose plots representing the (360 − θ) angle of *dnaaf1$^{+/+}$* and *dnaaf1$^{-/-}$* in untreated and

3 mM NACET-treated conditions. Embryos were treated from 27hpf to 2.5 dpf and θ

angles were measured at the end of the treatment. Untreated *dnaaf1$^{+/+}$* are

represented in (C − dark green), and treated ones are in (D − light green). Untreated

dnaaf1$^{-/-}$ are in (E − blue) and treated ones are in (F − red). NACET treatment is able

to rescue curly tail down phenotype in *dnaaf1$^{-/-}$* embryos without affecting the controls.

A

rpgrip1l⁻/⁻

5 dpf

5 wpf

12 wpf

B

Scoliosis appareance in clutches from
rpgrip1l⁻/+ incrosses

C

Scoliosis penetrance in adults

D Ctrl **E** *rpgrip1l⁻/⁻*

F Dorsal view

G Lateral view

ctrl *rpgrip1l⁻/⁻* *rpgrip1l⁻/⁻*

H

| Tol2 | Foxj1a enh. | 5XMyc-RPGRIP1L | polyA. | polyA | GFP. | cmlc2 enh. | Tol2 |
| Tol2 | Col2a1 enh. | 5XMyc-RPGRIP1L | polyA. | polyA | GFP. | cmlc2 enh. | Tol2 |

I

Genotype	*rpgrip1l⁻/⁻*	*rpgrip1l⁻/⁻ ; Tg Foxj1a :RPGRIP1L*	*rpgrip1l⁻/⁻*	*rpgrip1l⁻/⁻ ; Tg Col2a1 :RPGRIP1L*
Adult Scoliotic fish (n)	72/74	0/63	28/29	28/29
Clutch (N)	N=2	N=2	N=1	N=1

J

A ChP · CCe · CC · Tel · TeO · Rh · SCO

B ROI4

C ROI4.4

D ROI6

E

Control   ***rpgrip1l⁻/⁻* straight**   ***rpgrip1l⁻/⁻* scoliotic**

F

Ventricular volume (arbitrary units)

1500 · 1000 · 500

ns

ROI4 ctl · ROI4 -/-

100 · 80 · 60 · 40 · 20

*

ctl · *rpgrip1l⁻/⁻*
ROI4.4
= RhV a

*

ctl · *rpgrip1l⁻/⁻*
ROI6
= RhV b

G

Surface of RhV b
section (arbitrary units)

250 · 200 · 150 · 100 · 50

*

Control · Rpgrip1l-/-

H

Trunk · Central canal · Reissner Fiber · Lateral view

Posterior commissure · DiV · SCO · Transverse view

D · A ↔ P · V

rpgrip1l⁺/⁺ · ***rpgrip1l⁻/⁻* straight** · ***rpgrip1l⁻/⁻* scoliotic**

I Trunk · RF   **J**   **K**   Anti-RF

L Post commissure · DiV   **M**   **N**   Ant SCO   Tg(sspo:GFP)

O   **P**   **Q**   Post SCO

A

Fold change (*Urp2* expression)

●● Straight
●● Scoliotic

WT *rpgrip1l*⁻ᐟ⁻   WT *rpgrip1l*⁻ᐟ⁻

Stage: Juvenile   Adult

C

rpgrip1l⁻ᐟ⁻ ;urp2⁺ᐟ⁺

rpgrip1l⁻ᐟ⁻ ;urp2⁻ᐟ⁻

B

Curvature index

Radian/length in pixels

0.06
0.05
0.04
0.03
0.02
0.01
0.00

ns

****

rpgrip1l⁺ᐟ⁻;urp2⁺ᐟ⁺
rpgrip1l⁺ᐟ⁺;urp2⁻ᐟ⁻
rpgrip1l⁻ᐟ⁻ ; urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻ ; urp2⁺ᐟ⁻
rpgrip1l⁻ᐟ⁻ ; urp2⁻ᐟ⁻

D

urp2

2^{-dct}

0.15

0.10

0.05

0.00

***   ****
      **

rpgrip1l⁺ᐟ⁺;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁻
rpgrip1l⁻ᐟ⁻;urp2⁻ᐟ⁻

E

urp1

2^{-dct}

0.15

0.10

0.05

0.00

●● Straight
●● Scoliotic

**   ns

rpgrip1l⁺ᐟ⁺;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁺
rpgrip1l⁻ᐟ⁻;urp2⁺ᐟ⁻
rpgrip1l⁻ᐟ⁻;urp2⁻ᐟ⁻

A — Brain

B — Dorsal trunk

C

GO:0035082: axoneme assembly
GO:0001539: cilium or flagellum-dependent cell motility
GO:0002460: adaptive immune response
GO:0003351: epithelial cilium movement involved in extracellular fluid movement
GO:0007276: gamete generation
R-HSA-6785807: Interleukin-4 and Interleukin-13 signaling
GO:0045921: positive regulation of exocytosis

D — Up-regulated genes

Trunk   Brain
560   54   6

E

Process/Pathway	Gene	Trunk.LogFC	Trunk P adj.	Brain.LogFC	Brain P adj.
Foxj1a and	*tbata*	2,25	8,82E-22	0,84	1,80E-07
Its targets	*foxj1a*	1,90	3,77E-23	1,36	9,67E-21
	enkur	1,88	8,98E-24	0,76	2,13E-06
	stpg1	1,80	1,10E-09	0,69	3,00E-05
	ccdc65	1,73	8,65E-11	0,74	1,25E-05
Regulators of	*pkd1b*	3,20	2,21E-47	1,90	1,08E-45
axis curvature	*urp1*	1,92	1,32E-16	-	-
	urp2	0,96	1,22E-03	0,87	5,44E-09
	sspo	1,18	4,16E-05	-	-
Immune	*irg1l*	2,45	1,20E-16	-	-
Response	*il11a*	2,23	3,09E-16	0,67	1,21E-05
	c4	1,97	1,50E-12	1,54	5,63E-29
	anxa2a	1,52	8,15E-17	1,53	7,04E-26
	mmp9	1,00	3,94E-04	0,46	4,02E-02
	c7a	1,86	8,82E-22	0,80	9,38E-07
	stat3	0,76	3,63E-07	0,80	5,23E-10

F — Brain proteome

Adult rpgrip1l⁺/⁺

DiV

A · DAPI GFAP A' · ANXA2

RhV

C ▲ · C' ▲

Scoliotic adult rpgrip1l⁻/⁻

B · DAPI GFAP B' · ANXA2

D ▲ △ · D' ▲ △

rpgrip1l⁺/⁺ straight rpgrip1l⁻/⁻ rpgrip1l⁻/⁻ scoliosis onset

DiV

E F G H · DAPI GFAP

RhV (cerebellar lev.)

I J K L

M

% of GFAP(+) cells among SCO cells

ns *** **

rpgrip1l⁺/⁺ rpgrip1l⁻/⁻ str. rpgrip1l⁻/⁻ sco.

rpgrip1l⁺/⁺ straight rpgrip1l⁻/⁻ rpgrip1l⁻/⁻ scoliosis onset

N O P Q · DAPI LCP1

N' O' P' Q'

R

LCP1(+) cells density

P value 0,07

ns ns

WT rpgrip1l⁻/⁻ str. rpgrip1l⁻/⁻ sco.

A **Scoliosis penetrance in *rpgrip1l*⁺/⁻ incross**

B **Curvature index**

C **Scoliotic phenotype and associated curvature index of *rpgrip1l*⁻/⁻ adults**

D **Lcp1 cells in SCO region**

E

F % of GFAP(+) cells within the SCO

Control

cep290 -/-

A B C C' — Glu.Tub DAPI

D E F F'

Control

cep290 -/-

G J — Ac.Tub DAPI

H K — GFAP

I L — LCP1

M

Astrogliosis within SCO cells

% GFAP(+) SCO cells

-5

*

Cep290⁺/⁺ Cep290⁻/⁻

N

Immune cells density

Av. Lcp1 per field

*

Cep290⁺/⁺ Cep290⁻/⁻

Control

cep290 -/-

O Q — DAPI

P R — GFAP

Control

cep290 -/-

S U — DAPI

T V — GFAP

A

Exon 4   Exon 25

Start
Codon
CRISPR site   Deletion   CRISPR site

A'

544 bp : ex4F/ex25R for MUT*
364 bp : ex25F/ex25R for WT

B

	Heart looping	Kidney cysts	Total nb embryos
wt	1	0	900
rpgrip1l^ex4/ex4	0	0	73
rpgrip1l^Δ/Δ	3	0	694

C

Zpr3 Ac-tub DAPI

Supplementary Figure S1

Supplementary Figure S2

Supplementary Fig S3

A: Up-regulated genes in trunk

GO:0003341: cilium movement
GO:0070286: axonemal dynein complex assembly
GO:0003352: regulation of cilium movement
GO:0090660: cerebrospinal fluid circulation
GO:1904158: axonemal central apparatus assembly
GO:0002252: immune effector process
GO:0071345: cellular response to cytokine stimulus
GO:0007420: brain development
hsa04814: Motor proteins
GO:0002521: leukocyte differentiation
GO:0070663: regulation of leukocyte proliferation
WP2855: Dopaminergic neurogenesis
GO:0032735: positive regulation of interleukin-12 production
GO:0120229: protein localization to motile cilium
GO:0150063: visual system development
GO:0040017: positive regulation of locomotion
GO:0035239: tube morphogenesis
GO:0048598: embryonic morphogenesis
GO:0044703: multi-organism reproductive process
GO:0060541: respiratory system development

-log10(P)

B: Enriched proteins in adult brain

GO:0051336: regulation of hydrolase activity
GO:0010638: positive regulation of organelle organization
GO:0006897: endocytosis
GO:1905686: positive regulation of plasma membrane repair
GO:0006886: intracellular protein transport
GO:0030162: regulation of proteolysis
R-HSA-2454202: Fc epsilon receptor (FCERI) signaling
WP5323: Fatty Acids and Lipoproteins Transport in Hepatocytes
GO:0060627: regulation of vesicle-mediated transport
GO:2000644: regulation of receptor catabolic process
GO:0006954: inflammatory response
GO:0071363: cellular response to growth factor stimulus
GO:0016310: phosphorylation
GO:0031329: regulation of cellular catabolic process
GO:0006887: exocytosis
GO:0032944: regulation of mononuclear cell proliferation
GO:0030036: actin cytoskeleton organization
GO:0150105: protein localization to cell-cell junction
GO:0032386: regulation of intracellular transport
GO:0050778: positive regulation of immune response

-log10(P)

C: 25 highest P Value

Protein	FC	P value
anxa2a	7,28	4,82E-08
echdc2	4,97	1,99E-06
anxa1a	6,06	9,6856E-06
anxa5b	2,37	3,3303E-05
capsla	3,72	0,00011272
tagln2	2,53	0,00011315
psph	1,72	0,00011655
c7a	3,07	0,0002431
mapk9	2,49	0,00029106
vav3b	2,31	0,00036912
serpina7	2,46	0,0003851
gps1	1,92	0,00057247
synrg	2,08	0,00066171
c3 like	3,77	0,00068862
tut7	2,01	0,00102053
gsk3bb	2,40	0,00109597
tmed3	3,44	0,00148983
mrc1a	2,10	0,00152123
wdr12	2,78	0,00186884
wdfy2	1,97	0,00235225
plcd4b	2,24	0,00249521
itchb	1,81	0,00250875
ehd1b	1,94	0,00253596
exoc6	1,77	0,00298918
stat3	5,31	0,00319733

D: Activated astrocytes markers

Hum. Gene	Zf Gene	FC (trunk)	P.adj. Value
S100B	s100b	2,98	7,35711E-27
C4b	c4b	5,15	1,48224E-21
GFAP	gfap	2,49	1,99844E-09
CTSB	ctslb	3,60	1,29502E-08
CAPS	capsla	2,67	2,31406E-06
GLAST/SLC1A3	slc1a3b	2,68	1,46788E-05
PLCXD3	plcxd3	1,74	1,6557E-05
VIM	viml	2,14	8,30568E-05

Supplementary Figure S4

Supplementary Figure S5

Supplementary Figure S6

Subcommissural Organ (SCO)

SCO
RF

Posterior
Commissure

DiV

mcc

Defective CSF flow
Motility/Coordination defects

Defective Cilia signalling
Astrogliosis/
immune cells attraction

• Sco-spondin

GFAP+

Immune cells

SCO