ASTROGLIOSIS AND NEUROINFLAMMATION UNDERLIE SCOLIOSIS UPON CILIA DYSFUNCTION

Morgane Djebar[1], Isabelle Anselme[1], Guillaume Pezeron[2], Pierre-Luc Bardet[1], Yasmine Cantaut-Belarif[3], Alexis Eschstruth[1], Diego López Santos[1], Hélène Le Ribeuz[1], Arnim Jenett[4], Hanane Khoury[1], Joëlle Véziers[5], Caroline Parmentier[6], Aurélie Hirschler[7], Christine Carapito[7], Ruxandra Bachmann-Gagescu[8], Sylvie Schneider-Maunoury[1]* and Christine Vesque[1]*

Sorbonne Université, CNRS UMR7622, INSERM U1156, Institut de Biologie Paris Seine (IBPS) - Developmental Biology Unit, 75005, Paris, France
Molecular Physiology and Adaptation (PhyMA - UMR7221), Muséum National d'Histoire Naturelle, CNRS, Paris, France
Institut du Cerveau et de la Moelle épinière (ICM), Sorbonne Université, Inserm U 1127, CNRS UMR 7225, F-75013, Paris, France
TEFOR Paris-Saclay, CNRS UMS2010 / INRA UMS1451, Université Paris-Saclay.
Inserm UMR 1229, CHU Nantes PHU4 OTONN, SC3M facility, Inserm UMS 016, CNRS 3556, Université de Nantes, Nantes, France.
Sorbonne Université, CNRS UMR8246, INSERM U1130, Institut de Biologie Paris Seine (IBPS) – Neurosciences Paris Seine (NPS), 75005, Paris, France
Laboratoire de Spectrométrie de Masse Bio-Organique, IPHC, UMR 7178, Université de Strasbourg, CNRS, Infrastructure Nationale de Protéomique ProFI - FR2048, Strasbourg, France.
Institute of Medical Genetics, University of Zurich, Wagistrasse 12, 8952 Schlieren, Zurich, Switzerland; Institute of Molecular Life Sciences, University of Zurich, Winterthurerstrassse 190, 8057 Zurich, Switzerland
* Co-corresponding authors, co-last authors.
sylvie.schneider-maunoury@sorbonne-universite.fr
christine.vesque@sorbonne-universite.fr

ABSTRACT

Cilia defects lead to scoliosis in zebrafish, but the underlying pathogenic mechanisms are poorly understood and may diverge depending on the mutated gene. Here, we dissected the mechanisms of scoliosis onset in a zebrafish mutant for the *rpgrip1l* gene encoding a ciliary transition zone protein. *rpgrip1l* mutant fish developed scoliosis with near-total penetrance but asynchronous onset in juveniles. Taking advantage of this asynchrony, we found that curvature onset was preceded by ventricle dilations and was concomitant to the perturbation of Reissner fiber polymerization and to the loss of multiciliated tufts around the subcommissural organ. Rescue experiments showed that Rpgrip1l was exclusively required in *foxj1a*-expressing cells to prevent axis curvature. Genetic interactions investigations ruled out Urp1/2 levels as a main driver of scoliosis in *rpgrip1* mutants. Transcriptomic and proteomic studies identified neuroinflammation associated with increased Annexin levels as a potential mechanism of scoliosis development in *rpgrip1l* juveniles. Investigating the cell types associated with *annexin2* over-expression, we uncovered astrogliosis, arising in glial cells surrounding the diencephalic and rhombencephalic ventricles just before scoliosis onset and increasing with time in severity. Anti-inflammatory drug treatment reduced scoliosis penetrance and severity and this correlated with reduced astrogliosis and macrophage/microglia enrichment around the diencephalic ventricle. Mutation of the *cep290* gene encoding another transition zone protein also associated astrogliosis with scoliosis. Thus, we propose astrogliosis induced by perturbed ventricular homeostasis and associated with immune cell activation as a novel pathogenic mechanism of zebrafish scoliosis caused by cilia dysfunction.

INTRODUCTION

Idiopathic scoliosis (IS) is a 3D rotation of the spine without vertebral anomalies that affects about 3% of adolescents worldwide. Its etiology remained mysterious, mainly because of the lack of appropriate animal models and the complexity of its inheritance profile in large families presenting a severe scoliosis trait (Patten et al., 2015). Since the description of the first IS model in the zebrafish *ptk7* gene mutant (Hayes et al., 2014), a number of mutants for genes encoding ciliary proteins were shown to develop scoliosis at late larval and juvenile stage (4 to 12 weeks post-fertilization, wpf) without any vertebral fusion or fracture, highlighting the link between cilia function and straight axis maintenance in that species (Grimes et al., 2016; Wang et al., 2022).

Cilia are microtubular organelles with sensory and/or motile functions. Zebrafish mutants in genes involved in cilia motility or in intraflagellar transport display severe embryonic axis curvature and are usually not viable beyond larval stages (Grimes et al., 2016; Rose et al., 2020). In contrast, mutants in genes implicated in ciliary gating or ciliary trafficking encoding respectively components of the transition zone (TZ) or of the BBsome complex survive to adult stage and develop scoliosis with variable penetrance (Bentley-Ford et al., 2022; Masek et al., 2022; Song et al., 2020; Wang et al., 2022). The molecular basis for these differences is still poorly understood, even if progress has been made in our understanding of zebrafish scoliosis. Cilia-driven movements of the cerebrospinal fluid (CSF) are involved in zebrafish axis straightness, both in embryos and juveniles (Boswell and Ciruna, 2017; Grimes et al., 2016) and are tightly linked to the assembly and maintenance of the Reissner fiber (RF), a SCO-spondin polymer secreted by the subcommissural organ (SCO) and running down the brain and spinal cord CSF-filled cavities (Cantaut-Belarif et al., 2018). RF loss at embryonic stage in null *sspo* mutants generates ventral curvatures of the posterior axis of the body and is lethal while its loss at juvenile stage in hypomorphic *sspo* mutants triggers scoliosis with full penetrance (Cantaut-Belarif et al., 2018; Rose et al., 2020; Troutwine et al., 2020). Signaling downstream of the RF in embryos implicates Urp1 and Urp2, two neuropeptides of the Urotensin 2 family expressed in ventral CSF-contacting neurons (CSF-cNs), which act on dorsal muscles in embryos and larvae (Cantaut-Belarif et al., 2020; Gaillard et al., 2023; Lu

et al., 2020; Zhang et al., 2018). Their combined mutations or the mutation of their receptor gene *uts2r3* lead to scoliosis at adult stages (Bearce et al., 2022; Gaillard et al., 2023; Zhang et al., 2018). However, whether RF maintenance and Urp1/2 signaling are perturbed in juvenile scoliotic mutants of genes encoding TZ proteins ("TZ mutants") and how these perturbations are linked to scoliosis is unknown, especially because TZ mutants do not display any sign of embryonic or larval cilia motility defects (Wang et al., 2022). Moreover, two scoliotic models develop axial curvature while maintaining a RF showing that their curvature onset occurs independently of RF loss (Meyer-Miner et al., 2022; Xie et al., 2023). Finally, neuro-inflammation has been described in a small subset of zebrafish IS models for which anti-inflammatory/anti-oxidant treatments (with NAC or NACET) partially rescue scoliosis penetrance and severity (Rose et al., 2020; Van Gennip et al., 2018).

In this paper we dissect the mechanisms of scoliosis appearance in a novel deletion allele of the zebrafish TZ *rpgrip1l* gene. *RPGRIP1L* mutations have been found in neurodevelopmental ciliopathies, Meckel syndrome (MKS) and Joubert syndrome (JS) (Delous et al., 2007)(Arts et al., 2007)characterized by severe brain malformations. Other organs such as the kidney and liver are variably affected. Some JS patients also present with severe scoliosis, but this phenotypic trait is of variable penetrance (Delous et al., 2007)(Brancati et al., 2008). Functional studies performed in mouse and zebrafish have shown that *Rpgrip1l* is required for proper central nervous system patterning due to complex perturbations of Sonic Hedgehog signaling (in mice) and cilia planar positioning (in mice and zebrafish) (Andreu-Cervera et al., 2021; Mahuzier et al., 2012; Vierkotten et al., 2007). Zebrafish *rpgrip1l* mutants are viable and develop an axis curvature phenotype at juvenile stages with nearly full penetrance. Using a variety of approaches, we show that i) Rpgrip1l is required in ciliated cells surrounding ventricular cavities for maintaining a straight axis; ii) at juvenile stages, *rpgrip1l* mutants develop cilia defects associated with ventricular dilations and loss of the RF; iii) increased *urp1/2* expression in *rpgrip1l* mutants does not account for spine curvature defects; and iv) a pervasive astrogliosis process associated with neuroinflammation participates in scoliosis onset and progression. Finally, we show that astrogliosis is conserved in another zebrafish ciliary TZ mutant, *cep290,* identifying a novel mechanism involved in scoliosis upon ciliary dysfunction.

RESULTS

Rpgrip1l$^{-/-}$ fish develop scoliosis asynchronously at juvenile stage

To study the mechanisms of scoliosis appearance upon ciliary dysfunction in

zebrafish, we made use of a viable zebrafish deletion mutant in the *rpgrip1l* gene

encoding a ciliary transition zone protein (*rpgrip1l$^{\Delta}$*, identified in ZFIN as *rpgrip1l^{bps1}*)

Supplementary Figure S1A). We produced a 26 kb deletion after a CRISPR-Cas9

double-cut which removes 22 out of the 26 exons of the protein (Supplementary

Figure S1A) and triggers protein truncation at amino-acid (aa) position 88 out of

1256. 90-100% of *rpgrip1l$^{\Delta/\Delta}$* embryos and larvae were straight (0-10%, depending on

the clutch, displayed a sigmoid curvature). They did not display any additional

defects found in many ciliary mutant embryos or larvae such as randomized left-right

asymmetry, kidney cysts or retinal anomalies (Bachmann-Gagescu et al., 2011;

Kramer-Zucker et al., 2005; Tsujikawa and Malicki, 2004)(Figure 1A and

Supplementary Figure S1B, D, E). Another allele with a frameshift mutation in exon 4

(*rpgrip1l^{ex4},* identified in ZFIN as *rpgrip1l^{bps2}*) showed a similar phenotype

(Supplementary Figure S1A, B). Considering the large size of the deletion, we

considered the *rpgrip1l$^{\Delta}$* allele as a null and called it *rpgrip1l$^{-/-}$* thereafter. *rpgrip1l$^{-/-}$*

animals developed scoliosis during juvenile stages. Scoliosis appearance was

asynchronous between clutches, from 4 weeks post fertilization (wpf) (around 1 cm

length) to 11 wpf (1,8- 2,3 cm), and also within a clutch. It started with slight upward

bending of the tail (the *tail-up* phenotype) and progressed toward severe curvature

(Figure 1A, B) with 90% penetrance in adults (100% in females and 80% in males)

(Figure 1C). Micro-computed tomography (µCT) at two different stages (5 wpf and 23

wpf) confirmed that spine curvature was three-dimensional and showed no evidence

of vertebral fusion, malformation or fracture (Figure 1D-G and Supplementary Movie

1).

Thus, the ciliary *rpgrip1l$^{-/-}$* mutant shows almost totally penetrant, late-onset juvenile

scoliosis without vertebral anomalies, making it a valuable model for studying the

etiology of human idiopathic scoliosis. Furthermore, the asynchronous onset of

curvature is an asset for studying the chronology of early defects leading to scoliosis.

Reintroducing RPGRIP1L in foxj1a lineages suppresses scoliosis in rpgrip1l^{-/-} fish

Since *rpgrip1l* is ubiquitously expressed in zebrafish (Mahuzier et al., 2012), scoliosis could have different tissue origins, including a neurological origin as shown for the *ptk7* and *ktnb1* scoliotic fish (Meyer-Miner et al., 2022; Van Gennip et al., 2018). Indeed, no tissue-specific rescue has been performed in mutants for zebrafish ciliary TZ or BBS genes. We tested if introducing by transgenesis a tagged human RPGRIP1L protein under the control of the *col2a1a (Dale and Topczewski, 2011)* or *foxj1a* (Van Gennip et al., 2018) enhancers (Figure 1H) would reduce scoliosis penetrance and severity. The *foxj1a* enhancer drives expression in motile ciliated cells, among which ependymal cells lining CNS cavities (Van Gennip et al., 2018), while the *col2a1a* regulatory region drives expression in cartilage cells including intervertebral discs and semi-circular canals, tissues that have been proposed to be defective in mammalian scoliosis models (Hitier et al., 2015; Karner et al., 2015). After selecting F1 transgenic fish that expressed the Myc-tagged RPGRIP1L protein in the targeted tissue at 2.5 dpf, we introduced one copy of each transgene into the *rpgrip1l^{-/-}* background. The Foxj1a:5XMycRPGRIP1L transgene was sufficient to fully suppress scoliosis in *rpgrip1l^{-/-}* fish: none of the transgenic *rpgrip1l^{-/-}* fish became scoliotic while virtually all the non-transgenic *rpgrip1l^{-/-}* siblings did (Figure 1I, J). In contrast, RPGRIP1L expression triggered by the *col2a1a* promoter did not have any beneficial effect on scoliosis penetrance (Figure 1I, J). In conclusion, *rpgrip1l* expression in *foxj1a*-expressing cells is sufficient to maintain a straight axis.

Rpgrip1l^{-/-} adult fish display cilia defects along CNS cavities

We observed normal cilia in the neural tube of *rpgrip1l^{-/-}* embryos (Supplementary Figure S1C). In 3 month-adults, scanning electron microscopy (SEM) at the level of the rhombencephalic ventricle (RhV) showed cilia tufts of multiciliated ependymal cells (MCC) lining the ventricle in controls and straight *rpgrip1l^{-/-}* fish (Supplementary Figure S1F-G, F'-G'). In scoliotic *rpgrip1l^{-/-}* fish, cilia tufts were sparse and disorganized (Supplementary Figure S1H, H'). Cilia of mono-ciliated ependymal cells

were present in *rpgrip1l⁻/⁻* fish, some of them with irregular shape. (Supplementary
Figure S1, F'',G'', H'' and H''').

MCC defects at SCO level correlate with scoliosis appearance at juvenile
stages

We then took advantage of the asynchrony in scoliosis onset to correlate, at a given
stage, cilia defects at different levels of the CNS cavities with the spine curvature
status (straight or curved) of *rpgrip1l⁻/⁻* juveniles. We analyzed cilia defects in three
CNS regions implicated in scoliosis appearance: the spinal cord central canal (since
scoliosis develops in the spine), the forebrain choroid plexus (fChP) whose cilia
defects have been associated with scoliosis in *katnb1⁻/⁻* fish (Meyer-Miner et al.,
2022) and the subcommissural organ (SCO), which secretes SCO-spondin whose
absence leads to scoliosis (Rose et al., 2020; Troutwine et al., 2020).

In the spinal cord central canal of 8 wpf juvenile fish, cilia density was reduced to the
same extent in *rpgrip1l⁻/⁻* fish irrespective of the fish curvature status (Figure 2 A-C,
E). In addition, cilia length increased in straight mutants and was more heterogenous
than controls in scoliotic mutants (Figure 2 A'-C', D). Arl13b content appeared
severely reduced in mutants (Figure 2 A''-C''). In the fChP, mono- and multi-ciliated
cell populations were observed with a specific distribution along the anterior-posterior
axis, as previously described (D'Gama et al., 2021). At the level of the habenula
nuclei, cells of dorsal and ventral midline territories were monociliated, while lateral
cells presented multicilia tufts (Supplementary Figure S2 K-K''). *rpgrip1l⁻/⁻* juveniles
presented an incomplete penetrance of ciliary defects in the fChP, with no correlation
between cilia defects and the curvature status (Supplementary Figure S2, L-O'').

The SCO lies at the dorsal midline of the diencephalic ventricle (DiV), under the
posterior commissure (PC) (Figure 2 G, H, Supplementary Figure 2A, C). Cilia in and
around the SCO have not been described previously in zebrafish. We showed that, in
control animals, Sco-spondin (Sspo) secreting cells varied in number along SCO
length, from 24 cells anteriorly to 6-8 cells posteriorly (Supplementary Figure 2 A, A',
C) and were monociliated at all anteroposterior levels (Figure 2G-H, Supplementary
Figure 2 A'). The Sspo protein produced from one copy of the transgene Scospondin-
GFP[ut24] (Troutwine et al., 2020) formed cytoplasmic granules around SCO cell nuclei,
under emerging primary cilia (Supplementary Figure 2A'). Conversely, MCCs lateral

to the SCO did not secrete Sspo and were prominent towards posterior (Figure 2H,
Supplementary Figure 2C), as observed in the rat brain (Collins and Woollam, 1979).
Monocilia of Sspo secreting cells were still present but appeared longer in straight
rpgrip1l[-/-] juveniles and could not be individualized anymore at anterior SCO level in
scoliotic fish: we observed a weaker and denser acetylated and glutamylated tubulin
staining that likely corresponded to longer intertwined cilia (Figure 2 I, I', K, K').
Lateral multicilia tufts were preserved in straight *rpgrip1l*[-/-] juveniles (n=5) (Figure 2 J)
but were missing at posterior SCO level in tail-up or scoliotic *rpgrip1l*[-/-] fish (n=7)
(Figure 2L).
Thus, *rpgrip1l* mutants develop both monocilia and multicilia defects at juvenile
stages, but cilia defects in the spinal cord and in the fChP do not correlate with axis
curvature status of the fish. By contrast, the loss of MCC cilia at the SCO strictly
correlates with scoliosis onset, suggesting a causal relationship.
178

179 *Rpgrip1l*[-/-] juveniles show ventricular dilations and loss of the Reissner fiber at
180 scoliosis onset

Ciliary beating is an essential actor of CSF flow and of ventricular development in
zebrafish larvae (Kramer-Zucker et al., 2005) (Olstad et al., 2019). Thus, ciliary
defects in the brain and spinal cord of *rpgrip1l*[-/-] fish could lead to abnormal
ventricular and central canal volume and content. In the spinal cord, the lumen of the
central canal was enlarged at thoracic and lumbar levels in all *rpgrip1l*[-/-] mutant
juveniles, with no correlation to the axis curvature status (Figure 2 A-C; F). To
determine the timing of brain ventricular dilations relative to scoliosis, we analyzed
ventricular volume at the onset of spine curvature (Figure 3A-G). Ventricular
reconstruction was performed on cleared brains of 4 control and 4 *rpgrip1l*[-/-] (3 tail-up
and 1 straight) fish at 5 wpf, stained with ZO1 to highlight the ventricular border and
with DiI to outline brain shape, focusing on the RhV at the level of the posterior
midbrain and hindbrain (Figure 3A-D). We identified a significant increase in ventricle
volume in *rpgrip1l*[-/-] fish compared to controls, that was restricted to the ventral
regions of the RhV in regions ROI4.4 (green sphere in Figure 3C) and ROI6 (green
sphere in Figure 3D), as confirmed by numerical sections (Figure 3E, G). Ventricle

dilations were also present, although to a lesser extent, in the RhV of the unique
straight *rpgrip1l*[-/-] fish of this study (green squares and triangle in Figure 3F, G).
Defective ependymal ciliogenesis has been associated with abnormal CSF flow and
defective Reissner Fiber (RF) maintenance in several scoliotic zebrafish models
(Grimes et al., 2016)(Rose et al., 2020). The RF is mainly composed of Sspo
secreted by the SCO and floor plate (FP) and its loss at juvenile stages triggers
scoliosis in zebrafish (Rose et al., 2020)(Troutwine et al., 2020). To visualize the RF,
we used an antiserum that labels the RF in zebrafish embryos (Didier et al.,
1995)(Cantaut-Belarif et al., 2018) and we confirmed the results by introducing one
copy of the *sspo-GFP* knock-in allele, sco-spondin-GFP[ut24,] (Troutwine et al., 2020)
by genetic cross into *rpgrip1l*[-/-] animals. The RF formed normally in *rpgrip1l*[-/-]
embryos, as expected given the absence of embryonic curvature (Supplementary
Figure S1 C). To study its maintenance in *rpgrip1l*[-/-] juveniles, we immunostained
spinal cord longitudinal sections of 7-8 wpf curved or straight fish of the same clutch.
The RF was visualized as a 1 µm diameter rod in the central canal of the neural tube
and few dots were present in the apical cytoplasm of FP cells (Figure 3I) as well as in
the SCO of all controls (n=8, Figure 3L, O) and in ventricles caudal to the SCO
(Supplementary Figure 2H). The RF was also present in all *rpgrip1l*[-/-] straight fish
(n=9, Figure 3J, Supplementary Figure 2I). In contrast, in *rpgrip1l*[-/-] tail-up and
scoliotic fish, the RF was absent at all axis levels and a few SCO-spondin-positive
debris were present in the central canal (n=8, Figure 3K). Transverse sections at the
level of the SCO and caudal to the SCO in severely scoliotic fish revealed abnormally
packed material in the ventricle (Figure 3N, Q, Supplementary Figure 2J), while Sspo
secreted cells from *rpgrip1l*[-/-] at scoliosis onset, ie in juveniles with a slight tail-up
phenotype, appeared to produce Sspo-GFP cytoplasmic granules as in controls
(Supplementary Figure 2B,B',D). We also observed that Sspo aggregates spread
anteriorly into the diencephalic ventricle (DiV) at fChP level in scoliotic juveniles
(Supplementary Figure 2G, n=7/7), a situation never observed in controls
(Supplementary Figure 2E, n=0/6) or in straight juveniles (Supplementary Figure 2F,
n=0/5).
Thus, our study uncovers a strict temporal correlation between the loss of MCC at
posterior SCO level, impaired RF polymerization and scoliosis onset in *rpgrip1l*[-/-]
juvenile fish.

Urp1/2 upregulation does not contribute to scoliosis penetrance or severity in rpgrip1l$^{-/-}$ fish

In scoliotic fish where the RF is not maintained at juvenile stage, *urp1* and *urp2* levels are downregulated (Rose et al., 2020). *urp1* and *urp2* encode peptides of the Urotensin 2 family, expressed in ventral spinal CSF-cNs (Parmentier et al., 2011; Quan et al., 2015). Their expression levels have to be finely tuned to keep a straight axis since their combined loss-of-function induces larval kyphosis that evolves into adult scoliosis (Bearce et al., 2022; Gaillard et al., 2023), while an increased dose of Urp1/2 peptides induces an upward curvature in embryos (Zhang et al., 2018) and juveniles (Gaillard et al., 2023; Lu et al., 2020). We therefore monitored via qRT-PCR *urp1 and urp2* expression levels on individual straight or scoliotic *rpgrip1l$^{-/-}$* juvenile and/or adult fish. *urp2* was the most highly expressed gene among all *urotensin 2* members at juvenile stage (Figure 4A, D, E and Supplementary Figure 3A-E. *urp2* expression was upregulated in 5 weeks juveniles, both in straight and scoliotic fish and its upregulation was maintained in adult scoliotic fish (Figure 4A).

Given the early and long-lasting upregulation of *urp1/2* in *rpgrip1l$^{-/-}$* fish, we attempted to rescue axis curvature by decreasing global Urp1/2 production via genetic means by downregulating *urp2* expression level. To that end, we generated double (*rpgrip1l$^{+/-}$; urp2$^{+/-}$*) heterozygous fish using the recently generated *urp2* deletion allele (Gaillard et al., 2023) and crossed them to produce double mutants. Using a curvature quantification method on whole fish (Supplementary Figure 3I and methods), we observed that the removal of one or two *urp2* functional allele(s) in *rpgrip1l$^{-/-}$* fish did not lower their curvature index (Figure 4B), neither at two mpf (Supplementary Figure 3 F) nor at four mpf (Supplementary Figure 3 G-H). We observed that *urp2* mRNA amounts were reduced by removing one or two functional copies of *urp2*, probably as a consequence of mRNA decay (Gaillard et al., 2023) (Figure 4D). We also showed that none of the other *urotensin 2* family members was upregulated by a compensation mechanism upon *urp2* loss (Figure 4E for *urp1* and Supplementary Figure 3C-E for *urp, uts2a* and *uts2b*).

261 Thus, lowering global Urp1/2 expression level in *rpgrip1l* mutants did not have any
262 beneficial effect on the axis curvature phenotype, ruling out altered *urp1/2* signaling
263 as a main driver of scoliosis in *rpgrip1l* mutants.
264

Regulators of motile ciliogenesis, inflammation genes and annexins are
266 **upregulated in rpgrip1l$^{-/-}$ juveniles**
267

To get further insight into the early mechanisms driving spine curvature in *rpgrip1l*
mutants, we obtained the transcriptome of the brain and dorsal trunk of 7 *rpgrip1l$^{-/-}$*
juveniles at scoliosis onset (6 tail-up and 1 straight 5 wpf juveniles) and 5 control
siblings (2 *rpgrip1l$^{+/+}$* and 3 *rpgrip1l$^{+/-}$*). Differential gene expression (DGE) analysis
was performed with DSEQ2 software within the Galaxy environment. A complete
annotated gene list is displayed in Supplementary Table 1 (Trunk) and Table 2
(Brain). Filtering the gene list to select for P-adj < 0.05 and Log2 FC > 0.75 or < -0.75
recovered a vast majority of upregulated genes in the dorsal trunk (75 downregulated
vs 614 upregulated genes) and, to a lesser extent, in the brain (1 downregulated,
rpgrip1l, vs 60 upregulated genes) as illustrated on the volcano plots from DGE
analysis of the brain (Figure 5A) and trunk (Figure 5B). 54 out of 60 genes
upregulated in the brain were also upregulated in the trunk, suggesting that, in the
trunk, these genes were specific to the CNS (Figure 5D). In both the brain (Figure
5C) and the trunk (Supplementary Figure S4A), the most upregulated biological
processes in GO term analysis included cilium movement and cilium assembly.
Among the 20 most significant GO terms found for upregulated genes in the trunk, 8
were related to ciliary movement/assembly (Supplementary Figure S4A). The
upregulation in both tissues of *foxj1a,* encoding a master transcriptional activator of
genes involved in cilia motility (Yu et al., 2008) (Figure 4E) likely contributed to this
enrichment. To confirm this hypothesis, we compared our dataset with those of
targets of Foxj1 from (Choksi et al., 2014; Mukherjee et al., 2019; Quigley and
Kintner, 2017). We found that 106 out of 614 upregulated genes were targets of
Foxj1, among which 77 were direct targets (Supplementary Table 3). Several genes
encoding regulators of embryonic axis curvature were also upregulated in *rpgrip1l$^{-/-}$*
fish, among which *pkd1b* (Mangos et al., 2010), *urp1* and *urp2* (Bearce et al., 2022;
Gaillard et al., 2023; Zhang et al., 2018) and *sspo* (Cantaut-Belarif et al., 2018; Rose

et al., 2020; Troutwine et al., 2020) (Figure 5E). Finally, GO terms enrichment
analysis also highlighted processes associated with inflammation and innate and
adaptive immune responses, as described for the scoliotic *ptk7* model (Van Gennip
et al., 2018) sharing for example *irg1l* and complement genes up-regulation (Figure 5
C, E, and Supplementary Figure 4A).

In order to confirm these results and to identify processes affected at the post-
transcriptional level in *rpgrip1l* mutants, we performed a quantitative proteomic
analysis by mass spectrometry of brains from 5 *rpgrip1l*$^{-/-}$ adult fish and 5 control
siblings. This analysis detected 5706 proteins present in at least 4 over the 5
replicates, of which 172 are associated with a P value < 0.05 and a Log2FC > +0.75
and < -0.75. 127 proteins were present in higher and 45 in lower quantity, as
illustrated on the volcano plot (Figure 5F and Supplementary Table 4). Pathways
enriched in *rpgrip1l*$^{-/-}$ brains included regulation of vesicular trafficking and membrane
repair processes, as well as proteolysis and catabolic processes, suggesting
profound perturbation of cellular homeostasis (Supplementary Figure 4B). Among the
proteins that were present in highest quantity in mutants, several members of the
Annexin family were identified: Anxa2a, Anxa1a, Anxa5b (Figure 5F and
Supplementary Figure 4C. Annexins are associated with membranes from endo and
exovesicles as well as plasma membranes and involved in a wide array of
intracellular trafficking processes such as cholesterol endocytosis, plasma membrane
repair as well as modulation of immune system functions (Boye and Nylandsted,
2016; Grewal et al., 2016; Rentero et al., 2018). Of note, only four proteins of that list,
Anxa2a, C4, C7a and Stat3, were the products of genes up-regulated in the
transcriptomic study. In agreement with the common enriched GO terms in both
studies (Figure 5C and Supplementary Figure 4B), these four proteins are associated
with immune response. Overall, *rpgrip1l*$^{-/-}$ mutants at scoliosis onset present
deregulated pathways in common with other scoliotic models such as activation of
immune response genes, but also specific deregulated pathways such a marked
activation of the Foxj1a ciliary motility program and an increase of Annexin levels.

Astrogliosis precedes scoliosis onset of rpgrip1l*$^{-/-}$ *juvenile fish

Both transcriptomic and proteomic analysis highlighted the occurrence of innate and
adaptive immune responses in *rpgrip1l*[-/-] fish. Several upregulated genes, such as
irg1l, mmp9 and anxa2, are reported to be expressed in macrophages and/or
microglia but also in epithelia or neurons (Hall et al., 2014) (Kuhlmann et al., 2014)
(Shan et al., 2015). We therefore investigated in which cell type(s) Anxa2, the most
upregulated protein in the *rpgrip1l*[-/-] brain proteome, was expressed. Immunostaining
of brain sections revealed a strong Anxa2 staining on cells lining brain ventricles,
such as SCO cells and ventral ependymal cells of the rhombencephalic ventricle
(RhV) (Fig 6 B', D') in *rpgrip1l*[-/-] adult fish, which was barely detectable in controls
(Figure 6 A', C'), and to a lesser extent on other brain regions (Supplementary Fig 5
A'-F'), suggesting a deregulation of ependymal cell homeostasis. In mutants, Anxa2
staining spread over long cellular extensions reaching the pial surface of the
rhombencephalon (Figure 6 D' and Supplementary Figure 5F'). This cell shape is
characteristic of radial glial cells, a cell type that behaves as neural and glial
progenitors and also plays astrocytic functions in zebrafish, as anamniote brains are
devoid of stellate astrocytes (Jurisch☐Yaksi et al., 2020; Lyons and Talbot, 2015). To
confirm the glial nature of Anxa2-positive cells around brain ventricles in *rpgrip1l*[-/-]
fish, we double-labeled brain sections with GFAP, a glial cell marker, and Anxa2. In
adult *rpgrip1l*[-/-] fish, SCO cells were strongly double-positive for Anxa2 and GFAP
(Fig 6 B, B'), while controls displayed much weaker labeling for GFAP in the SCO
and no labeling for Anxa2 (Fig 6 A, A'). We made similar observations at
rhombencephalic ventricle (RhV) and other brain levels: ependymal cells were
positive for Anxa2 and strongly positive for GFAP, and GFAP labeling was much
higher in mutants than in controls (Fig 6 C-D'; supplementary Figure 5A-F'). GFAP
overexpression is a hallmark of astrogliosis, a process in which astroglial cells react
to the perturbed environment, as shown during brain infection, injury, ischemia or
neurodegeneration (Escartin et al., 2021; Matusova et al., 2023; Zamanian et al.,
2012). Thus, the strong increase of GFAP staining combined with Anxa2
overexpression in brain ependymal cells of *rpgrip1l*[-/-] adult fish strongly suggests that
these cells undergo astrogliosis as observed in other animal models following CNS
injury (Chen et al., 2017; Zamanian et al., 2012).
To assay whether an astrogliosis-like phenotype was already present at scoliotic
onset, we immunostained GFAP on juvenile brain sections of straight and tail-up

mutants and controls. Tail-up juveniles (n=3/3) presented a strong GFAP staining in
almost all cells of the SCO and along the ventral rhombencephalic ventricle, higher
than control levels (Figure 6 E, I, H, L, M) while the phenotype of straight fish was
heterogeneous: two out of four juveniles presented GFAP-positive cells (Figure 6G, K
and red dots on Figure 6M graph), while the two other did not (Figure 6J and green
dots on Figure 6M graph) as confirmed by cell quantification (Figure 6M). As scoliosis
onset is asynchronous in $rpgrip1l^{-/-}$, heterogenous GFAP staining among straight
mutant fish may support an early role of astrogliosis at SCO and RhV levels in
scoliosis onset.
368
369 ***Immune cell enrichment around the SCO and within tectum parenchyma is***
370 ***detected prior to scoliosis onset.***
371
Astrogliosis may be reinforced or induced by an inflammatory environment containing
cytokines released by microglia or macrophages or the presence of high levels of
ROS and NO (Sofroniew, 2015). We thus performed immunostaining for the LCP1/L-
Plastin marker, which labels both macrophages and microglia (Figure 6N-Q'), in
adjacent sections to those used for GFAP (Figure 6E-L). We found that the scoliotic
fish presented a higher number of LCP1-positive cells of ameboid shape within SCO
cells (Figure 6Q, Q', R). Strikingly, the straight mutants (2/4) that presented a high
GFAP staining within SCO cells (Figure 6G, K) also showed a higher number of
LCP1-positive cells around the SCO (Figure 6P, P'; red dots in Figure 6R) than in
controls (Figure 6N, N'), while the other straight mutants did not ((Figure 6O, O',
green dots in Figure 6M) suggesting positive regulation between astrogliosis and
macrophages/microglia recruitment. LCP1+ cells were also detected in other regions
of the brain (Supplementary Figure S5 G-K). In the optic tectum, where resident
LCP1-positive macrophages/microglia were present in controls, a two-fold increase in
the number of ramified LCP1^{+} cells was observed in straight $rpgrip1l^{-/-}$ juveniles
compared to controls (Supplementary Figure 5H-J). Surprisingly, scoliotic $rpgrip1l^{-/-}$
fish showed similar LCP1^{+} cell number as control fish in the tectum (Supplementary
Figure S5G), suggesting a transient increase of macrophage/microglia activation
preceding scoliosis.

Our data thus uncovers an early astrogliosis process initiated asynchronously at the
SCO and RhV levels of straight *rpgrip1l*$^{-/-}$ fish, favored by a transient inflammatory
state, and preceding the loss of multiciliated tufts around the SCO and that of the RF,
that we only detected in curved *rpgrip1l*$^{-/-}$ fish. Astrogliosis then spreads along the
brain ventricles and becomes prominent in brain of scoliotic fish. In addition, we
found among the upregulated genes in dorsal trunk at scoliosis onset, a number of
activated astrocyte markers (Matusova et al., 2023) such as *s100b*, *c4b*, *gfap*,
glast/slc1a3b, *viml*, *ctsb*, (Supplementary Figure S4D), showing that astrogliosis
spread over the whole central nervous system.

Brain neuroinflammation and astrogliosis contribute to scoliosis penetrance
and severity in rpgrip1l*$^{-/-}$ *fish

Our data suggest that astrogliosis and neuro-inflammation could play a role in
triggering scoliosis in *rpgrip1l* mutants. We thus aimed to counteract inflammation
before curvature initiation by performing drug treatment with the hope of reducing
scoliosis penetrance and severity. We used the NACET anti-oxidant and anti-
inflammatory drug that successfully reduced scoliosis of *sspo* hypomorphic mutant
fish (Rose et al., 2020). We first assayed the biological activity of our NACET batch
on the ciliary motility mutant *dnaaf1*$^{-/-}$ (Sullivan-Brown et al., 2008). Indeed, NACET
was shown to suppress the embryonic tail-down phenotype of the ciliary motility
mutant *ccdc151*^{ts} (Rose et al., 2020). We found that 3mM NACET treatment
suppressed the *dnaaf1*$^{-/-}$ tail-down phenotype (Supplementary Figure S6). To
suppress scoliosis, we treated the progeny of *rpgrip1l*$^{+/-}$ incrosses from 4 to 12 wpf
with 1.5 mM NACET as in (Rose et al., 2020).This long-term treatment reduced
scoliosis penetrance from 92% to 58% and markedly reduced spinal curvature index
(Figure 7A-C), measured and illustrated as shown in Supplementary Figure 3I.
We then asked whether NACET treatment suppressed astrogliosis and
neuroinflammation in *rpgrip1l*$^{-/-}$ treated fish. Indeed, the intensity and number of
GFAP+ cells within the SCO were reduced in slightly curved fish and alleviated in
straight fish (Figure 7E, F). Furthermore, the quantification of Lcp1-positive cells in
the SCO region showed a drastic reduction of these cells in NACET-treated fish,
back to levels found in *rpgrip1l*$^{+/+}$ fish (Figure 7D, E).

Our data thus show that *rpgrip1l$^{-/-}$* juvenile fish display brain astrogliosis and
inflammation at scoliosis onset and that an anti-inflammatory/anti-oxidant treatment
of *rpgrip1l$^{-/-}$* juveniles is able to maintain a straight axis in approximately one third of
the mutants and to slow down curve progression in the other two thirds. The
beneficial effect of NACET on axis curvature correlates with the combined reduction
of astrogliosis and immune cell density.

Brain astrogliosis is present in zebrafish *cep290 $^{fh297/fh297}$* scoliotic fish

To investigate whether abnormal SCO cilia function and brain astrogliosis may be a
general mechanism of scoliosis development in transition zone mutants, we studied
the *cep290$^{fh297/fh297}$* zebrafish mutant. Zebrafish *cep290* mutants display variable axis
curvature defects at embryonic stages as well as slow retinal degeneration, and
develop juvenile scoliosis (Lessieur et al., 2019; Wang et al., 2022). Like *RPGRIP1L*,
the human *CEP290* gene encodes a TZ protein and its mutation leads to severe
neurodevelopmental ciliopathies (Baala et al., 2007). All *cep290$^{fh297/fh297}$* juveniles
developed a fully penetrant scoliosis by 4 wpf in our fish facility (n: 30/30). We
analyzed cilia presence within the SCO by immunostaining on sections (Figure 8A-
F'). At 4 wpf, the SCO had not reached its final adult width (14 versus 24 cells wide in
the anterior SCO), and each of its cells harbored a short monocilium labeled by
glutamylated tubulin (Figure 8A-C). Multicilia tufts on SCO lateral walls were not as
clearly visible as in adults (Figure 2F) but patches of glutamylated tubulin staining
were detected on both sides of the posterior SCO (Figure 8B-C'). In *cep290$^{-/-}$*,
monocilia of the SCO appeared longer than in controls (N=3/3) and pointed towards
an abnormally elongated diencephalic ventricle (Figure 8D-F). Lateral patches of
glutamylated tubulin staining were present in *cep290$^{-/-}$* posterior SCO as in controls
(Figure 8E-F'). Thus, cilia morphology is perturbed within *cep290$^{-/-}$* SCO monociliated
cells, as well as the overall morphology of the diencephalic ventricle at this level.
To test for the presence of astrogliosis, we quantified the number of SCO cells
presenting high GFAP labeling on several sections along the A/P extent of the SCO
in 3 control and 3 *cep290$^{-/-}$* brains (Figure 8G-K). 50% of SCO cells presented strong
GFAP staining in fully scoliotic juveniles, which was not detectable in controls (Figure
8H,K, N=2/3), while a lower proportion (22%) of SCO cells was GFAP positive in the

juvenile with a milder curvature (Figure 8 M N=1/3). A strong GFAP staining was also
detected in some diencephalic radial glial cells (Figure 8K). On the same sections,
immune cells were quantified by LCP1 staining. A mean of 4 LCP1 positive cells per
SCO field was found in *cep290$^{-/-}$,* versus 1 in controls (Figure 8I-L and N). As in
rpgrip1l mutants, both abnormal monocilia, astrogliosis and immune cell enrichment
are evidenced at SCO levels in *Cep290* $^{-/-}$(3/3). Furthermore, astrogliosis was
observed within radial glial cells lining the RhV at cerebellar level (Figure 8P-R) and
at anterior hindbrain level (Figure 8T-V), as shown in *rpgrip1l$^{-/-}$* scoliotic brain (Figure
6D, K, L). This indicates that astrogliosis is shared by these two zebrafish scoliotic
models, even though they develop axis curvature at different stages (3-4 wpf for
cep290$^{-/-}$ versus 5-12 wpf for *rpgrip1l$^{-/-}$*).

468
469

DISCUSSION

In this paper we dissected the mechanisms of scoliosis appearance in a zebrafish
mutant for the TZ gene *rpgrip1l*, taking advantage of its asynchronous onset to
decipher the chronology of events leading to spine torsion. Transcriptome analysis of
the brain and trunk at scoliosis onset in *rpgrip1l$^{-/-}$* fish compared to control siblings
revealed increased expression of *urp1/2*, *foxj1a* and its target genes, as well as
genes linked to immune response. Genetic experiments ruled out an involvement of
increased Urp1/2 signaling in scoliosis development. Cilia defects appeared
asynchronously in different CNS regions, while RF loss strictly coincided with
scoliosis onset. Strikingly, we observed a strong increase of GFAP staining along the
diencephalic and rhombencephalic ventricles, indicative of astrogliosis. This
astrogliosis preceded scoliosis onset and then spread posteriorly and increased in
severity. In the region of the SCO, the appearance of astrogliosis and
neuroinflammation coincided temporally, preceding the onset of RF depolymerization
and spine torsion. NACET treatment showed that increased astrogliosis and
inflammation participated in the penetrance and severity of scoliosis in *rpgrip1l$^{-/-}$* fish.
Finally, we showed that another TZ gene mutant, *cep290,* also displayed astrogliosis
at the time of scoliosis onset. Thus, the cascade of events uncovered in the *rpgrip1l*

mutant sheds a new light on the origin of zebrafish scoliosis caused by ciliary
transition zone defects.

Cilia motility defects lead to RF loss and to scoliosis in several zebrafish mutants
(Grimes et al., 2016; Rose et al., 2020)). *Rpgrip1l* encodes a TZ protein important for
cilia integrity and function in many eukaryotic organisms (Gogendeau et al., 2020;
Jensen et al., 2015; Wiegering et al., 2018). Our data show that *rpgrip1l* deficiency in
zebrafish affects cilia maintenance differentially depending on the CNS territory and
the developmental stage, raising the question of which ciliated cells are responsible
for RF loss and scoliosis initiation in this mutant. The full rescue observed after re-
introducing RPGRIP1L in *foxj1a*-expressing cells points towards a crucial function of
the protein in cells lining CNS cavities. Foxj1a-positive cell populations encompass
both monociliated and multiciliated ependymal cells, radial glial progenitors, glial cells
of the SCO and the ChP, as well as CSF-contacting sensory neurons in the spinal
cord (D'Gama et al., 2021; Prendergast et al., 2023; Ribeiro et al., 2017; Van Gennip
et al., 2018). Recent work suggested that impairment of ciliogenesis in the fChP of
the *katnb1* mutant could play a role in inducing axis curvature (Meyer-Miner et al.,
2022). In the case of the *rpgrip1l* mutant, we temporally uncorrelated fChP cilia
defects from scoliosis onset. This, combined with the fact that the total loss of
multicilia in the fChP of ($foxj1b^{-/-}$, $gmnc^{-/-}$) double mutants does not trigger axis
curvature (D'Gama et al., 2021) suggests that another *foxj1a*-expressing population
is crucial for axis straightness. Our results strongly suggest that cilia defects in the
SCO trigger axis curvature in *rpgrip1l* mutants. Two cilia populations were affected in
the SCO of *rpgrip1l* mutant juveniles at the onset of scoliosis: cilia of monociliated
glial cells that secrete SCO-spondin, which appeared longer in mutants than in
controls, and multiciliated tufts at SCO exit, which were lost in scoliotic mutants. Both
defects could be responsible for the phenotype, by perturbing CSF flow and content
(presence of SCO-spondin aggregates in ventricles) (left side on the Graphical
Abstract) and leading to ventricular dilations or by perturbing primary cilia signaling
within SCO cells (right side of the Graphical Abstract). The observation that
multiciliated cells were not yet fully differentiated in *cep290* mutants at scoliosis onset
is in favor of a prominent role of monocilia. This is consistent with the literature
showing that most ciliary mutant models develop axis curvature at 3-4 weeks, a

stage when brain multiciliated cells are not yet differentiated (D'Gama et al., 2021).

Still, as scoliosis onset in *rpgrip1l* mutant is asynchronous from 5 to 12 weeks, MCC

loss at SCO exit could contribute to defective RF polymerization.

Our model for scoliosis appearance in the rpgrip1l mutant is presented in the

graphical abstract. Since we have observed several straight *rpgrip1l* [-/-] with

astrogliosis and increased immune cells presence around the SCO, we propose that

rpgrip1l [-/-] SCO and astroglial cells experience an altered signalling that activates

immune cells recruitment, producing an inflammatory environment. This would

promote multicilia loss as a secondary event (Graphical Abstract, orange arrow)

(Lattke et al., 2012) leading to RF depolymerization and spine curvature. However,

as *sspo* [-/-] hypomorphic mutants which present aggregates within ventricles have

been shown to present an inflammatory response (Rose et al., 2020), it is equally

possible that SCO astrogliosis is a downstream consequence of the presence of

these proteoglycans aggregates in contact with primary cilia or ventricular cell

surfaces (Graphical Abstract , green arrow). These two scenarios may also act in

parallel to reinforce scoliosis onset and ependymal cilia loss. Future experiments will

help to discriminate between these two scenarios.

Our transcriptome analysis and qRT-PCR data challenges a model proposed for axis

straightness of zebrafish embryos in which the loss of RF leads to a down-regulation

of *urp2* and *urp1* expression (Zhang et al., 2018) since we observed an upregulation

of both genes before and after RF loss. As forced expression of *urp1/2* induces a tail-

up phenotype in embryos and juveniles (Zhang et al., 2018) (Gaillard et al., 2023; Lu

et al., 2020), we attempted to rescue *rpgrip1l*[-/-] axis curvature by down-regulating

urp1/2 expression. No beneficial effect on scoliosis penetrance or severity was

observed, indicating that increased *urp1/2* expression level does not significantly

contribute to axis curvature in *rpgrip1l*[-/-] fish.

In this study, we found an unexpected and strong defect characterized by enhanced

GFAP and Anxa2 expression, which we identified as astrogliosis (also called reactive

astrogliosis). Astrogliosis is a reaction of astroglial cells to perturbed homeostasis of

the CNS, characterized by molecular and phenotypic changes in these cells (Escartin

et al., 2021; Matusova et al., 2023). In zebrafish, radial glial cells are endowed with
both neurogenic and astrocytic functions, and are thus also called astroglial cells
(Jurisch☐Yaksi et al., 2020). Astrogliosis in *rpgrip1l* mutants arose in a subdomain of
Foxj1a-positive cells, within the DiV and ventral ependymal cells lining the RhV. The
observation of astrogliosis in some straight juvenile mutants suggests an early role of
this defect in scoliosis onset.

Astrogliosis along the ventricular cavities could arise in response to local mechanical
or chemical perturbations caused by cilia motility defects, abnormal CSF flow and
content and ventricular dilations. In two murine models of primary ciliary dyskinesia,
decreased CSF flow was associated with gliosis at juvenile stage (Finn et al., 2014).
A response to CSF pertubation in *rpgrip1l* mutants is also suggested by the strong
upregulation of Foxj1a and its target genes. *foxj1a* expression is also reported to be
upregulated in the CNS of three other scoliotic models in addition to *rpgrip1l*, namely
in *ptk7, sspo* and *katnb1* mutants (Escartin et al., 2021; Meyer-Miner et al., 2022;
Van Gennip et al., 2018), suggesting that it constitutes a common response to similar
ventricular and CSF defects. Interestingly, *foxj1a* expression is also upregulated
upon zebrafish embryonic (Cavone et al., 2021; Hellman et al., 2010) or adult
(Ribeiro et al., 2017) spinal cord injury. The up-regulation in the *rpgrip1l* mutant
proteome of the membrane repair module Anxa2-S100a10-Ahnak (Bharadwaj et al.,
2021) (Figure 4C) together with the Foxj1-induced motility module detected in the
transcriptomic analysis may indicate ongoing reparation attempts of CNS ventricles
and associated neural tissues.

Moreover, the presence of few Lcp1+ cells around the mutant SCO suggests that
microglia/macrophages could participate in astrogliosis establishment and
reinforcement. This dialogue between immune Lcp1+ cells and astroglial cells is
evidenced during early stages of regeneration after acute injury or cellular damages
(Cavone et al., 2021). In our bulk RNAseq analysis performed at scoliosis onset, we
have indications of the upregulation of microglia and macrophage markers (*p2ry12*
and *mpeg1*) at trunk levels and of numerous markers of activated astrocytes
characterized in several mammalian astrogliosis models (Matusova et al., 2023). A
similar dialogue is detected in polycystic kidney disease models where compromised
primary cilia signalling leads to uncontrolled cytokines secretion by epithelial cells, a

situation that favors local immune cells recruitment and proliferation (Viau et al.,
2018).

What are the intracellular mechanisms involved in astrogliosis induction in ciliated
ventricular cells? Candidate pathways emerge from our multi-omics studies.
Proteomic data showed a 2.4-fold increase in GSK3bb amount in *rpgrip1l*$^{-/-}$ mutant
brains (P value < 0.001, Supplementary Table 3). In several neurodegenerative
murine animal models, GSK3b enzymatic activity was shown to promote
inflammation and gliosis (Jorge-Torres et al., 2018; Medina and Avila, 2010; Mines et
al., 2011). Another potential trigger of astrogliosis may be a defective mitochondrial
activity as the amounts of four proteins involved in electron transport chain activity
(*gpd1b, mt-nd6, pdia5 and dmgdh*) were reduced by 40% to 86% in the analysis of
mutant brain proteome (Supplementary Table 4). We think this is of particular interest
in light of a recent report demonstrating that impaired mitochondrial activity within
ciliated astrocytes leads to the induction of the Foxj1 ciliary motility program, the
elongation and distortion of astrocyte primary cilia as well as reactive astrogliosis
(Ignatenko et al., 2023), three phenotypes also observed in *rpgrip1l*$^{-/-}$ brains.

Finally, our observation of widespread astrogliosis in two TZ gene mutants, *rpgrip1l*
and *cep290*, suggests that it could represent a general mechanism involved in
scoliosis downstream of cilia dysfunction. Of note, astrogliosis markers such as
glast/slc1a3b, *vim*, *s100b*, *c4*, *timp2b*, *gfap* and *ctssb.2*, are also upregulated in the
transcriptome of *ptk7 mutants* (Van Gennip et al., 2018). We propose that sustained
astrogliosis might impair neuronal survival and activity, crucial for proper
interoception and locomotor control, thus leading to axis curvature in the context of a
rapidly growing axial skeleton. A similar context of astrocytosis associated with spinal
cord dilation has been observed in human patients with syringomyelia. This
population with a high incidence of scoliosis can present hyper-signals on T2 MRI
scans, which were proposed to reflect local astrocytosis and could be validated by
biopsies in rare cases (Sherman et al., 1987). Further imaging studies will need to be
performed to validate a potential link between developing astrogliosis and the onset
and progression of idiopathic scoliosis in humans.

ACKNOWLEDGEMENTS

We are grateful to the IBPS aquatic animal, imaging and bioinformatics facilities and

to the ICM sequencing facility for their technical assistance. We thank Michaeｌl

Trichet from the IBPS electron microscopy facility for participating in the MEB

experiments, the TACGENE facility for providing the Cas9 protein; the TEFOR Paris-

Saclay facility for the brain clearing experiment; Thierry Jaffredo and Pierre Charbord

for their precious help in transcriptome analysis; Martin Catala for helpful discussions

and documentation on patients with Chiari malformation presenting syringomyelia,

Nicolas Baylé for initial characterization of the *rpgrip1l* mutant; Claudia Hoffman for

providing *Cep290* fixed samples; Brian Ciruna for sharing the *foxj1a* expressing

constructs and detailed NACET treatment protocol; Ryan Gray for sharing the

scospondin-GFP^{ut24} zebrafish line. This work was supported by funding to SSM from

the Fondation pour la Recherche Médicale (Equipe FRM EQU201903007943) and

the Fondation Yves Cotrel.

REFERENCES

Affaticati P, Simion M, De Job E, Rivière L, Hermel J-M, Machado E, Joly J-S, Jenett A. 2017. zPACT: Tissue Clearing and Immunohistochemistry on Juvenile Zebrafish Brain. *BIO-PROTOCOL* **7**. doi:10.21769/BioProtoc.2636

Andreu-Cervera A, Anselme I, Karam A, Laclef C, Catala M, Schneider-Maunoury S. 2019. The ciliopathy gene *ftm/rpgrip1l* controls mouse forebrain patterning via region-specific modulation of hedgehog/gli signaling. *J Neurosci* 2199–18. doi:10.1523/JNEUROSCI.2199-18.2019

Andreu-Cervera A, Catala M, Schneider-Maunoury S. 2021. Cilia, ciliopathies and hedgehog-related forebrain developmental disorders. *Neurobiology of Disease* **150**:105236. doi:10.1016/j.nbd.2020.105236

Arts HH, Doherty D, Van Beersum SEC, Parisi MA, Letteboer SJF, Gorden NT, Peters TA, Märker T, Voesenek K, Kartono A, Ozyurek H, Farin FM, Kroes HY, Wolfrum U, Brunner HG, Cremers FPM, Glass IA, Knoers NVAM, Roepman R. 2007. Mutations in the gene encoding the basal body protein RPGRIP1L, a nephrocystin-4 interactor, cause Joubert syndrome. *Nat Genet* **39**:882–888. doi:10.1038/ng2069

Baala L, Audollent S, Martinovic J, Ozilou C, Babron M-C, Sivanandamoorthy S, Saunier S, Salomon R, Gonzales M, Rattenberry E, Esculpavit C, Toutain A, Moraine C, Parent P, Marcorelles P, Dauge M-C, Roume J, Merrer ML, Meiner V, Meir K, Menez F, Beaufrère A-M, Francannet C, Tantau J, Sinico M, Dumez Y, MacDonald F, Munnich A, Lyonnet S, Gubler M-C, Génin E, Johnson CA, Vekemans M, Encha-Razavi F, Attié-Bitach T. 2007. Pleiotropic Effects of CEP290 (NPHP6) Mutations Extend to Meckel Syndrome. *The American Journal of Human Genetics* **81**:170–179. doi:10.1086/519494

Bachmann-Gagescu R, Phelps IG, Stearns G, Link BA, Brockerhoff SE, Moens CB, Doherty D. 2011. The ciliopathy gene cc2d2a controls zebrafish photoreceptor outer segment development through a role in Rab8-dependent vesicle trafficking. *Human Molecular Genetics* **20**:4041–4055. doi:10.1093/hmg/ddr332

Bearce EA, Irons ZH, O'Hara-Smith JR, Kuhns CJ, Fisher SI, Crow WE, Grimes DT. 2022. Urotensin II-related peptides, Urp1 and Urp2, control zebrafish spine morphology. *eLife* **11**:e83883. doi:10.7554/eLife.83883

Bentley-Ford MR, LaBonty M, Thomas HR, Haycraft CJ, Scott M, LaFayette C, Croyle MJ, Andersen RS, Parant JM, Yoder BK. 2022. Evolutionarily conserved genetic interactions between *nphp-4* and *bbs-5* mutations exacerbate ciliopathy phenotypes. *Genetics* **220**:iyab209. doi:10.1093/genetics/iyab209

Bharadwaj A, Kempster E, Waisman DM. 2021. The Annexin A2/S100A10 Complex: The Mutualistic Symbiosis of Two Distinct Proteins. *Biomolecules* **11**:1849. doi:10.3390/biom11121849

Boswell CW, Ciruna B. 2017. Understanding Idiopathic Scoliosis: A New Zebrafish School of Thought. *Trends in Genetics* **33**:183–196. doi:10.1016/j.tig.2017.01.001

Bouyssié D, Hesse A-M, Mouton-Barbosa E, Rompais M, Macron C, Carapito C, Gonzalez De Peredo A, Couté Y, Dupierris V, Burel A, Menetrey J-P, Kalaitzakis A, Poisat J, Romdhani A, Burlet-Schiltz O, Cianférani S, Garin J, Bruley C. 2020. Proline: an efficient and user-friendly software suite for large-

scale proteomics. *Bioinformatics* **36**:3148–3155. doi:10.1093/bioinformatics/btaa118

Boye TL, Nylandsted J. 2016. Annexins in plasma membrane repair. *Biological Chemistry* **397**:961–969. doi:10.1515/hsz-2016-0171

Brancati F, Travaglini L, Zablocka D, Boltshauser E, Accorsi P, Montagna G, Silhavy JL, Barrano G, Bertini E, Emma F, Rigoli L, International JSRD Study Group, Dallapiccola B, Gleeson JG, Valente EM. 2008. RPGRIP1L mutations are mainly associated with the cerebello-renal phenotype of Joubert syndrome-related disorders. *Clin Genet* **74**:164–170. doi:10.1111/j.1399-0004.2008.01047.x

Cantaut-Belarif Y, Orts Del'Immagine A, Penru M, Pézeron G, Wyart C, Bardet P-L. 2020. Adrenergic activation modulates the signal from the Reissner fiber to cerebrospinal fluid-contacting neurons during development. *eLife* **9**:e59469. doi:10.7554/eLife.59469

Cantaut-Belarif Y, Sternberg JR, Thouvenin O, Wyart C, Bardet P-L. 2018. The Reissner Fiber in the Cerebrospinal Fluid Controls Morphogenesis of the Body Axis. *Current Biology* **28**:2479-2486.e4. doi:10.1016/j.cub.2018.05.079

Cavone L, McCann T, Drake LK, Aguzzi EA, Oprişoreanu A-M, Pedersen E, Sandi S, Selvarajah J, Tsarouchas TM, Wehner D, Keatinge M, Mysiak KS, Henderson BEP, Dobie R, Henderson NC, Becker T, Becker CG. 2021. A unique macrophage subpopulation signals directly to progenitor cells to promote regenerative neurogenesis in the zebrafish spinal cord. *Developmental Cell* **56**:1617-1630.e6. doi:10.1016/j.devcel.2021.04.031

Chen J, Cui Z, Yang S, Wu C, Li W, Bao G, Xu G, Sun Y, Wang L, Zhang J. 2017. The upregulation of annexin A2 after spinal cord injury in rats may have implication for astrocyte proliferation. *Neuropeptides* **61**:67–76. doi:10.1016/j.npep.2016.10.007

Choksi SP, Lauter G, Swoboda P, Roy S. 2014. Switching on cilia: transcriptional networks regulating ciliogenesis. *Development* **141**:1427–1441. doi:10.1242/dev.074666

Collins P, Woollam DH. 1979. The ventricular surface of the subcommissural organ: a scanning and transmission electron microscopic study. *J Anat* **129**:623–631.

Dale RM, Topczewski J. 2011. Identification of an evolutionarily conserved regulatory element of the zebrafish col2a1a gene. *Developmental Biology* **357**:518–531. doi:10.1016/j.ydbio.2011.06.020

Delous M, Baala L, Salomon R, Laclef C, Vierkotten J, Tory K, Golzio C, Lacoste T, Besse L, Ozilou C, Moutkine I, Hellman NE, Anselme I, Silbermann F, Vesque C, Gerhardt C, Rattenberry E, Wolf MTF, Gubler MC, Martinovic J, Encha-Razavi F, Boddaert N, Gonzales M, Macher MA, Nivet H, Champion G, Berthélémé JP, Niaudet P, McDonald F, Hildebrandt F, Johnson CA, Vekemans M, Antignac C, Rüther U, Schneider-Maunoury S, Attié-Bitach T, Saunier S. 2007. The ciliary gene RPGRIP1L is mutated in cerebello-oculo-renal syndrome (Joubert syndrome type B) and Meckel syndrome. *Nat Genet* **39**:875–881. doi:10.1038/ng2039

D'Gama PP, Qiu T, Cosacak MI, Rayamajhi D, Konac A, Hansen JN, Ringers C, Acuña-Hinrichsen F, Hui SP, Olstad EW, Chong YL, Lim CKA, Gupta A, Ng CP, Nilges BS, Kashikar ND, Wachten D, Liebl D, Kikuchi K, Kizil C, Yaksi E, Roy S, Jurisch-Yaksi N. 2021. Diversity and function of motile ciliated cell types within ependymal lineages of the zebrafish brain. *Cell Reports* **37**:109775. doi:10.1016/j.celrep.2021.109775

Didier R, Dastugue B, Meiniel A. 1995. The secretory material of the subcommissural organ of the chick embryo. Characterization of a specific polypeptide by two-dimensional electrophoresis. *Int J Dev Biol* **39**:493–499.

Escartin C, Galea E, Lakatos A, O'Callaghan JP, Petzold GC, Serrano-Pozo A, Steinhäuser C, Volterra A, Carmignoto G, Agarwal A, Allen NJ, Araque A, Barbeito L, Barzilai A, Bergles DE, Bonvento G, Butt AM, Chen W-T, Cohen-Salmon M, Cunningham C, Deneen B, De Strooper B, Díaz-Castro B, Farina C, Freeman M, Gallo V, Goldman JE, Goldman SA, Götz M, Gutiérrez A, Haydon PG, Heiland DH, Hol EM, Holt MG, Iino M, Kastanenka KV, Kettenmann H, Khakh BS, Koizumi S, Lee CJ, Liddelow SA, MacVicar BA, Magistretti P, Messing A, Mishra A, Molofsky AV, Murai KK, Norris CM, Okada S, Oliet SHR, Oliveira JF, Panatier A, Parpura V, Pekna M, Pekny M, Pellerin L, Perea G, Pérez-Nievas BG, Pfrieger FW, Poskanzer KE, Quintana FJ, Ransohoff RM, Riquelme-Perez M, Robel S, Rose CR, Rothstein JD, Rouach N, Rowitch DH, Semyanov A, Sirko S, Sontheimer H, Swanson RA, Vitorica J, Wanner I-B, Wood LB, Wu J, Zheng B, Zimmer ER, Zorec R, Sofroniew MV, Verkhratsky A. 2021. Reactive astrocyte nomenclature, definitions, and future directions. *Nat Neurosci* **24**:312–325. doi:10.1038/s41593-020-00783-4

Finn R, Evans CC, Lee L. 2014. Strain-dependent brain defects in mouse models of primary ciliary dyskinesia with mutations in Pcdp1 and Spef2. *Neuroscience* **277**:552–567. doi:10.1016/j.neuroscience.2014.07.029

Gaillard A-L, Mohamad T, Quan FB, De Cian A, Mosimann C, Tostivint H, Pézeron G. 2023. Urp1 and Urp2 act redundantly to maintain spine shape in zebrafish larvae. *Developmental Biology* **496**:36–51. doi:10.1016/j.ydbio.2023.01.010

Gogendeau D, Lemullois M, Le Borgne P, Castelli M, Aubusson-Fleury A, Arnaiz O, Cohen J, Vesque C, Schneider-Maunoury S, Bouhouche K, Koll F, Tassin A-M. 2020. MKS-NPHP module proteins control ciliary shedding at the transition zone. *PLoS Biol* **18**:e3000640. doi:10.1371/journal.pbio.3000640

Grewal T, Wason SJ, Enrich C, Rentero C. 2016. Annexins – insights from knockout mice. *Biological Chemistry* **397**:1031–1053. doi:10.1515/hsz-2016-0168

Grimes DT, Boswell CW, Morante NFC, Henkelman RM, Burdine RD, Ciruna B. 2016. Zebrafish models of idiopathic scoliosis link cerebrospinal fluid flow defects to spine curvature. *Science* **352**:1341–1344. doi:10.1126/science.aaf6419

Haeussler M, Schönig K, Eckert H, Eschstruth A, Mianné J, Renaud J-B, Schneider-Maunoury S, Shkumatava A, Teboul L, Kent J, Joly J-S, Concordet J-P. 2016. Evaluation of off-target and on-target scoring algorithms and integration into the guide RNA selection tool CRISPOR. *Genome Biol* **17**:148. doi:10.1186/s13059-016-1012-2

Hall CJ, Boyle RH, Sun X, Wicker SM, Misa JP, Krissansen GW, Print CG, Crosier KE, Crosier PS. 2014. Epidermal cells help coordinate leukocyte migration during inflammation through fatty acid-fuelled matrix metalloproteinase production. *Nat Commun* **5**:3880. doi:10.1038/ncomms4880

Hayes M, Gao X, Yu LX, Paria N, Henkelman RM, Wise CA, Ciruna B. 2014. ptk7 mutant zebrafish models of congenital and idiopathic scoliosis implicate dysregulated Wnt signalling in disease. *Nat Commun* **5**:4777. doi:10.1038/ncomms5777

Hellman NE, Liu Y, Merkel E, Austin C, Le Corre S, Beier DR, Sun Z, Sharma N, Yoder BK, Drummond IA. 2010. The zebrafish *foxj1a* transcription factor

781   regulates cilia function in response to injury and epithelial stretch. *Proc Natl*
782   *Acad Sci USA* **107**:18499–18504. doi:10.1073/pnas.1005998107
783 Hitier M, Hamon M, Denise P, Lacoudre J, Thenint M-A, Mallet J-F, Moreau S,
784   Quarck G. 2015. Lateral Semicircular Canal Asymmetry in Idiopathic Scoliosis:
785   An Early Link between Biomechanical, Hormonal and Neurosensory
786   Theories? *PLoS ONE* **10**:e0131120. doi:10.1371/journal.pone.0131120
787 Hughes CS, Foehr S, Garfield DA, Furlong EE, Steinmetz LM, Krijgsveld J. 2014.
788   Ultrasensitive proteome analysis using paramagnetic bead technology.
789   *Molecular Systems Biology* **10**:757. doi:10.15252/msb.20145625
790 Ignatenko O, Malinen S, Rybas S, Vihinen H, Nikkanen J, Kononov A, Jokitalo ES,
791   Ince-Dunn G, Suomalainen A. 2023. Mitochondrial dysfunction compromises
792   ciliary homeostasis in astrocytes. *Journal of Cell Biology* **222**:e202203019.
793   doi:10.1083/jcb.202203019
794 Jensen VL, Li C, Bowie RV, Clarke L, Mohan S, Blacque OE, Leroux MR. 2015.
795   Formation of the transition zone by Mks5/Rpgrip1L establishes a ciliary zone
796   of exclusion (CIZE) that compartmentalises ciliary signalling proteins and
797   controls PIP $_2$ ciliary abundance. *The EMBO Journal* **34**:2537–2556.
798   doi:10.15252/embj.201488044
799 Jorge-Torres OC, Szczesna K, Roa L, Casal C, Gonzalez-Somermeyer L, Soler M,
800   Velasco CD, Martínez-San Segundo P, Petazzi P, Sáez MA, Delgado-Morales
801   R, Fourcade S, Pujol A, Huertas D, Llobet A, Guil S, Esteller M. 2018.
802   Inhibition of Gsk3b Reduces Nfkb1 Signaling and Rescues Synaptic Activity to
803   Improve the Rett Syndrome Phenotype in Mecp2-Knockout Mice. *Cell Reports*
804   **23**:1665–1677. doi:10.1016/j.celrep.2018.04.010
805 Jurisch-Yaksi N, Yaksi E, Kizil C. 2020. Radial glia in the zebrafish brain: Functional,
806   structural, and physiological comparison with the mammalian glia. *Glia*
807   **68**:2451–2470. doi:10.1002/glia.23849
808 Karner CM, Long F, Solnica-Krezel L, Monk KR, Gray RS. 2015. *Gpr126/Adgrg6*
809   deletion in cartilage models idiopathic scoliosis and pectus excavatum in mice.
810   *Hum Mol Genet* **24**:4365–4373. doi:10.1093/hmg/ddv170
811 Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of
812   embryonic development of the zebrafish. *Developmental Dynamics* **203**:253–
813   310. doi:10.1002/aja.1002030302
814 Kramer-Zucker AG, Olale F, Haycraft CJ, Yoder BK, Schier AF, Drummond IA. 2005.
815   Cilia-driven fluid flow in the zebrafish pronephros, brain and Kupffer's vesicle
816   is required for normal organogenesis. *Development* **132**:1907–1921.
817   doi:10.1242/dev.01772
818 Kuhlmann K, Tschapek A, Wiese H, Eisenacher M, Meyer HE, Hatt HH, Oeljeklaus
819   S, Warscheid B. 2014. The Membrane Proteome of Sensory Cilia to the Depth
820   of Olfactory Receptors. *Molecular & Cellular Proteomics* **13**:1828–1843.
821   doi:10.1074/mcp.M113.035378
822 Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, Parant JM,
823   Yost HJ, Kanki JP, Chien C. 2007. The Tol2kit: A multisite gateway-based
824   construction kit for *Tol2* transposon transgenesis constructs. *Developmental*
825   *Dynamics* **236**:3088–3099. doi:10.1002/dvdy.21343
826 Lattke M, Magnutzki A, Walther P, Baumann B. 2012. Nuclear Factor κB Activation
827   Impairs Ependymal Ciliogenesis and Links Neuroinflammation to
828   Hydrocephalus Formation. *J Neurosci* **32**:11511–11523.
829   doi:10.1523/JNEUROSCI.0182-12.2012

Lessieur EM, Song P, Nivar GC, Piccillo EM, Fogerty J, Rozic R, Perkins BD. 2019.
Ciliary genes arl13b, ahi1 and cc2d2a differentially modify expression of visual
acuity phenotypes but do not enhance retinal degeneration due to mutation of
cep290 in zebrafish. *PLoS One* **14**:e0213960.
doi:10.1371/journal.pone.0213960
Lu H, Shagirova A, Goggi JL, Yeo HL, Roy S. 2020. Reissner fibre-induced urotensin
signalling from cerebrospinal fluid-contacting neurons prevents scoliosis of the
vertebrate spine. *Biology Open* **9**:bio052027. doi:10.1242/bio.052027
Lyons DA, Talbot WS. 2015. Glial Cell Development and Function in Zebrafish. *Cold
Spring Harb Perspect Biol* **7**:a020586. doi:10.1101/cshperspect.a020586
Mahuzier A, Gaudé H-M, Grampa V, Anselme I, Silbermann F, Leroux-Berger M,
Delacour D, Ezan J, Montcouquiol M, Saunier S, Schneider-Maunoury S,
Vesque C. 2012. Dishevelled stabilization by the ciliopathy protein Rpgrip1l is
essential for planar cell polarity. *Journal of Cell Biology* **198**:927–940.
doi:10.1083/jcb.201111009
Mangos S, Lam P, Zhao A, Liu Y, Mudumana S, Vasilyev A, Liu A, Drummond IA.
2010. The ADPKD genes *pkd1a/b* and *pkd2* regulate extracellular matrix
formation. *Disease Models & Mechanisms* **3**:354–365.
doi:10.1242/dmm.003194
Masek M, Etard C, Hofmann C, Hülsmeier AJ, Zang J, Takamiya M, Gesemann M,
Neuhauss SCF, Hornemann T, Strähle U, Bachmann-Gagescu R. 2022. Loss
of the Bardet-Biedl protein Bbs1 alters photoreceptor outer segment protein
and lipid composition. *Nat Commun* **13**:1282. doi:10.1038/s41467-022-28982-
6
Matusova Z, Hol EM, Pekny M, Kubista M, Valihrach L. 2023. Reactive astrogliosis in
the era of single-cell transcriptomics. *Front Cell Neurosci* **17**:1173200.
doi:10.3389/fncel.2023.1173200
Medina M, Avila J. 2010. Glycogen Synthase Kinase-3 (GSK-3) Inhibitors for the
Treatment of Alzheimers Disease. *CPD* **16**:2790–2798.
doi:10.2174/138161210793176581
Meyer-Miner A, Van Gennip JLM, Henke K, Harris MP, Ciruna B. 2022. Resolving
primary pathomechanisms driving idiopathic-like spinal curvature using a new
katnb1 scoliosis model. *iScience* **25**:105028. doi:10.1016/j.isci.2022.105028
Mines MA, Beurel E, Jope RS. 2011. Regulation of Cell Survival Mechanisms in
Alzheimer's Disease by Glycogen Synthase Kinase-3. *International Journal of
Alzheimer's Disease* **2011**:1–11. doi:10.4061/2011/861072
Mukherjee I, Roy S, Chakrabarti S. 2019. Identification of Important Effector Proteins
in the FOXJ1 Transcriptional Network Associated With Ciliogenesis and Ciliary
Function. *Front Genet* **10**:23. doi:10.3389/fgene.2019.00023
Olstad EW, Ringers C, Hansen JN, Wens A, Brandt C, Wachten D, Yaksi E, Jurisch-
Yaksi N. 2019. Ciliary Beating Compartmentalizes Cerebrospinal Fluid Flow in
the Brain and Regulates Ventricular Development. *Current Biology* **29**:229-
241.e6. doi:10.1016/j.cub.2018.11.059
Parmentier C, Hameury E, Dubessy C, Quan FB, Habert D, Calas A, Vaudry H,
Lihrmann I, Tostivint H. 2011. Occurrence of Two Distinct Urotensin II-Related
Peptides in Zebrafish Provides New Insight into the Evolutionary History of the
Urotensin II Gene Family. *Endocrinology* **152**:2330–2341.
doi:10.1210/en.2010-1500
Patten SA, Margaritte-Jeannin P, Bernard J-C, Alix E, Labalme A, Besson A, Girard
SL, Fendri K, Fraisse N, Biot B, Poizat C, Campan-Fournier A, Abelin-

880     Genevois K, Cunin V, Zaouter C, Liao M, Lamy R, Lesca G, Menassa R,
881     Marcaillou C, Letexier M, Sanlaville D, Berard J, Rouleau GA, Clerget-
882     Darpoux F, Drapeau P, Moldovan F, Edery P. 2015. Functional variants of
883     POC5 identified in patients with idiopathic scoliosis. *J Clin Invest* **125**:1124–
884     1128. doi:10.1172/JCI77262
885 Perez-Riverol Y, Bai J, Bandla C, García-Seisdedos D, Hewapathirana S,
886     Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M,
887     Walzer M, Wang S, Brazma A, Vizcaíno JA. 2022. The PRIDE database
888     resources in 2022: a hub for mass spectrometry-based proteomics evidences.
889     *Nucleic Acids Research* **50**:D543–D552. doi:10.1093/nar/gkab1038
890 Prendergast AE, Jim KK, Marnas H, Desban L, Quan FB, Djenoune L, Laghi V,
891     Hocquemiller A, Lunsford ET, Roussel J, Keiser L, Lejeune F-X, Dhanasekar
892     M, Bardet P-L, Levraud J-P, Van De Beek D, Vandenbroucke-Grauls CMJE,
893     Wyart C. 2023. CSF-contacting neurons respond to Streptococcus
894     pneumoniae and promote host survival during central nervous system
895     infection. *Current Biology* **33**:940-956.e10. doi:10.1016/j.cub.2023.01.039
896 Quan FB, Dubessy C, Galant S, Kenigfest NB, Djenoune L, Leprince J, Wyart C,
897     Lihrmann I, Tostivint H. 2015. Comparative Distribution and In Vitro Activities
898     of the Urotensin II-Related Peptides URP1 and URP2 in Zebrafish: Evidence
899     for Their Colocalization in Spinal Cerebrospinal Fluid-Contacting Neurons.
900     *PLoS ONE* **10**:e0119290. doi:10.1371/journal.pone.0119290
901 Quigley IK, Kintner C. 2017. Rfx2 Stabilizes Foxj1 Binding at Chromatin Loops to
902     Enable Multiciliated Cell Gene Expression. *PLoS Genet* **13**:e1006538.
903     doi:10.1371/journal.pgen.1006538
904 Rentero C, Blanco-Muñoz P, Meneses-Salas E, Grewal T, Enrich C. 2018.
905     Annexins—Coordinators of Cholesterol Homeostasis in Endocytic Pathways.
906     *IJMS* **19**:1444. doi:10.3390/ijms19051444
907 Ribeiro A, Monteiro JF, Certal AC, Cristovão AM, Saúde L. 2017. Foxj1a is
908     expressed in ependymal precursors, controls central canal position and is
909     activated in new ependymal cells during regeneration in zebrafish. *Open Biol*
910     **7**:170139. doi:10.1098/rsob.170139
911 Rose CD, Pompili D, Henke K, Van Gennip JLM, Meyer-Miner A, Rana R, Gobron S,
912     Harris MP, Nitz M, Ciruna B. 2020. SCO-Spondin Defects and
913     Neuroinflammation Are Conserved Mechanisms Driving Spinal Deformity
914     across Genetic Models of Idiopathic Scoliosis. *Current Biology* **30**:2363-
915     2373.e6. doi:10.1016/j.cub.2020.04.020
916 Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch
917     S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V,
918     Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for
919     biological-image analysis. *Nat Methods* **9**:676–682. doi:10.1038/nmeth.2019
920 Shan Y, Fang C, Cheng C, Wang Y, Peng J, Fang W. 2015. Immersion infection of
921     germ-free zebrafish with Listeria monocytogenes induces transient expression
922     of innate immune response genes. *Front Microbiol* **6**.
923     doi:10.3389/fmicb.2015.00373
924 Sherman J, Barkovich A, Citrin C. 1987. The MR appearance of syringomyelia: new
925     observations. *American Journal of Roentgenology* **148**:381–391.
926     doi:10.2214/ajr.148.2.381
927 Sofroniew MV. 2015. Astrogliosis. *Cold Spring Harb Perspect Biol* **7**:a020420.
928     doi:10.1101/cshperspect.a020420

Song P, Fogerty J, Cianciolo LT, Stupay R, Perkins BD. 2020. Cone Photoreceptor
Degeneration and Neuroinflammation in the Zebrafish Bardet-Biedl Syndrome
2 (bbs2) Mutant Does Not Lead to Retinal Regeneration. *Front Cell Dev Biol*
8:578528. doi:10.3389/fcell.2020.578528
Sullivan-Brown J, Schottenfeld J, Okabe N, Hostetter CL, Serluca FC, Thiberge SY,
Burdine RD. 2008. Zebrafish mutations affecting cilia motility share similar
cystic phenotypes and suggest a mechanism of cyst formation that differs from
pkd2 morphants. *Developmental Biology* **314**:261–275.
doi:10.1016/j.ydbio.2007.11.025
Troutwine BR, Gontarz P, Konjikusic MJ, Minowa R, Monstad-Rios A, Sepich DS,
Kwon RY, Solnica-Krezel L, Gray RS. 2020. The Reissner Fiber Is Highly
Dynamic In Vivo and Controls Morphogenesis of the Spine. *Current Biology*
30:2353-2362.e3. doi:10.1016/j.cub.2020.04.015
Tsujikawa M, Malicki J. 2004. Intraflagellar Transport Genes Are Essential for
Differentiation and Survival of Vertebrate Sensory Neurons. *Neuron* **42**:703–
716. doi:10.1016/S0896-6273(04)00268-5
Van Gennip JLM, Boswell CW, Ciruna B. 2018. Neuroinflammatory signals drive
spinal curve formation in zebrafish models of idiopathic scoliosis. *Sci Adv*
4:eaav1781. doi:10.1126/sciadv.aav1781
van Rooijen E, Giles RH, Voest EE, van Rooijen C, Schulte-Merker S, van Eeden FJ.
2008. LRRC50, a conserved ciliary protein implicated in polycystic kidney
disease. *J Am Soc Nephrol* **19**:1128–1138. doi:10.1681/ASN.2007080917
Viau A, Bienaimé F, Lukas K, Todkar AP, Knoll M, Yakulov TA, Hofherr A, Kretz O,
Helmstädter M, Reichardt W, Braeg S, Aschman T, Merkle A, Pfeifer D, Dumit
VI, Gubler M, Nitschke R, Huber TB, Terzi F, Dengjel J, Grahammer F,
Köttgen M, Busch H, Boerries M, Walz G, Triantafyllopoulou A, Kuehn EW.
2018. Cilia-localized LKB 1 regulates chemokine signaling, macrophage
recruitment, and tissue homeostasis in the kidney. *The EMBO Journal*
37:e98615. doi:10.15252/embj.201798615
Vierkotten J, Dildrop R, Peters T, Wang B, Rüther U. 2007. Ftm is a novel basal body
protein of cilia involved in Shh signalling. *Development* **134**:2569–2577.
doi:10.1242/dev.003715
Wang J, Thomas HR, Thompson RG, Waldrep SC, Fogerty J, Song P, Li Z, Ma Y,
Santra P, Hoover JD, Yeo NC, Drummond IA, Yoder BK, Amack JD, Perkins
B, Parant JM. 2022. Variable phenotypes and penetrance between and within
different zebrafish ciliary transition zone mutants. *Disease Models &*
Mechanisms **15**:dmm049568. doi:10.1242/dmm.049568
Wieczorek S, Combes F, Lazar C, Giai Gianetto Q, Gatto L, Dorffer A, Hesse A-M,
Couté Y, Ferro M, Bruley C, Burger T. 2017. DAPAR & ProStaR: software to
perform statistical analyses in quantitative discovery proteomics.
Bioinformatics **33**:135–136. doi:10.1093/bioinformatics/btw580
Wiegering A, Rüther U, Gerhardt C. 2018. The ciliary protein Rpgrip1l in
development and disease. *Developmental Biology* **442**:60–68.
doi:10.1016/j.ydbio.2018.07.024
Xie H, Kang Y, Liu J, Huang M, Dai Z, Shi J, Wang S, Li L, Li Y, Zheng P, Sun Y,
Han Q, Zhang J, Zhu Z, Xu L, Yelick PC, Cao M, Zhao C. 2023. Ependymal
polarity defects coupled with disorganized ciliary beating drive abnormal
cerebrospinal fluid flow and spine curvature in zebrafish. *PLoS Biol*
21:e3002008. doi:10.1371/journal.pbio.3002008

Yu X, Ng CP, Habacher H, Roy S. 2008. Foxj1 transcription factors are master
regulators of the motile ciliogenic program. *Nat Genet* **40**:1445–1453.
doi:10.1038/ng.263
Zamanian JL, Xu L, Foo LC, Nouri N, Zhou L, Giffard RG, Barres BA. 2012. Genomic
Analysis of Reactive Astrogliosis. *J Neurosci* **32**:6391–6410.
doi:10.1523/JNEUROSCI.6221-11.2012
Zhang X, Jia S, Chen Z, Chong YL, Xie H, Feng D, Wu X, Song DZ, Roy S, Zhao C.
2018. Cilia-driven cerebrospinal fluid flow directs expression of urotensin
neuropeptides to straighten the vertebrate body axis. *Nat Genet* **50**:1666–
1673. doi:10.1038/s41588-018-0260-3
Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C,
Chanda SK. 2019. Metascape provides a biologist-oriented resource for the
analysis of systems-level datasets. *Nat Commun* **10**:1523.
doi:10.1038/s41467-019-09234-6

 MATERIAL AND METHODS

Zebrafish

Wild-type, *rpgrip1l^ex4^* and *rpgrip1l^Δ^* zebrafish embryos and adults were raised, staged
and maintained as previously described (Kimmel et al., 1995). All our experiments
were made in agreement with the european Directive 210/63/EU on the protection of
animals used for scientific purposes, and the French application decree 'Décret
2013-118'. The projects of our group have been approved by our local ethical
committee 'Comité d'éthique Charles Darwin'. The authorization number is APAFIS
#31540-2021051809258003 v4. The fish facility has been approved by the French
'Service for animal protection and health' with approval number A750525. All
experiments were performed on Danio rerio embryos of mixed AB/TL background.
Animals were raised at 28.5°C under a 14/10 light/dark cycle.

Rpgrip1l mutant generation and genotyping
1008
<u>Guide RNA preparation and microinjection</u>
CRISPR target sites were selected for their high predicted specificity and efficiency
using the CRISPOR online tool (Haeussler et al., 2016). Real efficiency was
assessed on zebrafish embryos by T7E1 test. The two most efficient guides
(Rpgrip1l_x4_G1: GCTTACGGTCCTTCACCAGACGG and Rpgrip1l_x25_G3:
CCTCAGTTGACAGGTTTCAGCGG) respectively situated 24 nt from the beginning
of exon 4 and 82 nt downstream of exon 25 were kept for further experiments.
sgRNA transcription templates were obtained by PCR using T7_Rpgrip1l-x4_G1_Fw
primer (5'-
GAAATTAATACGACTCACTATAGGCTTACGGTCCTTCACCAGAGTTTTAGAGCTA
GAAATAGC- 3') or T7_Rpgrip1l-x25_G3_Fw (5'-
GAAATTAATACGACTCACTATAGGCCTCAGTTGACAGGTTTCAGGTTTTAGAGCT
AGAAATAG C-3') as forward primer and sgRNA_R universal primer (5'-
AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTT
AACTTGCTA TTTCTAGCTCTAAAAC-3') as reverse primer. sgRNAs were
transcribed using Megascript T7 Transcription Kit (Thermo Fisher Scientific,
Waltham, MA) and purified using NucleoSpin® RNA Clean up XS kit (Macherey
Nagel, Du☐ren, Germany). sgRNA:Cas9 RNP complex was obtain by incubating

Cas9 protein (gift of TACGENE, Paris, France) (7.5 µM) with sgRNA (10 µM) in 20

mM Hepes-NaOH pH 7.5, 150 mM KCl for 10 min at 28 °C. 1–2 nl was injected per

embryo. For deletion, Rpgrip1l_x4_G1 and Rpgrip1l_x25_G3 RNP complexes were

mixed half and half.

We kept two F1 fish, one with a 15 Kb deletion between exon 4 and 25 of *rpgrip1l*,

that we called *rpgrip1l*$^\Delta$ and that we submitted to ZFIN under the name *rpgrip1l*

^{bps1}and a second one with a stop codon within exon 4 called *rpgrip1l*^{ex4}, that is

submitted to ZFIN under the name *rpgrip1l*^{bps2}.

1035

<u>Screening and genotyping</u>

Injected (F0) fish were screened for germline transmission by crossing with wild type

fish and extracting genomic DNA from resulting embryos. For genomic DNA

extraction, caudal fin (juveniles/adults) or whole embryo DNA were used. Genomic

DNA was isolated with Proteinase K (PK) digestion in 40 of lysis buffer (100 mM Tris-

HCl pH 7.5, 1 mM EDTA, 250 mM NaCl, 0.2% SDS, 0.1 µg/µl Proteinase K (PK)) for

embryos (300 µl for adult fin) overnight at 37°C with agitation. PK was inactivated for

10 min at 90°C and a five-fold dilution was used as template for PCR amplification.

Combined genotyping of wild type and mutant alleles (Supplementary Figure S1A')

was performed by PCR using 3 primers, a common reverse primer for both alleles :

Rpgrip1l_ex25_R3 (GTTGTGTCTCTGCCATATATTG), a specific forward primer for

the deleted allele: Rpgrip1l-ex4-del (CCCACACTGCATACGCACTC) and a specific

forward primer for the wild type allele : Rpgrip1l_ex25F

(AGTGTGCGGTACATCTCCAA) at an annealing temperature of 60°C for 35 cycles.

Expected size of the amplified fragment is 364 nt for the wild type allele and 544nt for

the mutant allele. The deleted allele presents a frameshift at position N77 leading to

the generation of a STOP codon after 11 aas, thus leading to a N-terminally

truncated protein at position 88/1256 aas.

For genotyping the *urp2* mutant line, caudal fins from adult fish were extracted in

300µl of lysis solution and PCR was performed as above to detect exon 5 deletion

using the following primers: urp2-F1 (TGATTACTAGCCCTGTCCCAAC) and urp2-R1

(AGGTACAGTACACACGTCACAG).

1058

1059

1060

μCT scans

μCT scans

The samples were scanned on a Bruker micro scanner (Skyscan 1272) with a
resolution of 8.5 μm, a rotation step of 0.55° and a total rotation of 180°. For the
acquisition of adult fish (2.5 cm), a 0.25 mm aluminium filter was used, for a voltage
of 50 kV and an intensity of 180 mA, for juvenile fish (0.9 to 1.2 cm) the filter was
omitted, and a voltage of 60 kV was used with an intensity of 166 mA. Each image
contained 1008 x 672 pixels and was based on the average of 3 images. The 3D
reconstruction by backprojection was carried out by the NRecon software and the 2D
image overlays were then cleaned by the CT Analyser software. The Dataviewer
software allowed all fish skeletons to be oriented in the same way taking the otoliths
as landmarks. The CTvox software then allowed 3D visualizing of the samples.
Morphometric analysis was performed with the CT Analyser software.

1073

Scanning electron microscopy

Scanning electron microscopy

3 month fish (3 controls, 5 mutants) were euthanized using lethal concentration of
MS222 (0.028 mg/mL). The brains were quickly dissected in 1.22 X PBS (pH 7.4),
0.1 M sodium cacodylate and fixed overnight with 2% glutaraldehyde in 0.61 X PBS
(pH 7.4), 0.1 M sodium cacodylate at 4°C. They were sectioned along the dorsal
midline with a razor blade to expose their ventricular surfaces. Both halves were
washed four times in 1.22 X PBS and post-fixed for 15 minutes in 1.22 X PBS
containing 1% OsO_4. Fixed samples were washed four times in ultrapure water,
dehydrated with a graded series of ethanol and critical point dried (CPD 300, Leica)
at 79 bar and 38 °C with liquid CO_2 as the transition fluid and then depressurized
slowly (0,025 bar/s). They were then mounted on aluminum mounts with conductive
silver cement. Sample surfaces were coated with a 5 nm platinum layer using a
sputtering device (ACE 600, Leica). Samples were observed under high vacuum
conditions using a Field Emission Scanning Electron Microscope (Gemini 500, Zeiss)
operating at 5 kV, with a 20 μm objective aperture diameter and a working distance
around 3 mm. Secondary electrons were collected with an in-lens detector. Scan
speed and line compensation integrations were adjusted during observation.

1091

1092

1093

Histological sections of juvenile and immunofluorescence on sections

1094

Juvenile and adult zebrafish were euthanized using lethal concentration of MS222

(0.028 mg/mL). Pictures and size measurements were systematically taken before

fixation. Fish were fixed in Zamboni fixative [35 ml PFA 4 %, 7.5 ml saturated picric

acid (1.2 %), 7.5 ml 0.2M Phosphate Buffer (PB)] [55] overnight at 4°C under

agitation. Fish were washed with Ethanol 70% and processed for dehydration by

successive 1 h incubation in Ethanol (3 x 70% and 2 x 100%) at room temperature

under agitation, then for paraffin inclusion. 14 µm sagittal paraffin sections were

obtained using a Leica RM2125RT microtome. Sections were deparaffinized and

antigen retrieval was performed by incubation for 7 min in boiling citrate buffer (10

mM, pH 6). Immunofluorescence staining was performed as described previously

(Andreu-Cervera et al., 2019). The following primary antibodies were used: anti-RF,

anti-Acetylated Tubulin, anti-Glutamylated Tubulin, anti-Arl13b, anti-LCP1, anti-

AnnexinA2. Corresponding primary and secondary antibodies are described and

referenced in the Resources Table.

1109

Immunofluorescence on whole embryos

1110

Embryos from 24 to 40 hpf were fixed 4 hr to overnight in 4% paraformaldehyde

(PFA) at 4°C. For Reissner fiber staining, larvae at 48 and 72 hpf were fixed 2 hr in

4% PFA and 3% sucrose at 4°C, skin from the rostral trunk and yolk were removed.

Samples from 24 to 40 hpf embryos were blocked overnight in a solution containing

0.5% Triton, 1% DMSO, 10% normal goat serum and 2 mg/mL BSA. For older

samples (48 to 72 hpf) triton concentration was increased to 0.7%. Primary

antibodies were incubated one to two nights at 4°C in a buffer containing 0.5% Triton,

1% DMSO, 1% NGS and 1 mg/mL BSA. All secondary antibodies were from

Molecular Probes, used at 1:400 in blocking buffer, and incubated a minimum of 2.5

hr at room temperature. The primary antibodies chosen for in toto immuno-labeling

against Reissner fiber, Acetylated-tubulin, Myc, Arl13b and Gamma-tubulin as well as

the corresponding secondary antibodies are referenced in the Key Resources Table.

Whole mount zebrafish embryos (dorsal or lateral mounting in Vectashield Mounting

Medium) were imaged on a Leica SP5 confocal microscope and Zeiss LSM910

confocal microscope, both equipped with a 63X immersion objective. Images were then processed using Fiji (Schindelin et al., 2012).

Whole-mount brain clearing

Brains were dissected from 4-5 wpf size-matched (0.9-1.2 mm length) zebrafish after in toto fixation with formaldehyde. Whole–mount tissue clearing was performed following the zPACT protocol (Affaticati et al., 2017). In brief, brains were infused for 2 days in hydrogel monomer solution (4% acrylamide, 0.25% VA- 044, 1% formaldehyde and 5% DMSO in 1X PBS) at 4°C. Polymerization was carried out for 2h30 at 37°C in a desiccation chamber filled with pure nitrogen. Brains were transferred into histology cassettes and incubated in clearing solution (8% SDS and 200 mM boric acid in dH2O) at 37°C with gentle agitation for 8 days. Cleared brains were washed in 1X PBS with 0.1% Tween-20 (PBT) for 3 days at room temperature and kept in 0.5% formaldehyde, 0.05% sodium azide in PBT at 4°C until further processing. Brains were subsequently placed for 1h in depigmentation pre-incubation solution (0.5X SSC, 0.1% Tween-20 in dH2O) at room temperature. The solution was replaced by depigmentation solution (0.5X SSC, Triton X-100 0.5%, formamide 0.05% and H2O2 0.03% in dH2O) for 45 minutes at room temperature. Depigmented brains were washed for 4h in PBT and post-fixed (2% formaldehyde and 2% DMSO in PBT) overnight at 4°C.

Cleared brain immunostaining of whole adult brains

Whole-mount immunolabeling of cleared brains was performed as described in the zPACT protocol with slight modifications. Briefly, brains were washed in PBT for one day at room temperature and blocked for 10h in 10% normal goat serum, 10% DMSO, 5% PBS-glycine 1M and 0.5% Triton X-100 in PBT at room temperature. Brains were washed again in PBT for 1h prior to incubation with anti- ZO-1 antibody (ZO1-1A12, Thermofisher, 1:150) in staining solution (2% normal goat serum, 10% DMSO, 0.1% Tween-20, 0.1% Triton X-100 and 0.05% sodium azide in PBT) for 12 days at room temperature under gentle agitation. Primary antibody was renewed once after 6 days of incubation. Samples were washed three times in PBT and thereafter incubated with Alexa Fluor 488-conjugated secondary antibody (A11001, Invitrogen, 1:200) for 10 days in the staining solution at room temperature under gentle agitation. Secondary antibody was renewed once after 5 days of incubation.

Samples were washed three times in PBT prior to a counterstaining with DiIC18 (D282, Invitrogen, 1µM) in the staining solution for three days at room temperature with gentle agitation. Samples were washed in PBT for 3h before mounting procedure.

Mounting and confocal imaging

Samples were placed in 50% fructose-based high-refractive index solution (fbHRI, see (Affaticati et al., 2017) /50% PBT for 1h, then in 100% fbHRI. Brains were mounted in agarose-coated (1% in standard embryo medium) 60 mm Petri dish with custom imprinted niches to help orientation. Niche-fitted brains were embedded in 1% phytagel and the Petri dish filled with 100% fbHRI. fbHRI was changed three times before imaging until its refractive index matched 1.457. Images were acquired with a Leica TCS SP8 laser scanning confocal microscope equipped with a Leica HC FLUOTAR L 25x/1.00 IMM motCorr objective. Brains were scanned at a resolution of 1.74x1.74x1.74 µm (xyz) and tiled into 45 to 70 individual image stacks, depending on brain dimensions, subsequently stitched, using LAS X software.

Volumetric analysis of the posterior ventricles

The volumes of the posterior ventricles were segmented, reconstructed and analyzed using Amira for Life & Biomedical Sciences software (Thermo Fisher Scientific). In essence the ventricles volumes were manually segmented in Amira's segmentation editor and subsequently refined by local thresholding and simplification of the corresponding surfaces. Volumes, which were open to the environment were artificially closed with minimal surfaces by connecting the distal-most points of their surface to the contralaterally corresponding points using straight edges. Due to the biologic variability of the sample population the overall size of the specimens needed to be normalized to keep the measured volumes comparable. For this spatial normalization one of the specimens was randomly selected from the wildtype population as template (1664, grey) and the 'Registration' module in Amira was used to compute region-specific rigid registrations for the other specimens, allowing for isotropic scaling only [details in supp. mat.]. For excluding the influence of the ventricular volumes, the registration was computed on the basis of the independent reference stain (DiIC18) (details in supp. mat.). Region specific volume differences

1190 between the mutant and wildtype population were evaluated on seven subvolumes of
1191 the posterior ventricles (details in supp. mat.).

1192

1193

1194

RNA extraction for transcriptome analysis and quantitative RT-PCR
Juvenile and adult zebrafish were euthanized using lethal concentration of MS222
(0.28 mg/mL). Their length was measured and their fin cut-off for genotyping. For
transcriptomic analysis, brain and dorsal trunk from 1 month juvenile (0.9 to 1.0 cm)
zebrafish were dissected in cold PBS with forceps and lysed in QIAzol (QIAGEN)
after homogenization with plunger pistons and 1ml syringes. Samples were either
stored in QIAzol at -80°C or immediately processed. Extracts containing RNA were
loaded onto QIAGEN-mini columns, DNAse digested and purified in the miRNAeasy
QIAGEN kit (Cat 217004) protocol. Samples were stored at -80°C until use. RNA
concentration and size profile were obtained on the Tapestation of ICM platform. All
preparations had a RIN above 9.2. For Q-PCR from juveniles, whole fish minus
internal organs were lysed in Trizol (Life technologies) using the Manufacturer
protocol.

1208

Quantitative PCR from individual juvenile or adult fish
The cDNA from the isolated RNAs was obtained following GoScript Reverse
Transcription System protocol (Promega), using 3 to 4 µg of total RNA for each
sample. The 20 µL RT-reaction was diluted 4-fold and 4µl was used for each
amplification performed in duplicates. The primers used for qPCR were the following:
urp2 (F: ACCAGAGGAAACAGCAATGGAC; R: TGAGGTTTCCATCCGTCACTAC),
urp1 (F: ACATTCTGGCTGTGGTTTGTTC; R: CTCTTTTGCACCTCTCTGAAGC),
urp (F: GCGGAAAAATGTCATGCCTCTTC; R: TTGAGCTCCTTTTGAAGCTCCTG),
uts2a (F: CACTGCTCAACAGAGACAGTATCA; R:
CCAAAAGACCACTGGGAGGAAC), *uts2b* (F: TACCCGTCTCTCATCAGTGGAG; R:
TTTTCCAGCAAGGCCTCTTTTAC) where all of them are from Gaillard et al. 2023,
rpgrip1l (F: CAGACACCTGCTGGAGTTACA; R: TCCTGACTCACATCAAACGCA),
irg1l (F: TCCCTAAGAGGTTCCATCCTCC; R: AAGATGCAGCGATGGCCAAA),
stat3 (F: CAGGACTGGGCGTATGCG; R: GAAGCGGCTGTACTGCTGAT), *c4b* (F:

GGGTGTTCTTATGGCGGTGG and R: GCGCACAACAAGCTTTTCTCATC) *lsm12b*

(F: GAGACTCCTCCTCCTCTAGCAT and R: GATTGCATAGGCTTGGGACAAC).

The Q-PCR were performed using the StepOne real-time PCR system and following

its standard amplification protocol (Thermo Scientific). Relative gene expression

levels were quantified using the comparative CT method ($2^{-\Delta\Delta CT}$ method or $2^{-\Delta CT}$) on

the basis of CT values for target genes and *lsm12b* as internal control.

RNA-sequencing and analysis

RNA-seq libraries were constructed by the ICM Platform (PARIS) from 100 ng of total

RNA using the KAPA mRNA Hyperprep kit (Roche) that allows to obtain stranded

polyA libraries, and their quality controlled on Agilent Tape station. The sequencing

was performed on a NovaSeq 6000 Illumina for both strands. Sequences were

aligned against the GRCz11 version of zebrafish genome using **RNA Star function**

in "Galaxy" environment. The 12 Brain libraries reached between 22 to 26 Millions of

assigned reads, and the trunk libraries between 26 to 35 Millions of assigned reads.

Expression level for each gene was determined using **Feature counts function** from

RNA Star BAM files. **DESeq2 function** was used to calculate the log2 Fold change

and adjusted P values from the Wald test for each gene comparing the 7 mutants

raw count collection to the 5 controls raw counts collection. Volcano plots were

generated using the Log2 fold change and P adj. value of each Gene ID from the

DESeq2 table. A threshold of 5.10 E-2 was chosen for the Padj. values and a log2

fold change > to +0.75 or < to -0.75 to select significantly up or down-regulated

regulated genes. Both gene lists were used as input to search for enriched GO terms

and KEGGs pathways using Metascape software (Zhou et al., 2019) as well as to

compare up-regulated genes in the dorsal trunk versus the brain using Galaxy.

1247

Quantitative proteomics

Sample preparation: Adult zebrafish were euthanized using lethal concentration of

MS222 (0.28 mg/mL). For proteomic analysis, brain from 3 months zebrafish were

dissected in cold PBS with forceps and immediately flash frozen in liquid nitrogen

and stored at -80°C. Zebrafish brain samples were lysed in RIPA buffer (Sigma) with

1/100 antiprotease and sonicated for 5 minutes. After a 660 nm protein assay

(Thermo), samples were digested using a Single Pot Solid Phase enhanced Sample

Preparation (SP3) protocol (Hughes et al., 2014). In brief, 40 µg of each protein

extract was reduced with 12mM dithiothreitol (DTT) and alkylated using 40 mM

iodoacetamic acid (IAM). A mixture of hydrophilic and hydrophobic magnetic beads

was used to clean-up the proteins at a ratio of 20:1 beads:proteins (Sera-Mag Speed

beads, Fisher Scientific). After addition of ACN to a final concentration of 50%, the

beads were allowed to bind to the proteins for 18 minutes. Protein-bead mixtures

were washed twice with 80% EtOH and once with 100% ACN. The protein-bead

complexes were digested with a mixture of trypsin:LysC (Promega) at a 1:20 ratio

overnight at 37°C. Extracted peptides were cleaned-up using automated C18 solid

phase extraction on the Bravo AssayMAP platform (Agilent Technologies).

1265

NanoLC-MS-MS analysis

The peptide extracts were analysed on a nanoLC-TimsTof Pro coupling (Bruker

Daltonics). The peptides (200ng) were separated on an IonOpticks column (25cm X

75µm 1.6µm C18 resin) using a gradient of 2-37% B (2%ACN, 0.1%FA) in 100

minutes at a flow rate of 0.3 µl/min. The dual TIMS had a ramp time and

accumulation time of 166 ms resulting in a total cycle time of 1.89s. Data was

acquired in Data Dependent Acquisition-Parallel Accumulation Serial Fragmentation

(DDA-PASEF) mode with 10 PASEF scans in a mass range from 100 m/z to 1700

m/z. The ion mobility scan range was fixed from 0.7-1.25 Vs/cm^2. All samples were

injected using a randomized injection sequence. To minimize carry-over, one solvent

blank injection was performed after each sample.

1277

Data interpretation

Raw files were converted to .mgf peaklists using DataAnalysis (version 5.3, Bruker

Daltonics) and were submitted to Mascot database searches (version 2.5.1,

MatrixScience, London, UK) against a *Danio rerio* protein sequence database

downloaded from UniProtKB-SwissProt (2022_05_18, 61 732 entries, taxonomy ID:

7955), to which common contaminants and decoy sequences were added. Spectra

were searched with a mass tolerance of 15 ppm in MS mode and 0.05 Da in MS/MS

mode. One trypsin missed cleavage was tolerated. Carbamidomethylation of cysteine

residues was set as a fixed modification. Oxidation of methionine residues and

acetylation of proteins' n-termini were set as variable modifications. Identification

results were imported into the Proline software (version 2.1.2,

http://proline.profiproteomics.fr) (Bouyssié et al., 2020) for validation. Peptide
Spectrum Matches (PSM) with pretty ranks equal to one were retained. False
Discovery Rate (FDR) was then optimized to be below 1% at PSM level using
Mascot Adjusted E-value and below 1% at protein level using Mascot Mudpit score.
For label free quantification, peptide abundances were extracted without cross
assignment between the samples. Protein abundances were computed using the
sum of the unique peptide abundances normalized at the peptide level using the
median.
To be considered, proteins must be identified in at least four out of the five replicates
in at least one condition. Imputation of the missing values and differential data
analysis were performed using the open-source ProStaR software (Wieczorek et al.,
2017). Imputation of missing values was done using the approximation of the lower
limit of quantification by the 2.5% lower quantile of each replicate intensity distribution
("det quantile"). A Limma moderated t-test was applied on the dataset to perform
differential analysis. The adaptive Benjamini-Hochberg procedure was applied to
adjust the p-values and FDR. The mass spectrometry proteomics data have been
deposited to the ProteomeXchange Consortium via the PRIDE partner repository
(Perez-Riverol et al., 2022), with the dataset identifier PXD042283.
1307

Embryos drug treatment

1308
1309 27 hpf embryos from *dnaaf1*[tm317b/+] incross were dechorionated manually. N-acetyl
1310 cysteine ethyl ester (NACET) (BIOLLA Chemicals #59587-09-6) was prepared
1311 extemporaneously at the final concentration of 3 mM in pure water (pH: 7.2;
1312 conductivity 650 µS). Embryos were treated between 27 hpf to 60 hpf and embryos
1313 were anesthetized before being imaged laterally.
1314

Juvenile drug treatment and preparation of treated samples

1315
Two hundred larvae were housed off-system in 1.8 liter tanks with 20 fish per tank,
being fed twice a day throughout the experiment. 100 were treated with NACET
(BIOLLA Chemicals #59587-09-6), which was prepared extemporaneously at 1.5 mM
(286.5mg/L) in fish water (pH 7.2, Conductivity 650 µS) which was changed once per
day. Fish were treated from 30 dpf to 84 dpf and monitored for curvature onset once
per week. At 85 dpf, fish were euthanized, imaged to measure the curvature index.

Half of the fish were fixed for histology analysis in zamboni fixative [35 ml PFA 4 %,

15 ml saturated picric acid (1.2 %)] overnight at 4°C under agitation, while the other

half was processed for RT-qPCR analysis. To prepare dorsal trunk RNA, internal

organs were removed in cold PBS 1X, and the remaining tissue was cut in small

pieces before lysis in QIAzol (QIAGEN). Homogenization was achieved using

plunger pistons and passages through 1ml syringes.

Transgenesis

The Tol2 -5.2foxj1a:5xmyc-RPGRIP1L plasmid was produced using the Gateway

system by combining four plasmids. P5E-foxj1a enhancer was kindly donated by [3].

5xmyc-RPGRIP1L cDNA was amplified from the plasmid pCS2-5xmyc-RPGRIP1L

(Mahuzier et al., 2012) using CloneAmp HiFi PCR Premix (Takara # 639298) using

primers   :   forward   5'-

GGGGACAAGTTTGTACAAAAAGCAGGCTGCAGGATCCCATCGATTTAAAGCT-3'

and   reverse   5'

GGGGACCACTTTGTACAAGAAAGCTGGGTCTCCGAGCTCGGTTCAAGCCTCCA

AGTCATCTCTGT-3. The PCR product was gel purified using QIAEX II gel extraction

kit (QIAGEN, #20021) . BP recombination (Invitrogren "Gateway BP Clonase II

Enzyme Mix" #11789-020) was performed into pDONR221 #218 to generate pMe-

5xmyc-RPGRIP1L. The 3' entry polyadenylation signal plasmid was the one

described in [61]. The 3 plasmids were shuttled into the backbone containing the

cmcl2:GFP selection cassette (Kwan et al., 2007) using the LR recombination

(Invitrogen "Gateway LR Clonase II Plus Enzyme Mix" #12538-120). rpgrip1l[+/-] X AB

embryos outcrosses were injected at the one cell stage with 1nl of a mix containing

20 ng/µl plasmid and 25 ng/µl Tol2 transposase RNA and screened at 48 hpf for GFP

expression in the heart. Some F0 founders produced F1 zebrafish carrying one copy

of the transgene that were further checked for RPGRIP1L specific expression in

foxj1a territory, using Myc labelling at the base of FP cilia in 2,5 dpf embryos.

The Tol2-1.7kb col2a1a:5xMyc-RPGRIP1L plasmid was produced using the Gateway

System by combining four plasmids. Col2a1a enhancer was amplified from the

plasmid -1.7kbcol2a1a:EGFP-CAAX (Dale and Topczewski, 2011) using CloneAmp

HiFi premix (Takara #639298) with the primers: forward 5'-GGG GAC AAC TTT GTA

TAG AAA AGT TGG CCC TCT GAC ACC TGA TGC CAA TTG C-3' and reverse 5'-

GGG GAC TGC TTT TTT GTA CAA ACT TGC TTG CAG GTC CTA AGG GGT GAA

AGT CG-. The PCR product was gel purified and BP recombination was performed
into pDONR_P4-P1R-#219. The middle entry plasmid and the 3' plasmid were the
same as those used to generate the Tol-5.2foxj1a: 5xmyc-RPGRIP1L plasmid. F1
transgenic were first selected for GFP expression in the heart and then on Myc
labelling within notochordal cells at 2,5 dpf within their embryonic progeny.

1360

Skeletal preparations, Alizarin staining and imaging
Animals were euthanized using a lethal concentration of MS222 (0.28 mg/mL), skin
was removed and fixation was performed with 4% paraformaldehyde overnight at
4°C. After evisceration, samples were incubated in borax 5%, rinsed several times in
1% KOH and incubated in solution composed with Alizarin 0.01% (Sigma, A5533)
and KOH 1% during 2 days. Fish were rinsed several times in KOH 1% and
incubated in trypsin 1% and borax 2% for 2 days until cleared. Samples were rinsed
several times in distilled water and transferred in glycerol 80% (in KOH 1%) using
progressive dilutions. Samples were kept in glycerol 80% until imaging.

1370

Cobb angles measurements
To quantitatively evaluate the severity of spine curvature in *rpgrip1l$^{+/-}$; urp2$^{+/-}$*
incrosses, we used the zebrafish skeletal preparations stained with Alizarin. We drew
parallel lines to the top and bottom most displaced vertebrae on lateral and dorsal
views. The Cobb angle was then measured as the angle of intersection between
lines drawn perpendicular to the original 2 lines. This was conducted using FIJI
software. For each fish, the total Cobb angle was calculated by summing all Cobb
angles.

1379

Curvature index measurements
We implemented a novel procedure to quantify curvatures on both axes by drawing a
line along the body of the fish as shown in Figure supplementary 4I and curvature
was calculated in MATLAB (code available on demand). We decomposed the line in
a series of equidistant points and we measured the curvature at each of these points
using the LineCurvature2D function. We then added the absolute values of these
measures, which are expressed as an angle (in radian) per unit of length. The higher
the sum is for a given line, the more this line is curvated. We verified the accurancy

of this metric by comparing it with the visual assessment of curvatures by 3

independent observers. For *rpgrip1l*$^{+/-}$*; urp2*$^{+/-}$ incross and NACET experiment, a line

following the body deformation was drawn from the mouth to the base of the tail

following the midline of the fish in lateral position and another one along the dorsal

axis. Curvature index of both curves were summed for each fish. Curvature analysis

was performed blinded to fish genotype. *rpgri1l*$^{+/+}$ or *rpgrip1l*$^{+/-}$ siblings were used as

controls.

1395

Data acquisition for body-curvature analysis at embryonic stage

Zebrafish embryos were anesthetized and imaged laterally with the head pointing to

the left. The angle between the line connecting the center of the eye to the center of

the yolk and the line connecting the center of the yolk to the tip of the tail was

measured to evaluate the body curvature of the embryos. For quantitative analysis,

the angles of each embryo were put in the same concentric circles represented with

a Roseplot, with 0° pointing to the right and 90° pointing to the top. Each triangle

represents a 30° quadrant, and its height indicates the number of embryos within the

same quadrant.

1405

Quantification, statistical analysis and figure preparation

For all experiments the number of samples analyzed is indicated in the text and/or

visible in the figures. Statistical analysis was performed using the Prism software.

****: P value < 0.0001; ***: P value < 0.001; **: P value < 0.01; *: P value < 0.05.

Graph were made using Prism and Matlab and figures were assembled using

Photoshop software.

1412

1413 **REAGENT AND RESOURCE TABLE**

Reagent type or resource	Designation	Source or reference	Identifiers	Additional Infos

Antibody (antiserum)	Rabbit polyclonal IgG anti-bovine Reissner Fiber	Didier et al 1995	Courtesy of Dr. S.Gobron	1:200
Antibody	Mouse monoclonal IgG1k anti ZO1 -1A12	Thermofisher	#33-9100	1:150
Antibody	Mouse monoclonal IgG2b anti-acetylated-tubulin (clone 6-11B-1)	Sigma-Aldrich	#T 6793 RRID: AB_477585	1:400
Antibody	Mouse monoclonal IgG1 anti glutamylated Tubulin	Adipogen	#AG-20B-0020-C100	1:500
Antibody	Rabbit polyclonal anti-Arl13b	Proteintech	#17711-1-AP	1:200
Antibody	Mouse monoclonal IgG1 anti-GFAP	Sigma	#G3893	1:400
Antibody	Chicken monoclonal	AVES lab	#GFP-1020	1:200

	IgY anti-GFP			
Antibody	Mouse monoclonal IgG2a anti-myc (clone 9b11)	Cell signaling	#2276	1:200
Antibody	Rabbit polyclonal anti-LCP1	GeneTex	#GTX124420	1:200
Antibody	Mouse monoclonal IgG1 anti-gamma-tubulin	Sigma	#T6557	1:500
Antibody	Mouse polyclonal anti-annexin A2	Proteintech	#11256-1-AP	1:200
Antibody	Goat anti-mouse IgG1 Alexa633	Molecular probes	# A-21126 RRID:AB_2535768	1:400
Antibody	Goat anti-mouse IgG1 Alexa568	Molecular Probes	# A-21124 RRID: AB_2535766	1:400
Antibody	Goat anti-mouse IgG2a Alexa568	Molecular probes	# A-21134 RRID:AB_2535773	1:400

Antibody	Goat anti-mouse IgG2a Alexa488	Molecular probes	# A-21131 RRID:AB_141618	1:400
Antibody	Goat anti-mouse IgG2b Alexa633	Molecular probes	# A-21146 RRID:AB_2535782	1:400
Antibody	Goat anti-mouse IgG2b Alexa568	Molecular probes	# A-21144 RRID: AB_2535780	1:400
Antibody	Goat anti-rabbit IgG Alexa568	Molecular probes	# A-11011 RRID:AB_143157	1:400
Antibody	Goat anti-rabbit IgG Alexa488	Molecular probes	# A-11008 RRID: AB_143165	1:400
Antibody	FITC donkey Anti-Chicken IgY	Jackson ImmunoResearch	# 703-096-155	1:200
Chemical	Vectashield	Vector Laboratories	H-1000	
Strain (Danio rerio)	zebrafish wild-type AB or (TL x AB) hybrid strains	N/A	N/A	
Strain (Danio rerio)	Zebrafish *rpgrip1Δ*	This manuscript,		

	mutant strain	Zfin:*rpgrip1l*^{bps1}		
Strain (Danio rerio)	Zebrafish *rpgrip1l*^{ex4} mutant strain	This manuscript, Zfin:*rpgrip1l*^{bps2}		
Strain (Danio rerio)	Zebrafish *urp2*^{+/-}	(Gaillard et al., 2023)	ZDB-ALT-230207-5	
Strain (Danio rerio)	Zebrafish *dnaaf1*^{tm317b/+}	(van Rooijen et al., 2008)		
Strain (Danio rerio)	Zebrafish Scospondin-GFP^{ut24}	(Troutwine et al., 2020)		
Recomb. DNA	Foxj1 promoter-enhancer sequence	(Dale and Topczewski, 2011)		
Software	LASX	Leica		
Software	Amira for Life & Biomedical Sciences	Thermo Fisher Scientific		
Software	CRISPOR	(Haeussler et al., 2016)	http://crispor.tefor.net/	
Software	Fiji/ImageJ	(Schindelin et al., 2012)	https://imagej.net/Fiji/Downloads	
Software	PRISM	GraphPad	https://www.graphpad.com/	
Software	Metascape		https://metascape.org/gp/index.html#/mai	

- Zfin:*rpgrip1l* ^{bps1}
- *rpgrip1l* ^{ex4}, Zfin:*rpgrip1l* ^{bps2}
- *urp2* $^{+/-}$
- *dnaaf1* $^{tm317b/+}$
- Scospondin-GFP ^{ut24}

			n/step1	
Software	Matlab	The Mathworks, Inc.	https://fr.mathworks.com/products/matlab.html	
Software	NRecon reconstruction software	Micro Photonics Inc.		
Software	CTvox	Bruker		
Software	CT Analysis	Bruker		
Other				
Refractometer	Roth Sochiel EURL	DR 301-95		
Confocal microscope	Leica	SP5		
Confocal microscope	Zeiss	LSM 980 Inverted		
Confocal microscope	Zeiss	LSM 710		
Macroscope	Zeiss	AXIOZOOM V16		
Micro-scanner	Bruker	Skyscan 1272		
Microtome	Leica	RM2125RT		
Q-PCR apparatus	Applied Biosystems	Step One plus		

1414

1415

1416 **FIGURE LEGENDS**

1417

Figure 1: *rpgrip1l^{-/-}* juvenile fish develop scoliosis, which is rescued by
RPGRIP1L expression in *foxj1a*-positive cells.

(A) Representative *rpgrip1l^{-/-}* fish at 5 days post-fertilization (dpf) (4 mm body length,
larvae), 5 weeks pf (wpf) (1.2 cm body length, juveniles) and 12 wpf (2.5 cm body
length, adults), showing absence of morphological defects in embryos, onset of spine
curvature (tail-up) in juveniles and scoliosis in adults. **(B-C)** Graph showing the time
course of scoliosis appearance **(B)** in *rpgrip1l^{-/+}* incrosses (total 4 clutches, 252 fish)
and scoliosis penetrance **(C)** in adults. **(D-E)** Micro-computed tomography (µCT)
scans of control siblings (D) and *rpgrip1l^{-/-}* (E) adult fish. 4 fish were analyzed for
each condition. **(F-G)** Dorsal and lateral superimposed µCT views of the spines of
one control (yellow) and two *rpgrip1l^{-/-}* (pink, cyan) adult fish illustrating the 3D spine
curvatures in mutants. **(H)** Schematic representation of the two rescue constructs. **(I)**
Table presenting the number and phenotype of *rpgrip1l^{-/-}* fish generated for both
rescue experiments by stable transgenesis *(J)* Graph representing scoliosis
penetrance in transgenic and non-transgenic *rpgrip1l^{-/-}* siblings.

1433

Figure 2 Altered ciliogenesis at trunk and subcommissural organ (SCO) levels.

(**A-C'''**) Immunostaining of trunk sagittal sections at the level of the spinal cord
central canal in *rpgrip1l^{+/+}* (A1-A'''; n=5), straight *rpgrip1l^{-/-}* (B-B'''; n=4) and scoliotic
rpgrip1l^{-/-} fish (C-C'''; n=6)) at 8 weeks (N=2 independent experiments). Cilia were
immunostained with Arl13b (cyan, A'', B'', C'') and glutamylated-tubulin (magenta,
A''', B''', C'''). Panels A-C' are merged images with DAPI. Panels A', B' and C' show
higher magnification of the squared regions in A, B and C, respectively. Scale bars:
10 µm in A, 5 µm in A'. **(D)** Violin plot showing the distribution of cilia length along the
central canal of *rpgrip1l^{+/+}* siblings (n=300 cilia from 5 fish), straight *rpgrip1l^{-/-}* (n=267
from 4 fish) and scoliotic *rpgrip1l^{-/-}* (n=414 from 6 fish) juvenile fish (N=2). **(E)** Violin
plot showing the distribution of cilia density along the central canal of juvenile
rpgrip1l^{+/+} siblings (n=5), straight *rpgrip1l^{-/-}* (n=4) and scoliotic *rpgrip1l^{-/-}* fish (n=6)
(N=2). Density is the average number of ventral and dorsal cilia lining the CC over a
length of 100 µm from 3 to 5 sections for each fish. Each dot represents the mean

cilia length (D) or cilia density (E) of 4-8 sections analyzed at different antero-
posterior levels. Statistical analysis was performed using Tukey's multiple
comparisons test where * means P-value < 0.05, and ** means P-value < 0.01. **(F)**
Violin plot showing the distribution of width of the central canal (nuclear-free region)
(in μm) of juvenile *rpgrip1l*$^{+/+}$ (n=5), straight *rpgrip1l*$^{/-}$ (n=4) and scoliotic *rpgrip1l*$^{/-}$ fish
(n=6) (N=2). Each dot represents the mean of 3 measurements made at different AP
levels. Statistical analysis was performed using Tukey's multiple comparisons test
where ns means non-significant, * means P-value < 0.05. **(G-L)** Immunostaining of
cilia on transverse sections of the brain at SCO level using glutamylated-tubulin and
acetylated-tubulin antibodies (both in white), in *rpgrip1l*$^{+/+}$ (n=6) (G, G', H), straight
rpgrip1l$^{/-}$ (n=3) (I, I', J) and scoliotic *rpgrip1l*$^{/-}$ (n=7) (K, K', L) 8 wpf fish (N=2). Nuclei
were stained with DAPI (blue). G', I' and K' are higher magnifications of squared
regions in G, I and K, respectively. In H, J and L the bottom-right inset represents a
higher magnification of the squared region. Scale bars in G and G': 10 μm.

1462

Figure 3: *rpgrip1*$^{-/-}$ juveniles show ventricular dilations and loss of the Reissner
Fiber at scoliosis onset.

(A) Reconstruction of the RhV at the level of posterior midbrain and hindbrain in a
transparised 5 wpf control zebrafish brain, stained with ZO1 antibody (ventricular
surface) and DiI (global brain shape). Several brain regions are annotated on the
reconstruction where Tel means telencephalon, ChP forebrain choroid plexus, SCO
subcommissural organ, TeO optic tectum, CCe corpus cerebelli, CC crista
cerebellaris and Rh rhombencephalon. **(B-D)** Cyan circles represent the ROIs
measured in F. The orange line in D indicates the level of optical sections in E. **(E)**
Optical transverse sections showing the caudal part of reconstructed ventricles of a
control and two *rpgrip1l*$^{/-}$ fish, one straight and two tail-up. **(F)** Graphs of the ventricle
volume at the onset of scoliosis in 3 control and 4 *rpgrip1l*$^{/-}$ (one straight and three
tail-up) fish. The green dot corresponds to the straight mutant fish. Each dot
represents a fish. Statistical analysis was performed using unpaired t-test where ns
means non-significant, and * means P-value < 0.05. **(G)** Graph of the surface of the
optical sections as illustrated in E. The green dot corresponds to a straight mutant
fish. Each dot represents a fish. Statistical analysis was performed using unpaired t-
test where ns means non-significant, and * means P-value < 0.05. **(H)** Schematic

representation of a portion of the spinal cord central canal in a lateral view of the fish
trunk, showing the Reissner Fiber (RF), and of the SCO in a transverse view of the
brain at the level of diencephalic Ventricle (DiV). **(I-Q)** Visualization of the GFP
fluorescence in the *sspo-GFP^(uts24/+)* transgenic line (fire LUT) in sagittal sections of the
trunk (I-K) (N=3) and transverse sections of the brain at the level of the SCO (N=2)
(L-Q) in juvenile fish. The corresponding fish are 8 wpf *rpgrip1l^(+/+)* (I, L, O) (n=10);
straight *rpgrip1l^(-/-)* (J, M, P) (n=9) and scoliotic *rpgrip1l^(-/-)* (K, N, Q) (n=8). L-N and O-Q
show sections at anterior and posterior SCO levels, respectively. Scale bars: 5 µm in
I-K, 10 µm in L-Q.
1490

Figure 4: Upregulation of *urp1/2* expression does not contribute to axis
curvature of *rpgrip1l^(-/-)* fish.

(A) Quantitative reverse transcription polymerase chain reaction (qRT-PCR)
assaying *urp2* expression level in *rpgrip1l^(-/-)* and *rpgrip1l^(+/+)* (WT) siblings. Each dot
represents the value obtained for one fish. Green and purple dots represent straight
and scoliotic *rpgrip1l^(-/-)* fish, respectively. qRT-PCR analysis was performed at 5 wpf
(juvenile stage, 5 controls and 7 *rpgrip1l^(-/-)*) and 12 wpf (adult stage, 5 controls and 9
rpgrip1l^(-/-)). *urp2* expression levels are increased by approximately 3.5-fold in *rpgrip1l^*
^(-/-) at both stages. Statistical analysis was performed using unpaired t-test where **P <
0.01. Error bars represent s.d. **(B)** Quantification of body axis curvature from dorsal
and lateral views in 4 mpf fish issued from [*rpgrip1l^(+/-)*, *urp2^(+/-)*] incrosses. Each point
represents the value measured for one fish. Statistical analysis was performed using
Tukey's multiple comparisons test where **** P < 0,0001 with error bars represent
s.d. n=9 [*rpgrip1l^(+/+)*; *urp2^(+/+)*], 7 [*rpgrip1l^(+/+)*; *urp2^(-/-)*], 27 [*rpgrip1l^(-/-)*; *urp2^(+/+)*], 48 [*rpgrip1l^*
^(-/-); *urp2^(+/-)*] and 27 [*rpgrip1l^(-/-)*; *urp2^(-/-)*] fish. **(C)** Representative [*rpgrip1l^(-/-)*; *urp2^(+/+)*] and
sibling [*rpgrip1l^(-/-)*; *urp2^(-/-)*] fish on lateral and dorsal views at 12 wpf. Scale bar 5mm.
(D, E) Expression levels of *urp2* (D) and *urp1* (E) in adult fish (12 wpf). Each dot
represents the value for one fish. Green and purple dots represent straight and
scoliotic *rpgrip1l^(-/-)* fish, respectively (n=5 [*rpgrip1l^(+/+)*; *urp2^(+/+)*], 9 [*rpgrip1l^(+/+)*; *urp2^(-/-)*], 11
[*rpgrip1l^(-/-)*; *urp2^(+/-)*] and 10 [*rpgrip1l^(-/-)*; *urp2^(-/-)*]). Each gene expression level is
compared to the *lsm12b* housekeeping gene. Statistical analysis was performed
using Tukey's multiple comparisons test where ns means non-significant and ** P <
0.01, **** P < 0.0001. Error bars represent s.d.

1514

Figure 5: Comparative transcriptome and proteome analysis of control and *rpgrip1l^/-* fish.

(A-B) Volcano-plots showing differentially regulated genes in the transcriptomes of the brain (A) and dorsal trunk (B) comparing 5 controls (2 wt, 3 *rpgrip1l^/+*) and 7 *rpgrip1l^/-* (6 tail-up and 1 straight) fish at 5 wpf (0.9 cm body length). Each dot corresponds to one gene. A gene was considered deregulated if its associated P-adj. value was inferior to 0,05. Red and blue dots represent up-regulated (Log2FC > 0.75), and down-regulated (Log2FC < -0.75) genes, respectively. The names of the top 25 most deregulated genes are boxed. **(C)** GO term analysis of biological processes in the brain samples using Metascape. **(D)** Venn diagram of the upregulated genes in brain and trunk samples, showing many common genes between the two regions. **(E)** Selected genes of interest upregulated in the trunk and/or brain of *rpgrip1l^/-* fish. The four genes in red were also found upregulated in the proteome of adult *rpgrip1l^/-* fish. **(F)** Volcano-plot showing differentially expressed proteins in the brain of 5 *rpgrip1l^+/+* fish versus 5 *rpgrip1l^/-* adult scoliotic fish (3 months pf). Each dot corresponds to one protein. A protein was considered differentially expressed if its associated P-adj. value was inferior to 0.05. Red and blue dots represent proteins enriched (Log2FC > 0.75) or depleted (Log2FC < -0.75) in mutant brains, respectively. The names of the top 25 most differentially expressed proteins are boxed.

Figure 6: Anxa2 and GFAP upregulation and increased number of LCP1-positive cells around CNS ventricles of *rpgrip1l^/-* juvenile fish at scoliosis onset.

(A-D') Immunostaining for GFAP (A, B, C, D) and Anxa2 (A', B', C', D') on adult brain transverse sections at DiV (A-B') and RhV at cerebellar level (C-D') levels in controls (A, A', C, C') (n=3) and *rpgrip1l^/-* scoliotic (B, B', D, D') (n=4) fish. Nuclei are stained with DAPI (blue). White stars indicate the SCO, white triangles point to the ventral RhV region. Open arrows point to a long cellular process co-labelled by Anxa2 and GFAP. **(E-L)** Immunostaining for GFAP (magenta) on transverse sections of the brain at SCO (E-H) and RhV (I-L) levels in controls (E, I) (n=4), straight *rpgrip1l^/-* (F, G, J, K) (n=4) and scoliotic *rpgrip1l^/-* (H, L) (n=3) juvenile (8 wpf) fish. Nuclei are stained

with DAPI (white). Scale bar: 125µM for A-D', 20 µm for E-H and 100 µm for I-L. **(M)** Graph representing the percentage of GFAP-positive cells among the total number of SCO cells in control (n=4), straight *rpgrip1l$^{-/-}$* (n=4), and scoliotic *rpgrip1l$^{-/-}$* (n=3) fish at scoliosis onset (N=1). Each dot represents the mean ratio of GFAP-positive over -negative cells in 3-6 sections per fish. Green dots correspond to fish with no or very low percentage of GFAP+ cells, red dots to fish with a high percentage of GFAP+ cells. Statistical analysis was performed with Tukey's multiple comparisons test where ns means non-significant, ** means P-value < 0.01 and *** P-value < 0.001. Error bars represent s.d. "str": straight; "sco": scoliotic. **(N-Q')** Immunostaining for LCP1+ microglia/macrophages (magenta) on brain transverse sections at SCO levels in a rectangle of 100 µm x 60 µm located under the posterior commissure in controls (n=4) (N, N'), straight *rpgrip1l$^{-/-}$* (n=4) (O-P') and *rpgrip1l$^{-/-}$* at scoliosis onset (n=3) (Q, Q') 8 wpf fish (N=1). Nuclei are stained with DAPI (white). Scale bar: 20 µm. **(R)** Quantification of LCP1+ cell density around the SCO in controls (n=4), straight *rpgrip1l$^{-/-}$* (n=4) and scoliotic *rpgrip1l$^{-/-}$* (n=3) at 8 wpf. 23 sections were counted for control, 21 sections for *rpgrip1l$^{-/-}$* straight and 18 sections for *rpgrip1l$^{-/-}$* scoliotic fish. Each dot represents the mean value for one fish. Green dots correspond to the fish without any GFAP+ SCO cells, red dots correspond to the fish with GFAP+ SCO cells (SCO cells are defined as the dorsal cells located underneath the posterior commissure and above the diencephalic ventricle). Note that straight *rpgrip1l$^{-/-}$* fish with GFAP+ SCO cells present a higher number of LCP1+ cells. Statistical analysis was performed with Tukey's multiple comparisons test where ns means non-significant. Error bars represent s.e.m. "str":straight "sco": scoliotic.

Figure 7: NACET treatment reduces scoliosis penetrance and severity in *rpgrip1l$^{-/-}$* and decreases astrogliosis and LCP1+ cell number at SCO level.

(A) Kinetics of scoliosis development in *rpgrip1l$^{+/-}$* incrosses in the presence (n=150 fish) or absence (n=174 fish) of NACET treatment (1.5 mM) from 4 to 12 weeks (N=1). "str": straight; "sco": scoliotic. **(B)** Quantification of body axis curvature from dorsal and lateral views at 13 wpf, at the end of the NACET treatment. Each dot corresponds to one fish. Among the 174 fish raised in untreated condition, we measured 18 controls (*rpgrip1l$^{+/-}$*) and 33 *rpgrip1l$^{-/-}$*, and upon NACET treatment, 16 controls (*rpgrip1l$^{+/+}$*) and 23 *rpgrip1l$^{-/-}$*. *rpgrip1l$^{+/-}$* represents a pool of *rpgrip1l$^{+/+}$* and

rpgrip1l[+/-]. Statistical analysis was performed using Tukey's multiple comparisons test

where **** means P-value < 0.0001. Error bars represent s.d. **(C)** Representative

pictures of *rpgrip1l*[-/-] fish with their curvature index without or after NACET treatment,

at 13 wpf. Scale bar is 0,5 cm. Left panels correspond to an untreated *rpgrip1l*[-/-]

scoliotic fish on lateral (upper) and dorsal (lower) views. Middle panels represent

treated *rpgrip1l*[-/-,] with a lower curvature index than scoliotic fish on lateral (upper)

and dorsal (lower) views and right panels are pictures of rescued straight *rpgrip1l*[-/-]

fish on lateral (upper) and dorsal (lower) views. **(D)** Graph representing the number

of LCP1+ cells in wt (n=3), untreated *rpgrip1l*[-/-] (n=3) and NACET-treated *rgprip1l*[-/-]

(n=3) fish in the SCO region (N=1). **(E)** Immunostaining for GFAP (green), Acetylated

tubulin (white) and LCP1 (red) on adult brain transverse sections at SCO level in

untreated (upper panels) and NACET-treated (lower panels) fish. Nuclei are stained

with DAPI (blue). No GFAP staining was present in controls (n=3) (upper left), while

untreated mutants displayed strong GFAP labeling (upper right) (n=3). NACET

treatment totally suppressed gliosis in a straight *rpgrip1l*[-/-] fish (lower right panel)

(n=1) and diminished the intensity of GFAP labeling in slightly tail-up *rpgrip1l*[-/-] (n=2).

Scale bar: 12,5 µm. **(F)** Graph presenting the percentage of GFAP positive cells

among the total number of SCO secretory cells compared between controls (n=3),

untreated *rpgrip1l*[-/-] (n=3), and NACET treated *rpgrip1l*[-/-] adult fish (n=3) (N=1). Each

dot represents the mean of % of GFAP-positive cells present in 3-6 sections per fish.

1600

Figure 8: Altered ciliogenesis, GFAP upregulation and increased number of

LCP1-positive cells around ventricles of *cep290*[-/-] juvenile fish at 4 wpf.

(A-F') Immunostaining of the cilia marker glutamylated tubulin (white) on 4 wpf brain

transverse sections at 3 antero-posterior SCO levels in 4 wpf control (A-C', n=3) and

cep290[-/-] scoliotic (D-F', n=3) fish. Nuclei are stained with DAPI (blue). White stars

indicate the diencephalic ventricle (DiV), above which lies the SCO. White rectangles

in C and F delineate the close-up on cilia (C', F'). Scale bar: 10 µm for A-F, 2,5 µm

for C'-F'. **(G-L)** Immunostaining for Acetylated Tubulin (G, J), GFAP (H, K) and LCP1

(I, L) on transverse sections of the brain at SCO level in controls (G-I) (n=3), and

cep290[-/-] (J-L) (n=3) juvenile (4 wpf) fish. In G and J, nuclei are stained with DAPI

(blue). Scale bar: identical to A-F. **(M)** Graph representing the percentage of GFAP-

positive cells among the total number of SCO secretory cells in control (n=3) and

scoliotic *cep290*^{-/-} (n=3) fish at scoliosis onset (N=1). Each dot represents the mean

ratio of GFAP-positive over total SCO secretory cells in 5-7 sections per fish. **(N)**

Quantification of LCP1+ cell density around the SCO in controls (n=3) and curved

cep290^{-/-} (n=3) fish at 4 wpf. 16 sections were counted for control, 18 sections for

cep290^{-/-} scoliotic fish. Each dot represents the mean value for one fish. Statistical

analysis was performed with unpaired T test where * means P-value < 0.05. Error

bars represent s.d. **(O-V)** Immunostaining for GFAP (white) on brain transverse

sections at cerebellum levels (P, R) and anterior hindbrain level (T, V) in control (P,

T) (n=3), and *cep290*^{-/-} (R, V) (n=3) fish. Corresponding nuclear staining (DAPI in

white) is shown above (O, Q, S, U). Scale bar: 125 µm.

1623

 SUPPLEMENTARY FIGURE LEGENDS

 Supplementary figure S1: Production and characterization of the *rpgrip1ᐃ* and

 ***rpgrip1l^ex4^* fish lines.**

 (A) Schematic exon-intron structure of the *Danio rerio rpgrip1l* gene. Two alleles

 were produced by CRISPR/cas9-mediated genome engineering, *rpgrip1l^ex4^* with a

 nonsense mutation in exon 4 and *rpgrip1l^ᐃ4-25^* (also called *rpgrip1l^ᐃ^* or *rpgrip1l*) with

 a large deletion between exons 4 and 25, encompassing most of the protein coding

 sequence. Both alleles had identical embryonic phenotype (panel B and data not

 shown). (A') Electrophoresis gel showing the migration profile of amplicons for each

 genotype, *rpgrip1l^ᐃ/ᐃ^*, *rpgrip1l^ᐃ/+^*, *rpgrip1^+l/+^* produced by triplex PCR (B) Absence of

 laterality defects and kidney cysts in *rpgrip1l^ᐃ/ᐃ^* and *rpgrip1l^ex4/ex4^* embryos. **(C)**

 Immunostaining on whole 30 hpf control (left panels) and *rpgrip1l^-/-^* (right panels)

 embryos to visualize the RF (anti-RF antibody, middle panels) and cilia (acetylated

 tubulin antibody, lower panels) in the caudal region. Nuclei (upper panels) are

 stained with DAPI. The RF is present in the central canal of *rpgrip1l^-/-^* embryos, as

 well as cilia in the whole neural tube width. Scale bar: 6 µm for upper panels; 15 µm

 for lower panels. **(D-E)** Absence of retinal morphological defects in 5 dpf *rpgrip1l^-/-^*

 larvae. Retinal sections of control (D) and *rpgrip1l^-/-^* (E) larvae were immunostained

 with Zpr3 (red) to label the external segment of photoreceptors and Acetylated

 Tubulin (Ac Tub) (green) to label axonemes. D' and E' are insets in D and E focusing

 on the photoreceptor layer. Scale bar: 50 µm. (**F-H'''**) Scanning electron microscopy

 of brain ventricles in control (F-F''), straight (str.) *rpgrip1l^-/-^* (G-G'') and scoliotic (sco.)

 *rpgrip1l^-/-^* (H-H''') adult (3 months) fish. Ependymal multiciliated cells of the

 rhombencephalic ventricle (RhV) are almost totally absent in *rpgrip1l^-/-^* scoliotic fish

 (I-I'), while they are present in straight r*pgrip1l^-/-^*(H-H') and controls (G-G''). Cilia of

 hindbrain mono-ciliated cells in *rpgrip1l^-/-^* (F'', G'') and controls (G''). Scale bars: 10

 µm for F-I, 2 µm for F'-H', 1 µm for F'', G'', H'''' and H'''.

Supplementary figure 2: Presence of scospondin aggregates and heterogenous ciliary defects in the fChP of *rpgrip1l*^/- juvenile fish.

(A-B') Immunostaining on transverse sections of the SCO, at anterior level (A, A', B) or posterior level (C, D) in *rpgrip1l*^+/+ (n=3) and tail-up *rpgrip1l*^/- (n=3), transgenic for Sspo-GFP (N=2). Sspo-GFP forms small cytoplasmic aggregates within SCO cells of both genotypes (* in A' and B'), under the primary cilia extending towards the DiV, and accumulates at the apical surface of tail-up *rpgrip1l*^/-, creating a double positive area for cilia and Sspo-GFP staining (yellow colour in B'). Multiciliated cells lateral to the posterior SCO (mcc) are negative for Sspo-GFP staining in controls (C) and absent in tail-up *rpgrip1l*^/- (D). **(E-J)** GFP immunostaining on transverse section of the fChP (E-G) or at the level of the optic tectum (H-J) in *rpgrip1l*^+/+ (n=7), straight *rpgrip1l*^/- (n=4) and scoliotic *rpgrip1l*^/- (n=5) 8 wpf fish transgenic for *sspo-GFP* (N=2). Scoliotic *rpgrip1l*^/- fish present with filamentous material and aggregates within the ventricles (G, J) while a small RF fragment is present within the ventricle of controls (H) and straight mutants (I) at optic tectum level. * indicates aggregates. **(K-O'')** Representative maximum intensity Z-stack projections of confocal transverse brain section at the level of the fChP in *rpgrip1l*^+/+ (K-K'') (n=7), straight *rpgrip1l*^/- (L-M'') (n=4) and scoliotic *rpgrip1l*^/- (N-O'') (n=5) 8 wpf fish (N=2). Cilia were stained with glutamylated-tubulin and acetylated-tubulin antibodies (both in white) and nuclei were stained with DAPI (blue). Sections shown are at the level of the habenular nuclei. At this level, monocilia in the dorsal midline region (K') and multicilia on lateral sides (K'') (n=7/7) are present in controls. Straight *rpgrip1l*^/- exhibited either normal (L', L'') (n=2/4) or abnormal and sparser (M', M'') (n=2/4) monocilia and multicilia. Similarly, scoliotic *rpgrip1l*^/- presented either normal monocilia and multicilia (N', N'') (n=3/5) or long monocilia (O') and decreased cilia density of multi-ciliated tufts (O'') (n=2/5). Scale bars: 20 µm.

Supplementary figure 3: Genetic invalidation of *urp2* in *rpgrip1l*^/- fish does not have any beneficial effect on scoliosis penetrance or severity.

(A-E) Absence of compensatory upregulation of *urp* family members in *urp2*^/- fish. Expression levels of *urp2* (A), *urp1* (B), *urp* (C), *uts2a* (D) and *uts2b* (E) in zebrafish in 13 wpf adults. Each dot represents the value measured for one fish; green and purple dots represent straight and curved fish, respectively. n = 5 [*rpgrip1l*^+/+*; urp2*^+/+],

9 [*rpgrip1l$^{+/+}$; urp2$^{-/}$*], 11 [*rpgrip1l$^{-/}$; urp2$^{+/}$*] and 10 [*rpgrip1l$^{-/}$; urp2$^{-/}$*]. Expression

levels for each gene are compared to the *lsm12b* housekeeping gene. Statistical

analysis was performed using the Tukey test where ns means not significant, ** P <

0.01, **** P < 0.0001. Error bars represent s.d. None of the *uts2* family members is

significantly upregulated in [*rpgrip1l$^{-/}$*; urp2$^{-/}$] fish compared to *rpgrip1l$^{-/}$*, suggesting

an absence of compensation. **(F-G)** Quantification of body axis curvature on dorsal

and lateral positions at 2 (F) and 4 (G) mpf. Each dot represents the curvature index

of one fish. Statistical analysis was done using Tukey's multiple comparisons test

where ****P < 0.0001, with error bars representing s.d. At 2 mpf, n=30 [*rpgrip1l$^{+/+}$;

urp2$^{+/+}$*], 26 [*rpgrip1l$^{+/+}$; urp2$^{-/}$*], 27 [*rpgrip1l$^{-/}$; urp2$^{+/+}$*], 55 [*rpgrip1l$^{-/}$; urp2$^{+/}$*] and 32

[*rpgrip1l$^{-/}$; urp2$^{-/}$*]. At 4 mpf, n=9 [*rpgrip1l$^{+/+}$; urp2$^{+/}$*], 7 [*rpgrip1l$^{+/+}$; urp2$^{-/}$*], 27

[*rpgrip1l$^{-/}$; urp2$^{+/+}$*], 48 [*rpgrip1l$^{-/}$; urp2$^{+/}$*] and 27 [*rpgrip1l$^{-/}$; urp2$^{-/}$*]. **(H)** Rose plots

representing combined Cobb angle measurements at 4 mpf of [*rpgrip1l$^{+/+}$; urp2$^{+/+}$*]

(n=7) in green, [*rpgrip1l$^{+/+}$; urp2$^{-/}$*] (n=6) in blue, [*rpgrip1l$^{-/}$; urp2$^{+/+}$*] (n=10) in pink

and [*rpgrip1l$^{-/}$; urp2$^{-/}$*] (n=7) in yellow. Cobb angle measurements were performed on

Alizarin red fish skeletons. One example of Alizarin red skeletons for each genotype

is presented below its corresponding rose-plot. **(I)** Illustration of combined curvature

index measurement from traced lines on lateral (left) and dorsal (right) views of fish.

The curvature index of both traced lines was measured using a modified MATLAB

program (LineCurvature2D function) which provided a measurement in radian/length

in pixels. Scale bar: 5 mm.

1709

Supplementary Figure 4: Complementary comparative transcriptome and

proteomic analysis of *rpgrip1l$^{-/}$* versus control fish.

(A, B) GO term analysis of biological processes enriched in *rpgrip1l$^{-/}$* for genes

upregulated in the trunk transcriptome data (A) or enriched in *rpgrip1l$^{-/}$* for proteins in

the brain adult proteome data (B) using Metascape. **(C)** List of the top 25 most

enriched proteins in the *rpgrip1l$^{-/}$* vs control brain proteome. D) List of selected genes

upregulated in the zebrafish *rpgrip1l* trunk transcriptome and found in different

reactive astrogliosis models in single cell transcriptome studies (Matusova et al,

2023).

1719

Supplementary figure 5: Increased GFAP and Annexin 2 staining and LCP1+ cell density in different brain regions.

(A-F') Immunostaining of Annexin A2 (Anxa2) (A, B, C, D, E, F) and GFAP (A', B', C', D', E', F') on transverse brain sections at habenula (A, A', D, D'), tectum (B, B', E, E') and hindbrain (C, C', F, F') levels in controls (A-C') and scoliotic *rpgrip1l*$^{-/-}$ (D-F') fish. Nuclei are stained with DAPI (blue). Scoliotic *rpgrip1l*$^{-/-}$ fish upregulate Annexin A2 and GFAP in overlapping territories. Scale bar: 20 µm. **(G)** Quantification of LCP1+ cell density in the optic tectum region of *rpgrip1l*$^{+/+}$ (n=4), straight *rpgrip1l*$^{-/-}$ (n=4) and scoliotic *rpgrip1l*$^{-/-}$ (n=3) 8 wpf fish. Each dot represents a fish. 15-20 sections were counted for each brain level. Straight *rpgrip1l*$^{-/-}$ fish sections displayed around twice more Lcp1+ cells at tectum level than controls, while scoliotic *rpgrip1l*$^{-/-}$ fish had a slightly increased LCP1+ cell density compared to controls. Statistical analysis was performed using Tukey's multiple comparisons test where ns means non significative and * Pvalue < 0.05, ** Pvalue < 0.01 with error bars represent s.d. **(H-J')** Immunostaining of LCP1+ immune cells on transverse sections of the optic tectum region of *rpgrip1l*$^{+/+}$ (H, H') (n=4), straight *rpgrip1l*$^{-/-}$ (I, I') (n=4) and scoliotic *rpgrip1l*$^{-/-}$ (J, J') (n=3) 8 wpf fish. Immune cells are labelled with the LCP1 antibody (magenta) and nuclei with DAPI (blue). H', I' and J' are magnifications of the squared regions in H, I and J, respectively. Scale bars: 100 µm for H, I, J; 30 µm for H', I', J'. str: straight, sco: scoliotic. **(K)** Quantification of LCP1+ cell density in the cerebellum region of 8 wpf fish (N=1). The analysis compared straight *rpgrip1l*$^{-/-}$ (n=4) and scoliotic *rpgrip1l*$^{-/-}$ (n=3) fish to *rpgrip1l*$^{+/+}$ siblings (n=4). No significant difference was found between genotypes. Statistical analysis was performed using Tukey's multiple comparisons test where ns means not significant. Error bars represent s.d. "str" means straight and "sco" means scoliotic.

Supplementary figure 6: Biological assay of NACET activity on *dnaaf1*$^{-/-}$ embryos and expression level of three immune markers at adult stage. (A) Theta angle (θ) measurement between the eye, the caudal end of the yolk extension and the tail extremity, corresponding to the embryo curvature index. **(B)** Lateral views of 60 hpf representative embryos showing axial curvature phenotypes. In untreated conditions, *dnaaf1*$^{-/-}$ embryos show a characteristic curly tail-down (CTD) phenotype, whereas *dnaaf1*$^{+/+}$ siblings have a straight axis. Treatment at 27 hpf with

3 mM NACET suppresses the CTD phenotype of $dnaaf1^{-/-}$ embryos and does not

affect axis straightness of $dnaaf1^{+/+}$ siblings. Embryos were measured around 2.5 to

3 mm. **(C-F)** Rose plots representing the (360 – θ) angle of $dnaaf1^{+/+}$ and $dnaaf1^{-/-}$ in

untreated and 3 mM NACET-treated conditions. Embryos were treated from 27hpf to

2.5 dpf and θ angles were measured at the end of the treatment. Untreated $dnaaf1^{+/+}$

are represented in (C – dark green), and treated ones are in (D – light green).

Untreated $dnaaf1^{-/-}$ are in (E – blue) and treated ones are in (F – red). NACET

treatment is able to rescue curly tail down phenotype in $dnaaf1^{-/-}$ embryos without

affecting the controls.

1762

1763 **Supplementary Movie 1: Micro-CT of juvenile spines.**

1764 Superposition of 3D-reconstructed spines of 5 wpf juvenile fish, 1 control (white), 3

1765 rpgrip1lΔ/Δ of different severities: 2 tail-up (yellow, blue) and 1 scoliotic (pink)

A

rpgrip1l⁻/⁻

5 dpf

5 wpf

12 wpf

B

Scoliosis appearance in clutches from
rpgrip1l⁻/⁺ **incrosses**

C

Scoliosis penetrance in adults

D Ctrl   **E** *rpgrip1l⁻/⁻*

Dorsal view

Lateral view

ctrl   *rpgrip1l⁻/⁻*   *rpgrip1l⁻/⁻*

H

Tol2 | Foxj1a enh. | 5XMyc-RPGRIP1L | polyA. | polyA | GFP. | cmlc2 enh. | Tol2

Tol2 | Col2a1 enh. | 5XMyc-RPGRIP1L | polyA. | polyA | GFP. | cmlc2 enh. | Tol2

I

Genotype	*rpgrip1l⁻/⁻*	*rpgrip1l⁻/⁻ ; Tg Foxj1a :RPGRIP1L*	*rpgrip1l⁻/⁻*	*rpgrip1l⁻/⁻ ; Tg Col2a1 :RPGRIP1L*
Adult Scoliotic fish (n)	72/74	0/63	28/29	28/29
Clutch (N)	N=2	N=2	N=1	N=1

J

DAPI Arl13b Glu-tub

Arl13b

Glu-tub

rpgrip1l[+/+]

rpgrip1l[-/-] str.

rpgrip1l[-/-] sco.

A A' A'' A'''

B B' B'' B'''

C C' C'' C'''

D

Cilia length mean along central canal (µm)

P value 0,09

**

*

rpgrip1l +/+ -/- str. -/- sco.

E

Cilia density mean along central canal

****

***

ns

rpgrip1l +/+ -/- str. -/- sco.

F

Central canal width (µm)

P Value 0,07

*

ns

rpgrip1l +/+ -/- str. -/- sco.

rpgrip1l[+/+] *rpgrip1l*[-/-] straight *rpgrip1l*[-/-] scoliotic

Anterior SCO

Post SCO

G I K

G' I' K'

H J L

Ac.tub/Glu.Tub

DAPI

A Brain

B Dorsal trunk

C

GO:0035082: axoneme assembly
GO:0001539: cilium or flagellum-dependent cell motility
GO:0002460: adaptive immune response
GO:0003351: epithelial cilium movement involved in extracellular fluid movement
GO:0007276: gamete generation
R-HSA-6785807: Interleukin-4 and Interleukin-13 signaling
GO:0045921: positive regulation of exocytosis

D Up-regulated genes

Trunk   Brain

560   54   6

E

Process/Pathway	Gene	Trunk.LogFC	Trunk P adj.	Brain.LogFC	Brain P adj.
Foxj1a and	*tbata*	2,25	8,82E-22	0,84	1,80E-07
Its targets	*foxj1a*	1,90	3,77E-23	1,36	9,67E-21
	enkur	1,88	8,98E-24	0,76	2,13E-06
	stpg1	1,80	1,10E-09	0,69	3,00E-05
	ccdc65	1,73	8,65E-11	0,74	1,25E-05
Regulators of	*pkd1b*	3,20	2,21E-47	1,90	1,08E-45
axis curvature	*urp1*	1,92	1,32E-16	-	-
	urp2	0,96	1,22E-03	0,87	5,44E-09
	sspo	1,18	4,16E-05	-	-
Immune	*irg1l*	2,45	1,20E-16	-	-
Response	*il11a*	2,23	3,09E-16	0,67	1,21E-05
	c4	1,97	1,50E-12	1,54	5,63E-29
	anxa2a	1,52	8,15E-17	1,53	7,04E-26
	mmp9	1,00	3,94E-04	0,46	4,02E-02
	c7a	1,86	8,82E-22	0,80	9,38E-07
	stat3	0,76	3,63E-07	0,80	5,23E-10

F Brain proteome

Adult rpgrip1l⁺/⁺

Scoliotic adult rpgrip1l⁻/⁻

DiV

RhV

DAPI GFAP

ANXA2

rpgrip1l⁺/⁺

straight rpgrip1l⁻/⁻

rpgrip1l⁻/⁻ scoliosis onset

DiV

RhV (cerebellar lev.)

DAPI GFAP

M

% of GFAP(+) cells among SCO cells

ns

***

**

rpgrip1l⁺/⁺   rpgrip1l⁻/⁻ str.   rpgrip1l⁻/⁻sco.

rpgrip1l⁺/⁺

straight rpgrip1l⁻/⁻

rpgrip1l⁻/⁻ scoliosis onset

DAPI LCP1

R

LCP1(+) cells density

P value 0,07

ns   ns

WT   rpgrip1l⁻/⁻ str.   rpgrip1l⁻/⁻ sco.

A **Scoliosis penetrance in *rpgrip1l*+/- incross**

B **Curvature index**

C **Scoliotic phenotype and associated curvature index of *rpgrip1l*-/- adults**

D **Lcp1 cells in SCO region**

E

F **% of GFAP(+) cells within the SCO**