Stomatal patterning is differently regulated in adaxial and abaxial epidermis in Arabidopsis

Pirko Jalakas, Ingmar Tulva, Nele Malvīne Bērziņa, Hanna Hõrak

Institute of Technology, University of Tartu, Nooruse 1, 50411, Tartu, Estonia

Abstract

Stomatal pores in leaves mediate CO_2 uptake into the plant and water loss via transpiration. Most plants are hypostomatous with stomata present only in the lower leaf surface (abaxial epidermis). Many herbs, including the model plant *Arabidopsis thaliana*, have substantial numbers of stomata also on the upper (adaxial) leaf surface. Studies of stomatal development have mostly focused on abaxial stomata and very little is known of adaxial stomatal formation. We addressed the role of leaf number in determination of stomatal density and stomatal ratio, and studied adaxial and abaxial stomatal patterns in mutants deficient in known abaxial stomatal development regulators. We found that stomatal density in some genetic backgrounds varies between different fully developed leaves and recommend using defined leaves for analyses of stomatal patterning. Our results indicate that stomatal development is at least partly independently regulated in adaxial and abaxial epidermis, as i) plants deficient in ABA biosynthesis and perception have increased stomatal ratios, ii) the *epf1epf2*, *tmm* and *sdd1* mutants have reduced stomatal ratios, iii) *erl2* mutants have increased adaxial but not abaxial stomatal index, and iv) stomatal precursors preferentially occur in abaxial epidermis. Further studies of adaxial stomata can reveal new insights into stomatal form and function.

Keywords: abscisic acid, abaxial, adaxial, amphistomaty, Arabidopsis, stomatal density, stomatal development, stomatal patterning, stomatal ratio

Introduction

Stomatal pores in plant leaves mediate CO_2 uptake for photosynthesis and water loss via transpiration. Stomatal numbers and size determine the maximal potential for stomatal conductance, a trait often positively related with yield (Roche, 2015). At the same time, lower stomatal densities are associated with increased water use efficiency and drought tolerance (Hughes *et al.*, 2017; Caine *et al.*, 2019; Dunn *et al.*, 2019). Hence, understanding how stomatal numbers are determined in leaves can help in breeding for varieties with increased productivity or water use efficiency.

Knowledge on stomatal formation in dicot leaves mostly originates from studies of abaxial epidermal development in the model plant *Arabidopsis thaliana* (Arabidopsis). Stomatal differentiation is achieved via sequential activation of the transcription factors SPEECHLESS (SPCH, MacAlister *et al.*, 2007; Pillitteri *et al.*, 2007), MUTE (MacAlister *et al.*, 2007; Pillitteri *et al.*, 2007) and FAMA (Ohashi-Ito and Bergmann, 2006). The central differentiation path is regulated by cell-to-cell signaling via peptides (epidermal patterning factors, EPFs) perceived by cell surface receptors to ensure optimal stomatal spacing. EPF1 and EPF2 peptides (Hara *et al.*, 2007; Hunt and Gray, 2009) are perceived by a membrane complex formed by a leucine-rich repeat receptor-like kinase from the ERECTA (ER) family (Shpak *et al.*, 2005) and the

receptor-like protein TOO MANY MOUTHS (TMM) (Yang and Sack, 1995; Lee *et al.*, 2012; Lin *et al.*, 2017). The ER family includes the ER, ERECTA-LIKE 1 (ERL1), and ERL2 proteins that have partly overlapping and partly distinct functions (Shpak *et al.*, 2005). ER is considered a major receptor for EPF2 and ERL1 for EPF1 (Lee *et al.*, 2012). The binding of EPF1 or EPF2 to the receptor complex activates downstream signaling that ultimately suppresses stomatal differentiation. Cleavage of pro-peptides is necessary to form biologically active signaling peptides; an apoplastic subtilisin-like serine protease STOMATAL DENSITY AND DISTRIBUTION 1 (SDD1) that suppresses stomatal development has been suggested to carry out this function (Berger and Altmann, 2000; Von Groll *et al.*, 2002). Loss-of-function mutations in *EPF1* and *EPF2*, *ER*-family genes, *TMM,* or *SDD1* lead to increased stomatal densities in Arabidopsis.

Abscisic acid (ABA) is a phytohormone produced in response to dry air or soil water deficit in plants (McAdam *et al.*, 2016; Kuromori *et al.*, 2022). Recently, ABA was shown to suppress stomatal development in Arabidopsis via the SUCROSE NONFERMENTING 1-related protein kinases SnRK2.2, SnRK2.3, and SnRK2.6 (OST1) that phosphorylate SPCH, leading to its degradation (Yang *et al.*, 2022*a*). ABA biosynthesis-deficient mutants *nced3nced5*, *aba2-2*, and *aba3-1* have higher stomatal densities or indices (Tanaka *et al.*, 2013*a*; Chater *et al.*, 2015; Jalakas *et al.*, 2018), while the ABA catabolism mutant *cyp707a1cyp707a3* has lower stomatal numbers (Tanaka *et al.*, 2013*a*; Jalakas *et al.*, 2018).

While most plants are hypostomatous with stomata located only in the abaxial epidermis (Salisbury and Oliver, 1928; Muir, 2015), Arabidopsis is an amphistomatous plant that produces a substantial proportion of its stomata in the adaxial epidermis (Berger and Altmann, 2000; Pantin *et al.*, 2013; Hronková *et al.*, 2015; Vráblová *et al.*, 2017; Tulva *et al.*, 2023, Preprint; Watts *et al.*, 2023, Preprint). Relatively little is known of the mechanisms of adaxial stomatal development and whether it is regulated by the same components and in a similar manner as stomatal formation in the abaxial epidermis. Previous studies suggest that adaxial and abaxial stomatal development are at least partly independently regulated. For example, stomatal density and index in the adaxial epidermis seem to be more responsive to changes in light conditions and relative air humidity (Hronková *et al.*, 2015; Devi and Reddy, 2018; Tulva *et al.*, 2023, Preprint). Stomatal ratio is altered in some Arabidopsis mutants lacking or overexpressing known regulators of stomatal development (Berger and Altmann, 2000; Kong *et al.*, 2012; Dow *et al.*, 2014; Franks *et al.*, 2015; Hronková *et al.*, 2015; Vráblová *et al.*, 2017; Qi *et al.*, 2019), pointing at differences between stomatal developmental mechanisms in the upper and lower leaf surface.

Here we address the within-plant variation of stomatal density and ratio in Arabidopsis and use mutants deficient in known regulators of stomatal development to test whether these signaling components similarly affect adaxial and abaxial stomatal development. We also analyze mutants with different ABA levels to understand if ABA affects adaxial and abaxial stomatal formation in a similar manner.

Materials and Methods

Plant lines and growth conditions

Arabidopsis thaliana accession Col-0 (wild type) and mutants in the same background were used for experiments, details for mutants used in experiments are listed in Table 1. T-DNA insertion mutants were from the SALK (Alonso *et al.*, 2003) and GABI-Kat collections (Rosso *et al.*, 2003) and ordered from Nottingham Arabidopsis Stock Centre (NASC, (Scholl *et al.*, 2000)). Primers used for genotyping studied mutants are shown in Supplementary Table 1. We note that the *tmm* T-DNA insertion line SALK_115723C used in our study also harbors a T-DNA insertion in *STKR1* (Nietzsche *et al.*, 2018). Although this line is a *tmm stkr1* double mutant, its stomatal phenotypes are similar to previously published results for other *tmm* alleles, hence we refer to the mutant as *tmm*.

For experiments addressing differences between stomatal anatomical traits in different leaves (Figure 1), plants were grown in growth cabinets (Percival AR-22L, Percival Scientific, IA, USA) at 10 h light : 14 h dark photoperiod (light intensity 250 µmol m^{-2} s^{-1}), relative air humidity 60% during day and 80% during night and temperature 23°C and 19°C during day and night, respectively. For experiments addressing stomatal density, index and ratio in leaf 8 of different stomatal development mutants (Figures 2-4), plants were grown in a growth cabinet (Microclima Arabidopsis MCA1600-3LP6-E, Snijders Scientific, Tilburg, Netherlands) under the same conditions, with the exception of uniform relative air humidity of 70% during both day and night.

Stomatal patterning analyses

Leaves to be sampled for anatomical analyses were numbered according to the protocol of Farmer *et al.*, (2013). In experiments addressing stomatal patterning in different leaves, leaves 6 and 8 were sampled from plants at 7 weeks of age, and leaves 10 and 12 were sampled as plants had started bolting at 8 weeks of age from different plants grown in parallel. In experiments addressing stomatal patterning in leaf 8 of different mutants, leaves were sampled at 6 weeks of age. The sampled leaves were cut in half at the midvein and leaf impressions were collected from both the adaxial and abaxial leaf side, one from each leaf half (Figure 1A) with dental silicone (Speedex light body, Coltene/Whaledent AG; Alstätte, Switzerland and oranwash L, Zhermack; www.zhermack.com for adaxial and abaxial leaf side, respectively) as in Casson *et al.*, (2009). Secondary nail varnish imprints were taken from the silicone and transferred to microscope slides with transparent tape. One area of 0.26 mm^{-2} from each imprint was photographed under a microscope (Kern OBF 133; Kern & Sohn GmbH; www.kern-sohn.com) at 200x magnification. Stomatal density (stomata mm^{-2}, including both mature stomata and stomatal precursor cells), stomatal ratio (ratio of adaxial and abaxial stomatal densities), and stomatal index (proportion of stomata from all epidermal cells) were determined from the images using ImageJ software (National Institutes of Health, USA; Schneider et al. 2012).

Rosette area analysis

In experiments addressing stomatal patterning in leaf 8 of different mutants, plants were photographed at 4 weeks of age and the projected rosette area was determined with ImageJ software (Schneider *et al.*, 2012).

Statistical analyses

Pooled data from two independently grown batches of plants is shown for all experiments. Statistical analyses were carried out with GraphPad Prism version 10 (GraphPad Software, Boston, MA, USA). Two-way ANOVA with Tukey post hoc test, one-way Welch ANOVA with Dunnett T3 post hoc test, or linear regression were used as indicated in Figure legends. All effects were considered significant at $p < 0.05$.

Results

Stomatal density depends on leaf number and ABA preferentially suppresses adaxial stomatal development

Most studies addressing stomatal density in true leaves sample an unspecified fully developed leaf for analysis. As Arabidopsis fully-developed leaves vary in shape and size, it is possible that developmentally distinct leaves also differ in their stomatal patterning. To address the potential within-plant variation in stomatal densities and in the distribution of stomata between adaxial and abaxial leaf surfaces, we analyzed stomatal density and ratio in leaves 6, 8, 10, and 12 in Arabidopsis (Figure 1A). To test if ABA regulates stomatal distribution between adaxial and abaxial leaf surfaces, we included mutants with higher (*cyp707a1/a3*) or lower (*nced3/5*) ABA levels, or impaired ABA perception *(pyrpyl112458)* in our analysis.

Both adaxial and abaxial stomatal density was increased in all studied leaves of the *nced3/5* and *pyrpyl112458* mutants compared with wild type plants (Figure 1B,C), in line with generally increased abaxial stomatal density of ABA biosynthesis and perception mutants (Chater *et al.*, 2015; Hsu *et al.*, 2018; Jalakas *et al.*, 2018; Merilo *et al.*, 2018). Although other studies have reported a decrease in stomatal density in the *cyp707a1/a3* mutants (Tanaka *et al.*, 2013*a*; Jalakas *et al.*, 2018), in our experiment it was significant only in leaf 8 (Figure 1B, C). While stomatal densities of most mutants were similar in different leaves, the ABA perception mutant *pyrpyl112458* had higher adaxial and abaxial stomatal densities in leaves 10 and 12 compared with leaf 6 (Figure 1B, C). Thus, stomatal density depends on leaf number in some Arabidopsis genotypes.

Stomatal ratios were similar across different leaves in all studied mutants (Figure 1D). The ABA biosynthesis mutant *nced3/5* and ABA perception mutant *pyrpyl112458* had slightly higher stomatal ratios compared with Col-0 and *cyp707a1/a3* (significant main effect of genotype, Table 2), indicating that ABA preferentially suppresses stomatal development in adaxial epidermis.

SDD1, EPF1/2, ERL2, and TMM differently affect stomatal development in adaxial and abaxial epidermis

The increase in stomatal ratio in ABA biosynthesis- or perception-deficient mutants indicates that adaxial and abaxial stomatal development are at least partly independently regulated. We aimed to test whether major regulators of abaxial stomatal development similarly affect adaxial stomatal development. To this end, we obtained T-DNA insertion lines for key components of the stomatal development signaling pathway (Table 1, Figure 2A) and analyzed stomatal density and index in both adaxial and abaxial epidermis of leaf 8 in these mutants. The components studied in our experiment included the SDD1 protease, the signaling peptides EPF1 and EPF2, and the peptide receptor complex components ER, ERL1, ERL2, and TMM (Figure 2A). We also included mutants with lower or higher ABA levels (*nced3/5*, *cyp707a1a3*) and a mutant combining low ABA levels with lack of EPF1 and EPF2 (*nced3/5 epf1/2*).

Stomatal ratio was significantly lower than in wild type in the *epf1/2*, *sdd1-4* and *tmm* mutants (Figure 2B), indicating that respective signaling components more prominently suppress stomatal development in the abaxial than in the adaxial epidermis. Similarly reduced stomatal ratios have been found before in *epf1/2* (Franks *et al.*, 2015), *sdd1* (Berger and Altmann, 2000; Vráblová *et al.*, 2017), and *tmm* mutants (Hronková *et al.*, 2015; Vráblová *et al.*, 2017). Most studied mutations affected stomatal density in adaxial and abaxial epidermis in a similar manner: stomatal density was increased compared with wild type by the *nced3/5*, *epf1/2*, *nced3/5 epf1/2*, *sdd1* and *er* mutations, and not significantly affected by the *cyp707a1/a3*, *erl1* and *erl2* mutations (Figure 2C, 2D). In line with previous works (Hronková *et al.*, 2015; Vráblová *et al.*, 2017), stomatal density was increased only in the abaxial epidermis in the *tmm* mutant (Figure 2C, 2D), indicating a minor role for TMM in adaxial stomatal development. Stomatal index was higher than in wild type in both adaxial and abaxial epidermis in the *epf1/2*, *nced3/5 epf1/2*, *sdd1*, and *sdd1-4* mutants (Figure 2E, 2F), whereas it was higher only in the adaxial epidermis in both studied *erl2* mutants (Figure 2E), and higher only in the abaxial epidermis in the *tmm* mutant (Figure 2F). Stomatal index was not significantly affected in the *cyp707a1a3* and *nced3/5* mutants (Figure 2E, 2F), indicating that altered plant ABA levels do not affect the proportion of stomata from all epidermal cells in leaf 8. Together, our data indicate that SDD1, EPF1/2 and TMM are more important for regulating stomatal development in the abaxial than in the adaxial epidermis, whereas ERL2 has a relatively more important role in adaxial than in abaxial epidermal patterning.

Stomatal development rarely arrests in the precursor state in adaxial epidermis

In plant lines with aberrant stomatal development, stomata sometimes arrest in a precursor state. In most studied mutants, we did not detect more stomatal precursors than in wild type plants (Supplementary Figure 1). When stomatal precursors were found in fully-developed leaf 8, such cells usually only occurred in the abaxial epidermis. This was true for the *epf1/2*, and *er* mutants, as observed before (Hunt and Gray, 2009; Lee *et al.*, 2012), whereas we found stomatal precursors in both adaxial and abaxial epidermis in the *nced3/5 epf1/2* mutant (Figure 3). Thus, stomatal development rarely arrests in the precursor state in the adaxial epidermis, indicating different regulation of adaxial and abaxial stomatal differentiation.

Stomatal density is negatively related with plant growth

Total stomatal density (sum of adaxial and abaxial stomatal densities) was significantly increased in the *nced3/5*, *nced3/5 epf1/2*, *epf1/2*, *sdd1*, *er* and *tmm* mutants, with the most prominent effects in *epf1/2*, *sdd1* and *epf1/2 nced3/5* mutants (Figure 4A). The projected rosette area was significantly reduced in the mutant with the highest stomatal density – the *nced3/5 epf1/2* double mutant (Figure 4B). Across all studied mutants, there was a significant negative relationship between stomatal density and plant size (Figure 4C), similar to our previous study (Tulva *et al.*, 2023, Preprint).

Discussion

Here we addressed within-plant variation in stomatal density and ratio in Arabidopsis and the role of major stomatal developmental regulators in stomatal patterning in adaxial and abaxial epidermis. We show that while stomatal ratio is similar across developmentally different leaves, stomatal density in some Arabidopsis genotypes depends on the sampled leaf (Figure 1). We also found that a lack of certain stomatal development regulators leads to different developmental outcomes in upper and lower leaf surfaces (Figures 2, 3), indicating independent regulation of adaxial and abaxial stomatal development.

In wild type Arabidopsis, stomatal density was similar in developmentally different fully-developed leaves, whereas in the ABA perception mutant *pyrpyl112458*, stomatal densities were significantly higher in fully-developed younger leaves (Figure 1B, C). Thus, to reduce variation in stomatal density analyses in true leaves, we recommend sampling a defined fully-developed leaf for analyses of stomatal patterning. Stomatal ratio did not differ between leaves in any of our studied genotypes (Figure 1D). However, the stomatal ratio values that we measured in true leaves of wild-type plants were higher than reported before in cotyledons (Geisler *et al.*, 1998; Berger and Altmann, 2000; Delgado *et al.*, 2019), indicating that there is within-plant variation also in leaf stomatal ratio. Differences between stomatal characteristics depending on leaf number have been discussed before, stressing more severe effects of different mutations on stomatal development in cotyledons than in true leaves (Dow *et al.*, 2014). Understanding how stomatal development between cotyledons and true leaves differs is important for practical applications aiming to regulate plant water use efficiency or productivity via modulation of stomatal numbers, as cotyledons have a minor role in overall plant gas-exchange and physiology.

The lack of negative regulators of stomatal development increased stomatal density more strongly in the abaxial epidermis, leading to reduced stomatal ratios in *epf1/2*, *sdd1* and *tmm* mutants (Figure 2). Similarly reduced stomatal ratios have been found before for these mutants (Berger and Altmann, 2000; Dow *et al.*, 2014; Franks *et al.*, 2015; Hronková *et al.*, 2015; Vráblová *et al.*, 2017), indicating a more prominent role of respective signaling components in adjusting abaxial than adaxial stomatal development. More severely disrupted abaxial than adaxial stomatal patterns have also been found in plants with modified expression levels of SPCH (Dow *et al.*, 2014), a repressor of SPCH (IDD16, Qi *et a*l., 2019), or STOMAGEN/EPFL9 (Hronková *et al.*, 2015). As studies of stomatal form and function have largely focused on abaxial stomata, it is expected that the currently identified stomatal developmental signaling

components are of major importance in abaxial epidermal patterning. Focusing on the adaxial stomatal development in the future might reveal new signaling components or pathways in stomatal development. Several studies indicate that adaxial stomatal development is more sensitive to changes in environmental conditions, such as light levels (Hronková *et al.*, 2015) or relative air humidity (Devi and Reddy, 2018; Tulva *et al.*, 2023, Preprint). Between-species comparisons also suggest that adaxial stomatal numbers have been adjusted more freely during evolution (Muir *et al.*, 2023). The presence of adaxial stomata in amphistomatic leaves has been associated with improved CO_2 uptake (Parkhurst and Mott, 1990; Drake *et al.*, 2019), photosynthesis (Xiong and Flexas, 2020) and faster stomatal responsiveness (Haworth *et al.*, 2018). Understanding how adaxial stomatal development is adjusted in plants is necessary to better understand the physiology of amphistomatous plants.

We found that the lack of ERL2 alone was sufficient to increase adaxial but not abaxial stomatal index (Figure 2), indicating that ERL2 is more involved in suppressing adaxial than abaxial stomatal development. Indeed, similarly increased adaxial but not abaxial stomatal index was found before in the *erl1erl2* double mutant (Kong *et al.*, 2012). Our findings are in line with previous evidence showing distinct functions of specific ER family proteins in fine-tuning stomatal development. The specific roles of ER, ERL1 and ERL2 manifest in their different expression patterns (Shpak *et al.*, 2004), binding of different EPFs (Lee *et al.*, 2012), and different regulation of their subcellular localization (Yang *et al.*, 2022*b*). Our findings indicate that specific ER family protein members also contribute to a different extent to stomatal differentiation in adaxial and abaxial epidermis.

Stomatal precursor cell formation was shown to cease earlier in adaxial compared with abaxial epidermis in cotyledons (Geisler and Sack, 2002). Here, stomatal precursor cells, if present in fully-developed leaves of stomatal development mutants, were typically found only in the abaxial epidermis (Figure 3). This is in line with prolonged stomatal initiation in abaxial epidermis. As an exception, the *nced3/5 epf1/2* mutant had arrested stomatal precursor cells both in the adaxial and abaxial epidermis of fully-developed leaves (Figure 3). Thus, low ABA levels in the *epf1/2* mutant background led to increased numbers of stomatal precursors in the adaxial epidermis, indicating that normal ABA levels can rescue stomatal differentiation in the adaxial epidermis of the *epf1/2* mutant that only harbors stomatal precursors in the abaxial epidermis. The asymmetry of the presence of cells arrested in a stomatal precursor state between adaxial and abaxial epidermis indicates that different mechanisms control stomatal differentiation in adaxial and abaxial epidermis.

ABA is considered a suppressing signal for stomatal development largely based on studies of ABA biosynthesis and catabolism mutants. Here, we found no effect of the *cyp707a1/a3* mutations on stomatal density or index (Figure 2). This difference between our and previous studies (Tanaka *et al.*, 2013*a*; Jalakas *et al.*, 2018) might in part be explained by variable ABA levels in ABA catabolism mutants under different growth conditions as has been found before for some ABA biosynthesis mutants (Merilo *et al.*, 2018). The ABA-deficient *nced3/5* mutant had higher stomatal density in our experiments as observed before (Chater *et al.*, 2015) but as there was no change in stomatal index, the change in density might result from the effects of ABA-deficiency on cell growth rather than stomatal development. Stomatal ratio was slightly

increased in plants deficient in ABA biosynthesis or perception (Figure 1D, Table 2), indicating that ABA might preferentially suppress stomatal development in the adaxial epidermis. Together, our data suggest that in true leaves, ABA levels have a minor effect on the proportion of cells entering the stomatal lineage and on the distribution of stomata between leaf surfaces, whereas mutations in stomatal development pathway components have much larger effects.

Studies finding a role for ABA levels in regulating stomatal index have often been carried out in cotyledons of plate-grown seedlings subjected to 100% relative air humidity (Tanaka *et al.*, 2013*a*; Mohamed *et al.*, 2023). Leaf ABA levels depend on air humidity (McAdam *et al.*, 2016) and low relative air humidity primes ABA sensitivity in leaves (Pantin *et al.*, 2013). Thus, it is expected that the quantitative effects of ABA on stomatal patterning are different between plants that have not been subjected to naturally fluctuating air humidity and those that have. Our experiments suggest that in true leaves grown under moderate relative air humidity, modification of ABA levels in the range achieved by the ABA biosynthesis and catabolism mutants is of minor importance in regulating entry into the stomatal lineage.

We previously found a negative relationship between plant growth and stomatal density in plant lines, where stomatal density was altered by the *epf1* and/or *epf2* mutations (Tulva *et al.*, 2023, Preprint). Here, we show that this relationship is more universal, as it holds for a set of plant lines with varying stomatal density due to different mutations (Figure 4C). Despite the improved CO_2 uptake capacity and increased photosynthesis found before in high stomatal density mutants (Schlüter *et al.*, 2003; Tanaka *et al.*, 2013*b*; Tulva *et al.*, 2023, Preprint), such mutants fail to achieve better growth. Possibly, the resources obtained by increased net assimilation are used to carry out expensive cell divisions (Dow *et al.*, 2014; Franks *et al.*, 2015; Bowers, 2018), or for investment into roots to improve water uptake capacity, as discussed before (Hepworth *et al.*, 2016). Future studies of above- and below-ground growth in plants with altered stomatal conductance due to different stomatal apertures or densities are needed to test the latter hypothesis.

Acknowledgements

We thank Agnes Vaht, Marilin Poogen and Tamta Jashi for help with plant growth and sample collection, Elfi Aliyeva for help with sample imaging, and Kārlis Dieviņš for help with projected rosette area measurements.

Author contributions

Conceptualization - P.J., I.T., H.H., Data curation and analysis - all authors; Funding Acquisition - H.H.; Investigation - all authors; Supervision - I.T., H.H.; Visualization - P.J.; Writing – Original Draft - P.J., H.H.; Writing – Review & Editing - all authors

Conflict of interest

No conflict of interest declared.

Funding

This work was supported by the Estonian Research Council grants PSG404 and MOBERC104.

Data availability

Primary data for figures and tables presented in this manuscript are included as supplementary material.

References

Alonso JM, Stepanova AN, Leisse TJ, *et al.* 2003. Genome-Wide Insertional Mutagenesis of Arabidopsis thaliana. Science **301**, 653–657.

Berger D, Altmann T. 2000. A subtilisin-like serine protease involved in the regulation of stomatal density and distribution in Arabidopsis thaliana. Genes & Development **14**, 1119–1131.

Bowers RD. 2018. The Economics of Stomatal Development.

Caine RS, Yin X, Sloan J, *et al.* 2019. Rice with reduced stomatal density conserves water and has improved drought tolerance under future climate conditions. The New Phytologist **221**, 371–384.

Casson SA, Franklin KA, Gray JE, Grierson CS, Whitelam GC, Hetherington AM. 2009. phytochrome B and *PIF4* Regulate Stomatal Development in Response to Light Quantity. Current Biology **19**, 229–234.

Chater C, Peng K, Movahedi M, *et al.* 2015. Elevated CO_2-Induced Responses in Stomata Require ABA and ABA Signaling. Current Biology **25**, 2709–2716.

Delgado D, Sánchez-Bermejo E, de Marcos A, Martín-Jimenez C, Fenoll C, Alonso-Blanco C, Mena M. 2019. A Genetic Dissection of Natural Variation for Stomatal Abundance Traits in Arabidopsis. Frontiers in Plant Science **10**.

Devi MJ, Reddy VR. 2018. Transpiration Response of Cotton to Vapor Pressure Deficit and Its Relationship With Stomatal Traits. Frontiers in Plant Science **9**.

Dow GJ, Berry JA, Bergmann DC. 2014. The physiological importance of developmental mechanisms that enforce proper stomatal spacing in Arabidopsis thaliana. New Phytologist **201**, 1205–1217.

Drake PL, de Boer HJ, Schymanski SJ, Veneklaas EJ. 2019. Two sides to every leaf: water and CO_2 transport in hypostomatous and amphistomatous leaves. New Phytologist **222**, 1179–1187.

Dunn J, Hunt L, Afsharinafar M, Meselmani MA, Mitchell A, Howells R, Wallington E, Fleming AJ, Gray JE. 2019. Reduced stomatal density in bread wheat leads to increased water-use efficiency. Journal of Experimental Botany **70**, 4737–4748.

Farmer E, Mousavi S, Lenglet A. 2013. Leaf numbering for experiments on long distance signalling in Arabidopsis. doi: 10.1038/protex.2013.071.

Franks PJ, W. Doheny-Adams T, Britton-Harper ZJ, Gray JE. 2015. Increasing water-use efficiency directly through genetic manipulation of stomatal density. New Phytologist **207**, 188–195.

Frey A, Effroy D, Lefebvre V, Seo M, Perreau F, Berger A, Sechet J, To A, North HM, Marion-Poll A. 2012. Epoxycarotenoid cleavage by NCED5 fine-tunes ABA accumulation and affects seed dormancy and drought tolerance with other NCED family members. The Plant Journal **70**, 501–512.

Geisler MJ, Sack FD. 2002. Variable timing of developmental progression in the stomatal pathway in Arabidopsis cotyledons. New Phytologist **153**, 469–476.

Geisler M, Yang M, Sack FD. 1998. Divergent regulation of stomatal initiation and patterning in organ and suborgan regions of the *Arabidopsis* mutants *too many mouths* and *four lips*. Planta

205, 522–530.

Gonzalez-Guzman M, Pizzio GA, Antoni R, *et al.* 2012. Arabidopsis PYR/PYL/RCAR Receptors Play a Major Role in Quantitative Regulation of Stomatal Aperture and Transcriptional Response to Abscisic Acid. The Plant Cell Online **24**, 2483–2496.

Hara K, Kajita R, Torii KU, Bergmann DC, Kakimoto T. 2007. The secretory peptide gene EPF1 enforces the stomatal one-cell-spacing rule. Genes & Development **21**, 1720–1725.

Haworth M, Scutt CP, Douthe C, Marino G, Gomes MTG, Loreto F, Flexas J, Centritto M. 2018. Allocation of the epidermis to stomata relates to stomatal physiological control: Stomatal factors involved in the evolutionary diversification of the angiosperms and development of amphistomaty. Environmental and Experimental Botany **151**, 55–63.

Hepworth C, Turner C, Landim MG, Cameron D, Gray JE. 2016. Balancing Water Uptake and Loss through the Coordinated Regulation of Stomatal and Root Development. PLOS ONE **11**, e0156930.

Hronková M, Wiesnerová D, Šimková M, Skůpa P, Dewitte W, Vráblová M, Zažímalová E, Šantrůček J. 2015. Light-induced STOMAGEN-mediated stomatal development in *Arabidopsis* leaves. Journal of Experimental Botany **66**, 4621–4630.

Hsu P-K, Takahashi Y, Munemasa S, Merilo E, Laanemets K, Waadt R, Pater D, Kollist H, Schroeder JI. 2018. Abscisic acid-independent stomatal CO_2 signal transduction pathway and convergence of $_2$ and ABA signaling downstream of OST1 kinase. Proceedings of the National Academy of Sciences **115**, E9971–E9980.

Hughes J, Hepworth C, Dutton C, Dunn JA, Hunt L, Stephens J, Cameron D, Waugh R, Gray JE. 2017. Reducing stomatal density in barley improves drought tolerance without impacting on yield. Plant Physiology, pp.01844.2016.

Hunt L, Gray JE. 2009. The Signaling Peptide EPF2 Controls Asymmetric Cell Divisions during Stomatal Development. Current Biology **19**, 864–869.

Jalakas P, Merilo E, Kollist H, Brosché M. 2018. ABA-mediated regulation of stomatal density is OST1-independent. Plant Direct **2**, e00082.

Kong D, Karve R, Willet A, Chen M-K, Oden J, Shpak ED. 2012. Regulation of Plasmodesmatal Permeability and Stomatal Patterning by the Glycosyltransferase-Like Protein KOBITO1. Plant Physiology **159**, 156–168.

Kuromori T, Fujita M, Takahashi F, Yamaguchi-Shinozaki K, Shinozaki K. 2022. Inter-tissue and inter-organ signaling in drought stress response and phenotyping of drought tolerance. The Plant Journal **109**, 342–358.

Kushiro T, Okamoto M, Nakabayashi K, Yamagishi K, Kitamura S, Asami T, Hirai N, Koshiba T, Kamiya Y, Nambara E. 2004. The Arabidopsis cytochrome P450 CYP707A encodes ABA 8′-hydroxylases: key enzymes in ABA catabolism. The EMBO Journal **23**, 1647–1656.

Lee JS, Kuroha T, Hnilova M, Khatayevich D, Kanaoka MM, McAbee JM, Sarikaya M, Tamerler C, Torii KU. 2012. Direct interaction of ligand–receptor pairs specifying stomatal patterning. Genes & Development **26**, 126–136.

Lin G, Zhang L, Han Z, Yang X, Liu W, Li E, Chang J, Qi Y, Shpak ED, Chai J. 2017. A receptor-like protein acts as a specificity switch for the regulation of stomatal development. Genes & Development **31**, 927–938.

MacAlister CA, Ohashi-Ito K, Bergmann DC. 2007. Transcription factor control of asymmetric cell divisions that establish the stomatal lineage. Nature **445**, 537–540.

McAdam SAM, Sussmilch FC, Brodribb TJ. 2016. Stomatal responses to vapour pressure deficit are regulated by high speed gene expression in angiosperms. Plant, Cell & Environment **39**, 485–491.

Merilo E, Yarmolinsky D, Jalakas P, Parik H, Tulva I, Rasulov B, Kilk K, Kollist H. 2018. Stomatal VPD Response: There Is More to the Story Than ABA. Plant Physiology **176**, 851–864.

Mohamed D, Vonapartis E, Corcega DY, Gazzarrini S. 2023. ABA guides stomatal proliferation and patterning through the EPF-SPCH signaling pathway in Arabidopsis thaliana. Development, dev.201258.

Muir CD. 2015. Making pore choices: repeated regime shifts in stomatal ratio. Proceedings of the Royal Society B: Biological Sciences **282**, 20151498.

Muir CD, Conesa MÀ, Galmés J, Pathare VS, Rivera P, López Rodríguez R, Terrazas T, Xiong D. 2023. How Important Are Functional and Developmental Constraints on Phenotypic Evolution? An Empirical Test with the Stomatal Anatomy of Flowering Plants. The American Naturalist **201**, 794–812.

Nanda AK, El Habti A, Hocart CH, Masle J. 2019. ERECTA receptor-kinases play a key role in the appropriate timing of seed germination under changing salinity. Journal of Experimental Botany **70**, 6417–6435.

Nietzsche M, Guerra T, Alseekh S, Wiermer M, Sonnewald S, Fernie AR, Börnke F. 2018. STOREKEEPER RELATED1/G-Element Binding Protein (STKR1) Interacts with Protein Kinase SnRK1. Plant Physiology **176**, 1773–1792.

Ohashi-Ito K, Bergmann DC. 2006. *Arabidopsis* FAMA Controls the Final Proliferation/Differentiation Switch during Stomatal Development. The Plant Cell **18**, 2493–2505.

Okamoto M, Kuwahara A, Seo M, Kushiro T, Asami T, Hirai N, Kamiya Y, Koshiba T, Nambara E. 2006. CYP707A1 and CYP707A2, Which Encode Abscisic Acid 8′-Hydroxylases, Are Indispensable for Proper Control of Seed Dormancy and Germination in Arabidopsis. Plant Physiology **141**, 97–107.

Pantin F, Renaud J, Barbier F, *et al.* 2013. Developmental Priming of Stomatal Sensitivity to Abscisic Acid by Leaf Microclimate. Current Biology **23**, 1805–1811.

Parkhurst DF, Mott KA. 1990. Intercellular Diffusion Limits to CO_2 Uptake in Leaves 1: Studies in Air and Helox. Plant Physiology **94**, 1024–1032.

Pillitteri LJ, Sloan DB, Bogenschutz NL, Torii KU. 2007. Termination of asymmetric cell division and differentiation of stomata. Nature **445**, 501–505.

Qi S-L, Lin Q-F, Feng X-J, Han H-L, Liu J, Zhang L, Wu S, Le J, Blumwald E, Hua X-J. 2019. IDD16 negatively regulates stomatal initiation via trans-repression of *SPCH* in *Arabidopsis*. Plant Biotechnology Journal **17**, 1446–1457.

Roche D. 2015. Stomatal Conductance Is Essential for Higher Yield Potential of C3 Crops. Critical Reviews in Plant Sciences **34**, 429–453.

Rosso MG, Li Y, Strizhov N, Reiss B, Dekker K, Weisshaar B. 2003. An Arabidopsis thaliana T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. Plant Molecular Biology **53**, 247–259.

Salisbury EJ, Oliver FW. 1928. I. On the causes and ecological significance of stomatal frequency, with special reference to the woodland flora. Philosophical Transactions of the Royal Society of London. Series B, Containing Papers of a Biological Character **216**, 1–65.

Schlüter U, Muschak M, Berger D, Altmann T. 2003. Photosynthetic performance of an *Arabidopsis* mutant with elevated stomatal density (*sdd1‑1*) under different light regimes. Journal of Experimental Botany **54**, 867–874.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nature Methods **9**, 671–675.

Scholl RL, May ST, Ware DH. 2000. Seed and Molecular Resources for Arabidopsis. Plant Physiology **124**, 1477–1480.

Shpak ED, Berthiaume CT, Hill EJ, Torii KU. 2004. Synergistic interaction of three ERECTA-family receptor-like kinases controls Arabidopsis organ growth and flower development by promoting cell proliferation. Development **131**, 1491–1501.

Shpak ED, McAbee JM, Pillitteri LJ, Torii KU. 2005. Stomatal Patterning and Differentiation by Synergistic Interactions of Receptor Kinases. Science **309**, 290–293.

Tanaka Y, Nose T, Jikumaru Y, Kamiya Y. 2013*a*. ABA inhibits entry into stomatal-lineage

development in Arabidopsis leaves. The Plant Journal **74**, 448–457.

Tanaka Y, Sugano SS, Shimada T, Hara-Nishimura I. 2013*b*. Enhancement of leaf photosynthetic capacity through increased stomatal density in Arabidopsis. New Phytologist **198**, 757–764.

Tulva I, Koolmeister K, Horak H. 2023. Low relative air humidity and increased stomatal density independently hamper growth in young Arabidopsis. bioRxiv.

Von Groll U, Berger D, Altmann T. 2002. The subtilisin-like serine protease SDD1 mediates cell-to-cell signaling during Arabidopsis stomatal development. The Plant Cell **14**, 1527–1539.

Vráblová M, Vrábl D, Hronková M, Kubásek J, Šantrůček J. 2017. Stomatal function, density and pattern, and CO_2 assimilation in Arabidopsis thaliana tmm1 and sdd1-1 mutants. Plant Biology **19**, 689–701.

Watts JL, Dow GJ, Buckley TN, Muir CD. 2023. Does stomatal patterning in amphistomatous leaves minimize the CO_2 diffusion path length within leaves? bioRxiv.

Xiong D, Flexas J. 2020. From one side to two sides: the effects of stomatal distribution on photosynthesis. New Phytologist **228**, 1754–1766.

Yang X, Gavya S L, Zhou Z, Urano D, Lau OS. 2022*a*. Abscisic acid regulates stomatal production by imprinting a SnRK2 kinase–mediated phosphocode on the master regulator SPEECHLESS. Science Advances **8**, eadd2063.

Yang M, Sack FD. 1995. The *too many mouths* and *four lips* mutations affect stomatal production in Arabidopsis. The Plant Cell **7**, 2227–2239.

Yang K-Z, Zuo C-R, Leng Y-J, Yue J-L, Liu H-C, Fan Z-B, Xue X-Y, Dong J, Chen L-Q, Le J. 2022*b*. The functional specificity of ERECTA-family receptors in Arabidopsis stomatal development is ensured by molecular chaperones in the endoplasmic reticulum. Development **149**, dev200892.

Tables

Table 1. Mutants used in experiments

Mutant name	Mutant short name	Mutation/T-DNA insertion line name	Reference
nced3-2nced5-2	*nced3/5*	GK-129B08, GK-328D05	(Frey *et al.*, 2012)
epf1-1epf2-2	*epf1/2*	SALK_137549, GK-673E01	(Hunt and Gray, 2009)
nced3-2nced5-2epf1-1epf2-2	*nced3/5 epf1/2*	GK-129B08, GK-328D05, SALK_137549, GK-673E01	created by crossing *nced3-2nced5-2* and *epf1-1epf2-2* in this study
cyp707a1-1cyp707a3	*cyp707a1/a3*	SALK_069127, SALK_101566	(Kushiro *et al.*, 2004; Okamoto *et al.*, 2006)
pyr1-1pyl1-1pyl2-1pyl4-1pyl5 pyl8-1	*pyrpyl112458*	Q169stop, SALK_054640, GT_2864, SAIL_517_C08, SM3_3493, SAIL_1269_A02	(Gonzalez-Guzman *et al.*, 2012)
er-5	*er-5*	GK-182D08	isolated in this study
er-6	*er-6*	GK-364C05	isolated in this study
erl1	*erl1*	GK-109G04	isolated in this study
erl2-2	*erl2-2*	SALK_015275C	(Nanda *et al.*, 2019)
erl2-3	*erl2-3*	GK-486E03	(Yang *et al.*, 2022*b*)
sdd1	*sdd1*	GK-627D04	(Hara *et al.*, 2007)
sdd1-4	*sdd1-4*	GK-693D08	isolated in this study
tmm stkr1	*tmm*	SALK_115723C	isolated in this study, also homozygous for T-DNA insert in *STKR1* (Nietzsche *et al.*, 2018)

Table 2. ANOVA results for Figure 1D.

	df	SS	MS	F-value	P-value
genotype	3	1.54	0.514	15.53	**< 0.0001**
leaf	3	0.15	0.051	1.55	0.20
genotype x leaf	9	0.28	0.031	0.96	0.48
residuals	465	15.39	0.033		
total	480	17.37			

Figure legends

Figure 1. Stomatal density depends on the leaf number in some Arabidopsis genotypes. (A) A scheme depicting the Arabidopsis plant with leaves numbered according to the developmental order. Arrowheads indicate leaves sampled for analyses. Stomatal density in the adaxial (B) and abaxial (C) epidermis, and stomatal ratio (D) in leaves 6, 8, 10, and 12 of selected mutants, genotype (gt). The boxes represent the 25^{th} and 75^{th} percentiles, with the median indicated with the horizontal line; the whiskers show the range of values. Solid dots represent individual plants. Significant differences were determined by two-way ANOVA with Tukey post hoc test. Different letters above boxplots represent significant differences at $p < 0.05$, n = 28-32 plants for each genotype.

Figure 2. Lack of major regulators of stomatal development differently affects adaxial and abaxial stomatal patterns. (A) Schematic overview of regulators of stomatal development included in our study. Meristemoid mother cell (MMC); guard mother cell (GMC); guard cell (GC). Stomatal ratio (B), adaxial (C) and abaxial (D) stomatal density and stomatal index in the adaxial (E) and abaxial (F) epidermis. The boxes represent the 25^{th} and 75^{th} percentiles, with the median indicated with the horizontal line; the whiskers show the range of values. Solid dots represent individual plants. Significant differences were determined by one-way Welch ANOVA with Dunnett T3 post hoc test. Different letters above boxplots represent significant differences at $p < 0.05$, n = 9-10 plants for each genotype.

Figure 3. Stomatal precursors are mostly found in abaxial epidermis. Representative images of adaxial and abaxial stomatal impressions of analyzed mutants that harbor stomatal precursors in fully-developed leaves. Yellow arrowheads indicate stomatal precursors. The scale bar represents 50 μm.

Figure 4. Stomatal density is negatively related with plant growth. Total stomatal density (A) and projected rosette area (B) of stomatal developmental and ABA biosynthesis or catabolism mutants. (C) Relationship between total stomatal density and rosette area. The boxes represent the 25^{th} and 75^{th} percentiles, with the median indicated with the horizontal line; the whiskers show the range of values. Solid dots represent individual plants. Significant differences were determined by one-way Welch ANOVA with Dunnett T3 post hoc test (A, B) or linear regression (C). Different letters above boxplots represent significant differences at $p < 0.05$, n = 9-10 plants for each genotype.

Figure 1. Stomatal density depends on the leaf number in some Arabidopsis geno-types. (A) A scheme depicting the Arabidopsis plant with leaves numbered according to the developmental order. Arrowheads indicate leaves sampled for analyses. Stomatal density in the adaxial (B) and abaxial (C) epidermis, and stomatal ratio (D) in leaves 6, 8, 10, and 12 of selected mutants, genotype (gt). The boxes represent the 25th and 75th percentiles, with the median indicated with the horizontal line; the whiskers show the range of values. Solid dots represent individual plants. Significant differences were determined by two-way ANOVA with Tukey post hoc test. Different letters above boxplots represent significant differences at p < 0.05, n = 28-32 plants for each genotype.

Figure 2. Lack of major regulators of stomatal development differently affects adaxial and abaxial stomatal patterns. (A) Schematic overview of regulators of stomatal development included in our study. Meristemoid mother cell (MMC); guard mother cell (GMC); guard cell (GC). Stomatal ratio (B), adaxial (C) and abaxial (D) stomatal density and stomatal index in the adaxial (E) and abaxial (F) epidermis. The boxes represent the 25th and 75th percentiles, with the median indicated with the horizontal line; the whiskers show the range of values. Solid dots represent individual plants. Significant differences were determined by one-way Welch ANOVA with Dunnett T3 post hoc test. Different letters above boxplots represent significant differences at p < 0.05, n = 9-10 plants for each genotype.

Figure 3. Stomatal precursors are mostly found in abaxial epidermis. Representative images of adaxial and abaxial stomatal impressions of analyzed mutants that harbor stomatal precursors in fully-developed leaves. Yellow arrowheads indicate stomatal precursors. The scale bar represents 50 µm.

Figure 4. Stomatal density is negatively related with plant growth. Total stomatal density (A) and projected rosette area (B) of stomatal developmental and ABA biosynthesis or catabolism mutants. (C) Relationship between total stomatal density and rosette area. The boxes represent the 25th and 75th percentiles, with the median indicated with the horizontal line; the whiskers show the range of values. Solid dots represent individual plants. Significant differences were determined by one-way Welch ANOVA with Dunnett T3 post hoc test (A, B) or linear regression (C). Different letters above boxplots represent significant differences at $p < 0.05$, n = 9-10 plants for each genotype.