MRI mapping of hemodynamics in the human spinal cord

Kimberly J. Hemmerling[12], Mark A. Hoggarth[13], Milap S. Sandhu[45], Todd B. Parrish[26], Molly G. Bright[12]

Author Affiliations:
[1] Department of Physical Therapy and Human Movement Sciences, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA 60611
[2] Department of Biomedical Engineering, McCormick School of Engineering, Northwestern University, Evanston, IL, USA 60208
[3] Department of Physical Therapy, North Central College, Naperville, IL 60540
[4] Shirley Ryan Ability Lab, Chicago, IL, USA 60611
[5] Department of Physical Medicine and Rehabilitation, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA 60611
[6] Department of Radiology, Feinberg School of Medicine, Northwestern University, Chicago, IL, USA 60611

Abstract

Impaired spinal cord vascular function contributes to numerous neurological pathologies, making it important to be able to noninvasively characterize these changes. Here, we propose a functional magnetic resonance imaging (fMRI)-based method to map spinal cord vascular reactivity (SCVR). We used a hypercapnic breath-holding task to evoke a systemic vasodilatory response during concurrent blood oxygenation level-dependent fMRI. SCVR amplitude and hemodynamic delay were mapped at the group level as proof-of-concept of the approach, and in two highly-sampled participants to probe feasibility/stability of individual SCVR mapping. Across the group and individuals, a strong ventral SCVR amplitude was initially observed without accounting for local regional variation in the timing of the vasodilatory response. Shifted breathing traces were used to account for temporal differences in the vasodilatory response across the cord, producing maps of SCVR delay. These maps demonstrate distinct gray matter regions concordant with territories of arterial supply. The SCVR fMRI methods described here enable robust mapping of spatiotemporal hemodynamic properties of the human spinal cord. This noninvasive approach has the potential to provide early insight into pathology-driven vascular changes in the cord, which may precede and predict future irreversible tissue damage and guide the treatment of several neurological pathologies involving the spine.

Acknowledgements: This work was supported by the Center for Translational Imaging at Northwestern University. The authors would like to thank Robert L. Barry for contributions to the development of our data preprocessing pipeline and Sameer Faruquee for contributions to data preprocessing. Research supported by the Craig H. Neilsen Foundation (595499). KJH was supported by an NIH NIBIB-funded training program (T32EB025766) and NINDS-funded predoctoral fellowship (F31NS134222). This research was supported in part through the computational resources and staff contributions provided for the Quest high performance computing facility at Northwestern University which is jointly supported by the Office of the Provost, the Office for Research, and Northwestern University Information Technology.

Corresponding Authors:
Kimberly J. Hemmerling: khemmerling@northwestern.edu
Molly G. Bright: molly.bright@northwestern.edu

Introduction

Impaired vascular function is a major contributor to multiple neurological pathologies involving the spinal cord, including traumatic spinal cord injury [1] and degenerative cervical myelopathy [2]. It is therefore of great importance, for both basic research and clinical translation, to characterize these changes in blood supply to the spinal cord tissue. However, existing methods to noninvasively characterize vascular function in the human brain have not yet been successfully translated to the cord. In general, physiological and functional magnetic resonance imaging (fMRI) of the spinal cord has lagged the development of such methods in the brain. This has largely been due to the challenges typically associated with these methods, including the small size of the spinal cord, nearby sources of physiological noise, magnetic field inhomogeneities, low signal to noise ratio (SNR), and lower widespread availability of specialized radiofrequency coils [3,4]. More recently, technological and methodological developments have enabled more robust fMRI studies of the spinal cord, including mapping of sensory, motor, and resting-state neural activation patterns [5].

Despite these advances, MRI techniques for characterizing vascular properties of the spinal cord have remained somewhat limited. Dynamic susceptibility contrast (DSC) MRI, following injection of a gadolinium-based contrast agent, presents the gold standard for quantifying tissue perfusion and blood volume in the brain. Existing studies of vascular changes in the spinal cord of people with degenerative cervical myelopathy use this technique, showing sensitivity to the ischemic effects of cord compression [6,7]. However, due to the low SNR of DSC data in the cord, these works report a single non-quantitative value for the entire cord region, rather than quantifying and mapping absolute vascular metrics. Repeated scans may boost data quality, but they are not readily feasible in DSC due to the long wash-out time of the contrast agent and the growing safety concerns over repeated exposure to gadolinium-based contrast agents [8]. Dynamic contrast enhanced (DCE) MRI measures the contrast agent accumulation [9]; although more commonly used to look at tumors in the spine, preliminary work demonstrates the potential of DCE to evaluate spinal cord perfusion after compression, but the technique is still in early development [10].

A contrast-free alternative, arterial spin labeling (ASL) MRI, shows excellent promise in the spinal cord injured rat [11,12], and in the human spinal cord in early conference proceedings [13,14]. However, despite rapid growth of ASL for perfusion mapping of the brain, the unique challenges of spinal cord ASL have continued to hinder progress (e.g., low SNR, variable vascular anatomy for labelling and delay modeling, B_0 inhomogeneity influencing labeling efficiency) [11]. Recent exciting advances in pre-clinical studies are renewing interest and enthusiasm for spinal cord ASL in humans, but substantial methodological development is still needed before this technique can reliably shed light on human spinal cord pathophysiology [11]. Another current research area is high-field intra-voxel incoherent motion (IVIM) MRI, which has shown promising maps of healthy human spinal cord perfusion, albeit in a single slice [15].

Here, we propose an alternative technique to map vascular function throughout the
human spinal cord using blood oxygenation level-dependent (BOLD) fMRI, a well-
established neuroimaging technique that is sensitive to changes in regional blood flow, and
which has been successfully adapted and optimized for spinal cord neural activation
mapping in recent years [5]. Specifically, we use fMRI to map spinal cord vascular reactivity
(SCVR), the cord's blood flow responsiveness to a systemic vasodilatory stimulus. In the
brain, vascular reactivity and hemodynamic delay measured with BOLD fMRI have been
established as robust metrics of cerebrovascular function [16,17]. However, techniques for
adapting these cerebrovascular reactivity methods to the spinal cord are not yet well
established. Two prior spinal cord fMRI studies have used gas inhalation challenges to
modulate the vascular system; however, these are smaller studies (<10 participants) without
more modern acquisition and modeling innovations, and do not map the distribution of
vascular reactivity and delay [18,19].

In this study, we use advanced spinal cord fMRI methods to measure the local
vasodilatory response to hypercapnia and map SCVR across the cervical spinal cord in
healthy adults. We evoke systemic hypercapnia (elevated CO_2) with simple breath-holding
tasks (see reviews on breath-hold cerebrovascular reactivity [16,17]), precluding the need for
gas delivery systems. By recording end-tidal CO_2 ($P_{ET}CO_2$) levels during scanning, we form
a subject- and scan-specific model of the hypercapnia achieved by our breath-holding
protocol. By determining the voxelwise optimal temporal shift for this model, we produce
more accurate estimates of SCVR amplitude and an additional complementary metric of
hemodynamic delay. This approach is becoming standard in the field, and BOLD fMRI-based
cerebrovascular reactivity studies have increasingly employed a transfer function analysis
(gas inhalation) or voxelwise shift optimization procedure (breathing task) to model local
hemodynamics [20–30].

In this pioneering work, we explore SCVR from two approaches. In the first **(Group
SCVR)**, we aim to establish proof-of-concept of our SCVR approach in the cervical spinal
cord, mapping the group-level BOLD response from a cohort of 27 healthy participants.
Participants completed 2 breath-holding task fMRI runs across 2 sessions. However, to
provide more clinically relevant information, SCVR needs to be characterized in individuals.
Therefore, in the second study **(Individual SCVR)** we focus on 2 highly-sampled participants
to evaluate feasibility of individual SCVR mapping. Each participant completed 18 breath-
holding task fMRI runs across 3 MRI sessions.

In both of these approaches, we mapped SCVR and the associated hemodynamic delay
using a lagged general linear model (GLM) strategy. To map SCVR amplitude, $P_{ET}CO_2$ task
regressors were fit to the fMRI timeseries in a GLM (also incorporating denoising via
nuisance regression). Output parameter estimates (%BOLD per change in $P_{ET}CO_2$ in mmHg)
were combined across the group (Group SCVR) or across individuals' runs (Individual
SCVR). Hemodynamic delay mapping was achieved by temporally shifting the $P_{ET}CO_2$
regressor and fitting each shifted regressor to the fMRI timeseries. Then, the shift with the

best fit across participants (Group SCVR) or runs (Individual SCVR) was used to create a
hemodynamic delay map, representing regional variation in the timing of this vascular
response across the spinal cord. The SCVR amplitude estimate at this optimal shift was
considered the "delay-corrected" SCVR.

We present maps of SCVR amplitude and hemodynamic delay at the group-level and in
highly-sampled individuals, in the C5–C8 segments of the human cervical spinal cord.
Furthermore, by thresholding the hemodynamic delay maps, we produce noninvasive maps
of vascular territories in the human spinal cord, showing agreement with territory maps
extrapolated from invasive and preclinical studies. Our scanning methods are readily
implemented on most 3T MRI scanners, making this approach to characterizing spinal cord
hemodynamics immediately available for basic science and translational research into
human spinal cord neurovascular physiology and pathology.

140
141
142 **Results**
143
Group SCVR amplitude. The breath-hold response was modeled with a $P_{ET}CO_2$ task
regressor in first-level GLMs, allowing SCVR to be mapped in standard semi-quantitative
units of %BOLD/mmHg. Group SCVR amplitude was mapped in PAM50 template space
(Fig. 1). The group-level SCVR response is distributed predominantly within the ventral
spinal hemicord, primarily in or near the gray matter ventral horns, and spread throughout
the longitudinal extent of the cervical spinal cord.

151
152 **Fig. 1. Group-level SCVR amplitude map.** Significant SCVR is shown in units of
153 %BOLD/mmHg ($p < 0.05$, FWE-corrected). Two sagittal, two coronal, and three axial slices
154 are shown (green lines indicate location of slices). Approximate centers of spinal cord
155 segments are indicated. (FWE=family-wise error, S=superior, I=inferior, D=dorsal, V=ventral,
156 L=left, R=right).

Individual SCVR amplitude. The $P_{ET}CO_2$ regressors show high task compliance across
18 fMRI task runs for the two highly-sampled subjects **(Fig. 2A,E)**. First-level GLMs were
calculated, and then the 18 output parameter estimate maps were collated. Individual SCVR
amplitude was mapped in PAM50 template space **(Fig. 2B,F)**. SCVR amplitude for both
highly-sampled subjects has a similar spatial distribution to the group-level maps, with a high
density of significantly responding voxels in the ventral gray matter tissue. However, the
amplitude for Subject 1 is visibly higher than for Subject 2.

Fig. 2. SCVR mapping in two highly-sampled subjects. (A,E) $P_{ET}CO_2$ regressors
(convolved with the canonical HRF and demeaned) for 18 breath-holding task runs and their
average. **(B,F)** Maps of SCVR amplitude in axial, sagittal, and coronal planes (FWE-
corrected, $p < 0.05$). Spinal cord segments and slice locations (green lines) are
indicated. **(C,G)** Split-halves spatial correlation of gray matter SCVR amplitude for varying
number of runs combined (1-9) vs. other half combined (9 runs). **(D,H)** Average SCVR as
runs are sequentially combined in the order they were acquired (red line) and in randomly
shuffled orders (shaded areas; ±1SD and ±2SD).

To investigate the reliability of SCVR amplitude estimates generated from subsets of the
highly-sampled individual datasets, a split halves analysis was performed. Modeled after
Laumann et al. 2015 [31], this was done by randomly and repeatedly splitting gray matter

SCVR estimates for the 18 runs into two halves and comparing combinations of one half (1-9
runs) to the other half combined (9 runs). A spatial Pearson correlation within ventral gray
matter voxels was used to compare the two halves **(Fig. 2C,G)**. In both subjects, comparing
SCVR estimates derived from only 1 run versus from 9 runs combined yields moderate
spatial correlation (median $r < 0.5$). When comparing SCVR estimates from 6 runs combined
versus 9 runs combined, this yields a strong spatial correlation (median $r \approx$ 0.7-0.8). Looking
at average SCVR as runs are incrementally combined, the variability initially rapidly
decreases, and then more gradually **(Fig. 2D,H)**.

Group SCVR delay. The distinct ventral distribution of significant SCVR responses
raises the question of whether variation in the timing of the SCVR response may be
obscuring this effect in more dorsal regions. To investigate and account for temporal delays,
first-level models were recalculated with shifted versions of the $P_{ET}CO_2$ task regressor (±10s,
in increments of 2s) **(Fig. 3A)**. Group-level SCVR was calculated for each shift using the
same method as above. The optimal shifts were identified, and the corresponding delay map
and delay-corrected SCVR amplitude map were generated **(Fig. 3B, Supplementary Fig.**
S1). (Note, this approach assumes that the timing of the voxelwise SCVR response is the
same across all subjects.)

Fig. 3. SCVR hemodynamic delay mapping at the group-level. (A) Example of temporally
shifted $P_{ET}CO_2$ regressors. The unshifted (no delay) $P_{ET}CO_2$ regressor (thick green line) and
each tested shift ±10 s, in 2s increments, are shown. Line colors correspond to the "Delay"
colorbar in panel B. Full traces are truncated at 100 TRs for ease of visualization. **(B)** SCVR
hemodynamic delay map (ranging ±10 s, in increments of 2 s). The same slices as Fig. 1 are
shown. **(C)** Gray matter regional segmentation based on delay threshold.

As anticipated, the delay-corrected SCVR amplitude map shows a more diffuse
distribution of significantly responding voxels. Delay-corrected SCVR (mean ± SD) is 0.04 ±
0.02 %BOLD/mmHg in the cord and 0.05 ± 0.02 %BOLD/mmHg in the ventral gray matter.
Across all participants, the distribution of Group SCVR hemodynamic delay shows an earlier
response in the ventral cord, and later in the dorsal cord. Moreover, these earlier and later
responding regions appear to resemble the shape of the ventral and dorsal gray matter

horns, respectively. The mean (± SD) delay between the dorsal and ventral horns is
approximately 6.2 ± 4.9 seconds (about 3 times our sampling TR).

Using a histogram of gray matter SCVR delay, a delay threshold was manually defined;
each voxel was labeled as Region 1 (sub-threshold, earlier response) or Region 2 (supra-
threshold, later response), which comprises the delay-based gray matter segmentation **(Fig.**
3C). The approximate boundary between the ventral and dorsal horns can be clearly
observed.

Individual SCVR delay. Hemodynamic delay was mapped with the same methods as
for Group SCVR. The SCVR delay maps for the two highly-sampled individuals show a
similar ventral-dorsal spatial distribution as for Group SCVR **(Fig. 4A,E)**. The delay between
dorsal and ventral gray matter is approximately 6.3 ± 5.2 seconds and 7.6 ± 8.0 seconds for
Subject 1 and Subject 2, respectively. A delay-corrected SCVR amplitude map was created
using the parameter estimates associated with the optimal shift for each voxel **(Fig. 4B,F)**,
and shows a more diffuse spread of significantly responding voxels, (compared to Fig. 2B,F),
although there are still dorsal regions without a significant response. The mean delay-
corrected SCVR in the spinal cord for Subject 1 and Subject 2 is 0.08 ± 0.06 %BOLD/mmHg
and 0.07 ± 0.06 %BOLD/mmHg, respectively. And in the ventral gray matter for Subject 1
and Subject 2 is 0.10 ± 0.04 %BOLD/mmHg and 0.07 ± 0.04 %BOLD/mmHg, respectively.

Fig. 4. SCVR delay mapping in two highly-sampled subjects. (A,E) SCVR delay map
(ranging ±10 s, in increments of 2 s). Spinal cord segments and slice locations (solid green
lines) are indicated. The three solid and dotted green lines in the sagittal view indicate axial
slices in the lower panels. **(B,F)** Delay-corrected SCVR amplitude (FWE-corrected, *p* < 0.05,
Šidák corrected). The same slices as Fig. 2 are shown. **(C,G)** SCVR delay map within gray
matter voxels. **(D,H)** Gray matter regional segmentation based on delay threshold.

The individual subject SCVR hemodynamic delay maps, restricted to show gray matter
voxels, clearly recapitulate the earlier and later responses in the ventral and dorsal horns,
respectively **(Fig. 4C,G)**. Applying the thresholding approach described above, the ventral
and dorsal boundary is defined, and agrees with the group-level results **(Fig. 4D,H)**.

Discussion

In this work, we used a breath-holding task to evoke systemic vasodilation and a corresponding BOLD fMRI response in the spinal cord, and then modeled SCVR amplitude and hemodynamic delay. Applying these methods in a cross-sectional cohort and in two highly-sampled individuals, we produced robust spatiotemporal maps of vascular function in the human cervical spinal cord. Our results reveal hemodynamic differences in ventral versus dorsal gray matter regions, consistent across individuals, suggesting that SCVR delay could delineate vascular territories in the cord.

SCVR modeled with $P_{ET}CO_2$ provides semi-quantitative information about the regional spinal cord hemodynamic response to a vasoactive stimulus. Cerebrovascular reactivity has already proven to deliver clinically relevant information about cerebrovascular health [16,17], indicating the potential utility of similar measures for the spinal cord. In the healthy cohort examined in this study, the distribution of significant SCVR (without delay correction) is restricted to more ventral tissue, although extending throughout the longitudinal extent of our imaging field of view. Sensitivity of the SCVR technique in this region may be particularly relevant for early detection of vascular changes in patients with degenerative cervical myelopathy, as age-related cord compression and subsequent neurodegeneration is most prevalent in the ventral regions [32].

The significant ventral SCVR response is predominantly, but not strictly, localized to gray matter voxels. This agrees with observations of vascular reactivity in the brain, where the cerebrovascular reactivity amplitude is greater in gray matter versus white matter voxels due to it being more highly vascularized and having higher SNR [17].

Early reports of BOLD signal changes in the spinal cord achieved with a controlled hypercapnic gas inhalation challenge were roughly 0.17 %BOLD/mmHg (based on a reported 0.6 ± 0.3 mean percent signal change with a 3.5 ± 0.8 mmHg mean $P_{ET}CO_2$ change [18]), as compared to 0.04 ± 0.02 %BOLD/mmHg for the group delay-corrected SCVR across the whole spinal cord region-of-interest in this study. This difference may be due to smaller sample sizes inflating parameter estimates [33], or because of the different methods of averaging – our measure is obtained from the group level parameter estimates. Additionally, the response to breath-holding is transient, while gas inhalation evokes a more prolonged response, making delay correction particularly important for breath-holding SCVR. Delay correction performed at the group-level, however, is sub-optimal due to anticipated variability across participants. This is evidenced by the higher mean SCVR estimates for the two highly-sampled subjects. Furthermore, breath-hold induced hypercapnia is associated with a confounding effect of hypoxia, while controlled hypercapnia is not [34,35]. Based on Schulman and Uludağ's simulation of cerebrovascular reactivity with and without hypoxia (to mimic breath-holding and controlled hypercapnia, respectively) breath-holding is expected to underestimate cerebrovascular reactivity measurements, which agrees with reports in the literature [34]. The mean group delay-corrected SCVR in ventral gray matter is 0.05 ± 0.02

%BOLD/mmHg, which is about four to five times lower than similar mean gray matter
cerebrovascular reactivity measures [36,37] and is to be expected because of higher SNR in the
brain.

The SCVR maps derived from the highly-sampled individuals reveal higher SCVR
amplitude for Subject 1 compared to Subject 2. There is also a conspicuous absence of
significant SCVR estimates in the superior spinal cord for Subject 2. The underlying reason
for these differences in SCVR amplitude is unclear and may represent differences in
anatomy, physiology, or regional fMRI signal quality.

Individual SCVR amplitude maps represent the 126 breath-holds performed across the
18 fMRI runs. Considering our long-term goal to use SCVR to provide clinical insight into
neurological pathology development and progression, it is not reasonable to expect patients
to undergo over 100 minutes of breath-hold fMRI. To assess the feasibility of clinical
translation using our current SCVR acquisition strategy, we implemented a split halves
analysis to probe the number of fMRI runs necessary to achieve reliable SCVR estimates. As
shown in **Fig. 2**, the voxelwise spatial correlation in ventral gray matter begins to plateau at
approximately 5-6 runs, indicating data from one scan session (34-41 min) may be sufficient
to map SCVR. The high initial variability in average ventral gray matter SCVR is also
substantially reduced at 5-6 runs combined. Although this scan duration may limit the use of
SCVR methodology to research studies, we anticipate that future methodological
improvements in acquisition, denoising, and machine learning will further decrease this scan
time requirement and make SCVR mapping a viable clinical scan technique. Furthermore,
there is relatively low variability in voxelwise SCVR within gray matter in these healthy
individuals, and spatial correlations may be much higher in subjects with greater
heterogeneity in SCVR values due to pathology.

Since the hypercapnic breath-hold challenge evokes a systemic vasodilatory response,
we expected a more widespread fMRI response throughout both ventral and dorsal spinal
cord tissue. The absence of significant dorsal responses in **Figs. 1** and **2** allude to
spatiotemporal variability in the relationship between our $P_{ET}CO_2$ model and the associated
BOLD response across the spinal cord. Differences in timing could arise because of arterial
transit time differences and local variability in the vasodilatory blood flow response and
associated BOLD contrast changes. Maps of SCVR hemodynamic delay reveal local
regional variation in the timing of the blood flow response that are consistent across our
cohort, with earlier and later responses in the ventral and dorsal gray matter, respectively.
Although BOLD fMRI is primarily sensitive to venous effects, the response to CO_2 is driven
by arteriolar vasodilation. So, local timing differences may (at least in part) represent
territories of arterial supply to the spinal cord.

Supplying approximately the ventral two-thirds of the cord is the central system, derived
from the anterior spinal artery (ASA) [1]. From the ASA, central sulcal arteries penetrate into
the anterior median fissure [1,38]. Blood flows centrifugally, supplying the ventral gray matter,

intermediate zone, ventral aspect of the dorsal gray matter, and other white matter regions
[1,39]. This aligns well with the spatial distribution of the earlier ventral gray matter response.

The approximate remaining one third of the cord is supplied by the peripheral system,
derived from the paired posterior spinal arteries (PSAs) and pial arterial plexus [1]. The ASA
and PSAs contribute to an anastomosed network on the periphery of the spinal cord, the pial
arterial plexus, from which vessels penetrate the spinal cord perpendicular to its surface [1,39].
Blood flows centripetally, supplying the dorsal aspect of the dorsal gray matter and other
white matter regions [1,39]. This aligns well with the spatial distribution of the later SCVR
responses. Additionally, by applying a simple threshold to the SCVR hemodynamic delay
maps, we can cleanly delineate these two gray matter vascular territories: (1) areas with an
earlier response of likely ASA supply (ventral horns and intermediate zone) and (2) areas
with a later response of likely PSAs supply (dorsal horns).

The observed difference in delay between the ventral and dorsal horns was on the order
of several seconds for both the group and individual SCVR maps. In interpreting the spatial
distribution of delay across the spinal cord, it is important to remember that the SCVR
response does not only represent an arterial transit time difference but also represents
variation in the local vasodilatory response as well as variation in the BOLD signal evolution.
While differences in arterial transit time measured across the brain are about 1-2 seconds
[40,41], maps of cerebrovascular reactivity delay using BOLD fMRI in healthy participants have
shown larger regional differences of several seconds across brain regions [21,24,27,28]. It is also
relevant to consider how these estimates may be influenced by the spatial resolution,
particularly in the dorsal horn. Specifically, there is the potential for partial volume effects
with the nearby white matter, which may be exacerbated during co-registration and the
transformation of images from the subject space to the template space. The capillary bed in
white matter is much less dense as compared to the gray matter [1,42,43], similar to what is
observed in the brain [17]. Furthermore, a portion of the dorsal horn is not well supplied by the
capillary bed [1]. These factors may adversely affect SCVR amplitude and delay estimation [17].
Higher magnetic field strengths are an appealing choice for imaging white matter
hemodynamic signals [44], and future ultra-high field MRI work may offer improved
characterization of dorsal horn or white matter SCVR.

These local timing differences in the BOLD response to hypercapnia are driven by
multiple sources such as vascular transit delays, variation in the local vasodilatory response,
and variation in the BOLD signal response [20,21,24,27]. Incorporating a voxelwise temporal shift
has demonstrated differences in delay across the healthy or injured brain [21,22,24] and
accounting for these delays can provide for more accurate local estimation of
cerebrovascular reactivity, which is particularly relevant for measurement in pathology.
Similarly, when accounting for SCVR hemodynamic delay by "correcting" the timing of the
$P_{ET}CO_2$ regressor, we have a more optimal model of vascularly active voxels across the
entire cord. After delay correction, significantly responding voxels are more diffusely spread
over both ventral and dorsal regions, as compared to the original SCVR amplitude map. This

agrees with work in the brain showing increased cerebrovascular reactivity estimates after
delay correction [20]. This effect is more pronounced in the highly-sampled individuals than for
the group-level results; this is likely due to inter-individual variation in hemodynamic delay,
which makes it sub-optimal to apply the same delay correction to an entire group. However,
the limited data available for each subject in the cross-sectional analysis are not sufficient to
robustly estimate and correct for their unique voxelwise hemodynamic delay patterns using
our current methods.

We demonstrate that our fMRI acquisition and modeling approach can produce
consistent and meaningful insight into hemodynamics of the human spinal cord. However,
there are numerous challenges in robustly mapping SCVR amplitude and delay that should
be considered and addressed in future research.

A breathing task, such as breath-holding, is relatively simple to implement and should
cause large, detectable variations in the fMRI signal, but is also likely to cause respiration-
induced B_0 artifacts, time-locked to the task. Fourier-based respiratory RETROICOR [45]
nuisance regressors were included to aid fMRI denoising. There may be concern that some
of the signal of interest would be fitted by respiratory RETROICOR regressors, instead of the
task. Ultimately, these regressors were included because physiological noise modeling
techniques with RETROICOR have been shown to be beneficial with noisy spinal cord fMRI
data [46]. Additionally, these regressors were included in the first-level model, rather than as a
separate denoising regression step. The inclusion of RETROICOR denoising in our study
may therefore cause SCVR amplitude to be slightly underestimated, and the impact on delay
estimation is not yet clear.

Another consideration for breathing tasks is the potential for motion confounds that are
time-locked to respiration [28,47]. Mitigation of these effects was done through a spinal cord-
specific motion correction procedure [48] and then by using those output motion traces in the
GLM (see Methods). However, we acknowledge that effects of motion may not be entirely
accounted for by these procedures. In future work, hypercapnia could be achieved using gas
inhalation challenges, which may result in a more controlled hypercapnic stimulus that is not
time-locked to ventilation, chest position changes, and subsequent B_0 effects. Gas inhalation
challenges could also increase task compliance, because participants do not need to actively
comply with the voluntary breathing task. Indeed, the early spinal cord fMRI work involving a
hypercapnic stimulus used different gas inhalation techniques [18,19]. However, the requisite
facemask may be less comfortable for participants and less compatible with some coils (e.g.,
the 64-channel head/neck coil used in this study). Although literature from cerebrovascular
reactivity mapping in the brain suggests that these two methods generate similar results [49],
future work should be done to compare breath-holding task-based SCVR measurements
with those derived from gas inhalation challenges.

The ground truth of our SCVR measurements across spinal cord vascular territories is
unknown; it will be important in future work to refine the range and step size of the shifted
$P_{ET}CO_2$ regressors. The ±10s $P_{ET}CO_2$ range was informed by previous work from our lab

mapping cerebrovascular reactivity using a shift range of ±9s [27,50]. It is pertinent to note that
histograms of gray matter delay showed a high density at the upper boundary of the range
(+10s) as well as the presence of voxels with -10s delays directly adjacent to voxels with
+10s delays. These voxels may not reflect physiologically plausible SCVR delays.
Additionally, in brain literature, with finer (~0.3s) increments, shifts at the upper and lower
boundary of the range are sometimes not included because they are not considered
optimized [20,27]. Considering the coarse 2s shift increment used in this study, boundary delays
were not removed. Refinement of the range and step size will be an important target for
future work, and evidence from brain literature suggests this may be particularly important in
individuals with pathologically delayed blood supply to the cord [27]. Additionally, Bayesian
modeling of SCVR amplitude and delay, using our initial work to inform spatial priors, could
also improve our ability to robustly map individual subject hemodynamics [25].

Finally, the $P_{ET}CO_2$ regressors used throughout this study were convolved with the two-
gamma variate canonical HRF [51]. While this HRF was designed to convert a neural task
design into a physiological response, the canonical HRF convolved with $P_{ET}CO_2$ is often
used throughout the cerebrovascular reactivity literature [16,27,36,52]. In a direct comparison of a
set of models, $P_{ET}CO_2$ convolved with the canonical HRF plus its temporal derivative was
shown to be the best model of the BOLD response to breath-holding [53]. While the present
study did not incorporate a temporal derivative, temporal offsets in the local BOLD response
were instead accounted for by the analysis of hemodynamic delay. Aside from the delay,
there may still be differences in the shape of the response that should be accounted for. In
fact, a recent preprint comparing models with voxelwise lag optimization suggests either non-
convolved $P_{ET}CO_2$ or single gamma HRF-convolved $P_{ET}CO_2$ as the best model. Given the
temporal resolution and low SNR of the spinal cord fMRI data, it is not possible to estimate
the response shape in this initial study. Future work should explore an appropriate response
function specifically for measures of vascular reactivity in the spinal cord.

427
428
429 **Conclusion**
430
SCVR amplitude and delay mapping has the potential to provide clinically relevant
information about spinal cord vascular health. Robust maps of SCVR amplitude and delay
were presented at the group-level and in two highly-sampled individuals. Hemodynamic
delay maps represent the temporal difference in the BOLD response across the cord and
resemble expected territories of arterial supply. Future work to refine these promising
noninvasive methods will facilitate our long-term goal to establish SCVR as a clinically
relevant metric of spinal cord vascular health.

Methods

Participants. This study was approved by the Northwestern University Institutional Review Board, and informed consent was obtained for all participants; all methods were performed in accordance with relevant guidelines and regulations. Two datasets were collected for this study and will be referred to as Group SCVR and Individual SCVR. Group SCVR dataset: MRI data were collected from 30 healthy participants (26±5y, 19M). Data were excluded for 3 participants due to excessive image artifacts (N=2) and an incidental finding (N=1). All subsequent Group SCVR analyses represent data from the remaining 27 participants (26±5y, 18F). Individual SCVR dataset: MRI data were collected from 3 healthy participants (24±3y, 2F). Data were excluded for one participant due to radiofrequency image artifacts. Subsequent Individual SCVR analyses represent data from the remaining 2 participants (26±2y, 2F).

Data Collection. Spinal MRI data were acquired on a 3T Siemens Prisma MRI system (Siemens Healthcare, Erlangen, Germany) with a 64-channel head/neck coil. A SatPad™ cervical collar (SatPad Clinical Imaging Solutions, West Chester, PA, USA) was positioned around the posterior of the neck to increase the homogeneity of the magnetic field around the imaging region. Group SCVR: MRI was acquired before and after the administration of a 30-minute acute intermittent hypoxia (AIH) protocol, as part of a larger research study. Individual SCVR: Anatomical and functional scans were acquired in 3 sessions across 2 days during the same time of day for each participant. Sessions 1 and 2 took place on the first scan day and were in the morning and afternoon, respectively; session 3 took place in the morning on the second scan day. The two scan days were one or two days apart.

Acute Intermittent Hypoxia (AIH). AIH has been effective at improving motor function in spinal cord injured cohorts [54,55]. The data collected for the Group SCVR study was a part of a broader study to investigate with MRI the mechanisms of neural plasticity associated with AIH. This AIH protocol was administered after the first MRI session with a HYP 123 oxygen generator (Hypoxico, Inc., New York, NY, USA) in 15 2-minute cycles. Each cycle consisted of brief exposures to a hypoxic gas mixture (9% FiO_2, 30-60 seconds), alternating with normal room air (21% FiO_2, 60-90 seconds), targeting 85% SpO_2 during each bout. The second MRI session occurred 45-60 minutes after AIH. Note, AIH had a non-significant impact on SCVR (via a two-tailed paired t-test using threshold-free cluster enhancement (TFCE) and 5000 permutations (*randomise*) [56–58]. Pre- and post-AIH MRI scans were also considered independently to further probe any possible impact of AIH. For details of this analysis see Supplementary Information Section 1. The SCVR maps and average values were consistent (**Supplementary Fig. S2, S3, and S4**). Thus, there was no evidence that AIH modulates SCVR in healthy individuals and the impact of AIH was considered negligible in further analyses.

Imaging Protocol. A high resolution anatomical T2-weighted scan, covering the brainstem to upper thoracic spine, was acquired with the following parameters: repetition time

(TR)/echo time (TE) = 1500/135 ms, sagittal slice thickness = 0.8 mm, in-plane resolution

0.39 mm^2, 64 slices, flip-angle (FA) = 140°, field-of-view (FOV) = 640 mm^2. Spinal cord fMRI

scans were collected using a gradient-echo echo-planar imaging (EPI) sequence and

ZOOMit selective excitation (TR/TE = 2000/30 ms, axial in-plane resolution = 1 mm^2, axial

slice thickness = 3 mm, 25 ascending interleaved slices, FA = 90°, FOV = 128 x 44 mm^2).

The ZOOMit acquisition reduces the field-of-view around the spinal cord. The functional

acquisition volume was positioned perpendicular to the spinal cord; the bottom of the volume

was positioned at the bottom of the C7 vertebral level. Cervical coverage was approximately

from the C4 to C7 vertebral level, but the exact coverage varied due to participant height and

spinal anatomy. For each scan session, the task paradigm was displayed to participants via

a mirror on the head coil, reflecting a monitor placed behind the bore of the magnet. Group

SCVR: A spinal cord anatomical and breath-holding task functional scan were acquired

during each of the 2 MRI sessions. Additional spinal cord and brain scans were also

acquired during the sessions and were not analyzed in this study. Individual SCVR: A spinal

cord anatomical and 5-7 breath-holding task functional scans were acquired during each of

the 3 MRI sessions. (It was intended that 6 runs would be collected during each session but

due to time constraints during one session, five runs were collected during session 1 and 7

runs were collected during session 2 for Subject 1.) The total number of task runs was 18 for

each participant.

Breath-holding Task Paradigm. A hypercapnic breath-holding task was used to modulate

participants CO_2. The fMRI acquisition was 6 minutes and 50 seconds (205 volumes). The

task began with a 20 second rest, followed by 7 breath-hold cycles, and ended with a 30

second rest period. Each breath-hold cycle was as follows: 24 seconds paced breathing (3

seconds in, 3 seconds out), 18 second end-expiration breath-hold, 2 second exhalation, 6

second recovery.

Physiological Monitoring. Physiological data were collected throughout the fMRI scans,

including exhaled CO_2 and O_2 through a nasal cannula and breathing via respiratory belt. A

pulse transducer was placed either on the dorsalis pedis artery of the foot (for the Group

SCVR study because participants were holding hand-grips for another task) or on the right

index finger (for the Individual SCVR study). These signals were fed through a Gas Analyzer

(CO_2, O_2 only) and PowerLab and recorded with LabChart (ADInstruments, Dunedin, New

Zealand). The scanner trigger was also recorded through the same system to facilitate

alignment of all recordings with fMRI timeseries data.

fMRI Preprocessing Pipeline. Images were converted from DICOM to NIFTI format

(dcm2niix_afni [59]). Motion Correction: 2D slicewise motion correction was applied with the

Neptune Toolbox [48] (version 1.211227) and three steps of the toolbox were used to complete

motion correction, including the manual definition of a "not cord" mask around the spinal cord

and cerebrospinal fluid (CSF) region (#5), computation of a Gaussian weighting mask of the

spinal cord, derived from the "not cord" mask (#7), and the application of motion correction

(#8). The motion correction algorithm used the Gaussian mask as a weight for each voxel, a

temporal median image as the target image for the correction algorithm, and used AFNI
(version 22.0.05) to compute and apply the motion parameters to the data (*3dWarpDrive*,
3dAllineate) [60,61]. Temporal median filtering was de-selected. Motion corrected functional
data and slicewise X and Y motion traces were output. <u>Registration:</u> Co-registration between
native, anatomical, and PAM50 template space [62] was performed with the Spinal Cord
Toolbox (version 5.3.0) [63]. (Note, the Spinal Cord Toolbox version 6.1.0 included an update
to the spinal cord segments and this new version was used to add correct labels to figures.)
A binary spinal cord mask in the functional image space was manually drawn (excluding the
top/bottom slices) and is used in several steps. The anatomical T2-weighted image of each
subject was segmented (*sct_deepseg_sc*) [64] and then registered to PAM50 template space
(*sct_register_to_template*). The binary spinal cord mask in functional image space and
anatomical-template warping field were used to inform registration of the motion corrected
functional data to the PAM50 template space (*sct_register_multimodal*). These warping fields
were calculated but not applied until after subject-level modeling. <u>Smoothing:</u> Functional data
were smoothed using a "restricted smoothing" technique [65], within a mask of the spinal cord.
A 2 x 2 x 6 mm^3 full-width half-maximum (FWHM) Gaussian smoothing kernel was used
(*3dBlurInMask*). <u>Trimming fMRI Volumes:</u> The first and last 5 volumes were trimmed from the
functional data to allow for shifts of the task regressor for hemodynamic delay mapping
(discussed later).

Task and Nuisance Regressors. Physiological data (CO_2, pulse, and respiratory belt) were
preprocessed in a bespoke MATLAB (MathWorks, Natick, MA, R2019b) script. The results of
a peak-finding algorithm were manually verified. <u>Breath-holding Task:</u> From the
preprocessed CO_2 data, an end-tidal CO_2 ($P_{ET}CO_2$) task regressor was calculated and
convolved with the two-gamma variate canonical hemodynamic response function (HRF) [51].
<u>Respiratory and Cardiac:</u> From the preprocessed respiration belt and pulse transducer data,
8 respiratory and 8 cardiac slicewise RETROICOR [45,46] regressors were calculated. Only
cardiac regressors were calculated for one subject in the Group SCVR study without belt
data. <u>CSF:</u> An initial CSF mask was defined; high variance voxels (top 20% in each slice)
were retained in the mask and the average timeseries within this mask was used to create
slicewise CSF regressors [46,66].

fMRI Analysis and Statistics. <u>First-level Analysis:</u> FSL [67] (version 6.0.3) was used to
calculate the first-level fMRI models (FEAT [68]). The first-level models contained the $P_{ET}CO_2$
task regressor and 19 nuisance regressors (2 motion, 16 RETROICOR, 1 CSF). All input
regressors were demeaned. In FEAT, a lenient high-pass filter cutoff of 100 seconds to
chosen to remove the low frequency noise and FMRIB's Improved Linear Model (FILM)
prewhitening was used. One statistical contrast was defined: $P_{ET}CO_2 > 0$. The output "contrast
of parameter estimate" (COPE) maps were normalized by the voxelwise mean to convert to
standard units (i.e., %BOLD/mmHg). Output files were warped from subject functional space
to the PAM50 template space. <u>Higher-level Analysis:</u> The COPE maps were averaged
across the two sessions within each subject (Group SCVR). A group spinal cord mask was

calculated as the consensus region of the co-registered first-level results, and higher-level
analyses were performed within this mask, only. This mask was calculated separately for the
Group SCVR and Individual SCVR studies. A non-parametric one-sample t-test using TFCE
and 5000 permutations was used to calculate the higher-level activation maps with family-
wise error (FWE) rate correction (*randomise*) [56–58]. The average parameter estimates are the
SCVR amplitude.

Reliability of Individual SCVR Estimates. To probe the reliability of SCVR estimates for
different amounts of data, a split-halves analysis was used [31]. For each subject, the 18
COPE maps were randomly split into 2 halves, comparing subsets of one half (1-9 runs) to
the other 9 runs combined via spatial Pearson correlation in the ventral gray matter. This was
repeated 1000 times to build a distribution. Average SCVR in the ventral gray matter was
also considered by incrementally combining runs from 1 run to all 18 runs. This was done in
the originally acquired order of fMRI runs and then for 100 randomly shuffled orders.

SCVR Hemodynamic Delay. 10 temporally shifted versions of the $P_{ET}CO_2$ regressor were
created to allow for timing differences of ±10 s in the increment of the TR (2 s). First-level
models were repeated identically to the original except for the temporally shifted $P_{ET}CO_2$
regressor. The higher-level analyses in PAM50 space were also repeated for each shift. For
each voxel, the highest t-statistic across the delay range was designated as the optimal shift
(the "delay" for that voxel) (*getBestFits)*. Delay-corrected SCVR Amplitude: These optimal
shifts were then used to create a delay-corrected SCVR map. For each run, delay-corrected
COPE maps were generated using the parameter estimates associated with the optimal shift
(*delayCorrectedSCVR*). This resulted in delay-corrected COPE maps for each run, which
were then processed with the same higher-level analyses used previously. The average
parameter estimates are the delay-corrected SCVR amplitude. Šidák correction was used to
account for the multiple tests in the delay-correction procedure [69,70]. Delay-based
Segmentation: A histogram of gray matter delays was manually thresholded. Voxels with
delays on either side of the threshold were designated as *Region 1* (Earlier) or *Region 2*
(Later).

References

1.   Martirosyan, N. L. *et al.* Blood supply and vascular reactivity of the spinal cord under
normal and pathological conditions: A review. *J Neurosurg Spine* **15**, (2011).

2.   Nouri, A., Tetreault, L., Singh, A., Karadimas, S. K. & Fehlings, M. G. Degenerative
cervical myelopathy: Epidemiology, genetics, and pathogenesis. *Spine (Phila Pa 1976)*
40, (2015).

3.   Eippert, F., Kong, Y., Jenkinson, M., Tracey, I. & Brooks, J. C. W. Denoising spinal cord
fMRI data: Approaches to acquisition and analysis. *Neuroimage* **154**, 255–266 (2017).

4.   Cohen-Adad, J. & Wald, L. L. Array Coils. in *Quantitative MRI of the Spinal Cord* 59–67
(Elsevier, 2014). doi:10.1016/B978-0-12-396973-6.00005-8.

5.   Kinany, N., Pirondini, E., Micera, S. & Van De Ville, D. Spinal Cord fMRI: A New Window
into the Central Nervous System. *Neuroscientist* **29**, 715–731 (2022).

6.   Ellingson, B. M., Woodworth, D. C., Leu, K., Salamon, N. & Holly, L. T. Spinal Cord
Perfusion MR Imaging Implicates Both Ischemia and Hypoxia in the Pathogenesis of
Cervical Spondylosis. *World Neurosurg* **128**, (2019).

7.   Uemura, K. *et al.* Perfusion-weighted magnetic resonance imaging of the spinal cord in
cervical spondylotic myelopathy. *Neurol Med Chir (Tokyo)* **46**, (2006).

8.   Kanda, T. *et al.* Gadolinium-based contrast agent accumulates in the brain even in
subjects without severe renal dysfunction: Evaluation of autopsy brain specimens with
inductively coupled plasma mass spectroscopy. *Radiology* **276**, 228–232 (2015).

9.   Jahng, G. H., Li, K. L., Ostergaard, L. & Calamante, F. Perfusion magnetic resonance
imaging: A comprehensive update on principles and techniques. *Korean J Radiol* **15**, 554–
577 (2014).

10.   Frébourg, G. *et al.* Dynamic Contrast Enhanced MR perfusion of the Spinal Cord with
radial streaking artefacts reduction at 3T: Preliminary results and applications. in *Proc Intl
Soc Mag Reson Med 30* (2022).

11.   Meyer, B. P. *et al.* Optimized cervical spinal cord perfusion MRI after traumatic injury in
the rat. *Journal of Cerebral Blood Flow and Metabolism* **41**, (2021).

12.   Lee, S., Wilkins, N., Schmit, B. D., Kurpad, S. N. & Budde, M. D. Relationships between
spinal cord blood flow measured with flow-sensitive alternating inversion recovery (FAIR)
and neurobehavioral outcomes in rat spinal cord injury. *Magn Reson Imaging* **78**, 42–51
(2021).

13.   Nair, G. & Hu, X. P. Perfusion Imaging of the Human Cervical Spinal Cord. in *Proc Intl Soc
Mag Reson Med 18* (2010).

14.   Girard, O. M., Callot, V., Robert, B., Cozzone, P. J. & Duhamel, G. Perfusion MRI of the
Human Cervical Spinal Cord using Arterial Spin Labeling. in *Proc Intl Soc Mag Reson Med
21* (2013).

15.   Lévy, S. *et al.* Intravoxel Incoherent Motion at 7 Tesla to quantify human spinal cord
perfusion: limitations and promises. *Magn Reson Med* **84**, 1198–1217 (2020).

16.   Pinto, J., Bright, M. G., Bulte, D. P. & Figueiredo, P. Cerebrovascular Reactivity Mapping
Without Gas Challenges: A Methodological Guide. *Front Physiol* **11**, (2021).

17.   Urback, A. L., MacIntosh, B. J. & Goldstein, B. I. Cerebrovascular reactivity measured by
functional magnetic resonance imaging during breath-hold challenge: A systematic review.
Neurosci Biobehav Rev **79**, 27–47 (2017).

18.   Cohen-Adad, J. *et al.* BOLD signal responses to controlled hypercapnia in human spinal
cord. *Neuroimage* **50**, 1074–1084 (2010).

19. Barry, R. L. *et al.* Multi-shot acquisitions for stimulus-evoked spinal cord BOLD fMRI.
Magn Reson Med **85**, 2016–2026 (2021).

20. Moia, S. *et al.* Voxelwise optimization of hemodynamic lags to improve regional CVR
estimates in breath-hold fMRI. *Annu Int Conf IEEE Eng Med Biol Soc* **2020**, 1489–1492
(2020).

21. Chang, C., Thomason, M. E. & Glover, G. H. Mapping and correction of vascular
hemodynamic latency in the BOLD signal. *Neuroimage* **43**, (2008).

22. Geranmayeh, F., Wise, R. J. S., Leech, R. & Murphy, K. Measuring vascular reactivity with
breath-holds after stroke: A method to aid interpretation of group-level BOLD signal
changes in longitudinal fMRI studies. *Hum Brain Mapp* **36**, 1755–1771 (2015).

23. Cohen, A. D. & Wang, Y. Improving the Assessment of Breath-Holding Induced Cerebral
Vascular Reactivity Using a Multiband Multi-echo ASL/BOLD Sequence. *Sci Rep* **9**,
(2019).

24. Donahue, M. J. *et al.* Time delay processing of hypercapnic fMRI allows quantitative
parameterization of cerebrovascular reactivity and blood flow delays. *Journal of Cerebral
Blood Flow & Metabolism* **36**, (2016).

25. Pinto, J. *et al.* Evaluating cerebrovascular reactivity dynamics through a Bayesian
inference approach. in *Proc Intl Soc Mag Reson Med 30* (2022).

26. Bhogal, A. A. Medullary vein architecture modulates the white matter BOLD
cerebrovascular reactivity signal response to CO2: Observations from high-resolution T2*
weighted imaging at 7T. *Neuroimage* **245**, (2021).

27. Stickland, R. C. *et al.* A practical modification to a resting state fMRI protocol for improved
characterization of cerebrovascular function. *Neuroimage* **239**, (2021).

28. Bright, M. G., Bulte, D. P., Jezzard, P. & Duyn, J. H. Characterization of regional
heterogeneity in cerebrovascular reactivity dynamics using novel hypocapnia task and
BOLD fMRI. *Neuroimage* **48**, 166–175 (2009).

29. Leung, J., Duffin, J., Fisher, J. A. & Kassner, A. MRI-based cerebrovascular reactivity
using transfer function analysis reveals temporal group differences between patients with
sickle cell disease and healthy controls. *Neuroimage Clin* **12**, (2016).

30. Duffin, J. *et al.* The dynamics of cerebrovascular reactivity shown with transfer function
analysis. *Neuroimage* **114**, (2015).

31. Laumann, T. O. *et al.* Functional System and Areal Organization of a Highly Sampled
Individual Human Brain. *Neuron* **87**, 657–670 (2015).

32. Kang, K. C., Jang, T. S., Choi, S. H. & Kim, H. W. Difference between Anterior and
Posterior Cord Compression and Its Clinical Implication in Patients with Degenerative
Cervical Myelopathy. *J Clin Med* **12**, (2023).

33. Hemmerling, K. J., Hoggarth, M. A., Sandhu, M. S., Parrish, T. B. & Bright, M. G. Spatial
distribution of hand-grasp motor task activity in spinal cord functional magnetic resonance
imaging. *Hum Brain Mapp* **44**, 5567–5581 (2023).

34. Schulman, J. B. & Uludağ, K. Problems and solutions in quantifying cerebrovascular
reactivity using BOLD-MRI. *Imaging Neuroscience* **3**, (2025).

35. Tancredi, F. B. & Hoge, R. D. Comparison of Cerebral Vascular Reactivity Measures
Obtained Using Breath-Holding and CO$_2$ Inhalation. *Journal of Cerebral Blood Flow &
Metabolism* **33**, 1066–1074 (2013).

36. Bright, M. G. & Murphy, K. Reliable quantification of BOLD fMRI cerebrovascular reactivity
despite poor breath-hold performance. *Neuroimage* **83**, 559–568 (2013).

37.   Pinto, J., Jorge, J., Sousa, I., Vilela, P. & Figueiredo, P. Fourier modeling of the BOLD
681   response to a breath-hold task: Optimization and reproducibility. *Neuroimage* **135**, 223–
682   231 (2016).

683   38.   Mautes, A. E., Weinzierl, M. R., Donovan, F. & Noble, L. J. Vascular Events After Spinal
684   Cord Injury: Contribution to Secondary Pathogenesis Spinal Cord Injury Special Series.
685   *Phys Ther* **80**, 673–687 (2000).

686   39.   Tator, C. H. & Koyanagi, I. Vascular mechanisms in the pathophysiology of human spinal
687   cord injury. *J Neurosurg* **86**, 483–492 (1997).

688   40.   Qiu, M. *et al.* Arterial transit time effects in pulsed arterial spin labeling CBF mapping:
689   Insight from a PET and MR study in normal human subjects. *Magn Reson Med* **63**, (2010).

690   41.   Chen, Y., Wang, D. J. J. & Detre, J. A. Comparison of arterial transit times estimated
691   using arterial spin labeling. *Magnetic Resonance Materials in Physics, Biology and*
692   *Medicine* **25**, (2012).

693   42.   Thron, A. K. *Vascular Anatomy of the Spinal Cord*. (Springer US, 2016). doi:10.1007/978-
694   3-319-27440-9.

695   43.   Turnbull, I. M. Chapter 5. Blood supply of the spinal cord: normal and pathological
696   considerations. *Clin Neurosurg* **20**, (1973).

697   44.   Bhogal, A. A. *et al.* Examining the regional and cerebral depth-dependent BOLD
698   cerebrovascular reactivity response at 7T. *Neuroimage* **114**, 239–248 (2015).

699   45.   Glover, G. H., Li, T. Q. & Ress, D. Image-based method for retrospective correction of
700   physiological motion effects in fMRI: RETROICOR. *Magn Reson Med* **44**, 162–167 (2000).

701   46.   Brooks, J. C. W. *et al.* Physiological noise modelling for spinal functional magnetic
702   resonance imaging studies. *Neuroimage* **39**, 680–692 (2008).

703   47.   Power, J. D., Plitt, M., Laumann, T. O. & Martin, A. Sources and implications of whole-
704   brain fMRI signals in humans. *Neuroimage* **146**, 609–625 (2017).

705   48.   Deshpande, R. & Barry, R. Neptune: a toolbox for spinal cord functional MRI data
706   processing and quality assurance. in *Proc Intl Soc Mag Reson Med 30* (2022).
707   doi:https://doi.org/10.58530/2022/0396.

708   49.   Kastrup, A., Krüger, G., Neumann-Haefelin, T. & Moseley, M. E. Assessment of
709   cerebrovascular reactivity with functional magnetic resonance imaging: Comparison of
710   CO_2 and breath holding. *Magn Reson Imaging* **19**, 13–20 (2001).

711   50.   Zvolanek, K. M. *et al.* Comparing end-tidal CO_2, respiration volume per time (RVT), and
712   average gray matter signal for mapping cerebrovascular reactivity amplitude and delay
713   with breath-hold task BOLD fMRI. *Neuroimage* **272**, (2023).

714   51.   Friston, K. J. *et al.* Event-related fMRI: Characterizing differential responses. *Neuroimage*
715   **7**, (1998).

716   52.   Lipp, I., Murphy, K., Caseras, X. & Wise, R. G. Agreement and repeatability of vascular
717   reactivity estimates based on a breath-hold task and a resting state scan. *Neuroimage*
718   **113**, 387–396 (2015).

719   53.   Murphy, K., Harris, A. D. & Wise, R. G. Robustly measuring vascular reactivity differences
720   with breath-hold: Normalising stimulus-evoked and resting state BOLD fMRI data.
721   *Neuroimage* **54**, (2011).

722   54.   Sandhu, M. S. & Rymer, W. Z. Brief exposure to systemic hypoxia enhances plasticity of
723   the central nervous system in spinal cord injured animals and man. *Curr Opin Neurol* **34**,
724   (2021).

55. Tan, A. Q., Barth, S. & Trumbower, R. D. Acute Intermittent Hypoxia as a Potential
Adjuvant to Improve Walking Following Spinal Cord Injury: Evidence, Challenges, and
Future Directions. *Curr Phys Med Rehabil Rep* **8**, (2020).

56. Winkler, A. M., Ridgway, G. R., Webster, M. A., Smith, S. M. & Nichols, T. E. Permutation
inference for the general linear model. *Neuroimage* **92**, 381–397 (2014).

57. Smith, S. M. & Nichols, T. E. Threshold-free cluster enhancement: Addressing problems
of smoothing, threshold dependence and localisation in cluster inference. *Neuroimage* **44**,
83–98 (2009).

58. Nichols, T. E. & Holmes, A. P. Nonparametric Permutation Tests For Functional
Neuroimaging: A Primer with Examples. *Hum Brain Mapp* **15**, 1–25 (2002).

59. Li, X., Morgan, P. S., Ashburner, J., Smith, J. & Rorden, C. The first step for neuroimaging
data analysis: DICOM to NIfTI conversion. *J Neurosci Methods* **264**, 47–56 (2016).

60. Cox, R. W. AFNI: Software for analysis and visualization of functional magnetic resonance
neuroimages. *Computers and Biomedical Research* **29**, 162–173 (1996).

61. Cox, R. W. & Hyde, J. S. Software tools for analysis and visualization of fMRI data. *NMR
Biomed* **10**, 171–178 (1997).

62. De Leener, B. *et al.* PAM50: Unbiased multimodal template of the brainstem and spinal
cord aligned with the ICBM152 space. *Neuroimage* **165**, 170–179 (2018).

63. De Leener, B. *et al.* SCT: Spinal Cord Toolbox, an open-source software for processing
spinal cord MRI data. *Neuroimage* **145**, 24–43 (2017).

64. Gros, C. *et al.* Automatic segmentation of the spinal cord and intramedullary multiple
sclerosis lesions with convolutional neural networks. *Neuroimage* **184**, 901–915 (2019).

65. Hemmerling, K. & Bright, M. Restricted smoothing of spinal cord fMRI data resolves
structured temporal variation in heatmaps. in *Proc Organ Hum Brain Mapp 27* (2021).

66. Kong, Y., Jenkinson, M., Andersson, J., Tracey, I. & Brooks, J. C. W. Assessment of
physiological noise modelling methods for functional imaging of the spinal cord.
Neuroimage **60**, 1538–1549 (2012).

67. Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W. & Smith, S. M. FSL.
Neuroimage **62**, 782–790 (2012).

68. Woolrich, M. W., Ripley, B. D., Brady, M. & Smith, S. M. Temporal Autocorrelation in
Univariate Linear Modeling of FMRI Data. *Neuroimage* **14**, 1370–1386 (2001).

69. Bright, M. G., Tench, C. R. & Murphy, K. Potential pitfalls when denoising resting state
fMRI data using nuisance regression. *Neuroimage* **154**, 159–168 (2017).

70. Šidák, Z. Rectangular Confidence Regions for the Means of Multivariate Normal
Distributions. *J Am Stat Assoc* **62**, (1967).

Author Contributions: M.G.B., M.S.S., and T.B.P. designed research; K.J.H. and M.A.H.
performed research; K.J.H. analyzed data; K.J.H. wrote the initial draft; all authors reviewed
the manuscript.

Code availability statement: The code to perform the SCVR delay analysis is available at
https://github.com/BrightLab-ANVIL/SCVR.

Additional Information: The authors declare no competing interests.