Positioning of negative feedback loops within immune signaling pathways influences gene expression noise

Danial Asgari[1,2], Ann T. Tate[1,2*]

[1]*Department of Biological Sciences, Vanderbilt University, Nashville, Tennessee*

[2]*Evolutionary Studies Initiative, Vanderbilt University, Nashville, TN, USA*

[*]*Email: a.tate@vanderbilt.edu*

Abstract

Signaling pathways depend on negative and positive feedback loops (NFLs and PFLs) to regulate internal noise. Across diverse organisms, signaling is regulated by NFLs that function at different cellular locations. These range from NFLs functioning upstream near signal-receiving receptors to those downstream within the nucleus. While previous studies have examined the relationship between NFLs, internal noise in signaling pathways, and network topology, none have directly addressed how the cellular location of NFLs impacts noise regulation. This is significant given the almost ubiquitous presence of multi-level regulation systems within signaling pathways. Here, we use stochastic models inspired by Imd and Toll signaling to address this gap within the context of immune signaling. We use both mechanistic and evolutionary models to demonstrate how noise is regulated by NFLs and how this, in turn, affects the host's ability to fight off infection while minimizing immunopathologic effects of excessive immune gene expression. We found that downstream NFLs reduce noise in antimicrobial peptides (AMP) expression for some parameter values. On the other hand, upstream NFLs amplify the noise, but the presence of a strong PFL can reduce this noise. Our evolutionary simulations suggest that the mechanisms through which the downstream NFL operates within the cell can affect the evolution of the upstream NFLs. The results of our study provide insight into why distinct signaling pathways are regulated by varying numbers of NFLs, which operate in different cellular locations and employ diverse mechanisms to control gene expression.

Author Summary

Signaling pathways are noisy biological circuits. This noise is caused by random fluctuations in the number of proteins that function within these pathways. To properly respond to external

stimuli, the ratio of noise to information needs to be minimized. To regulate noise, biological

pathways produce proteins that either reduce (negative feedback) or amplify (positive feedback)

signaling following stimulation. Negative feedback loops can shut down signaling by interfering

with the first steps of signaling, which entail the detection of stimuli. Conversely, signaling might

be left intact, and instead, negative feedback loops might interfere with the last step, which is

the production of output. These regulatory differences can affect noise within signaling

pathways. Here, we examined this using stochastic (inherently random) models to simulate

immune signaling in response to pathogens. We found that negative feedback loops that

function at later stages of signaling can decrease noise in the output, while negative feedback

loops that act at earlier stages amplify the noise, but this noise can be reduced by the presence

of a strong positive feedback loop.

Introduction

Organisms rely on signaling pathways to monitor environmental changes and generate

appropriate responses [1–4]. Noise or stochasticity (inherent unpredictability) is a fundamental

characteristic of signaling pathways [5], which originates from random fluctuations in the small

number of signaling proteins and interactions between them [6]. Biological noise is influenced by

many factors such as the sequence of the promoter, nucleosome occupancy, and the number of

transcription factor binding sites [7]. Increased noise in gene expression is linked to cancer and

aging [8,9]. Therefore, proper homeostasis demands tight control over noise.

Signaling pathways regulate noise through several mechanisms. One such mechanism

involves feedback loops [10,11]. Feedback loops activate proteins that either amplify the signal

(positive feedback loops, or PFLs) or attenuate it (negative feedback loops, or NFLs). The

modulation of noise through feedback loops depends on several factors. These include the

source of the noise, which can be either due to environmental fluctuations (extrinsic) or due to

inherent fluctuations of the number of proteins involved in a signaling pathway (intrinsic) [12].

Autorepression, in which a gene suppresses its own activity, has been shown to reduce noise

within signaling pathways [13,14], however, this is often linked to extrinsic, not intrinsic, noise

[11,15]. The network topology and the specific mechanisms by which feedback operates can

further complicate noise modulation by NFLs. For example, Chakravarty et al. showed that, in

circadian networks, NFLs reduce intrinsic noise, while PFLs reduce extrinsic noise [16]. The

magnitude of noise modulation also varies depending on how the feedback functions within the

pathway. For example, mRNA-mediated feedback—where mRNA suppresses its own

translation (e.g., microRNA)—provides better control over noise at the protein level compared to

protein-mediated feedback, in which the protein downregulates the expression of its own gene

[17]. In reality, many signaling pathways employ NFLs at multiple levels of the transduction

cascade, which attenuate response through diverse mechanisms. However, it remains an open

question whether this layering mainly provides redundancy or whether each NFL within a

pathway contributes something unique or context-dependent to signal and noise control.

Immune signaling networks provide particularly salient case studies of the interaction of

layered NFLs in mediating noise. The immune signaling pathways of most multicellular

eukaryotes share a common architecture: signaling activates a central transcription factor that

induces target genes against pathogens as well as genes encoding NFLs to turn the response

back down, allowing the organism to conserve energy and mitigate immunopathological effects

[18,19]. Although there are biological differences across immune signaling pathways, NFLs that

regulate them generally fall into two categories: those that decrease input into the pathway by

acting upstream and those that function downstream to reduce the output while maintaining the

signaling process (Table 1).

Table 1. NFLs[*] at different steps of signaling in immune-related pathways. The Toll pathway in

insects is only regulated by an NFL downstream of the pathway.

Signaling pathways	Receptor	Upstream NFL	target	Downstream NFL	target
NF-κB (vertebrate)	TLR4 (innate) TIR (innate) TCR/BCR (adaptive)	A20[1,2]	RIP (outside the nucleus)	IκBα[3]	NF-κB (inside the nucleus)
Imd (invertebrate)	PGRP-LC	Pirk[4]	PGRP-LC (outside the nucleus)	Repressosome complex[5]	NF-κB (inside the nucleus)
Toll (invertebrate)	Toll	NA	NA	Cactus	NF-κB (inside the nucleus)
Jasmonate (plants)	JAT1	CYP94B3[6]	JA-Ile (outside the nucleus)	JAM[7]	MYC (inside the nucleus)

1: [20]; 2: [21] 3: [22]; 4: [23]; 5: [24]; 6: [25]; 7: [26]
[*]Here, NFLs are listed that are activated upon infection. These differ from negative regulators that control

the baseline expression of genes.

Simple theoretical models that consider shared versus unique topological attributes can

be used to clarify the relationship between noise, the response to stimuli, and the positioning of

NFLs and PFLs within pathways while also shedding light on their evolution. For example,

Asgari et al. [27] used a theoretical model of the fruit fly immune response to show that

upstream and downstream NFLs contribute differently to defense against pathogens with

varying proliferation rates. In the fruit fly, *Drosophila melanogaster*, the Toll pathway primarily

reacts against gram-positive bacteria and fungi [28,29] while the Imd pathway responds to

gram-negative bacteria [30], but both stimulate the expression of antimicrobial peptides (AMPs)

through the action of NF-κB transcription factors [31]. In addition, the NF-kB activates a PFL by

upregulating genes encoding receptors, which boosts the signal to provide a stronger response

to the pathogen [36]. Analogously to vertebrate NF-kB signaling (Table 1), the Imd pathway

includes two major NFLs: the induction of Pirk instigates the upstream removal of immune

receptors from the cell surface [23] while the Repressosome acts downstream [24]. The Toll

pathway, on the other hand, lacks an obvious upstream NFL and relies on the downstream

Cactus protein to provide negative feedback [32,33]. In both pathways, downstream NFLs inhibit

the activity of NF-κB transcription factors and reduce the output while maintaining signaling,

similar to JAM transcription factors competing with MYC2 to dampen Jasmonate signaling in

Arabidopsis [26]. Differences in the cellular location of NFLs (e.g., Pirk vs Repressosome) and

mechanistic differences in the function of NFLs operating at the same location (Cactus vs

Repressosome) could potentially yield distinct effects on noise regulation and fitness.[37].

In this study, we built stochastic models inspired by Imd and Toll signaling that contain

modules representing upstream NFLs and different types of downstream NFLs. We examined

how different NFLs contribute, individually and in tandem, to AMP output and noise during

repeated pathogen encounters. Considering that too much immune signaling can promote

pathology or impose other costs to host fitness, we conducted evolutionary simulations to

examine how NFLs functioning at different locations evolve to modulate immune response, and

how this, in turn, affects noise in AMP expression.

Methods

Models

We designed stochastic models containing only the fundamental components of an

immune signaling pathway. These entail a receptor that detects the pathogen and activates a

central transcription factor, which in turn induces AMPs that kill the pathogen, along with NFLs

that attenuate the response. Inspired by the Imd pathway, we included two NFL modules, acting

upstream (Pirk that removes receptors) and downstream (Repressosome that competes with

the central transcription factor Relish over binding to the promoter of AMP genes). Inspired by

Toll topology, we supplied a cactus-like module that could be swapped for the Imd

repressosome module to provide an alternative downstream NFL. Unlike the repressosome,

cactus inhibits the response by degrading Relish before it enters the nucleus (Fig.1). We

implemented the two models using the Gillespie algorithm [38], where reactions capture the

change of proteins within an immune signaling pathway. At each step, a reaction is

probabilistically chosen and carried out. The likelihood of a reaction depends on its rate relative

to the total rates of all reactions. In both models, time is discrete (T = 1,000). The complete

models are presented in detail in the S1 text. Both models entail two sets of reaction rates:

reaction rates for bacteria and reaction rates controlling immune signaling. If both sets are

empty (i.e., all reaction rates are 0), no reactions occur. If both sets contain non-zero rates, two

reactions occur in one step (one for bacteria, which is either elimination or proliferation, and one

for a randomly chosen protein of the immune network). Otherwise, one reaction occurs.

Fig. 1. Simplified diagrams of models inspired by the Imd and Toll pathway. Bacteria

proliferate at a rate k_0 and bind to receptors. NF-κB transcription factors are activated by

bacteria-receptor complexes at a rate β_2. NF-κB induces the production of receptors (β_1), Pirk,

which is the upstream NFL(β_3), Repressosome (in Imd-like model), and Cactus (in Toll-like

model), which are the downstream NFLs (β_4), and AMPs (β_5). Pirk reduces receptors in both

models. Repressosome competes with NF-κB for binding to the promoter of the AMP gene, and

Cactus inhibits the entry of NF-κB to the nucleus. The binding energies of the NF-κB and

Repressosome are Z_n and Z_s, respectively. The colors of the arrows are based on the affected

components in each reaction.

The stochastic reactions described above simulate inherent noise within the Imd and

Toll-inspired systems. To understand how these systems function in the absence of noise, we

developed deterministic models of the systems described above (S1 Text) using systems of

ordinary differential equations (ODEs). To solve these equations, we employed the Backward

Differentiation Formula (BDF) method via the solve_ivp function in Python, setting atol=10^{-6}

and rtol=10^{-8} to ensure numerical accuracy.

Parameter regimes

We considered four parameter regimes to examine the effect of NFLs on the dynamics

of Imd and Toll-inspired models (Table 2). The four regimes are based on two parameters that

specify the production rate of NFLs. The upstream ($\boldsymbol{\beta_3}$) and downstream ($\boldsymbol{\beta_4}$) NFLs are either

highly expressed (β_3 or $\beta_4 = \boldsymbol{10}$) or knocked down ($\beta_3$ or $\beta_4 = \boldsymbol{0}$). We chose these values based

on the qualitative assessments of several simulations before conducting the systematic analysis

of the models. We conduct additional simulations for other values of β_3 and β_4 for specific

scenarios as discussed in the Results section.

Table 2. Four parameter regimes capturing

different NFL activities upstream and

downstream of Imd and Toll signaling

Regime	parameters
Both knocked down	β_3 and $\beta_4 = 0$
Upstream active	$\beta_3 = 10$; $\beta_4 = 0$
Downstream active	$\beta_3 = 0$; $\beta_4 = 10$
Both active	β_3 and $\beta_4 = 10$

There are four background parameters $(\beta_1, \beta_2, \beta_5, \lambda)$ in Imd and Toll-like models. These

are distinct from parameters that determine the dynamics of NFLs (β_3, β_4, Z_s) and the binding

energy of NF-κB (Z_n). For each of the background parameters, we considered 10 different

values. The background parameter values range from 1 to 10 for $\beta_1, \beta_2, \beta_5$ and from 0.1 to 1 for

λ. We chose smaller values for λ because larger values result in a highly inefficient immune

response (i.e., the immune proteins degrade before the transmission of the signal). This

constitutes a total of 10,000 simulations to explore all combinations of background parameters

for one parameter regime (10,000 x 4 = 40,000 parameters for the four regimes in Table 2). In

the Imd-like model, across all parameter regimes, we maintained the following condition: $Z_s =$

$Z_n = 1$, unless otherwise stated. This ensures that the binding of the Repressosome is only

influenced by the relative number of NF-κB to Repressosome, which in turn is influenced by β_4

that varies across the four regimes. For consistency in the Toll-like model, we set $Z_n = 1$.

Calculation of AMP noise, average AMP expression, and fitness

For each parameter set, we conducted 10,000 replicate simulations, and quantified

noise in AMP expression at every time point of the host life time (T = 1,000) by calculating the

coefficient of variation [39] (Eq. 1). We then determined the magnitude of the noise by

calculating the arithmetic average of the noise across all time points (Eq. 2). We measured the

average AMP expression for each parameter set by calculating the arithmetic mean of average

AMP expression (average across 10,000 replicate simulations) across the lifetime of the host (T

= 1,000) (Eq. 3).

$(1)\ CV_t = \frac{\sigma_t}{\mu_t}$ $(2)\ Noise = \frac{\sum_{t=0}^{T} CV_t}{T}$ $(3)\overline{AMP} = \frac{\sum_{t=0}^{T} \mu_t}{T}$

Here μ_t and σ_t are the average and standard deviation across 10,000 replicate simulations.

If $\mu_t = \sigma_t = 0$, CV_t is set to 0.

We estimate the fitness by either calculating the host's ability to reduce bacterial load (Eq.4) or

by the host's ability to fight off infection while minimizing the AMP expression (Eq.5). Eq. 4

reflects a scenario where the pathogen virulence far outweighs immunopathology, while Eq. 5

weights them more equally. In Eqs. 4 and 5, A and B are the average value for these variables

across the lifespan of the host. Eq.5 is used to run evolutionary simulations of Imd and Toll

signaling.

$(4)\ e^{-[B]}$ $(5)\ e^{-[B+A]}$

Repeated encounters with pathogens create strong evolutionary pressure, which drives the

evolution of an optimized immune signaling network. Here, for all simulations, the host

encounters bacteria periodically at every 200 time points within the host lifetime (T = 1000, five

total encounters during the host lifetime). This captures the real-life condition of multiple

encounters with the pathogen [40,41].

Evolutionary simulations

We ran evolutionary simulations to find optimum parameters controlling the expression

rates of NFLs upstream and downstream of both Imd and Toll-like models. Each evolutionary

simulation consists of 500 replicates, each starting with a random set of parameter values, and

lasting for 5,000 generations. We are interested in the final distribution of parameters at the end

of each simulation. At each generation, a single mutation happens in a randomly selected

parameter among six parameters, which include four background parameters and two that

regulate upstream and downstream NFLs. The mutation changes the parameter value by either

increasing or decreasing it by 1 unit with equal probability, except for the degradation rate,

which changes by 0.1 unit. The boundary condition keeps parameters within the range [0.1,1]

for the degradation rate and between [1,10] for other parameters. If the mutation increases

fitness estimated using Eq.5, the mutation is accepted, otherwise, it's rejected. The simulations

do not consist of populations of evolving individuals; rather, they entail changes in parameter

values to optimize the immune response to pathogens according to Eq.5.

Results

The downstream NFL can suppress noise in gene expression, while

the upstream NFL reduces noise through its interaction with the PFL

Here, we use models inspired by Imd and Toll signaling pathways to understand the

effect of NFLs on noise in the final product of immune signaling (i.e., AMP) and ultimately host

fitness. Our model has parameters controlling the strength of NFLs, and background

parameters that specify other reactions, such as NF-κB activation rate, AMP expression rate,

and degradation rates of signaling proteins. The upstream NFL in both models is Pirk, which

removes cell surface receptors. The downstream NFL in the Imd-like model is the

Repressosome complex, which antagonistically competes with NF-κB for binding to the

promoter of the AMP gene. The downstream NFL in Toll-like model is Cactus, which prevents

entry of NF-κB to the nucleus.

To understand how upstream and downstream NFLs change AMP noise, for both

models, we compared noise in AMP expression in a circuit without NFLs (i.e., both knocked

down) to one with an upstream NFL, a downstream NFL, or both (Table.1). This yields three

pair-wise noise level comparisons (Δ noise). To avoid confounding noise (coefficient of

variation) with the average AMP expression, we only compared noise levels between

background parameter sets with the same average AMP expression, following Borri et al. (2016)

[42]. We found that in both models the upstream NFL amplifies the noise (Fig.2A). On the other

hand, the downstream NFL suppresses noise in AMP expression, but only for some background

parameter values (Fig.2B). When both NFLs are active, the effect of the upstream NFL (noise

amplification) is stronger than the downstream NFL (Fig.2C). For the Toll-like model and the

stronger binding of the downstream NFL in the Imd-like model, the downstream NFLs suppress

noise only when AMP expression is small (second row in Fig.2B and Fig.S1).

Fig.2. The downstream NFL suppresses noise (Δ noise < 0) in AMP expression for some

background parameters, while the upstream NFL amplifies noise for both Imd and Toll-

like models. Here, the Y-axis shows the difference in AMP noise between a signaling pathway

without NFLs (i.e., both NFLs are knocked down) and one with one NFL (first two columns to

the left) or both (third column to the right) NFLs, when the average AMP expression between

the two circuits (X-axis) are almost identical ($|\Delta$ average$| \leq 0.001$). The first row corresponds to

the Imd-inspired model, while the second row corresponds to the Toll-inspired model. The Y-

axis is tanh normalized. Here, the bacterial proliferation rate is low ($k_0 = 0.1$). For $k_0 = 0.5$, see

261   Fig.S2.

263   Lestas et al. (2010) [43] used information theory to show that noise suppression by

negative feedback has a fundamental limit. Therefore, we explored the limit of noise

suppression by calculating min[Δ noise] for different strengths of the downstream NFL. To this

end, we measured min[Δ noise] for weak ($\beta_4 = 1$), strong ($\beta_4 = 10$), and very strong ($\beta_4 = 20$)

expression of the downstream NFL, as well as for strong expression with 10× stronger binding

to the promoter ($\beta_4 = 10$ and $Z_s = 0.1$). In all other cases, binding strength was kept at $Z_s = 1$. We

found that maximum noise suppression occurs at strong downstream NFL expression ($\beta_4 = 10$),

while both lower and higher levels result in a weaker suppression (Fig.2D).

271   In biological circuits, NFLs and PFLs interact to adjust the level of noise [10]. Therefore,

we explored how the presence of one or both NFLs influences AMP noise across varying levels

of PFL. In both models, the receptor that detects the pathogen is activated by its downstream

protein (NF-κB) as a PFL. For the four parameter regimes, we plotted AMP noise as a function

of receptor activation rate (receptor activation ranging from 1 to 10) (Fig.3A), and calculated the

slope of the resulting line, while keeping other background parameters (other than receptor

activation rate) fixed. We did this for 1,000 different combinations of fixed background

parameter values. Unlike previous analyses, here we focus on changes in noise within each

regime for different receptor activation rates, whereas before (Fig.2) we compared the noise of

each regime to a regime with no NFLs. We observed that, for most parameter values, noise is

reduced for higher levels of PFL when both NFLs are present (Fig.3B, bottom right panel).

When only the upstream NFL is present, noise is reduced for some parameter values and

increased for others, while it is almost unchanged when only the downstream NFL is present

(Fig.3B). For the Toll-inspired model we observed the same results except that when only the

upstream NFL is present noise is almost always reduced (Fig.S3). Overall, when both NFLs are

active, the presence of a strong PFL reduces noise, but this effect is mostly linked to the

presence of the upstream NFL.

Fig.3. Within each regime, the noise in AMP expression is reduced for higher intensity of

the PFL when both NFLs are highly expressed, primarily due to the upstream NFL. Panel

A shows 10 trajectories of noise (y-axis) against receptor activation rates (x-axis) within two

regimes: both NFLs active (top) and only the downstream NFL active (bottom). In Panel B, the

X-axis shows the slope of the line between the strength of the PFL (R activation rate), which

ranges from 1 to 10 (x-axis in Panel A), and noise in AMP expression (y-axis in Panel A). When

the slope is negative, AMP noise is reduced for higher rates of receptor activation. The results

are shown for Imd signaling and $k_0 = 0.1$.

We next explored the effect of increasing NF-κB activation rate (β_2) rather than the

receptor which activates NF-κB as a PFL (i.e., focusing on positive regulator rather than the

positive feedback loop) on AMP noise within each of the four regimes. The results closely

mirrored those observed when changing the PFL (Fig.S4). It is worth noting that here we do not

control for the average expression when comparing noise levels. Thus, the reduction of noise

could be partly due to an increase in gene expression. However, the key observation is that

amplifying the signal, whether through the PFL (i.e., the receptor) or the positive regulator of the

signaling pathway (NF-κB) reduces noise, especially when both NFLs are present, and this

effect is mostly linked to the presence of the upstream NFL.

Both NFLs evolve for the Imd-inspired model, while the Toll-inspired model only evolves toward a downstream NFL

To understand the effect of AMP noise on fitness, we first evaluated the effect of AMP

noise on the organism's ability to fight off infection. To this end, we estimated fitness based on

the bacterial load ($F = e^{-B}$; as $B \to \infty, F \to 0$). We compared host fitness under conditions

with (via Gillespie simulation) and without noise (via ODE simulation). For optimal parameters

(i.e., high fitness for the ODE system—that is, in the absence of noise), we observed that the

fitness from the Gillespie algorithm highly depends on the AMP noise level. Specifically, optimal

parameters with high noise have low fitness, while those with low noise have high fitness

(Fig.S5). This shows that noise in AMP expression impairs the host's ability to fight off infection.

Given that noise is regulated differently by upstream and downstream NFLs (Fig.2 and

Fig.3) and that noisy AMP expression impairs the host response to the pathogen (Fig.S5), we

next asked how these NFLs evolve to optimize host fitness. To this end, we conducted

evolutionary simulations starting with random parameters that mutate by adding or subtracting a

small value to one of the six evolving parameters. These include background parameters and

parameters controlling the expression levels of the two NFLs. We assumed that the fitness

depends on the host's ability to reduce bacterial load while minimizing AMP expression to avoid

immunopathology $(e^{-[A+B]})$. Under this scenario, NFLs are expected to evolve and optimize

the level of AMP expression. We observed that when the downstream NFL competes with NF-

κB (Imd-inspired model), most simulations evolved toward either a strong downstream NFL (and

a weak upstream NFL) or toward a circuit with a strong expression of both NFLs. On the other

hand, when the downstream NFL affects the translocation of NF-κB (Toll-inspired model), the

system evolved only toward a strong downstream NFL (Fig.4A). This is interesting given that

Toll signaling in *D. melanogaster* lacks an upstream NFL. For both models, in some simulations,

AMP noise increased during evolution (red stars in Fig.4A). These results were robust to

changes in the bacterial proliferation rate (Fig.S6). When we used $e^{-[B]}$ as the fitness function,

noise did not evolve. This suggests that the evolution of noise is driven by the signaling

pathway's attempt to lower average AMP expression, which minimizes energetic costs and

immunopathology (Fig.S7).

Fig.4. Imd signaling evolves toward a system with a downstream NFL or a system

regulated by both NFLs, while Toll solely evolves toward a system with a downstream

NFL. Panel A shows the final values of parameters controlling downstream (X-axis) and

upstream (Y-axis) NFLs after 5,000 generations. Stars show simulations in which noise

increases across generations. The size of the circles and the darkness of the color for every

combination of NFLs are determined by the proportion of simulations that converge to that

combination of NFLs. Panel B shows the difference in noise values between average-matched

parameters across two regimes: both NFLs vs just upstream for Toll and Imd signaling.

Because the only difference between Toll and Imd-inspired models is in the mechanism

of action of the downstream NFL, we hypothesized that compared to Toll-inspired model, the

downstream NFL in Imd-inspired model is more effective in reducing the noise, which is

amplified by the upstream NFL (Fig.2). To test this, we asked how the presence of both NFL in

the two models affect the noise compared to a circuit with only the upstream NFL. To this end,

we compared AMP noise between average-matched parameters across these two regimes

(Fig.4B). We observed that the Imd-inspired model is slightly more effective in suppressing

noise than Toll signaling. This could partly explain the evolution of both NFLs in the Imd-inspired

model.

To further explore the effect of AMP noise on the evolution of NFLs, we ran deterministic

simulations of both Imd and Toll signaling for two bacterial proliferation rates ($k_0 = 0.1, 0.5$),

while allowing the six parameters to evolve. We observed that both signaling networks evolve

toward a broader range of parameter values (Fig.S8). This is because for most parameters,

ODE has a high fitness due to its rapid response to the pathogen. However, for a lower bacterial

proliferation rate, both signaling pathways evolved toward an upstream-biased NFL, consistent

with the results from Asgari et al. [27]. The evolution of the upstream-biased signaling pathways

in deterministic but not stochastic simulations confirms the deleterious effect of the upstream

NFL caused by high noise in AMP expression (Fig.2), which should be more pronounced for the

Toll-like model (Fig.4B).

Discussion

We estimated the impact of upstream and downstream NFLs on the noise level in AMP

expression using stochastic models inspired by Imd and Toll signaling. We found that,

compared to a circuit with no NFL, the upstream NFL amplifies noise, whereas, for some

parameter values, the downstream NFL reduces noise (Fig.2). Although the upstream NFL

amplifies noise, this noise can be reduced by the presence of a strong PFL, likely due to a

concomitant increase in gene expression (Fig.3). This is important given that noisy AMP

expression reduces the host's ability to fight off infection (Fig.S5). We have also shown that the

noisy expression of AMP evolves when AMP expression incurs a cost and needs to be

minimized (Fig.4A and Fig.S7). The evolutionary pattern of NFLs was consistent with the

structures of Imd and Toll signaling in *D. melanogaster* (Fig.4A)*.* Specifically, we found that the

Imd-inspired model evolves toward both upstream and downstream NFLs, whereas the Toll-

inspired model only converges toward the downstream NFLs. We suggest that this is due to the

effectiveness of the downstream NFL within the Imd-inspired model in reducing noise (Fig.4B).

The observation that an upstream NFL amplifies noise makes intuitive sense because

the upstream NFL reduces the input to the signaling pathway (e.g., by reducing receptors that

receive environmental signals). This stops the signaling before it propagates, which significantly

reduces the expression of the target gene (here AMP). A low expression results in significant

fluctuations in gene expression. On the other hand, a downstream NFL keeps signaling intact

but only regulates the target gene expression either directly (Repressosome in Imd) or indirectly

(Cactus in Toll). Borri et al. (2016) [42] showed that stronger protein-promoter binding in a

negative feedback system reduces noise in enzymatic reactions. This is consistent with the

effect of the downstream NFL on noise reduction in Imd signaling, where the Repressosome

binds to the promoter of AMP as an NFL. Because we observed a similar effect for the

downstream NFL in the Toll-inspired model, where the downstream NFL does not bind to the

promoter but instead reduces transcription factor entry into the nucleus, we suggest that

downstream NFLs, regardless of their mechanism of action, are capable of reducing noise in

biological circuits for at least some parameter values.

Signaling networks usually take advantage of both positive and negative feedback loops

to provide an appropriate response to environmental changes [10]. For example, a PFL can

contribute to plasticity by inducing a switch-like behavior [44] or promoting oscillations [45]. Here

we identify another interesting role for PFLs in biological circuits. We show that a strong PFL

can reduce the noise induced by the presence of an upstream NFL. In the Imd signaling

pathway, Pirk (or PGRP-LB) reduces the input to the system by preventing an interaction

between the bacterial peptidoglycan and the receptor [23,46]. This, according to the predictions

of our model, could increase noise in AMP expression, which reduces host's ability to fight off

infection, even compared to the same level of AMP expression without noise. NF-κB, however,

not only activates Pirk and PGRP-LB but also upregulates the receptor production itself [37].

This serves as a PFL, which attenuates noise in AMP expression and possibly improves the

host's ability to respond to pathogens.

Signaling pathways show variation in both the number of NFLs and in the mechanism by

which NFLs regulate signaling. For example, Wnt signaling, which is conserved across

multicellular organisms and controls different aspects of cellular development [47], is controlled

by both upstream (DKK3) and downstream (AXIN2) NFLs [48,49]. Hippo signaling, which is also

crucial for the developmental process across species, is regulated by downstream NFLs

(LATS1/2), which interact with transcription factors that activate target genes [50,51]. The

interaction between an NFL and its target protein can be direct or indirect. For example, in NF-

κB signaling of mammals, A20 indirectly interacts with the receptor (through RIP-1), whereas

Pirk in Imd signaling directly interacts with the receptor. These differences in the number and

function of NFLs can have consequences on the evolution of NFLs. For example, Asgari and

Tate [52] have shown that downstream NFLs are more resistant to evolutionary changes due to

direct interaction with the target genes. Here, by incorporating noise in our model we show that t

the downstream NFL in Imd signaling (Repressosome), which directly binds to the promoter of

the target gene, is more efficient in noise reduction than the downstream NFL in Toll signaling

(Cactus), which only inhibits the entry of the transcription factor to the nucleus. Given that we

found upstream NFLs amplify noise (Fig.2), this could partly explain why Toll signaling in

organisms such as *D. melanogaster* lacks an upstream NFL.

In conclusion, by using stochastic models inspired by the two main immune signaling

networks in invertebrates, we estimate the effect of upstream and downstream NFLs on noise

and ultimately on host fitness. Our research provides insights into why different signaling

pathways vary in the number and placement of upstream and downstream NFLs (and PFLs),

helping us better understand and appreciate the diversity of signaling architectures.

Acknowledgments

This work was supported by the National Institute of General Medical Sciences at the National

Institutes of Health (grant number R35GM138007 to A.T.T.).

This work used Purdue Anvil resources at Purdue University through allocation BIO240182 from

the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS)

program, which is supported by U.S. National Science Foundation grants #2138259, #2138286,

#2138307, #2137603, and #2138296.

Author Contributions

D.A. and A.T.T. conceived and designed the study. D.A. performed the simulations and wrote

the first draft. D.A. and A.T.T. contributed to the final draft.

Data Availability Statement

We performed all simulations in Python. All codes are accessible at:

https://github.com/danialasg74/Positioning-of-negative-feedback-loops-and-noise/tree/main

References

[1]   Kawasaki T, Kawai T. Toll-like receptor signaling pathways. Front Immunol 2014;5:461.

[2]   Komiya Y, Habas R. Wnt signal transduction pathways. Organogenesis 2008;4:68–75.

[3]   De Meyts P. The Insulin Receptor and Its Signal Transduction Network. MDText.com, Inc.; 2016.

[4]   Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. Cell 2017;168:960–76.

[5]   Tsimring LS. Noise in biology. Rep Prog Phys 2014;77:026601.

[6]   Ladbury JE, Arold ST. Noise in cellular signaling pathways: causes and effects. Trends Biochem Sci 2012;37:173–8.

[7]   Sanchez A, Choubey S, Kondev J. Regulation of noise in gene expression. Annu Rev Biophys 2013;42:469–91.

[8]   Bahar R, Hartmann CH, Rodriguez KA, Denny AD, Busuttil RA, Dollé MET, et al. Increased cell-to-cell variation in gene expression in ageing mouse heart. Nature 2006;441:1011–4.

[9]   Kemkemer R, Schrank S, Vogel W, Gruler H, Kaufmann D. Increased noise as an effect of haploinsufficiency of the tumor-suppressor gene neurofibromatosis type 1 in vitro. Proceedings of the National Academy of Sciences of the United States of America 2002;99. https://doi.org/10.1073/pnas.212386999.

[10] Zhang H, Chen Y, Chen Y. Noise Propagation in Gene Regulation Networks Involving Interlinked Positive and Negative Feedback Loops. PLOS ONE 2012;7:e51840.

[11] Hooshangi S, Weiss R. The effect of negative feedback on noise propagation in transcriptional gene networks. Chaos 2006;16:026108.

[12] Thomas P. Intrinsic and extrinsic noise of gene expression in lineage trees. Scientific Reports 2019;9:1–16.

[13] Becskei A, Serrano L. Engineering stability in gene networks by autoregulation. Nature 2000;405. https://doi.org/10.1038/35014651.

[14] Shahrezaei V, Ollivier JF, Swain PS. Colored extrinsic fluctuations and stochastic gene expression. Molecular Systems Biology 2008. https://doi.org/10.1038/msb.2008.31.

[15] Dublanche Y, Michalodimitrakis K, Kümmerer N, Foglierini M, Serrano L. Noise in transcription negative feedback loops: simulation and experimental analysis. Molecular Systems Biology 2006. https://doi.org/10.1038/msb4100081.

[16] Chakravarty S, Hong CI, Csikász-Nagy A. Systematic analysis of negative and positive feedback

loops for robustness and temperature compensation in circadian rhythms. Npj Systems Biology and
Applications 2023;9:1–10.
[17] Singh A. Negative Feedback Through mRNA Provides the Best Control of Gene-Expression Noise
n.d. https://ieeexplore.ieee.org/abstract/document/6048013 (accessed March 21, 2025).
[18] DiAngelo JR, Bland ML, Bambina S, Cherry S, Birnbaum MJ. The immune response attenuates
growth and nutrient storage in Drosophila by reducing insulin signaling. Proceedings of the National
Academy of Sciences 2009;106:20853–8.
[19] Gabryšová L, Nicolson KS, Streeter HB, Verhagen J, Sabatos-Peyton CA, Morgan DJ, et al. Negative
feedback control of the autoimmune response through antigen-induced differentiation of IL-10–
secreting Th1 cells. The Journal of Experimental Medicine 2009;206:1755.
[20] Pujari R, Hunte R, Khan WN, Shembade N. A20-mediated negative regulation of canonical NF-κB
signaling pathway. Immunol Res 2013;57:166–71.
[21] Coornaert B, Carpentier I, Beyaert R. A20: central gatekeeper in inflammation and immunity. J Biol
Chem 2009;284:8217–21.
[22] Zabel U, Baeuerle PA. Purified human I kappa B can rapidly dissociate the complex of the NF-kappa
B transcription factor with its cognate DNA. Cell 1990;61:255–65.
[23] Kleino A, Myllymäki H, Kallio J, Vanha-aho L-M, Oksanen K, Ulvila J, et al. Pirk is a negative
regulator of the *Drosophila* Imd pathway. J Immunol 2008;180:5413–22.
[24] Kim LK, Choi UY, Cho HS, Lee JS, Lee W-B, Kim J, et al. Down-regulation of NF-kappaB target
genes by the AP-1 and STAT complex during the innate immune response in *Drosophila*. PLoS Biol
2007;5:e238.
[25] Koo AJK, Cooke TF, Howe GA. Cytochrome P450 CYP94B3 mediates catabolism and inactivation of
the plant hormone jasmonoyl-L-isoleucine. Proc Natl Acad Sci U S A 2011;108:9298–303.
[26] Sasaki-Sekimoto Y, Jikumaru Y, Obayashi T, Saito H, Masuda S, Kamiya Y, et al. Basic helix-loop-
helix transcription factors JASMONATE-ASSOCIATED MYC2-LIKE1 (JAM1), JAM2, and JAM3 are
negative regulators of jasmonate responses in *Arabidopsis*. Plant Physiol 2013;163:291–304.
[27] Asgari D, Stewart AJ, Meisel RP. The role of uncertainty and negative feedback loops in the evolution
of induced immune defenses. G3 (Bethesda, Md) 2024;14. https://doi.org/10.1093/g3journal/jkae182.
[28] Valanne S, Wang J-H, Rämet M. The Drosophila Toll signaling pathway. J Immunol 2011;186:649–
56.
[29] Buchon N, Broderick NA, Poidevin M, Pradervand S, Lemaitre B. Drosophila intestinal response to
bacterial infection: activation of host defense and stem cell proliferation. Cell Host Microbe
2009;5:200–11.
[30] Myllymäki H, Valanne S, Rämet M. The Drosophila imd signaling pathway. J Immunol
2014;192:3455–62.
[31] De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. The Toll and Imd pathways are the
major regulators of the immune response in Drosophila. The EMBO Journal 2002;21:2568.
[32] Belvin MP, Anderson KV. A conserved signaling pathway: the *Drosophila* toll-dorsal pathway. Annu
Rev Cell Dev Biol 1996;12:393–416.
[33] Cai Q, Guo H, Fang R, Hua Y, Zhu Y, Zheng X, et al. A Toll-dependent Bre1/Rad6-cact feedback
loop in controlling host innate immune response. Cell Rep 2022;41:111795.
[34] Shi X, Zhan X, Wu J. A positive feedback loop between Gli1 and tyrosine kinase Hck amplifies shh
signaling activities in medulloblastoma. Oncogenesis 2015;4:e176–e176.
[35] Harris SL, Levine AJ. The p53 pathway: positive and negative feedback loops. Oncogene
2005;24:2899–908.
[36] Georgopoulos NT, Kirkwood LA, Southgate J. A novel bidirectional positive-feedback loop between
Wnt–β-catenin and EGFR–ERK plays a role in context-specific modulation of epithelial tissue
regeneration. Journal of Cell Science 2014;127:2967.
[37] Liu Y, Ye N, Chen M, Zhao H, An J. Structural and Functional Analysis of PGRP-LC Indicates
Exclusive Dap-Type PGN Binding in Bumblebees. International Journal of Molecular Sciences
2020;21. https://doi.org/10.3390/ijms21072441.
[38] Gillespie DT. A general method for numerically simulating the stochastic time evolution of coupled
chemical reactions. J Comput Phys 1976;22:403–34.
[39] Wang Z, Zhang J. Impact of gene expression noise on organismal fitness and the efficacy of natural
selection. Proc Natl Acad Sci U S A 2011;108:E67–76.
[40] Dutt A, Andrivon D, Le May C. Multi-infections, competitive interactions, and pathogen coexistence.

Plant Pathology 2022;71:5–22.
[41] Nayduch D, Burrus RG. Flourishing in Filth: House Fly–Microbe Interactions Across Life History. Ann
Entomol Soc Am 2017;110:6–18.
[42] Borri A, Palumbo P, Singh A. Impact of negative feedback in metabolic noise propagation. IET
Systems Biology 2016;10:179–86.
[43] Lestas I, Vinnicombe G, Paulsson J. Fundamental limits on the suppression of molecular fluctuations.
Nature 2010;467:174–8.
[44] Shinohara H, Behar M, Inoue K, Hiroshima M, Yasuda T, Nagashima T, et al. Positive feedback
within a kinase signaling complex functions as a switch mechanism for NF-κB activation. Science
2014;344:760–4.
[45] Ananthasubramaniam B, Herzel H. Positive Feedback Promotes Oscillations in Negative Feedback
Loops. PLOS ONE 2014;9:e104761.
[46] Zaidman-Rémy A, Hervé M, Poidevin M, Pili-Floury S, Kim MS, Blanot D, et al. The Drosophila
amidase PGRP-LB modulates the immune response to bacterial infection. Immunity 2006;24.
https://doi.org/10.1016/j.immuni.2006.02.012.
[47] Hayat R, Manzoor M, Hussain A. Wnt signaling pathway: A comprehensive review. Cell Biology
International 2022;46:863–77.
[48] Mao B, Wu W, Davidson G, Marhold J, Li M, Mechler BM, et al. Kremen proteins are Dickkopf
receptors that regulate Wnt/β-catenin signalling. Nature 2002;417:664–7.
[49] Bernkopf DB, Hadjihannas MV, Behrens J. Negative-feedback regulation of the Wnt pathway by
conductin/axin2 involves insensitivity to upstream signalling. Journal of Cell Science 2015;128.
https://doi.org/10.1242/jcs.159145.
[50] Fu M, Hu Y, Lan T, Guan K-L, Luo T, Luo M. The Hippo signalling pathway and its implications in
human health and diseases. Signal Transduction and Targeted Therapy 2022;7:1–20.
[51] Moroishi T, Park HW, Qin B, Chen Q, Meng Z, Plouffe SW, et al. A YAP/TAZ-induced feedback
mechanism regulates Hippo pathway homeostasis. Genes & Development 2015;29.
https://doi.org/10.1101/gad.262816.115.
[52] Asgari D, Tate AT. How the Structure of Signaling Regulation Evolves: Insights from an Evolutionary
Model. Mol Biol Evol 2025:msaf104.

Supporting information

S1 Text. Supplementary information file, including a detailed description of the Gillespie algorithm and the deterministic model. The first section contains a detailed description of simulations of Imd and Toll signaling using the Gillespie algorithm. The second section shows the system of ODE modeling deterministic signaling pathways.

Fig.S1. In Imd signaling, upon strong binding of the downstream NFL to the promoter of AMP genes, AMP noise is suppressed only when gene expression is small. The change in noise is shown for a downstream biased regime with a strong expression of the downstream NFL ($\beta_4 = 10$) and 10x stronger binding ($Z_s = 0.1$) compared to Fig.2 is compared to a circuit with no NFL. The comparison is done for parameters with the same average AMP expression.

Fig.S2. For a high bacterial proliferation rate, the downstream NFL reduces AMP noise for some parameters, while the upstream NFL amplifies noise. The change in noise across three regimes (upstream biased, downstream biased, both from left to right) compared to a circuit with no NFLs. The comparison is done for parameters with the same average AMP expression. Here, the bacterial proliferation is set to $k_0 = 0.5$.

Fig.S3. The combination of the upstream NFL and a strong expression of PFL reduces AMP noise. The slope between receptor activation rate and AMP noise is plotted for Toll signaling ($k_0 = 0.1$) at the top, Imd signaling ($k_0 = 0.5$) at the bottom left, and Toll ($k_0 = 0.5$) at the bottom right. Positive slope indicates that increasing receptor activation rate increases noise, whereas negative values indicate that increasing the receptor activation rate reduces noise.

Fig.S4. The combination of the upstream NFL and a strong expression of the positive regulator (NF-κB) reduces AMP noise. The slope between NF-kB activation rate and noise is plotted for Imd signaling on the left and Toll signaling on the right. Positive slope indicates that increasing NFkB activation rate increases noise, whereas negative values indicate that increasing NFkB activation rate reduces noise. Here, we set $k_0 = 0.1$.

Fig.S5. Noise reduces fitness even for optimal parameters (high fitness in the ODE model). Fitness (e^{-B}) values for different combinations of background parameters are shown using the Gillespie algorithm (Y-axis) and the ODE system (X-axis). The heatmap color represents the level of noise. Four parameter regimes are shown: No NFL, downstream NFL only (Down), upstream NFL only (Up), and both NFLs. To enhance the visibility of the pattern, the noise values are log-normalized. Imd signaling results are shown on the left, and the results for Toll signaling are on the right. For each pathway, in the top two panels, the fitness of the ODE system is always near 1, regardless of the background parameter value. Here we set $k_0 = 0.1$.

Fig.S6. When bacterial proliferation is high, immune signaling pathways evolve either
towards a downstream NFL or both NFLs. Evolution of Imd (left) and Toll signaling when
bacterial proliferation rate is high ($k_0 = 0.5$). Plots show the final values of parameters
controlling downstream (X-axis) and upstream (Y-axis) NFLs after 5,000 generations. Stars
show simulations in which noise increases across generations. The size of the circles and the
darkness of the color for every combination of NFLs are determined by the proportion of
simulations that converge to that combination of NFLs.

Fig.S7. Noise only evolves when signaling pathways must reduce the cost of AMP
expression. The correlation plots on the left are shown for e-[A+B] and e-[B] fitness functions.
Here, we measured the Spearman correlation between noise (and average expression) and
fitness across 5,000 generations of Imd and Toll evolution. Since fitness only increases during
the evolution of both pathways (i.e., mutations that decrease fitness are rejected), a positive
correlation with fitness indicates that the parameters (noise or average expression) is increasing
across generations, whereas a negative correlation means the parameter is decreasing as the
evolution continues. Here, we only see negative correlation for noise values (red plots) when we
have e-[B]. On the other hand, average expression shows positive correlation when we have e-
[B] to eliminate the pathogen. The results for e-[A+B] is more complicated because the signaling
pathway should minimize both B and A. Minimizing A increases noise, and results in positive
correlation values in some simulations when we have e-[A+B]. On the right the final
combinations of upstream (Y-axis) and downstream (X-axis) NFLs are shown after 5000
generations when we have e-[B]. For all simulations presented here, we set $k_0 = 0.1$.

Fig.S8. For a lower bacterial proliferation rate, both Imd and Toll signaling evolve toward an upstream-biased NFL. The evolution of a deterministic Imd (on the top) and Toll (on the bottom) signaling for two bacterial proliferation rates (k_0=0.1 on the left and k_0=0.5 on the right). The brighter colors show combinations of downstream and upstream NFLs that evolve more often.