Effects of childhood hearing loss on the subcortical and cortical representation

of speech

Axelle Calcus[1, 2, 3] & Stuart Rosen[1]

[1] Department of Speech, Hearing and Phonetic Sciences, UCL, London WC1N 1PF, UK

[2] Laboratoire des Systèmes Perceptifs, CNRS UMR 8248, École Normale Supérieure, Paris

Sciences et Lettres Research University, Paris, France

[3] Center for Research in Cognition & Neurosciences (CRCN), ULB Neuroscience Institute

(UNI), Université Libre de Bruxelles (ULB), Brussels, Belgium

Corresponding author: Axelle Calcus (axelle.calcus@ulb.be)

Abbreviated title: Speech processing in childhood hearing loss

Number of figures: 7; Number of supplementary figures: 2; Number of tables: 2 (submit with

Tables_v9.docx – supplementary tables)

The authors declare no conflict of interest

Abstract

Little is known about the effects of childhood mild-to-moderate sensorineural hearing loss (MM

HL) on the function of the auditory pathway. We aimed to examine the effect of childhood MM

HL and the benefit of frequency-specific amplification on both subcortical and cortical auditory

processing, and to relate it to speech-perceptual abilities. We recorded subcortical and cortical

responses to speech syllables in nineteen children with congenital MM HL (unamplified and

amplified), and sixteen children with typical hearing (unamplified sounds only). Speech

perception was measured behaviourally. Congenital HL led to smaller subcortical and cortical

responses to unamplified speech sounds. There was a significant benefit of amplification on

subcortical and early, but not late, cortical responses, with some effects differing across age.

No relationship was found between the neural and behavioural measures. Childhood MM HL

affects both subcortical and cortical processing of speech. Amplification mostly benefits

subcortical processing of speech in younger children. Childhood HL leads to functional changes

in the processing of sounds, with amplification differentially affecting subcortical and cortical

levels of the auditory pathway.

Introduction

Animal studies have shown biological and physiological alterations in the properties of neurons in the cochlea, subcortex, and cortex following even a temporary, mild-to-moderate (MM) hearing loss (HL; Caras & Sanes, 2015; Mowery et al., 2015). Similar changes have been documented in humans with congenital deafness (Daly et al. 1976; Ponton et al. 2000; Sharma et al. 1997), or in older adults with age-related MM HL (for a review see Peelle & Wingfield 2016). Nowadays, most individuals with MM HL who seek treatment receive auditory stimuli via hearing aids, which amplify the incoming auditory signal. To date, the bulk of studies investigating the effects of MM HL on the functional integrity of the auditory pathway (and the potential benefit of hearing aid amplification) has focused on older adults. However, congenital hearing loss affects up to 2 live births in 1000 (Lieu et al. 2020), milder degrees of loss being more prevalent yet more often overlooked. Here, we evaluate the effect of childhood MM HL and the benefit of amplification on simultaneously recorded subcortical and cortical responses to speech sounds.

In the auditory system, speech sounds travel from the cochleae to the auditory cortices, undergoing increasingly complex levels of processing. At the level of the cochleae, sounds undergo a frequency analysis. The output of each channel can be separated into their relatively slowly fluctuating modulated signals known as the envelope (ENV), imposed upon a rapidly varying carrier waveform, known as the temporal fine structure (TFS). Both ENV and TFS cues are thought to be important in decoding speech. ENV cues support the robust identification of speech in quiet (Shannon et al. 1995), whereas TFS cues are thought to be especially useful in noisy backgrounds (Qin & Oxenham 2003; Zeng et al. 2005). Indeed, individuals with typical hearing (TH) typically benefit from the TFS when they glimpse into the "dips" (i.e., local improvements in SNR) of a fluctuating background noise (Hopkins & Moore 2009; Cooke 2006). Therefore, speech intelligibility is much better in the presence of fluctuating than steady

background noise, a phenomenon termed "masking release" (Miller & Licklider 1950; Festen

& Plomp 1990). However, adults with HL show reduced ability to use TFS (Lorenzi et al.

2006), a difficulty that limits the amount of masking release they experience (Hopkins et al.

2008; Strelcyk & Dau 2009). Similarly to adults, children and adolescents with MM HL

experience poorer speech intelligibility in noise (Crandell 1993; Lewis et al. 2015; Goldsworthy

& Markle 2019). They show reduced sensitivity to TFS compared to children/adolescents with

TH (Halliday et al. 2019), which likely limits masking release of speech in the presence of a

fluctuating background noise, even when they wear hearing aids (Brennan et al. 2016).

At the subcortical level of the auditory pathway, the frequency following response

(FFR) reflects the periodicity of speech sounds (Kraus et al. 2017). Adding the neural responses

to speech sounds presented in alternating polarities is thought to enhance the ENV of the

response (henceforth FFR_{ENV}, here used as an index of fundamental frequency encoding),

whereas subtracting them is considered to enhance the TFS (henceforth FFR_{TFS}, an index of

spectral harmonic encoding (Aiken & Picton 2008)). Although the primary generator of the

FFR appears to vary according to the fundamental frequency (F0) of the input stimulus, the

current view is that the FFR emerges from multiple generators, including the cochlear nucleus,

inferior colliculus and thalamus (Bidelman 2018; Coffey et al. 2019). Early studies suggested

that the auditory brainstem reached maturity within the first 18 months of life (Hecox &

Galambos 1974). However, more recent studies indicate that the FFR evoked by speech sounds

keeps maturing well into the childhood years (Krizman et al. 2015; Krizman et al. 2019).

At the cortical level, components of the late auditory event-related potentials (LAERs)

arise which are thought to reflect the initial detection (P1, N1, and P2), classification and

discrimination (P2 and N2) of auditory stimuli, reflecting processing for auditory events

ranging from isolated sounds to complex auditory scenes (Alain & Tremblay 2007). LAERs

are thought to be generated in the auditory cortex (for a review, see Eggermont & Ponton,

2002), with additional contributors from non-auditory (e.g., prefrontal, premotor) areas (Scherg et al. 1989). The mismatch negativity (MMN) is thought to reflect the discrimination of a deviant sound, and is typically evoked by a comparatively rare deviant in a stream of repeated standards (i.e. an oddball paradigm; for a review see Näätänen et al. 2012). The MMN is thought to have several additional generators, including bilateral temporal cortex, right inferior frontal gyrus, and bilateral frontal and centro-parietal regions (e.g., Alho 1995; Rinne et al. 2000; Zhang et al. 2018). In addition to distinct topographies, both LAERs and MMNs show distinct developmental trajectories. In typical development, P1 and N2 responses are thought to decrease in amplitude as children grow older, while N1 and P2 progressively appear in childhood, growing in amplitude until adolescence (Bishop et al. 2007; Mahajan & McArthur 2012). MMN amplitude is thought to increase throughout childhood (Wetzel et al. 2006), but to be stable from around 10 years of age to adulthood (Mahajan & McArthur 2015; Mueller et al. 2008) - although some studies report a prolonged increase in MMN amplitude throughout adolescence (Bishop et al., 2011; Oades et al., 1997).

Both subcortical and cortical levels of the auditory pathway thus appear to follow a protracted trajectory, maturing until late adolescence. How MM HL affects the development of subcortical and cortical speech processing remains largely unknown. So far, studies have focused on *either* subcortical *or* cortical levels of processing, only providing fragmented information about the physiology of the auditory pathway in children/adolescents with MM HL. Yet using the right set of recording/analysis parameters, it is now possible to obtain subcortical and cortical responses during the same recording session (Krishnan et al. 2012; Bidelman, 2015). Therefore, the first aim of this project was to evaluate the effect of childhood hearing loss on simultaneously recorded subcortical and cortical levels of processing. Noteworthily, the existing literature on this topic can be classified according to the strategy used to manage intensity levels across listener groups.

Neural processing of speech presented at fixed intensity levels across groups with and without

HL

When studying the effect of hearing loss on auditory processing, researchers often

choose to investigate the unaided performance of individuals with HL. Stimuli are thus

presented at a fixed intensity level for all individuals, with HL or not. Adult studies using this

strategy to evaluate the effect of HL on subcortical processing have led to contradictory findings

(for a review, see Jacxsens et al. 2024). While some authors reported smaller FFR_{ENV} and

FFR_{TFS} in adults with HL than adults with TH (Plyler & Ananthanarayan 2001;

Ananthakrishnan et al. 2016), others did not show such group differences (Anderson et al. 2013;

Presacco et al. 2019). Note that a possible explanation for this discrepancy lies in the age-

matching across groups (Anderson et al. 2013; Presacco et al. 2019), or lack thereof (Plyler &

Ananthanarayan 2001; Ananthakrishnan et al. 2016). To our knowledge, no study has focused

on the effects of childhood MM HL on subcortical processing of speech, using fixed intensity

stimuli across groups of children/adolescents.

At the cortical level, studies using fixed intensity levels across groups suggest that

childhood MM HL leads to delayed and/or smaller LAERs evoked by speech or pure tones

(Calcus et al. 2019; Ji et al. 2023). Developmental changes following MM HL were also

apparent in the MMN: While 8-11 year old children with MM HL showed age-appropriate

MMN responses to sounds, this was no longer the case in 12-16 year old adolescents with MM

HL, as shown both cross-sectionally and longitudinally (Calcus et al. 2019). This suggests that

some effects of childhood MM HL might only reveal themselves in adolescence, highlighting

the need to take age into account in developmental studies.

Neural processing of speech presented at similar sensation levels across groups

A common limitation of the above-mentioned studies using fixed intensity levels for all

participants is that individuals with HL would have perceived the stimuli at reduced sensation

levels (SL) relative to controls. Therefore, some researchers have attempted to equate for the

effects of peripheral hearing loss by presenting listeners with TH and those with SNHL are

presented with sounds at different intensities – more intense for individuals with SNHL (either

by simulating the frequency-specific gain provided by individuals' hearing aids, or by recording

the neurophysiological response while listeners wear hearing aids). With such adjustments for

HL, adults with acquired MM HL show larger FFR_{ENV} to an amplified speech stimulus

compared to an unamplified one (BinKhamis et al. 2019; Easwar et al. 2015) and in one study,

than the FFR_{ENV} evoked by unamplified speech presented to TH listeners (Anderson et al.

2013). A similar benefit of amplification on the FFR_{ENV} evoked by the speech token /suʃi/ was

recently reported in 18 children with HL aged 6 to 17 years, compared to the unamplified

stimulus (Easwar et al. 2023). Note that older adults with acquired MM HL did not show a

significant benefit of amplification on FFR_{TFS} (Anderson et al., 2013). To our knowledge, the

effect of amplification and childhood HL on FFR_{TFS} has remained unexplored so far.

At the cortical level, studies looking at the effect of childhood HL on LAERs by

accounting for the effects of peripheral HL have led to contradictory findings. Pilot data from

a small sample of 5 children aged 2 to 6 years show age-appropriate P1 and N2 evoked by

speech stimuli when the children wore their hearing aids (Martinez et al. 2013). Another study

focused on older, unaided children (9-12 years) with MM HL, to whom speech stimuli were

presented on average 18 dB greater than in children with TH (Koravand et al. 2013). Note that

here, the amplification provided to the group with HL was not frequency-specific – unlike

hearing aids. Results indicate age-appropriate P1 and MMN responses in children with MM

HL, but smaller (but not later) N2 responses compared to children with TH. However, 5 to 8

year-old children who wore their hearing aids (n = 7) showed significantly smaller LAERs

evoked by complex tones than age-matched children with TH (Engström et al. 2021).

Interestingly, when re-tested three years later, the same children with HL showed age-

appropriate LAERs. At both time-points, MMN amplitudes were age-appropriate in children

with MM HL. To our knowledge, no study has directly investigated the intra-individual benefit

of amplification on the cortical processing of speech in children/adolescents with MM HL.

Relationship between speech processing and perception

A main stake of auditory neurophysiological studies is to relate neural processing to

speech perception in a variety of contexts and clinical populations. Because the FFR reflects

encoding of F0 and lower harmonics of the signal, that are particularly helpful in adverse

listening environments, it is thought to relate to speech perception in noise. Better speech

perception in noise has been associated with subcortical processing of complex sounds in quiet,

as observed in children (Anderson et al. 2010; Hornickel et al. 2009), young (Bidelman &

Momtaz 2021) and older adults with TH (Mai et al. 2018; Mai & Howell 2023); but see

Presacco et al., 2016 for a study that did not find such an association). In fact, FFR_{ENV} evoked

by amplitude-modulated tones presented in quiet even seems to predict sentence understanding

in noise in adults with TH (Mepani et al. 2021). Recent evidence suggests that FFR_{ENV} (evoked

by speech in quiet) correlates with speech perception in noise in unaided (but not in aided) 4 to

9 year-olds with congenital mild to moderate HL (van Hirtum et al. 2023). Whether subcortical

processing relates to speech perception in noise in older children/adolescents with congenital

HL remains an open question.

Cortical processing of sounds is also thought to relate to behavioural speech perception,

mostly in quiet. In 4-month old children with TH or HL, components of the LAER relate to

parents' reports of auditory functional performance (Ching et al. 2023). In adults, the MMN is

thought to relate to perceptual speech discrimination, at least at the group level (for a review,

see Näätänen 2001). However, no clear relationship has emerged between MMN amplitude and

speech discrimination in children with TH so far (Sharma et al. 2006; Jost et al. 2015; Shafer

et al. 2005). To our knowledge, it remains unclear whether cortical EEG measures relate to

speech perception in children/adolescents with congenital HL.

Study objectives

The main goal of this study was twofold: to evaluate (i) the effect of childhood HL and

(ii) the benefit of amplification on both subcortical and cortical auditory processing. (i) So far,

no study has investigated the effect of childhood MM HL on subcortical processing presented

at fixed intensity levels across age-matched groups. Additionally, results from (older) adult

studies have led to contradictory results. Therefore, this study will be a first investigation of the

effect of MM HL on the processing of unamplified sounds throughout development. At the

cortical level, we predict smaller/later LAER and MMN responses in children/adolescents with

HL than those with TH, with a possible interaction between age and group on the MMN. (ii)

Regarding the second objective, we expect larger FFR_{ENV} in children with HL when sounds are

amplified than unamplified. At the cortical level, results from the literature are mixed, but have

never compared LAER/MMN with and without amplification in the same children with HL.

We will thus study the effect of amplification on the cortical responses evoked in

children/adolescents with HL.

Given that subcortical and cortical processing have both been related to speech

perception, an exploratory objective was (iii) to investigate how these neurophysiological

responses relate to speech intelligibility in quiet and in noise.

Methods

Participants

Thirty-five children aged 8 to 16 years were included in the study. Nineteen (8 female) had a diagnosis of mild-to-moderate SNHL (HL group) and sixteen (8 female) were age-matched children with typical hearing (TH group). All participants were monolingual British English natives and obtained a nonverbal IQ (NVIQ) score of at least 85 (Block Design subtest of the Weschler Abbreviated Scale of Intelligence). No child had any known medical, psychological, neurological or developmental disorders other than SNHL. Individual air-conduction audiometric thresholds were measured bilaterally in all children at octave frequencies from 0.125 to 8 kHz using an Interacoustics AC33 audiometer. The study was conducted with the verbal assent of the participants, the written consent of their parents, and was approved by the UCL Research Ethics Committee (2109/004). It was performed in accordance with the Declaration of Helsinki.

Figure 1: Pure-tone air-conduction audiometric thresholds for children with HL (orange) and children with TH (blue and white). Audiometric thresholds are shown across octave frequencies from 0.25 to 8 kHz in the left and right ears. Individual thresholds for the HL group are shown as lines, and the group mean is shown as a bold line. Mean thresholds for the TH group are marked in white, with the shaded blue area representing the range for the TH group.

Participants in the HL group had better-ear pure tone average (BEPTA) thresholds of

21-70 dB HL across 0.25-8 kHz (see Figure 1), with on average 7 dB difference in PTA between

the ears. All children in the HL group were spoken language users (i.e., non-signers), and

sixteen had prescription hearing aids, but these were not used in this study. Children in the HL

group were individually matched in age (± 6 months) to those in the TH group, and mean age

did not differ significantly between the two groups (see Table 1). None of the children in the

TH group had a known history of hearing loss (including otitis media with effusion),

educational difficulties, or speech and language problems (based on parent/guardian report).

All had mean PTA thresholds ≤ 20 dB HL across octave frequencies 0.25-8 kHz in both ears

and obtained thresholds no higher than 20 dB HL at any given frequency.

Group comparisons are presented in Table 1. Note that some children did not complete

all behavioural and electrophysiological tasks. With respect to the behavioural tasks, two

children (1 TH and 1 HL) did not perform the speech identification in noise tasks. Due to

technical difficulties, two children (1 TH and 1 HL) did not perform the speech identification

in a fluctuating noise task. Two children with HL did not perform the speech discrimination in

quiet task. With respect to the EEG measurement, one child (HL) was only tested in the

amplified, but not the unamplified EEG condition. The sampling frequency of the EEG

recording was accidentally set to 2048 Hz for another child (HL). For this child, we could obtain

the cortical measures, but not the subcortical measures. In all these cases, results were treated

as missing data.

274 **Table 1** : Mean (SD) participant characteristics for the HL and TH groups. Comparisons are
275 independent-samples *t*-tests. Effect sizes are Cohen's *d*. Significant effects are shown in boldface.

Variable	TH (n = 15)	HL (n = 19)	*t*(df)	*p*-val	Effect size	95 % CI
Age (years)	12.6 (2.43)	12.4 (2.44)	0.26 (32.1)	.795	0.08	[-1.46, 1.89]
Nonverbal IQ	119.1 (12.6)	102.5 (12.5)	3.83 (31.44)	**<.001**	1.32	[-25.4, -7.7]
Maternal education	20.9 (2.59)	18.3 (2.78)	2.76 (30.99)	**.009**	0.96	[-4.49, -0.67]
BEPTA threshold (dB HL)	3.7 (2.87)	45.0 (13.30)	-13.18 (19.97)	**<.001**	4.12	[-47.86, -34.78]
Age of HL detection (years)	-	3.6 (3.03)				

276
277

Data acquisition and pre-processing

Stimuli were digitised /bɑ/ (standard) and /dɑ/ (deviant) syllables (see Figure 2)

originally spoken by a native British female speaker, and identical to those used in Halliday et

al., 2019; Calcus et al., 2019; Bishop et al., 2011. The intonation contours were made monotone

at 220 Hz using Praat (Boersma & Weenink, 2001) and the final stimuli were RMS equalised

with GoldWave (Craig, 2008). Vowel formant frequencies were approximately 800, 1340 and

2700 Hz for formants F_1, F_2 and F_3 respectively, although there were small differences between

the two syllables as the stimuli were based on natural utterances. Both syllables had rising F_1s

over ~50 ms with a much larger transition for the /dɑ/ (~230 Hz) than the /bɑ/ (~115 Hz). The

phonetic contrast was cued primarily by differences in the properties of the initial release burst

(more intense for the /dɑ/, especially at higher frequencies) and on the formant transitions into

the vowel. No F_2 transition was present for the /bɑ/ but the /dɑ/ had a significant falling

transition for F_2 of ~350 Hz.

Each condition consisted of a total of 1320, 175-ms stimuli, presented in a passive

oddball paradigm (deviant probability: 10%). Half of all stimuli were presented in positive

polarity and the other half in negative polarity. The order of presentation of each polarity was

randomized. The inter-stimulus interval was jittered randomly between 275 and 375 ms to limit

neural adaptation (Malmierca et al., 2014).

Figure 2: Top row: Spectrograms of the standard and deviant speech syllables in their unamplified form. Bottom
row: Spectra of the vocalic portion (20 to 170 ms) of the standard syllable /bɑ/ in both unamplified and NAL-R
amplified versions. Children with TH were only presented with unamplified versions of the stimuli. Children with
HL were presented with both versions of the stimuli, with the amplification being dependent upon each
individual's audiogram. For visualisation purposes, the amplified stimulus here was generated based on the
average audiogram of the HL group. It can clearly be seen that, as expected, the level of the harmonics below ~2
kHz is unchanged because there is minimal hearing loss at those frequencies, with amplification of the higher
harmonics in the frequency region in which there is substantial HL.

Children with TH were presented with only one condition, consisting of unamplified

stimuli that were presented binaurally at 70 dB SPL through shielded insert earphones (ER-2,

Etymotic Research) – henceforth referred to as children with TH_U. Children with HL undertook

two conditions - unamplified and amplified – presented in counter-balanced order (henceforth:

HL_U and HL_A). Amplification used the National Acoustic Laboratory-Revised algorithm

(NAL-R, Byrne & Dillon, 1986) to provide a frequency-specific gain (without compression),

based on each individual child's audiogram. This strategy minimizes the risk of potential

confounds related to sound-field testing as well as potential inter-individual variability in the

hearing aid processors of individual children (Billings, 2013). Stimuli in the amplified condition

were limited in intensity such that they did not exceed 85 dB SPL. Each condition lasted

approximately 20 minutes, with three short breaks offered every 5 minutes. During the

recording, participants were seated comfortably in a shielded, sound-attenuated booth, and

watched a silent movie.

EEG was recorded using a BioSemi ActiveTwo system at a sampling rate of 8192 Hz,

from 32 scalp electrodes in the standard 10/20 configuration (Jasper, 1958). Additional

electrodes were placed on each mastoid and recordings were re-referenced offline to the average

of activity at the mastoid electrodes. Electrode offsets were consistently < 50 mV.

Subcortical response analysis

Subcortical response analyses were performed only on the standard (/bɑ/), to avoid the

confound of small spectral differences between the standard and the deviant syllables. 1188

epochs (standard probability: 90%) were obtained by applying a band-pass filter from 90 to

1600 Hz (zero-phase, finite impulse response, -6dB/octave) to the EEG data evoked by the

standard syllables (/bɑ/) at Cz, and extracting data 25-175 ms relative to trigger onset, which

covers the duration of the steady portion of the vowel. Trials exceeding ± 100 µV were rejected

before averaging. There was no significant difference in the proportion of trials that were

rejected across groups [TH_U: 18.5%; HL_U: 19.1%; HL_A: 20.3%; $F(2, 47) = 0.723$, $p = 0.491$, η^2

$= 0.03$]. The averaged responses to positive and negative polarities were then added together to

enhance the envelope of the response (FFR_{ENV}), and subtracted from each other (i.e. positive

minus negative polarity) to enhance the TFS of the response (FFR_{TFS}). Spectral amplitudes were

calculated using a fast Fourier transform (FFT), after applying zero-padding to the sampling

rate to obtain an FFT with a resolution of 1 Hz. Spectral peaks at F0 (220 Hz) and harmonics

(H2: 440 Hz; H3: 660 Hz; H4: 880 Hz; H5: 1120 Hz; and H6: 1320 Hz) were computed as the

maximum amplitude within an 8-Hz bin centred on F0 and subsequent harmonics, following

visual inspection of the grand average subcortical response, and in line with previous studies

(e.g. Schoof & Rosen, 2016). Neural background noise amplitudes were calculated by taking

the mean amplitude of the fast Fourier transform across 100-Hz windows surrounding each

spectral peak (50 Hz on each side, excluding the ±70 Hz adjacent bins).

Cortical response analysis

Evoked potentials considered to be cortical in origin were obtained by band-pass

filtering the EEG recorded at electrode Fz between 0.5 Hz and 35 Hz (zero-phase, finite impulse

response, -6dB/octave) and creating epochs from -100 ms to 500 ms relative to each stimulus

onset time. 1320 epochs (1188 standards and 132 deviants) were baseline corrected using the

mean value from -100 to 0 ms. Trials exceeding ± 100 µV were rejected before averaging.

There was no significant difference in the proportion of trials that were rejected across groups

[TH_U: 12.7%; HL_U: 13.4%; HL_A: 15.75%; $F(2, 50) = 0.43$, $p = 0.654$, $\eta^2 = 0.01$].

P1 and N2 grand average latency were quantified by means of the half-area latency

measure of the standard waveform recorded in the TH group (Zhang & Luck, 2023), a

computation that is less sensitive to high-frequency noise than traditional latency peak measures

(Finke et al., 2016; Luck, 2014). The window for computation of the area under the curve of

the P1 ranged from 55 to 135 ms and from 250 to 350 ms for the N2, based on visual inspection

of the TH grand average. P1 and N2 amplitudes were then computed for each participant and

condition as the mean amplitude within a 40-ms window centred around the half-area latency

(P1: 96 ms post-stimulus onset; N2: 329 ms post-stimulus onset). P1 and N2 latencies were

computed for each participant and condition as the half-area latency for the local peak to appear,

within the 40-ms window centred around the TH grand average evoked by standards. Note that

N1 and P2 were not analysed because they could not be reliably identified in the younger

participants, which is likely due to a protracted maturation of these responses until adolescence

(Bishop et al., 2011; Sussman et al., 2008).

The MMN was computed as the differential wave obtained by subtracting the neural

response evoked by the standard to that evoked by the deviant, for each participant and

condition. The MMN was then statistically assessed in two ways. First, to determine whether

there was evidence for neural discrimination of speech deviants, the presence of an MMN was

evaluated for the three combinations of group and amplification conditions. Point-to-point

comparison of the differential wave amplitudes was performed to determine the latency period

over which the waveforms were significantly smaller than zero, if any. One-sided *t*-tests were

computed within the 100-500 ms post-stimulus-onset time window which was identified as the

region most likely to contain the MMN by visual inspection of the grand average waveforms.

Because adjacent points in the waveform are highly correlated, potentially leading to spurious

significant values in short intervals, an MMN was considered present when $p < .01$ (one-tailed)

for more than 20 ms at adjacent time-points (Kraus et al., 1993; McGee et al., 1997). Half-area

MMN latency was computed on the differential waveform recorded in the TH group. MMN

amplitude for each participant and condition were then computed as the mean amplitude in a

100-ms window centred around the grand averaged differential waveform half-area latency

peak observed in children with TH (at 426 ms post-stimulus onset; Zhang & Luck, 2023). MMN

individual latencies were computed as the half-area latency within a 100-ms window centred

around the grand average differential waveform of the children with TH.

Speech discrimination in quiet

The speech discrimination in quiet task was the same as that used in previous studies

(Halliday et al., 2017; 2019). The two endpoints of the continuum were based on the same

digitised /bɑ/ and /dɑ/ used in the electrophysiological recordings. A continuum of 100 stimuli

was then constructed (including the endpoints), using the morphing capabilities of the

programme STRAIGHT (Kawahara et al., 1999). The task was delivered via a child-friendly

computer game, incorporating an adaptive, three-interval, three-alternative, forced-choice

paradigm. Stimuli were presented binaurally via headphones (Sennheiser 25 HD) at a fixed

intensity (70 dB SPL), on a tablet, meaning identical levels presented in the two groups.

Children were required to detect the odd-one-out, where two of the intervals corresponded to

the /bɑ/ end of the continuum, and the third, target stimulus fell somewhere along the rest of

the continuum. Participants received visual feedback regarding the accuracy of their responses.

A three-down, one-up procedure was used to select, on each trial, the appropriate target sound

from a continuum, tracking a performance level of 79.4% correct (Levitt, 1971). This was

preceded by an initial one-down, one-up procedure until the first reversal (Baker & Rosen,

2001). The initial step size was 15 stimulus places along the continuum, which reduced to five

after the first reversal. The threshold was the arithmetic mean of the last four reversals in

direction of the adaptive track, expressed as the stimulus number (0-99) of the target stimulus

along the continuum. Therefore, smaller numbers indicate better performance. The task was

preceded by five practice trials which contained standard-target difference that had previously

been deemed suprathreshold for adults with TH (Halliday et al. 2019).

Speech perception in noise

The rationale for choosing the stimuli for this task was to provide the most direct

comparison to the neurophysiological stimuli, which were speech syllables. Another advantage

of syllables is that they minimise the role of linguistic and cognitive factors on auditory

performance, allowing to focus more on lower level auditory properties. Speech reception

thresholds (SRTs) were thus measured adaptively for 13 consonants presented in vowel-

consonant-vowel (VCV) logatomes, e.g. /imi/, /apa/. Vowels were /i/, /ɑ/, and /u/, and

consonants were /p/, /k/, /b/, /d/, /g/, /f/, /m/, /n/, /l/, /v/, /w/, /y/, and /z/. Both the vowels and

consonants varied from trial to trial, but only the response for the consonant was scored. Stimuli

were recorded by a native British English female speaker in an anechoic chamber and were

digitized via a 16-bit analogue-to-digital converter at a 44.1-kHz sampling frequency. The

logatomes were presented together with two types of background noise: steady (i.e.

unmodulated, but see Stone & Moore, 2014) or fluctuating in amplitude sinusoidally at 8 Hz,

which corresponds to a syllabic modulation rate (Greenberg et al., 2003). The modulation depth

was fixed at 1.0 and the starting phase was randomized between 0° and 360° on each trial. The

noise was shaped on-and-off using a raised-cosine with 50-ms rise-fall times.

Participants were seated in a quiet room and the stimuli were presented binaurally over

Etymotics ER-2 earphones at an overall level of 70 dB SPL, again meaning identical levels

presented in the two groups. They were asked to repeat the logatomes as best as they could.

The experimenter scored the participants' responses using a graphical interface which showed

the 13 possible consonants. No feedback was provided.

The expected wide variability in intelligibility across consonants at a given SNR would

not allow their reliable use in a simple adaptive procedure. Therefore, SRTs for each

combination of consonant and vowel were estimated from a prior study of young adults with

TH (unpublished), allowing the calculation of adjustment factors in SNR to give all stimuli the

same nominal SRT (Plomp & Mimpen, 1979). These factors were then used during the adaptive

procedure to adjust the presented SNR. The signal-to-noise ratio (SNR) was varied adaptively

by varying the level of the target and fixing the level of the noise at 70 dB SPL. The first

logatome was presented at a nominal SNR of -10 dB (taking into account that token's

adjustment factor). Depending on the participant's response (correct/incorrect), the SNR of the

second logatome was decreased/increased by 10 dB. Again, depending on the participant's

response (correct/incorrect), the SNR of the third logatome was decreased/increased by 6 dB.

For each subsequent logatome, the SNR increased/decreased by 2 dB for incorrect/correct

responses, respectively. Measurement stopped after either 12 reversals or 78 trials, whichever

occurred first. The SRT was computed by taking the mean SNR (dB) across all track reversals.

Speech perception in quiet

As a baseline and familiarisation for the speech perception in noise task, we also

included a speech perception in quiet task. Percentage correct responses were measured for 78

consonants presented at a fixed intensity of 70 dB SPL, using the same VCV logatomes and

procedure for recording the participants' responses as those used in the speech perception in

noise task.

Statistical analyses

Regarding the subcortical data, we first ran an outlier detection algorithm. Data points

that fell outside the mean $\pm$ 3 *SD* (within each combination of group, amplification, FFR

component and harmonic) were considered outliers and excluded from further analyses. In total,

2.7% of data points were excluded from the analyses, most of them from the HL group. After

excluding outliers, the subcortical data were normally distributed (Kolmogorov-Smirnov test:

all ps > .05). Note that for both FFR_{ENV} and FFR_{TFS}, there were significant differences in

variance between groups as evidenced by Levene's test for equality of variance (both ps <

0.001).

Second, paired two-tailed *t*-tests with Bonferroni-Holm correction were conducted

separately for each group (TH_U, HL_U, HL_A), FFR component (FFR_{ENV} and FFR_{TFS}) and

harmonic (F0 to H6) to determine whether the FFR responses were significantly above the

neural noise floor. FFR responses are considered present if at least one of the three groups

showed a significant difference between target frequency peak and spectral noise floor in each

component, at each harmonic. For FFR_{ENV}, peaks were significantly larger than the noise floor

only at F0 and H2 (all ps < 0.001). For FFR_{TFS}, peaks were significantly larger than the noise

floor at H3, H4, H5 and H6 (all ps < 0.05).

Third, a linear mixed model was used to determine whether amplitude of the spectral

noise floor differed across groups (TH_U, HL_U, HL_A), FFR component (FFR_{ENV} and FFR_{TFS}) or

harmonic (F0 to H6). As expected, the only significant predictor was harmonic ($p < 0.001$). In

the absence of a significant group effect on the baseline ($p > 0.10$), subsequent analyses were

performed on the target frequency peak amplitude.

Regarding the cortical data, we found no outliers (± 3 SD from the mean) in the

amplitude of either P1 or MMN. The cortical data were normally distributed (Kolmogorov-

Smirnov test: all $ps > .05$). For both P1 and MMN amplitudes, there were no significant

differences in variances between groups (Levene's test for equality of variance: both $ps > .05$).

Linear mixed models (lmerTest package of R[92]) were used to evaluate group differences

in both cortical (P1, N2 and MMN) and subcortical (FFR_{ENV} and FFR_{TFS}) measures. Fixed

effect predictors and all their interactions (all orders) were included in the initial model. In all

models, the factor listener was used as a random intercept, and random slopes for all the

appropriate predictors (harmonic and amplification) that did not lead to convergence issues or

inestimable models. Backward stepwise reduction was then used to simplify the models based

on F-tests, eliminating insignificant terms, but always keeping lower-order terms that were

significant in higher-order interactions. This was done first for the random effects, and then the

fixed effects. Only significant results are reported below. Analyses of group differences aimed

to answer three main questions.

-　First, is there a significant group difference in electrophysiological responses to

unamplified speech stimuli? Here, the predictors were group (TH_U vs. HL_U) and age

(continuous variable), with harmonic as an additional predictor (FFR_{ENV}: F0 and

H2; FFR_{TFS}: H3 to H6) to model the subcortical data.

-　Second, for children with HL only, is there a benefit of amplification on the neural

encoding of sounds? Here, the predictors were amplification (HL_U vs. HL_A),

BEPTA and age, again with harmonic as an additional predictor for the subcortical

data.

- Third, does sound amplification in children with HL lead to electrophysiological

responses that are similar to the control group (TH_U vs. HL_A)? Again, the predictors

were group (TH_U vs. HL_U) and age, with harmonic as an additional predictor for the

subcortical data.

With respect to the speech intelligibility measure in quiet, whose outcome variables was

binomial, generalized linear mixed models (logistic regressions) were fitted to the data. Group

(TH, HL) and age, as well as their two-way interaction were included in the model as between-

subjects factors, with both listener and consonant included as random effects. For speech

perception in noise, a linear mixed model was computed with group (TH, HL), noise type

(steady, fluctuating), age, and all their interactions as fixed factors, and listener as a random

effect. Backward stepwise reduction was then applied, as for the EEG measurements.

Finally, the relationship between subcortical and cortical measures, as well as that

between electrophysiological and behavioural measures were investigated using correlations,

conducted separately on the TH and HL groups, and separately for unamplified and amplified

conditions in the latter group.

Table 2: Summary of the main effects of hearing loss (HL_U versus TH_U) and amplification
(HL_A versus HL_U) for subcortical, cortical and behavioural measures. * $p < 0.05$, ** $p < 0.01$,
*** $p < 0.001$.

		Effect of hearing loss (HL_U versus TH_U)	**Benefit of amplification** (HL_U versus HL_A)	**Benefit of amplification** (HL_A versus TH_U)
Subcortical measures				
FFR_{ENV}	Amplitude	$HL_U < TH_U$ ***	$HL_U < HL_A$*	
FFR_{TFS}	Amplitude	$HL_U < TH_U$ *	Amplification × Harmonic ***	Amplification × Harmonic ***
Cortical measures				
P1	Amplitude	$HL_U < TH_U$ **	Amplification × BEPTA *	
	Latency	TH_U earlier than HL_U *	HL_A earlier than HL_U*	
N2	Amplitude	$HL_U < TH_U$ ***		$HL_A < TH_U$ **
	Latency		Amplification × BEPTA *	
MMN	Amplitude	$HL_U < TH_U$ **		$HL_A < TH_U$ **
	Latency			

Results

A summary of the main effects of hearing loss (HL_U versus TH_U) and of amplification

(HL_A versus HL_U) is provided in Table 2, for subcortical, cortical and behavioural measures.

1. Subcortical responses

Figure 3 shows the grand-average time waveforms and spectra of the subcortical responses

evoked at electrode Cz by unamplified speech for children with TH (left panels), and

unamplified and amplified speech for children with HL (middle and right panels, respectively).

Grand-averages were derived from the time waveforms and spectra of the individual

waveforms. Scatterplots of the amplitude of the various components are shown as a function of

age in Figure 4.

First, group differences in the subcortical processing of unamplified speech were examined

(TH_U vs. HL_U; see Supplementary Table 1). FFR_{ENV} were significantly smaller (by 0.082 μV)

in children with HL_U than in children with TH_U (p = .001) for both harmonics, with no effect

of age. Similarly, FFR_{TFS} were significantly smaller (by 0.028 μV) in children with HL_U than

in children with TH_U (p = .015). Note that in neither analysis was there a significant group x

harmonic interaction.

Next, the effects of amplification were examined by comparing the subcortical responses

of children with HL presented with unamplified versus amplified speech stimuli (HL_U vs. HL_A;

see Supplementary Table 2). Overall, FFR_{ENV} were significantly larger (by 0.118 μV) in

response to amplified compared to unamplified speech sounds (p = 0.011). The amplification

× age interaction got excluded during model reduction (p = 0.051), yet we observed a

marginally significant decreasing benefit of amplification with age on FFR_{ENV} amplitude (p =

0.019). With respect to the FFR_{TFS}, we observed a significant amplification × harmonic

534 interaction. FFR$_{TFS}$ were significantly larger in response to amplified than unamplified speech

535 sounds at H4, H5 and H6, but not at H3.

Figure 3: Grand average of the subcortical responses at Cz. The upper row represents the FFR$_{ENV}$ in the time
domain. The middle and lower rows represent the FFR$_{ENV}$ and the FFR$_{TFS}$ evoked by the standard sound /bɑ/,
presented either unamplified to children with TH (70 dB SPL, left panel), unamplified to children with HL (70 dB
SPL, middle column), or amplified to children with HL (with a frequency-specific gain tailored to each child's
hearing thresholds, right panel). The grand average spectra were computed by averaging the individual amplitude
spectra.

545 Last, we compared the amplitude of the responses obtained when amplified sounds were

546 presented to children with HL versus unamplified sounds presented to children with TH (HL$_A$

547 vs. TH$_U$; see Supplementary Table 3). The amplitude of the FFR$_{ENV}$ did not significantly differ

548 between children with HL$_A$ and those with TH$_U$ ($p = 0.168$). With respect to the FFR$_{TFS}$, we

549 observed a group × harmonic interaction ($p < 0.001$). FFR$_{TFS}$ were significantly larger in

550 children with HL$_A$ than those with TH$_U$ at both H4 and H6, but not at H3 and H5.

Figure 4: Scatterplots of the amplitude (µV) of the FFR$_{ENV}$ (F0 and H2) and FFR$_{TFS}$ (H3 to H6) over age, for each group (TH$_U$, HL$_U$ and HL$_A$). Regression lines are fit on the basis of each combination of group, FFR component and harmonic.

2. Cortical responses

Figure 5: Grand average waveforms at Fz, evoked by the standard (lighter line) and deviant (darker line) presented either unamplified to both children with TH$_U$ (70 dB SPL, left column) and children with HL$_U$ (70 dB SPL, middle column) or amplified to children with HL$_A$ (right column). The thin, gray line represents the differential wave between standards and deviants (MMN). Voltage maps show the mean MMN activity during the 100-500 ms post-stimulus time window. Negative values of the MMN are shown in blue, and positive values in red. The horizontal line represents the duration of a significant MMN in the period of the 100-500 ms post-stimulus-onset epoch evoked in children with TH. No significant MMNs were observed in either the unamplified or amplified conditions for the children with HL as a group.

Figure 5 shows the grand-average cortical waveforms evoked by standard and deviant

stimuli at electrode Fz for both groups. In children with TH, the response to the standard sound

was dominated by a large positivity occurring around 85 ms post-stimulus onset (the P1),

followed by a large and prolonged negativity peaking around 300 ms (the N2). As we and others

have reported previously, this pattern of results is typical of children with TH in this age range

(Bishop et al., 2011; Calcus et al., 2019). For children with HL, the overall pattern of responses

to the standard and deviant sounds was similar, albeit reduced in amplitude compared to

children with TH. Note that there is no reliable evidence of an MMN in children with HL in

either unamplified or amplified conditions, but strong evidence for one in children with TH

(illustrated by a horizontal line covering the range from 330 to 500 ms in the left panel of Figure

5).

First, we assessed group differences in the cortical processing of unamplified speech

(TH$_U$ vs. HL$_U$; see Supplementary Table 4). As shown in Figure 6, P1 was smaller (by 2.05 µV)

and later (by 2.79 ms) in children with HL$_U$ than those with TH$_U$ (both $ps < 0.05$). N2 was also

smaller (by 3.79 µV; $p < 0.001$) but not later in children with HL$_U$ than those with TH$_U$.

Additionally, children with HL$_U$ had significantly smaller MMN responses (by 3.47 µV) than

children with TH$_U$ ($p = 0.003$).

Next, we sought to determine whether amplification benefited the cortical processing

of speech sounds in children with HL (HL$_U$ vs. HL$_A$; see Supplementary Table 5). For P1

amplitude, we found a significant amplification × BEPTA interaction ($p = 0.028$; see

Supplementary Figure 1). Perhaps not surprisingly, the P1 amplitude in children with better

BEPTA thresholds was not changed by the relatively small amplification they received,

whereas the P1 amplitude for children with poorer BEPTA thresholds (and who received greater

degrees of amplification) was increased so as to make P1 amplitude more-or-less independent

of BEPTA. P1 latency was longer in the unamplified than the amplified condition ($p = 0.027$).

It was also longer for individuals with higher BEPTA thresholds ($p = 0.024$) and for younger

children ($p = 0.005$), irrespective of the condition for both. For N2 latency, there was a

significant main effect of amplification ($p = 0.033$). Even though the amplification × BEPTA

interaction was also significant ($p = 0.030$), post-hoc contrasts were not significant ($p = 0.842$).

There was no significant effect of amplification on MMN amplitude or latency (both ps > 0.10).

Note that irrespective of the condition, MMN amplitude decreased with age in children with

HL ($p = 0.040$).

Supplementary Figure 1: Amplitude (µV) of the P1 for the HL group as a function of BEPTA. Individual (shapes)
and group (lines) data are shown for both unamplified (HL$_U$, plain lines) and amplified (HL$_A$, dotted lines) raw.
Shaded lines represent the 95% confidence intervals.
604

Last, we evaluated the potential benefit of amplification for cortical processing of

speech in children with HL to that of children with TH (HL$_A$ vs. TH$_U$; see Supplementary Table

6). There was no significant effect of group on P1 amplitude or latency (both ps > 0.10). Yet

there remained a significant group difference when looking at N2. Despite amplification, N2

amplitude was smaller (by 2.43 µV; $p = 0.004$), but not later, in the group with HL$_A$ than the

group with TH$_U$. Similarly, children with HL$_A$ showed significantly smaller MMN than the

group with TH$_U$ ($p = 0.002$). Note that, although the group × age interaction did not come out

significant (p = 0.146), post-hoc comparisons indicate a significant decrease of MMN

amplitude with age in children with HL$_A$ (p = 0.038) but not in children with TH$_U$ (p > 0.10).

Figure 6: Amplitude (µV; left column) and latency (ms; right column) of the P1, N2 and MMN for each group
and condition. Individual (shapes) and group (colour) data are shown for both unamplified (HL$_U$, orange) and
amplified (HL$_A$, red) data. Shaded lines represent the 95% confidence intervals. Latencies (right column) are
illustrated as group boxplots because there was no effect of age.

620
621
622
3. Behavioural data

3. 1. Speech discrimination in quiet

Figure 7 shows the behavioural performance of the two groups on the speech perception

tasks. For speech discrimination of the speech sounds /bɑ/ vs /dɑ/ in quiet (Figure 7A), there

was no significant effect of group, age, or their interaction on response thresholds (all ps > .10;

see Supplementary Table 7A).

Figure 7: Boxplots of (A) stimulus distance (B) percentage correct consonant identification in quiet and (C) SRTs
for consonant identification in steady (light grey) and fluctuating (dark grey) background noises. Children with
TH are shown in blue, HL children are shown in orange. The boxes show the interquartile range (25^{th}-75^{th}
percentile) of the data with horizontal line indicating the median. The whiskers indicate values that fall within 1.5
times the interquartile range.

3. 2. Speech identification in quiet and noise

Figure 7B presents the percentage correct consonant identification in quiet for the two

groups (also see Supplementary Table 7B). Children with HL were significantly poorer than

children with TH ($p < 0.001$). Note that for both groups, performance significantly improved

with age ($p = .003$), although ceiling effects meant only a very small increase in percentage

correct for the TH group.

Finally, Figure 7C presents the SRTs in steady and fluctuating background noise for the

two groups (also see Supplementary Table 7C). The significant main effects of group and noise

type (both $ps < .001$) need to be considered in light of the significant group × noise type

interaction ($p = .006$). As expected, both groups experienced masking release when they were

presented with fluctuating noise (both $ps < 0.001$). However, the magnitude of this effect was

larger in children with TH (8.5 dB SNR) than in children with HL (3.9 dB SNR).

4.   Relationship between the neural and behavioural responses

The relationship between electrophysiological activity and behavioural responses was

investigated by means of a set of correlations. Holm-Bonferroni correction was applied to

correct for multiple comparisons where appropriate.

Given the evidence that perception of speech in quiet is dominated by the encoding of

the envelope of the signal, we investigated the relationship between the amplitude of the

FFR_{ENV} at both F0 and H2 and behavioural performance of speech identification in quiet,

separately for children with TH_U and children with HL_U. None of the correlations were

significant (all ps > 0.05).

Adequate perception of speech in noise requires precise spectro-temporal processing of

speech, particularly in the dips of a fluctuating background noise. In fact, the amplitude of

speech-evoked FFR in quiet relates to the perception of speech in noise (see introduction).

Therefore, we investigated the relationship between amplitude of the speech-evoked FFR_{TFS} at

H4 and H6 (in quiet) and behavioural speech perception in steady and fluctuating noise, as well

as masking release, separately for children with TH and children with HL (in both unamplified

and amplified conditions). None of the correlations were significant (all ps > 0.05).

The speech stimuli used for the electrophysiological recordings were the endpoints /bɑ/

and /dɑ/ of the continuum used for the speech discrimination in quiet behavioural task. As such,

we hypothesised that a relationship might be observed between the amplitude of the MMN and

the stimulus distance threshold measured in the speech discrimination in quiet behavioural task.

However, there was no significant correlation between speech discrimination in quiet and

MMN amplitude, as evaluated separately for both groups, and separately for the amplified and

unamplified conditions in children with HL (all ps > 0.05).

Last, we investigated the relationship between subcortical and cortical responses,

separately for both groups (and separately for HL_A and HL_U). In children with TH, larger

676 FFR$_{ENV}$ at F0 were associated with smaller N2 responses ($p = 0.004$). There was no other

677 significant correlation between FFR$_{ENV}$ at F0 and P1, N2 or MMN (all ps > 0.05).

679 **Discussion**

680   This study investigated the effect of childhood MM HL and the benefit of amplification

681 on both subcortical and cortical processing of speech sounds. When sounds were presented at

682 the same intensity across groups, both subcortical and cortical responses were smaller in

683 children with HL than those with TH. When sounds were amplified, the subcortical responses

684 increased in children with HL$_A$ compared to HL$_U$. In fact, they could even be larger in children

685 with HL$_A$ than TH$_U$ (although only significantly so for the FFR$_{TFS}$). At the cortical level, an

686 amplification benefit was also observed on P1 amplitude but not on the amplitude of later

687 responses (N2 or MMN). In fact, at the group level, children with HL did not show a significant

688 MMN (unlike children with TH), whether the sounds were amplified or not. When looking at

689 individual responses, the amplitude of the MMN tended to decrease with age in children with

690 HL but not in those with TH. Behaviourally, children with HL had poorer speech intelligibility

691 than those with TH both in quiet and in noise, and they showed less benefit from background

692 noise fluctuations. Yet, the relationship between the neurophysiological responses and the

693 behavioural performance remains unclear.

694   *Subcortical processing*

695   When presented with unamplified sounds, children with MM HL showed significantly

696 smaller FFR$_{ENV}$ and FFR$_{TFS}$ than those with TH. This finding replicates and extends previous

697 results from older adults with age-related HL (Plyler & Ananthanarayan 2001;

698 Ananthakrishnan et al. 2016) – but contrasts with some studies that did not find group

699 differences on unamplified FFR measures (Anderson et al. 2013; Presacco et al. 2019). The

700 etiology of the HL might thus modulate its effect on subcortical processing of unamplified

sounds. When presented with amplified sounds, children with MM HL showed a clear benefit

from the frequency-specific gain applied to the sounds. In fact, the amplified FFR_{ENV} of

children with MM HL did not significantly differ from that of children with TH_U. Furthermore,

the amplified FFR_{TFS} was even larger than that of children with TH_U. Our findings thus fit with

existing results from the adult and child literature to indicate a clear benefit of amplification at

the subcortical level (Anderson et al., 2013; BinKhamis et al., 2019; Easwar et al., 2015, 2023,

2024; Jenkins et al., 1993).

We were initially surprised by the spectacular increase in amplitude of the FFR_{TFS} in

the higher harmonics, especially H6, following amplification. However, it is crucial to keep in

mind that the components reflected in this spectrum are not directly present on the scalp-

recorded EEG, but arise through a subtraction of the waves of the two stimulating polarities.

Consider Supplementary Figure 2, which depicts the spectra of the positive and negative

polarity stimuli separately. For children with HL_A, the spectra do indeed show a large

component at H6 (the largest component in the FFR_{TFS}), but one which is roughly the size of

the one at F0. This is not such a different pattern from that shown by children with TH_U,

although the F0 is somewhat larger than H6 in this latter case. The large differences between

these two groups in the FFR_{TFS} spectra arise not only from the component spectral amplitudes,

but also because the *phases* of the original components change over frequency. F0 and H2 have

very similar phases in the negative and positive polarities (thus leading to their enhancement in

the FFR_{ENV} spectra and cancellation in the FFR_{TFS} spectra), whereas, H3-H6 have phases that

are close to 180° out of phase with one another (thus leading to their cancellation in the FFR_{ENV}

spectra and enhancement in the FFR_{TFS} spectra). In that respect, the size of the higher frequency

components is roughly doubled in the FFR_{TFS} spectra, as compared to their actual measured

sizes in the individual polarity waves.

We can also exclude the possibility of electrical artefacts from the stimuli by noting that

the latency of the FFR_{TFS} and FFR_{ENV} responses for all groups was very similar, even though

any artefacts would be cancelled in the FFR_{ENV} responses. Note too that high amplitudes of

high harmonics have been seen before in FFR_{TFS} spectra. Krishan (2002), in particular, for a

similar /ɑ/ vowel to the one used here (but based on a male voice with F0 ≈ 125 Hz), found, at

the highest level tested, that the largest spectral components was at H7 (very close to the first

formant frequency).

We believe these high amplitudes of H4-H6 (at 880 - 1320 Hz) stem from applying the

individualized, frequency-specific NAL gains calculated from audiograms with sloping hearing

losses (leading to increasing gains for higher frequency harmonics), but also because the vowel

has a broad spectral peak in that region (see Fig 2) due to the F1 and F2 frequencies of 800 and

1340 Hz. To our knowledge, the only other study[47] that used the NAL formula analysed FFR_{TFS}

responses only up to ~700 Hz. In this frequency range, there was no significant difference in

the FFR_{TFS} of older adults with HL (in quiet, with or without amplification) and age-matched

adults with TH. In our study, FFR_{TFS} was only above the noise floor at frequencies at H3 (660

Hz) and above. In fact, our results show significantly larger FFR_{TFS} in children with HL_A

compared to both children with HL_U and children with TH_U at H4 and H6, which are close to

the first and second formant peaks of our stimulus. In adults with TH, FFR_{TFS} are most clearly

detected at harmonics that are close to formant peaks, up to 1500 Hz[24]. In short, especially large

FFR_{TFS} at H4 - H6 might therefore reflect the interaction between frequency-specific NAL

amplification and the formants of our speech stimuli. Yet a *larger* FFR_{TFS} in children with HL_A

than those with TH_U might indicate hyperactivity at the subcortical level following childhood

HL.

Hyperactivity has been widely reported as a consequence of SNHL (for a review, see

Herrmann & Butler, 2021), and is thought to compensate for reduced inputs from damaged

peripheral structures to central brain regions, to maintain sensation (Chambers et al., 2016).

Such compensation fits within the theoretical framework of homeostatic plasticity, which states

that neurons regulate their physiological properties to compensate for persistent changes in their

activity level (Turrigiano & Nelson, 2004). Homeostatic mechanisms are thought to operate

from synaptic to network levels of sensory systems. That FFR amplitude was larger in children

with HL than those with TH, but only when sounds were amplified suggests that compensatory

activity is intensity-dependent at the brainstem level. In mice, changes in auditory cortex

responsiveness following conductive HL were only observed in sounds above a certain

threshold, or in certain frequency ranges (Teichert et al., 2017). Future studies are warranted to

investigate the effect of audibility on homeostatic plasticity of subcortical processing.

Cortical processing

Despite the medium to large benefit of amplification on subcortical responses, there was

only a small benefit of amplification on cortical responses (P1), or no benefit at all (N2, MMN).

Individualized frequency-specific gain increased P1 amplitude (which reflects improved sound

detectability, Eggermont & Ponton 2002), but only for children with poorer BEPTA.

Irrespective of BEPTA, P1 was earlier in HL_A than HL_U. In fact, the P1 amplitude and latency

of HL_A did not significantly differ from that of children with TH_U. This is in line with studies

that show age-appropriate P1 when sounds are presented at similar sensation level to children

with HL and those with TH (Koravand et al. 2013; Martinez et al. 2013; Engström et al. 2021).

Despite preserved P1, our results indicate that childhood HL leads to smaller (but not

later) N2 even when sounds are amplified, in line with Koravand et al., 2013, but contrary to

Martinez et al., 2013 or Engström et al., 2021. Furthermore, we did not observe a significant

MMN at the group level on children with HL, even when sounds were amplified. At the

individual level, the MMN amplitude of children with HL_A remained smaller than those with

TH_U. Note however that there was no group difference in MMN latency. Our finding of

absent/smaller MMNs in 8 to 16 year-olds with HL contrast with those of age-appropriate

MMNs in 5-12 year-old children (Koravand et al., 2013; Engström et al., 2021). Part of the

explanation for this discrepancy might lie in the developmental effects of childhood HL on the

central auditory pathway (see below for a discussion of the developmental effects observed in

this study).

Why does the P1 amplitude benefit from amplification, but not the N2 or MMN? One

possible explanation might lie in the different generators of the P1, N2 and MMN. The P1

generator has been reliably located within the secondary auditory cortex (Godey et al., 2001;

Liegeois-Chauvel et al., 2004), whereas MMN generators likely include an extended

thalamocortical circuitry projecting to primary auditory cortex (Lakatos et al., 2020). Little is

known about the generators of the N2, but early intracranial recordings also seem to point

towards a subcortical (thalamic) source (Velasco et al., 1989). Although anatomically close, the

primary and secondary auditory cortices receive input from distinct portions of the medial

geniculate body. Whereas the lemniscal pathway connects the ventral medial geniculate body

to the primary auditory cortex, the nonlemniscal pathway connects the medial/dorsal geniculate

body to both primary and secondary auditory cortices (Webster, 1992). Given that our results

indicate smaller N2 and MMN even when children with HL were presented with amplified

sounds, these could indicate a specific alteration of the nonlemniscal pathway connecting the

medial portion of the geniculate body to the primary auditory cortex. Our cortical findings thus

contrast with the subcortical evidence of hyperactivity following childhood HL.

The effect of HL on the development of the central auditory pathway

In fact, our cortical findings also contrast with findings of hyperactivity in the auditory

cortex as a consequence of age-related HL. Indeed, studies have reported some degree of

enhancement of the cortical responses to sounds in older adults with age-related HL compared

to age-matched individuals with TH (Alain et al., 2014; Tremblay et al., 2003; Millman et al.,

2017). To our knowledge, none of the pediatric studies investigating cortical processing of (amplified or unamplified) sounds in children with HL found such indications of increased cortical responsiveness (Calcus et al., 2019; Engström et al., 2021; Ji et al., 2023; Martinez et al., 2013; Koravand et al., 2013; Rance et al., 2002; Glista et al., 2012; Uhler et al., 2023). In mice, the same population of neurons showed distinct mechanisms of homeostatic adaptation if visual deprivation occurred in juvenile or adult animals (Wen & Turrigiano, 2021). Such an age-dependent effect of homeostatic plasticity might also be observed in the central auditory pathway following auditory deprivation.

Noteworthily, these age-dependent effects might already be operating between 8 and 16 years of age. Indeed, we observed a significant decrease in the benefit of amplification with age on FFR_{ENV} amplitude. Additionally, the amplitude of the MMN decreased with age in children with HL_A (but not in those with TH_U) – although note that the group $\times$ age interaction was not significant. Both results seem to indicate that adolescence could be a period of heightened plasticity for functional changes following auditory deprivation. Future studies are warranted to investigate this claim in a larger sample size, ideally taking puberty into account, as it could act as a biological trigger to those alterations (Gracia-Tabuenca et al., 2021; Laube et al., 2020; Vijayakumar et al., 2021).

Multiple gain mechanisms within the auditory system

To our knowledge, only three studies so far have combined subcortical and cortical measures from the same individuals to investigate the effects of HL on auditory processing. Speech-evoked subcortical and cortical responses were recorded successively in older adults, with and without hearing aids. When sounds were presented at 65 dB SPL, hearing aid amplification lead to significantly larger subcortical, but not cortical responses (Jenkins et al, 2017). In gerbils with developmental conductive HL, age-appropriate subcortical, but not cortical responses were recorded to amplitude modulated sounds presented at similar sensation

level as in animals with TH (Yao & Sanes, 2018). More recently, Hutchison et al. (2023)

recorded subcortical and cortical neurophysiological responses in young adults with TH before,

throughout and after fitting them with earplugs. The results indicate an increase in the

subcortical response, but a reduction of cortical activity ipsilateral to the deprived ear. Together

with our observations, the literature suggests that multiple gain mechanisms might underlie

homeostatic plasticity throughout the auditory pathway (Munro et al., 2014). Neurons from

each level of the auditory pathway are thought to display idiosyncratic changes following HL

(for a review, see Sanes, 2013).

Relationship between speech processing & perception

An exploratory objective of our study was to investigate how changes in

neurophysiological responses following HL might relate to speech intelligibility in quiet and in

noise. To do so, we chose stimuli that were as similar as possible for both neurophysiological

and behavioural tasks. However, two methodological choices are worth mentioning here:

Unlike the neurophysiological task, stimuli were only presented unamplified in the behavioural

tasks. Additionally, syllables were only presented in quiet in the neurophysiological tasks; yet

speech perception was investigated in quiet and in noise.

Behaviourally, children/adolescents with HL have higher (i.e. poorer) thresholds in

noise than children/adolescents with TH (Lewis et al. 2015; Goldsworthy & Markle 2019).

Their thresholds improved in the presence of fluctuating compared to steady background noise,

but to a lesser extent than the group with TH. Similarly to adults with HL (Lorenzi et al. 2006;

Hopkins et al. 2008; Strelcyk & Dau 2009), children with MM HL seem less able to listen in

the "valleys" of a fluctuating background noise to improve their speech perception. However,

we did not find any significant relationship between measures of speech processing and

perception. This contrasts with numerous studies that report small to medium correlations

between subcortical responses and speech intelligibility, especially in noise (Anderson et al.,

2010; Bidelman & Momtaz, 2021; Hornickel et al., 2009; Mai et al., 2018; Mepani et al., 2021;

van Hirtum et al., 2023). Some correlations have also been observed between cortical responses

and speech discrimination (Ching et al., 2023; Näätänen, 2001), although the nature of this

relationship remains unclear (Jost et al., 2015; Sharma et al., 2006; Shafer et al., 2005). Again,

our sample size might be too small to adequately investigate such possible correlations. Future

studies including larger sample sizes are warranted to clarify the relationship between speech

processing and perception in the context of congenital HL. These studies would benefit from a

direct investigation of the effect of childhood HL on subcortical processing of speech in noise

and its relationship with performance.

Conclusions

When presented with unamplified sounds, children with HL show impaired subcortical

and cortical responses to speech syllables. Amplification was beneficial for subcortical and

early (P1) but not late (N2, MMN) cortical responses. Our results support the existence of

multiple gain mechanisms that could underlie homeostatic plasticity throughout the auditory

pathway. Responses elicited at different levels of the auditory pathway appear to display

idiosyncratic changes following HL, and may be age- and level- dependent. Even though speech

perception was poorer in the group with HL than TH (in quiet and in noise), there was no clear

relationship between the neurophysiological and behavioural measures. Even mild to moderate

congenital HL induces functional alterations to the central auditory pathway, each level

showing a specific alteration of its typical response. Future studies are needed to clarify the

relationship between these functional changes and the behavioural performance.

Supplementary Figure 2: Grand average of the subcortical responses at Cz, as elicited by stimulus in each polarity. The upper row represents the spectra evoked by the standard syllable /bɑ/, presented in positive polarity; the lower row represents the spectra evoked by the same stimulus presented in negative polarity. Grand average spectra were computed by averaging the individual amplitude spectra.

Acknowledgments

We are very grateful for the time and efforts of the children who participated in this study and to their parents. We would like to thank Dr Lorna Halliday for her input throughout this project, and for her comments on previous versions of this manuscript. We would also like to thank Gaston Hilkhuysen for his help with aspects of the methods. We thank two anonymous reviewers for their constructive feedback on an earlier version of this manuscript. This work was supported by the FP7 people programme (Marie Curie Actions), grant agreement no FP7-607139 (improving Children's Auditory REhabilitation, iCARE). Axelle Calcus was supported by a MSCA Individual Fellowship (798093, EAR-DNA) and a FNRS MIS (F.4508.22).

Author contributions

A.C. and S.R. conceived the idea, A.C. collected data, A.C. and S.R. analysed the data, A.C. and S.R. interpreted the data, A.C. and S.R. wrote the manuscript.

Data availability

Unidentifiable data, stimuli and statistical analyses scripts are publicly available on: https://github.com/acalcus/MMHL_amplification

Competing interests

The authors declare no competing interests

References

Aiken, S.J., Picton, T.W. (2008). Envelope and spectral frequency-following responses to vowel sounds. *Hearing Res*, 245, 35–47.

Alain, C., Tremblay, K. (2007). The role of event-related brain potentials in assessing central auditory processing. *J Am Acad Audiol*, 18, 573–589.

Alho, K. (1995). Cerebral Generators of Mismatch Negativity (MMN) and Its Magnetic Counterpart (MMNm) Elicited by Sound Changes. *Ear Hearing*, 16, 38–51.

Ananthakrishnan, S., Krishnan, A., Bartlett, E.L. (2016). Human frequency following response: neural representation of envelope and temporal fine structure in listeners with normal hearing and sensorineural hearing loss. *Ear and hearing*, 37, e91–e103.

Anderson, S., Anderson, S., Parbery-Clark, A., et al. (2013). Effects of hearing loss on the subcortical representation of speech cues. *J Acoust Soc Am*, 133, 3030–3038.

Anderson, S., Skoe, E., Chandrasekaran, B., et al. (2010). Neural Timing Is Linked to Speech Perception in Noise. *J Neurosci*, 30, 4922–4926.

Bidelman, G.M. (2018). Subcortical sources dominate the neuroelectric auditory frequency-following response to speech. *Neuroimage*, 175, 56–69.

Bidelman, G.M. (2015). Towards an optimal paradigm for simultaneously recording cortical and brainstem auditory evoked potentials. *J Neurosci Meth*, 241, 94–100. w

Bidelman, G.M., Momtaz, S. (2021). Subcortical rather than cortical sources of the frequency-following response (FFR) relate to speech-in-noise perception in normal-hearing listeners. *Neurosci Lett*, 746, 135664.

BinKhamis, G., Forte, A.E., Reichenbach, T., et al. (2019). Speech Auditory Brainstem Responses in Adult Hearing Aid Users: Effects of Aiding and Background Noise, and Prediction of Behavioral Measures. *Trends Hear*, 23, 233121651984829–20.

Bishop, D.V.M., Hardiman, M., Uwer, R., et al. (2007). Maturation of the long-latency auditory ERP: step function changes at start and end of adolescence. *Developmental Sci*, 10, 565–575.

Boersma, P., and Weenink, D. "Praat: A system for doing phonetics by computer," *Glot. Int.* **5**, 341–347. (2001).

Brennan, M., McCreery, R., Kopun, J., et al. (2016). Masking Release in Children and Adults With Hearing Loss When Using Amplification. *J. Speech, Lang., Hear. Res.*, 59, 110–121.

Calcus, A., Tuomainen, O., Campos, A., et al. (2019). Functional brain alterations following mild-to-moderate sensorineural hearing loss in children. *eLife*, 1–37.

Caras, M.L., Sanes, D.H. (2015). Sustained Perceptual Deficits from Transient Sensory Deprivation. *J Neurosci*, 35, 10831–10842.

Ching, T.Y.C., Zhang, V.W., Ibrahim, R., et al. (2023). Acoustic change complex for assessing speech
discrimination in normal-hearing and hearing-impaired infants. *Clin. Neurophysiol.*, 149, 121–132.

Coffey, E.B.J., Nicol, T., White-Schwoch, T., et al. (2019). Evolving perspectives on the sources of
the frequency-following response. *Nat Commun*, 10, 5036.

Cooke, M. (2006). A glimpsing model of speech perception in noise. *J Acoust Soc Am*, 119, 1562.

Craig, C. GoldWave (Version 5.23) [Sound Editing Program] St Johns, Newfoundland: GoldWave
Inc. (2008).

Crandell, C.C. (1993). Speech Recognition in Noise by Children with Minimal Degrees of
Sensorineural Hearing loss. *Ear Hearing*, 14, 210–216.

Daly, D.M., Roeser, R.J., Moushegian, G. (1976). The frequency-following response in subjects with
profound unilateral hearing loss. *Electroencephalogr. Clin. Neurophysiol.*, 40, 132–142.

Easwar, V., Purcell, D., Wright, T. (2023). Predicting Hearing aid Benefit Using Speech-Evoked
Envelope Following Responses in Children With Hearing Loss. *Trends Hear.*, 27.

Easwar, V., Purcell, D.W., Aiken, S.J., et al. (2015). Evaluation of speech-evoked envelope following
responses as an objective aided outcome measure: Effect of stimulus level, bandwidth, and
amplification in adults with hearing loss. *Ear Hearing*, 36, 635–652.

Eggermont, J.J., Ponton, C.W. (2002). The Neurophysiology of Auditory Perception: From Single
Units to Evoked Potentials. *Audiology Neurotol*, 7, 71–99.

Engström, E., Kallioinen, P., Mentzer, C.N. von, et al. (2021). Auditory event-related potentials and
mismatch negativity in children with hearing loss using hearing aids or cochlear implants – A
three-year follow-up study. *Int J Pediatr Otorhi*, 140, 110519.

Festen, J.M., Plomp, R. (1990). Effects of fluctuating noise and interfering speech on the speech-
reception threshold for impaired and normal hearing. *J Acoust Soc Am*, 88, 1725–1736.

Goldsworthy, R.L., Markle, K.L. (2019). Pediatric Hearing Loss and Speech Recognition in Quiet and
in Different Types of Background Noise. *J Speech Lang Hear Res*, 62, 758–767.

Halliday, L.F., Rosen, S., Tuomainen, O., et al. (2019). Impaired frequency selectivity and sensitivity
to temporal fine structure, but not envelope cues, in children with mild-to-moderate sensorineural
hearing loss. *J Acoust Soc Am*, 146, 4299–4314.

Hecox, K., Galambos, R. (1974). Brain Stem Auditory Evoked Responses in Human Infants and
Adults. *Arch Otolaryngol*, 99, 30–33.

Hirtum, T.V., Somers, B., Dieudonné, B., et al. (2023). Neural envelope tracking predicts speech
intelligibility and hearing aid benefit in children with hearing loss. *Hear. Res.*, 439, 108893.

Hopkins, K., Moore, B.C.J. (2009). The contribution of temporal fine structure to the intelligibility of
speech in steady and modulated noise. *J Acoust Soc Am*, 125, 442–446.

Hopkins, K., Moore, B.C.J., Stone, M.A. (2008). Effects of moderate cochlear hearing loss on the
ability to benefit from temporal fine structure information in speech. *J Acoust Soc Am*, 123, 1140–
1153.

Hornickel, J., Skoe, E., Nicol, T., et al. (2009). Subcortical differentiation of stop consonants relates to
reading and speech-in-noise perception. *Proc. Natl. Acad. Sci. USA*, 106, 13022–13027.

Jacxsens, L., Biot, L., Escera, C., et al. (2024). Frequency-Following Responses in Sensorineural
Hearing Loss: A Systematic Review. *J. Assoc. Res. Otolaryngol.*, 25, 131–147.

Jasper, H. The ten-twenty system of the International Federation. *Electroencephalogr Clin
Neurophysiol*, Suppl **52**: 3-6. (1958).

Ji, H., Yu, X., Xiao, Z., et al. (2023). Features of Cognitive Ability and Central Auditory Processing of
Preschool Children With Minimal and Mild Hearing Loss. *J Speech Lang Hear Res*, 66, 1867–
1888.

Jost, L.B., Eberhard-Moscicka, A.K., Pleisch, G., et al. (2015). Native and non-native speech sound
processing and the neural mismatch responses: A longitudinal study on classroom-based foreign
language learning. *Neuropsychologia*, 72, 94–104.

Koravand, A., Jutras, B., Lassonde, M. (2013). Auditory event related potentials in children with
peripheral hearing loss. *Clin Neurophysiol*, 124, 1439–1447.

Kraus, N., Anderson, S., White-Schwoch, T. (2017). The Frequency-Following Response: A Window
into Human Communication. In Human Auditory Development. (*pp. 1–15*). Cham: Springer
Science & Business Media.

Krishnan, A., Bidelman, G.M., Smalt, C.J. (2012). Relationship between brainstem, cortical and
behavioral measures relevant to pitch salience in humans. *Neuropsychologia*, 50, 2849–2859.

Krizman, J., Bonacina, S., Kraus, N. (2019). Sex differences in subcortical auditory processing emerge
across development. *Hear. Res.*, 380, 166–174.

Krizman, J., Tierney, A., Fitzroy, A.B., et al. (2015). Continued maturation of auditory brainstem
function during adolescence: A longitudinal approach. *Clin. Neurophysiol.*, 126, 2348–2355.

Lewis, D.E., Valente, D.L., Spalding, J. (2015). Effect of minimal/mild hearing loss on children's
speech understanding in a simulated classroom. *Ear Hearing*, 36, 136–144.

Lieu, J.E.C., Kenna, M., Anne, S., et al. (2020). Hearing Loss in Children. *JAMA*, 324, 2195–2205.

Lorenzi, C., Gilbert, G., Carn, H., et al. (2006). Speech perception problems of the hearing impaired
reflect inability to use temporal fine structure. *Proc National Acad Sci*, 103, 18866–18869.

Mahajan, Y., McArthur, G. (2012). Maturation of auditory event-related potentials across adolescence.
Hearing Res, 294, 82–94.

Mahajan, Y., McArthur, G. (2015). Maturation of mismatch negativity and P3a response across
adolescence. *Neurosci Lett*, 587, 102–106.

Mai, G., Howell, P. (2023). The possible role of early-stage phase-locked neural activities in speech-
in-noise perception in human adults across age and hearing loss. *Hearing Res*, 427, 108647.

Mai, G., Tuomainen, J., Howell, P. (2018). Relationship between speech-evoked neural responses and
perception of speech in noise in older adults. *J Acoust Soc Am*, 143, 1333–1345.

Martinez, A., Eisenberg, L., Boothroyd, A. (2013). The Acoustic Change Complex in Young Children
with Hearing Loss: A Preliminary Study. *Seminars Hear*, 34, 278–287.

Mepani, A.M., Verhulst, S., Hancock, K.E., et al. (2021). Envelope following responses predict
speech-in-noise performance in normal-hearing listeners. *J Neurophysiol*, 125, 1213–1222.

Miller, G.A., Licklider, J. (1950). The intelligibility of interrupted speech. *J Acoust Soc Am*, 22, 167–
173.

Mowery, T.M., Kotak, V.C., Sanes, D.H. (2015). Transient Hearing Loss Within a Critical Period
Causes Persistent Changes to Cellular Properties in Adult Auditory Cortex. *Cereb Cortex*, 25,
2083–2094.

Mueller, V., Brehmer, Y., Oertzen, T. von, et al. (2008). Electrophysiological correlates of selective
attention: A lifespan comparison. *Bmc Neurosci*, 9, 18–21.

Näätänen, R. (2001). The perception of speech sounds by the human brain as reflected by the
mismatch negativity (MMN) and its magnetic equivalent (MMNm). *Psychophysiology*, 38, 1–21.

Näätänen, R., Kujala, T., Escera, C., et al. (2012). The mismatch negativity (MMN) - A unique
window to disturbed central auditory processing in ageing and different clinical conditions. *Clin
Neurophysiol*, 123, 424–458.

Oades, R., Dittman-Balcar, A., & Zerbin, D. (1997). Development and topography of auditory event-
related potentials (ERPs): Mismatch and processing negativity in individuals 8-22 years of age.
Psychophysiology, 34, 677-693.

Peelle, J.E., Wingfield, A. (2016). The Neural Consequences of Age-Related Hearing Loss. *Trends
Neurosci*, 1–12.

Plyler, P., Ananthanarayan, A.K. (2001). Human Frequency Following Responses: Representation of
second formant transitions in normal-hearing and hearing-impaired listeners. *J A A A*, 37, e91–
e103.

Ponton, Eggermont, Don, et al. (2000). Maturation of the Mismatch Negativity: Effects of Profound
Deafness and Cochlear Implant Use. *Audiology Neurotol*, 5, 167–185.

Presacco, A., Simon, J.Z., Anderson, S. (2016). Evidence of degraded representation of speech in
noise, in the aging midbrain and cortex. *J. Neurophysiol.*, 116, 2346–2355.

Presacco, A., Simon, J.Z., Anderson, S., et al. (2019). Speech-in-noise representation in the aging
midbrain and cortex: Effects of hearing loss. *Plos One*, 14, e0213899-26.

Qin, M.K., Oxenham, A.J. (2003). Effects of simulated cochlear-implant processing on speech
reception in fluctuating maskers. *J Acoust Soc Am*, 114, 446–454.

Rinne, T., Alho, K., Ilmoniemi, R.J., et al. (2000). Separate Time Behaviors of the Temporal and
Frontal Mismatch Negativity Sources. *Neuroimage*, 12, 14–19.

Scherg, M., Vajsar, J., Picton, T. (1989). A source analysis of the late human auditory evoked
potentials. *Journal of Cognitive Neuroscience*.

Shafer, V., Morr, M., Datta, H., et al. (2005). Neurophysiological indexes of speech processing
deficits in children with specific language impairment. *Journal of Cognitive Neuroscience*, 17.

Shannon, R.V., Zeng, F.-G., Kamath, V., et al. (1995). Speech Recognition with Primarily Temporal
Cues. *Science*, 270, 303–304.

Sharma, A., Kraus, N., McGee, T.J. (1997). Developmental changes in P1 and N1 central auditory
responses elicited by consonant-vowel syllables. *Electroencephalogr Clin Neurophysiology*
Evoked Potentials Sect, 104, 540–545.

Sharma, M., Sharma, M., Purdy, S.C., et al. (2006). Electrophysiological and behavioral evidence of
auditory processing deficits in children with reading disorder. *Clin Neurophysiol*, 117, 1130–1144.

Strelcyk, O., Dau, T. (2009). Relations between frequency selectivity, temporal fine-structure
processing, and speech reception in impaired hearing. *J Acoust Soc Am*, 125, 3328.

Wetzel, N., Widmann, A., Berti, S., et al. (2006). The development of involuntary and voluntary
attention from childhood to adulthood: A combined behavioral and event-related potential study.
Clin. Neurophysiol., 117, 2191–2203.

Zeng, F.-G., Nie, K., Stickney, G.S., et al. (2005). Speech recognition with amplitude and frequency
modulations. *Proc. Natl. Acad. Sci.*, 102, 2293–2298.

Zhang, Y., Yan, F., Wang, L., et al. (2018). Cortical Areas Associated With Mismatch Negativity: A
Connectivity Study Using Propofol Anesthesia. *Front Hum Neurosci*, 12, 38–10.

1062

1063

1064