HIPPOMAPS: MULTISCALE CARTOGRAPHY OF HUMAN HIPPOCAMPAL ORGANIZATION

Jordan DeKraker[1], Donna Gift Cabalo[1], Jessica Royer[1], Ali R. Khan[2], Bradley Karat[2], Oualid Benkarim[1], Raul Rodriguez-Cruces[1], Birgit Frauscher[3], Raluca Pana[1], Justine Y. Hansen[1], Bratislav Misic[1], Sofie L. Valk[4,5,6], Jonathan C. Lau[2], Matthias Kirschner[7], Andrea Bernasconi[1], Neda Bernasconi[1], Sascha Muenzing[5], Markus Axer[5], Katrin Amunts[5,6], Alan C. Evans[1]*, Boris C. Bernhardt[1]*

[1] McConnell Brain Imaging Centre, Montreal Neurological Institute and Hospital, McGill University, Canada
[2] Robarts Research Institute, University of Western Ontario, Canada
[3] Department of Neurology, Duke University Medical Center, Durham, NC, United States
[4] Otto Hahn Group Cognitive Neurogenetics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany
[5] Institute of Neuroscience and Medicine INM-1, Research Centre Jülich, Germany
[6] C. & O. Vogt Institute for Brain Research, University Hospital Düsseldorf, Heinrich-Heine-University, Germany
[7] Division of Adult Psychiatry, Department of Psychiatry, University Hospitals of Geneva, 2, Chemin du Petit-Bel-Air, CH-1226, Thonex, Switzerland
*co-senior authors

ABSTRACT

The hippocampus has a unique microarchitecture, is situated at the nexus of multiple macroscale functional networks, contributes to numerous cognitive as well as affective processes, and is highly susceptible to brain pathology across common disorders. These features make the hippocampus a model to understand how brain structure covaries with function, in both health and disease. Here, we introduce HippoMaps, an open access toolbox and online data warehouse for the mapping and contextualization of hippocampal data in the human brain (http://hippomaps.readthedocs.io). HippoMaps capitalizes on a novel hippocampal unfolding approach as well as shape intrinsic registration capabilities to allow for cross-subject and cross-modal data aggregation. We initialize this repository with data spanning 3D *post-mortem* histology, *ex-vivo* 9.4 Tesla MRI, as well as *in-vivo* structural MRI and resting-state functional MRI (rsfMRI) obtained at 3 and 7 Tesla, together with intracranial encephalography (iEEG) recordings in epilepsy patients. HippoMaps also contains validated tools for spatial map association analysis in the hippocampus that correct for autocorrelation. All code and data are compliant with community standards, and comprehensive online tutorials facilitate broad adoption. Applications of this work span methodologies and modalities, spatial scales, as well as clinical and basic research contexts, and we encourage community feedback and contributions in the spirit of open and iterative scientific resource development.

INTRODUCTION

The hippocampus has long been regarded as a model to understand how brain structure spatially covaries with function (Bahr, 1995; Eichenbaum, 2000). On the one hand, hippocampal anatomy has been recognized to be organized in both anterior-posterior and proximal-distal dimensions (Duvernoy et al., 2013; Olsen et al., 2019). Anterior-posterior organization is emphasized in foundational descriptions of its

anatomical segments (*i.e.,* head, body, and tail) as well as functional gradients along the hippocampal
long axis (Bouffard et al., 2023; Poppenk et al., 2013; Przeździk et al., 2019; Strange et al., 2014; Vogel
et al., 2020; Vos de Wael et al., 2018). Perpendicular to this, there is a preserved arrangement of
hippocampal subfields along the proximal-distal (also referred to as medio-lateral) axis (Genon et al.,
2021; Insausti & Amaral, 2004; Olsen et al., 2019; Paquola, Benkarim, et al., 2020; Ramón y Cajal, 1904;
Yushkevich et al., 2015). These macroanatomical and microstructural features have been suggested to
directly relate to hippocampal circuit organization and its embedding within macroscale functional
networks (Knierim & Neunuebel, 2016; S. Leutgeb & Leutgeb, 2007; Rolls, 2016), contributing to
specific hippocampal computations and its role as a nexus connecting paralimbic, sensory, and
heteromodal association systems, notably the default mode network (Andrews-Hanna, Reidler, Sepulcre,
et al., 2010; Buckner et al., 2008; Smallwood et al., 2021; Vos de Wael et al., 2018). It broad involvement
in multiple macroscale networks is clearly compatible with the key role the hippocampus plays in
numerous cognitive and affective processes, including memory and language function, together with
affective reactivity, stress as well as spatial navigation (Barnett et al., 2024; O'Keefe & Nadel, 1978;
Stachenfeld et al., 2014, 2017; Whittington et al., 2022). Notably, the hippocampus is also recognized as
one of the proximate evolutionary origins of the neocortex (Puelles et al., 2019; Sanides, 1969), making it
a candidate structure to investigate principles of evolutionary conservation and innovation in the primate
lineage (Eichert *et al.*, 2023). Collectively, these insights contribute to the notion that the hippocampus is
a microcosm of the brain, and that an assessment of its sub-regional organization provides key insights
into human neural architectures.
60
The fine-grained subregional organization of the hippocampus contrasts the somewhat coarse assessment
of this structure by most contemporary neuroimaging investigations, which often still treat this complex
archicortical structure as a single entity (Jordan DeKraker et al., 2021), or even erroneously label it as
'subcortical'. This is, in part, due to technical limitations: since the hippocampus is thinner and more
tightly convoluted than the neocortex, it is difficult to appreciate its cortical architecture in magnetic
resonance imaging (MRI) or the extent of its 3D convolutions in sparse histology slices. More recently,
relatively few studies have compared its microstructural to mesoscale structural and functional features
directly, with most studies opting instead to apply subfield parcellation as a proxy (Caldairou et al., 2016;
Iglesias et al., 2015; Kulaga-Yoskovitz et al., 2015; Olsen et al., 2019; Romero et al., 2017; Yushkevich
et al., 2010). Here, we introduce HippoMaps, an open access toolbox and online data warehouse for *(i)*
the surface based mapping and analytical unfolding of hippocampal subregional features, *(ii)* the
contextualization of a given hippocampal map (derived from *e.g.,* a typical task-based functional MRI
experiment or structural abnormality map in disease) with respect to normative hippocampal data
obtained from histology and imaging, and for offering *(iii)* a non-parametric statistical framework to
establish the correlation across standardized surface maps, while controlling for spatial autocorrelation.
HippoMaps adopts best practices and methods developed throughout the neocortical mapping community
(Alexander-Bloch et al., 2018; Glasser et al., 2013; Lepage et al., 2017; Markello et al., 2022), and we
provide a set of tools, tutorials, and guidelines for broad adoption and continued development.
79
HippoMaps benefits from multiple recent technical innovations in hippocampal image processing and
analysis. First, it leverages a unified hippocampal segmentation and surface mapping approach using deep
learning-based image processing (Jordan DeKraker et al., 2022), imposing a known prior topology
(Jordan DeKraker et al., 2018) and shape-inherent inter-subject alignment (DeKraker et al., 2023). Similar

to neocortical surface extraction and registration procedures (Boucher et al., 2009; Dale et al., 1999;
Fischl, Sereno, & Dale, 1999; Fischl, Sereno, Tootell, et al., 1999; Kim et al., 2005; Lyttelton et al., 2007;
MacDonald et al., 2000), this allows for topology-informed inter-subject registration to a standardized
unfolded space (Jordan DeKraker et al., 2023). This has begun a new wave of high-sensitivity
hippocampally-focused studies in topics including the mapping of histology features (J. DeKraker et al.,
2020; Paquola, Benkarim, et al., 2020), blood perfusion (Haast *et al.*, 2023; Ngo *et al.*, 2023),
biophysically-constrained diffusion (Karat *et al.*, 2023), hippocampal sclerosis (Ripart *et al.*, 2023),
neurodevelopmental trajectories (Hanson *et al.*, 2023), functional connectivity (Cabalo *et al.*, 2023;
Lariviere *et al.*, 2023; Xie *et al.*, 2023), visual receptive field mapping (Leferink *et al.*, 2023), and cross-
species comparison (Eichert *et al.*, 2023). With the increasing aggregation of hippocampal features in a
common reference space, it is now possible to devise repositories that allow for a broad contextualization
of hippocampal findings. Such work may aid in the interpretation of findings from new studies and
experiments, for example by allowing for the cross-referencing of these results against established
features of hippocampal functional and structural organization.

At the level of the neocortex, there has been an increasing repertoire of comprehensive open tools for
contextualization of findings, including BALSA (David C. Van Essen et al., 2017), NeuroVault
(Gorgolewski *et al.*, 2015), and NeuroMaps (Markello *et al.*, 2022), as well as other contextualization
methods incorporated in statistical software such as BrainStat (Lariviere et al 2022) and the ENIGMA
toolbox (Lariviere et al. 2022). With HippoMaps, we expand anatomy-driven neuroinformatics to the
hippocampus, and provide a high-quality and broad multimodal online repository of normative maps,
using a common folded and unfolded surface representational space. We initialize the HippoMaps
repository with a spectrum of data spanning 3D histology, structural MRI and resting-state functional
MRI (rsfMRI) obtained at field strengths of 3 and 7 Tesla from healthy individuals, as well as intracranial
encephalography (iEEG) collected from epilepsy patients. We provide tools for high-definition
hippocampal visualization and contextualization. Moreover, we incorporate adapted methods to control
for autocorrelation when assessing spatial maps to one another, a key for accurate enrichment analysis in
the hippocampus (Alexander-Bloch et al., 2018; Karat et al., 2023; Vos de Wael et al., 2020). Finally,
provide an example of how future hippocampal mapping studies can use spatial correlation with
HippoMaps to contextualize results, linking structure and function. This is supported by online tutorials to
reproduce all results shown here (https://github.com/HippAI/hippomaps or https://github.com/MICA-
MNI/hippomaps), with extensibility so future studies may contribute their methods and mapped data
(https://osf.io/92p34/).

METHODS

Datasets
To provide broad coverage of many areas of hippocampal research, we initialize HippoMaps with 30
novel minimally processed but spatially aligned data spanning 3D *post-mortem* histology, high field *in-*
vivo structural as well as resting-state functional MRI (rsfMRI), and intracranial electroencephalography
(iEEG). These data originate from open source resources including BigBrain (Amunts *et al.*, 2013),
AHEAD (Alkemade *et al.*, 2022), MICs (Royer *et al.*, 2022), the MNI open iEEG atlas (Frauscher *et al.*,
2018), and are further supplemented with locally collected data including further healthy structural and

functional MRI obtained at 3 Tesla and 7 Tesla, as well as iEEG data obtained in epilepsy patients that
also underwent pre-implantation multimodal MRI. See the **Supplementary Materials** for details of each
dataset and preprocessing.

Surface mapping

Data processing details are available in the **Supplementary Methods**. Briefly, minimal preprocessing
was applied to each dataset using *micapipe v0.2.2* (https://github.com/MICA-MNI/micapipe) for
structural and functional MRI (Cruces *et al.*, 2022) and custom code for other data. Though the
processing of each data modality differs, they were each mapped to a standardized folded and unfolded
surface space using *HippUnfold v1.3.0* (Jordan DeKraker et al., 2022). Briefly, this entails tissue type
segmentation using a deep UNet neural network, fitting of inner, outer, and midthickness surfaces to
hippocampal gray matter, mapping to a standardized unfolded rectangular space, and then registration in
unfolded space to a standard, histology-derived generated atlas (Jordan DeKraker et al., 2023). This
standardized space is, thus, made equivalent across all subjects. Notably, despite surface meshes having
differing tessellations (**Figure 1A**), they can be interpolated in unfolded space to match microscale
features (*e.g.,* 3D reconstructed histological stains) to MRI or *vice versa*, spanning a scale of micrometers
to millimeters.
141

Figure 1. Mapping multiscale data to standardized hippocampal surfaces. **A)** Surface folding and density are matched to the sample shape and resolution. Mapping to a standardized unfolded space enables registration and interpolation across scales, which can then be followed by parcellation, averaging, or comparison by spatial correlation. **B)** Depth-wise microstructural profiles are calculated by fitting surfaces at multiple depths (yellow-orange-red), including extrapolated surfaces over surrounding tissues (green and pink). Profiles at a given vertex are then translated vertically to maximize alignment to the average before being cut off at the gray matter boundary. **C)** Sparse data like depth electrodes or tissue punches are mapped to a hippocampal surface, and their data are extrapolated across the hippocampal surface proportionally to geodesic distance.

In addition to inner, midthickness, and outer surfaces, any number of intermediate surfaces can be generated at different depths or linearly extrapolated around the outer bounds of the hippocampus (**Figure 1B**) (Marcus *et al.*, 2011). This is especially useful for sub-millimetric data, where laminar or microstructural profile information can be extracted. In this case, we illustrate a function for refining

155   alignment of such profiles using vertical (that is, in the laminar direction) translations. This data-driven
156   refinement leverages image intensity to mitigate slight differences in boundary criteria or segmentation
157   errors between different parts of the hippocampus, without making strong assumptions about how
158   different laminae are stained (provided they are consistent within the sample).
159
Using a similar approach, even sparsely sampled data can be spatially mapped across the hippocampus
(**Figure 1C**). In this case, we map the centroids of iEEG channels to their nearest corresponding
hippocampal vertices. However, in principle, this could also apply to other sparse (or scattered) data such
as tissue punches, other invasive recording devices, small resections, or other irregularly spaced sampling
methods. We then map iEEG channel data to all vertices within <5mm of the channel centroid, and
average data across all channels from all patients with a weighting proportional to geodesic distance from
those vertices. This extrapolation method is more robust than a linear or nearest-neighbour extrapolation,
which would be strongly driven by only one or a few nearby vertices with data mapped to them, while
also still preserving some spatial preference for data from nearby channels.

Significance testing

Spatial autocorrelation can compromise statistical significance testing when testing correlations between
continuous maps (Alexander-Bloch *et al.*, 2018). HippoMaps provides two types of permutation test to
ensure robustness against this issue: Moran spectral randomization (Wagner & Dray, 2015) and "spin"
test randomization (Alexander-Bloch et al., 2018; Karat et al., 2023; Vos de Wael et al., 2020). To make
it suitable for the hippocampus, spin permutation tests include wrapping of the anterior-posterior and
proximal-distal edges of the hippocampus, making the topology of a torus (see (Karat *et al.*, 2023) for
details).

RESULTS

We present novel hippocampal maps in a standardized folded and unfolded space for each of the datasets
outlined above. This includes 30 distinct group-averaged maps which have been attentively preprocessed
and curated. Within each methodology, some interpretation and summarization via dimensionality
reduction is offered, and finally we compare all maps across methodologies in the "Feature combinations"
section.

Histology

Histology is considered a neuroanatomical gold standard, and is the basis for most parcellations and
descriptions of brain regions (Amunts et al., 2020; Brodmann, 1909; Eickhoff et al., 2018; Paquola et al.,
2019). Here we examined data collected from BigBrain Merker staining for cell bodies (Amunts *et al.*,
2013), 3D polarized light imaging (PLI) of neural processes (M. Axer et al., 2011), and the AHEAD
dataset with different stains serving as markers of neurons, myelin, and subtypes of interneurons
(Alkemade *et al.*, 2022) (**Figure 2A**). Most features showed banding in the proximal-distal direction, in
alignment with the subfield atlas shown in **Figure 1**.

Figure 2. Histology mapping, depth-wise microstructural profiles, and dimensionality reduction. **A)** Sample slices and averaged 3D maps of histological features. Maps are averaged across depths and, where possible, samples. **B)** Example of microstructural profile shapes from five evenly spaced bins across the proximal-distal axis of the Merker stain map. Grey indicates points outside of the gray matter mask. **C)** Correlation between microstructural profiles, concatenated across modalities, at each vertex (left). Dimensionality reduction into primary diffusion embedding gradients 1-3 (right).

Microstructural (or laminar) profiles are shown for five ROIs across the proximal-distal axis of the BigBrain Merker stain (**Figure 2B**). They show a tight unimodal distribution in the distal CA fields, and a more bimodal distribution in the subiculum as expected based on their known laminar architectures (Duvernoy et al., 2013). Profiles for all vertices were concatenated across all stains to make multimodal profiles, a common method for characterizing laminar structure (Schleicher *et al.*, 1999). Next, multimodal covariance matrices between vertices were calculated (mMPC matrix) (**Figure 2C**). Diffusion map embedding, a non-linear dimensionality reduction technique (Coifman *et al.*, 2005; Margulies *et al.*, 2016; Vos de Wael *et al.*, 2020), decomposed the mMPC matrix into primary components, or gradients, that highlighted the differences between vertices with respect to all modalities and depths. In the first gradient, a sharp boundary was seen between the subicular complex and proximal CA1 and the rest of the hippocampus. The second and third gradients in turn highlighted the CA2-3 regions and CA1 with parts of the subiculum, respectively. This is data-driven evidence that subfields across the proximal-distal extent of the hippocampus, rather than anterior-posterior or other patterns, account for structural variance in the hippocampus with respect to these stains. These data-driven decompositions, thereby, echo classical and recent neuroanatomy descriptions of hippocampal microstructure (Ding & Van Hoesen, 2015; Duvernoy et al., 2013; Olsen et al., 2019).

Structural MRI

MRI is a key tool for studying human neuroanatomy and structure-function relations due to its non-invasive nature and potential for biomarker discovery. 7 Tesla (7T) and *ex-vivo* 9.4T scanning are

218   especially powerful, achieving greater resolution and contrast than typical 3T or 1.5T clinical scans
219   (Duyn, 2012; Opheim *et al.*, 2021). Here, we provide healthy normative maps for such scans (**Figure 3A**)
220   including popular acquisitions: quantitative T1 relaxometry (qT1) and its non-quantitative *ex-vivo*
221   inverse: R1, T2* and its inverse R2*, proton density, diffusion weighted imaging (DWI) estimates of
222   fractional anisotropy (FA) and apparent diffusivity coefficient (ADC), and magnetic transfer ratio (MTR).

223
224   **Figure 3.** Structural MRI mapping, inter-sample consistency, and dimensionality reduction. **A)** Sample slices and averaged 3D
225   maps of histological features. Maps are averaged across depths and, where possible, samples. **B)** Consistency, as measured by the
226   correlation between all pairs of individual sample maps, **C)** Correlation between microstructural profiles, concatenated across
227   modalities, at each vertex (*left*). Dimensionality reduction into primary diffusion embedding gradients 1-3 (*right*).

228
Multiple scans were available for averaging (n=4 left+right hippocampi at 9.4T and n=20 left+right
hippocampi at 7T), enabling a calculation of consistency across samples via Pearson's R (**Figure 3B**).
DWI and qT1 maps were also calculated in a second validation dataset, consisting of 82 locally scanned
healthy participants (including the subset from the MICA-MICs dataset) with a 3T scanner, which
showed similar patterns (**Figure S1**). mMPCs were generated as above and were reduced using diffusion
map embedding into primary gradients, which again highlighted differences across subfields. Only the
third gradient showed anterior-posterior differences, largely within the CA1 subfield.

Functional MRI

Functional MRI during the resting state (rsfMRI) allows interrogation of intrinsic brain function via the
analysis of spontaneous activity and its statistical dependencies, and has become a key technique in the
mapping of functional-anatomical systems (Biswal et al., 1997; Buckner et al., 2008; Smith et al., 2009).
Here, we examined several features of rsfMRI in 88 healthy participants scanned at 3T. Intrinsic
timescale is a measure of the time it takes for the temporal autocorrelation to drop below a threshold
(Golesorkhi *et al.*, 2021; Wolff *et al.*, 2022) (**Figure 4A**). On a functional level, this is thought to be
driven in part by recurrent connections that maintain activity patterns on the order of seconds (Fallon *et*
al., 2020). Regional homogeneity considers the similarity between adjacent vertices' time series, which is

thought to indicate the extent of horizontal (*i.e.,* between cortical columns) excitatory connectivity (Zang
et al., 2004) (**Figure 4B**). Finally, macroscale functional connectivity is by far the most popular rsfMRI
feature, with many rich properties that have been explored with respect to white matter connections
(Damoiseaux & Greicius, 2009; Greicius et al., 2009; Honey et al., 2009), network properties
(Schmittmann et al., 2015; van den Heuvel & Sporns, 2013), organizational gradients (Bernhardt et al.,
2022; Margulies et al., 2016; Paquola et al., 2019; Park et al., 2021), and many other summary metrics.
For simplicity, we examined connectivity between all hippocampal vertices and neocortical parcels from
the Schaeffer400 parcellation (Schaefer *et al.*, 2018) (**Figure 4C**). The consistency of maps was examined
as above, and all measures were significantly greater than zero. Repetition of these analyses in a smaller
sample of 7T rsfMRI data showed consistent results (**Figure S2**).

255
256 **Figure 4.** Functional MRI properties. Resting state (rsfMRI) data were used to calculate **A)** intrinsic timescale (recurrence), **B)**
257 regional homogeneity (short range connectivity), and **C)** functional connectivity (long range; to the neocortex). **D)**
258 Decomposition of functional connectivity patterns across hippocampal vertices into primary diffusion map embedding gradients.
259
As mentioned above, functional connectivity is a rich measure that can be summarized in many ways.
Here, we identified gradients of intrinsic hippocampal connectivity variations (**Figure 4D**) using the
aforementioned non-linear decomposition techniques. Consistent with previous work (Genon et al., 2021;
Poppenk et al., 2013; Przeździk et al., 2019; Strange et al., 2014; Vogel et al., 2020; Vos de Wael et al.,
2018), we found anterior-posterior differentiation in the first hippocampal gradient, together with
proximal-distal banding with CA1 in particular differing from the other subfields. Neocortical
counterparts of this gradient show that anterior and CA1 regions shared more connectivity with temporal
pole, insula, and frontal regions whereas more posterior and non-CA1 subfields shared connectivity with
more posterior parietal and visual areas, again consistent with previous findings (Vos de Wael *et al.*,
2018). The second gradient also showed differentiation of CA1 from subiculum and CA2-3 in the more

middle and posterior regions, with neocortical correspondences to medial prefrontal and posterior
cingulate regions for CA1 and more visual areas for CA2-3 and posterior subiculum.

Intracranial EEG

Invasive recording methods such as iEEG provide a direct measure of neural activity at high temporal
resolution, but typically have lower spatial coverage and are limited to neurological patient populations.
In that sense, they can be considered as scattered spatial data, which can be interpolated or extrapolated
for hippocampal mapping as described in **Figure 1C**, or following previous approaches (Frauscher *et al.*,
2018). We employ common measures of the periodic component of iEEG data, as shown by power
spectrum density and additionally further simplified to Delta, Theta, Alpha, Beta, and Gamma band
powers from low to high frequencies, respectively. Power spectrum densities and band powers derived
from hippocampal channels resembled those derived from all channels (**Figure 5A**). Extrapolating
channel information across neighbouring vertices from a given hippocampus, a spatial pattern emerged in
which both proximal-distal and anterior-posterior differences were seen (**Figure 5B**). Band power is a
limited measure of the full power spectrum density though, and so in **Figure 5C** we performed gradient
mapping of the full power spectrum density. This showed a primary anterior-posterior gradient driven by
higher Theta and Alpha power in the posterior and higher Delta power in the anterior hippocampus. The
second gradient showed increased Delta power in the anterior and posterior hippocampus, while the third
gradient showed a slight increase in Delta and decrease in Theta in the subiculum. Results were consistent
when using an open iEEG atlas (Frauscher *et al.*, 2018) or locally collected data in patients (Paquola,
Seidlitz, *et al.*, 2020), showing largely conserved patterns in **Figure S3**.

290
Figure 5. Intracranial EEG (iEEG) properties from time periods deemed "normal" in implanted patients assessed during resting
state. **A)** (left) power spectrum density plots of all channels (n=4279) and hippocampal channels (<5mm from any hippocampal
midthickness vertex) (n=81), standard deviation shaded. (right) lognormal power within each band for each hippocampal channel,
with vertical lines indicating the median and with corresponding bands from all channels in gray. **B)** Spatial extrapolation
weighted by geodesic distance shows largely anterior-posterior differences in band powers. **C)** Power spectrum densities reduced
into primary diffusion map embedding gradients.

Feature combinations

The biggest advantage of a common hippocampal mapping space is that it allows for direct spatial correlation between features from different scales and methods. In Figure **6A**, we examined relationships between all maps shown above using Pearson's R with an adapted spin test significance testing to control for spatial autocorrelation in the data. This revealed many greater-than-chance correlations, both within methodologies and between. Finally, we additionally compared morphological measures of thickness, gyrification, and curvature which are generated within the *HippUnfold* workflow (**Figure S4**). Previous work (J. DeKraker et al., 2020) showed that these features differed between MRI and histology, with the latter showing greater detail including more gyrification and lower thickness. After group-averaging, each of these features was significantly spatially correlated between histology and MRI.

Figure 6. Relationship between all hippocampal maps. **A)** correlation matrix of all features, after resampling to a common 0.5mm vertex-spacing surface. **B)** Diffusion map embeddings 1-3 across all features. **C)** Alignment of gradients 1 and 2 to hippocampal subfields, proximal-distal, and anterior-posterior axes. **D)** Absolute correlation between each feature map and the anterior-posterior axis (Pearson's R) and the maximum permuted subfield labels (Spearman's R).

We performed a dimensionality reduction across all features from all figures using diffusion map gradient embeddings. For visualization, we plotted components 1 and 2 with colour coding according to subfield and continuous anterior-posterior and proximal-distal gradients (**Figure 6C**). The proximal-distal and anterior-posterior axes of the hippocampus are closely aligned to gradients 1 and 2, respectively, with gradient 1 explaining approximately twice the variance (**Figure 6B**). This suggests that while these two

318   axes emerge as natural summaries of many hippocampal feature maps, the proximal-distal direction is
319   stronger.
320
Figure 6D provides a summary of which measures are most correlated with the anterior-posterior and
subfield axes of the hippocampus. As expected, the strongest subfield relationships were observed in
histological features such as Calbindin and Calretinin staining, or thickness measures at a histological
level of precision. Many structural 9.4T and 7T features also showed strong subfield correlations,
especially qT1 and qR1. This is encouraging given the increasing availability and adoption of quantitative
T1 sequences (Bidhult *et al.*, 2016; Haast *et al.*, 2016; van der Weijden *et al.*, 2021). The employed
rsfMRI and iEEG features were still moderately correlated with subfield division, but iEEG and rsfMRI
gradients in particular showed strong correlations with the anterior-posterior hippocampal axis. Some
caution should be exercised here: iEEG data were sparsely sampled and so after extrapolation each band
power map was very smooth, which could amplify correlation values (but not significance, since spin test
permutations were used to control for spatial autocorrelation). Note also that laminar profiles were not
used in this analysis, and histological measures in particular can benefit from the information added by
such methods due to their high precision.

Usability experiment and documentation

HippoMaps as an open toolbox and online data warehouse paves the way for multiple new research
avenues, examples of which are shown in **Figure 7**. We anticipate that as hippocampal mapping studies
are performed in other research areas, authors can use the initial maps provided here as comparisons and
will upload their own maps in the spirit of open and reproducible science, and also to boost the visibility
of their work. To this end, we provide a set of Python tools, well documented example code to reproduce
the maps shown here (labeled as tutorials), and guidelines for how other experimenters should upload
their maps to this repository. We have and will continue to answer questions and create community
resources   via   GitHub   (https://github.com/HippAI/hippomaps   or   https://github.com/MICA-
MNI/hippomaps),   and   all   current   maps   are   available   on   the   Open   Science   Framework
(https://osf.io/92p34/).

Figure 7. Examples of HippoMaps usage. **A)** New experiments can utilize HippoMaps tools and resources while also contributing new maps to the repository. Only example repository maps are shown, with descriptive naming and additional details to be provided in README files. **B)** task-fMRI during the Mnemonic Similarity Task (MST) to probe the haemodynamic response function (HRF) magnitudes during successful pattern separation and novel trials. These maps are then compared to all others (right), listing the top three strongest correlations (black lines). **C)** Morphological differences between ipsilateral temporal lobe epilepsy (TLE) patients and healthy controls

Figure 7A shows a generic use case of HippoMaps wherein a new finding is contextualized by comparison to other maps in HippoMaps, and data is in turn contributed to HippoMaps to extend its utility in future work. **Figure 7B** illustrates an example experiment with task-fMRI using the Mnemonic

Similarity Task (MST) designed to probe pattern separation, a task thought to preferentially involve
hippocampal subregions (Pishdadian et al., 2020; Stark et al., 2019). This can be seen most strongly in
subiculum for the successful pattern separation trials, whereas trials with novel stimuli showed anterior-
posterior differentiation. Comparing these maps directly to microcircuit features provides context for the
demands of these two task conditions: pattern separation was strongly correlated to detailed maps of
curvature, thickness, and neocortical connectivity, whereas novelty was moderately correlated to intrinsic
timescale, beta band power, and gamma band power (**Figure 7B**, right). Further task-fMRI results from
an object-pairing memory task, as well as replication data of the MST at 7T, are shown in **Figure S5**.

Figure 7C illustrates an example experiment comparing 35 temporal lobe epilepsy (TLE) patients to 81
healthy, age- and sex-matched controls scanned at 3T MRI. Reductions in hippocampal thickness and
gyrification are seen, with the greatest changes in CA1 and CA4 subfields, which have previously been
identified as vulnerable areas (Blümcke et al., 2012, 2013; Duvernoy et al., 2013; Steve et al., 2020).
Comparing thickness reduction patterns to other maps shows moderate correlations with rsfMRI
properties of intrinsic timescale, neocortical connectivity, and histological Bieloschowsky staining.
Gyrification loss was strongly correlated with healthy gyrification in histology and 7T MRI, and iEEG
delta band power.

DISCUSSION

Despite its critical role in human brain organization in both health and disease, the field lacks a
standardized framework to aggregate, represent, and compare structural and functional features of the
hippocampus. The current work presented HippoMaps as a centralized toolbox and online data warehouse
for hippocampal subregional analysis and contextualization. HippoMaps is based on a standardized
hippocampal reference space for data aggregation, sharing, and analysis, which leverages recent advances
in automated hippocampal segmentation and computational unfolding (Jordan DeKraker et al., 2022), as
well as improvements for cross-modal and cross-subject alignment (DeKraker et al., 2023). This
repository is initialized with 30 novel maps of hippocampal subregional organization, aggregating a broad
array of features from 3D *post-mortem* histology, *ex-vivo* 9.4 Tesla MRI, alongside with *in-vivo* structural
and resting-state functional MRI (rsfMRI) obtained at 3 and 7 Tesla, as well as intracranial
encephalography (iEEG) collected from a large cohort of epilepsy patients. This is further extended by a
host of tools for visualization and contextualization, as well as online tutorials that recreate the maps
shown here and demonstrate how new data can be incorporated and analyzed. HippoMaps will provide
key guidance to: *(i)* compare hippocampal features derived from different methods, in particular to cross-
reference *in-vivo* imaging measures with *ex-vivo* and *post-mortem* data, *(ii)* interrogate structure-function
relationships, for example by contextualizing task-based fMRI findings or intracranial neural recording
against spatial patterns obtained from anatomical and microstructural measures, *(iii)* contextualizing case-
control deviations in clinical populations against established principles of subregional hippocampal
organization, and *(iv)* refining our understanding of hippocampal circuitry, by mapping its functional
connectivity and microstructure for a better understanding of its computational operations and transfer
functions at the subregional level. HippoMaps is fully open access and designed according to community
standards (http://hippomaps.readthedocs.io), to facilitate its dissemination and usability. As such, we
anticipate that HippoMaps will represent a powerful analytical ally for fundamental and clinical
neuroscientists alike. Considering the unique role the hippocampus plays in human neuroanatomy and

cognition (Duvernoy et al., 2013; O'Keefe & Nadel, 1978) and its likely important computational
properties (Knierim & Neunuebel, 2016; S. Leutgeb & Leutgeb, 2007), it may furthermore provide key
insights and guidance into the design and validation of emerging bio-inspired AI architectures.
401

We anticipate that surface-based registration will become the standard for hippocampal mapping, as it has
in the neocortex (Fischl, Sereno, Tootell, et al., 1999; Glasser et al., 2013; Ma et al., 2023; Robinson et
al., 2014; D. C. Van Essen et al., 1998). HippoMaps is a major step in advancing the usability of this
methodology, generating utilities, scientific context, and an open community for examining the
hippocampus in detail. Moreover, our repository is designed to employ the same data standards that have
already been extensively developed for neocortical brain imaging data including Brain Imaging Data
Standards (BIDS) (Gorgolewski *et al.*, 2016); NIfTI/GIfTI file formatting (Glasser *et al.*, 2013); and
Findability, Accessibility, Interoperability, and Reusability (FAIR) principles (Gorgolewski & Poldrack,
2016; Wilkinson et al., 2016). Despite its demonstrated benefits, surface-based alignment is not yet
universal for the neocortex and certainly still in its infancy for the hippocampus. Thus, while we
encourage the use of surface-based methods, we also provide code and examples of how to map
volumetrically aligned hippocampal data (*e.g.,* in a standard volumetric space such as MNI152 or others)
to hippocampal surfaces for comparison and contribution to HippoMaps. In the field, work progresses at
the level of hippocampal subfield parcellation at the level of histology, for example to derive additional
subregional divisions (González-Arnay et al., 2024; Henriksen et al., 2010; Igarashi et al., 2014).
Moreover, there have been ongoing efforts by the neuroimaging community to harmonize boundary
heuristics (Olsen et al., 2019; Yushkevich et al., 2015). Under the HippoMaps framework, descriptions go
beyond typical unitary descriptions of the hippocampus and beyond its parcellation into subfields to the
level of mapping vertex-wise or columnar structure of hippocampal archicortex. The columnar level
represents an important structural and functional modularization of the brain (Mountcastle, 1997), and has
the potential to unlock new facets of hippocampal computation. As such, different subfield parcellations
can also be converted to surface format and integrated seamlessly within the HippoMaps warehouse.
Thus, we apply considerable futureproofing, and we encourage the broader hippocampal research
community to upload their own maps to this repository under our support, curation, and online guidelines
and tutorials.
427

Multi-feature aggregation as in the HippoMaps repository provides extensive opportunities to assess
relationships between hippocampal structure and function, to cross-validate *in-vivo* measures with *ex-vivo*
and *post-mortem* imaging as well as histological data. Structural and microstructural data derived from
3D histology and MRI currently aggregated support a close alignment of many feature maps with the
classic subfields account of the hippocampal circuitry. Moreover, several measures, particularly those
derived from functional modalities such as rsfMRI or iEEG, lend additional evidence for anterior-
posterior differentiation of the hippocampal formation. Specifically, gradient decomposition of
hippocampal rsfMRI connectivity and iEEG power spectrum densities showed that anterior-posterior
differentiation captured most inter-regional variance, whereas histological and structural MRI measures
showed primarily proximal-distal or subfield-related differentiation. It is notable that some features
showed extreme intensity values at the anterior and posterior edges - these are relatively small in native
space and so have limited constituent data and are prone to misalignment artifacts. Thus, the anterior and
posterior edges of each map should be interpreted with some caution. Nevertheless, the consistently
repeated structural motifs across the anterior-posterior axis of the hippocampus are suggestive of parallel

repeated computations being performed on different input and output information across the anterior-
posterior hippocampal axis, in line with prior accounts (Poppenk *et al.*, 2013; Strange *et al.*, 2014). These
two dimensions have also been suggested to topographically represent the functional embedding of the
broader mesiotemporal region in large-scale functional networks, in particular default mode and multiple
demand networks (Andrews-Hanna, Reidler, Sepulcre, et al., 2010; Buckner et al., 2008; Duncan, 2010),
which provides a potential substrate for the parametric mixing of both functional systems in macroscale
brain function (Paquola, Benkarim, *et al.*, 2020). It is, therefore, not surprising that two axes explain the
greatest proportion of the variance across all maps in the current repository as well, consolidating the
notion that a two dimensional organization may serve as a powerful summary descriptor for a broad array
of hippocampal structural and functional features (Genon *et al.*, 2021).
452
We provide adapted methods to control for autocorrelation when comparing spatial maps to one another
in the hippocampus. We specifically adapted Moran's spectral randomization and "spin test" permutation
testing that have previously been introduced to study neocortical data (Alexander-Bloch et al., 2018;
Karat et al., 2023; Vos de Wael et al., 2020; Wagner & Dray, 2015). These methods reveal robust
correlations between many of the maps included here. Many of these relationships support the validity of
the methods being applied, for example between *in-vivo* qT1 and *ex-vivo* R1 which are inverses of one
another. Another example is that functional connectivity of the hippocampus was strong to default mode
neocortical areas, as shown in previous work (Andrews-Hanna, Reidler, Huang, et al., 2010; Norman et
al., 2021; Vos de Wael et al., 2018; Ward et al., 2014), with connectivity being strongest in the
subiculum. This recapitulates the role of the subiculum as the primary output structure of the
hippocampus, and contributions of the hippocampus to functions typically ascribed to the default mode
network such as mind-wandering, episodic recall, or future-thinking that are frequent during rest (Bellana
et al., 2017; Buckner, 2010; Christoff et al., 2016; Fox et al., 2015; Ross & Easton, 2022; Schacter et al.,
2017; Yang et al., 2020). Some relationships reveal novel information about the methods themselves: PLI
transmittance is thought to reflect many microscopic structures under the broad heading of "neural
processes" or "nerve fibers" (H. Axer et al., 2001; Dammers et al., 2012). Across the extent of the
hippocampus, this feature correlated with Beilsochowsky and Thionin staining, R2*, average neocortical
functional connectivity, and, most significantly, rsfMRI intrinsic timescale. Intrinsic timescale is
hypothesized to relate to recurrent connections (Chaudhuri et al., 2014), which could indeed be supported
by dense neural processes. Finally, we illustrate contextualization via nonlinear gradient decomposition
across maps. When applied to all maps, we show data-driven separation of subfields, in line with previous
work. We also note that in this latent space, CA4 closely resembles CA1, even though they are not
adjacent topologically. This fits descriptions of CA4 as having a wide pyramidal layer with large and
dispersed neurons, similar to CA1 (Duvernoy et al., 2013), and indeed in some cases these two areas have
similar disease vulnerabilities (Blümcke *et al.*, 2012). Future work may determine more selectively what
features make these two regions similarly vulnerable or examine why in some disease subtypes one is
affected without the other.
480
At the level of the neocortex, several packages already exist to facilitate the contextualization of results
(Larivière et al., 2023, 2021; Markello et al., 2022). With HippoMaps, such an approach is now also
possible for the hippocampal region, and we demonstrate the contextualization of task fMRI maps during
an episodic memory paradigm as well morphological alterations in patients with temporal lobe epilepsy
relative to healthy individuals. Such approaches can help to clarify the hypothetical role of

microstructural features in specific hippocampal computations, such as pattern separation (Bakker et al.,
2008; J. K. Leutgeb et al., 2007; Schmidt et al., 2012), pattern completion (Guzman et al., 2016; S.
Leutgeb & Leutgeb, 2007), and novelty detection (Chen *et al.*, 2011; Larkin *et al.*, 2014). These
previously assumed relations of function to microstructure have generally relied on parcellations of the
hippocampus into stereotyped subfields; with HippoMaps, it is instead possible to compare functional and
microstructural maps directly without any predefined subfield labelling. In addition to offering potential
increases in anatomical specificity, this representation may also lend itself more naturally to sensitive
spatial correlation with autocorrelation control through permutation testing. One area for future work will
lie in consolidating mesoscale connectivity with detailed descriptions of the internal hippocampal
circuity, which will not only help to further understand the computations of specific hippocampal
subregions but which may also clarify the different substrates of computation (Beaujoin et al., 2018;
Bennett & Stark, 2016; Berron et al., 2016; Karat et al., 2023; Lacy et al., 2011; Ly et al., 2020). Indeed,
hippocampal circuitry has inspired the basic ways in which we think about biological computation,
spurring principles such as long-term potentiation (Hebb, 2005), and carrying important inventions like
the Boltzmann machine (Ackley et al., 1985) and Tolman Eichenbaum machine (Whittington *et al.*,
2020). Even recent theories and computational models still center around hippocampal structure as told
through a stereotyped subfield architecture (Gandolfi et al., 2023; Whittington et al., 2020). Formal
mapping, rather than stereotyped descriptions, can extend this work, building up biological plausibility of
such models and scaffolding our understanding of these systems. For this reason, HippoMaps may also
provide precise macro-, meso- and micro-scale hippocampal features in a common same space to further
identify and harness computational properties of its circuitry.
507
508
509
510

References

Ackley, D. H., Hinton, G. E., & Sejnowski, T. J. (1985). A learning algorithm for boltzmann machines. *Cognitive Science*, *9*(1), 147–169.

Alexander-Bloch, A. F., Shou, H., Liu, S., Satterthwaite, T. D., Glahn, D. C., Shinohara, R. T., Vandekar, S. N., & Raznahan, A. (2018). On testing for spatial correspondence between maps of human brain structure and function. *NeuroImage*, *178*, 540–551.

Alkemade, A., Bazin, P.-L., Balesar, R., Pine, K., Kirilina, E., Möller, H. E., Trampel, R., Kros, J. M., Keuken, M. C., Bleys, R. L. A. W., Swaab, D. F., Herrler, A., Weiskopf, N., & Forstmann, B. U. (2022). A unified 3D map of microscopic architecture and MRI of the human brain. *Science Advances*, *8*(17), eabj7892.

Amunts, K., Lepage, C., Borgeat, L., Mohlberg, H., Dickscheid, T., Rousseau, M.-É., Bludau, S., Bazin, P.-L., Lewis, L. B., Oros-Peusquens, A.-M., Shah, N. J., Lippert, T., Zilles, K., & Evans, A. C. (2013). BigBrain: an ultrahigh-resolution 3D human brain model. *Science*, *340*(6139), 1472–1475.

Amunts, K., Mohlberg, H., Bludau, S., & Zilles, K. (2020). Julich-Brain: A 3D probabilistic atlas of the human brain's cytoarchitecture. *Science*, *369*(6506), 988–992.

Andrews-Hanna, J. R., Reidler, J. S., Huang, C., & Buckner, R. L. (2010). Evidence for the default network's role in spontaneous cognition. *Journal of Neurophysiology*, *104*(1), 322–335.

Andrews-Hanna, J. R., Reidler, J. S., Sepulcre, J., Poulin, R., & Buckner, R. L. (2010). Functional-anatomic fractionation of the brain's default network. *Neuron*, *65*(4), 550–562.

Avants, B. B., Tustison, N., Song, G., & Others. (2009). Advanced normalization tools (ANTS). *The Insight Journal*, *2*(365), 1–35.

Axer, H., Axer, M., Krings, T., & Keyserlingk, D. G. (2001). Quantitative estimation of 3-D fiber course in gross histological sections of the human brain using polarized light. *Journal of Neuroscience Methods*, *105*(2), 121–131.

Axer, M., Grässel, D., Kleiner, M., Dammers, J., Dickscheid, T., Reckfort, J., Hütz, T., Eiben, B., Pietrzyk, U., Zilles, K., & Amunts, K. (2011). High-resolution fiber tract reconstruction in the human brain by means of three-dimensional polarized light imaging. *Frontiers in Neuroinformatics*, *5*, 34.

Bahr, B. A. (1995). Long-term hippocampal slices: a model system for investigating synaptic mechanisms and pathologic processes. *Journal of Neuroscience Research*, *42*(3), 294–305.

Bakker, A., Kirwan, C. B., Miller, M., & Stark, C. E. L. (2008). Pattern separation in the human hippocampal CA3
and dentate gyrus. *Science*, *319*(5870), 1640–1642.

Barnett, A. J., Nguyen, M., Spargo, J., Yadav, R., Cohn-Sheehy, B. I., & Ranganath, C. (2024). Hippocampal-
cortical interactions during event boundaries support retention of complex narrative events. *Neuron*, *112*(2),
319-330.e7.

Beaujoin, J., Palomero-Gallagher, N., Boumezbeur, F., Axer, M., Bernard, J., Poupon, F., Schmitz, D., Mangin, J.-
F., & Poupon, C. (2018). Post-mortem inference of the human hippocampal connectivity and
microstructure using ultra-high field diffusion MRI at 11.7 T. *Brain Structure & Function*, *223*(5), 2157–
2179.

Bellana, B., Liu, Z.-X., Diamond, N. B., Grady, C. L., & Moscovitch, M. (2017). Similarities and differences in the
default mode network across rest, retrieval, and future imagining. *Human Brain Mapping*, *38*(3), 1155–
1171.

Bennett, I. J., & Stark, C. E. L. (2016). Mnemonic discrimination relates to perforant path integrity: An ultra-high
resolution diffusion tensor imaging study. *Neurobiology of Learning and Memory*, *129*, 107–112.

Bernhardt, B. C., Smallwood, J., Keilholz, S., & Margulies, D. S. (2022). Gradients in brain organization.
NeuroImage, *251*, 118987.

Berron, D., Schütze, H., Maass, A., Cardenas-Blanco, A., Kuijf, H. J., Kumaran, D., & Düzel, E. (2016). Strong
Evidence for Pattern Separation in Human Dentate Gyrus. *The Journal of Neuroscience: The Official
Journal of the Society for Neuroscience*, *36*(29), 7569–7579.

Bidhult, S., Kantasis, G., Aletras, A. H., Arheden, H., Heiberg, E., & Hedström, E. (2016). Validation of T1 and T2
algorithms for quantitative MRI: performance by a vendor-independent software. *BMC Medical Imaging*,
16(1), 46.

Biswal, B. B., Van Kylen, J., & Hyde, J. S. (1997). Simultaneous assessment of flow and BOLD signals in resting-
state functional connectivity maps. *NMR in Biomedicine*, *10*(4–5), 165–170.

Blümcke, I., Coras, R., Miyata, H., & Ozkara, C. (2012). Defining clinico-neuropathological subtypes of mesial
temporal lobe epilepsy with hippocampal sclerosis. *Brain Pathology* , *22*(3), 402–411.

Blümcke, I., Thom, M., Aronica, E., Armstrong, D. D., Bartolomei, F., Bernasconi, A., Bernasconi, N., Bien, C. G.,
Cendes, F., Coras, R., Cross, J. H., Jacques, T. S., Kahane, P., Mathern, G. W., Miyata, H., Moshé, S. L.,

Oz, B., Özkara, Ç., Perucca, E., … Spreafico, R. (2013). International consensus classification of

hippocampal sclerosis in temporal lobe epilepsy: a Task Force report from the ILAE Commission on

Diagnostic Methods. *Epilepsia*, *54*(7), 1315–1329.

Boucher, M., Whitesides, S., & Evans, A. (2009). Depth potential function for folding pattern representation,

registration and analysis. *Medical Image Analysis*, *13*(2), 203–214.

Bouffard, N. R., Golestani, A., Brunec, I. K., Bellana, B., Park, J. Y., Barense, M. D., & Moscovitch, M. (2023).

Single voxel autocorrelation uncovers gradients of temporal dynamics in the hippocampus and entorhinal

cortex during rest and navigation. *Cerebral Cortex* , *33*(6), 3265–3283.

Brodmann, K. (1909). *Vergleichende Lokalisationslehre der Grosshirnrinde in ihren Prinzipien dargestellt auf*

Grund des Zellenbaues. Barth.

Buckner, R. L. (2010). The role of the hippocampus in prediction and imagination. *Annual Review of Psychology*,

61(1), 27–48, C1-8.

Buckner, R. L., Andrews-Hanna, J. R., & Schacter, D. L. (2008). The brain's default network: anatomy, function,

and relevance to disease. *Annals of the New York Academy of Sciences*, *1124*, 1–38.

Cabalo, D. G., DeKraker, J., Royer, J., Xie, K., Tavakol, S., Rodríguez-Cruces, R., Bernasconi, A., Bernasconi, N.,

Weil, A., Pana, R., Frauscher, B., Caciagli, L., Jefferies, E., Smallwood, J., & Bernhardt, B. C. (2023).

Differential Reorganization of Episodic and Semantic Memory Systems in Epilepsy-Related

Mesiotemporal Pathology. In *bioRxiv* (p. 2023.09.28.560002). https://doi.org/10.1101/2023.09.28.560002

Caldairou, B., Bernhardt, B. C., Kulaga-Yoskovitz, J., Kim, H., Bernasconi, N., & Bernasconi, A. (2016). A Surface

Patch-Based Segmentation Method for Hippocampal Subfields. *Medical Image Computing and Computer-*

Assisted Intervention – MICCAI 2016, 379–387.

Chaudhuri, R., Bernacchia, A., & Wang, X.-J. (2014). A diversity of localized timescales in network activity. *ELife*,

3, e01239.

Chen, J., Olsen, R. K., Preston, A. R., Glover, G. H., & Wagner, A. D. (2011). Associative retrieval processes in the

human medial temporal lobe: hippocampal retrieval success and CA1 mismatch detection. *Learning &*

Memory , *18*(8), 523–528.

Christoff, K., Irving, Z. C., Fox, K. C. R., Spreng, R. N., & Andrews-Hanna, J. R. (2016). Mind-wandering as

spontaneous thought: a dynamic framework. *Nature Reviews. Neuroscience*, *17*(11), 718–731.

Coifman, R. R., Lafon, S., Lee, A. B., Maggioni, M., Nadler, B., Warner, F., & Zucker, S. W. (2005). Geometric

diffusions as a tool for harmonic analysis and structure definition of data: diffusion maps. *Proceedings of*

the National Academy of Sciences of the United States of America, *102*(21), 7426–7431.

Cruces, R. R., Royer, J., Herholz, P., Larivière, S., Vos de Wael, R., Paquola, C., Benkarim, O., Park, B.-Y., Degré-

Pelletier, J., Nelson, M. C., DeKraker, J., Leppert, I. R., Tardif, C., Poline, J.-B., Concha, L., & Bernhardt,

B. C. (2022). Micapipe: A pipeline for multimodal neuroimaging and connectome analysis. *NeuroImage*,

263, 119612.

Dale, A. M., Fischl, B., & Sereno, M. I. (1999). Cortical surface-based analysis. I. Segmentation and surface

reconstruction. *NeuroImage*, *9*(2), 179–194.

Dammers, J., Breuer, L., Axer, M., Kleiner, M., Eiben, B., Grässel, D., Dickscheid, T., Zilles, K., Amunts, K., Shah,

N. J., & Pietrzyk, U. (2012). Automatic identification of gray and white matter components in polarized

light imaging. *NeuroImage*, *59*(2), 1338–1347.

Damoiseaux, J. S., & Greicius, M. D. (2009). Greater than the sum of its parts: a review of studies combining

structural connectivity and resting-state functional connectivity. *Brain Structure & Function*, *213*(6), 525–

533.

DeKraker, J., Lau, J. C., Ferko, K. M., Khan, A. R., & Köhler, S. (2020). Hippocampal subfields revealed through

unfolding and unsupervised clustering of laminar and morphological features in 3D BigBrain. *NeuroImage*,

206, 116328.

DeKraker, Jordan, Ferko, K. M., Lau, J. C., Köhler, S., & Khan, A. R. (2018). Unfolding the hippocampus: An

intrinsic coordinate system for subfield segmentations and quantitative mapping. *NeuroImage*, *167*, 408–

418.

DeKraker, Jordan, Haast, R. A. M., Yousif, M. D., Karat, B., Lau, J. C., Köhler, S., & Khan, A. R. (2022).

Automated hippocampal unfolding for morphometry and subfield segmentation with HippUnfold. *ELife*,

11. https://doi.org/10.7554/eLife.77945

DeKraker, Jordan, Köhler, S., & Khan, A. R. (2021). Surface-based hippocampal subfield segmentation. *Trends in*

Neurosciences, *44*(11), 856–863.

DeKraker, Jordan, Palomero-Gallagher, N., Kedo, O., Ladbon-Bernasconi, N., Muenzing, S. E. A., Axer, M.,

Amunts, K., Khan, A. R., Bernhardt, B. C., & Evans, A. C. (2023). Evaluation of surface-based

hippocampal registration using ground-truth subfield definitions. *ELife*, *12*.

https://doi.org/10.7554/eLife.88404

Ding, S.-L., & Van Hoesen, G. W. (2015). Organization and detailed parcellation of human hippocampal head and

body regions based on a combined analysis of cyto- and chemoarchitecture. *The Journal of Comparative*

Neurology, *523*(15), 2233–2253.

Drouin, S., Kochanowska, A., Kersten-Oertel, M., Gerard, I. J., Zelmann, R., De Nigris, D., Bériault, S., Arbel, T.,

Sirhan, D., Sadikot, A. F., Hall, J. A., Sinclair, D. S., Petrecca, K., DelMaestro, R. F., & Collins, D. L.

(2017). IBIS: an OR ready open-source platform for image-guided neurosurgery. *International Journal of*

Computer Assisted Radiology and Surgery, *12*(3), 363–378.

Duncan, J. (2010). The multiple-demand (MD) system of the primate brain: mental programs for intelligent

behaviour. *Trends in Cognitive Sciences*, *14*(4), 172–179.

Duvernoy, H. M., Cattin, F., & Risold, P.-Y. (2013). *The Human Hippocampus*. Springer Berlin Heidelberg.

Duyn, J. H. (2012). The future of ultra-high field MRI and fMRI for study of the human brain. *NeuroImage*, *62*(2),

1241–1248.

Eichenbaum, H. (2000). A cortical–hippocampal system for declarative memory. *Nature Reviews. Neuroscience*,

1(1), 41–50.

Eichert, N., DeKraker, J., Howard, A. F. D., Huszar, I. N., Zhu, S., Sallet, J., Miller, K. L., Mars, R. B., Jbabdi, S., &

Bernhardt, B. C. (2023). Hippocampal connectivity patterns echo macroscale cortical evolution in the

primate brain. In *bioRxiv* (p. 2023.09.08.556859). https://doi.org/10.1101/2023.09.08.556859

Eickhoff, S. B., Yeo, B. T. T., & Genon, S. (2018). Imaging-based parcellations of the human brain. *Nature*

Reviews. Neuroscience, *19*(11), 672–686.

Fallon, J., Ward, P. G. D., Parkes, L., Oldham, S., Arnatkevičiūtė, A., Fornito, A., & Fulcher, B. D. (2020).

Timescales of spontaneous fMRI fluctuations relate to structural connectivity in the brain. *Network*

Neuroscience (Cambridge, Mass.), *4*(3), 788–806.

Fischl, B., Sereno, M. I., & Dale, A. M. (1999). Cortical surface-based analysis. II: Inflation, flattening, and a

surface-based coordinate system. *NeuroImage*, *9*(2), 195–207.

Fischl, B., Sereno, M. I., Tootell, R. B., & Dale, A. M. (1999). High-resolution intersubject averaging and a

coordinate system for the cortical surface. *Human Brain Mapping*, *8*(4), 272–284.

Fox, K. C. R., Spreng, R. N., Ellamil, M., Andrews-Hanna, J. R., & Christoff, K. (2015). The wandering brain:
meta-analysis of functional neuroimaging studies of mind-wandering and related spontaneous thought
processes. *NeuroImage*, *111*, 611–621.

Frauscher, B., von Ellenrieder, N., Zelmann, R., Doležalová, I., Minotti, L., Olivier, A., Hall, J., Hoffmann, D.,
Nguyen, D. K., Kahane, P., Dubeau, F., & Gotman, J. (2018). Atlas of the normal intracranial
electroencephalogram: neurophysiological awake activity in different cortical areas. *Brain: A Journal of*
Neurology, *141*(4), 1130–1144.

Gandolfi, D., Mapelli, J., Solinas, S. M. G., Triebkorn, P., D'Angelo, E., Jirsa, V., & Migliore, M. (2023). Full-scale
scaffold model of the human hippocampus CA1 area. *Nature Computational Science*, *3*(3), 264–276.

Genon, S., Bernhardt, B. C., La Joie, R., Amunts, K., & Eickhoff, S. B. (2021). The many dimensions of human
hippocampal organization and (dys)function. *Trends in Neurosciences*, *44*(12), 977–989.

Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L., Xu, J., Jbabdi, S.,
Webster, M., Polimeni, J. R., Van Essen, D. C., Jenkinson, M., & WU-Minn HCP Consortium. (2013). The
minimal preprocessing pipelines for the Human Connectome Project. *NeuroImage*, *80*, 105–124.

Golesorkhi, M., Gomez-Pilar, J., Zilio, F., Berberian, N., Wolff, A., Yagoub, M. C. E., & Northoff, G. (2021). The
brain and its time: intrinsic neural timescales are key for input processing. *Communications Biology*, *4*(1),
970.

González-Arnay, E., Pérez-Santos, I., Jiménez-Sánchez, L., Cid, E., Gal, B., de la Prida, L. M., & Cavada, C.
(2024). Immunohistochemical field parcellation of the human hippocampus along its antero-posterior axis.
Brain Structure & Function. https://doi.org/10.1007/s00429-023-02725-9

Gorgolewski, K. J., Auer, T., Calhoun, V. D., Craddock, R. C., Das, S., Duff, E. P., Flandin, G., Ghosh, S. S.,
Glatard, T., Halchenko, Y. O., Handwerker, D. A., Hanke, M., Keator, D., Li, X., Michael, Z., Maumet, C.,
Nichols, B. N., Nichols, T. E., Pellman, J., … Poldrack, R. A. (2016). The brain imaging data structure, a
format for organizing and describing outputs of neuroimaging experiments. *Scientific Data*, *3*, 160044.

Gorgolewski, K. J., & Poldrack, R. A. (2016). A Practical Guide for Improving Transparency and Reproducibility in
Neuroimaging Research. *PLoS Biology*, *14*(7), e1002506.

Gorgolewski, K. J., Varoquaux, G., Rivera, G., Schwarz, Y., Ghosh, S. S., Maumet, C., Sochat, V. V., Nichols, T.
E., Poldrack, R. A., Poline, J.-B., Yarkoni, T., & Margulies, D. S. (2015). Neurovault.org: a web-based

repository for collecting and sharing unthresholded statistical maps of the human brain. *Frontiers in*

Neuroinformatics, *9*, 8.

Greicius, M. D., Supekar, K., Menon, V., & Dougherty, R. F. (2009). Resting-state functional connectivity reflects

structural connectivity in the default mode network. *Cerebral Cortex* , *19*(1), 72–78.

Guzman, S. J., Schlögl, A., Frotscher, M., & Jonas, P. (2016). Synaptic mechanisms of pattern completion in the

hippocampal CA3 network. *Science*, *353*(6304), 1117–1123.

Haast, R. A. M., Ivanov, D., Formisano, E., & Uluda☐, K. (2016). Reproducibility and Reliability of Quantitative

and Weighted T1 and T2∗ Mapping for Myelin-Based Cortical Parcellation at 7 Tesla. *Frontiers in*

Neuroanatomy, *10*, 112.

Haast, R. A. M., Kashyap, S., Ivanov, D., Yousif, M. D., DeKraker, J., Poser, B. A., & Khan, A. R. (2023). Novel

insights into hippocampal perfusion using high-resolution, multi-modal 7T MRI. *BioRxiv : The Preprint*

Server for Biology. https://doi.org/10.1101/2023.07.19.549533

Hanson, J. L., Adkins, D. J., Nacewicz, B. M., & Barry, K. R. (2023). Impact of Socioeconomic Status on Amygdala

and Hippocampus Subdivisions in Children and Adolescents. *BioRxiv : The Preprint Server for Biology*.

https://doi.org/10.1101/2023.03.10.532071

Hebb, D. O. (2005). *The Organization of Behavior: A Neuropsychological Theory*. Psychology Press.

Henriksen, E. J., Colgin, L. L., Barnes, C. A., Witter, M. P., Moser, M.-B., & Moser, E. I. (2010). Spatial

representation along the proximodistal axis of CA1. *Neuron*, *68*(1), 127–137.

Honey, C. J., Sporns, O., Cammoun, L., Gigandet, X., Thiran, J. P., Meuli, R., & Hagmann, P. (2009). Predicting

human resting-state functional connectivity from structural connectivity. *Proceedings of the National*

Academy of Sciences of the United States of America, *106*(6), 2035–2040.

Huntenburg, J., Abraham, A., Loula, J., Liem, F., Dadi, K., & Varoquaux, G. (2017). Loading and plotting of

cortical surface representations in Nilearn. *Research Ideas and Outcomes*, *3*, e12342.

Igarashi, K. M., Ito, H. T., Moser, E. I., & Moser, M.-B. (2014). Functional diversity along the transverse axis of

hippocampal area CA1. *FEBS Letters*, *588*(15), 2470–2476.

Iglesias, J. E., Augustinack, J. C., Nguyen, K., Player, C. M., Player, A., Wright, M., Roy, N., Frosch, M. P.,

McKee, A. C., Wald, L. L., Fischl, B., Van Leemput, K., & Alzheimer's Disease Neuroimaging Initiative.

(2015). A computational atlas of the hippocampal formation using ex vivo, ultra-high resolution MRI:

Application to adaptive segmentation of in vivo MRI. *NeuroImage*, *115*, 117–137.

Insausti, R., & Amaral, D. G. (2004). Hippocampal formation. *The Human Nervous System*.

https://www.researchgate.net/profile/Ricardo-Insausti-

2/publication/279431811_Hippocampal_Formation/links/5f05ac7c92851c52d6208202/Hippocampal-

Formation.pdf

Karat, B. G., DeKraker, J., Hussain, U., Köhler, S., & Khan, A. R. (2023). Mapping the macrostructure and

microstructure of the in vivo human hippocampus using diffusion MRI. *Human Brain Mapping*, *44*(16),

5485–5503.

Kim, J. S., Singh, V., Lee, J. K., Lerch, J., Ad-Dab'bagh, Y., MacDonald, D., Lee, J. M., Kim, S. I., & Evans, A. C.

(2005). Automated 3-D extraction and evaluation of the inner and outer cortical surfaces using a Laplacian

map and partial volume effect classification. *NeuroImage*, *27*(1), 210–221.

Knierim, J. J., & Neunuebel, J. P. (2016). Tracking the flow of hippocampal computation: Pattern separation, pattern

completion, and attractor dynamics. *Neurobiology of Learning and Memory*, *129*, 38–49.

Kulaga-Yoskovitz, J., Bernhardt, B. C., Hong, S.-J., Mansi, T., Liang, K. E., van der Kouwe, A. J. W., Smallwood,

J., Bernasconi, A., & Bernasconi, N. (2015). Multi-contrast submillimetric 3 Tesla hippocampal subfield

segmentation protocol and dataset. *Scientific Data*, *2*, 150059.

Lacy, J. W., Yassa, M. A., Stark, S. M., Muftuler, L. T., & Stark, C. E. L. (2011). Distinct pattern separation related

transfer functions in human CA3/dentate and CA1 revealed using high-resolution fMRI and variable

mnemonic similarity. *Learning & Memory* , *18*(1), 15–18.

Larivière, S., Bayrak, Ş., Vos de Wael, R., Benkarim, O., Herholz, P., Rodriguez-Cruces, R., Paquola, C., Hong, S.-

J., Misic, B., Evans, A. C., Valk, S. L., & Bernhardt, B. C. (2023). BrainStat: A toolbox for brain-wide

statistics and multimodal feature associations. *NeuroImage*, *266*, 119807.

Larivière, S., Paquola, C., Park, B.-Y., Royer, J., Wang, Y., Benkarim, O., Vos de Wael, R., Valk, S. L.,

Thomopoulos, S. I., Kirschner, M., Lewis, L. B., Evans, A. C., Sisodiya, S. M., McDonald, C. R.,

Thompson, P. M., & Bernhardt, B. C. (2021). The ENIGMA Toolbox: multiscale neural contextualization

of multisite neuroimaging datasets. *Nature Methods*, *18*(7), 698–700.

Lariviere, S., Park, B.-Y., Royer, J., DeKraker, J., Ngo, A., Sahlas, E., Chen, J., Rodriguez-Cruces, R., Weng, Y.,

Frauscher, B., & Others. (2023). Connectome reorganization associated with temporal lobe pathology and

its surgical resection. *MedRxiv*, 2023–2011.

Larkin, M. C., Lykken, C., Tye, L. D., Wickelgren, J. G., & Frank, L. M. (2014). Hippocampal output area CA1

broadcasts a generalized novelty signal during an object-place recognition task. *Hippocampus*, *24*(7), 773–

783.

Leferink, C. A., DeKraker, J., Brunec, I. K., Köhler, S., Moscovitch, M., & Walther, D. B. (2023). Organization of

pRF size along the AP axis of the hippocampus and adjacent medial temporal cortex is related to

specialization for scenes versus faces. *Cerebral Cortex* . https://doi.org/10.1093/cercor/bhad429

Lepage, C., Lewis, L., Jeun, S., Bermudez, P., Khalili-Mahani, N., Omidyegaheh, M., Zijdenbos, A., Vincent, R. D.,

Adalat, R., & Evans, A. C. (2017). Human MR evaluation of cortical thickness using CIVET v2. 1.

Organization   for   Human   Brain   Mapping.

https://archive.aievolution.com/2017/hbm1701/index.cfm?do=abs.viewAbs&abs=3292

Leutgeb, J. K., Leutgeb, S., Moser, M.-B., & Moser, E. I. (2007). Pattern separation in the dentate gyrus and CA3 of

the hippocampus. *Science*, *315*(5814), 961–966.

Leutgeb, S., & Leutgeb, J. K. (2007). Pattern separation, pattern completion, and new neuronal codes within a

continuous CA3 map. *Learning & Memory* , *14*(11), 745–757.

Ly, M., Foley, L., Manivannan, A., Hitchens, T. K., Richardson, R. M., & Modo, M. (2020). Mesoscale diffusion

magnetic resonance imaging of the ex vivo human hippocampus. *Human Brain Mapping*, *41*(15), 4200–

4218.

Lyttelton, O., Boucher, M., Robbins, S., & Evans, A. (2007). An unbiased iterative group registration template for

cortical surface analysis. *NeuroImage*, *34*(4), 1535–1544.

Ma, Q., Li, L., Robinson, E. C., Kainz, B., Rueckert, D., & Alansary, A. (2023). CortexODE: Learning Cortical

Surface Reconstruction by Neural ODEs. *IEEE Transactions on Medical Imaging*, *42*(2), 430–443.

MacDonald, D., Kabani, N., Avis, D., & Evans, A. C. (2000). Automated 3-D extraction of inner and outer surfaces

of cerebral cortex from MRI. *NeuroImage*, *12*(3), 340–356.

Marcus, D. S., Harwell, J., Olsen, T., Hodge, M., Glasser, M. F., Prior, F., Jenkinson, M., Laumann, T., Curtiss, S.

W., & Van Essen, D. C. (2011). Informatics and data mining tools and strategies for the human connectome

project. *Frontiers in Neuroinformatics*, *5*, 4.

Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., Bezgin, G., Eickhoff, S.

B., Castellanos, F. X., Petrides, M., Jefferies, E., & Smallwood, J. (2016). Situating the default-mode

network along a principal gradient of macroscale cortical organization. *Proceedings of the National*

Academy of Sciences of the United States of America, *113*(44), 12574–12579.

Markello, R. D., Hansen, J. Y., Liu, Z.-Q., Bazinet, V., Shafiei, G., Suárez, L. E., Blostein, N., Seidlitz, J., Baillet,

S., Satterthwaite, T. D., Chakravarty, M. M., Raznahan, A., & Misic, B. (2022). neuromaps: structural and

functional interpretation of brain maps. *Nature Methods*, *19*(11), 1472–1479.

Mountcastle, V. B. (1997). The columnar organization of the neocortex. *Brain: A Journal of Neurology*, *120 (Pt 4)*,

701–722.

Ngo, A., Royer, J., Rodríguez-Cruces, R., Xie, K., DeKraker, J., Auer, H., Tavakol, S., Lam, J., Schrader, D.,

Dudley, R. W. R., Bernasconi, A., Bernasconi, N., Frauscher, B., Larivière, S., & Bernhardt, B. C. (2023).

Cerebral perfusion alterations in temporal lobe epilepsy: Structural underpinnings and network disruptions.

In *bioRxiv* (p. 2023.08.22.553552). https://doi.org/10.1101/2023.08.22.553552

Norman, Y., Raccah, O., Liu, S., Parvizi, J., & Malach, R. (2021). Hippocampal ripples and their coordinated

dialogue with the default mode network during recent and remote recollection. *Neuron*, *109*(17), 2767-

2780.e5.

O'Keefe, J., & Nadel, L. (1978). *The Hippocampus as a Cognitive Map*. Clarendon Press.

Olsen, R. K., Carr, V. A., Daugherty, A. M., La Joie, R., Amaral, R. S. C., Amunts, K., Augustinack, J. C., Bakker,

A., Bender, A. R., Berron, D., Boccardi, M., Bocchetta, M., Burggren, A. C., Chakravarty, M. M., Chételat,

G., de Flores, R., DeKraker, J., Ding, S.-L., Geerlings, M. I., … Hippocampal Subfields Group. (2019).

Progress update from the hippocampal subfields group. *Alzheimer's & Dementia: The Journal of the*

Alzheimer's Association, *11*, 439–449.

Opheim, G., van der Kolk, A., Markenroth Bloch, K., Colon, A. J., Davis, K. A., Henry, T. R., Jansen, J. F. A.,

Jones, S. E., Pan, J. W., Rössler, K., Stein, J. M., Strandberg, M. C., Trattnig, S., Van de Moortele, P.-F.,

Vargas, M. I., Wang, I., Bartolomei, F., Bernasconi, N., Bernasconi, A., … Guye, M. (2021). 7T Epilepsy

Task Force Consensus Recommendations on the Use of 7T MRI in Clinical Practice. *Neurology*, *96*(7),

327–341.

Paquola, C., Benkarim, O., DeKraker, J., Larivière, S., Frässle, S., Royer, J., Tavakol, S., Valk, S., Bernasconi, A.,

Bernasconi, N., Khan, A., Evans, A. C., Razi, A., Smallwood, J., & Bernhardt, B. C. (2020). Convergence

of cortical types and functional motifs in the human mesiotemporal lobe. *ELife*, *9*.

https://doi.org/10.7554/eLife.60673

Paquola, C., Seidlitz, J., Benkarim, O., Royer, J., Klimes, P., Bethlehem, R. A. I., Larivière, S., Vos de Wael, R.,

Rodríguez-Cruces, R., Hall, J. A., Frauscher, B., Smallwood, J., & Bernhardt, B. C. (2020). A multi-scale

cortical wiring space links cellular architecture and functional dynamics in the human brain. *PLoS Biology*,

18(11), e3000979.

Paquola, C., Vos De Wael, R., Wagstyl, K., Bethlehem, R. A. I., Hong, S.-J., Seidlitz, J., Bullmore, E. T., Evans, A.

C., Misic, B., Margulies, D. S., Smallwood, J., & Bernhardt, B. C. (2019). Microstructural and functional

gradients are increasingly dissociated in transmodal cortices. *PLoS Biology*, *17*(5), e3000284.

Park, B.-Y., Vos de Wael, R., Paquola, C., Larivière, S., Benkarim, O., Royer, J., Tavakol, S., Cruces, R. R., Li, Q.,

Valk, S. L., Margulies, D. S., Mišić, B., Bzdok, D., Smallwood, J., & Bernhardt, B. C. (2021). Signal

diffusion along connectome gradients and inter-hub routing differentially contribute to dynamic human

brain function. *NeuroImage*, *224*, 117429.

Pishdadian, S., Hoang, N. V., Baker, S., Moscovitch, M., & Rosenbaum, R. S. (2020). Not only memory:

Investigating the sensitivity and specificity of the Mnemonic Similarity Task in older adults.

Neuropsychologia, *149*, 107670.

Poppenk, J., Evensmoen, H. R., Moscovitch, M., & Nadel, L. (2013). Long-axis specialization of the human

hippocampus. *Trends in Cognitive Sciences*, *17*(5), 230–240.

Przeździk, I., Faber, M., Fernández, G., Beckmann, C. F., & Haak, K. V. (2019). The functional organisation of the

hippocampus along its long axis is gradual and predicts recollection. *Cortex; a Journal Devoted to the*

Study of the Nervous System and Behavior, *119*, 324–335.

Puelles, L., Alonso, A., García-Calero, E., & Martínez-de-la-Torre, M. (2019). Concentric ring topology of

mammalian cortical sectors and relevance for patterning studies. *The Journal of Comparative Neurology*,

527(10), 1731–1752.

Ramón y Cajal, S. (1904). Textura del Sistema Nervioso del Hombre y de los Vertebrados, tomo II, primera parte.

Imprenta y Libreria de Nicolas Moya, Madrid, Reprinted.

Ripart, M., DeKraker, J., Eriksson, M. H., Piper, R. J., Mo, J.-J., Su, T.-Y., Kochi, R., Wang, I., Winston, G. P.,

Clark, C. A., D'Arco, F., Mankad, K., Khan, A. R., Baldeweg, T., Adler, S., & Wagstyl, K. (2023).

Automated and Interpretable Detection of Hippocampal Sclerosis in temporal lobe epilepsy: AID-HS. In

820   *bioRxiv*. https://doi.org/10.1101/2023.10.13.23296991

821   Robinson, E. C., Jbabdi, S., Glasser, M. F., Andersson, J., Burgess, G. C., Harms, M. P., Smith, S. M., Van Essen,

822   D. C., & Jenkinson, M. (2014). MSM: a new flexible framework for Multimodal Surface Matching.

823   *NeuroImage*, *100*, 414–426.

824   Rolls, E. T. (2016). Pattern separation, completion, and categorisation in the hippocampus and neocortex.

825   *Neurobiology of Learning and Memory*, *129*, 4–28.

826   Romero, J. E., Coupé, P., & Manjón, J. V. (2017). HIPS: A new hippocampus subfield segmentation method.

827   *NeuroImage*, *163*, 286–295.

828   Ross, T. W., & Easton, A. (2022). The Hippocampal Horizon: Constructing and Segmenting Experience for

829   Episodic Memory. *Neuroscience and Biobehavioral Reviews*, *132*, 181–196.

830   Royer, J., Rodríguez-Cruces, R., Tavakol, S., Larivière, S., Herholz, P., Li, Q., Vos de Wael, R., Paquola, C.,

831   Benkarim, O., Park, B.-Y., Lowe, A. J., Margulies, D., Smallwood, J., Bernasconi, A., Bernasconi, N.,

832   Frauscher, B., & Bernhardt, B. C. (2022). An Open MRI Dataset For Multiscale Neuroscience. *Scientific

833   Data*, *9*(1), 569.

834   Sanides, F. (1969). Comparative architectonics of the neocortex of mammals and their evolutionary interpretation.

835   *Annals of the New York Academy of Sciences*, *167*(1), 404–423.

836   Schacter, D. L., Benoit, R. G., & Szpunar, K. K. (2017). Episodic Future Thinking: Mechanisms and Functions.

837   *Current Opinion in Behavioral Sciences*, *17*, 41–50.

838   Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T.

839   (2018). Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity

840   MRI. *Cerebral Cortex* , *28*(9), 3095–3114.

841   Schleicher, A., Amunts, K., Geyer, S., Morosan, P., & Zilles, K. (1999). Observer-independent method for

842   microstructural parcellation of cerebral cortex: A quantitative approach to cytoarchitectonics. *NeuroImage*,

843   *9*(1), 165–177.

844   Schmidt, B., Marrone, D. F., & Markus, E. J. (2012). Disambiguating the similar: the dentate gyrus and pattern

845   separation. *Behavioural Brain Research*, *226*(1), 56–65.

Schmittmann, V. D., Jahfari, S., Borsboom, D., Savi, A. O., & Waldorp, L. J. (2015). Making Large-Scale Networks

from fMRI Data. *PloS One*, *10*(9), e0129074.

Smallwood, J., Bernhardt, B. C., Leech, R., Bzdok, D., Jefferies, E., & Margulies, D. S. (2021). The default mode

network in cognition: a topographical perspective. *Nature Reviews. Neuroscience*, *22*(8), 503–513.

Smith, S. M., Fox, P. T., Miller, K. L., Glahn, D. C., Fox, P. M., Mackay, C. E., Filippini, N., Watkins, K. E., Toro,

R., Laird, A. R., & Beckmann, C. F. (2009). Correspondence of the brain's functional architecture during

activation and rest. *Proceedings of the National Academy of Sciences of the United States of America*,

106(31), 13040–13045.

Stachenfeld, K. L., Botvinick, M., & Gershman, S. J. (2014). Design principles of the hippocampal cognitive map.

Advances   in   Neural   Information   Processing   Systems,   *27*.

https://proceedings.neurips.cc/paper_files/paper/2014/hash/dfd7468ac613286cdbb40872c8ef3b06-

Abstract.html

Stachenfeld, K. L., Botvinick, M. M., & Gershman, S. J. (2017). The hippocampus as a predictive map. *Nature*

Neuroscience, *20*(11), 1643–1653.

Stark, S. M., Kirwan, C. B., & Stark, C. E. L. (2019). Mnemonic Similarity Task: A Tool for Assessing

Hippocampal Integrity. *Trends in Cognitive Sciences*, *23*(11), 938–951.

Steve, T. A., Gargula, J., Misaghi, E., Nowacki, T. A., Schmitt, L. M., Wheatley, B. M., & Gross, D. W. (2020).

Hippocampal subfield measurement and ILAE hippocampal sclerosis subtype classification with in vivo

4.7 tesla MRI. *Epilepsy Research*, *161*, 106279.

Strange, B. A., Witter, M. P., Lein, E. S., & Moser, E. I. (2014). Functional organization of the hippocampal

longitudinal axis. *Nature Reviews. Neuroscience*, *15*(10), 655–669.

Tournier, J.-D., Calamante, F., & Connelly, A. (2012). MRtrix: Diffusion tractography in crossing fiber regions.

International Journal of Imaging Systems and Technology, *22*(1), 53–66.

van den Heuvel, M. P., & Sporns, O. (2013). Network hubs in the human brain. *Trends in Cognitive Sciences*,

17(12), 683–696.

van der Weijden, C. W. J., García, D. V., Borra, R. J. H., Thurner, P., Meilof, J. F., van Laar, P.-J., Dierckx, R. A. J.

O., Gutmann, I. W., & de Vries, E. F. J. (2021). Myelin quantification with MRI: A systematic review of

accuracy and reproducibility. *NeuroImage*, *226*, 117561.

Van Essen, D. C., Drury, H. A., Joshi, S., & Miller, M. I. (1998). Functional and structural mapping of human

cerebral cortex: solutions are in the surfaces. *Proceedings of the National Academy of Sciences of the*
United States of America, *95*(3), 788–795.

Van Essen, David C., Smith, J., Glasser, M. F., Elam, J., Donahue, C. J., Dierker, D. L., Reid, E. K., Coalson, T., &
Harwell, J. (2017). The Brain Analysis Library of Spatial maps and Atlases (BALSA) database.
NeuroImage, *144*(Pt B), 270–274.

Vogel, J. W., La Joie, R., Grothe, M. J., Diaz-Papkovich, A., Doyle, A., Vachon-Presseau, E., Lepage, C., Vos de
Wael, R., Thomas, R. A., Iturria-Medina, Y., Bernhardt, B., Rabinovici, G. D., & Evans, A. C. (2020). A
molecular gradient along the longitudinal axis of the human hippocampus informs large-scale behavioral
systems. *Nature Communications*, *11*(1), 960.

Vos de Wael, R., Benkarim, O., Paquola, C., Lariviere, S., Royer, J., Tavakol, S., Xu, T., Hong, S.-J., Langs, G.,
Valk, S., Misic, B., Milham, M., Margulies, D., Smallwood, J., & Bernhardt, B. C. (2020). BrainSpace: a
toolbox for the analysis of macroscale gradients in neuroimaging and connectomics datasets.
Communications Biology, *3*(1), 103.

Vos de Wael, R., Larivière, S., Caldairou, B., Hong, S.-J., Margulies, D. S., Jefferies, E., Bernasconi, A.,
Smallwood, J., Bernasconi, N., & Bernhardt, B. C. (2018). Anatomical and microstructural determinants of
hippocampal subfield functional connectome embedding. *Proceedings of the National Academy of Sciences*
of the United States of America, *115*(40), 10154–10159.

Wagner, H. H., & Dray, S. (2015). Generating spatially constrained null models for irregularly spaced data using
Moran spectral randomization methods. *Methods in Ecology and Evolution / British Ecological Society*,
6(10), 1169–1178.

Ward, A. M., Schultz, A. P., Huijbers, W., Van Dijk, K. R. A., Hedden, T., & Sperling, R. A. (2014). The
parahippocampal gyrus links the default-mode cortical network with the medial temporal lobe memory
system. *Human Brain Mapping*, *35*(3), 1061–1073.

Whittington, J. C. R., McCaffary, D., Bakermans, J. J. W., & Behrens, T. E. J. (2022). How to build a cognitive
map: insights from models of the hippocampal formation. In *arXiv [q-bio.NC]*. arXiv.
http://arxiv.org/abs/2202.01682

Whittington, J. C. R., Muller, T. H., Mark, S., Chen, G., Barry, C., Burgess, N., & Behrens, T. E. J. (2020). The
Tolman-Eichenbaum Machine: Unifying Space and Relational Memory through Generalization in the

Hippocampal Formation. *Cell*, *183*(5), 1249-1263.e23.

Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-

W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I.,

Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., … Mons, B. (2016). The FAIR Guiding Principles for

scientific data management and stewardship. *Scientific Data*, *3*, 160018.

Wolff, A., Berberian, N., Golesorkhi, M., Gomez-Pilar, J., Zilio, F., & Northoff, G. (2022). Intrinsic neural

timescales: temporal integration and segregation. *Trends in Cognitive Sciences*, *26*(2), 159–173.

Xie, K., Royer, J., Larivière, S., Rodriguez-Cruces, R., Frässle, S., Cabalo, D. G., Ngo, A., DeKraker, J., Auer, H.,

Tavakol, S., Weng, Y., Abdallah, C., Horwood, L., Frauscher, B., Caciagli, L., Bernasconi, A., Bernasconi,

N., Zhang, Z., Concha, L., & Bernhardt, B. C. (2023). Atypical connectome topography and signal flow in

temporal lobe epilepsy. *BioRxiv : The Preprint Server for Biology*.

https://doi.org/10.1101/2023.05.23.541934

Yang, Y., Chen, Z., Zhang, R., Xu, T., & Feng, T. (2020). Neural substrates underlying episodic future thinking: A

voxel-based morphometry study. *Neuropsychologia*, *138*, 107255.

Yushkevich, P. A., Amaral, R. S. C., Augustinack, J. C., Bender, A. R., Bernstein, J. D., Boccardi, M., Bocchetta,

M., Burggren, A. C., Carr, V. A., Chakravarty, M. M., Chételat, G., Daugherty, A. M., Davachi, L., Ding,

S.-L., Ekstrom, A., Geerlings, M. I., Hassan, A., Huang, Y., Iglesias, J. E., … Hippocampal Subfields

Group (HSG). (2015). Quantitative comparison of 21 protocols for labeling hippocampal subfields and

parahippocampal subregions in in vivo MRI: towards a harmonized segmentation protocol. *NeuroImage*,

111, 526–541.

Yushkevich, P. A., Wang, H., Pluta, J., Das, S. R., Craige, C., Avants, B. B., Weiner, M. W., & Mueller, S. (2010).

Nearly automatic segmentation of hippocampal subfields in in vivo focal T2-weighted MRI. *NeuroImage*,

53(4), 1208–1224.

Zang, Y., Jiang, T., Lu, Y., He, Y., & Tian, L. (2004). Regional homogeneity approach to fMRI data analysis.

NeuroImage, *22*(1), 394–400.