***Agrobacterium*-mediated *Cuscuta campestris* transformation as a tool for understanding**

plant-plant interactions

Supral Adhikari[1,†], Asha Mudalige[1,†], Lydia Phillips[1], Hyeyoung Lee[2], Vivian Bernal-Galeano[3],

Hope Gruszewski[3], James H. Westwood[3], So-Yon Park*[1]

[1] Division of Plant Science and Technology, University of Missouri, Columbia, MO, 65211,

USA

[2] Division of Animal Science, University of Missouri, Columbia, MO, 65211, USA

[3] School of Plant and Environmental Sciences, Virginia Tech, Blacksburg, VA, 24061, USA

*Author for correspondence: parksoy@missouri.edu

[†] These authors contributed equally to this article.

ORCIDs:

Supral Adhikari https://orcid.org/0000-0002-9914-2986

Asha Mudalige https://orcid.org/0009-0002-2842-1758

Lydia Phillips

Hyeyoung Lee

Vivian Bernal-Galeano https://orcid.org/0000-0002-0438-2937

Hope Gruszewski

James H. Westwood https://orcid.org/0000-0001-5127-7294

So-Yon Park https://orcid.org/0000-0002-2126-3382

Summary

Cuscuta campestris, a stem parasitic plant, has served as a valuable model plant for the exploration of plant-plant interactions and molecular trafficking. However, a major barrier to *C. campestris* research is that a method to generate stable transgenic plants has not yet been developed. Here, we describe the development of a *Cuscuta* transformation protocol using various reporter genes (GFP, GUS, or RUBY) and morphogenic genes (*CcWUS2* and *CcGRF/GIF*), ultimately leading to a robust protocol for *Agrobacterium*-mediated *C. campestris* transformation. The stably transformed and regenerated RUBY *C. campestris* plants produced haustoria, the signature organ of parasitic plants, and these were functional in forming host attachments. Transformed *C. campestris* also produced flowers and transgenic seeds exhibiting betalain pigment, providing proof of germline transmission of the RUBY transgene. The integration locations of T-DNAs were confirmed through TAIL-PCR. Furthermore, the RUBY reporter is not only a useful selectable marker for the *Agrobacterium*-mediated transformation but also provides insight into the movement of molecules from *C. campestris* to the host during parasitism. Thus, the protocol to generate transgenic *C. campestris* reported here overcomes a major obstacle to *Cuscuta* research and opens up new possibilities for studying parasitic plant interactions.

Key words: *Cuscuta*, parasitic plant, *Agrobacterium*-mediated transformation, plant-plant interactions, RUBY, mobile molecules.

Introduction

Cuscuta spp. are obligate stem parasitic plants that infect a broad range of host plants (Dawson *et al.*, 1994). The thread-like plant consists of leafless, twining stems that can attach to all aerial parts of herbaceous plants, woody shrubs, and even trees. The parasitic invasion of a *Cuscuta* begins by coiling around the stem of its host plant and initiating points of growth toward the host. The epidermal cells of the adhesive disk secrete a bioadhesive substance that anchors the parasite to the host (Vaughn, 2002; Galloway *et al.*, 2020). Next, a multicellular feeding structure termed the haustorium develops from the parasite stem and breaches the host tissues using a combination of mechanical pressure and enzymatic digestion of host cell walls, ultimately resulting in vascular connections with the host (Nagar *et al.*, 1984; Johnsen *et al.*, 2015; Kaiser *et al.*, 2015).

Cuscuta spp. are among the most agriculturally destructive parasitic weeds, as well as interesting subjects for studies of plant-plant interaction. The economic impact is due to their wide range of potential host species, which includes many dicotyledonous crops and ornamental plants (Lanini & Kogan, 2005), as well as their habit of extracting water and nutrients that the host needs for productivity. For example, *C. campestris* reduced tomato fresh weight by 98% in a greenhouse study (Goldwasser *et al.*, 2012), and *C. indecora* decreased alfalfa yield by 57% (Cudney *et al.*, 1992). Although systematic, chemical, and mechanical control methods have been developed to suppress the growth of *Cuscuta*, there is currently no efficient method to eliminate established *Cuscuta* without also causing damage to their hosts (Dechassa & Regassa, 2021).

Cuscuta spp. are useful for plant biology because their parasitic lifestyle provides insights into the mechanisms by which plants detect and interact with other plants. Examples of this include *Cuscuta* detection of neighboring plants by volatile signals (Runyon *et al.*, 2006), *Cuscuta*-to-host bidirectional mobility of mRNAs, small RNAs, and proteins (Kim *et al.*, 2014; Shahid *et al.*, 2018; Liu *et al.*, 2020; Park *et al.*, 2022), and elucidation of host resistance mechanisms against parasites (Hegenauer *et al.*, 2016; Hegenauer *et al.*, 2020). Therefore, both applied and basic plant sciences would benefit from greater ability to investigate the molecular mechanisms underlying interactions between *Cuscuta* and its hosts.

Plant genetic transformation is an essential tool for functional genomic studies as it enables the delivery of foreign genes to generate transgenic plants such as knockout or overexpression

mutant lines. This approach has been used for selected species of parasitic plants in the family

Orobanchaceae to understand the mechanisms by which they develop and interact with their

hosts. Methods for the transient and stable transformation of these root parasitic plants have been

published for *Triphysaria versicolor, Phtheirospermum japonicum*, and *Phelipanche aegyptiaca*

(Tomilov *et al.*, 2007; Fernández-Aparicio *et al.*, 2011; Ishida *et al.*, 2011; Bandaranayake &

Yoder, 2018).

As for *Cuscuta* transformation, several efforts have led to partial success. Regeneration of

shoots from callus of *C. reflexa* (Srivastava & Dwivedi, 2001) has been reported, and

transformation was accomplished in cultured *C. trifolii* (Borsics *et al.*, 2002) and *C. europeae*

(Švubová & Blehová, 2013). Transformation of haustorial structures was recently accomplished

in *C. reflexa* (Lachner *et al.*, 2020). However, despite these attempts with diverse *Cuscuta*

species, the inability to regenerate a functional *Cuscuta* plant from transformed explant tissues

has blocked progress toward achieving stably transformed *Cuscuta*.

The failure to regenerate plants from transgenic callus is a common bottleneck in plant

transformation, but progress has come from the use of genes involved in morphogenesis, which

promote the formation of plant organs in callus of plant species that are recalcitrant to

transformation or regeneration (Lee & Wang, 2023). For example, recent reports have shown

improved regeneration and transformation efficiency with the representative morphogenic

transcription factors such as *WUSCHEL* (*WUS*) and *BABY BOOM* (*BBM*). *WUS*, previously

known to play a vital role in maintaining the stem cell population, was also found to enhance

embryogenesis in *Arabidopsis* (Zuo *et al.*, 2002). When used with a transformation recalcitrant

maize line, *BBM-WUS* enhanced transformation efficiency by over 40% (Lowe *et al.*, 2016). In

another approach, a chimera consisting of *GROWTH-REGULATING FACTOR* (*GRF*) and its co-

factor *GRF-INTERACTING FACTOR* (*GIF*) was shown to promote transformation and

regeneration efficiency in wheat and citrus (Debernardi *et al.*, 2020). *GRF* and *GIF* are

transcription factors regulating growth and development of plants, and their ability to enhance

the efficiency of transformation has been extended to maize, watermelon, and lettuce. The

combination of *GRF-GIF* with *BBM* improved the transformation efficiency of maize seedlings

from 26% to 37% (Chen *et al.*, 2022). Furthermore, in watermelon, overexpression of the *GRF-*

GIF chimeric gene resulted in higher transformation efficiency than control conditions (Feng *et*

al., 2021). The *GRF-GIF* chimera with mutated miRNA396 improved the regeneration and

shooting efficiency in lettuce (Bull *et al.*, 2023).

Here we describe a transformation protocol for *C. campestris* that explored several different

reporter genes and morphogenic genes. Although *Cuscuta* species have been broadly regarded as

recalcitrant to transformation, the technique readily generates transgenic *C. campestris* plants

that show normal coiling and haustorium formation, and ultimately produce flowers and seeds.

This approach unlocks new possibilities for conducting functional studies on *Cuscuta* and

facilitates investigations into understanding of plant-plant interactions.

Materials and methods

***Cuscuta* seed scarification and germination**

C. campestris seedlings were attached to 3-week-old beets (*Beta vulgaris*) and grown in a greenhouse with a temperature of 25°C - 30°C and day/night length of 16/8 hours. Three months after the inoculation, *Cuscuta* seeds were collected and cleaned. To promote germination, clean seeds were scarified as described previously (Bernal-Galeano *et al.*, 2022). Scarified seeds were rinsed with autoclaved water three times, stirred in 3% sodium hypochlorite solution for 15 minutes for the sterilization. The treated seeds were washed by autoclaved water five times and incubated at 4C° overnight. For seed germination, 100 *Cuscuta* seeds were placed on a Petri dish with a wet 3M paper and incubated in a 28C° incubator without light for two days. The Petri dish containing *Cuscuta* seedlings was transferred to a walk-in chamber (25C° and 16 light/8 dark light conditions) and incubated for one day.

***Agrobacterium*-mediated *Cuscuta* transformation**

To develop a proper *Cuscuta* transformation protocol by *Agrobacterium*-mediated, we conducted three different protocols, such as optimization of growth hormone concentration, comparison of inoculation methods, and stable generation of *Cuscuta* transgenic lines.

First protocol-*Optimization of growth hormone concentration*: Starting with 3-day-old seedlings (Figure 1A), the shoot apical meristem end and root were cut from each plant (Figure 1B), and submerged into liquid modified Murashige and Skoog (MMS) with *A. tumefaciens* EHA105 harboring the construct *35S:GUS-pMAS:GFP* (Figure 1C and Supp table 1). The hypocotyls were placed on MMS media containing 3 mg/L NAA and 1 mg/L BAP for co-cultivation at 15 C° over 7 days with long-day condition (Figure 1D and Supp table 1) (Srivastava & Dwivedi, 2001). Subsequently, the infected explants were transferred to MMS media with the antibiotic cefotaxime and various types and concentrations of cytokinin and auxin to further optimize the growth (Figure 1E).

Second protocol-*Optimization of growth hormone concentration*: Using a sterilized surgical blade, 3-day-old seedlings were chopped into small pieces to make around 5mm segments, which greatly facilitated the tissue preparation step and produce large number of transformable sections at once. Also, instead of a construct with GUS and GFP reporters, we used a construct containing *35S:RUBY* (He *et al.*, 2020). Transformed tissues expressing RUBY were counted around a month after inoculation.

Third protocol-*Stable generation of Cuscuta transgenic lines*: Four-day-old *Cuscuta*

seedlings were chopped into small pieces as described in the second protocol. After cocultivation

with *A. tumefaciens* solutions, *Cuscuta* seedlings were uniformly distributed and incubated on

co-cultivation media for five days at 25 C° with long-day condition (Supp table 2). After the co-

cultivation stage, infected seedlings were transferred to Callus Induction Media (CIM)

containing 5 mg/L BAP and 250 mg/L Cefotaxime (Supp table 2). For the GUS staining assay,

calli were harvested 30-35 days after infection. To generate stably transformed *Cuscuta*, calli

were incubated on CIM for at least 8 weeks, shoot induction media (SIM) for 8 weeks, and shoot

elongation media for 3-4 weeks (Supp table 2).

GUS staining and GFP observation

159 For the GUS staining, *Cuscuta* calli were incubated in X-Gluc solution for up to 24 hours and

160 destained in 95 % Ethanol (Park *et al.*, 2022). For GFP detection, explants were placed on Laxco

161 stereomicroscope (Fisher) and observed with blue excitation light (Nightsea, SFA-RB-GO).

Vibratome sectioning and imaging

Cuscuta haustoria expressing RUBY grown on Rutgers tomatoes or *Arabidopsis* Col-0 were

harvested and embedded in 5 % Agarose (Park *et al.*, 2022). Tissue sectioning was performed

using a Leica series Vibratome 3000 Plus. Longitudinal and transverse sections were cut with

thickness ranging from 250μm to 400μm, depending on the softness of the tissue. Sections were

stained by 0.02% toluidine blue o (Pradhan Mitra & Loqué, 2014). For imaging, sections were

placed on microscope slides and imaged by Leica M205 FA automated stereo microscope.

Thermal asymmetric interlaced (TAIL)-PCR

To confirm T-DNA insertions in transgenic *Cuscuta*, genomic DNA was extracted using

Cetyltrimethylammonium-bromide (CTAB) (Sigma), and TAIL PCRs were conducted using

insertion specific primers (RB1 and RB2) and arbitrary degenerate primers (AD primers) (Supp

tables 3 and 4). Specific bands from amplification with AD and RB2 primers were extracted and

sequenced. The presence of T-DNA insertions in transgenic RUBY *Cuscuta* plants was verified

by aligning the sequencing results with a *Cuscuta campestris* genome sequence (Vogel et al.,

2018).

Vector constructions

35S:GUS-pMAS:GFP was modified by replacing the 35S promoter with the MAS promoter

from *pGFPGUSPlus* (Plasmid #64401, Addgene) (He *et al.*, 2020).

CcGRF and *CcGIF* were identified from the *C. campestris* genome (Vogel et al., 2018) via

sequence alignments and phylogenetic trees including wheat homologs (Debernardi et al., 2020),

performed in the Geneious Prime 2022.2.2 software. The selected *CcGRF* homolog Cc010427

and *CcGIF* homolog Cc021394 were cloned and sequenced using gene specific primers (Supp

table 5). The confirmed clones were ligated to make a *CcGRF/CcGIF* fusion via the Golden Gate

method using NEB's NEBridge kit and re-cloned into pCR8GWTOPO vector (Invitrogen).

CcWUS2 was synthesized by Azenta life science (www.azenta.com) and re-cloned into

pCR8GWTOPO vector.

CcWUS2 and *CcGRF/GIF* in entry clone pCR8GWTOPO were incubated with LR clonase

and pMDC139 to generate destination vectors with genes of interest (*35S:CcWUS2-GUS* and

35S:CcGRF/CcGIF-GUS). *35S:GUS-MAS:GFP*, *35S:CcWUS2-GUS*, and *35S:CcGRF/GIF-GUS*

were transformed to *A. tumefaciens* EHA105 strain.

Results

Test hormones and media compositions for *C. campestris* tissue culture

To establish the transformation protocol for *C. campestris*, we initially conducted screenings involving various concentrations of NAA and cytokinin (BAP/TDZ) in a medium designed for both co-cultivation and callus induction (Supp table 1). A negative control, consisting of the same medium without growth hormones, was also employed. GFP expression patterns were observed thirty days post-inoculation with *A. tumefaciens* EHA105-*35S:GUS-pMAS:GFP* (Figure 1). A callus showing fluorescence with the blue excitation light (Figure 1F) but not with the yellow excitation light (Figure 1G) was counted as transformed tissue (Supp table 6). Initially, we conducted a comparison of transformation efficiencies between groups with the presence or absence of plant growth regulators. The statistical analysis indicated a significantly higher GFP expression in the media containing plant growth regulators compared to that without hormones (Figure 2). *Cuscuta* calli cultivated on CIM media with BAP or TDZ exhibited significantly higher GFP expression than those on CIM media with NAA or the no hormone media, indicating that the presence of cytokinins (BAP or TDZ) had a noteworthy impact on efficiency. However, there was no significant difference among treatments with different concentration of BAP or TDZ. Thus, we concluded that the presence of BAP or TDZ without NAA in the CIM culture media significantly enhanced the *C. campestris* transformation efficiency. For further analysis, we used 5 mg/L BAP to induce *Cuscuta* callus in CIM media.

Comparing the cut *vs* the chop method using RUBY reporter

While approximately 50% of *Cuscuta* calli expressed GFP 30 days after *A. tumefaciens* inoculation and cultivated on 5mg/L BAP media (Figure 2), the preparation of individual meristems through the cut method (Figure 1B) is labor-intensive, as it involves the individual handling of each seedling. To simplify the inoculation process, we developed a bulk chopping method that is less precise but quicker and we conducted a comparative assessment of transformation efficiencies between the cut and bulk chop methods for preparing *Cuscuta* segments.

In this comparison, both cut and bulk chopped samples were infected with *A. tumefaciens* EHA105-*35S:RUBY* and incubated on MMS with 5mg/L BAP for 30 days. RUBY encodes enzymes for the synthesis of betalain and is useful for visually monitoring transformation events without extra equipment. Calli derived from the cut method exhibited 100% RUBY expression,

whereas calli from the bulk chop method displayed relatively lower RUBY expression (65%)

(Supp table 7). Previously we found that only the meristem region of *Cuscuta* was transformable

tissue, while neither the roots nor the shoots of *Cuscuta* exhibited such expression (Mudalige et

al., under preparation). Thus, we hypothesize that the bulk chop method had lower RUBY

expression than the cut method because the non-transformable tissues including root or shoot tips

were included with the transformable tissues incubated on the media. In addition, calli

expressing RUBY from the bulk chop method (Figure 3D-F) did not exhibit any phenotypic

differences when compared to calli from the cut method (Figure 3A-C). Even four-month-old

calli from the bulk chop method displayed vigorously growing RUBY tissues (Figure 3G-I). As

RUBY has been used as a visual selection marker in other plants (Yu *et al.*, 2023), we decided to

employ it as a visual selection marker during the early co-cultivation stage, eliminating the need

for a selection reagent.

***Agrobacterium*-mediated *Cuscuta* transformation**

To conduct the *Agrobacterium*-mediated *Cuscuta* transformation, 4-day-old *C. campestris*

seedlings were chopped and infected by *A. tumefaciens* EHA105-*35S:RUBY*. *Agrobacterium*

infected segments were incubated on the co-cultivation media without growth hormones for five

days (Figure 4A). Subsequently, *Cuscuta* segments expressing RUBY were transferred to CIM

media (Figure 4B). The explants were cultured in CIM media for at least 8 weeks before being

transferred to shoot induction media (Figure 4C). Larger and darker red calli were obtained after

another 8 weeks of incubation on the shoot induction media (Figure 4D) and then transferred to

shoot elongation media when the red calli reached a size of 1 cm (Figure 4E). Calli grown on the

shoot elongation media started to produce shoots in two weeks (Figure 4F).

During the callus induction stage, the number of visible red spots on *A. tumefaciens*

EHA105-*35S:RUBY* infected *Cuscuta* segments were counted on 5, 15, 25, 35, and 45 days after

infection to track the transformation efficiency. Since we selected *Cuscuta* segments expressing

RUBY to callus induction media after 5-day co-cultivation, transformed calli represented 100%

on the fifth day of inoculation. The number of transformed calli slowly declined to 93.2%,

86.0%, 76.7 %, and 62.1% on the 15, 25, 35, and 45 days after inoculation, respectively (Figure

5A and Supp table 8).

Further declines in the number of stably transformed calli occurred over subsequent growth

stages. Percentages of RUBY callus growing on shoot elongation media, callus producing RUBY

Cuscuta shoots, and RUBY *Cuscuta* shoots coiled on host were 16%, 5.3%, and 2.8%,

respectively (Figure 5B and Supp table 9).

Successful parasitism of transgenic RUBY *Cuscuta*

Once the RUBY *Cuscuta* shoots reached a size of 3 cm on the shoot elongation media, calli

with RUBY *Cuscuta* shoots were moved to magenta boxes containing the shoot elongation

media (Figure 6A). To observe the characteristic wrapping behavior of *Cuscuta*, a toothpick was

inserted into the media to serve as a substitute host. Within a day of introducing the toothpick,

RUBY *Cuscuta* coiled (Figure 6B) and produced haustoria (Figure 6C). We also introduced

RUBY *Cuscuta* stems to live host plants. Within 24 hours post-inoculation, the RUBY *Cuscuta*

stems exhibited coiling and initiated haustoria development on 1-week-old tomato (Figure 6D).

All host plants, both tomatoes (Figure 6E) and *Arabidopsis* (Figure 6F), were also successfully

parasitized by the RUBY *Cuscuta*. Approximately 2 months after transfer to beet hosts, RUBY

Cuscuta produced flowers (Figure 6G-H) and seeds (Figure 6I).

Event-specific detections of transgenic lines

From 8 attempts of the *Agrobacterium*-mediated *Cuscuta* transformation, 11 transgenic

events were generated (Supp table 9). To validate the T-DNA insertions, gDNA was extracted

from 2 well-developed transgenic events. Subsequently, TAIL-PCRs using right border (RB)

primers located in the 35S promoter of the *35S:RUBY* were conducted (Figure 7A). After

sequencing the amplicons, results showed that both transgenes lacked the RB sequences (gray

box, Figure 7A). Furthermore, the transgene of the first event contained a complete 35S

promoter, whereas the transgene from the second event had a truncated 35S promoter (Figure 7B

and Supp figure 1). By aligning TAIL-PCR sequences with *C. campestris* genome data, the

integration locations of transgenes were located on *C. campestris* Scaffold 32B and *C.*

campestris Scaffold 104, indicating that those events were independent transgenic plants (Figure

7B).

Diffused RUBY pigments in host stems

To compare the structural differences of haustoria between WT and transgenic RUBY

Cuscuta, WT *Cuscuta* and RUBY *Cuscuta* stems were inoculated on 2-week-old tomato

seedlings. The haustoria of wild-type *C. campestris* exhibited the typical hypersensitive response

near the epidermis without any visible color on the host stem (Figure 8A and E). When RUBY

Cuscuta was successfully coiled on the host stem but failed to penetrate, the red pigment was not

detected in host stems or vascular bundles (Figure 8B, F). Similarly, no RUBY pigment was
observed in the host stem during the early haustorium penetration stage of RUBY *Cuscuta*
(Figure 8C, G). However, we observed diffuse RUBY pigments (Figure 8D and white circles of
Figure 8H) in the area where successful haustoria development occurred, along with developed
searching hyphae (black arrows, Figure 8H).

The RUBY diffusion was also observed when RUBY *Cuscuta* was inoculated on *Arabidopsis*
Col-0 plants. After staining tissues with toluidine blue, we found that RUBY pigments were
mainly detected on cortex and phloem of host *Arabidopsis* stem (Figures 8I-K). RUBY was even
detected in regions away from *Cuscuta* haustoria (white circles of Figure 8I, L). Besides the
RUBY diffusion, there was no obvious physical difference between WT *Cuscuta* and RUBY
Cuscuta haustoria.

Enhanced transformation efficiency using *CcGRF/GIF Cuscuta*

Since morphogenic genes have enhanced plant transformation efficiency (Lowe *et al.*, 2016;
Chen *et al.*, 2022), we tested whether *C. campestris WUS2* or *C. campestris GRF/GIF* can
promote transformation efficiency. Four-day-old *C. campestris* seedlings (Figure 9A) were
chopped and infected by EHA105 with *GUS*, *CcWUS2-GUS*, and *CcGRF/GIF-GUS* (Figure 9B).
After the co-cultivation, 50 infected meristem-containing-segments were placed on CIM media
(Figure 9C). Around 35-day-old *Cuscuta* segments infected by EHA105 with *GUS*, *CcWUS2-
GUS*, and *CcGRF/GIF-GUS* were harvested and stained (Figure 9D-F). More specifically,
segments expressing GUS were quantified and classified by sizes of the GUS expressing sectors,
such as no GUS, less than 0.2 mm GUS, between 0.2 to 2 mm GUS, and larger than 2 mm GUS,
aiming to distinguish the extent of GUS expression in a segment (Supp figure 2).

In terms of presence and absence of GUS expression, the transformation efficiencies using
GUS, *CcWUS2-GUS*, and *CcGRF/GIF-GUS* were 78.8%, 51.7%, and 74.8%, respectively
(Figure 10A). Notably, the transformation efficiency associated with *CcWUS2-GUS* was
significantly lower than *GUS* or *CcGRF/GIF-GUS*. In contrast, there was no significant
difference in transformation efficiency between GUS and *CcGRF/GIF-GUS*. Data from
categorized segments indicated that *Cuscuta* segments infected with *GUS* only showed a
significantly higher proportion in the below 0.2 mm category compared to *CcWUS2-GUS* and
CcGRF/GIF-GUS (Figure 10B). *Cuscuta* with *CcGRF/GIF-GUS* had a significantly higher
percentage in the over 2mm group than *GUS* only. These findings suggest that *GUS* only and

317   *CcGRF/GIF-GUS* had similar levels of transformation efficiency based on the presence or

318   absence of GUS expression. However, *CcGRF/GIF-GUS* significantly enhanced regeneration

319   efficiency during the callus induction stage, leading to more significant GUS expression

320   compared to *GUS* only.

Discussion

The lack of a protocol for transformation and regeneration of *Cuscuta* has hindered progress

in studying parasite gene functions and the mechanisms of *Cuscuta*-host interactions. Despite

previous studies demonstrating transformation of *Cuscuta* tissues (Srivastava & Dwivedi, 2001;

Borsics *et al.*, 2002; Švubová & Blehová, 2013; Lachner *et al.*, 2020), none have succeeded in

regenerating stably transformed *Cuscuta* plants that can be used to study the full range of

parasite biology. Here we describe a method for the relatively fast and efficient stable

transformation of *C. campestris* plants that behave as normal parasites, producing haustoria,

flowers, and seeds (Figure 6). The process hinges on *A. tumefaciens*-mediated transformation

with appropriate media for inducing callus and shoots. We estimate that the process yields about

2.7% of transformed plants and takes about 5 months. While this transformation efficiency may

not be high compared to easily transformable plants like tobacco, it stands out as remarkably

successful for the initial attempt with a transformation recalcitrant species.

An important innovation was use of the recently developed RUBY reporter (He *et al.*, 2020).

RUBY encodes enzymes for synthesizing betalain from tyrosine and is useful for visually

monitoring transformation events without using fluorescence microscopy or a chemical selection

process. However, recent studies reported that RUBY may cause low regeneration efficiency and

poor fertility during seed development due to the lack of the essential amino acid tyrosine. For

example, maize expressing a strong RUBY reporter had growth retardation (Lee *et al.*, 2023).

While we are mindful that the RUBY reporter could be detrimental to *Cuscuta*, evidence to date

suggests that transgenic RUBY *Cuscuta* shows normal, though somewhat slower, growth

compared to non-transgenic plants. But importantly, the vivid RUBY reporter was a valuable

asset in the development of the protocol and may not be easily replaced by other selectable

markers.

The use of morphogenic genes appears to contribute positively to the *Cuscuta* transformation

protocol. We observed an enhancement in transformation efficiency with *Cuscuta GRF-GIF*, a

phenomenon similar to previously reported findings in wheat and citrus transformation

(Debernardi *et al.*, 2020). In *Agrobacterium*-mediated *Cuscuta* transformation, *CcGRF-GIF*

showed minimal impact on the overall presence/absence ratios of GUS transformation efficiency

(Figure 10A). However, the number of GUS segments larger than 2mm was significantly

increased in *CcGRF-GIF-GUS* infected *Cuscuta* compared to *Cuscuta* segments with *GUS* only

(Figure 10B). This suggests that *CcGRF-GIF* may not significantly influence *Agrobacterium-*

mediated transformation but could enhance the regeneration of transformed tissue, as previously

reported (Debernardi *et al.*, 2020). Further investigation is needed regarding the use of *CcGRF-*

GIF with *Cuscuta* transformation, as well as whether Ccu-miR396 can improve the efficiency of

stable transformation. In contrast, using *CcWUS-GUS* in transformations led to lower

presence/absence ratios of GUS transformation efficiency than *GUS* alone or *CcGRF-GIF-GUS*

(Figure 10A). We hypothesize that, similar to previous studies, the constitutive *CcWUS2*

expression by the 35S promoter may interfere with the *Cuscuta* development genes, resulting in

the decreased transformation efficiency (Rashid *et al.*, 2007; Lowe *et al.*, 2016; Hoerster *et al.*,

2020; Kadri *et al.*, 2021).

In this study we observed what appears to be movement of RUBY pigment from the

transformed parasite into host cells in the area around developed *Cuscuta* searching hyphae

inside vascular structures of the host stem (Figure 8H, K). However, the RUBY diffusion was

not observed during the early invasion stage (asterisk, Figure 8G). This is intriguing given the

ability of *Cuscuta* to exchange large numbers of mobile small RNAs, mRNAs and proteins

between *Cuscuta* and host plants. Here, toluidine staining suggest that RUBY pigments were

detected in host phloem and cortex (Figure 8K). Studies of long distance movement of mRNAs

suggest trafficking occurs in parasite and host phloem (Kim *et al.*, 2014; Park *et al.*, 2022), while

small RNAs appear to concentrate in haustorial regions (Shahid *et al.*, 2018). Consistent with

these findings, we detect the presence of RUBY pigment in both tissues, perhaps reflecting

general movement of molecules from parasite to host. It is interesting to note the differences in

patterns between the two host species, with a significant amount of RUBY found in the

Arabidopsis cortex, as compared to a distinct pattern around vascular bundles in tomato stems.

The existence of a partially effective defense in tomato may contribute to this. Thus, this

observation suggests that the RUBY reporter system serves not only as a selection marker for

transformation, but also as an indicator of events during parasitism at the molecular level.

Specifically, the RUBY movement prompts speculation about potential mechanisms of transfer:

1) translation of mobile RNA, wherein mRNAs of betalain synthesis enzymes move from

Cuscuta to the host stem and are translated near the haustoria; 2) mobile proteins from *Cuscuta*

to host, involving the export of three betalain synthesis enzymes from *Cuscuta* that produced

betalain in the host stem; or 3) the movement of betalain pigment itself from *Cuscuta* to the host.

Fortunately, we now have the tools to validate these hypotheses. It would be interesting to

generate transgenic *Cuscuta* expressing a fluorescence reporter gene fused with endoplasmic

reticulum (ER) targeting peptide, such as ER-YFP (Sager *et al.*, 2020), which produces a non-

mobile product. But regardless of the details, the ability to generate transgenic *Cuscuta* as

described here will enable many more experiments, and much greater understanding, of *Cuscuta*

biology.

Acknowledgement

This work was supported by the USDA-AFRI- 2023-67013-39896 to SP and JW, CAFNR-JOY,

University of Missouri to SP, Research Council, University of Missouri to SP and NSF award IOS-

1645027 to JW. We thank Duy Trinh who made the *CcGRF-GIF* construction. Grammarly generated

responses to the following AI prompts: Prompts created by Grammarly- "Improve it".

Conflicting interests

All authors declare that they have no conflicting interests**.**

Author contributions

SP, HL, and JW conceived the ideas and designed methodology. SA, AM, VB, HG, and SP conducted

lab-work and analyzed the data. SA, AM, SP, and JW led the writing of the manuscript. All authors

contributed critically to the drafts and gave final approval for publication.

Data availability

The datasets used and/or analyzed during the current study are available from the corresponding author

upon request.

References

Bandaranayake PCG, Yoder JI. 2018. Factors affecting the efficiency of Rhizobium rhizogenes root transformation of the root parasitic plant Triphysaria versicolor and its host Arabidopsis thaliana. *Plant Methods* **14**: 61.

Bernal-Galeano V, Beard K, Westwood JH. 2022. An artificial host system enables the obligate parasite Cuscuta campestris to grow and reproduce in vitro. *Plant Physiology* **189**(2): 687-702.

Borsics T, Mihálka V, Oreifig AS, Bárány I, Lados M, Nagy I, Jenes B, Toldi O. 2002. Methods for genetic transformation of the parasitic weed dodder (*Cuscuta trifolii* Bab. et Gibs) and for PCR-based detection of early transformation events. *Plant Science* **162**(2): 193-199.

Bull T, Debernardi J, Reeves M, Hill T, Bertier L, Van Deynze A, Michelmore R. 2023. GRF–GIF chimeric proteins enhance in vitro regeneration and Agrobacterium-mediated transformation efficiencies of lettuce (Lactuca spp.). *Plant Cell Reports* **42**(3): 629-643.

Chen Z, Debernardi JM, Dubcovsky J, Gallavotti A. 2022. Recent advances in crop transformation technologies. *Nature Plants* **8**(12): 1343-1351.

Cudney DW, Orloff SB, Reints JS. 1992. An Integrated Weed Management Procedure for the Control of Dodder (Cuscuta indecora) in Alfalfa (Medicago sativa). *Weed Technology* **6**(3): 603-606.

Dawson JH, Musselman LJ, Wolswinkel P, Dörr I. 1994. Biology and control of Cuscuta. *Reviews of Weed Science* **6**: 265-317.

Debernardi JM, Tricoli DM, Ercoli MF, Hayta S, Ronald P, Palatnik JF, Dubcovsky J. 2020. A GRF–GIF chimeric protein improves the regeneration efficiency of transgenic plants. *Nature Biotechnology* **38**(11): 1274-1279.

Dechassa N, Regassa B. 2021. Current Status, Economic importance and Management of Dodders (Cuscuta Spp) of Important Crops.

Feng Q, Xiao L, He Y, Liu M, Wang J, Tian S, Zhang X, Yuan L. 2021. Highly efficient, genotype-independent transformation and gene editing in watermelon (Citrullus lanatus) using a chimeric ClGRF4-GIF1 gene. *J Integr Plant Biol* **63**(12): 2038-2042.

Fernández-Aparicio M, Rubiales D, Bandaranayake PC, Yoder JI, Westwood JH. 2011. Transformation and regeneration of the holoparasitic plant Phelipanche aegyptiaca. *Plant Methods* **7**(1): 36.

Galloway AF, Knox P, Krause K. 2020. Sticky mucilages and exudates of plants: putative microenvironmental design elements with biotechnological value. *New Phytologist* **225**(4): 1461-1469.

Goldwasser Y, Sazo MRM, Lanini WT. 2012. Control of Field Dodder (Cuscuta campestris) Parasitizing Tomato with ALS-Inhibiting Herbicides. *Weed Technology* **26**(4): 740-746.

He Y, Zhang T, Sun H, Zhan H, Zhao Y. 2020. A reporter for noninvasively monitoring gene expression and plant transformation. *Horticulture Research* **7**(1): 152-152.

Hegenauer V, Fürst U, Kaiser B, Smoker M, Zipfel C, Felix G, Stahl M, Albert M. 2016. Detection of the plant parasite Cuscuta reflexa by a tomato cell surface receptor. *Science* **353**(6298): 478-481.

Hegenauer V, Slaby P, Körner M, Bruckmüller J-A, Burggraf R, Albert I, Kaiser B, Löffelhardt B, Droste-Borel I, Sklenar J. 2020. The tomato receptor CuRe1 senses a cell wall protein to identify Cuscuta as a pathogen. *Nature communications* **11**(1): 5299.

Hoerster G, Wang N, Ryan L, Wu E, Anand A, McBride K, Lowe K, Jones T, Gordon-Kamm B. 2020. Use of non-integrating Zm-Wus2 vectors to enhance maize transformation: non-integrating WUS2 enhances transformation. *In Vitro Cellular & Developmental Biology-Plant* **56**(3): 265-279.

Ishida JK, Yoshida S, Ito M, Namba S, Shirasu K. 2011. Agrobacterium rhizogenes-Mediated Transformation of the Parasitic Plant Phtheirospermum japonicum. *PLoS ONE* **6**(10): e25802.

Johnsen HR, Striberny B, Olsen S, Vidal-Melgosa S, Fangel JU, Willats WGT, Rose JKC, Krause K. 2015. Cell wall composition profiling of parasitic giant dodder (Cuscuta reflexa) and its hosts: a priori differences and induced changes. *New Phytologist* **207**(3): 805-816.

Kadri A, Grenier De March G, Guerineau F, Cosson V, Ratet P. 2021. WUSCHEL Overexpression Promotes Callogenesis and Somatic Embryogenesis in Medicago truncatula Gaertn. *Plants (Basel)* **10**(4).

Kaiser B, Vogg G, Fürst UB, Albert M. 2015. Parasitic plants of the genus Cuscuta and their interaction with susceptible and resistant host plants. *Frontiers in Plant Science* **6**.

Kim G, LeBlanc ML, Wafula EK, dePamphilis CW, Westwood JH. 2014. Genomic-scale exchange of mRNA between a parasitic plant and its hosts. *Science* **345**(6198): 808-811.

Lachner LA-M, Galstyan L, Krause K. 2020. A highly efficient protocol for transforming Cuscuta reflexa based on artificially induced infection sites. *Plant Direct* **4**(8): e00254.

Lanini W, Kogan M. 2005. Biology and management of Cuscuta in crops. *International Journal of Agriculture and Natural Resources* **32**(3): 127-141.

Lee K, Kang M, Ji Q, Grosic S, Wang K. 2023. New T-DNA binary vectors with NptII selection and RUBY reporter for efficient maize transformation and targeted mutagenesis. *Plant Physiology* **192**(4): 2598-2603.

Lee K, Wang K. 2023. Strategies for genotype-flexible plant transformation. *Current Opinion in Biotechnology* **79**: 102848.

Liu N, Shen G, Xu Y, Liu H, Zhang J, Li S, Li J, Zhang C, Qi J, Wang L, Wu J. 2020. Extensive Inter-plant Protein Transfer between Cuscuta Parasites and Their Host Plants. *Mol Plant* **13**(4): 573-585.

Lowe K, Wu E, Wang N, Hoerster G, Hastings C, Cho MJ, Scelonge C, Lenderts B, Chamberlin M, Cushatt J, et al. 2016. Morphogenic Regulators Baby boom and Wuschel Improve Monocot Transformation. *Plant Cell* **28**(9): 1998-2015.

Nagar R, Singh M, Sanwal GG. 1984. Cell Wall Degrading Enzymes in Cuscuta reflexa and Its Hosts. *Journal of Experimental Botany* **35**(8): 1104-1112.

Park SY, Shimizu K, Brown J, Aoki K, Westwood JH. 2022. Mobile Host mRNAs Are Translated to Protein in the Associated Parasitic Plant Cuscuta campestris. *Plants (Basel)* **11**(1).

Pradhan Mitra P, Loqué D. 2014. Histochemical staining of Arabidopsis thaliana secondary cell wall elements. *J Vis Exp*(87).

Rashid SZ, Yamaji N, Kyo M. 2007. Shoot formation from root tip region: a developmental alteration by WUS in transgenic tobacco. *Plant Cell Reports* **26**(9): 1449-1455.

Runyon JB, Mescher MC, De Moraes CM. 2006. Volatile chemical cues guide host location and host selection by parasitic plants. *Science* **313**(5795): 1964-1967.

Sager R, Wang X, Hill K, Yoo B-C, Caplan J, Nedo A, Tran T, Bennett MJ, Lee J-Y. 2020. Auxin-dependent control of a plasmodesmal regulator creates a negative feedback loop modulating lateral root emergence. *Nature Communications* **11**(1): 364.

Shahid S, Kim G, Johnson NR, Wafula E, Wang F, Coruh C, Bernal-Galeano V, Phifer T, dePamphilis CW, Westwood JH, Axtell MJ. 2018. MicroRNAs from the parasitic plant Cuscuta campestris target host messenger RNAs. *Nature* **553**: 82.

Srivastava S, Dwivedi UN. 2001. Plant regeneration from callus of Cuscuta reflexa–an angiospermic parasite–and modulation of catalase and peroxidase activity by salicylic acid and naphthalene acetic acid. *Plant Physiology and Biochemistry* **39**(6): 529-538.

Švubová R, Blehová A. 2013. Stable transformation and actin visualization in callus cultures of dodder (Cuscuta europaea). *Biologia* **68**(4): 633-640.

Tomilov A, Tomilova N, Yoder JI. 2007. Agrobacterium tumefaciens and Agrobacterium rhizogenes transformed roots of the parasitic plant Triphysaria versicolor retain parasitic competence. *Planta* **225**(5): 1059-1071.

Vaughn KC. 2002. Attachment of the parasitic weed dodder to the host. *Protoplasma* **219**(3): 227-237.

507   **Yu J, Deng S, Huang H, Mo J, Xu Z-F, Wang Y. 2023.** Exploring the Potential Applications of the

508   Noninvasive Reporter Gene RUBY in Plant Genetic Transformation. *Forests* **14**(3): 637.

509   **Zuo J, Niu Q-W, Frugis G, Chua N-H. 2002.** The WUSCHEL gene promotes vegetative-to-embryonic

510   transition in Arabidopsis. *The Plant Journal* **30**(3): 349-359.

Figure 1. Process of *Agrobacterium*-mediated *Cuscuta* transformation using *35S:GUS-pMAS:GFP* construct. *Cuscuta* seedlings (A) were cut by scissors (B) and dipped in liquid MMS containing *A. tumefaciens* EHA105-*35S:GUS-pMAS:GFP* (C). An excessive *Agrobacteria* was cultivated from infected tissue five days after inoculation on the co-cultivation media (D). Explants were incubated on CIM (E). An explant imaged under blue excitation light (F), yellow excitation light (G), and bright field (H). Images from F-H were merged (I).

Figure 2. Effect of cytokinins and auxin on GFP positive calli formation after inoculation with *Agrobacterium tumefaciens* EHA105 containing *35S:GUS-pMAS:GFP*. Different types and concentrations of cytokinins (BAP and TDZ) and auxin (NAA) were tested (n=10). Six independent biological replicates were performed. Data are presented as the mean ± standard deviation. Groups with different letters vary significantly (two-way ANOVA-Tukey test, P,0.05).

Figure 3. RUBY expression in explants. RUBY expression in cut explants (A, B, C) and bulk chopped explants (D, E, F) 40 days after inoculation. Callus expressing RUBY developed in bulk chopped explants 4 months after inoculation (G, H, I).

Figure 4. Process of *Agrobacterium*-mediated *Cuscuta* transformation using *35S:RUBY* construct. *Cuscuta* seedlings were infected by EHA105-*35S:RUBY* and incubated on co-cultivation media for 5 days (A) and separated on CIM (B). RUBY-expressing calli were grown on CIM (C) and SIM (D). Calli expressing RUBY were selected and transferred to SEM (E) and produced shoots (F).

Figure 5. *C. campestris* transformation efficiency with the RUBY reporter gene. (A) Transformation rate on various days following inoculation. Sixteen independent replicates were performed (n=50). (B) The callus induction rate, shoot induction rate, and coiling rate were measured for RUBY *Cuscuta*. Eight independent replicates were performed (n=50). Data are presented as the mean ± standard deviation.

Figure 6. Transgenic RUBY *Cuscuta* attaching to non-living and living hosts. A shoot from transgenic callus (A) coiled on a toothpick (B-C). (C) Zoomed-in image of the black box in (B) showing haustoria formation. RUBY *Cuscuta* coiled on (D) and parasitizing (E) the stem of a 2-week-old tomato. (F) RUBY *Cuscuta* was successfully grown on *Arabidopsis*. (I) Transgenic RUBY *Cuscuta* produced flowers on beets (G-H) and seeds.

Figure 7. Alignment of sequences from TAIL PCRs and the *35S:RUBY* construct. (A) Diagram of 35S promoter and RUBY near RB region in *35S:RUBY*. Red arrow, yellow arrow, and gray box are RUBY, 35S promoter, and right border, respectively. Black arrows (RB1 and RB2) are primers used for the TAIL PCRs. (B) Schematic designs of T-DNA sequences and insertion points into the *C. campestris* genome. Yellow boxes indicate 35S promoter of *35S:RUBY* and white boxes are *C. campestris* genome sequences.

Figure 8. Cross sections of *Cuscuta* haustoria on tomato or *Arabidopsis* stems. Haustoria of WT *C. campestris* successfully penetrated tomato stems and established a connection with the host vascular bundle (A, E). RUBY *Cuscuta* coiling around a host tomato stem but failing to penetrate (B, F). RUBY *Cuscuta* successfully attaching to tomato is shown at early (C, G) and mature (D, H) haustoria development stages. (E-H): Zoomed-in images of the white box region of corresponding images in (A-D). RUBY *Cuscuta* penetrated and developed haustoria on *Arabidopsis* Col-0 stem (I). Images of the white box in (I) before staining (J) and after toluidine staining (K). Vertical section of *Cuscuta* haustorium on *Arabidopsis* Col-0 (L). Asterisks (*), white circles, and black arrows indicate the presence of *Cuscuta* haustorium, area of diffused RUBY pigment, and searching hyphae, respectively. xy, xylem; ixf, xylem fibres; ph, phloem; cor, cortex.

Figure 9. *Agrobacterium*-mediated *Cuscuta* transformation using chopped seedlings. Four-day-old seedlings (A) were chopped (B) and inoculated with *Agrobacterium tumefaciens* strain EHA105 harboring *35S:GUS-pMAS:GFP*. (C) Infected segments were incubated on cocultivation media. (D-F) Approximately 35 days after inoculation, segments infected by *35S:GUS-pMAS:GFP, 35S:CcWUS2-GUS* and *35S: CcGRF-CcGIF-GUS* were stained with X-gluc solution.

Figure 10. Effect of *CcWUS2* and *CcGRF/GIF* on *Agrobacterium*-mediated *Cuscuta* transformation. (A) The transformation efficiencies of *GUS* (blue bar), *CcWUS2-GUS* (purple bar)*,* and *CcGRF/GIF-GUS* (pink bar) were compared. (B) GUS expressions were classified into four groups according to the size of GUS expression. Thirty independent replicates were performed (n=50). Data are presented as the mean ± standard deviation.

Supplementary Figure 1. Results of TAIL-PCRs to confirm genomic integration. Sequences are details of the diagram in Figure 8B. The *35S:RUBY* vector sequences (yellow highlights) are linked to *C. campestris* genome scaffolds 32B and 104.

577 **Supplementary Figure 2.** Various sizes of GUS expression sectors in transformed tissues. Following the

578 GUS staining assay, the segments were classified into four groups: No GUS (A-C), less than 0.2 mm

579 GUS (D-F), between 0.2 to 2 mm GUS (G-I), and larger than 2 mm GUS (J-L). The bars represent a scale

580 of 500 μm.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

Figure 7

Figure 8

A 4-day-old seedlings   **B** Inoculation   **C** Callus induction

D *35S:GUS-pMAS:GFP*   **E** *35S:CcWUS2-GUS*   **F** *35S:CcGRF-CcGIF-GUS*

Figure 9

Figure 10

35S promoter and scaffolds 32B

ATGCTCCTCGTGGGTGGGGGTCCATCTTTGGGACCACTGTCGGCAGAGGCATCTTGAAC
GATAGCCTTTCCTTTATCGCAATGATGGCATTTGTAGGTGCCACCTTCCTTTTCTACTGTCC
TTTTGATGAAGTGACAGATAGCTGGGCAATGGAATCCGAGGAGGTTTCCCGATATTACCCT
TTGTAGAAAAGTCTCAAATCATCCATATATTCGATTGATTCTTTGCAGTACCCTTACCCGAC
TTATGGTGATACTTTATCTTCTAAAGGATTTCATATTCTCAGTGAGAATATGAAACGATCCA
AACGTACGAGACTTCCGACAACTTCTATGGATGTTATTTCACATTATCATCCTACAACAATT
CAAGTTGATAGCATTCCAATCGATCGCCAAATCTCTATCGGGTCAATCAACTAGCCCTTTG
ATCGCCTTCTTATTGGCCCTATCTGGGTGGACCCCTTCTGACCAGTCGTTCATCCCTCCGT
GGCGCGGCACACTTGTCCTACCGTCACCGAGTCAAGTCCCAAGCTGGTTAGCCATGGGTT
TACACCCATCACTTGGTACCCAATCTGGTGTGATTATTTGATATTGTTCTTATTTTTGTACTC
CCTACATCCTATTAGGACTTTCCAAATTCAAAGAAAAAATGTGTTGTCTATTATTTTTTCTCA
ATAATGAAATTTAATCACACTCTTC

35S promoter and scaffolds 104

ATGCTCCTCGTGGGTGGGGGTCCATCTTTGGGACCACTGTCGGCAGAGGCATCTTGGAC
GATAGCCTTTCCTTTATCGCAATGATGGCATTTGTAGGTGCCACCTTCCTTTTCTACTGTCC
TTTTGATGAAGTGACAGATAGCTGGGCAATGGAATCCGAGGAGGTTTCCCGATATTACCCT
ATCCATGAAACTTAGGAGTTCGCACCCCGCAGTTTCGTCCACCATTTGATCTGGACTGGG
GAGGGGGTATGAATCCATTGGACAAGCCTTGTTCAAGTCCGTATAATCCACACACATTCTC
CACGTGTCGCCCTTGGGAGCAAGGACAACGTTGGCCAGCCAATCCGGGTACTTGACTTC
CCGGATGTGACCAACACTGAGGAGGTCTCTCACTTCCTTCTTGACGAAGTCTCGCCGGTC
ACTCGCCAGGTACCGCTTCTTTTGGCGTACCGGTCGACTTCCCTCCCTCACGGCCAACCG
GTGGCAGATAACAGAGCGATCTATCCCGGGCATATCCGCCACGCTCCATGCGAATATGCC
CGCGTATTCTTGGAGGACCCGTGTAATCTGCTCCTTGAGATCCTCGGGAAGATTTGCTCC
AATTCTGACCATCCTCGAAGAATCACCCGGGTGAAGTTCGATTTCCACCGTCTCAGCCTC
GGGCTCTAGTTTCTCCTGGGTCTCTCCCTTGGTGATCTCTTGGGAGATGACATCGACCCC
CCTGGTCATTTTCTTGACTGCCCGGGCATAGCACTCCCGGGCGTTCCTCTGGTCGCCCTT
CATCTCACTCGACCA

Supp Figure 1

Supp Figure 2

Supplementary table 1. Media compositions for the first and second protocols.

Modified MS Media (MMS)		Co-Cultivation Media (CCM)		Callus Induction Media (CIM)	
Component	Amount	Component	Amount	Component	Amount
NH_4NO_3	300 mg/l	NH_4NO_3	300 mg/l	NH_4NO_3	300 mg/l
KNO_3	800 mg/l	KNO_3	800 mg/l	KNO_3	800 mg/l
$CaCl_2.2H_2O$	250 mg/l	$CaCl_2.2H_2O$	250 mg/l	$CaCl_2.2H_2O$	250 mg/l
$MgSO_4.7H_2O$	260 mg/l	$MgSO_4.7H_2O$	260 mg/l	$MgSO_4.7H_2O$	260 mg/l
KH_2PO_4	120 mg/l	KH_2PO_4	120 mg/l	KH_2PO_4	120 mg/l
H_3BO_3	6.2 mg/l	H_3BO_3	6.2 mg/l	H_3BO_3	6.2 mg/l
$MnSO_4.4H_2O$	22.3 mg/l	$MnSO_4.4H_2O$	22.3 mg/l	$MnSO_4.4H_2O$	22.3 mg/l
$ZnSO_4.7H_2O$	8.6 mg/l	$ZnSO_4.7H_2O$	8.6 mg/l	$ZnSO_4.7H_2O$	8.6 mg/l
KI	0.83 mg/l	KI	0.83 mg/l	KI	0.83 mg/l
$NaMO_4.2H_2O$	0.25 mg/l	$NaMO_4.2H_2O$	0.25 mg/l	$NaMO_4.2H_2O$	0.25 mg/l
$CuSO_4.5H_2O$	0.025 mg/l	$CuSO_4.5H_2O$	0.025 mg/l	$CuSO_4.5H_2O$	0.025 mg/l
$CoCl_2.6H_2O$	0.025 mg/l	$CoCl_2.6H_2O$	0.025 mg/l	$CoCl_2.6H_2O$	0.025 mg/l
Myo-inositol	100 mg/l	Myo-inositol	100 mg/l	Myo-inositol	100 mg/l
Nicotinic acid	1 mg/l	Nicotinic acid	1 mg/l	Nicotinic acid	1 mg/l
Pyridoxine hydrochloride	1 mg/l	Pyridoxine hydrochloride	1 mg/l	Pyridoxine hydrochloride	1 mg/l
Thiamine hydrochloride	10 mg/l	Thiamine hydrochloride	10 mg/l	Thiamine hydrochloride	10 mg/l
Glucose	3% (w/v)	Glucose	3% (w/v)	Glucose	3% (w/v)
		*Plant growth regulators		Cefotaxime	250 mg/l
				*Plant growth regulators	

* Different amounts of plant growth regulators were applied as described in the methods section.

Supplementary table 2. Media compositions for the third protocol.

Co-cultivation media (CCM)		Callus Induction Media (CIM)		Shoot Induction Media (SIM)		Shoot Elongation Media (SEM)	
Component	Amount	Component	Amount	Component	Amount	Component	Amount
NH_4NO_3	300 mg/l	NH_4NO_3	300 mg/l	NH_4NO_3	825 mg/l	NH_4NO_3	825 mg/l
KNO_3	800 mg/l	KNO_3	800 mg/l	KNO_3	950 mg/l	KNO_3	950 mg/l
$CaCl_2.2H_2O$	250 mg/l	$CaCl_2.2H_2O$	250 mg/l	$CaCl_2.2H_2O$	166.1 mg/l	$CaCl_2.2H_2O$	166.1 mg/l
$MgSO_4.7H_2O$	260 mg/l	$MgSO_4.7H_2O$	260 mg/l	$MgSO_4.7H_2O$	180.7 mg/l	$MgSO_4.7H_2O$	180.7 mg/l
KH_2PO_4	120 mg/l	KH_2PO_4	120 mg/l	KH_2PO_4	170 mg/l	KH_2PO_4	170 mg/l
H_3BO_3	6.2 mg/l	H_3BO_3	6.2 mg/l	H_3BO_3	6.2 mg/l	H_3BO_3	6.2 mg/l
$MnSO_4.4H_2O$	22.3 mg/l	$MnSO_4.4H_2O$	22.3 mg/l	$MnSO_4.H_2O$	16.9 mg/l	$MnSO_4.H_2O$	16.9 mg/l
$ZnSO_4.7H_2O$	8.6 mg/l	$ZnSO_4.7H_2O$	8.6 mg/l	$ZnSO_4.7H_2O$	8.6 mg/l	$ZnSO_4.7H_2O$	8.6 mg/l
KI	0.83 mg/l	KI	0.83 mg/l	KI	0.83 mg/l	KI	0.83 mg/l
$NaMO_4.2H_2O$	0.25 mg/l	$NaMO_4.2H_2O$	0.25 mg/l	$NaMO_4.2H_2O$	0.25 mg/l	$NaMO_4.2H_2O$	0.25 mg/l
$CuSO_4.5H_2O$	0.025 mg/l	$CuSO_4.5H_2O$	0.025 mg/l	$CuSO_4.5H_2O$	0.025 mg/l	$CuSO_4.5H_2O$	0.025 mg/l
$CoCl_2.6H_2O$	0.025 mg/l	$CoCl_2.6H_2O$	0.025 mg/l	$CoCl_2.6H_2O$	0.025 mg/l	$CoCl_2.6H_2O$	0.025 mg/l
Myo-inositol	100 mg/l	Myo-inositol	100 mg/l	$FeSO_4.7H_2O$	27.8 mg/l	$FeSO_4.7H_2O$	27.8 mg/l
Nicotinic acid	1 mg/l	Nicotinic acid	1 mg/l	$Na_2EDTA.2H_2O$	37.26 mg/l	$Na_2EDTA.2H_2O$	37.26 mg/l
Pyridoxine hydrochloride	1 mg/l	Pyridoxine hydrochloride	1 mg/l	Myo-inositol	100 mg/l	Myo-inositol	100 mg/l
Thiamine hydrochloride	10 mg/l	Thiamine hydrochloride	10 mg/l	Nicotinic acid	1 mg/l	Nicotinic acid	1 mg/l
Glucose	3% (w/v)	Glucose	3% (w/v)	Pyridoxine hydrochloride	1 mg/l	Pyridoxine hydrochloride	1 mg/l
NAA	3 mg/l	BAP	5 mg/l	Thiamine hydrochloride	10 mg/l	Thiamine hydrochloride	10 mg/l
BAP	1 mg/l	Cefotaxime	250 mg/l	Glucose	3% (w/v)	Glucose	3% (w/v)
				BAP	5 mg/l	BAP	5 mg/l
				Cefotaxime	250 mg/l	Cefotaxime	250 mg/l
				Timentin	100 mg/l	Timentin	100 mg/l

Supplementary table 3. TAIL-PCR conditions.

Primary TAIL-PCR	Secondary TAIL-PCR
95C – 3 min	95C – 3 min
5 cycles	5 cycles
95C – 30s	95C – 10s
62C – 1 min	64C – 1min
72C – 2:30 min	72C – 1:30min
2 cycles	15 cycles
95C – 30s	95C – 10s
25C – 3min (50% ramp)	64C – 1 min
72C – 2:30 min (32% ramp)	72C – 1:30 min
	95C -10s
	64 – 1 min
15 cycles	72C – 1:30 min
95C – 10s	95C -10s
68 – 1min	44C – 1 min
72C – 2:30 min	72C – 1:30 min
95C -10s	
68 – 1min	5 cycles
72C – 2:30 min	95C – 10s
95C – 10s	44C – 1 min
44C – 1 min	72C – 2min
72C – 2:30 min	
	72C – 5min
72C – 5min	4C 2min
4C – 2min	

Supplementary table 4. Primers used for TAIL PCR.

	Name	Sequence 5'-3'
T-DNA specific primers	RB1-RUBY	GTGTCCTCTCCAAATGAAATGAACTTCCTTATATAGAG
	RB2-RUBY	CCTTTGGTCTTCTGAGACTGTATCTTTGATATTCTTG
Arbitrary primers	AD1	NGTCGASWGANAWGAA
	AD2	TGWGNAGSANCASAGA
	AD3	AGWGNAGWANCAWAGG
	AD6	WGTGNAGWANCANAGA

Supplementary table 5. Primers used for cloning.

	Forward primers	Reverse primers
CcGRFGIF	ATGATGAGCAGTGGAAGAGGCAGG	tTTGATTCCCGCCCGCCGATC
CCWUS2	ATGGAGCCTCAACACTACTATC	tGGCGAAATGGTGAGACGAC

Supplementary table 6. Transformation efficiency on different callus induction media

NAA (mg/L)	BAP (mg/L)	TDZ (mg/L)	Plate no	Number of calli	Total explants per plate	Average of a plate	Average (%)
0	5	0	1	6	10	60	
0	5	0	2	6	10	60	
0	5	0	3	5	10	50	53.3
0	5	0	4	4	10	40	
0	5	0	5	5	10	50	
0	5	0	6	6	10	60	
0	7.5	0	1	5	10	50	
0	7.5	0	2	5	10	50	
0	7.5	0	3	7	10	70	56.7
0	7.5	0	4	6	10	60	
0	7.5	0	5	4	10	40	
0	7.5	0	6	7	10	70	
0	10	0	1	3	10	30	
0	10	0	2	3	10	30	
0	10	0	3	4	10	40	31.7
0	10	0	4	4	10	40	
0	10	0	5	2	10	20	
0	10	0	6	3	10	30	
0	25	0	1	6	10	60	
0	25	0	2	5	10	50	
0	25	0	3	6	10	60	43.3
0	25	0	4	5	10	50	
0	25	0	5	2	10	20	
0	25	0	6	2	10	20	
0	0	0.05	1	5	10	50	
0	0	0.05	2	3	10	30	
0	0	0.05	3	4	10	40	43.3
0	0	0.05	4	6	10	60	
0	0	0.05	5	3	10	30	
0	0	0.05	6	5	10	50	

0	0	0.1	1	7	10	70	
0	0	0.1	2	2	10	20	
0	0	0.1	3	3	10	30	36.7
0	0	0.1	4	5	10	50	
0	0	0.1	5	3	10	30	
0	0	0.1	6	2	10	20	
0	0	0.3	1	2	10	20	
0	0	0.3	2	4	10	40	
0	0	0.3	3	2	10	20	33.3
0	0	0.3	4	4	10	40	
0	0	0.3	5	4	10	40	
0	0	0.3	6	4	10	40	
0	0	0.5	1	6	10	60	
0	0	0.5	2	5	10	50	
0	0	0.5	3	6	10	60	50.0
0	0	0.5	4	7	10	70	
0	0	0.5	5	5	10	50	
0	0	0.5	6	1	10	10	
0.5	5	0	1	5	10	50	
0.5	5	0	2	2	10	20	
0.5	5	0	3	4	10	40	26.7
0.5	5	0	4	3	10	30	
0.5	5	0	5	2	10	20	
0.5	5	0	6	0	10	0	
0.5	7.5	0	1	2	10	20	
0.5	7.5	0	2	2	10	20	
0.5	7.5	0	3	3	10	30	26.7
0.5	7.5	0	4	3	10	30	
0.5	7.5	0	5	3	10	30	
0.5	7.5	0	6	3	10	30	
0.5	10	0	1	1	10	10	
0.5	10	0	2	4	10	40	
0.5	10	0	3	5	10	50	31.7
0.5	10	0	4	4	10	40	
0.5	10	0	5	2	10	20	

0.5	10	0	6	3	10	30	
0.5	25	0	1	1	10	10	
0.5	25	0	2	1	10	10	
0.5	25	0	3	2	10	20	13.3
0.5	25	0	4	1	10	10	
0.5	25	0	5	2	10	20	
0.5	25	0	6	1	10	10	
0.5	0	0.05	1	5	10	50	
0.5	0	0.05	2	2	10	20	
0.5	0	0.05	3	3	10	30	26.7
0.5	0	0.05	4	2	10	20	
0.5	0	0.05	5	3	10	30	
0.5	0	0.05	6	1	10	10	
0.5	0	0.1	1	1	10	10	
0.5	0	0.1	2	2	10	20	
0.5	0	0.1	3	1	10	10	16.7
0.5	0	0.1	4	3	10	30	
0.5	0	0.1	5	2	10	20	
0.5	0	0.1	6	1	10	10	
0.5	0	0.3	1	2	10	20	
0.5	0	0.3	2	5	10	50	
0.5	0	0.3	3	3	10	30	33.3
0.5	0	0.3	4	4	10	40	
0.5	0	0.3	5	2	10	20	
0.5	0	0.3	6	4	10	40	
0.5	0	0.5	1	3	10	30	
0.5	0	0.5	2	1	10	10	
0.5	0	0.5	3	2	10	20	25.0
0.5	0	0.5	4	3	10	30	
0.5	0	0.5	5	3	10	30	
0.5	0	0.5	6	3	10	30	
0	0	0	1	2	10	20	
0	0	0	2	2	10	20	8.3
0	0	0	3	1	10	10	
0	0	0	4	0	10	0	

0	0	0	5	1	10	10
0	0	0	6	1	10	10
0	0	0	1	0	10	0
0	0	0	2	1	10	10
0	0	0	3	0	10	0
0	0	0	4	0	10	0
0	0	0	5	1	10	10
0	0	0	6	1	10	10

Supplementary table 7. Different transformation efficiency between the cut and the bulk chop method. DAI is days after inoculation.

Wounding method	Incubation dates	Total explants per plate	Replicates					Average number of red spots in a plate	Average efficiency (%)
			plate 1	plate 2	plate 3	plate 4	plate 5		
Cut	20 DAI	10	10	10	10	10	10	10	100
Chop	20 DAI	20	15	17	10	9	14	13	65

Supplemental table 8. Transformation efficiency with RUBY reporter. Number of segments expressing RUBY were scored from 4DAI to 45 DAI. Sixteen independent biological repeats were performed (n=50). DAI: Days after Agro inoculation.

No. of reps.	5 DAI (%)	15 DAI (%)	25 DAI (%)	35 DAI (%)	45 DAI (%)
1	100	88.0	82.0	78.0	68.0
2	100	88.5	80.8	76.9	69.2
3	100	94.6	80.4	73.2	60.7
4	100	92.3	82.7	69.2	65.4
5	100	98.0	84.0	74.0	50.0
6	100	100.0	94.0	82.0	60.0
7	100	88.1	78.0	74.6	45.8
8	100	98.1	94.2	78.8	57.7
9	100	94.0	94.0	80.0	80.0
10	100	90.0	90.0	74.0	74.0
11	100	84.0	80.0	74.0	62.0
12	100	96.0	94.0	74.0	72.0
13	100	94.7	86.0	84.2	52.6
14	100	96.2	90.4	88.5	65.4
15	100	96.2	84.6	73.1	53.8
16	100	92.9	80.4	73.2	57.1
Average	100	93.2	86.0	76.7	62.1

Supplemental table 9. Transformation efficiency with RUBY reporter. Number of callus expressing RUBY, number of shoots expressing RUBY and number of RUBY *Cuscuta* coiled on a host were counted. Eight independent biological repeats were performed (n=50).

No. of reps.	% of callus on Shoot elongation	% of callus producing shoot	% of shoots coiled on a host
1	24.0	10.0	8.0
2	14.0	4.0	4.0
3	12.0	4.0	2.0
4	24.0	8.0	4.0
5	18.0	8.0	2.0
6	6.0	0.0	0.0
7	10.0	0.0	0.0
8	20.0	8.0	2.0
Average	16.0	5.3	2.8