IFNB induced by non-lytic virus immunotherapy promotes improved survival in hepatocellular carcinoma, mediated by MHCII-independent cytotoxic CD4[+] T-cells.

AUTHORS: Russell Hughes[1]*, Amy Moran[1], Samantha Garcia-Cardenas[1], Karen J. Scott[1], Elizabeth Appleton[1], Matthew J. Bentham[1], Elizabeth Ilett[1], Alan Melcher[3], Andrew Macdonald[2], Adel Samson[1,]*, and Stephen Griffin[1,]*

AFFILITATIONs:

[1]Leeds Institute of Medical Research, St. James's University Hospital, University of Leeds; Leeds, UK.

[2]School of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds; Leeds, UK.

[3]The Institute for Cancer Research, University of London; London, UK.

*Corresponding author(s): s.d.c.griffin@leeds.ac.uk; a.samson@leeds.ac.uk; r.hughes1@leeds.ac.uk

Word count: 6552

Number of figures: 7

COMPETING INTERESTS

All authors declare that there are no conflicts of interest.

FINANCIAL SUPPORT STATEMENT

This research was funded by the Medical Research Council (Award No: MR/T016205, awarded to SG), and Cancer Research UK (Award No: 29039, awarded to AS).

Key words: Reovirus, MHCII-independent, IFNB, Granzyme B, Perforin, CD4, cytotoxic.

AUTHOR CONTRIBUTIONS

Conceptualization, R.H., S.G, A.S., A.M[2]., and A.M[3]; *Methodology*, R.H., A.M[1]., S.G.C., and S.G; *Investigation*, R.H., A.S., S.G.C., K.J.S., E.A., M.B., and A.M[1]; *Writing – Original*, R.H. and S.G; *Writing – Review & Editing*, R.H., A.S., A.M[1]., A.M[2]., A.M[3]., E.A., K.J.S., M.B., and S.G; *Funding Acquisition*, A.S., A.M[2], A.M[3]., E.I., and S.G.

ABSTRACT

Hepatocellular carcinoma (HCC) is the third most common cause of cancer deaths worldwide. Combination immunotherapy is now standard of care for advanced HCC, improving patient outcomes. However, a considerable number of patients remain unresponsive, or are unable to tolerate therapy. Tyrosine kinase inhibitors (TKIs), such as the former first-line agent sorafenib, remain an option for such patients, yet provide only marginal efficacy. We hypothesised that a clinically advanced immunogenic "oncolytic" virus, namely, human *Orthoreovirus*, might improve TKI mediated therapy. Surprisingly, *uv*-inactivated, replication-deficient reovirus, but not live virus, significantly extended survival when combined with sorafenib in preclinical immunocompetent HCC models. Favourable responses were dependent upon adaptive immunity, mediated by IFNB-induced skewing of the infiltrating T-cell ratio in favour of cytotoxic CD4$^+$ T-cells expressing granzyme B and perforin. Interestingly, this subset effectively killed tumours via both contact juxtacrine and paracrine processes, the former being MHCII independent. Moreover, efficacy correlated with more rapid and robust IFN production by inactivated virus due to the absence of innate viral antagonists. Thus, we reveal a means to improve TKI-HCC outcomes through an alternative virus-driven immunotherapy, underpinned by non-classical immunological mechanisms.

IMPACT AND IMPLICATIONS

Immune checkpoint immunotherapy is revolutionising cancer treatment, yet considerable numbers of patients still fail to respond and must resort to older, more toxic and less effective therapies, including Sorafenib for the management of HCC.

We demonstrate that burgeoning virus-driven immunotherapy can be successfully combined with Sorafenib to extend preclinical HCC survival, but only when the virus is *uv*-inactivated to prevent already attenuated innate immune antagonism, specifically increasing the magnitude of tumour IFNB responses. IFNB was essential to promote tumour infiltration of cytotoxic CD4+ cells during therapy, which was a hallmark of long-term survival mediated by ensuing adaptive responses.

We anticipate this work will be of interest to clinicians and cancer immunology researchers, promoting closer inspection of the immune microenvironment and cancer-specific responses to OV therapy, specifically those driven by non-canonical anti-cancer mechanisms involving IFNB and cytotoxic CD4+ T-cells.

INTRODUCTION

Hepatocellular carcinoma (HCC) is the third leading cause of cancer deaths

worldwide. A significant proportion of HCC patients (~40%) present with advanced disease

at diagnosis, excluding them from potentially curative surgery. Consequently, ~88% of

patients succumb within 5-years post-diagnosis across disease stages. Recently,

immunotherapy combinations, including licensed treatments targeting PD-L1 and VEGF-A

have dramatically improved clinical outcomes, dependent upon a favourable immunological

tumour microenvironment. Patients failing to respond revert to prior systemic TKI therapy,

most commonly sorafenib. The survival benefit gained following sorafenib treatment is

limited[1], as is patient compliance. Thus, considerable need exists for novel therapeutic

approaches that augment existing TKI efficacy.

Oncolytic viruses (OV) are a promising area of cancer immunotherapy and are well tolerated

by patients. Numerous OVs are in clinical trials for a range of solid and hematological

malignancies. OV-therapies are known to exert their anti-cancer activity by both direct

tumour cell lysis and stimulation of anti-tumour immunity. This arises due to innate

responses, direct leukocyte stimulation, and the liberation of PAMPs, DAMPs and other cell

components including tumour antigens. The balance between how these processes combine

to achieve efficacy logically varies according to both the OV, as well as the tumour in

question.

Oncolytic human *Orthoreovirus* (type-3, Dearing Strain, supplied as Pelareorep, Oncolytics

Biotech, Calgary, CA. Referred to herein as "Reo") has been both widely and safely used in

numerous cancer clinical trials and is currently under fast-track review for both metastatic

breast cancer and advanced/metastatic pancreatic cancer[2]. Although Reo exerts direct lytic

effects following replication in cancerous cells, there is considerable interest in its

pronounced ability to activate the immune system[3-7]. The segmented, dsRNA genome of

Reovirus, including its terminal diphosphate, can be detected by innate cellular pattern

recognition receptors (PRR) including endosomal Toll-like receptors (TLR3) and cytosolic

RIG-like receptors[8] (RIG-I and MDA-5). Upon activation, PRRs trigger signalling cascades

that drive host cells to produce a wide variety of inflammatory cytokines, particularly

interferons. These play a major role inducing an anti-viral state by upregulating the

expression of interferon-stimulated genes (ISGs), but also stimulate and enhance immune

cell function. In the context of cancer, Reo stimulates antigen-specific cytotoxic $CD8^+$ T-cell

responses[4-7] capable of breaking tumour immune tolerance in pre-clinical mouse models[5, 6].

Immune activation is also observed in Reo treated patients, evidenced by increased

expression of the interferon gamma-inducible immune checkpoint molecule, PD-L1[9],

increased levels of anti-Reo neutralising antibodies in peripheral blood[10, 11], and the

accumulation of intra-tumoural T-cells[12].

$CD4^+$ T-helper cells (T_H-cells) are critical master co-ordinators of the adaptive immune

system. In addition, under specific conditions T_H-cells exert direct toxicity against a range of

cell types. Cytotoxic $CD4^+$ T_H-cells (CTHs) are frequently observed in patients with chronic

viral infections, including both human cytomegalovirus (hCMV)[13] and human

immunodeficiency virus (HIV)[14], as well as in pre-clinical models of lymphocytic

choriomeningitis (LCMV) [15].

CTHs resemble *bona fide* cytotoxic $CD8^+$ T-cells (CTLs) by expressing granzyme-B and

perforin, although rather than MHCI, they exert MHCII-dependent, antigen-specific cell

killing. T_H-cells also deploy granzyme/perforin-independent cytotoxicity, including FasL/Fas

and TNF-related apoptosis-inducing ligand (TRAIL), which are involved in maintenance of

peripheral tolerance through activation-induced cell death (AICD), and the elimination of

malignant or virus-infected cells[16, 17].

Here, we describe a striking observation whereby *uv*-inactivated, replication-deficient Reo

(*uv*-Reo), but *not* live Reo, dramatically extended survival in a pre-clinical, syngeneic,

immunocompetent HCC model when combined with sorafenib. Whilst long-term protection

relied upon adaptive responses, its inception was critically dependent upon expression of

IFNB, and a T_H1-dominated anti-tumour immune response underpinned, surprisingly, by

multiple MHCII-*independent* modes of killing.   Mechanistically, tumour-borne expression of

the T_H1-cell tropic chemokine, CCL5, led to a markedly increased CD4:CD8 ratio amongst

tumour-infiltrating lymphocytes (TIL) during therapy. *uv*-Reo/sorafenib induced CTH

exhibited both paracrine as well as proximity-dependent tumour cell killing. Sorafenib

sensitised tumour cells to secreted TNFA and IFNG produced by CTHs whereas IFNB

engaged a proximity-dependent mode of killing that was reliant on granzyme B and perforin,

but *not* MHCII. The superiority of the *uv*-Reo response was attributable to the absence of

viral IFN antagonists, raising questions over the possible limitations of other, less attenuated,

virus-driven immunotherapies.

MATERIALS & METHODS

Mouse models

All *in vivo* experiments were conducted with the approval of the University of Leeds

Applications and Amendments (Ethics) Committee and in accordance with UK Home Office

regulations (PP1816772). BALB/c, SCID, and SCID/Beige mice were housed in isolator

cages with 12-h light/dark cycles at 22^oC with access to food and water *ad libitum.* For

overall survival studies, female mice, aged 7 – 8 weeks, were implanted subcutaneously

(*s.c.,*) with 1MEA cells in 100 μL of PBS. Once palpable (~2 – 3 mm in diameter), mice were

treated with either Reo or *uv*-Reo ($1x10^7$ pfu) *via* intra-tumoural injection (*i.t.,*), three times

per week for six weeks. Sorafenib (10 mg/Kg) or vehicle (PBS, 25% PEG-400, 5% Tween-

20, 5% ethanol) were administered by oral gavage (*o.g.,*), daily for 4 four weeks. Tumour

diameter was measured in two dimensions daily and mice were culled when they reached 15

mm in diameter as a proxy for cancer-induced mortality. For histological, proteomic and

RNA analyses, female mice were implanted with murine HCC cells and treated with live- or

uv-Reo alone or in combination with sorafenib or vehicle as described above, for two weeks.

Twenty-four hours after the last treatment the mice were culled by an approved Schedule 1

method and tissue processed accordingly.

Primary cell cultures and cell lines

Human (HLE) and murine (1ME.A7.7R.1 [1MEA]) HCC cell lines, and primary immune

cells, were maintained in humidified incubators with 5% CO_2 in DMEM supplemented with

2 mM *l*-glutamine, 10% FBS, and 1% non-essential amino acids. Human and murine T-cells

were isolated by a combination of positive selection using antibody-conjugated magnetic

beads directed against CD4 and CD8 for whole blood (Human) or negative selection from

spleens and lymph nodes (Mouse). Activation of human and murine T-cells was performed

using plate-bound anti-CD3 and medium supplemented with anti-CD28 for three days, with

or without recombinant IL2 and IL12 for T_H1 polarisation.

Chemotaxis assay

$CD4^+$ T-cells were serum-starved for two hours in chemotaxis buffer (RPMI-1640 and 0.5 %

BSA) prior to assay. For each experiment, $3x10^5$ $CD4^+$ T-cells, labelled with CFSE, were

seeded into tissue culture inserts (5 μm pore size), in 24-well plates, in chemotaxis buffer in

the presence or absence a CCR5 inhibitor (Maraviroc, 1 μM) or vehicle. The lower

compartment contained either chemotaxis buffer alone or was supplemented with CCL5 (100

ng/mL) with or without Maraviroc or vehicle. Chemotaxis assays were stopped after 1.5

hours and cells counted using flow cytometry.

Immunofluorescence

Tumour cryosections 14 μm thick were fixed with either ice-cold acetone or 4%

paraformaldehyde, blocked in appropriate serum and incubated with fluorophore-conjugated

primary antibodies (2 – 5 μg/mL) in PBS, overnight at 4^oC. Nuclei were counter-stained

using DAPI and labelled sections were mounted in ProLong Diamond antifade reagent.

Random fields of view (F.O.V.) were acquired using both a Nikon A1R and Zeiss LSM 980

confocal microscopes. Image processing and quantification was performed using 'Fiji'

Image.

Proteomics

Cytokine arrays were performed using tumour protein extracts generated from snap-frozen

biopsies using a combination of bead mill and freeze/fracture in PBS. Mouse cytokine arrays

were incubated with 1 mg pooled total protein. Membranes were developed using Pierce^{TM}

chemiluminescent substrate with a ChemiDoc^{TM} imaging system and image quantification

was performed using 'Fiji' Image J. For ELISAs, clarified supernatants were generated from

human and mouse HCC cell lines following incubation overneght with Reo/*uv*-Reo (2

PFU/cell) in the presence or absence of sorafenib (7µM), and from T-cells at three days post-

activation.

Flow cytometry

Antibody labelling of T-cells and HCC cell lines was performed in staining buffer (HBSS +

0.5% BSA), on ice, using directly-conjugated antibodies (1 – 5 µg/mL). For cryopreserved

tumour biopsies, single cell suspensions were generated by passing tumours through a 70 µm

nylon mesh with subsequent labelling of both cell surface and intra-cellular antigens using a

CytoFix/CytoPerm Kit. Data were collected using a CytoFLEX S flow cytometer.

HCC/T-cell co-culture killing assays

For direct co-culture killing assays, human and murine CFSE-labelled HCC target cells

($1.5x10^4$) were co-cultured with near infra-red dye-labelled T-cells at a ratio of 50:1 in the

presence or absence of IFNB, with or without neutralising antibodies, EGTA, GZMB

inhibitor (z-AAD-CMK) or Caspase inhibitor (z-VAD-FMK), where indicated. Following

overnight incubation, all cells were collected and stained with Zombie UV viability dye then

fixed in 4% PFA prior to analysis. Target cell killing ($CFSE^+NIR^{neg}$) was determined using a

CytoFLEX S flow cytometer. For indirect co-culture killing assays, T-cells and target cells

were separated using tissue culture inserts with a 0.4 µm pore size.

RNASeq and immune deconvolution

RNA samples were extracted from 1MEA tumours using an RNeasy mini kit and subjected to

Illumina sequencing (Novogene UK Ltd), with a sequencing depth of 20 million reads.

RNASeq data were uploaded to the TIMER2.0 online immune estimation resource and the

xCell immune deconvolution algorithm was applied[18-20].

Statistics

All figures and statistical analyses were performed using Prism software (GraphPad, San

Diego, CA). All data presented are expressed as means ± standard error of the mean (SEM)

and were analysed by one-tailed or two-tailed unpaired Student's t-test where appropriate. P

values less than 0.05 were considered statistically significant and marked as follows; *

$p < 0.05$, ≠ $p < 0.01$, + < 0.001, ^ < 0.0001. Sample size (n) is indicated where appropriate in

figure legends.

RESULTS

Suppression of HCC tumour growth during *uv*-Reo/sorafenib therapy involves the

skewing of intra-tumoural T-cell ratios (T_H:CTL) in favour of CD4$^+$ T_H-cells.

We investigated the therapeutic effect of combining sorafenib with Reo in Balb/c

mice bearing syngeneic, subcutaneous 1MEA HCC tumours, controlling for virus gene

expression-mediated effects by including *uv*-inactivated, replication-deficient virus (*uv*-Reo -

Fig. S1A-**B**). Surprisingly, the combination of *uv*-Reo and sorafenib significantly extended

the survival of tumour-bearing mice well beyond that of all other treatment groups (**Fig. 1A**),

a response that was greatly reduced in SCID mice and lost entirely in SCID/Beige, supporting

an immune-mediated mechanism significantly driven by the adaptive response (**Fig. 1B**).

Thus, we examined the leukocyte composition of 1MEA tumours, grown in Balb/c hosts,

harvested when responding to therapy (Therapy phase – **Fig 1A** & **Fig. 1C**), to identify

changes in infiltrating immune cell(s) that contribute to the subsequent improved survival

phenotype. The only leukocyte population found to be increased in tumours treated with *uv*-

Reo/sorafenib combination therapy relative to control animals was the CD3$^+$CD4$^+$ T_H-cell,

but not CD3$^+$CD8$^+$ CTLs (**Fig. 2A – C** & **Fig. S2**). This significantly skewed the T-cell ratio

in favour of CD4$^+$ cells (**Fig. 2D**) implying that the response to therapy that underpinned the

ensuing improved survival was dominated by T_H-cells.

Cytokine array analysis (**Fig. S3A**) revealed increased intra-tumoural IFNG and TNFA (**Fig.**

3A) coincident with the accumulation of $CD3^+CD4^+$ T-cells. These cytokines are known to

be expressed by T_H1-activated T-cells and, accordingly, were detected in tumour-infiltrating

$CD3^+CD4^+$ cells in mice treated with *uv*-Reo/sorafenib therapy (**Fig. 3B**). By contrast, levels

of cytokines associated with T_H2 and T_H17 T-cell subsets did not mirror the pattern of $CD4^+$

T-cell recruitment (**Fig. S3B**).

We also observed elevated levels of the T_H1-cell chemokine CCL5 coincident with the

increased abundance of $CD3^+CD4^+$ T-cells (**Fig. 3C**). *In vitro*, the expression of CCL5 by

human and murine HCC cells was significantly increased by treatment with *uv*-Reo compared

to Reo, even in the presence of sorafenib (**Fig. S3C - D**). CCL5 exerts a potent chemotactic

effect on T_H1-cells, but not naïve $CD4^+$ T-cells, and a specific small molecule inhibitor

supported that this chemotaxis was mediated by the chemokine receptor CCR5 (**Fig. 3D**).

This was readily detected on the surface of T_H1-activated T-cells *in vitro,* and on intra-

tumoural $CD3^+CD4^+$ T-cells from mice treated with *uv*-Reo/sorafenib (**Fig. 3E**). These data

indicate that the combination of *uv*-Reo and sorafenib exerts a suppressive effect on

continuing HCC growth by skewing the intra-tumoural T-cell ratio in favour of $CD4^+$ T_H1-

cells, not $CD8^+$ CTLs.

T_H1-activated $CD4^+$ T-cells exert contact-independent tumouricidal activity against

HCC cells via a TNFA-dependent mechanism.

We co-cultured HCC cells with $CD4^+$ T-cells to determine whether the latter

possessed anti-tumour properties. T_H1-cells displayed measurable tumouricidal activity (**Fig.**

4A) that did not require the two cell types to be in physical proximity, demonstrating that a

soluble factor was responsible (**Fig. 4B**). Thus, we examined the tumouricidal activity of

soluble mediators released by T_H1-cells known to have cytotoxic properties. We discovered

that TNFA (**Fig. 4C – D**), not IFNG (**Fig. S4A**), was a direct-acting tumouricidal factor

produced by T_H1-cells. However, IFNG enhanced the sensitivity of HCC cells to TNFA

killing by T_H1-cells in co-culture assays (**Fig. S4B, Fig. 4E** – *left panel*).

Next, we treated HCC cells with sorafenib, using a clinically relevant dose, to determine

whether this could further modify the responses to TNFA and IFNG. Sorafenib significantly

enhanced the anti-tumour activity of TNFA against human HCC cells (**Fig. 4E** – *right panel*),

but not IFNG (**Fig. 4F** and **Fig. S4D**), and these findings were replicated using the mouse

HCC cell line used for *in vivo*. A contribution from T_H1-derived factors TRAIL and LTA was

excluded based on antibody neutralisation experiments in co-culture assays (**Fig S4E – F**).

Together, these data demonstrated that TNFA was the dominant soluble mediator of T_H1-

cellular cytotoxicity towards HCC, with a tumour-sensitising role for both IFNG and

sorafenib. However, because this difference alone did not explain the improved efficacy of

the *uv*-Reo/Sorafenib combination, we reasoned that additional mechanisms likely enhance

tumour killing for this treatment combination.

IFNB induces a close-contact MHCII-independent tumouricidal activity in T_H1-cells

mediated by granzyme B/perforin.

Type-I interferons are important mediators of anti-viral immunity *via* upregulation of

interferon-stimulated genes (ISGs), as well as enhancing immune-mediated killing of infected

cells. We, and others, have shown that Reo and *uv*-Reo are potent inducers of the type-I

interferon, 'IFNB'[7, 21] particularly in the context of both HCC and primary liver tissue[21].

Thus, we compared the levels of IFNB produced by both human and mouse HCC cells *in*

vitro, and mouse tumours *ex vivo,* when treated with Reo/*uv*-Reo monotherapy or in

combination with sorafenib. Human and mouse HCC cells responded to *uv*-Reo (alone and

in combination with sorafenib) with a robust induction of IFNB expression, which,

surprisingly, was significantly larger than the response to Reo (**Fig. 5A - B**).   Next, we

examined the intracellular signalling events in HCC cells treated with Reo/*uv*-Reo alone and

in the presence of sorafenib to better understand the differential induction of IFNB. The

transcription factors NF*k*B p65 (RELA) and IRF3 were phosphorylated to a greater extent in

HCCs treated with *uv*-Reo compared to Reo, both in the presence and absence of sorafenib,

as determined by Western blot (**Fig. 5C**). Therefore, we compared the contribution made by

each of these factors to the expression of IFNB using luciferase reporter assays containing

either IRF3 (PD116) or NF*k*B p65 (PRDII) binding elements from the IFNB promoter. The

IRF3 reporter was more strongly activated in HCC cells infected with *uv*-Reo (in both the

presence and absence of sorafenib) compared to Reo (**Fig. S5A**). However, the opposite was

true for the NF*k*B reporter with more robust activation in the presence of Reo, not *uv*-Reo

(**Fig. S5B**). These data indicated that the differential activation and functional output from

these pathways contributed the difference in IFNB induction by Reo/*uv*-Reo.

Viruses dedicate considerable proportions of their coding capacity towards the production of

interferon antagonists within infected cells. Reovirus is no exception[22, 23] so we hypothesised

this may account for the differential induction of IFNB observed in HCC cells. Consistently,

infecting HCC cells with Reo prior to *uv*-Reo significantly dampened the induction of IFNB

compared to *uv*-Reo alone (**Fig. 5D**). In addition, the kinetics of virus-induced IFNB

expression were slower with Reo compared to *uv*-Reo, indicating a delay in viral sensing

(**Fig. 5E**). Critically, this effect was dependent upon Reo replication, and so presumably gene

expression, as a nucleotide analogue inhibitor of the RNA-dependent RNA polymerase, 2'-C-

methylcytidine, ameliorated the suppressive effect (**Fig. 5D**). These data indicated that a

virulence factor could account for the differential induction of IFNB between Reo and *uv*-

Reo, *in vitro*. Accordingly, Reo μNS protein is known to redirect IRF3 to viral replication

factories[22], consistent with decreased levels of phosphorylated protein within Reo infected

HCC cells (**Fig. S5A**). Thus, whilst attenuation of the Type 3 Dearing strain may be severe

compared to other *Orthoreoviruses*, it is by no means absent.

Interestingly, the highest level of intra-tumoural IFNB was found in tumours from mice

treated with *uv*-Reo/sorafenib therapy (**Fig. 6A**), affirming that the magnitude of the IFNB

response might play an important mechanistic role in suppressing tumour growth. Thus, we

added recombinant IFNB to co-culture assays comprising HCC cells and T_H1-cells to

determine how this affected target cell killing. IFNB significantly enhanced the tumouricidal

activity of T_H1-cells (**Fig. 6B**) but, unlike TNFA, did so *via* a mechanism that required

proximity between the two cell types (**Fig. 6C - D**). Next, we examined the effect of pre-

treating HCC target cells with IFNB to determine whether the enhanced killing was a

consequence of either increased target cell sensitivity or the augmented killing capacity of

T_H1-cells. Consistent with the latter, target cell killing was only enhanced when IFNB was

added concurrently with T_H1-cells (**Fig. 6E**) suggesting that the IFNB acted either directly on

CD4$^+$ T-cells alone or simultaneously on both cell types. Moreover, shRNA-mediated

knockdown of IFNB in mouse 1MEA cells eliminated the efficacy of *uv*-Reo/Sorafenib

therapy, *in vivo* (**Fig. 6F**) and this correlated with a failure to recruit CD4$^+$ T-cells (**Fig. 6G**).

We next investigated potential juxtacrine mediators of IFNB-enhanced T_H1-mediated killing

of HCC cells, namely TNF-related apoptosis-inducing ligand (TRAIL) and FasL/Fas.

However, inhibiting these pathways in co-culture assays did not inhibit target cell killing

(**Fig. S6A – G**). Consequently, we considered the involvement of MHCII-dependent

degranulation, described previously for cytotoxic CD4$^+$ T-cells[24]. However, MHCII was

absent on both human and mouse HCC targets in culture, and addition of MHCII-neutralising

antibody to co-culture assays had no effect (**Fig. 7A - B**). Despite the lack of MHCII

involvement, addition of EGTA (a calcium chelating inhibitor of degranulation) to co-culture

assays dose-dependently inhibited IFNB-enhanced target cell killing (**Fig. 7C**). Hence, we

next quantified the proportion of CD4$^+$ T-cells with surface expression of the degranulation

marker CD107a and intra-cellular expression of both granzyme B and perforin, important

components of the apoptosis-inducing machinery deployed by cytotoxic T-cells. We

observed a marked increase in the proportion of human T-cells with surface CD107a

expression and intra-cellular granzyme B (GZMB) following T_H1-activation and this was

further enhanced upon stimulation with IFNB (**Fig. 7D – E**). A smaller, but statistically

significant, change in intra-cellular perforin (PRF) was also detected. Similar results were

obtained for mouse CD4$^+$ T-cells (**Fig. S7A**). GZMB and PRF were readily detected in

supernatants from T_H1-activated T-cells, and levels were further increased upon treatment

with IFNB (**Fig. 7F**). Accordingly, addition of a small molecule GZMB inhibitor to co-

culture assays dose dependently inhibited IFNB-enhanced killing (**Fig. 7G** – *left panel*). The

same response was also seen in the presence of a pan-caspase inhibitor (**Fig. 7G** – *right*

panel). Taken together, these data demonstrate that IFNB enhanced the tumouricidal activity

of CD4$^+$ T-cells by increasing the expression and/or licensing of GZMB/PRF activity against

MHCII-negative HCC cells.

Finally, given that CD8$^+$ cells were also present, but not enriched, within tumours treated

with *uv*-Reo and sorafenib (**Fig. 2**), we compared their ability to engage in antigen-

independent HCC killing with that of CD4$^+$ T-cells, *in vitro*. Consistent with the notion that

CD8$^+$ T-cells played a lesser role in tumour response to therapy, whilst they could indeed kill

HCCs upon activation and this was further enhanced by IFNB, their tumouricidal activity was

significantly lower than that of CD4$^+$ T-cells (**Fig. S7B**). Furthermore, CD8$^+$ T-cell

tumouricidal activity required neither TNFA (**Fig. S7C**) nor degranulation (**Fig. S7D**) and

was therefore distinct mechanistically from that of CD4$^+$ T-cells.

Thus, we conclude that the magnitude of the initial IFNB response, resulting from a lack of

viral antagonist, underpins the improved efficacy of   the *uv*-Reo/Sorafenib therapy in

preclinical HCC models. The induction of IFNB was associated with increased expression of

CCL5 and supported the accumulation of T$_H$1-activated CD4$^+$ T-cells with basal tumouricidal

activity, facilitated by their expression of TNFA and IFNG. Crucially, IFNB licenses and/or

focusses the cytolytic activity of GZMB and PRF, derived from T$_H$1-cells, against MHCII-

negative tumours.

DISCUSSION

This study demonstrates the superior therapeutic efficacy of a unique immunotherapy

combining *uv*-inactivated reovirus with the targeted agent sorafenib over either sorafenib

monotherapy or combination with live Reo, as a treatment for HCC. This combination

therapy significantly extended the survival of mice bearing HCC tumours by engaging a

multi-faceted, MHCII-independent response from cytotoxic $CD4^+$ T_H-cells (CTHs), induced

by IFNB.

The *uv*-Reo/sorafenib therapy specifically induced increased expression of the chemokine

CCL5 and cytokine IFNB relative to other treatment modalities. This was accompanied by

increased abundance of intra-tumoural $CCR5^+$ T_H1-activated $CD4^+$ T-cells, skewing the T-cell

ratio in favour of T_H1-cells. Tumour-infiltrating lymphocytes (TILs) exerted basal levels of

paracrine tumouricidal activity *via* TNFA, which was enhanced in HCC cells by both IFNG

and sorafenib. Concomitantly, increased expression of IFNB in HCC cells following

stimulation with *uv*-Reo/sorafenib was a crucial mechanistic switch, stimulating MHCII-

independent degranulation involving GZMB and PRF, requiring close cell-cell contact.

Hence, our findings support that *uv*-inactivated reovirus could be used to enhance the efficacy

of sorafenib during the treatment of HCC by engaging MHCII-independent $GZMB^+PRF^+$

T_H1-cells.

The role of TNF in liver cancer is complex and studies can be contradictory. A considerable

body of evidence details a pro-tumourigenic role for TNF in the development of liver

cancer[25]. However, in pre-clinical cancer models, direct intra-tumoural injection of TNF

resulted in widespread necrosis, an effect attributed to its anti-vascular activity, rather than

direct tumour cytolysis[26-28]. The consensus is, therefore, that TNF is a weak direct-acting

cytolytic, consistent with findings herein. However, this limited tumouricidal activity can be

significantly enhanced by co-administration of IFNG, a phenomenon demonstrated in a

variety of tumour types, including melanoma, breast, and colon cancer[26, 27], confirmed again

here. We now add that sorafenib also sensitised HCCs to TNFA-induced cell death,

reminiscent of responses described previously for other TNF family members including

TRAIL and Fas, attributed to downregulation of the anti-apoptotic protein, Mcl-1[29] in tumour

cells. Although it is likely that the response to TNFA is enhanced by sorafenib in a similar

way, the precise mechanism has yet to be determined. Interestingly, we found that

TNFA/IFNG-mediated HCC killing by T_H1-cells occurred in the presence of the pan-caspase

inhibitor Z-VAD-FMK, implicating a caspase-independent pathway.

These data provide an important mechanistic insight and suggest that the efficacy of *uv*-

Reo/sorafenib therapy is mediated, at least in part, by a skewing of T-cell ratios in favour of

$IFNG^+$ $TNFA^+$ T_H1-cells. However, IFNG and TNFA alone do not fully explain how T_H1-

cells control tumour growth in response to *uv*-Reo/sorafenib and not in other treatment

groups where they are also equally abundant.

We, and others, have demonstrated that *uv*-Reo elicits a significantly more robust induction of

IFNB in treated cells compared with Reo[21, 30], confirmed again, here, in both human and

murine HCC cells. Our analysis indicates a Reo-derived antagonist virulence factor is

responsible for the differential induction of IFNB by Reo/*uv*-Reo, as well as the ability of

Reo to suppress ensuing responses (**Fig 5C**). Several factors have been reported including

the non-structural mu protein (μNS)[22] and the outer capsid σ3[23]. It may be that one or both

contribute to suppression of IFNB in HCCs infected by Reo, although increased IRF3

phosphorylation in cells exposed to *uv*-Reo is consistent with a lack of μNS function, rather

than σ3 which predominantly antagonises PKR.

The genetic determinants of reovirus strain variability, specifically relating to interferon

antagonism and apoptotic potential, have been identified[31, 32]. Type-3 Dearing strain (T3D)

shows a reduced ability to suppress type-I interferon responses[31, 32] and demonstrates

enhanced pro-apoptotic activity in infected cells[31]. This strain has been extensively evaluated

as a oncolytic agent in clinical trials. However, in the context of HCC, T3D's attenuation

appears insufficient, failing to elicit a robust interferon response or effectively control tumour

growth. This finding suggests that many natural or genetically attenuated OVs currently in

clinical trials might not be adequately attenuated for all cancer types, particularly those that

are considered immunologically cold. This raises the question of whether the enhanced IFNB

response induced by *uv*-inactivated viruses could lead to superior clinical efficacy in these

cancer types. Interestingly, the TKI sorafenib was key to the robust induction of IFNB that

underpinned the $CD4^+$ T-cell response, despite being naturally immunosuppressive. This

raises the question of whether *uv*-inactivated viruses might be effective in other cancer types

treated with sorafenib and similar TKIs.

This study demonstrates that IFNB induces a unique GZMB/PRF-mediated tumouricidal

activity in T_H1-cells, independent of MHCII expression by target HCCs. Although the

expression of GZMB and PRF by CD4[+] T-cells is reported in the context of viral infection,

this mode of killing is both antigen-specific and reliant on MHCII[33, 34]. How an immune

synapse is able to form between IFNB-stimulated T_H1-cells and MHCII-negative tumour

cells, thereby inducing target cell death, is not fully understood but, a mechanism involving

the upregulation of such stress-induced NKG2D (present upon activated T_H1-cells) ligands,

in conjunction with ICAM1/LFA1, has been proposed[35-39]. Interestingly, NKG2D ligands can

by induced within tumour cells following viral infection and in response to type-I

interferon[38, 40].

CD8[+] T-cells were also present in tumours treated with *uv*-Reo/sorafenib therapy, although

their abundance was unchanged relative to controls. Furthermore, CD8[+] T-cells were

surprisingly less tumouricidal than their CD4[+] counterparts in the context of antigen-

independent immunity, and utilised neither TNFA nor GZMB. These data indicate that the

immediate response to therapy with *uv*-Reo/sorafenib is dominated by T_H1-activated T-cells.

Despite this, we do not discount a major role for antigen-specific immunity during the post-

therapy phase, which contributed significantly to the overall survival advantage (Fig 1). How

initial MHCII-independent responses translate into adaptive immunity remains the focus of

ongoing research.

Taken together, our data demonstrate that *uv*-inactivated reovirus and the targeted agent,

sorafenib, combine to drive IFNB and CCL5 mediated tumour infiltration by MHCII-

independent GZMB[+]PRF1[+] T_H1-cells, capable of establishing subsequent long-lasting tumour

survival despite the cessation of therapy. This also highlights an overlooked role for CD4[+]

T_H1-cells in mediating MHC-independent anti-tumour immunity. Further mechanistic

elaboration is needed to fully understand how non-canonical effector cells can be further

exploited as a therapeutic tool for the treatment of HCC.

ACKNOWLEDGEMENTS

Funding: The authors thank the Medical Research Council (Award No: MR/T016205,

awarded to SG), and Cancer Research UK (Award No: 29039, awarded to AS) for funding

this research. We also thank the NHS Blood and Transplant Service for the provision of donor

blood samples, the St. James's Biological Services Unit (SBS) for their support with animal

welfare and husbandry, and Oncolytics Biotech Inc. for supplying clinical grade human

Orthoreovirus, type-3 Dearing strain, Pelareorep.

REFERENCES

1.   Llovet, J.M. *et al.* Sorafenib in advanced hepatocellular carcinoma. *N Engl J Med*
359, 378-390 (2008).

2.   Bernstein, V. *et al.* A randomized phase II study of weekly paclitaxel with or without
pelareorep in patients with metastatic breast cancer: final analysis of Canadian
Cancer Trials Group IND.213. *Breast Cancer Res Treat* **167**, 485-493 (2018).

3.   Clements, D.R. *et al.* Newly recruited CD11b+, GR-1+, Ly6C(high) myeloid cells
augment tumor-associated immunosuppression immediately following the therapeutic
administration of oncolytic reovirus. *J Immunol* **194**, 4397-4412 (2015).

4.   Errington, F. *et al.* Reovirus activates human dendritic cells to promote innate
antitumor immunity. *J Immunol* **180**, 6018-6026 (2008).

5.   Gujar, S.A., Marcato, P., Pan, D. & Lee, P.W. Reovirus virotherapy overrides tumor
antigen presentation evasion and promotes protective antitumor immunity. *Mol*
Cancer Ther **9**, 2924-2933 (2010).

6.   Prestwich, R.J. *et al.* Tumor infection by oncolytic reovirus primes adaptive antitumor
immunity. *Clin Cancer Res* **14**, 7358-7366 (2008).

7.   Prestwich, R.J. *et al.* Reciprocal human dendritic cell-natural killer cell interactions
induce antitumor activity following tumor cell infection by oncolytic reovirus. *J*
Immunol **183**, 4312-4321 (2009).

8.   Goubau, D. *et al.* Antiviral immunity via RIG-I-mediated recognition of RNA bearing
5'-diphosphates. *Nature* **514**, 372-375 (2014).

9.   Mahalingam, D. *et al.* A Phase II Study of Pelareorep (REOLYSIN((R))) in
Combination with Gemcitabine for Patients with Advanced Pancreatic
Adenocarcinoma. *Cancers (Basel)* **10** (2018).

10.   Noonan, A.M. *et al.* Randomized Phase 2 Trial of the Oncolytic Virus Pelareorep
(Reolysin) in Upfront Treatment of Metastatic Pancreatic Adenocarcinoma. *Mol Ther*
24, 1150-1158 (2016).

11.   Sborov, D.W. *et al.* A phase I trial of single-agent reolysin in patients with relapsed
multiple myeloma. *Clin Cancer Res* **20**, 5946-5955 (2014).

12.   Mahalingam, D. *et al.* Pembrolizumab in Combination with the Oncolytic Virus
Pelareorep and Chemotherapy in Patients with Advanced Pancreatic
Adenocarcinoma: A Phase Ib Study. *Clin Cancer Res* **26**, 71-81 (2020).

13.   Zaunders, J.J. *et al.* Identification of circulating antigen-specific CD4+ T lymphocytes
with a CCR5+, cytotoxic phenotype in an HIV-1 long-term nonprogressor and in CMV
infection. *Blood* **103**, 2238-2247 (2004).

14.   Norris, P.J. *et al.* Beyond help: direct effector functions of human immunodeficiency
virus type 1-specific CD4(+) T cells. *J Virol* **78**, 8844-8851 (2004).

15.   Jellison, E.R., Kim, S.K. & Welsh, R.M. Cutting edge: MHC class II-restricted killing in
vivo during viral infection. *J Immunol* **174**, 614-618 (2005).

16.   Brunner, T. *et al.* Cell-autonomous Fas (CD95)/Fas-ligand interaction mediates
activation-induced apoptosis in T-cell hybridomas. *Nature* **373**, 441-444 (1995).

17.   Simon, A.K. *et al.* Tumor necrosis factor-related apoptosis-inducing ligand in T cell
446   development: sensitivity of human thymocytes. *Proc Natl Acad Sci U S A* **98**, 5158-
447   5163 (2001).

448   18.   Aran, D., Hu, Z. & Butte, A.J. xCell: digitally portraying the tissue cellular
449   heterogeneity landscape. *Genome Biol* **18**, 220 (2017).

450   19.   Li, B. *et al.* Comprehensive analyses of tumor immunity: implications for cancer
451   immunotherapy. *Genome Biol* **17**, 174 (2016).

452   20.   Sturm, G. *et al.* Comprehensive evaluation of transcriptome-based cell-type
453   quantification methods for immuno-oncology. *Bioinformatics* **35**, i436-i445 (2019).

454   21.   Samson, A. *et al.* Oncolytic reovirus as a combined antiviral and anti-tumour agent
455   for the treatment of liver cancer. *Gut* **67**, 562-573 (2018).

456   22.   Stanifer, M.L., Kischnick, C., Rippert, A., Albrecht, D. & Boulant, S. Reovirus inhibits
457   interferon production by sequestering IRF3 into viral factories. *Sci Rep* **7**, 10873
458   (2017).

459   23.   Imani, F. & Jacobs, B.L. Inhibitory activity for the interferon-induced protein kinase is
460   associated with the reovirus serotype 1 sigma 3 protein. *Proc Natl Acad Sci U S A*
461   **85**, 7887-7891 (1988).

462   24.   Oh, D.Y. *et al.* Intratumoral CD4(+) T Cells Mediate Anti-tumor Cytotoxicity in Human
463   Bladder Cancer. *Cell* **181**, 1612-1625 e1613 (2020).

464   25.   Park, E.J. *et al.* Dietary and genetic obesity promote liver inflammation and
465   tumorigenesis by enhancing IL-6 and TNF expression. *Cell* **140**, 197-208 (2010).

466   26.   Balkwill, F.R. *et al.* Human tumor xenografts treated with recombinant human tumor
467   necrosis factor alone or in combination with interferons. *Cancer Res* **46**, 3990-3993
468   (1986).

469   27.   Brouckaert, P.G., Leroux-Roels, G.G., Guisez, Y., Tavernier, J. & Fiers, W. In vivo
470   anti-tumour activity of recombinant human and murine TNF, alone and in combination
471   with murine IFN-gamma, on a syngeneic murine melanoma. *Int J Cancer* **38**, 763-
472   769 (1986).

473   28.   Mantovani, A. & Dejana, E. Cytokines as communication signals between leukocytes
474   and endothelial cells. *Immunol Today* **10**, 370-375 (1989).

475   29.   Chen, K.F. *et al.* Sorafenib overcomes TRAIL resistance of hepatocellular carcinoma
476   cells through the inhibition of STAT3. *Clin Cancer Res* **16**, 5189-5199 (2010).

477   30.   Henderson, D.R. & Joklik, W.K. The mechanism of interferon induction by UV-
478   irradiated reovirus. *Virology* **91**, 389-406 (1978).

479   31.   Baty, C.J. & Sherry, B. Cytopathogenic effect in cardiac myocytes but not in cardiac
480   fibroblasts is correlated with reovirus-induced acute myocarditis. *J Virol* **67**, 6295-
481   6298 (1993).

482   32.   Zurney, J., Kobayashi, T., Holm, G.H., Dermody, T.S. & Sherry, B. Reovirus mu2
483   protein inhibits interferon signaling through a novel mechanism involving nuclear
484   accumulation of interferon regulatory factor 9. *J Virol* **83**, 2178-2187 (2009).

33.   Adhikary, D. *et al.* Control of Epstein-Barr virus infection in vitro by T helper cells
specific for virion glycoproteins. *J Exp Med* **203**, 995-1006 (2006).

34.   Aslan, N. *et al.* Cytotoxic CD4 T cells in viral hepatitis. *J Viral Hepat* **13**, 505-514
(2006).

35.   Homma, S., Komita, H., Sagawa, Y., Ohno, T. & Toda, G. Antitumour activity
mediated by CD4+ cytotoxic T lymphocytes against MHC class II-negative mouse
hepatocellular carcinoma induced by dendritic cell vaccine and interleukin-12.
Immunology **115**, 451-461 (2005).

36.   Revy, P., Sospedra, M., Barbour, B. & Trautmann, A. Functional antigen-independent
synapses formed between T cells and dendritic cells. *Nat Immunol* **2**, 925-931
(2001).

37.   Somersalo, K. *et al.* Cytotoxic T lymphocytes form an antigen-independent ring
junction. *J Clin Invest* **113**, 49-57 (2004).

38.   Esteso, G., Guerra, S., Vales-Gomez, M. & Reyburn, H.T. Innate immune recognition
of double-stranded RNA triggers increased expression of NKG2D ligands after virus
infection. *J Biol Chem* **292**, 20472-20480 (2017).

39.   Markiewicz, M.A. *et al.* Costimulation through NKG2D enhances murine CD8+ CTL
function: similarities and differences between NKG2D and CD28 costimulation. *J*
Immunol **175**, 2825-2833 (2005).

40.   Zhang, C. *et al.* Opposing effects of interferon-alpha and interferon-gamma on the
expression of major histocompatibility complex class I chain-related A in tumors.
Cancer Sci **99**, 1279-1286 (2008).

FIGURE LEGENDS

Figure 1. The survival of mice bearing syngeneic hepatocellular carcinomas is

significantly extended by an immune-mediated mechanism induced by a combination of

sorafenib and *uv*-Reo. (**A**) Overall survival analysis of 1MEA tumour-bearing mice treated

with Reo (*left*) or uv-Reo (*right*) alone or in combination with sorafenib. All treatments were

significant relative to vehicle ($p \leq 0.05$) and *uv*-Reo/Sorafenib was significant compared to *uv*-

Reo ($p < 0.0189$) and sorafenib ($p < 0.0027$) monotherapies. (**B**) Overall survival analysis of

SCID (*left*) and SCID/Beige (*right*) mice bearing 1MEA tumours treated as with *uv*-Reo

alone or in combination with sorafenib. *uv*-Reo/sorafenib was significant compared to all

groups in SCID mice only ($p \leq 0.01$). (**C**) Tumour volumetric data from immunocompetent

Balb/c mice bearing 1MEA tumours treated with Reo/*uv*-Reo monotherapy or in combination

with sorafenib and culled after two-weeks of therapy for IF analysis (n = 5 mice per

condition).

Figure 2. CD4$^+$ T-helper cells, but not CD8$^+$ CTLs, accumulate in murine hepatocellular

carcinomas responding to the combination of *uv*-reovirus and sorafenib. (**A** – **B**)

Representative images of 1MEA tumour cryosections taken from tumours harvested midway

through the "therapy" phase, stained for CD3 and CD8 (*left*) or CD4 (*right*) then

counterstained with DAPI. (**C**) Quantification of CD3$^+$CD8$^+$ (*left*) and CD3$^+$CD4$^+$ (*centre*) T-

cells in random fields of view (FOV) taken from tumour cryosections in the indicated groups.

(**D**) Comparison of CD4$^+$:CD8$^+$ T-cell ratio from quantification in 'C' (Scale bar = 50 µm; n =

5 mice per condition).

Figure 3. Treatment of murine HCC tumours with *uv*-Reo/sorafenib therapy elicits at CD4$^+$ T$_H$1-cell response. (**A**) Cytokine array data for the T$_H$1-cytokines IFNG (*left*) and TNFA (*right*) from pooled tumour protein samples and from the indicated treatment groups. (**B**) ELISA-based quantification of IFNG and TNFA in supernatants from human CD4$^+$ T-cells, *in vitro* (*left*), and flow cytometric detection of intra-cellular IFNG and TNFA in tumour-infiltration CD4$^+$ T-cells in mice undergoing treatment with *uv*-Reo/sorafenib therapy (*right*). (**C**) Cytokine array data as described in 'A' but for CCL5. (**D**) Chemotaxis assay comparing the migratory potential of human CD4$^+$ T-cells towards CCL5 and the dependency on CCR5, using the antagonist Maraviroc (CCR5i – 1μM) or vehicle (DMSO). (**E**) Flow cytometric detection of cell surface CCR5 on human T$_H$1-activated CD4$^+$ T-cells, *in vitro* (*left*), and tumour-infiltrating CD4$^+$ T-cells in mice undergoing treatment with *uv*-Reo/sorafenib therapy (*right* – gated on CD45$^+$CD3$^+$CD4$^+$ cells).

Figure 4. T$_H$1-activated CD4$^+$ T-cells exert a TNFA-dependent tumouricidal activity that is enhanced by tumour cell exposure to IFNG or sorafenib. (**A**) Flow cytometric quantification of target HLE killing by human CD4$^+$ T-cells in direct co-culture or (**B**) following separation of T-cells/target cells by porous (0.4 µm) tissue culture inserts. (**C**) Quantification of HLE target killing in direct co-culture with CD4$^+$ T-cells in the presence of neutralising antibodies against TNFA or IgG control. (**D**) Quantification of HLE cell viability in the presence of increasing concentrations of TNFA alone or (**E**) following pre-treatment with IFNG (*left* - 100 U/mL) or sorafenib (*right* - 7 μM). (**F**) Quantification of 1MEA cell

viability in response to TNFA alone or following pre-treatment with IFNG (*left* – 100 U/mL)

or sorafenib (*right* – 7 µM) (n = 4 – 8 per condition).

Figure 5. The differential induction of IFNB by Reo/*uv*-Reo can be attributed to

virulence factor(s). Quantification of IFNB in supernatants from (**A**) human HLE cells or

(**B**) mouse 1MEA cells, by ELISA. (**C**) Representative Western blots revealing activation of

intracellular signalling pathways in HLEs following infection with Reo/*uv*-Reo alone or in

combination with sorafenib, 24 hours post-infection (**D**) *IFNB* expression in HLE cells

treated with Reo or *uv*-Reo, alone or sequentially, in the presence or absence of 2'-C-

Methylcytidine. (**E**) Time-course of *IFNB* upregulation in HLE cells treated with Reo or *uv*-

Reo (n = 3 per condition).

Figure 6. IFNB induces a proximity-dependent mode of tumour cell killing by T$_H$1-cells

and delays tumour growth, *in vivo*. (**A**) Quantification of IFNB in protein lysates from

tumour-bearing mice treated with Reo/*uv*-Reo (2 pfu/cell) alone or in combination with

sorafenib (7 µM). (**B**) Quantification of IFNB-induced HLE target cell killing by T$_H$1-

activated CD4$^+$ T-cells in direct co-culture or following separation of T-cells/target cells by

porous tissue culture inserts in (**C**) human and (**D**) mouse systems. (**E**) Quantification of

IFNB-induced killing of HLE target cells when co-administered with T-cells or given to

target cells as a pre-treatment. (n = 4 – 8 per condition). (**F**) Quantification of IFNB released

by 1MEA cells carrying Scrambled or IFNB-targeted shRNAs following treatment with *uv*-

Reo (*left*) and their response to *uv*-Reo/Sorafenib therapy, *in vivo* (*right*). (**G**) Quantification

(*left*) and representative images (*right*) of CD3$^+$CD4$^+$ T-cell abundance in 1MEA tumours

carrying Scrambled or IFNB-targeted shRNA constructs treated with *uv*-Reo/Sorafenib

therapy.

Figure 7. IFNB induces an antigen and perforin-independent, granzyme B-dependent mode of tumouricidal activity in T_H1-activated CD4$^+$ T-cells. (**A**) Flow cytometric detection of cell surface MHCII proteins on human and murine HCC cells, *in vitro*. (**B**) Quantification of HLE target cell killing by human T_H1-activated CD4$^+$ T-cells induced by IFNB in the presence of neutralising antibodies against MHCII or (**C**) the degranulation inhibitor EGTA. (**D**) Representative flow cytometry plots showing detection of cell surface CD107a (*left*) and intra-cellular GZMB (*centre*) or PRF (*right*) in human CD4$^+$ T-cells alongside (**E**) quantification of the proportion cells positive for each marker. (**F**) Quantification of cell-free GZMB and PRF in supernatants from human CD4$^+$ T-cells, by ELISA. (**G**) Quantification of HLE target cell killing by human T_H1-activated CD4$^+$ T-cells in the presence of IFNB and a GZMB inhibitor - z-AAD-CMK (*left*) or caspase inhibitor - z-VAD-FMK (*right*). (n = 4 – 8 per condition).

A.

Overall survival experimental time-line

B.

C.

A. HLE/CD4: Co-culture

B. HLE/CD4: Transwell assay

C. HLE/CD4: TNFA neutralization

D. HLE: TNFA viability assay

E. HLE: IFNG sensitization assay

HLE: Sorafenib sensitization assay

F. 1MEA: IFNγ sensitization assay

1MEA: Sorafenib sensitization assay

A. 1MEA Tumour: IFNB

B. HLE/CD4: Co-culture

C. HLE/CD4: Transwell assay

D. 1MEA/CD4: Transwell assay

E. HLE/CD4: IFNB pre-treatment

F. 1MEA: IFNB knockdown

G.