Title

A circadian behavioral analysis suite for real-time classification of daily rhythms in complex behaviors

Authors

Logan J. Perry[1], Blanca E. Perez[1], Larissa Rays Wahba[2], KL Nikhil[2], William C. Lenzen[1], and Jeff R. Jones[1,3,4]

Affiliations

[1]Department of Biology, Texas A&M University, College Station, TX

[2]Department of Biology, Washington University in St. Louis, St. Louis, MO

[3]Institute for Neuroscience, Texas A&M University, College Station, TX

[4]Center for Biological Clocks Research, Texas A&M University, College Station, TX

Abstract

Measuring animal behavior over long timescales has been traditionally limited to behaviors that are

easily measurable with real-time sensors. More complex behaviors have been measured over time, but these

approaches are considerably more challenging due to the intensive manual effort required for scoring behaviors.

Recent advances in machine learning have introduced automated behavior analysis methods, but these often

overlook long-term behavioral patterns and struggle with classification in varying environmental conditions. To

address this, we developed a pipeline that enables continuous, parallel recording and acquisition of animal

behavior for an indefinite duration. As part of this pipeline, we applied a recent breakthrough self-supervised

computer vision model to reduce training bias and overfitting and to ensure classification robustness. Our

system automatically classifies animal behaviors with a performance approaching that of expert-level human

labelers. Critically, classification occurs continuously, across multiple animals, and in real time. As a proof-of-

concept, we used our system to record behavior from 97 mice over two weeks to test the hypothesis that sex and

estrogen influence circadian rhythms in nine distinct home cage behaviors. We discovered novel sex- and

estrogen-dependent differences in circadian properties of several behaviors including digging and nesting

rhythms. We present a generalized version of our pipeline and novel classification model, the "circadian

behavioral analysis suite," (CBAS) as a user-friendly, open-source software package that allows researchers to

automatically acquire and analyze behavioral rhythms with a throughput that rivals sensor-based methods,

allowing for the temporal and circadian analysis of behaviors that were previously difficult or impossible to

observe.

Introduction

Understanding the genetic, neural, and ethological mechanisms that temporally organize behavior is a fundamental goal of fields including circadian biology, neuroscience, and ecology. However, the temporal analysis of behavior has been largely limited to behaviors that can be accurately measured with low latency and at high throughput. For instance, optical and electrical sensors enable such analysis of eating, drinking, and locomotor behaviors. These behaviors have been widely studied (although usually independently) at high temporal resolution for experimental durations of weeks, months, or even years (Schwartz and Zimmerman, 1990; Jud et al., 2005; Pendergast et al., 2013; Yamanaka et al., 2013; Metzger et al., 2020). Other more complex behaviors such as rearing, nesting, and grooming can often be measured simultaneously using video recording and manual behavior scoring by trained human observers (van der Veen et al., 2008; Gaskill et al., 2009; van Oosterhout et al., 2012; Fujita et al., 2017; Robinson-Junker et al., 2018; Shuboni-Mulligan et al., 2021). Over long timescales, however, this method becomes impractical because video acquisition inevitably outpaces human labeling, leading to an ever-increasing latency between data acquisition and data analysis. For example, it may take an observer less than a minute to label behaviors in a minute long video recording but, due to the tedium of the task and the limits of the human attention span, it would take that observer much longer than a week to classify behaviors in a week-long video recording (Segalin et al., 2020; Muller et al., 2021). Consequently, these behaviors have been studied infrequently at low temporal resolution for limited experimental durations, such as hourly over the course of a single day. To better understand how animal behavior changes over time, ethologically relevant behaviors (regardless of "measurability") must be measured in individual animals simultaneously, automatically, and, critically, in real time over multiple days and conditions at high temporal resolution (Peters et al., 2015; Grieco et al., 2021; Kahnau et al., 2023).

Over the last two decades, methods have been developed to classify animal behavior from video recordings, ranging from computer vision algorithms, such as centroid tracking, to machine learning-based approaches that use markerless pose estimation (e.g., DeepLabCut or DLC) or raw pixel values (e.g., DeepEthogram or DEG) to quantify behavior (Mathis et al., 2018; Pereira et al., 2020; Zhang et al., 2020;

Bohnslav et al., 2021). While previous studies have used machine learning to analyze temporal variation in more naturalistic "home cage" behaviors, these methods have faced several challenges (Steele et al., 2007; Goulding et al., 2008; Jhuang et al., 2010; Adamah-Biassi et al., 2014; Salem et al., 2015). For instance, existing methods tend to disregard long-range temporal information by simplifying analysis to frame-wise positional and motion values. A more holistic approach is needed to capture the temporal dynamics of both the recorded animal and potentially changing in-scene objects on both short and long time scales (Xie et al., 2017). Additionally, existing methods are often constrained by specific environmental conditions, such as video perspective, lighting condition, or subject coloration, which greatly limits their applicability. The development of an adaptable, condition-agnostic system is therefore essential for robust temporal analysis. Perhaps most importantly, existing methods have not been typically used for real-time behavior analysis and have not been used to analyze behavior on a "circadian" timescale of days to weeks or longer. Together, these challenges have prevented the widespread adoption of machine learning classification to the long-term or circadian analysis of behavior. This is likely exacerbated by the lack of user-friendly tools that facilitate acquisition, training, validation, and analysis of key behavioral metrics. To solve this problem, we developed a streamlined approach that allows users to extend modern deep learning methods to emulate the functionality of traditional sensor-based analyses of behavior.

Here, we introduce a versatile, high-throughput, real-time behavior acquisition and analysis pipeline for the temporal analysis of behavior. To do this, we created the software infrastructure to automatically acquire behavior data from video recording streams in real time, in parallel (here, from 24 mice simultaneously) for an essentially unlimited experimental duration (here, for up to two weeks continuously at 10 frames per second). Next, we developed a joint long short-term memory and linear layer model to integrate the visual and motion features output by DINOv2, a state-of-the-art self-supervised computer vision feature extractor. Finally, we combined this model with our recording pipeline to facilitate indefinite recording and behavioral analysis. As a proof-of-concept, we trained our classification model to identify nine home cage behaviors (eating, drinking, rearing, climbing, digging, nesting, resting, grooming, and locomotion; (Garner, 2017)) in male and female

mice to test the hypothesis that sex and estrogen influence circadian rhythms in home cage behaviors. Previous

studies have identified subtle sex differences in wheel-running activity rhythms (Krizo and Mintz, 2014; Joye

and Evans, 2022). However, despite the global regulation of behavior by the circadian system, sex differences

in other behavioral rhythms have not yet been identified due to technological limitations. Our automatic

inference system allowed us to discover novel sex- and estrogen-dependent differences in the phase and

amplitude of several behaviors, including, notably, digging and nesting rhythms. Finally, we developed our

DINOv2 model and automatic inference software into a user-friendly, open-source Python package called

CBAS, the "circadian behavioral analysis suite." CBAS allows researchers to automatically acquire and analyze

behavioral rhythms with a throughput that greatly exceeds manual video labeling and rivals sensor-based

methods.

Results

<u>Machine learning classification of behaviors approaches human level performance</u>

We first recorded continuous videos of individually-housed mice for >24 h at 10 fps in both a 12 h:12 h

light:dark (LD, where dark is defined as dim 850 nm infrared light) cycle and in constant darkness (DD) (**Fig.**

1a). From these videos, we used strict criteria (**Supplementary Table 1**) to define and manually label nine

ethologically-relevant behaviors that encompass the majority of an individual singly-housed mouse's daily

behavioral repertoire, including maintenance, exploratory, and inactive behaviors: eating, drinking, rearing,

climbing, grooming, exploring, digging, nesting, and resting (**Fig. 1b**) (Garner, 2017). For each behavior, we

identified the average length of time for a "bout," or behavioral instance (**Fig. 1c**). This allowed us to generate

balanced training and test sets from segments of videos sampled from 30 mice that contained a balanced

number of unique instances of each behavior (**Fig. 1d**). To control for lighting conditions, we sampled video

segments such that there was a balanced representation of each behavior during both the animal's active and

inactive phases.

We next used these training and test sets to train a previously published deep learning behavior

classification model, DeepEthogram (DEG) (Bohnslav et al., 2021), and our own DINOv2+ model (**Fig. 2a**).

We constructed DINOv2+ using the state-of-the-art DINOv2 vision transformer model (Oquab et al., 2023) as a

"frozen," or immutable, feature extractor 'backbone' with a trainable joint long short-term memory and linear

layer classification network 'head.' DEG and DINOv2+ are each capable of producing behavior classifications

from a video frame's raw pixel values as binary output matrices ("ethograms") that indicate if a behavior is

present or absent in a given frame. This temporally sequenced ethogram output is ideal for quantifying

behavioral rhythms because it is readily analyzed using field-standard circadian analysis methods that are

optimized for time series data. However, while DEG is trained using a supervised learning scheme, the

backbone feature extractor of our DINOv2+ model is pretrained using a self-supervised approach that has been

shown to be more generalizable (Tendle and Hasan, 2021). Thus, training and testing both models allows us to

directly compare the performance of these two different underlying learning schemes on visual feature

extraction.

If we wanted to use our models to automatically infer days of video – millions or potentially billions of

frames that would never be seen by a human – it was critical that our models were extensively validated. Model

performances quantified across all measured behaviors of existing commercial (e.g., HomeCageScan) and non-

commercial methods used for the temporal analysis of home cage behaviors are either unreported or, typically,

mediocre. Thus, after training our models, we performed rigorous validation of the model's predictions on our

labeled test sets with stringent model performance thresholds. Importantly, we did not adjust model

hyperparameters based on our model's test set performance. We compared the performance of our DEG and

DINOv2+ models with that of a trained human classifier. Each of these groups were given a 15-31 frame (1.5-

3.1 s) window to predict behaviors from our test set.

First, because most machine learning performance metrics require us to define a specific threshold value

at which behavior probabilities are converted into a binary prediction, we generated precision-recall curves

136 across different probability thresholds for our DEG and DINOv2+ models (**Fig. 2b**). We did not generate

137 human classifier precision-recall curves because in our training set human labels are inherently binary, not

138 probabilistic. We found that the areas under the precision-recall curves (AUPRC, a summarization of model

139 performance as a function of probability cut-off threshold) for our DINOv2+ model greatly outperformed DEG

140 on rearing and exploring behaviors and slightly, but significantly, outperformed DEG on climbing and resting

141 behaviors. DEG slightly, but significantly, outperformed DINOv2+ on digging behavior classification.

143   Next, we used multiple demanding metrics (F1 score, balanced accuracy, and normalized Matthews

144 correlation coefficient; (Brzezinski et al., 2020; Chicco and Jurman, 2020; Grandini et al., 2020)) to test the

145 performance of each of our models (**Fig. 2c**). Our predetermined criteria for a "successful" model was a score of

146 at least 0.80 for each behavior on each metric. A successful model would also ideally meet or exceed the

147 performance of a trained human classifier labeling the same test set. Our DINOv2+ model met or exceeded our

148 predefined threshold on all performance metrics, whereas our DEG model failed to meet this F1 and nMCC

149 score threshold for rearing and exploring behaviors. Notably, DINOv2+ exhibited greater performance than

150 DEG even on behaviors that had F1, balanced accuracy, or nMCC scores of >0.80, such as grooming and

151 resting. DINOv2+ also met or exceeded human classifier performance on metrics for most behaviors, including

152 eating, drinking, climbing, grooming, exploring, digging, and resting, while DEG only met or exceeded human

153 classifier performance on eating, drinking, climbing, and digging behaviors. Together, these results demonstrate

154 that our DINOv2+ model's performance on our test set approaches that of expert-level human classifiers. These

155 performance results confidently indicate a high level of reliability of our model, which would allow us to

156 perform behavior inference on a circadian timescale of days to weeks.

158   Finally, to assess the differences between supervised and self-supervised learning approaches in DEG

159 and DINOv2, we trained two additional linear probe heads on top of the frozen outputs of a pretrained DEG

160 model and the DINOv2 model. First, we trained a linear probe to classify our nine mouse behaviors using a

161 training and test set comprising mouse behavior frames that were simply rotated 90 degrees from the original

orientation of each frame used to pretrain the DEG model (**Fig. 2d**). We found that rotation had a negligible

impact on the DINOv2 model's performance. However, surprisingly, our DEG model's performance dropped

nearly 20% after a single rotation, even though DEG uses rotation as part of its image augmentation process

(Bohnslav et al., 2021). Next, we trained a linear probe on a completely novel task in which both models must

count the number of mice in a given frame using a training and test set comprising video frames containing

zero, one, or two mice in their home cage with and without the presence of a running wheel (**Fig. 2e**).

Unsurprisingly, because DINOv2 is a foundational model that can be adapted to a wide range of classification

tasks, DINOv2 greatly outperformed DEG on this counting task. These results demonstrate the difference in

visual feature robustness between supervised and self-supervised learning schemes and strongly suggest that the

DINOv2 model can serve as a powerful pretrained backbone for a wide variety of classification tasks.

Behavior classification occurs in real time

Regardless of our DINOv2+ model's exceptional performance, the complexity of machine learning

models often translates into poor usage speeds and low throughput in practice. Using DINOv2+ as an

enhancement to (or replacement for) traditional sensor-based behavior analysis requires us to use it to infer

videos in real time. That is, a video clip of n second duration must be recorded, processed, and automatically

inferred by DINOv2+ before n seconds have elapsed and the next video segment is ready to be processed and

inferred. To match the high throughput of sensor-based analysis (e.g., many running wheels can be recorded in

parallel), we also need to be able to record, process, and automatically infer behaviors from videos recorded

from multiple mice simultaneously. To solve this problem, we developed a hardware and software pipeline that

allowed us to automatically and continuously record and infer behaviors in real time from up to 24 mice in

parallel (**Fig. 3a**). Our system comprises power over ethernet (PoE) IP cameras connected in parallel to Gigabit

switches. These switches stream video data that is binned into constant length time segments onto dedicated

machine learning computers for inference and network-attached storage devices for backup.

Before using our system, we needed to identify the video segment length (in minutes) such that video data from x cameras can be inferred within that temporal window. To do this, we first calculated the single camera inference times for several potential models including DEG, DINOv2+, and, for comparison, a skeletal pose estimation model without behavior classification (DLC) (**Fig. 3b**) (Mathis et al., 2018). We found that while all models were able to infer video data from a single camera within each of the temporal windows tested, DINOv2+ was significantly faster at video inference than either DEG or DLC.

Next, to apply real-time inference to multiple animals in parallel, we first needed to identify the maximum number of cameras that could infer behaviors simultaneously within a reasonable video segment length. We again used DEG, DINOv2+, and DLC to calculate behavior inference times for various combinations of time segment lengths (5 min, 10 min, or 30 min) and numbers of cameras used to simultaneously stream video segments (10 or 20; **Fig. 3c**). We found that all models were able to infer video data from 10 cameras simultaneously regardless of video segment length. However, when our DEG model was used to infer video data from 20 cameras simultaneously, inference time exceeded the length of the video segment regardless of video segment length. This indicated a failure of real-time inference. In addition, our DINOv2+ model was significantly faster at video inference than either DEG or DLC at all time segment length and camera number combinations tested. Based on these results (and our experimental setup in which our behavior cabinets can hold a maximum of 12 mouse cages each), we chose to proceed with using our DINOv2+ model to infer videos with a video segment length of 30 min on a system comprising two sets of 12 cameras networked to individual machine learning computers (**Fig. 3a**). To test the efficacy of our system, we recorded videos from 24 mice simultaneously over 48 h in a 12h:12h LD cycle (**Fig. 3e, Supplementary Video 1**). Our system was able to successfully process video clips, infer behaviors, and plot time series activity profiles for each behavior over the duration of the recordings, "filling in" over time similarly to how wheel-running activity profiles are plotted by commercial circadian activity monitoring software. Together, these results demonstrate that our model can be used to automatically and continuously classify behaviors from multiple animals for an essentially unlimited experimental duration.

Sex influences circadian rhythms in home cage behaviors

Next, we applied our DINOv2+ model and automatic inference system to a fundamental question in

circadian biology: how do sex and estrogen influence circadian rhythms in behavior? Subtle sex differences in

wheel-running activity rhythms have been previously identified (Lee et al., 2004; Kuljis et al., 2013, 2016;

Krizo and Mintz, 2014; Joye and Evans, 2022; Anderson et al., 2023). However, because of technological

constraints, whether (and how) males and females differ in other behavioral rhythms is unknown. To address

this problem, we continuously recorded videos, inferred behaviors, and generated actograms (a field-standard

method of plotting activity profiles over multiple days) from male (n = 24) and female (n = 27) mice over 5 d in

LD and over 5 to 9 d in DD (**Figs. 4a, b**). Female mice underwent estrous staging prior to beginning recording,

allowing us to sort them into groups adjusted such that their actograms were aligned by their first day of

proestrus. We used these actograms to determine key circadian properties of each behavioral rhythm including

phase, amplitude, and period.

To quantify phase, we measured the acrophase (peak time of activity) for each behavior on each day in

LD and in DD (**Fig. 4c; Supplementary Figs. 1a, 2a**). We averaged these acrophases in LD and in DD to more

readily compare phase across behaviors and groups. For male mice, we averaged acrophases across each day.

We divided female mice into two groups based on their estrous cycle. For "proestrus/estrus" (P/E) female mice,

which have relatively high levels of endogenous estrogen, we averaged acrophases over each of the projected

days of proestrus based on pre-recording estrous staging. For "metestrus/diestrus" (M/D) female mice, which

have relatively low levels of endogenous estrogen, we averaged acrophases over all other days of recording.

Our first goal was to determine if any specific behaviors peaked at distinct times from other behaviors

and, if so, whether this pattern was observed in both males and females. To do this, we compared phase markers

for all nine behaviors separately within male and female groups (**Supplementary Fig. 2a**). In LD, we found that

for all groups of mice, all behaviors except resting and grooming peaked around the same time in the middle of the night (ZT or zeitgeber time 18, where ZT 0 is defined as lights on). As expected for nocturnal animals, resting peaked around the middle of the day, ZT 6. Curiously, grooming behavior in LD in male and P/E (but not M/D) female mice peaked about 30 min and 1.5 h earlier, respectively, than other non-resting behaviors. In DD, for M/D and P/E female mice, all behaviors except resting peaked around the same time, approximately 15 to 30 min earlier than their peak time in LD, as expected for "free-running" nocturnal animals with a shortened period of activity in DD. Surprisingly, for male mice, digging and nesting behaviors were greatly phase delayed in DD. Compared to all other non-resting behaviors, digging and nesting peaked about 30 min and 1 h later, respectively.

Next, to determine if individual behaviors peaked at distinct times in male and female mice, we compared phase markers for each behavior separately across male and female groups. In LD, we found that most behaviors in P/E female mice were phase delayed: eating, drinking, climbing, exploring, and resting behaviors peaked between 30 min to 1 h later compared to these behaviors in M/D females (**Supplementary Fig. 1a**). We also found that in male mice, some behaviors (drinking, grooming, and resting) peaked at similar times to those behaviors in M/D females. However, intriguingly, all other behaviors in male mice (eating, rearing, climbing, exploring, digging, and nesting) peaked at times in between the times those behaviors peaked in M/D and P/E females. In DD, we again found that most behaviors in P/E female mice were phase delayed: drinking, climbing, exploring, digging, nesting, and resting peaked between 30 min to 1 h later compared to these behaviors in M/D females (**Fig. 4c**). Most behaviors in male mice (eating, drinking, rearing, climbing, grooming, exploring, and resting) peaked at similar times to those behaviors in M/D female mice. However, digging and nesting behaviors in male mice instead peaked at similar times to those behaviors in P/E female mice because digging and nesting were phase delayed compared to all other behaviors in male mice in DD. Together, these results demonstrate that behavior rhythms in male and female mice exhibit distinct phase profiles. Specifically, we found that estrous state fundamentally alters behavior phase in female mice and that, in DD, nesting and digging behaviors are significantly delayed in male mice.

We next measured the amplitudes of each behavior rhythm in male, M/D female, and P/E female mice

by fitting a cosine wave to their averaged activity profiles in both LD and DD (**Fig. 4d; Supplementary Figs.**

1b, 3a). We found that the amplitudes of all behavior rhythms in male and female mice were dampened in DD

compared to LD, consistent with prior reports describing how light cycle influences wheel-running activity

amplitude (Li et al., 2006; Pasquali et al., 2010). We also observed that the amplitudes for most behavior

rhythms (rearing, climbing, exploring, digging, nesting, and resting) were significantly greater in P/E mice

compared to those behaviors in male mice in DD, but not in LD. We found that the amplitudes of some

behavior rhythms (climbing, exploring, nesting) were also greater in M/D mice compared to male mice in DD.

Finally, we calculated the periods of each behavior rhythm in males and females across all days in both LD and

DD (**Supplementary Fig. 4a**). We found that, as expected for nocturnal rodents, the free-running periods in DD

for all behavior rhythms in both male and female mice were shorter than the entrained periods in LD (averaged

across all behaviors: males LD 24.02 ± 0.03 h; males DD 23.73 ± 0.04 h; females LD 24.04 ± 0.03 h; females

DD 23.82 ± 0.03 h). Surprisingly, sex had little effect on period. Digging in females exhibited a slightly

lengthened period in DD, but no other behaviors showed significant period differences. Together, these results

demonstrate that biological sex has a profound effect on the amplitude of most behavioral rhythms but has little

to no effect on periodicity.

Estrogen replacement phenocopies multiple behavior rhythm changes seen in proestrus female mice.

To determine whether these observed sex differences in circadian behavior could be explained by

differences in endogenous estrogen levels, we again continuously recorded videos, inferred behaviors, and

generated actograms from ovariectomized (OVX; n = 24) and ovariectomized, estradiol-supplemented (OVXE;

n = 22) female mice over 5 d in LD and over 5 d in DD (**Figs. 5a, b**) (Ström et al., 2012). OVX mice have

chronically low levels of estrogen similar to the levels found in male mice or during metestrus/diestrus in intact

females, and OVXE mice have chronically elevated levels of estrogen similar to the levels found during

proestrus in intact females. We used these actograms to again determine key circadian properties of each

rhythm including phase, amplitude, and period.

To quantify phase, we again measured the acrophase (peak time of activity) for each behavior on each

day in OVX and OVXE mice in LD and in DD (**Fig. 5c; Supplementary Figs. 1c, 2b**). We averaged these

acrophases in LD and in DD to more readily compare phase across behaviors and groups. First, to determine if

any specific behaviors peaked at distinct times from other behaviors in OVX and OVXE mice, we compared

phase markers for all nine behaviors separately within estrogen replacement groups (**Supplementary Fig. 2b**).

In LD, we found that for both OVX and OVXE mice, all behaviors except resting peaked around the same time

in the middle of the night (ZT 18); resting peaked around the middle of the day, ZT 6. In DD, for both OVX and

OVXE mice, most non-resting behaviors peaked around the same time, about 30 min earlier than their peak

time in LD as expected for nocturnal rodents. However, grooming in OVX mice peaked about 30 min later, and

nesting and resting in OVXE mice peaked about 30 min and 1 h later, respectively, compared to all other non-

resting behaviors.

Next, to determine if individual behaviors peaked at distinct times in OVX and OVXE mice, we

compared phase markers for each behavior separately across estrogen replacement groups. In LD, we found no

difference between OVX and OVXE mice in the peak times of any behavior (**Supplementary Fig. 1c**). In DD,

similar to what we observed with intact P/E female mice, most behaviors (rearing, climbing, exploring, digging,

nesting, and resting) in OVXE mice were phase delayed, peaking between 30 min and 1 h later compared to

these behaviors in OVX mice (**Fig. 5c**). However, surprisingly, eating, drinking, and grooming behaviors in

OVXE mice peaked at the same time as those behaviors in OVX mice. Together, these results demonstrate that

behavior rhythms in OVX and OVXE mice exhibit distinct phase profiles. Specifically, we found that estrogen

replacement significantly phase delays most, but, importantly, not all behaviors in DD, but not in LD.

We next measured the amplitudes of each behavior rhythm in OVX and OVXE mice by fitting a cosine

wave to their averaged activity profiles in both LD and DD (**Fig. 5d; Supplementary Figs. 1d, 3b**). We found

that the amplitudes of all behavior rhythms except eating and nesting in OVX and OVXE mice were dampened

in DD compared to LD. We also observed that the amplitudes for some, but not all, behavior rhythms (drinking,

rearing, climbing, and exploring) were significantly greater in OVXE mice compared to those behaviors in

OVX mice in DD, but not in LD. Finally, we calculated the periods of each behavior rhythm in OVX and

OVXE mice across all days in both LD and DD (**Supplementary Fig. 4b**). We found that, as expected, the free-

running periods in DD for all behavior rhythms but nesting in both OVX and OVXE mice were shorter than the

entrained periods in LD (averaged across all behaviors: OVX LD 24.07 ± 0.02 h; OVX DD 23.70 ± 0.06 h;

OVXE LD 24.04 ± 0.04 h; OVXE DD 23.66 ± 0.06 h). Estrogen replacement had little effect on period: only

drinking and grooming behaviors in OVXE mice had slightly different periods (longer and shorter, respectively)

than in OVX mice. Together, these results demonstrate that estrogen replacement greatly influences both the

phase and amplitude of multiple behavior rhythms. Specifically, estrogen replacement increases behavior

rhythm amplitudes and mimics the phase delays we observed in intact female mice during proestrus.

A generalizable circadian behavioral analysis suite

Our DINOv2+ model and automatic inference system allowed us to thoroughly investigate how

circadian behaviors are influenced by sex and estrogen levels at an unprecedented throughput and acquisition

rate. We realized that introducing our software infrastructure to the broader scientific community could be

revolutionary to fields seeking to understand the temporal characteristics of animal behavior, including,

particularly, circadian biology. We therefore developed a "circadian behavioral analysis suite" (CBAS), a

Python package aimed at generalizing the software and DINOv2+ model used in our experiments (**Fig. 6a**).

CBAS is equipped to handle automated, continuous video acquisition, automated inference using the DINOv2

feature extractor and joint long short-term memory (LSTM) and linear layer models, and visualization of

behavior actograms in real time (**Fig. 6b**). Briefly, CBAS is divided into three modules: an acquisition module,

a classification and visualization module, and an optional training module. The acquisition module is capable of batch processing streaming video data from any number of network-configured real-time streaming protocol (RTSP) IP cameras. The classification and visualization module enables real-time inference on streaming video and displays acquired behavior time series data in real time as actograms that can be readily exported for offline analysis in a file format compatible with ClockLab Analysis, a widely-used circadian analysis software. Users wanting to fully replicate our recording setup (see the Jones Lab Github for a full parts list and assembly instructions) and nine behaviors of interest can immediately begin classification using our DINOv2+ joint LSTM and linear layer model that is included in the Python package. Importantly, because the DINOv2 visual backbone is kept static in our training model, users can also quickly and easily adapt CBAS to accommodate a diverse array of classification tasks, animal species, and video environments. The training module allows the user to create balanced training sets of behaviors of interest, train joint LSTM and linear layer model heads, and validate model performance on a naive test set of behavior instances. Importantly, CBAS's acquisition module is essentially machine learning model agnostic, allowing for future models to be easily incorporated into the CBAS pipeline. Together, these modules present an intuitive, accessible software interface that will allow for the rapid adoption of CBAS by end users with any level of programming ability.

Discussion

Here, we developed and validated a novel system for transforming existing machine learning classifiers into real-time sensors capable of phenotyping circadian rhythms in complex behaviors for an indefinite length of time. We used this pipeline to thoroughly characterize the effects of biological sex and estrogen levels on circadian behavior across 97 individual mouse recordings with a minimum duration of 10 d per recording at 10 frames per second. We then developed this toolkit into an open-source, user-friendly Python package – CBAS – for use by the broader circadian biology community and beyond. CBAS has the potential to reveal temporal variations in behavior that have previously gone undetected in a diverse range of animal models. In addition,

CBAS provides scientists with the tools needed to build, adequately validate, and automate highly reliable machine learning classifiers for any complex behavior(s) of interest.

CBAS's extensive model validation, classification power, and customizable, open-source nature set it apart from previous commercial (e.g., HomeCageScan, (Adamah-Biassi et al., 2013, 2014)) and non-commercial (e.g., (Steele et al., 2007; Goulding et al., 2008; Jhuang et al., 2010; Salem et al., 2015)) home cage behavior acquisition tools. For instance, (Jhuang et al., 2010) uses background masking and motion features to classify behaviors with a Hidden Markov Model Support Vector Machine, an outdated, but theoretically capable, architecture. While the authors point to a high classification accuracy, there is little to no information provided regarding more standard machine learning classification metrics such as precision, recall, F1 score, etc. With some assumptions, some of these values can be calculated from the provided confusion matrices, but for most behaviors they fail to meet our strict model performance threshold. Critically, the class balance of the training and test sets used for their model verification is unreported, and neither set remains available to the public. Similarly, (Goulding et al., 2008) fails to perform model performance validation for their supervised learning technique, although they do show that their system is capable of recording behaviors on a circadian timescale. However, because their model only identifies active and inactive behavior states, it is incapable of automatically identifying complex behaviors or animal-environment interactions. Adequate performance metrics, training sets, and testing sets of the HomeCageScan system are likewise scarce, a problem that is exacerbated by its closed-source nature. Importantly, none of these systems has user-friendly customizability to extend classification to other animal models, environments, and novel behaviors. Our goal with CBAS is to allow users with any level of programming ability to integrate completely customizable machine learning models, extensively validate model performance, and record behaviors in real time, indefinitely.

Although there have been several recent advances in supervised and even unsupervised pose-based behavior classification (e.g., SimBA, B-SOiD, A-SOiD, (Nilsson et al., 2020; Hsu and Yttri, 2021; Tillmann et al., 2024)), pose-based classifiers sacrifice critical learnable information with their sweeping dimensional

reduction to pose dynamics. For example, the subtle home cage environment differences that characterize digging versus nesting behavior in our recording setup would be completely lost in a reduction to pose time series. Furthermore, labeling pose data is significantly more difficult than labeling classification data, especially in dynamic video environments or with moving subjects. In contrast, DeepEthogram (DEG) models and our proposed DINOv2+ model have the capacity to learn specific high-dimensional features from spatial and temporal dynamics derived from raw pixel values (Bohnslav et al., 2021). Importantly, DEG models and DINOv2 are fundamentally different in how they are trained. DEG, which we previously used to analyze circadian behavior in a proof-of-concept study (Wahba et al., 2022), uses a feature extractor that is trained in a supervised manner where the model receives direct classification feedback from labeled data throughout training. In contrast, the DINOv2 feature extractor is pretrained in a self-supervised manner where the model is encouraged to produce a rich, often clustered, visual feature space that can then be used as a frozen basis for subsequent small supervised classification models. The DINOv2 backbone model has several benefits because of its self-supervised learning strategy. Most notably, this strategy generalizes the model's feature space to data and tasks which it has never been trained to recognize or accomplish. For example, DINOv2 is broadly capable of semantic segmentation, depth estimation, image/video classification, or object tracking/recognition with the minor addition of a trainable linear network layer (Oquab et al., 2023). Reducing bias is closely linked to enhancing feature generalizability. While supervised learning often optimizes for unstable visual heuristics, self-supervised training disregards these heuristics in favor of robust visual characteristics (Caron et al., 2021; Shwartz-Ziv and LeCun, 2023). Finally, self-supervised models help bridge the growing gap in access to computing resources and data science expertise needed to fully train and optimize high-performing vision models.

In addition to using our DINOv2+ model for inference, CBAS also leverages our standardized video recording pipeline to allow for real-time behavior classification that can rival the throughput of sensor-based recording systems (Siepka and Takahashi, 2005; Verwey et al., 2013). CBAS is also incredibly cost effective. The total cost of the hardware we used in this study, including our custom-built mouse cages and circadian

behavior cabinets, is only two-thirds the estimated cost of standard commercial systems. Moreover, our setup

can be used to record and infer any number of behaviors in parallel, whereas standard circadian acquisition

hardware is only capable of recording locomotor behavior by measuring wheel-running activity or infrared

beam breaks. Another major advantage of CBAS is its accessibility, presenting a low barrier to entry for the

broader circadian research community, including those with limited programming skills. Additionally, CBAS

mirrors the functionality of field-standard circadian analysis systems by plotting behavior data as actograms in

real time, providing immediate feedback about the state of ongoing experiments. CBAS also outputs these

behavior actograms in formats compatible with Actimetrics' ClockLab Analysis software, which ensures that

researchers can adapt familiar analyses to CBAS-generated behavior data. The open-source nature of CBAS

essentially democratizes the circadian analysis of complex behaviors, allowing a greater number of researchers

to investigate long-term behavior dynamics.

Previous studies have identified numerous sex differences in the temporal patterning of physiology. For

example, male and female mice exhibit distinct circadian rhythms in glucocorticoid production, cardiovascular

function, body temperature, and immune function (Griffin and Whitacre, 1991; Atkinson and Waddell, 1997;

Sanchez-Alavez et al., 2011; Barsha et al., 2016; Walton et al., 2022). However, the question of whether there

are pronounced sex differences in the temporal patterning of behavior has, to date, been mostly unanswered.

Subtle sex differences have been observed in wheel-running activity rhythms (Lee et al., 2004; Kuljis et al.,

2016; Anderson et al., 2023). For instance, male mice show a greater precision of wheel-running activity onsets

in LD and female mice show a longer wheel-running activity duration on the day of proestrus in DD (Albers et

al., 1981; Kuljis et al., 2013). However, given the presumed global regulation by the circadian system of

multiple brain circuits that control distinct behaviors, more work is needed to reveal and understand sex

differences in other behavioral rhythms (Starnes and Jones, 2023). We used CBAS to address this by testing the

hypothesis that circadian rhythms in behavior differ between males and females. We identified differences in

behavioral rhythms including, notably, that nesting and digging rhythms exhibit a distinct phase delay in male,

but not female, mice during constant darkness that had not been previously reported. We also observed that

most behavioral rhythms in female mice had higher peak-to-trough amplitudes, which is suggestive of more

robust circadian organization. This is consistent with previous work showing that the amplitude of wheel-

running activity rhythms is greater in female mice compared to male mice (Anderson et al., 2023). Previous

studies have also identified that the duration of wheel-running activity is extended (that is, it ends later) on the

day of proestrus (Albers et al., 1981). We confirm this finding and extend it to demonstrate that the temporal

organization of nearly all behaviors changes across the estrous cycle. Our findings reveal critical unseen sex

differences in many, but, importantly, not all behavioral rhythms, which emphasizes the importance of

measuring circadian rhythms in behaviors other than locomotor activity.

The limited number of previously identified sex differences in circadian behavior have been speculated

to be due to differences in levels of circulating sex hormones and/or sex hormone receptor expression (Walton

et al., 2022). Our experiments were designed to allow us to distinguish between differences in behavioral

rhythms that are due to biological sex and those that are due to the presence or absence of estrogen. We found

that estrogen replacement recapitulates most, but not all, of our observed sex differences in behavioral rhythms.

For instance, we found that P/E females exhibit higher amplitude circadian rhythms in most behaviors

compared to males and M/D females. Similarly, many behavioral rhythms in OVXE females are more robust

than in OVX females. We also found that the peak time of most behaviors in "high estrogen" P/E and OVXE

females was delayed compared to "low estrogen" M/D and OVX females. One possible explanation for this is

the relative distribution of estrogen receptors in brain circuits that regulate different behaviors. Exogenous

estrogen has been shown to increase the amplitude of wheel-running activity rhythms through the activation of

estrogen receptor (ER)α but to delay the phase of wheel-running activity rhythms through the activation of ERβ

(Royston et al., 2014). Indeed, a recent study determined that the lateral hypothalamus (LH), which has

subpopulations of both ERα-positive and ERβ-positive neurons, regulates nest-building behavior

(Merchenthaler et al., 2004; Sotelo et al., 2022). If these ERβ-expressing LH neurons are preferentially

activated during nesting behavior, this could explain why estradiol delays nesting rhythms in OVXE and P/E

mice. However, this does not explain why male mice, which have low levels of endogenous estrogen, exhibit

delayed digging and nesting rhythms that peak at similar times to those rhythms in OVXE and P/E mice, which

have high levels of endogenous estrogen. Further studies will need to determine whether this finding is due to

sex differences in developmental circuit wiring, differences in estrogen receptor distribution and/or expression

levels, or other factors.

In this study, we used our circadian behavioral analysis suite (CBAS) to automatically quantify

differences in the circadian regulation of behavior between male and female, and OVX and OVXE female,

mice. This approach can be readily expanded to address other critical questions in circadian biology,

neuroscience, and ecology, including the ethological investigation of other behavioral rhythms in videos of mice

recorded in the laboratory and, potentially, in the wild. Notably, CBAS can also be used for the rapid circadian

phenotyping of mice with different genotypes or disorders (Richardson, 2015). Current approaches almost

universally measure changes to wheel-running activity rhythms as evidence that a mutation, gene, or drug

influences circadian behavior. Here, we found that some, but, critically, not all, behavioral rhythms differ by

biological sex and by estrogen levels. It is therefore highly likely that any given experimental treatment could

cause circadian alterations in behaviors other than, or in addition to, wheel-running activity. CBAS aims to

extend the modern toolkit of machine learning classification into any and all long-term behavior assays, greatly

expanding the scope of potential hypotheses and impact of future studies.

Materials and methods

<u>Animals</u>

Prior to recording, we group-housed male (n = 24) and female (n = 51; 24 of which were subsequently

ovariectomized, see next section) wild-type mice in their home cages in a 12h:12h light:dark cycle (LD, where

lights on is defined as zeitgeber time (ZT) 0; light intensity ~2 x 10^{14} photons/cm^2/s) at constant temperature

(~23°C) and humidity (~40%) with food and water provided ad libitum. All mice were between 6 and 12 weeks

old at the time of the recording. To determine the estrous stage of female mice, we performed vaginal lavage for

four consecutive days prior to beginning long-term recording (Byers et al., 2012). All experiments were

approved by and performed in accordance with the guidelines of Texas A&M University's Institutional Animal

Care and Use Committee.

Ovariectomy and estradiol capsule implantation

We ovariectomized a cohort of female mice (OVX, n = 24) using standard methods (Ström et al., 2012).

Briefly, we made a sterile ~2 cm bilateral incision through the skin and peritoneum immediately dorsal to the

ovaries. After ligating and removing each ovary, we sutured the peritoneum and skin incisions. We provided the

mice with buprenorphine-SR (1 mg/kg; subcutaneous) and enrofloxacin (0.25 mg/ml; ad libitum in their

drinking water) and allowed them to recover in their home cages for at least 1 week prior to beginning long-

term recording. After recording OVX mice, we implanted them with a sterile 2 cm silastic capsule containing

17-β-estradiol (36 μg/ml in sesame oil) subcutaneously between the shoulder blades (OVXE; n = 22) (Ström et

al., 2012). We excluded two OVX mice from capsule implantation because they had excessive barbering around

their ovariectomy incision site. We allowed OVXE mice to recover in their recording cages for 1 d prior to

beginning long-term recording.

Experimental housing

We transferred individual mice from their home cages to custom-built recording cages inside custom-

built light-tight, temperature-and humidity controlled circadian cabinets for the duration of our experiments. We

built the cages (external dimensions, length x width x height: 22.9 cm x 20.3 cm x 21.6 cm; internal

dimensions: 20.3 cm x 17.8 cm x 20.3 cm) out of transparent and opaque acrylic panels (thickness, walls and

floor: 6.4 mm; lid, 3.2 mm) and T-slot aluminum extrusions (25.4 mm^2) (**Fig. 1a**). We 3D printed custom water

bottle holders and food hoppers out of PLA filament, coated them in food safe clear-cast epoxy resin

(Alumilite), and affixed them to the acrylic walls. To continue our recordings throughout the dark phase, and to

prevent potential glare and shadows from ceiling-mounted lights, we affixed dim infrared (850 nm) light strips

to the cage lid using a custom 3D printed cage topper. Prior to recording, we added ~7 mm wood chip bedding

and a 25 mm by 50 mm square of cotton nestlet to the cage bottom, added food pellets to the food hopper, and

attached a standard water bottle filled with water to the water bottle holder such that its metal spout protruded

about 1.5 cm into the cage. Our circadian cabinets were built to hold twelve of our custom-built mouse cages

across three vertical shelves, with four cages per shelf. We controlled the ceiling-mounted lights in the cabinets

(broad-spectrum white light, ~6 x 10^{13} photons/cm^2/s measured at the cage floor) using ClockLab Data

Collection hardware and software (Actimetrics) that communicated via a 5V transistor-transistor logic signal

with a high-power power relay (Digital Loggers). We performed daily animal welfare checks using dim red

light (650 nm).

<u>Automated video recording</u>

We positioned power-over-internet (PoE) IP cameras without infrared filters (I706-POE, Revotech)

equipped with 6 mm lenses (Xenocam) 47.5 cm above the recording cages such that all four corners of the cage,

the food hopper, and the water spout were each visible in the recorded video and the mouse and nesting material

were in focus. We recorded all videos at 10 frames per second (fps) with in-camera image settings set to a

contrast of 130/255, brightness of 140/255, saturation of 0/255, and sharpness of 128/255. We streamed videos

at a main stream bitrate of 2048 kilobits per second (kb/s) and a secondary stream bitrate of 256 kb/s. We

disabled audio streams to reduce bandwidth. We recorded our mice in cohorts of 8 to 12 mice split between two

circadian cabinets. We paused our recordings briefly between light settings (LD and DD) to allow for cage

changes, if necessary.

We used FFmpeg, a standard open-source video processing tool, to develop a custom video acquisition

system capable of streaming live video and storing successively binned segments of video for each network

camera simultaneously. During a recording, FFmpeg automatically handles cropping the video to a region of

interest and scaling the video to a desired size (here to a scale of 256x256 pixels). To do this, we connected our

cameras in parallel via 10 gigabits per second (Gbps) Cat6 ethernet cables to a Gigabit PoE switch (Aruba

JL684A#ABB). We then connected our switches via 40 Gbps Cat8 ethernet cables to our custom machine-

learning computers (12-core AMD Ryzen 9 5900X CPU, 32 GB RAM, NVIDIA GeForce RTX 3090 with 24

GB VRAM) (**Fig. 3a**). During a recording, our software creates two threads to monitor the creation of streamed

video segments and orchestrate the inference of incoming data (the "storage" and "inference" threads). The

storage thread records information about each video segment (creation time, segment length, camera-specific

settings), moves the video segments to the corresponding camera directories on the computers, and notifies the

inference thread that new videos are available for inference (see below).

<u>Behavior definitions</u>

We defined a list of nine home cage behaviors (eating, drinking, rearing, climbing, grooming, exploring,

digging, nesting, and resting, (Garner, 2017)) with the goal of identifying the visual and motion characteristics

of each behavior that our DINOv2+ model would be capable of learning (**Supplementary Table 1**). As such,

our definitions do not aim to ascribe intent to a behavior (as humans are often inclined to do), but rather contain

references to particular features that strictly define behavioral classes. These include spatial features that are

necessary constraints on a behavior and temporal features that are split into two groups indicative of the start

and stop of a behavior sequence. To further enforce these rigorous criteria defining behaviors, our entire set of

training instances were generated by a single labeler.

<u>Model training and inference</u>

To train our baseline DeepEthogram (DEG) classifier, we needed to individually train three components,

a "flow generator" that estimates optic flow across video frames, a "feature extractor" that determines the

probability of a behavior being present on a given frame based on a low-dimensional set of temporal and spatial

features, and a "sequence model" that further refines model predictions using a temporal gaussian mixture

(TGM) model with a larger temporal receptive field. We trained our flow generator on a set of videos consisting

of approximately 500,000 frames of videos from 8 mice recorded at 10 fps. We then trained our feature

extractor using the medium model size preset (deg_m, (Bohnslav et al., 2021)) and our TGM sequence model

using a temporal window of 15 frames. We trained both the feature extractor and TGM sequence models on an

identical balanced training set used for subsequent training of our DINOv2+ model (see below). Importantly,

we include our model configuration files for all DEG models, our DINOv2+ model, and all trained model

weights at the Jones lab Google Drive repository (see Data Availability section below).

Our DINOv2+ model architecture was designed to take as input sequenced outputs from the DINOv2

feature extractor and produce a robust, frame-to-frame stable, and accurate classification time series (**Fig. 2a**).

The DINOv2 feature extractor model outputs one 768 length vector for each given video frame encoding the

relevant visual information about the image scene. Our joint long short-term memory (LSTM) and linear layer

classification head integrates visual and motion information from a sequence of vectors (here 31 frames)

centered at the frame of interest into a behavioral classification. During a forward pass of our classification

head, noise is randomly injected into a normalized version of the sequence of DINOv2 outputs, transformed

through a single linear layer into the output size, and then averaged over an 11 frame, centered sub-window.

Simultaneously, the mean of the original input sequence is subtracted from the input sequence, compressed by a

linear layer to a latent dimension, and then passed through a single layer bidirectional LSTM network. The

logits of the LSTM layer are condensed to the output size and added to the outputs of the linear layer. A

softmax of the summed output results in the model's behavior classification confidence for each frame. The

softmax function is defined as

585
$$s(y_{oj}) = \frac{e^{y_{oj}}}{\sum_{k=1}^{n} e^{y_{ok}}}$$

586 where n is vector length (here, 9), y_{oj} is the output vector at position j, and y_{ij} is the input vector at position j.

587 We next trained DINOv2+ classification head on a balanced set of behavior instances sampled across

588 the light and dark phases from 30 unique mice and cages. For this task, we trained our model using a cross

589 entropy loss function, defined as:

590
$$L_{CE} = -\sum_{j=1}^{n} y_{ij} log(s(y_{oj}))$$

where, again, n is vector length (here, 9), y_{oj} is the output vector at position j, and y_{ij} is the input vector at

position j (Ciampiconi et al., 2023). Additionally, we added a covariance loss to discourage covariance of our

LSTM output features. Our covariance loss was defined as the off-diagonal sum of the absolute covariance

matrix constructed using the raw latent dimensional outputs of the LSTM layer divided by our latent dimension

size. This approach was inspired by the elegant loss function employed in the VICReg learning scheme, and it

consistently improved our classification performance (Bardes et al. 2021). We identified optimal

hyperparameters that minimized the total loss to be a latent dimension of 256, an LSTM latent dimension of 64,

and a linearly decreased learning rate of 5e-4 to 1e-5 over 10 epochs of training. During classification training,

model states are selectively saved by maximizing for the weighted average F1 score of model performance on a

test set.

Model validation

To validate the performance of our behavior classifier, we used a naive, balanced test set of behavior

sequences. Prior to model training, we randomly selected each unique behavior sequence (or "instance") from

our annotated dataset while preserving class balance. Importantly, to prevent misleading or skewed model

performance results, we did not use the instances in this test set during any form of model training or

adjustment.

From this balanced test set, we randomly sampled 1,000 sequences with a maximum length of 31

frames. After we used our model to infer all sampled sequences, we calculated precision, recall, F1 score,

specificity, and balanced accuracy using the *sklearn.metrics* library in Python. We also calculated the

normalized Matthews correlation coefficient (nMCC) using a custom Python implementation (Chicco and

Jurman, 2020). We repeated this random sampling for a total of ten iterations before calculating the mean and

standard deviations of each metric.

To cross validate our DINOv2+ model with the DEG TGM sequence model and human annotators, we

first trained a TGM sequence model with a temporal window of 15 frames on the equivalent training set to that

of our DINOv2+ model. We then repeated the sampling and metric calculation detailed above to determine

means and standard deviations for the TGM model metrics. Using both models' output prediction probabilities,

we calculated the precision-recall curves for each classifier. To determine if the change between the area under

the precision-recall curves (AUPRC) was significant, we used a Python version of a bootstrapping method

originally implemented in R to create a normal distribution of area differences between random subsamples of

the two curves with 10,000 sampling iterations (Zobolas, n.d.). We then compared the area difference of the two

total precision-recall curves to the mean and standard deviation of this distribution to determine significance.

Finally, we designed a custom GUI in Python that allows human annotators to classify 10,000 randomly

sampled 15 frame sequences of video frames from our test set by replaying the sequence until it is classified as

a behavior. Importantly, the GUI does not give the human annotator performance feedback over the course of

the annotation so (much like our machine learning models) they are unable to learn as they annotate. Using

these annotations, we repeated the sampling and metric calculations to determine means and standard deviations

for human labeler metrics.

Automated inference

At the beginning of a recording, our software automatically bins the video stream into segments of time.

In these experiments, we chose to record in thirty minute intervals. For each new video bin, a subprocess infers

the video using the frozen DINOv2 feature extractor model. In this manner, CBAS continuously automates

model inference until the recording is terminated by the user. Users can also add pre-recorded videos to the

project directory to begin the DINOv2 inference of these videos. If a joint LSTM and linear layer model is

trained and ready for use in inference (as in our experiments), CBAS also coordinates the automated inference

of the DINOv2 features into sequenced behavior classes.

Analysis

We produced behavior actograms by binning the number of frames predicted as a given behavior over a

30 min period. To account for differing estrous states in female mice, we shifted their behavior actograms such

that their projected day of proestrus (as determined by estrous scoring) was aligned for each mouse that we

recorded. In LD, adjustments and group ns were: 0 d (n = 8 mice), -1 d (n = 5 mice) , -2 d (n = 7 mice), and -3 d

(n = 7 mice). In DD, adjustments and group ns were: 0 d (n = 5 mice), -1 d (n = 4 mice), -2 d (n = 6 mice), -3 d

(n = 13 mice).

To calculate circadian parameters (phase, period, and amplitude), we used CBAS to export each

actogram as an .awd file, a file format compatible with ClockLab Analysis (Actimetrics), a widely-used

circadian analysis software. To calculate phase, we identified acrophases by calculating the midpoint between

onset and offset times determined by a standard template matching algorithm that searches for a 12 h period of

inactivity (or activity) followed by a 12 h period of activity (or inactivity). For analysis, acrophases for male,

OVX, and OVXE mice were averaged across each day in LD and DD. Acrophases for proestrus/estrus (P/E)

female mice were averaged on the projected days of proestrus: days 1 and 5 in LD and days 1, 5, and 9 in DD.

Acrophases for metestrus/diestrus (M/D) female mice were averaged on all other days (days 2-4 in LD and days

2-4 and 6-8 in DD). To calculate period, we used a Lomb-Scargle periodogram with a range of 20 to 28 hours

and a significance level of 0.001. To calculate amplitude, we measured the peak-to-peak amplitude of a sine

wave fitted to the average activity profile calculated across all days in LD or all days in DD.

We performed the following statistical tests in Prism 10.0 (Graphpad): one-way ANOVA, unpaired t-

test, two-way ANOVA, Tukey's multiple comparisons test, Dunnett's multiple comparisons test. We performed

a bootstrapping test in Python (Zobolas, n.d.). Because no phase markers occurred at or near the 24 h modulus,

we performed statistical comparisons without using circular statistics. We used Shapiro-Wilk and Brown-

Forsythe tests to test for normality and equal variance, defined α as 0.05, and presented all data as mean ± SEM.

Data availability

All data generated in this study that support our findings are presented within this paper or its Supplementary

Materials or at the Jones lab Google Drive repository at http://tinyurl.com/jones-lab-tamu. CBAS is also

available to the public at the Jones lab Github page at https://github.com/jones-lab-tamu.

Acknowledgments

We thank the members of the Jones lab for discussion and comments on the manuscript and V. Fisher for assistance with ovariectomies. This work was supported by National Institutes of Health Grant R35GM151020 (J.R.J.) and a Research Grant from the Whitehall Foundation (J.R.J.).

Figures

Figure 1. Recording and classification standardization of nine home cage behaviors. a) Schematic of the

home cage recording setup. **b)** Representative examples of individual frames depicting each of nine behaviors

(eating, orange; drinking, yellow; rearing, blue; climbing, red; grooming, green; exploring, brown; digging,

magenta; nesting, purple; resting, gray). First frame depicts the behavior occurring in the full field of view,

subsequent frames are zoomed in to better illustrate behaviors. **c)** Bout length (duration of a behavioral

instance) for each behavior within a maximum window size of 360 s. n $\geq$ 38 bouts from 29 to 30 mice per

behavior. Box and whiskers depict median and interquartile range. **d)** Number of unique instances of each

behavior in the 8.1 h human-labeled dataset broken down by training and test sets.

Figure 2. DINOv2+ approaches expert-level performance on behavior classification. a) Schematic of performance and generalization tests. Features from a frozen pretrained DeepEthogram (DEG) model and a frozen pretrained DINOv2 model were used to evaluate the ability of each visual feature extractor to successfully classify mouse behavior using our DINOv2+ joint LSTM and linear layer model head (performance; **Figs. 2b,c**), classify mouse behavior on behavior frames rotated 90° using a single layer linear network head (generalization; **Fig. 2d**), and count the number of mice in a cage using a single layer linear network head (generalization; **Fig. 2e**). **b)** Precision-recall curves for each behavior calculated for the DINOv2+ (colored lines) and DEG (dashed lines) models by varying the decision threshold of each binary classifier. Shading depicts the area under the precision-recall curve (AUPRC) for each behavior for each model. Bootstrap test; **, $p < 0.01$; ***, $p < 0.001$. **c)** Performance metrics for each behavior calculated for a trained human classifier (green), the DEG model (blue), and the DINOv2+ model (red). n = 10 sets of 1,000 randomly sampled

test set frames per behavior. Dashed line depicts a predefined performance threshold of 0.80. Lines and error

bars depict mean ± SEM. F1, F1 score; nMCC, normalized Matthews correlation coefficient. **d)** Relative

performance for the DEG (blue) and DINOv2 (red) pretrained models when tested on a rotated version of a

baseline behavior sequence test set using a single layer linear network head on top of the baseline models. **e)** F1

score calculated for both DEG and DINOv2 on a classification task involving counting the number of mice in a

cage using a single layer linear network head on top of the baseline models.

Figure 3. DINOv2+ allows for real-time behavior classification. a) Schematic of the real-time video

recording, processing, and inferring system comprising two sets of 12 PoE (power over ethernet) IP cameras

networked to a switch that passes streaming video data to a machine learning computer for video inference and

a network-attached storage device for video backup. **b)** Single-video inference times for video segments of

various lengths calculated for a skeletal pose estimation model without behavior classification (green, DLC),

DEG (blue), and DINOv2+ (red). n = 3 replicates per model. Two-way ANOVA with post-hoc Tukey's

multiple comparison's test; *, $p < 0.05$; ***, $p < 0.001$. **c)** Inference times for combinations of video segment

length and number of cameras used to simultaneously stream video segments calculated for each model. Dashed

lines depict the times at which inference time equals the length of the video segment. Failure of real-time

inference for a particular combination of segment length, camera number, and inference model is represented by

a black X above the bar. **d)** Representative activity profiles for each behavior from an individual mouse

recorded in a 12 h:12 h light:dark (LD) cycle for 48 h. 30 min segments of continuously recorded video were

automatically processed, inferred, and plotted over the duration of the recording, "filling in" over time. For

visualization, plots shown here are only updated every 6 h. ZT, zeitgeber time.

Figure 4. Male and female mice exhibit distinct circadian rhythms in home cage behaviors. a,b)

Representative double-plotted actograms depicting behaviors (colored lines on each row) averaged across eight

male mice or eight female mice that started the experiment in the same estrous state recorded over 5 d in a

12h:12h light:dark (LD) cycle (gray and yellow shading) and 9 d in constant darkness (DD; gray and light gray

shading). **c)** Behavior phase comparison plots depicting the acrophases (peak times in circadian time, where CT

18 is subjective midnight and CT 6 is subjective noon) for male (teal, n = 24), metestrus/diestrus (M/D; pink),

and proestrus/estrus (P/E; purple) female (n = 27) mice recorded in DD. Lines and error bars depict mean ±

SEM. Asterisks indicate behaviors with significant differences in acrophase across groups. One-way ANOVA

with post-hoc Tukey's multiple comparisons test; *, $p < 0.05$; **, $p < 0.01$. **d)** Normalized amplitude for each

behavior rhythm for male (teal), M/D (pink), and P/E (purple) female mice measured in DD. Two-way ANOVA

with post-hoc Tukey's multiple comparisons test; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure 5. Ovariectomized and ovariectomized, estradiol-supplemented female mice exhibit distinct circadian rhythms in home cage behaviors. a,b) Representative double-plotted actograms depicting behaviors (colored lines on each row) averaged across eight ovariectomized (OVX) female mice or eight ovariectomized, estradiol-supplemented (OVXE) female mice recorded over 5 d in a 12h:12h light:dark (LD) cycle (gray and yellow shading) and 5 d in constant darkness (DD; gray and light gray shading). **c)** Behavior phase comparison plots depicting the acrophases (peak times in circadian time, where CT 18 is subjective midnight and CT 6 is subjective noon) for OVX (pink, n = 24) and OVXE (purple; n = 22) female mice recorded in DD. Lines and error bars depict mean ± SEM. Asterisks indicate behaviors with significant differences in acrophase across groups. One-way ANOVA with post-hoc Tukey's multiple comparisons test; *, $p < 0.05$; **, $p < 0.01$. **d)** Normalized amplitude for each behavior rhythm for OVX (pink) and OVXE (purple) female mice measured in DD. Two-way ANOVA with post-hoc Tukey's multiple comparisons test; **, $p < 0.01$.

Figure 6. CBAS: a circadian behavioral analysis suite. a) CBAS is a user-friendly GUI-enabled Python

package that allows for the automated acquisition, classification, and visualization of behaviors over time. **b)**

Schematic of the CBAS pipeline. Red; acquisition module; blue, training module; green, classification and

visualization classification module.

References

References

Adamah-Biassi EB, Hudson RL, Dubocovich ML (2014) Genetic deletion of MT1 melatonin receptors alters
spontaneous behavioral rhythms in male and female C57BL/6 mice. Horm Behav 66:619–627.

Adamah-Biassi EB, Stepien I, Hudson RL, Dubocovich ML (2013) Automated video analysis system reveals
distinct diurnal behaviors in C57BL/6 and C3H/HeN mice. Behav Brain Res 243:306–312.

Albers HE, Gerall AA, Axelson JF (1981) Effect of reproductive state on circadian periodicity in the rat.
Physiol Behav 26:21–25.

Anderson ST, Meng H, Brooks TG, Tang SY, Lordan R, Sengupta A, Nayak S, Mřela A, Sarantopoulou D,
Lahens NF, Weljie A, Grant GR, Bushman FD, FitzGerald GA (2023) Sexual dimorphism in the response
to chronic circadian misalignment on a high-fat diet. Sci Transl Med 15:eabo2022.

Atkinson HC, Waddell BJ (1997) Circadian variation in basal plasma corticosterone and adrenocorticotropin in
the rat: sexual dimorphism and changes across the estrous cycle. Endocrinology 138:3842–3848.

Barsha G, Denton KM, Mirabito Colafella KM (2016) Sex- and age-related differences in arterial pressure and
albuminuria in mice. Biol Sex Differ 7:57.

Bohnslav JP, Wimalasena NK, Clausing KJ, Dai YY, Yarmolinsky DA, Cruz T, Kashlan AD, Chiappe ME,
Orefice LL, Woolf CJ, Harvey CD (2021) DeepEthogram, a machine learning pipeline for supervised
behavior classification from raw pixels. Elife 10 Available at: http://dx.doi.org/10.7554/eLife.63377.

Brzezinski D, Stefanowski J, Susmaga R, Szczech I (2020) On the Dynamics of Classification Measures for
Imbalanced and Streaming Data. IEEE Trans Neural Netw Learn Syst 31:2868–2878.

Byers SL, Wiles MV, Dunn SL, Taft RA (2012) Mouse estrous cycle identification tool and images. PLoS One
7:e35538.

Caron M, Touvron H, Misra I, Jégou H, Mairal J, Bojanowski P, Joulin A (2021) Emerging Properties in Self-
Supervised Vision Transformers. arXiv [csCV] Available at: http://arxiv.org/abs/2104.14294.

Chicco D, Jurman G (2020) The advantages of the Matthews correlation coefficient (MCC) over F1 score and
accuracy in binary classification evaluation. BMC Genomics 21:6.

Ciampiconi L, Elwood A, Leonardi M, Mohamed A, Rozza A (2023) A survey and taxonomy of loss functions
in machine learning. arXiv [csLG] Available at: http://arxiv.org/abs/2301.05579.

Fujita M, Hagino Y, Takeda T, Kasai S, Tanaka M, Takamatsu Y, Kobayashi K, Ikeda K (2017) Light/dark
phase-dependent spontaneous activity is maintained in dopamine-deficient mice. Mol Brain 10:49.

Garner J (2017) Mouse Ethogram. Available at: https://mousebehavior.org/ethogram/ [Accessed August 17,
2022].

Gaskill BN, Rohr SA, Pajor EA, Lucas JR, Garner JP (2009) Some like it hot: Mouse temperature preferences
in laboratory housing. Appl Anim Behav Sci 116:279–285.

Goulding EH, Schenk AK, Juneja P, MacKay AW, Wade JM, Tecott LH (2008) A robust automated system
elucidates mouse home cage behavioral structure. Proc Natl Acad Sci U S A 105:20575–20582.

Grandini M, Bagli E, Visani G (2020) Metrics for Multi-Class Classification: an Overview. arXiv [statML]

Available at: http://arxiv.org/abs/2008.05756.

Grieco F et al. (2021) Measuring Behavior in the Home Cage: Study Design, Applications, Challenges, and
Perspectives. Front Behav Neurosci 15:735387.

Griffin AC, Whitacre CC (1991) Sex and strain differences in the circadian rhythm fluctuation of endocrine and
immune function in the rat: implications for rodent models of autoimmune disease. J Neuroimmunol
35:53–64.

Hochreiter S, Schmidhuber J (1997) Long Short-Term Memory. Neural Comput 9:1735–1780.

Hsu AI, Yttri EA (2021) B-SOiD, an open-source unsupervised algorithm for identification and fast prediction
of behaviors. Nat Commun 12:5188.

Jhuang H, Garrote E, Mutch J, Yu X, Khilnani V, Poggio T, Steele AD, Serre T (2010) Automated home-cage
behavioural phenotyping of mice. Nat Commun 1:68.

Joye DAM, Evans JA (2022) Sex differences in daily timekeeping and circadian clock circuits. Semin Cell Dev
Biol 126:45–55.

Jud C, Schmutz I, Hampp G, Oster H, Albrecht U (2005) A guideline for analyzing circadian wheel-running
behavior in rodents under different lighting conditions. Biol Proced Online 7:101–116.

Kahnau P et al. (2023) A systematic review of the development and application of home cage monitoring in
laboratory mice and rats. BMC Biol 21:256.

Krizo JA, Mintz EM (2014) Sex differences in behavioral circadian rhythms in laboratory rodents. Front
Endocrinol 5:234.

Kuljis DA, Gad L, Loh DH, MacDowell Kaswan Z, Hitchcock ON, Ghiani CA, Colwell CS (2016) Sex
Differences in Circadian Dysfunction in the BACHD Mouse Model of Huntington's Disease. PLoS One
11:e0147583.

Kuljis DA, Loh DH, Truong D, Vosko AM, Ong ML, McClusky R, Arnold AP, Colwell CS (2013) Gonadal-
and sex-chromosome-dependent sex differences in the circadian system. Endocrinology 154:1501–1512.

Lee TM, Hummer DL, Jechura TJ, Mahoney MM (2004) Pubertal development of sex differences in circadian
function: an animal model. Ann N Y Acad Sci 1021:262–275.

Lever J, Krzywinski M, Altman N (2016) Model selection and overfitting. Nat Methods 13:703–704.

Li J-D, Hu W-P, Boehmer L, Cheng MY, Lee AG, Jilek A, Siegel JM, Zhou Q-Y (2006) Attenuated circadian
rhythms in mice lacking the prokineticin 2 gene. J Neurosci 26:11615–11623.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018) DeepLabCut:
markerless pose estimation of user-defined body parts with deep learning. Nat Neurosci 21:1281–1289.

Merchenthaler I, Lane MV, Numan S, Dellovade TL (2004) Distribution of estrogen receptor alpha and beta in
the mouse central nervous system: in vivo autoradiographic and immunocytochemical analyses. J Comp
Neurol 473:270–291.

Metzger J, Wicht H, Korf H-W, Pfeffer M (2020) Seasonal Variations of Locomotor Activity Rhythms in
Melatonin-Proficient and -Deficient Mice under Seminatural Outdoor Conditions. J Biol Rhythms 35:58–
71.

Muller M, Wolf CT, Andres J, Desmond M, Joshi NN, Ashktorab Z, Sharma A, Brimijoin K, Pan Q, Duesterwald E, Dugan C (2021) Designing Ground Truth and the Social Life of Labels. In: Proceedings of the 2021 CHI Conference on Human Factors in Computing Systems, pp 1–16 CHI '21. New York, NY, USA: Association for Computing Machinery.

Nilsson SRO, Goodwin NL, Choong JJ, Hwang S, Wright HR, Norville ZC, Tong X, Lin D, Bentzley BS, Eshel N, McLaughlin RJ, Golden SA (2020) Simple Behavioral Analysis (SimBA) – an open source toolkit for computer classification of complex social behaviors in experimental animals. bioRxiv:2020.04.19.049452 Available at: https://www.biorxiv.org/content/10.1101/2020.04.19.049452v2 [Accessed February 22, 2024].

Oquab M et al. (2023) DINOv2: Learning Robust Visual Features without Supervision. arXiv [csCV] Available at: http://arxiv.org/abs/2304.07193.

Pasquali V, Capasso A, Renzi P (2010) Circadian and ultradian rhythms in locomotory activity of inbred strains of mice. Biol Rhythm Res 41:63–74.

Pendergast JS, Branecky KL, Yang W, Ellacott KLJ, Niswender KD, Yamazaki S (2013) High-fat diet acutely affects circadian organisation and eating behavior. Eur J Neurosci 37:1350–1356.

Pereira TD, Shaevitz JW, Murthy M (2020) Quantifying behavior to understand the brain. Nat Neurosci 23:1537–1549.

Peters SM, Pothuizen HHJ, Spruijt BM (2015) Ethological concepts enhance the translational value of animal models. Eur J Pharmacol 759:42–50.

Richardson CA (2015) The power of automated behavioural homecage technologies in characterizing disease progression in laboratory mice: A review. Appl Anim Behav Sci 163:19–27.

Robinson-Junker AL, O'hara BF, Gaskill BN (2018) Out Like a Light? The Effects of a Diurnal Husbandry Schedule on Mouse Sleep and Behavior. J Am Assoc Lab Anim Sci 57:124–133.

Royston SE, Yasui N, Kondilis AG, Lord SV, Katzenellenbogen JA, Mahoney MM (2014) ESR1 and ESR2 differentially regulate daily and circadian activity rhythms in female mice. Endocrinology 155:2613–2623.

Salem GH, Dennis JU, Krynitsky J, Garmendia-Cedillos M, Swaroop K, Malley JD, Pajevic S, Abuhatzira L, Bustin M, Gillet J-P, Gottesman MM, Mitchell JB, Pohida TJ (2015) SCORHE: a novel and practical approach to video monitoring of laboratory mice housed in vivarium cage racks. Behav Res Methods 47:235–250.

Sanchez-Alavez M, Alboni S, Conti B (2011) Sex- and age-specific differences in core body temperature of C57Bl/6 mice. Age 33:89–99.

Schwartz WJ, Zimmerman P (1990) Circadian timekeeping in BALB/c and C57BL/6 inbred mouse strains. J Neurosci 10:3685–3694.

Segalin C, Williams J, Karigo T, Hui M, Zelikowsky M, Sun JJ, Perona P, Anderson DJ, Kennedy A (2020) The Mouse Action Recognition System (MARS): a software pipeline for automated analysis of social behaviors in mice. Cold Spring Harbor Laboratory:2020.07.26.222299 Available at: https://www.biorxiv.org/content/10.1101/2020.07.26.222299v1 [Accessed December 7, 2020].

Shuboni-Mulligan DD, Young DL Jr, De La Cruz Minyety J, Vera E, Munasinghe J, Gall AJ, Gilbert MR, Armstrong TS, Smart DK (2021) Impact of age on the circadian visual system and the sleep-wake cycle in mus musculus. NPJ Aging Mech Dis 7:10.

Shwartz-Ziv R, LeCun Y (2023) To Compress or Not to Compress- Self-Supervised Learning and Information
Theory: A Review. arXiv [csLG] Available at: http://arxiv.org/abs/2304.09355.

Siepka SM, Takahashi JS (2005) Methods to Record Circadian Rhythm Wheel Running Activity in Mice. In:
Methods in Enzymology (Young MW, ed), pp 230–239. Academic Press.

Sotelo MI, Tyan J, Markunas C, Sulaman BA, Horwitz L, Lee H, Morrow JG, Rothschild G, Duan B, Eban-
Rothschild A (2022) Lateral hypothalamic neuronal ensembles regulate pre-sleep nest-building behavior.
Curr Biol Available at: http://dx.doi.org/10.1016/j.cub.2021.12.053.

Starnes AN, Jones JR (2023) Inputs and Outputs of the Mammalian Circadian Clock. Biology 12:508.

Steele AD, Jackson WS, King OD, Lindquist S (2007) The power of automated high-resolution behavior
analysis revealed by its application to mouse models of Huntington's and prion diseases. Proc Natl Acad
Sci U S A 104:1983–1988.

Ström JO, Theodorsson A, Ingberg E, Isaksson I-M, Theodorsson E (2012) Ovariectomy and 17β-estradiol
replacement in rats and mice: a visual demonstration. J Vis Exp:e4013.

Tendle A, Hasan MR (2021) A study of the generalizability of self-supervised representations. Machine
Learning with Applications 6:100124.

Tillmann JF, Hsu AI, Schwarz MK, Yttri EA (2024) A-SOiD, an active-learning platform for expert-guided,
data-efficient discovery of behavior. Nat Methods Available at: http://dx.doi.org/10.1038/s41592-024-
02200-1.

van der Veen R, Abrous DN, de Kloet ER, Piazza PV, Koehl M (2008) Impact of intra- and interstrain cross-
fostering on mouse maternal care. Genes Brain Behav 7:184–192.

van Oosterhout F, Lucassen EA, Houben T, vanderLeest HT, Antle MC, Meijer JH (2012) Amplitude of the
SCN clock enhanced by the behavioral activity rhythm. PLoS One 7:e39693.

Vaswani A, Shazeer N, Parmar N, Uszkoreit J, Jones L, Gomez AN, Kaiser L, Polosukhin I (2017) Attention Is
All You Need. arXiv [csCL] Available at: http://arxiv.org/abs/1706.03762.

Verwey M, Robinson B, Amir S (2013) Recording and analysis of circadian rhythms in running-wheel activity
in rodents. J Vis Exp Available at: http://dx.doi.org/10.3791/50186.

Wahba LR, Perez B, Nikhil KL, Herzog ED, Jones JR (2022) Circadian rhythms in multiple behaviors depend
on sex, neuropeptide signaling, and ambient light. bioRxiv:2022.08.18.504454 Available at:
https://www.biorxiv.org/content/10.1101/2022.08.18.504454v1 [Accessed September 21, 2022].

Walton JC, Bumgarner JR, Nelson RJ (2022) Sex differences in circadian rhythms. Cold Spring Harb Perspect
Biol 14 Available at:
https://cshperspectives.cshlp.org/content/early/2022/01/31/cshperspect.a039107.short?casa_token=o8jmv7
0ohpsAAAAA:PRsQUgDmTXWRoMV1VShAAsMPEHyzVKZhhxTbcLPrbc8i61ioEFq40E6_CoCWdF
C6y89C6RCr.

Xie S, Sun C, Huang J, Tu Z, Murphy K (2017) Rethinking Spatiotemporal Feature Learning: Speed-Accuracy
Trade-offs in Video Classification. arXiv [csCV] Available at: http://arxiv.org/abs/1712.04851.

Yamanaka Y, Honma S, Honma K-I (2013) Daily exposure to a running wheel entrains circadian rhythms in
mice in parallel with development of an increase in spontaneous movement prior to running-wheel access.
Am J Physiol Regul Integr Comp Physiol 305:R1367–R1375.

Ying X (2019) An Overview of Overfitting and its Solutions. J Phys Conf Ser 1168:022022.

Zhang C, Li H, Han R (2020) An open-source video tracking system for mouse locomotor activity analysis.
BMC Res Notes 13:48.

Zobolas J (n.d.) usefun. https://github.com/bblodfon/usefun Available at: doi.org/10.5281/zenodo.10694717
[Accessed February 22, 2024].