TITLE:

Characterization of natural product inhibitors of quorum sensing in *Pseudomonas aeruginosa* reveals competitive inhibition of RhlR by *ortho*-vanillin

AUTHORS:

Kathryn E. Woods[a], Sana Akhter[b], Blanca Rodriguez[a], Kade A. Townsend[a], Nathan Smith[a], Ben Smith[a], Alice Wambua[a], Vaughn Craddock[a], Rhea G. Abisado-Duque[a], Emma E. Santa[c], Daniel E. Manson[c], Berl R. Oakley[a], Lynn E. Hancock[a], Yinglong Miao[a,b,d], Helen E. Blackwell[c], and Josephine R. Chandler[a#]

[a]Department of Molecular Biosciences, University of Kansas, Lawrence, KS 66045

[b]Center for Computational Biology, University of Kansas, Lawrence, KS 66045

[c]Department of Chemistry, University of Wisconsin–Madison, Madison, WI 53706

[d]Current location: Department of Pharmacology and Computational Medicine Program, University of North Carolina–Chapel Hill, Chapel Hill, NC 27599

Running title: Benzaldehyde activities on LasR and RhlR

[#]Address correspondence to Josephine R. Chandler, jrchandler@ku.edu.

ABSTRACT

Quorum sensing (QS) is a cell-cell signaling system that enables bacteria to coordinate

population density-dependent changes in behavior. This chemical communication pathway is

mediated by diffusible *N*-acyl L-homoserine lactone signals and cytoplasmic signal-responsive

LuxR-type receptors in Gram-negative bacteria. As many common pathogenic bacteria use QS to

regulate virulence, there is significant interest in disrupting QS as a potential therapeutic

strategy. Prior studies have implicated the natural products salicylic acid, cinnamaldehyde and

other related benzaldehyde derivatives as inhibitors of QS in the opportunistic pathogen

Pseudomonas aeruginosa, yet we lack an understanding of the mechanisms by which these

compounds function. Herein, we evaluate the activity of a set of benzaldehyde derivatives using

heterologous reporters of the *P. aeruginosa* LasR and RhlR QS signal receptors. We find that

most tested benzaldehyde derivatives can antagonize LasR or RhlR reporter activation at

micromolar concentrations, although certain molecules also caused mild growth defects and

nonspecific reporter antagonism. Notably, several compounds showed promising RhlR or LasR

specific inhibitory activities over a range of concentrations below that causing toxicity. *ortho*-

Vanillin, a previously untested compound, was the most promising within this set. Competition

experiments against the native ligands for LasR and RhlR revealed that *ortho*-vanillin can

interact competitively with RhlR but not with LasR. Overall, these studies expand our

understanding of benzaldehyde activities in the LasR and RhlR receptors and reveal potentially

promising effects of *ortho*-vanillin as a small molecule QS modulator against RhlR.

IMPORTANCE

Quorum sensing (QS) regulates many aspects of pathogenesis in bacteria and has attracted

interest as a target for anti-virulence therapies. As QS is regulated by low molecular weight

chemical signals, the development of chemical strategies that can interfere with this cell-cell

communication pathway has seen considerable scrutiny over the past 25 years. Much of this

research has focused on common human pathogens, including the LasR and RhlR QS receptors

in *Pseudomonas aeruginosa*. Potent and selective chemical agents capable of blocking the

activity of these receptors remain relatively scarce, however. Natural products have provided a

bounty of chemical scaffolds with anti-QS activities, but their molecular mechanisms are poorly

characterized. The current study serves to fill this void by examining the activity of an important

and wide-spread class of natural product QS modulators, benzaldehydes and related derivatives,

in LasR and RhlR. We demonstrate that *ortho*-vanillin can act as a competitive inhibitor of RhlR,

a receptor that has emerged and may supplant LasR in certain settings as a target for QS control

in *P. aeruginosa*. The results and insights provided herein will advance the design of chemical

tools to study QS with improved activities and selectivities.

INTRODUCTION

Many bacteria sense and respond to changes in population density using a gene regulation system called quorum sensing (QS). QS can regulate diverse behaviors including light production in marine bioluminescent bacteria, virulence factor production in plant and animal pathogens, and motility in many soil bacteria (1). In Proteobacteria, one type of QS system involves N-acyl L-homoserine lactone (AHL) signals (for reviews, see refs. (2, 3)). AHLs are produced by LuxI-type signal synthases, and detected by LuxR-type signal receptors, which are cytoplasmic transcriptional factors. At low population densities, AHLs are produced at low levels and accumulate in the local environment with increasing population density. The AHLs diffuse in and out of the cell, although active efflux can also contribute to the export of certain long chain AHLs (4). AHLs bind to the LuxR-type receptor protein and—for most of the known associative-type receptors—when they reach a critical concentration, they cause conformational changes to the protein that enable binding and activation of target gene promoters. AHLs interact with their cognate LuxR protein by making a series of hydrogen-bonding and hydrophobic contacts with residues in the ligand-binding pocket. AHL binding pockets vary structurally among LuxR family members to ensure specific responses to cognate AHLs, which differ in acyl chain structure.

Pseudomonas aeruginosa is an opportunistic pathogen that can cause debilitating infections in immunocompromised patients and is difficult to treat due to its multi-drug resistance. *P. aeruginosa* has two LuxI/R-type systems, LasI/R and RhlI/R. The LasI/R system produces and responds to N-(3-oxo)-dodecanoyl L-homoserine lactone (3OC12-HSL), and the RhlI/R system produces and responds to N-butanoyl L-homoserine lactone (C4-HSL). Upon AHL binding, LasR and RhlR activate distinct and overlapping regulons (5, 6). Among those are

the genes encoding factors with known roles in virulence, such as the secreted toxins phenazine

and hydrogen cyanide, proteases, and biofilm matrix proteins. These systems have been shown to

be important for *P. aeruginosa* virulence in numerous acute animal infection models (7-11).

Thus, *P. aeruginosa* QS has been proposed as an attractive target for the development of novel

anti-virulence therapeutics (12).

Over the past 30 years, there has been considerable effort to identify molecules that block

QS in *P. aeruginosa* and other bacteria. These prior studies have identified several promising

approaches such as inhibiting LuxI-type synthases (13), destroying or sequestering AHLs (14),

or inhibiting LuxR-type receptors (15). The latter strategy has received the most attention to date

in *P. aeruginosa*, with much focus on the LasR receptor, and more recently RhlR, in *P.*

aeruginosa. As a result, several promising molecules have been identified that inhibit these

receptors (16-19). These molecules have potencies in the high-nM to mid- to low-μM range. In

general, the most potent molecules have been identified as a result of high-throughput screens of

small molecule libraries or by making targeted changes to the native AHL or other promising

lead compounds via chemical synthesis.

In addition to these synthetic agents, there also has been widespread study of readily

available molecules that can be re-purposed as QS inhibitors. Many of these compounds are

natural products and were initially identified because of their ability to block QS-dependent

phenotypes in the native species, not via studies of their ability to target specific QS pathways.

These compounds include halogenated furanones (20), flavonoids such as baicalein (21, 22) and

several benzaldehydes such as cinnamaldehyde (23-28). Despite the widespread use of these

molecules as chemical tools for studies of QS inhibition, relatively little is known of the

specificity, potency, and mechanism of action for most of these compounds. New tools to study

QS are of considerable interest, as many of the known chemical modulators have limitations,

including relatively low potencies, efficacies, solubilities in aqueous media, and/or chemical

stabilities. Consequently, re-purposed bioactive agents and readily-available natural products

(and analogs) with promising QS inhibitory activities represents a valuable space to search for

new chemical probes to study bacterial signaling.

In this study, we used *E. coli* reporters to evaluate the ability of several naturally

occurring benzaldehydes and related derivatives to inhibit the *P. aeruginosa* QS receptors LasR

and RhlR. We focused on compounds reported to disrupt QS-dependent phenotypes in *P.*

aeruginosa, such as cinnamaldehyde and salicylic acid, along with several previously unstudied

compounds with some structural similarity, such as orsellinaldehyde and *ortho*-vanillin (Fig. 1).

We observed antagonism of the *E. coli* LasR and RhlR reporters at concentrations in the mid- to

low-µM range, with *ortho*-vanillin showing the most promising effects. The compounds also

caused mild reductions in growth and could nonspecifically antagonize a constitutive reporter at

higher concentrations; however, at lower concentrations there was a suitable window of activity

allowing for LasR and RhlR antagonism without any observable toxicity. In follow-up structure-

function studies using LasR mutants, we found that critical AHL-binding residues in LasR were

not required for *ortho*-vanillin to antagonize LasR. However, our results support that *ortho*-

vanillin might specifically interact with RhlR. Together, our results indicate that naturally

occurring benzaldehydes could have utility in QS inhibition and motivate future studies to

develop this chemical scaffold into small-molecule tools to explore LuxR-type protein function

and QS pathways.

Fig. 1. Structures of compounds examined in this study. 2H-4MB = 2-

RESULTS

Construction of cell-based *E. coli* bioreporters for LasR

To characterize compounds for their potential activity as LasR antagonists, we used a

heterologous *Escherichia coli* strain expressing LasR from an arabinose-inducible promoter

(P*ara-lasR)* on plasmid pJN105-L and a second plasmid with the LasR-inducible *lasI* promoter

fused to a promoterless *lacZ* reporter (P*lasI-lacZ*) on plasmid pSC11-L (Fig. 2A). In this strain,

lacZ expression required LasR and the LasR signal 3OC12-HSL (Fig. 2A), with a half-maximal

activation concentration (i.e., EC_{50} value) of 65 nM. As a control, we also constructed an *E. coli*

strain carrying a plasmid with *lacZ* expressed from the constitutive *aphA-3* promoter (29, 30),

pVT19. With this strain, *lacZ* expression is fully activated in the absence of LasR or 3OC12-

HSL (Fig. 2B).

Fig. 2. General schematic and initial characterization of *E. coli* bioreporter strains used for these studies. A. An *E. coli* bioreporter of LasR activity carries plasmid pSC11-L with a LasR-inducible P*lasI* promoter fused to the *lacZ* reporter and plasmid pJN105-L with an arabinose-inducible LasR. This strain produces β-galactosidase in response to 3OC12-HSL and 0.4% arabinose. The 3OC12-HSL induction is dose-responsive with a half-maximal concentration of 65 nM. **B.** An *E. coli* strain constitutively expressing *lacZ* carries the plasmid pVT19 with *lacZ* fused to the constitutive *aphA-3* promoter. This strain produces β-galactosidase in the presence or absence of 3OC12-HSL. Results are the averages of two (A) or three (B) independent experiments and the error bars represent the standard deviation.

E. coli reporter assays indicate orsellinaldehyde antagonizes reporter activation nonspecifically.

152
153
We utilized our *E. coli* reporters to evaluate the activity of the natural products and

related derivatives in Fig. 1 as LasR antagonists, and we initiated our study with

orsellinaldehyde, a metabolite produced by the fungus *Aspergillus nidulans* (31). Given its

structural similarity to several known QS inhibitors, such as vanillin and salicylic acid, we were

interested to examine orsellinaldehyde's activity as a LasR antagonist. In the presence of 100 nM

3OC12-HSL (Fig. 3A and Table 1), we found that the concentration of orsellinaldehyde needed

to inhibit P$lasI$-$lacZ$ activity by 50% (i.e., its IC_{50} value) was 2374 µM (Fig. 3A, black line),

indicating weak antagonist activity toward LasR. However, we observed that orsellinaldehyde

caused a dose-dependent reduction of growth yield by about 10-20% at the highest

concentrations (Fig. 3A, grey line). Further, orsellinaldehyde-dependent antagonism of the $lasI$-

$lacZ$ reporter correlated with its increasing effects on growth (correlation coefficient r=0.9877,

p<0.0001, Fig. 3B). These results suggest antagonism of the LasR bioreporter by

orsellinaldehyde may be due to the generalized effects of this compound on growth. To address

this possibility, we generated a dose-response curve with orsellinaldehyde and our constitutive

$lacZ$-producing control $E. coli$ strain (with plasmid pVT19). We found orsellinaldehyde

antagonized the constitutive $lacZ$ reporter in a dose-responsive manner with an IC_{50} of 2308 µM

(Fig. 3A, red line), which was similar to that of the LasR bioreporter (2374 µM). These results

support the conclusion that orsellinaldehyde antagonizes $lacZ$ reporter activation in a nonspecific

manner.

Fig. 3. Activity of orsellinaldehyde with *E. coli* LasR and constitutive reporters. A. Grey line, orsellinaldehyde-dependent growth inhibition as a percent of cells with no orsellinaldehyde. Dose-response curve of orsellinaldehyde in competition with 100 nM 3OC12-HSL in cultures of *E. coli* with arabinose-inducible LasR and a LasR-dependent *lasI-lacZ* reporter (black line) or of *E. coli* with a constitutive *aphA-3-lacZ* reporter (red line). IC50 values are given in Table 1. Results show the averages of five (LasR) or three (LacZ control) independent experiments and the error bars represent the standard deviation. **B**. Average values from A (% growth reduction vs. % inhibition of the *lasI-lacZ* reporter), which were used to determine Pearson's correlation coefficient (r value) and significance (p) and generate a fitted line using a simple linear regression model.

To test whether these effects were specific to the *lacZ* reporter or general to other

reporters, we generated an orsellinaldehyde dose-response curve using a strain constitutively

expressing GFP (*E. coli* carrying a constitutive GFP-producing plasmid pUC18T-mini-Tn7T-

Gm-gfpmut3). Orsellinaldehyde also antagonized the constitutive GFP reporter with an IC_{50} of

1057 µM for GFP (Fig. S1), which was similar to that of the *lacZ* reporter. These results support

the conclusion that the effects of orsellenaldehyde on our LasR bioreporter are related to a

generalized effect on gene expression or other cellular processes and not specific to LasR.

Evaluation of other benzaldehyde derivatives in *E. coli* LasR reporters.

We next examined compounds structurally related to orsellinaldehyde and previously reported to

modulate QS for antagonistic activity in LasR. In view of the results above, we questioned

whether some of the reported inhibitory activities were also largely due to nonspecific toxic

effects. We selected several such compounds; cinnamaldehyde (25), salicylic acid (25-28), and

the as-yet untested, but related compounds *ortho*-vanillin, 2-hydroxy-5-methylbenzaldehyde and

2-hydroxy-4-methoxybenzaldehyde (Fig. 1). The results (Fig. 4, Fig. S2 and Table 1) show that

each of the compounds can antagonize the LasR-dependent *lasI-lacZ* reporter with IC_{50}s ranging

from 437 µM for vanillin to 1674 µM for salicylic acid. We also observed decreases in growth

like that of orsellinaldehyde by ~25% at the highest concentrations (Fig. 4). The effects on

growth and inhibition of the *lasI-lacZ* reporter were significantly correlated for each of the

compounds (Fig. 4, Fig. S2 and Table 1), although there was a weaker correlation for *ortho*-

vanillin and cinnamaldehyde because the effects on growth were minimal at the lower

concentrations (Fig. 4A and C, right side). We also generated dose-response curves of each

compound with the control constitutive *lacZ* reporter strain (Fig. 4). All of these compounds

were less potent with the constitutive reporter than that of the LasR reporter, by 1.3-fold for 2-

hydroxy-5-methylbenzaldehyde to almost 3-fold lower for *ortho*-vanillin. These results suggest

that, while all of the compounds also have nonspecific effects at higher concentrations, certain

compounds—i.e., *ortho*-vanillin and cinnamaldehyde—have some specific activity against LasR

at lower concentrations.

Fig. 4. Activities of cinnamaldehyde (A), salicylic acid (B) and vanillin (C) with *E. coli* LasR and constitutive reporters. Left column: dose-response curves for each indicated compound in competition with 100 nM 3OC12-HSL in cultures of *E. coli* with arabinose-inducible LasR and a LasR-dependent *lasI-lacZ* reporter (black line) or of *E. coli* with a constitutive *aph-lacZ* reporter (red line). Results show the averages of four independent experiments and the error bars represent the standard deviation. IC_{50} values from the fit curves are given in Table 1. Right column shows average values from the graphs on the left (% growth reduction vs. % inhibition of the *lasI-lacZ* reporter), which were used to determine Pearson's correlation coefficient (r value) and significance (p) and generate a fitted line using a simple linear regression model (B) or a second-order polynomial nonlinear regression model (A and C).

Table 1. Potency of benzaldehydes using *E. coli* LasR and constitutive reporters[a]

Compound	$IC_{50} \pm CI$ (μM)[b, c]	
	LasR reporter	**Constitutive reporter**
Orsellinaldehyde	2374 (2283-2469)	2308 (2177-2452)
Salicylic Acid	1674 (1496-1865)	3645 (3451-3855)
Cinnamaldehyde	851 (761-943)	1459 (1395-1528)
ortho-Vanillin	437 (358-523)	1261 (1171-1354)
2-hydroxy-4-methoxybenzaldehyde	1040 (944-1139)	1955 (1801-2121)
2-hydroxy-5-methylbenzaldehyde	1469 (1298-1639)	2019 (1888-2157)

Results of LasR mutant reporters support *ortho*-vanillin not contacting specific residues in

the LasR ligand-binding domain

As *ortho*-vanillin was the most potent LasR antagonist identified above, we sought to

further characterize the nature of potential *ortho*-vanillin/LasR interactions. To our knowledge,

no other studies have experimentally addressed the molecular mechanism by which

benzaldehyde derivatives antagonize LuxR-type receptors. We began by asking whether *ortho*-

vanillin is acting as a competitive LasR antagonist, similar to the synthetic compound V-06-018

(19), and binding in the native ligand (i.e., 3OC12-HSL) binding site. To this end, we applied an

approach of competing *ortho*-vanillin with 3OC12-HSL at varying concentrations using our

LasR reporter assay described above. The reporter experiments in the heterologous *E. coli* host

provide a proxy to assess LasR interaction with AHLs in the absence of other host regulation

effects. The ability of *ortho*-vanillin to antagonize LasR should vary when the 3OC12-HSL

concentration is increased if both molecules are competing for binding to the same site in LasR.

We generated antagonism dose-response curves for *ortho*-vanillin competed against 3OC12-HSL

at 100 nM, 10 μM and 100 μM (Fig. 5A). Although there was a small difference in the *ortho*-

vanillin IC_{50} at 100 nM and 10 μM, this difference was not significant (p>0.07). These results do

not support the conclusion that *ortho*-vanillin is a competitive antagonist of LasR.

In addition, we performed *in silico* docking studies of *ortho*-vanillin within the ligand-

binding site (LBD) of LasR using the reported full-length LasR structure (PDB ID: 6V7X; see

Methods) and found that this compound could be accommodated. Three residues were identified

that could be important for the *ortho*-vanillin/LasR interaction: Thr75, Thr115 and Ser129 (Fig.

S3). These residues were predicted to hydrogen bond with the phenol and aldehyde substituents

of *ortho*-vanillin. Several other residues, such as Tyr56, Trp60, and Tyr93 were also predicted to

form close contacts with *ortho*-vanillin. Ser129 and several other of these residues (e.g., Arg61,

Tyr56 and Asp73) were also found to be important for LasR interaction with 3OC12-HSL and

other ligands (19, 32, 33) (Fig. S3).

To examine these putative interactions between *ortho*-vanillin and LasR, we determined

the activity of *ortho*-vanillin in several LasR mutants. In prior studies in our laboratories, a set of

LasR mutants were generated in which residues within the ligand-binding pocket were mutated

to a different residue of similar steric size but without the capability to hydrogen bond (e.g., Tyr

→ Phe). These mutants were introduced into *E. coli* to generate *lasI-lacZ* reporters analogous to

the wild-type LasR reporter above (see Table S1 and Methods). From that set, we selected five

LasR mutant reporters to test *ortho*-vanillin (W60F, Y56F, T75V, Y93F, and S129A), which

included the Thr75 and Ser129 residues predicted to be important for *o*-vanillin interaction in our

in silico study. Each of these mutants showed varying degrees of activation by 3OC12-HSL in

our reporter experiment (Fig. S4), which was consistent with prior results (33). We generated

dose-response curves with *ortho*-vanillin competed against 3OC12-HSL at the concentration

needed to cause half-maximal LasR activation (EC$_{50}$) for each mutant (Fig. S4). In our

experiments, *ortho*-vanillin antagonism of the LasR mutants was indistinguishable from that of

the wild type LasR (Fig. 5C and S6). As a control, we also tested the ability of V-06-018 to

antagonize the LasR S129A mutant. Consistent with prior results (19), we showed that V-06-018

was significantly less potent with the S129A mutant compared with wild type LasR (Fig. 5D).

Together, these results show that some of the LasR ligand binding site residues that make

important contacts with other ligands (AHL and non-AHL agonists or antagonists) are not

required for *ortho*-vanillin activity and are consistent with the idea that *ortho*-vanillin does not

interact with the LasR ligand-binding domain in a mode analogous to other ligands.

Fig. 5. LasR mutant antagonism data for *ortho*-vanillin and V-06-018. A. Dose-response curves of *o*-vanillin in competition with 3OC12-HSL at its EC$_{50}$ value of 65 nM, 1 μM or 10 μM in *E. coli* with arabinose-inducible LasR and a LasR-dependent *lasI-lacZ* reporter. Each curve shows results of three independent experiments with the standard deviation represented by horizontal bars. **B.** Data show averages of IC$_{50}$ values from each curve shown in panel A. Error bars show the standard deviation. There were no statistical differences between any of the conditions by one-way ANOVA (p>0.3). **C.** *ortho*-Vanillin was tested at varying concentrations (25 nM to 50 μM) against 3OC12-HSL at its EC$_{50}$ value in the specific *E. coli* reporter strain as indicated on the X-axis. The *ortho*-vanillin dose-response curves (shown in Fig. S6) were used to determine the IC$_{50}$ values, which are shown as the % of the IC$_{50}$ of *ortho*-vanillin for *E. coli* with wild type LasR. There were no significant differences of any of the LasR mutant results from that of wild type by one-way ANOVA. **D.** The LasR inhibitor V-06-018 was tested at 100 μM against 3OC12-HSL at its EC$_{50}$ value (65 nM) in *E. coli* with wild type or the S129A LasR as indicated on the X-axis. Values are reported as the % of reporter activation with the EC$_{50}$ of 3OC12-HSL with no other compound. *, statistical significance by students t-test (p<0.05).

Evaluation of benzaldehyde derivatives using *E. coli* RhlR reporters indicate vanillin can

antagonize RhlR

We hypothesized that the benzaldehyde derivatives in our studies (Fig. 1) could be poor

antagonists of LasR because they have very short or no acyl tail functionality, which has been

shown to be important for LasR interactions in studies of the native ligand 3OC12-HSL and

other inhibitors, such as V-06-018 (19, 34). We thus turned our attention to RhlR from *P.*

aeruginosa, which is regulated by an AHL with a much shorter 4-carbon tail, C4-HSL. We

performed *in silico* docking studies analogous to those for LasR above using the recently

published RhlR structure, which was purified with a non-native agonist meta-bromothiolactone

(mBTL) (PDB ID: 8DQ0) (35). We examined docking of *ortho*-vanillin and the native ligand

C4-HSL to RhlR (Fig. S6), and found that both could be accommodated. The phenol moiety of

ortho-vanillin was predicted to hydrogen bond with Asp81 of RhlR, supporting the idea that this

compound could possibly interact with RhlR. The docking score calculations were similar for

ortho-vanillin and the native ligand C4-HSL (about -5.2 kcal/mol), although the specific

interactions with RhlR appeared to be different for C4-HSL, which was predicted to have close

contact with residues Tyr69, Trp93 and Ala108, but not with Asp81.

To test the ability of our set of benzaldehydes and related compounds to antagonize

RhlR, we generated dose-response curves with these compounds using an *E. coli* RhlR reporter

strain (Fig. 6 and Table 2). This strain is analogous to the LasR reporter above but it carries

plasmid pECP61.5 expressing RhlR from the IPTG-inducible P*lac* promoter as well as the *rhlA-*

lacZ reporter (36). We also utilized our constitutive *lacZ* reporter plasmid pVT19 to generate

dose-response curves using the RhlR assay conditions (Fig. 6 and Table 2). The tested

compounds caused maximal ~20% growth reduction for the RhlR conditions. The potencies of

our compounds with the RhlR reporter ranged from an IC_{50} of 151 μM for *ortho*-vanillin to ~10

mM for salicylic acid. With the constitutive *lacZ* reporter, there was a similar spread in potencies

for our compounds, with *ortho*-vanillin having the lowest IC_{50} and salicylic acid as the highest.

However, the IC_{50} for *ortho*-vanillin was 5-fold lower with RhlR than with the constitutive *lacZ*

reporter. There were also no observed effects of *ortho*-vanillin on growth until concentrations at

which there was >50% antagonism of the RhlR reporter, and there was no antagonism of the

constitutive *lacZ* reporter until concentrations >500 μM. These results support the idea that

ortho-vanillin may specifically antagonize RhlR at concentrations below 500 μM.

Fig. 6. Activities of orsellinaldehyde (A) and *o*-vanillin (B) with *E. coli* RhlR and constitutive reporters. Left column: dose-response curves for each indicated compound in competition with 414 nM C4-HSL in cultures of *E. coli* with IPTG-inducible RhlR and a RhlR-dependent *rhlA-lacZ* reporter (black symbols) or of *E. coli* with a constitutive *aph-lacZ* reporter (red symbols). Results show the averages of three independent experiments and the error bars represent the standard deviation. IC$_{50}$ values from the fit curves are given in Table 2. Right column shows average values from the graphs on the left (% growth reduction vs. % *rhlA-lacZ* reporter antagonism), which were used to determine Pearson's correlation coefficient (r value) and significance (p) and generate fitted lines using a second-order polynomial nonlinear regression model.

Table 2. Potency of benzaldehyde derivatives using *E. coli* RhlR and constitutive reporters[a]

	$IC_{50} \pm CI$ (μM)[b, c]	
Compound	**RhlR reporter**	**Constitutive reporter**
Orsellinaldehyde	956 (826-1056)	2745 (2465-3100)
Salicylic Acid	10,049 (8666-11698)	6260 (4842-8302)
Cinnamaldehyde	1308 (1196-1426)	2847 (2641-3086)
ortho-Vanillin	151 (136.1 to 167.7)	872 (749-?)[d]
2-hydroxy-4-methoxybenzaldehyde	458 (409-511)	1355 (1194-1553)
2-hydroxy-5-methylbenzaldehyde	1082 (974-1210)	1789 (1259-2775)

[a]The *E. coli* reporter strain for RhlR carried plasmid pECP61.5 (carrying the inducible P*tac-rhlR* and the *rhlA-lacZ* reporter. The *E. coli* constitutive reporter strain carried plasmid pVT19 expressing *lacZ* constitutively from the *aph* promoter. Results with both reporters were from experiments carried out in the conditions described for the RhlR reporter in the *Materials and Methods*.

[b]Experiments were performed by competing the compounds at a range of concentrations (25 μM – 50 mM) against 400 nM C4-HSL using conditions described for the RhlR reporter in the Materials and Methods. IC_{50} values were determined using a nonlinear best-fit curve with variable parameters with the top and bottom constrained to 100% and 0%, respectively (in all cases the bottom of the computed dose-response curve or maximum inhibition was near 0%). Best-fit curve and IC_{50} calculations were using Prism v10. Full dose-response curves used to generate these data are shown in Fig. 6 and S7.

[c]CI = 95% confidence interval.

[d]CI and slope could not be calculated with a variable parameter model. With a Hill slope set at 1.0, the calculated IC_{50} was 774 with a CI of 531 to 1108.

RhlR reporter data support a competitive mechanism of RhlR antagonism by *ortho-vanillin.*

We were interested to determine whether *o*-vanillin was acting as a competitive RhlR antagonist. As with LasR, we tested whether competing with C4-HSL at different concentrations could elicit changes in the ability of *ortho*-vanillin to antagonize RhlR in the *E. coli lacZ* reporter. We generated antagonism dose-response curves for *ortho*-vanillin competed against C4-HSL at 400 nM, 10 μM, and 100 μM (Fig. 7). We observed a significant C4-HSL concentration-dependent decrease in the potency of *ortho*-vanillin. These differences were most apparent at the lowest concentrations of *ortho*-vanillin, which were below the concentration at

324   which nonspecific antagonism of the *lacZ* reporter were observed. These results are congruent

325   with the ability of *ortho*-vanillin can act as a competitive antagonist of RhlR.

Fig. 7. Dose-response RhlR antagonism by *ortho*-vanillin competed with varying C4-HSL concentrations. A. Dose-response curves of *ortho*-vanillin in competition with 400 nM, 10 µM or 100 µM C4-HSL in cultures of *E. coli* with IPTG-inducible RhlR and a RhlR-dependent *rhlA-lacZ* reporter. Each curve shows results of three independent experiments with the standard deviation represented by horizontal bars. **B.** Data show averages of IC$_{50}$ values from each curve shown in panel A. Error bars show the standard deviation. Statistical significance by one-way anova; *, p<0.05; ** p<0.01; *** p<0.001.

DISCUSSION

The contribution of QS to a wide array of phenotypes, including virulence, in *P.*

aeruginosa has attracted significant attention to the identification of QS inhibitors for use as

chemical probes and in therapeutic development. Despite considerable work in this area, there

are relatively few highly potent and selective QS inhibitors in *P. aeruginosa* and related

proteobacteria. Most of these compounds target LuxR-type receptor proteins, including V-06-

018 that antagonizes LasR in *P. aeruginosa* (18) and the chlorolactone AHL analog (CL) that

antagonizes CviR from *Chromobacterium violaceum* (37, 38). Beyond these classes of synthetic

compounds, there are many naturally derived compounds or extracts that have reported activities

as QS inhibitors in bacteria. For example, salicylic acid can downregulate production of the QS-

controlled virulence factors pyocyanin and elastase and attenuate the ability of *P. aeruginosa* to

infect plants (28). However, detailed studies to determine the molecular mechanisms by which

these natural products elicit their effects on QS are limited. In this study, we evaluate the ability

of salicylic acid, cinnamaldehyde, and several related benzaldehyde derivatives, to antagonize

the *P. aeruginosa* LuxR-type receptors LasR and RhlR using heterologous reporters in *E. coli*.

We provide evidence that one of these compounds, namely *ortho*-vanillin, can specifically

antagonize these receptors within a lower range of concentrations in which they are not generally

toxic. These results provide a basis to guide the use of these compounds in QS studies and

suggest chemical scaffolds to advance for the design of new QS receptor antagonists.

The investigations described here indicate that *ortho*-vanillin can specifically antagonize

LasR and that it does so through a non-competitive mechanism (Fig. 5). There are prior reports

of other compounds that might inhibit LuxR-type receptors noncompetitively. Halogenated

furanones, such as bromofuranone, have been shown to inhibit the *Vibrio fischeri* LuxR receptor

noncompetitively (39). Inhibition might involve a mechanism of increasing turnover of the

receptor protein in the cell (39), although bromofuranone can also be broadly toxic at inhibitory

concentrations (40). Some flavonoids also have been reported to inhibit LasR noncompetitively,

such as baicalein, although in the case of baicalein the mechanism is not known (22). Our

discovery that *ortho*-vanillin can antagonize LasR noncompetitively adds to this list of

noncompetitive antagonists.

In the case of RhlR, *ortho*-vanillin appears to act as a specific, competitive antagonist in

the *E. coli* reporter (Fig. 7). Competitive inhibition is by far the most invoked mechanism for

known LuxR-type inhibitors; the crystal structure of chlorolactone (CL) bound to CviR and

stabilizing an inactive conformation provides perhaps the most compelling support for this

mechanism (38). There are several other known competitive inhibitors of RhlR, most of which

closely resemble its native ligand C4-HSL, and our prior detailed structure-function studies have

revealed portions of the molecules that are essential for strong inhibitory activity (41). With the

recently determined crystal structure of RhlR (35), it is now possible to carry out more detailed

studies to better understand RhlR-ligand binding interactions, including with the native ligand

C4-HSL. Such studies will be interesting to reveal important insight into the mechanism of

RhlR-ligand interactions and advance the design of compounds that can modulate RhlR activity.

Our results with *E. coli* reporters show that *ortho*-vanillin is more potent against RhlR

than LasR. This difference could be due to the relatively small size of this molecule and/or its

lack of an acyl tail. The natural ligand of LasR, 3OC12-HSL, has a long 12-carbon acyl tail,

whereas the RhlR ligand C4-HSL has a much shorter 4-carbon acyl tail. Prior structure-function

studies of LasR and 3OC12-HSL reveal that there are important hydrophobic contacts formed

between the long tail of 3OC12-HSL and residues within the LasR binding pocket (42). These

contacts contribute to the strength and specificity of the interaction with LasR. In addition,

studies with V-06-018 analogs showed that shorter acyl tails weaken LasR interactions (43). In

turn, we have shown that RhlR is both activated and inhibited by AHLs analogs with shorter

tails. *ortho*-Vanillin largely lacks such a hydrophobic tail (Fig. 1), which might weaken its ability

to antagonize LasR, while enhance its ability to engage with RhlR. Our results support the idea

that the hydrophobic tails of ligands play a critical role in the specificity and strength of

interactions with LuxR proteins. As this competitive activity for *ortho*-vanillin in RhlR, and its

non-competitive activity in LasR, were observed in *E. coli* reporter systems, additional

experiments including *in vitro* studies will be necessary to provide further clarity into its

molecular mechanisms of action and the hypotheses outlined here. The relative simplicity of the

ortho-vanillin scaffold suggest straightforward routes to alter its structure and examine impact on

potency and specificity, along with reducing any associated toxicity. Overall, these studies

illustrate the importance of performing rigorous studies to determine the specificity and function

of small molecule QS inhibitors to inform their use as research tools and other applications.

MATERIALS AND METHODS

Culture conditions and reagents

Unless otherwise noted, bacteria were grown at 37 °C in Lysogeny broth (LB; 10 g

tryptone, 5 g tryptone and 5 g NaCl per L), or on LB agar (LBA; 1.5% (weight per volume)

Bacto-Agar). For RhlR bioreporter experiments, growth was at 30 °C and in A medium (44)(60

mM K_2HPO_4, 33 mM KH_2PO_4, 7.5 mM $(NH_4)2SO_4$, 1.7 mM sodium citrate $\cdot 2H_2O$, 0.4%

glucose, 0.05% yeast extract, 1 mM $MgSO_4$). All *E. coli* broth cultures were grown with shaking

at 250 rpm, 18 mm test tubes (for 5 ml cultures) or 125 ml baffled flasks (for 10 ml cultures)

unless otherwise specified. For selection, 100 µg ml^{-1} ampicillin, 10 µg ml^{-1} gentamicin, or 150

µg ml^{-1} spectinomycin were used. For experiments with the RhlR bioreporter strain, A medium

was used as described (36, 45). When needed for induction of LasR or RhlR, we added IPTG

(isopropyl β-D-1-thiogalactopyranoside) at 1 µM final concentration and L-(+)-arabinose at

0.25% final concentration. Native HSLs were suspended in ethyl acetate acidified with 0.01%

glacial acetic acid and added to culture tubes and dried down prior to adding growth medium for

experiments.

We measured β-galactosidase activity with a Tropix Galacto-Light Plus

chemiluminescence kit according to the manufacturer's protocol (Applied Biosystems, Foster

City, CA). Native HSLs (3oxoC12-HSL and C4-HSL) were purchased from Cayman Chemicals

(MI, USA), gentamicin was purchased from GoldBio (MO, USA) and ampicillin and

spectinomycin were purchased from Sigma Aldrich (MO, USA). DMSO (solvent for inhibitor

compounds), IPTG, and L-(+)-arabinose were purchased from Fisher Scientific (PA, USA).

Natural products and benzaldehyde derivatives were purchased from Sigma Aldrich (MO, USA).

V-06-018 was synthesized as previously described (19).

Strains and plasmids.

Strains and plasmids are listed in Table S1. To assess LasR activation of *lasR* expression

in recombinant *E. coli*, we used *E. coli* strain DH5α carrying two plasmids; plasmid pJN105-L

(46) with an arabinose-inducible *P. aeruginosa lasR* and plasmid pSC11-L (47) with the

promoter of the LasR-responsive gene *lasI* fused to a *lacZ* reporter. For some studies, pJN105-L

was replaced with derivatives of this plasmid encoding LasR mutants with single amino acid

substitutions (32). To assess RhlR activation of *rhlR* expression in recombinant *E. coli*, we used

E. coli DH5α with plasmid pECP61.5 (36) with an IPTG-inducible p*tac-rhlR* and a RhlR-

responsive gene *rhlA* fused to the *lacZ* reporter. For constitutive expression of the *lacZ* reporter,

we used *E. coli* DH5α with plasmid pVT19, which has the *lacZ* gene fused to the constitutive

aphA-3 promoter. To construct pVT19, the constitutive *aphA-3* promoter (29, 30) was amplified

from a pTCV-lac derivative using primers Vlac1 and Vlac2 (30). The resulting amplicon was

digested with EcoRI and BamHI and ligated into similarly digested pKS12A (48). The resulting

plasmid with the *aphA-3* promoter transcriptionally fused to *lacZ* was designated pVT19. For

constitutive expression of the *gfp* reporter, we used *E. coli* DH5α with plasmid pUC18T-mini-

Tn7T-Gm-gfpmut3 (49).

Transcription reporter assays in *E. coli*.

To assess LasR activation of *lasR* expression in recombinant *E. coli*, overnight cultures of

E. coli DH5α pSC11-L, pJN105-L were diluted 1:100 into LB containing selection antibiotics

gentamicin and ampicillin in 10 ml cultures. When the cultures reached an OD_{600} of 0.2-0.3, L-

(+)-arabinose was added to a final concentration of 0.25%. The control did not receive L-(+)-

arabinose. The cultures were then grown to an OD_{600} of 0.5-0.6 and 500 µL was added to 1.5 mL

micro centrifuge tubes containing dried 3OC12-HSL. Aliquots (5 µL) of increasing

concentrations of inhibitor test compound stock solution in DMSO were then added to the

designated micro centrifuge tubes containing culture. Tubes containing *E. coli* with just signal

and DMSO were included as controls. After 3 h at 37 °C with shaking, OD_{600} was measured

using a plate reader and β-galactosidase activity was measured as described above.

To assess RhlR activation of *rhlR* expression in recombinant *E. coli*, overnight cultures of

E. coli DH5α pECP61.5 grown at 30 °C in A medium containing antibiotic selection (ampicillin)

and IPTG to induce RhlR expression were diluted to an OD_{600} of 0.1 and 1 mL was added to

culture tubes containing dried C4-HSL. Aliquots (5 µL) of DMSO containing increasing

concentrations of inhibitor test compound or DMSO with no test compound were added to the

designated Eppendorf tubes containing culture. Tubes containing *E. coli* with signal and DMSO

were included as a vehicle control. After 5 h at 30 °C with shaking, OD_{600} was measured using a

plate reader and β-galactosidase activity was measured as described above.

Experiments with the LasR mutants and the constitutive *lacZ* expression plasmid pVT19

or constitutive *gfp* expression plasmid pUC18T-mini-Tn7T-Gm-gfpmut3 were carried out

identically as described above for the LasR or RhlR bioreporter experiments. Results with the

pBT19 constitutive reporter strain was different for the LasR vs. RhlR bioreporter protocols

likely due to differences in growth conditions (temperature and/or growth media).

Computational modeling.

The structure of LasR (PDB ID: 6V7X) and RhlR (PDB ID: 8DQ0) was used for docking studies

using the Lamarckian protocol and the empirical free energy function in AutoDock version 4.2.

The hit search was refined using an improved docking method. The α-β-α sandwich located near

the N-terminal ligand binding domain (LBD) was used as the binding location for docking

calculations. The protein target was prepared using AutoDock 4.2. Hydrogen atoms were added

and the water molecules were removed using the AutoDock Tools (ADT) module included in

AutoDock. Charges were adjusted using AutoDock's Gasteiger charges module for proteins, and

atom type was modified to ADT type for calculations. In our calculations, we dock the ligand

(natural or *ortho*-vanillin) with ligand-free LasR or RhlR. For each type of atom in the ligand

being docked, AutoDock needs a pre-calculated grid map. These maps are calculated using

AutoGrid. The Gasteiger-Marsili method was used to determine the atomic charges of the

protein. The AutoGrid application created mass-centered grid maps with 80 grid points in each

direction and 0.375 spacing. Ten different docking runs for the ligand were carried out, followed

by the evaluation of docking results for the binding mechanism and conserved interactions, such

as hydrogen bonds and hydrophobic interactions, between the hits and the LasR or RhlR binding

site. The common interactions of the ligand-docked complexes were analyzed and the one with

the best binding score based on the binding free energy was reported.

Statistical analyses

All statistical analyses (one-way anova, students t-test) were done using Prism v10. IC_{50} and EC_{50} curves

were fitted using a nonlinear regression model with a variable slope unless otherwise stated.

ACKNOWLEDGEMENTS

This work was supported by the NIH through grant R35GM133572 to J.R.C and R35 GM131817

to H.E.B and by Inez Jay Fund to J.R.C. V.D.C. was supported by an Undergraduate Research

Award from the KU Center for Undergraduate Research and a K-INBRE fellowship (P20

GM103418). K.A.T. was supported by KU Center for Undergraduate research Emerging

Scholars program, U.S. Department of Education McNair Scholars Program, and the NIH

Maximizing Access to Research Careers program (MARC) (T34GM136453-01). V.D.C. was

supported by an Undergraduate Research Award from the KU Center for Undergraduate

Research and a K-INBRE fellowship (P20 GM103418). R.G.A. was supported by the Fulbright

Foreign Student Program (15160174).

REFERENCES

1. Abisado RG, Benomar S, Klaus JR, Dandekar AA, Chandler JR. 2018. Bacterial quorum sensing and microbial community interactions. mBio 9:e02331-17.

2. Fuqua WC, Winans SC, Greenberg EP. 1994. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. J Bacteriol 176:269-75.

3. Papenfort K, Bassler BL. 2016. Quorum sensing signal-response systems in Gram-negative bacteria. Nat Rev Microbiol 14:576-88.

4. Moore JD, Gerdt JP, Eibergen NR, Blackwell HE. 2014. Active efflux influences the potency of quorum sensing inhibitors in *Pseudomonas aeruginosa*. Chembiochem 15:435-42.

5. Schuster M, Lostroh CP, Ogi T, Greenberg EP. 2003. Identification, timing, and signal specificity of *Pseudomonas aeruginosa* quorum-controlled genes: a transcriptome analysis. J Bacteriol 185:2066-79.

6. Wagner VE, Bushnell D, Passador L, Brooks AI, Iglewski BH. 2003. Microarray analysis of *Pseudomonas aeruginosa* quorum-sensing regulons: effects of growth phase and environment. J Bacteriol 185:2080-95.

7. Rumbaugh KP, Griswold JA, Hamood AN. 1999. Contribution of the regulatory gene *lasR* to the pathogenesis of *Pseudomonas aeruginosa* infection of burned mice. J Burn Care Rehabil 20:42-9.

8. Tang HB, DiMango E, Bryan R, Gambello M, Iglewski BH, Goldberg JB, Prince A. 1996. Contribution of specific *Pseudomonas aeruginosa* virulence factors to pathogenesis of pneumonia in a neonatal mouse model of infection. Infect Immun 64:37-43.

9. Tan MW, Mahajan-Miklos S, Ausubel FM. 1999. Killing of *Caenorhabditis elegans* by *Pseudomonas aeruginosa* used to model mammalian bacterial pathogenesis. Proc Natl Acad Sci U S A 96:715-20.

10. Mahajan-Miklos S, Tan MW, Rahme LG, Ausubel FM. 1999. Molecular mechanisms of bacterial virulence elucidated using a *Pseudomonas aeruginosa-Caenorhabditis elegans* pathogenesis model. Cell 96:47-56.

11. Tan MW, Rahme LG, Sternberg JA, Tompkins RG, Ausubel FM. 1999. Pseudomonas aeruginosa killing of Caenorhabditis elegans used to identify *P. aeruginosa* virulence factors. Proc Natl Acad Sci U S A 96:2408-13.

12. Greenberg EP. 2003. Bacterial communication and group behavior. The Journal of Clinical Investigation 112:1288-1290.

13. Christensen QH, Grove TL, Booker SJ, Greenberg EP. 2013. A high-throughput screen for quorum-sensing inhibitors that target acyl-homoserine lactone synthases. Proc Natl Acad Sci U S A 110:13815-20.

14. Amara N, Krom BP, Kaufmann GF, Meijler MM. 2011. Macromolecular inhibition of quorum sensing: enzymes, antibodies, and beyond. Chem Rev 111:195-208.

15. Welsh MA, Blackwell HE. 2016. Chemical probes of quorum sensing: from compound development to biological discovery. FEMS Microbiol Rev 40:774-94.

16. O'Loughlin CT, Miller LC, Siryaporn A, Drescher K, Semmelhack MF, Bassler BL. 2013. A quorum-sensing inhibitor blocks *Pseudomonas aeruginosa* virulence and biofilm formation. Proc Natl Acad Sci U S A 110:17981-6.

17. O'Reilly MC, Blackwell HE. 2016. Structure-based design and biological evaluation of triphenyl scaffold-based hybrid compounds as hydrolytically stable modulators of a LuxR-Type quorum sensing receptor. ACS Infect Dis 2:32-38.

18. Muh U, Schuster M, Heim R, Singh A, Olson ER, Greenberg EP. 2006. Novel *Pseudomonas aeruginosa* quorum-sensing inhibitors identified in an ultra-high-throughput screen. Antimicrob Agents Chemother 50:3674-9.

19. Manson DE, O'Reilly MC, Nyffeler KE, Blackwell HE. 2020. Design, synthesis, and biochemical characterization of non-native antagonists of the *Pseudomonas aeruginosa* quorum sensing receptor LasR with nanomolar IC50 values. ACS Infect Dis 6:649-661.

20. Givskov M, de Nys R, Manefield M, Gram L, Maximilien R, Eberl L, Molin S, Steinberg PD, Kjelleberg S. 1996. Eukaryotic interference with homoserine lactone-mediated prokaryotic signalling. J Bacteriol 178:6618-22.

21. Luo J, Dong B, Wang K, Cai S, Liu T, Cheng X, Lei D, Chen Y, Li Y, Kong J, Chen Y. 2017. Baicalin inhibits biofilm formation, attenuates the quorum sensing-controlled virulence and enhances Pseudomonas aeruginosa clearance in a mouse peritoneal implant infection model. PLoS One 12:e0176883.

22. Paczkowski JE, Mukherjee S, McCready AR, Cong JP, Aquino CJ, Kim H, Henke BR, Smith CD, Bassler BL. 2017. Flavonoids suppress *Pseudomonas aeruginosa* virulence through allosteric inhibition of quorum-sensing receptors. J Biol Chem 292:4064-4076.

23. Shastry RP, Ghate SD, Sukesh Kumar B, Srinath BS, Kumar V. 2022. Vanillin derivative inhibits quorum sensing and biofilm formation in *Pseudomonas aeruginosa:* a study in a *Caenorhabditis elegans* infection model. Nat Prod Res 36:1610-1615.

24. Dua T, Mangal S, Akshita G, Harshdeep, Atri AK, Sharma P, Harjai K, Singh V. 2023. Novel Vanillin-based hybrids inhibit quorum sensing and silences phenotypical expressions in Pseudomonas aeruginosa. Drug Development Research 84:45-61.

25. Ahmed SAKS, Rudden M, Smyth TJ, Dooley JSG, Marchant R, Banat IM. 2019. Natural quorum sensing inhibitors effectively downregulate gene expression of *Pseudomonas aeruginosa* virulence factors. Applied Microbiology and Biotechnology 103:3521-3535.

26. Yang L, Rybtke MT, Jakobsen TH, Hentzer M, Bjarnsholt T, Givskov M, Tolker-Nielsen T. 2009. Computer-aided identification of recognized drugs as *Pseudomonas aeruginosa* quorum-sensing inhibitors. Antimicrob Agents Chemother 53:2432-43.

27. Tan SY, Chua SL, Chen Y, Rice SA, Kjelleberg S, Nielsen TE, Yang L, Givskov M. 2013. Identification of five structurally unrelated quorum-sensing inhibitors of *Pseudomonas aeruginosa* from a natural-derivative database. Antimicrob Agents Chemother 57:5629-41.

28. Prithiviraj B, Bais HP, Weir T, Suresh B, Najarro EH, Dayakar BV, Schweizer HP, Vivanco JM. 2005. Down regulation of virulence factors of *Pseudomonas aeruginosa* by salicylic acid attenuates its virulence on *Arabidopsis thaliana* and *Caenorhabditis elegans*. Infection and Immunity 73:5319-5328.

572   29.   Trieu-Cuot P, Klier A, Courvalin P. 1985. DNA sequences specifying the transcription of the
573   streptococcal kanamycin resistance gene in *Escherichia coli* and *Bacillus subtilis*. Mol Gen Genet
574   198:348-52.

575   30.   Poyart C, Trieu-Cuot P. 1997. A broad-host-range mobilizable shuttle vector for the construction
576   of transcriptional fusions to beta-galactosidase in gram-positive bacteria. FEMS Microbiol Lett
577   156:193-8.

578   31.   Ahuja M, Chiang Y-M, Chang S-L, Praseuth MB, Entwistle R, Sanchez JF, Lo H-C, Yeh H-H,
579   Oakley BR, Wang CCC. 2012. Illuminating the diversity of aromatic polyketide synthases in
580   *Aspergillus nidulans*. Journal of the American Chemical Society 134:8212-8221.

581   32.   Gerdt JP, McInnis CE, Schell TL, Rossi FM, Blackwell HE. 2014. Mutational analysis of the
582   quorum-sensing receptor LasR reveals interactions that govern activation and inhibition by
583   nonlactone ligands. Chem Biol 21:1361-1369.

584   33.   Gerdt JP, McInnis CE, Schell TL, Blackwell HE. 2015. Unraveling the contributions of
585   hydrogen-bonding interactions to the activity of native and non-native ligands in the quorum-
586   sensing receptor LasR. Org Biomol Chem 13:1453-62.

587   34.   Passador L, Tucker KD, Guertin KR, Journet MP, Kende AS, Iglewski BH. 1996. Functional
588   analysis of the *Pseudomonas aeruginosa* autoinducer PAI. J Bacteriol 178:5995-6000.

589   35.   Feathers JR, Richael EK, Simanek KA, Fromme JC, Paczkowski JE. 2022. Structure of the RhlR-
590   PqsE complex from *Pseudomonas aeruginosa* reveals mechanistic insights into quorum-sensing
591   gene regulation. Structure 30:1626-1636.e4.

592   36.   Pearson JP, Gray KM, Passador L, Tucker KD, Eberhard A, Iglewski BH, Greenberg EP. 1994.
593   Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes.
594   Proc Natl Acad Sci U S A 91:197-201.

595   37.   Swem LR, Swem DL, O'Loughlin CT, Gatmaitan R, Zhao B, Ulrich SM, Bassler BL. 2009. A
596   quorum-sensing antagonist targets both membrane-bound and cytoplasmic receptors and controls
597   bacterial pathogenicity. Mol Cell 35:143-53.

598   38.   Chen G, Swem LR, Swem DL, Stauff DL, O'Loughlin CT, Jeffrey PD, Bassler BL, Hughson FM.
599   2011. A strategy for antagonizing quorum sensing. Mol Cell 42:199-209.

600   39.   Manefield M, Rasmussen TB, Henzter M, Andersen JB, Steinberg P, Kjelleberg S, Givskov M.
601   2002. Halogenated furanones inhibit quorum sensing through accelerated LuxR turnover.
602   Microbiology (Reading) 148:1119-1127.

603   40.   Moore JD, Rossi FM, Welsh MA, Nyffeler KE, Blackwell HE. 2015. A comparative analysis of
604   synthetic quorum sensing modulators in *Pseudomonas aeruginosa*: new insights into mechanism,
605   active efflux susceptibility, phenotypic response, and next-generation ligand design. J Am Chem
606   Soc 137:14626-39.

607   41.   Boursier ME, Moore JD, Heitman KM, Shepardson-Fungairino SP, Combs JB, Koenig LC, Shin
608   D, Brown EC, Nagarajan R, Blackwell HE. 2018. Structure-function analyses of the N-butanoyl
609   l-homoserine lactone quorum-sensing signal define features critical to activity in RhlR. ACS
610   Chem Biol 13:2655-2662.

611   42.   McCready AR, Paczkowski JE, Henke BR, Bassler BL. 2019. Structural determinants driving
612   homoserine lactone ligand selection in the *Pseudomonas aeruginosa* LasR quorum-sensing
613   receptor. Proc Natl Acad Sci U S A 116:245-254.

614   43.   Blackwell HE, Fuqua C. 2011. Introduction to bacterial signals and chemical communication.
615   Chem Rev 111:1-3.

616   44.   Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular cloning: A laboratory manual. Cold Spring
617   Harbor Laboratory.

618   45.   Pearson JP, Pesci EC, Iglewski BH. 1997. Roles of *Pseudomonas aeruginosa las* and *rhl* quorum-
619   sensing systems in control of elastase and rhamnolipid biosynthesis genes. J Bacteriol 179:5756-
620   67.

621   46.   Lee JH, Lequette Y, Greenberg EP. 2006. Activity of purified QscR, a *Pseudomonas aeruginosa*
622   orphan quorum-sensing transcription factor. Mol Microbiol 59:602-9.

623   47.   Chugani SA, Whiteley M, Lee KM, D'Argenio D, Manoil C, Greenberg EP. 2001. QscR, a
624   modulator of quorum-sensing signal synthesis and virulence in *Pseudomonas aeruginosa*. Proc
625   Natl Acad Sci U S A 98:2752-7.

626   48.   Varahan S, Iyer VS, Moore WT, Hancock LE. 2013. Eep confers lysozyme resistance to
627   *Enterococcus faecalis* via the activation of the extracytoplasmic function sigma factor SigV. J
628   Bacteriol 195:3125-34.

629   49.   Choi KH, Schweizer HP. 2006. mini-Tn7 insertion in bacteria with single attTn7 sites: example
630   *Pseudomonas aeruginosa*. Nat Protoc 1:153-61.