Ilamycins are dual inhibitors of ClpX and ClpC1 involved in proteostasis in Mycobacteria

Yamin Gao[1,2,3,4#], Cuiting Fang[1,2,3,4#], Biao Zhou[1,5,6], H.M. Adnan Hameed[1,2,3,4], Changli Sun[3,7], Xirong Tian[1,2,3,4], Jing He[1,2,4,8], Xingli Han[1,2,3,4], Han Zhang[1,2,4,9], Jianhua Ju[3,7], Xinwen Chen[6], Nanshan Zhong[6], Junying Ma[3,7*], Xiaoli Xiong[1,2,3,6*], Tianyu Zhang[1,2,3,4,6*]

[1] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[2] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious Diseases, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[3] University of Chinese Academy of Sciences, Beijing 100049, China

[4] China-New Zealand Joint Laboratory on Biomedicine and Health, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[5] Guangzhou Medical University, Guangzhou 510180, China

[6] Guangzhou National Laboratory, Guangzhou 510005, China

[7] CAS Key Laboratory of Tropical Marine Bio-Resources and Ecology, Guangdong Key Laboratory of Marine Materia Medica, RNAM Center for Marine Microbiology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China.

[8] Institute of Physical Science and Information Technology, Anhui University, Hefei 230601, China

[9] School of Life Sciences, University of Science and Technology of China, Hefei 230026, China

[#] These authors contributed equally to this work.

[*]Corresponding authors: Tianyu Zhang (Tel: +86-20-32015270, E-mail: zhang_tianyu@gibh.ac.cn); Xiaoli Xiong (Tel:+8617815390976, E-mail: xiong_xiaoli@gibh.ac.cn); Junying Ma (Tel: +86-20-34066449, Email: majunying@scsio.ac.cn).

Abstract

Combatting drug-resistant *Mycobacterium tuberculosis* (Mtb) necessitates the discovery of novel anti-Mtb agents targeting unexploited molecular pathways. The Clp protease system, integral to protein homeostasis in Mtb, has emerged as a promising target. We report that ilamycins exhibit potent antimycobacterial activity by specifically targeting ClpX and ClpC1. Spontaneous resistance to ilamycin E (ILE) and ilamycin F (ILF) in Mtb mutants correlated with mutations in ClpC1 and ClpX. Molecular docking simulations suggest these mutations occur at critical residues in the N-terminal domain (NTD) of MtbClpC1 and MtbClpX. Functional analyses, including gene over-expression in *Mycobacterium marinum* and targeted gene editing in *Mycobacterium abscessus* (Mab) and *Mycobacterium smegmatis* (Msm), substantiated the significance of these mutations in conferring ILE and ILF resistance. Gene silencing heightened the susceptibility of recombinant Mab and Msm strains to ILE and ILF, concurrently impairing their growth. Differential scanning fluorimetry confirmed the stabilization of MtbClpC1-NTD by ILE and ILF. Moreover, ILE markedly impeded the proteolytic function of the ClpC1P1P2 complex without diminishing ClpC1 ATPase activity. Collectively, our findings demonstrate that ilamycins exert mycobactericidal effects by disrupting ClpC1 and ClpX functions, leading to proteostasis dysregulation. This study underscores the therapeutic potential of targeting the Clp protease system in the treatment of tuberculosis.

Introduction

Tuberculosis (TB), caused by *Mycobacterium tuberculosis* (Mtb), is among the most lethal bacterial infections worldwide, accounting for over 1.4 million fatalities annually[1]. Despite the availability of treatments, TB remains a major public health threat on a global scale, aggravated by the proliferation of drug-resistant strains, including multidrug-resistant, extensively drug-resistant, and total drug-resistant Mtb[2,3]. Furthermore, infections caused by nontuberculous mycobacteria (NTM) are on the rise, presenting significant clinical challenges due to their inherent resistance to many available antibiotics[4]. NTM infections lead to serious respiratory, dermatological, and mucosal conditions, particularly in immunocompromised individuals[5]. Therefore, discovery and development of drugs of novel molecular mechanisms with efficacy against drug-resistant mycobacteria are urgently needed for the management of mycobacterial diseases.

Mycobacterial Clp proteases, vital multi-subunit protein complexes involved in intracellular protein degradation, are emerging as promising therapeutic targets[6-9]. The Clp proteases are composed of ClpP1 and ClpP2 peptidases, and an ATP-dependent AAA+ unfoldase (ClpX or ClpC1)[9]. In mycobacteria, ClpP1 and ClpP2 form homo-heptameric rings that subsequently assemble into an active hetero-tetradecamer protease. ClpX and ClpC1, each forming homo-hexameric rings, associate with ClpP2 to facilitate protein unfolding and translocation into the protease chamber via ATP hydrolysis[10,11]. Indispensable for mycobacterial survival[9,12,13], these enzymes not only maintain intracellular protein quality but also modulate responses to environmental stressors and contribute to the pathogenicity of virulent strains[7,14].

Ilamycins, also known as rufomycins, are a group of cycloheptapeptides synthesized by *Streptomyces atratus* SCSIO ZH16 derived from the South China Sea[15]. These compounds, particularly ilamycin E (ILE) and ilamycin F (ILF), have demonstrated potent activity against Mtb H37Rv[15]. Despite their promising anti-Mtb properties, the mode of action of ilamycins, the activity against NTM and other non-mycobacteria pathogenic bacteria, and potential drug interactions are still unknown.

52 Moreover, the efficacy of ilamycins in macrophages and against drug-resistant clinical

53 Mtb strains is yet to be elucidated. In this study, we comprehensively investigated the

54 mechanism of action of ilamycins and examined their activity extensively.

Results

Selective and potent *in vitro* and *ex vivo* antibacterial activity of ILE and ILF

ILE exhibited strong inhibitory effects on Mtb, *Mycobacterium marinum* (Mmr), *Mycobacterium abscessus* (Mab), and *Mycobacterium smegmatis* (Msm) (Table 1), while ILF was notably active against Mtb, Mmr, and Msm, with less pronounced activity against Mab. This variation in efficacy may be attributed to structural differences between ILE and ILF (Fig. 1) and the distinct cell wall permeabilities of the mycobacterial species[16,17]. The minimal inhibitory concentrations (MICs) of ILE to $H_{37}Rv$ and clinically isolated drug-resistant Mtb strains ranged from 0.01 to 0.05 μg mL^{-1}, consistent with that of the laboratory standard Mtb $H_{37}Rv$ strain (0.05 μg mL^{-1}). Synergistic effects were observed when ILE was combined with rifampicin (RIF) or the new anti-TB drug TB47 currently in the preclinical stage[18], and ILF with isoniazid (INH) or TB47, while partial synergy was noted when ILE combined with INH and ILE with ethambutol (EMB). Additive effects were seen when ILE combined with streptomycin (STR), ILF with EMB, and ILF with STR. No antagonism was observed between any of the drugs selected and ILE or ILF. In mouse RAW264.7 macrophage, ILE effectively inhibited autoluminescent Mtb H37Rv (AlRv, MIC, 0.2 μg mL^{-1}). However, neither ILE nor ILF showed activity against common non-mycobacterial clinical isolates, such as *Staphylococcus aureus*, *Acinetobacter baumannii*, *Enterococcus faecium*, *Klebsiella pneumoniae*, and *Pseudomonas aeruginosa* (MICs > 128 μg mL^{-1}).

Ilamycin E₁, R = αOH
Ilamycin E₂, R = βOH

Ilamycin F

Fig. 1 | The structure of ilamycin E and ilamycin F.

Table 1 | Ilamycin E and ilamycin F spectrum of activities

Strains	ILE MIC (μg mL^{-1})	ILF MIC (μg mL^{-1})
AlRa	0.05[a]	1.25[a]
AlRv	0.05[a]	-
AlMsm	0.0625[a]	2.5[a]
AlMab	1[a]	20[a]
AlMmr	0.08[a]	0.625[a]
H$_{37}$Rv	0.025[b]	2.5[b]
K11*	0.025-0.05[b]	-
K12*	0.025[b]	-
K13*	0.01[b]	-
K14*	0.01[b]	-
K16*	0.05[b]	-
K17*	0.01[b]	-
K18*	0.025-0.05[b]	-
K19*	0.01[b]	-
K20*	0.05[b]	-
Staphylococcus aureus	> 128	> 128
Acinetobacter baumannii	> 128	> 128
Enterococcus faecium	> 128	> 128
Klebsiella pneumoniae	> 128	> 128
Pseudomonas aeruginosa	> 128	> 128

AlRa, autoluminescent Mtb H$_{37}$Ra; AlRv, autoluminescent Mtb H$_{37}$Rv; AlMsm, autoluminescent Msm; AlMab, autoluminescent Mab; AlMmr, autoluminescent Mmr; *, clinical resistant Mtb; [a], MICs obtained by detection of relative light units; [b], MICs obtained by the microplate alamar blue assay (MABA) method.

Mutation patterns of *clpC1* and/or *clpX* in ILE- or ILF-resistant Mtb, Mab, Msm, and Mmr strains

We screened and isolated spontaneously generated mutants resistant to ILE or ILF. The spontaneous mutation rate screened for ILE-resistant Mtb at 0.5 μg mL^{-1} was approximately 2.5×10^{-9}, notably lower than the rates for RIF or INH[19]. Whole genome sequencing (WGS) revealed predominant mutations at P30 of ClpX and F80 of ClpC1 in ILE-resistant Mtb strains (Table 2). Whereas, ILE-resistant Mab had mutations at the same site in ClpX and different sites in ClpC1, including F2, Q17, H77, L88, an insertion leading to an M1-FERF-F2 sequence (Table 2). However, ILE-resistant Mmr strains carried mutations exclusively in ClpC1, with no alterations in ClpX (Table 2). Similarly, ILF-resistant Msm mutants showed mutations only in ClpC1, at F2, F5, F80, L88, I197, and an insertion between M1 and F2 (Table 2), respectively. Cross-resistance was observed between ILE and ILF, with all ILE-resistant Mab strains showing resistance to ILF and vice versa for ILF-resistant Msm strains. These findings suggest that both ILE and ILF may target the ClpX and ClpC1 proteases.

Table 2 | Mutations of ClpC1 and ClpX in ILE- and ILF-resistant mutants

Strains No.	Drug/MICs ($\mu g\ mL^{-1}$)	Mutation in ClpC1	Mutation in *clpC1*	Mutation in ClpX	Mutation in *clpX*
Mtb^R1#	ILE 0.625	-	-	P30H	CCC→CAC
Mtb^R2#	ILE 20	F80I	TTT→ATT	P30H	CCC→CAC
Mtb^R3#	ILE 20	F80V	TTT→GTT	P30H	CCC→CAC
Mtb^R4#	ILE 0.1	F80I	TTT→ATT	-	-
Mtb^R5#	ILE 0.1	F80V	TTT→GTT	-	-
Mtb^R6#	ILE 0.1	F80L	TTT→CTT	-	-
Mtb^R7#	ILE 20	F80C	TTT→TGT	-	-
Mab^R1#	ILE > 128 ILF > 128	-	-	P30H	CCC→CAC
Mab^R2#	ILE 4 ILF > 128	F2L	TTC→TCC	-	-
Mab^R3#	ILE 4 ILF > 128	M1-FERF-F2	Ins: TTCGAGAGATTC	-	-
Mab^R4#	ILE 4 ILF > 128	Q17R	CAA→CGA	-	-
Mab^R5#	ILE > 128 ILF > 128	H77D	CAC→GAC	-	-
Mab^R6#	ILE > 128 ILF > 128	L88R	CTG→CGG	-	-
Mmr^R1#	ILE 1 ILF 4	F2L	TTC→TCC	-	-
Mmr^R2#	ILE 10 ILF 40	H77D	CAC→GAC	-	-
Mmr^R3#	ILE 10 ILF 40	L88R	CTC→CGC	-	-
Msm^R1*	ILE 0.5 ILF 20	F2S	TTT→TCT	-	-
Msm^R2*	ILE 0.25 ILF 10	F2C	TTT→TGT	-	-
Msm^R3*	ILE 1 ILF 20	M1-L-F2	Ins: GTT	-	-
Msm^R4*	ILE 1 ILF 10	F5I	TTT→ATT	-	-
Msm^R5*	ILE 0.5 ILF 40	F80S	TTC→TCC	-	-
Msm^R6*	ILE 0.25 ILF 10	L88P	CTC→CCC	-	-
Msm^R7*	ILE 0.25 ILF 20	L88R	CTC→CGC	-	-

Msm^{R8*}	ILE 0.25 ILF 10	F2S; I197N	TTT→TCT; ATC→AAC	-	-

[Table note — italic] #, Resistant mutants were selected from ILE-containing plates; *, Resistant mutants were selected from ILF-containing plates; Ins, insertion; Underlines indicate mutated bases; - showed no mutation.

Molecular docking simulation between ILE and MtbClpC1-N-terminal domain (MtbClpC1-NTD) as well as MtbClpX-NTD

Through molecular docking, we simulated the interaction between ILE and the

MtbClpC1-NTD (Fig. 2a) as well as the MtbClpX-NTD (Fig. 2b). The simulations

identified key binding residues within MtbClpC1-NTD and MtbClpX-NTD, aligning

with mutation sites such as Q17, H77, F80, and L88 in ClpC1-NTD, and the P30 in

MtbClpX-NTD. Given the structural similarities between ILE and ILF, and their

observed cross-resistance, we did not perform separate docking simulations for ILF.

The docking outcomes support the hypothesis that both ILE and ILF target ClpC1 and

ClpX.

Fig. 2 | The graphical representation of crucial residues in the interaction between ILE and MtbClpC1-NTD as well as MtbClpX-NTD. a Crucial residues in the interaction between ILE and MtbClpC1-NTD. **b** Crucial residues in the interaction between ILE and MtbClpX-NTD. MtbClpC1-NTD, N-terminal domain of Mtb-ClpC1 (residues 1–145); MtbClpX-NTD, N-terminal domain of Mtb-ClpX (residues 1–112).

Resistance to ILE and ILF detected in *clpC1* or *clpX* edited Mab and Msm

We obtained 4 gene-edited strains successfully, 2 in Mab (wild-type ClpC1 (ClpC1[wt]) to ClpC1[F2L]; ClpX[wt] to ClpX[P30H]) and 2 in Msm (ClpC1[wt] to ClpC1[M1-L-F2]; ClpC1[wt] to ClpC1[I197N]) (Table 3) by CRISPR/Cpf1-mediated gene editing[20,21]. These edited strains exhibited different resistance levels (Table 3). Notably, the ClpX[wt] to ClpX[P30H] mutation in Mab conferred the highest resistance level , with MICs exceeding 128 μg mL[-1] for ILE and ILF. The ClpC1[wt] to ClpC1[F2L] mutation in Mab resulted in a

119   lower resistance increase, with an 8-fold MIC increase for ILE compared to the parent

120   strain. In Msm, the ClpC1^{wt} to ClpC1$^{M1-L-F2}$ mutation raised the MICs of ILE and ILF

121   to 4 and 6.4 times that of the wild-type strains, respectively, while the ClpC1^{wt} to

122   ClpC1^{I197N} mutation did not affect the MICs.

123   **Table 3 | MICs of ILE and ILF to gene edited strains and corresponding spontaneous mutants**

Strains	Amino acid change	ILE MIC (μg mL^{-1})/ (fold change)	ILF MIC (μg mL^{-1})/ (fold change)
Mab^{wt}	/	1	20
Mab$^{R2\#}$	ClpC1^{wt} to ClpC1^{F2L}	4 / (4)	> 128 / (> 6)
Mab-C1^{F2L}	ClpC1^{wt} to ClpC1^{F2L}	8 / (8)	> 128 / (> 6)
Mab$^{R1\#}$	ClpX^{wt} to ClpC1^{P30H}	> 128 / (> 128)	> 128 / (> 6)
Mab-X^{P29H}	ClpX^{wt} to ClpC1^{P30H}	> 128 / (> 128)	> 128 / (> 6)
Msm^{wt}	/	0.0625	2.5
Msm^{R3*}	ClpC1^{wt} to ClpC1$^{M1-L-F2}$	1 / (16)	20 / (8)
Msm-C1$^{M1-L-F2}$	ClpC1^{wt} to ClpC1$^{M1-L-F2}$	0.25 / (4)	16 / (6.4)
Msm^{R8*}	ClpC1^{wt} to ClpC1^{F2S} ClpC1^{wt} to ClpC1^{I197N}	0.25 / (4)	10 / (4)
Msm-C1^{I197N}	ClpC1^{wt} to ClpC1^{I197N}	0.0625 / (1)	2.5 / (1)

$^{\#}$, Resistant mutants were selected from ILE-containing plates. *, Resistant mutants were selected from ILF-containing plates. -C1 or -X means ClpC1 or ClpX was edited.

Sensitivity of Mab, Mmr or Msm overexpressing genes encoding subunit(s) of

Clp proteases to ILE or ILF

The *clpX*^{wt} and *clpC1*^{wt} genes and their mutant variants were cloned into the extra-

chromosome plasmids with different resistance marker, p60A or pMVA

(Supplementary Fig. 1), under the strong promoter *hsp60* for overexpression in

autoluminescent strains of Mab (AlMab)[22], Msm (AlMsm)[23], and Mmr (AlMmr)[24].

Neither overexpression of the wild-type nor mutated ClpC1 (F2L, Q17R, H77D, and

L88R) or ClpX (P30H) in AlMab altered sensitivity to ILE. Even concurrent

overexpression of all three genes encoding the corresponding subunits (*clpX*, *clpP1*,

clpP2 or *clpC1*, *clpP1*, *clpP2*) in AlMab did not confer ILE resistance. Similarly, in

AlMsm, overexpression of either wild-type or mutated ClpC1 (F2S, F2C, F5I, F80S,

L88P, and L88R) did not change sensitivity to ILF. In AlMmr, overexpressing ClpC1^{F2L}

or ClpC1^{H77D} conferred no resistance to both ILE and ILF. However, AlMmr strain

overexpressing ClpC1[L88R] showed resistance to both ILE and ILF compared to the

parent strain (Table 4).

Table 4 | MICs of ILE and ILF to AlMmr strains overexpressing ClpC1 variants

Strains	ILE MIC (µg mL^{-1}) / (fold change)	ILF MIC (µg mL^{-1}) / (fold change)
AlMmr[wt]	0.08	0.625
AlMmr::ClpC1[wt]	0.08 / (1)	0.625 / (1)
AlMmr::ClpC1[L88R]	0.625 / (8)	5 / (8)
AlMmr::ClpC1[F2L]	0.08 / (1)	0.625 / (1)
AlMmr::ClpC1[H77D]	0.08 / (1)	0.625 / (1)

Sensitivity of the *clpX* or *clpC1* gene knocking down Mab or Msm strains to ILE

or ILF

Knockdown of *clpX* or *clpC1* in Mab (Fig. 3a) or Msm (Fig. 3b) strains resulted in

heightened sensitivity to ILE and ILF. When induced with anhydrotetracycline (aTc) in

the absence of ilamycins, growth inhibition was observed in both the *clpX* or *clpC1*

knockdown strains of Msm and Mab, underscoring the crucial role of these proteins in

cellular proliferation. This confirms previous reports of their essentiality in Msm and

Mab[25,26].

Fig. 3 | Effect of knocking down *clpX* or *clpC1* genes on the sensitivity of Mab and Msm to ILE

or ILF. a Effect of knocking down *clpX* or *clpC1* genes on the sensitivity of Mab to ILE or ILF. **b**

Effect of knocking down *clpX* or *clpC1* genes on the sensitivity of Msm to ILE or ILF. aTc,

anhydrotetracycline, the inducer. [*], Mab strains contained different plasmids: vector, the empty

plasmid pLJR962; *clpC1,* pLJR962-clpC1 for inducible inhibition of *clpC1* gene; *clpX*, pLJR962-

clpX, for inducible inhibition of *clpX* gene. [#], Msm strains contained different plasmids: vector,

pLJR962, the empty plasmid; *clpC1,* pLJR962-clpC1, for inducible inhibition of *clpC1* gene; *clpX*,

pLJR962-clpX, for inducible inhibition of *clpX* gene.

Stablization of NTD of ClpC1 by ILE or ILF

Since all mutations occurred in the NTD of ClpC1 or ClpX in Mtb, Mab, Msm, and

Mmr, and due to their identical in NTD sequences (Supplemantary Fig. 2), we

specifically expressed and purified the NTD of ClpC1 and ClpX of Mtb, respectively.

Differential scanning fluorimetry (DSF) assay was performed to identify the
interactions of ILE or ILF with the purified MtbClpC1-NTD or MtbClpX-NTD. As
shown in Fig. 4a, ILE elevated the melting temperature (Tm) of MtbClpC1-NTD from
78.5℃ to 81.5℃, indicating a notable increase in thermal stability (ΔTm of 3℃). On
the other hand, ILF increased the thermal stability of the MtbClpC1-NTD slightly with
only 1℃ in a thermal shift (Fig. 4b) compared with ILE, which achieved a more evident
thermal shift. These results suggested the potential interactions of ILE and ILF with
MtbClpC1-NTD. We confirmed that ILE made MtbClpC1-NTD more stable and
shifted its melting temperature stronger than ILF did. However, our test results showed
that ILE and ILF did not change the dissolution temperature of MtbClpX-NTD (Data
not shown).

Fig. 4 | The melt curves of MtbClpC1-NTD with ILE and ILF. a The melt curves of MtbClpC1-
NTD with ILE. **b** The melt curves of MtbClpC1-NTD with ILF. Control, DMSO; Normalized
fluorescence intensity, the fluorescence intensity was normalized to the first time point after
exication for each well.

Inhibition of proteolytic activity of the ClpC1P1P2 by ILE but not the proteolytic
activity of ClpXP1P2 or the ATPase activity of ClpC1 or ClpX

179   Considering the superior affinity of ILE with ClpC1 over ILF according to the
above results, ILE was selected to represent ilamycins in subsequent biochemical
assays. Initial assays evaluated the effect of ILE to the ATP-dependent ATPase activity
of ClpC1 and ClpX, revealing no significant alteration in enzyme activity at varying
ILE concentrations relative to controls (Supplementary Fig. 3). Subsequent analysis of
the effect of ILE to the proteolytic activity of ClpC1P1P2 or ClpXP1P2 complex was

assessed by hydrolysis of the model substrate FITC-casein or GFP-SsrA, respectively.

ILE substantially inhibited the proteolytic capabilities of the ClpC1^{wt}P1P2 complex to

degrade casein with a significant difference as compared to the control group ($P < 0.05$

for 1, and $P < 0.0001$ for 10 and 20 µg mL^{-1} ILE groups) (Fig. 5a). Contrastingly, the

proteolytic function of multiple mutant ClpC1P1P2 complexes was unaffected at 10 µg

mL^{-1} ILE compared to the control with $P > 0.05$, while some inhibition was noted at 20

µg mL^{-1} ILE with $P < 0.01$ (Fig. 5c-5f). For the ClpXP1P2 complex, ILE did not affect

its proteolytic activity (Fig. 5b). Although ILE did not affect the ATPase activity of

ClpX and the hydrolytic activity of ClpXP1P2, it may have other ways of exerting its

antimicrobial effects through ClpX or ClpXP1P2, which need to be explored further.

Fig. 5 | Proteolytic activity of the ClpC1P1P2 and ClpXP1P2 complexes in response to ILE

treatment. a, c-f Proteolytic activity of variant ClpC1P1P2 complexes in response to ILE treatment.

b Proteolytic activity of the ClpXP1P2 complexes in response to ILE treatment. ClpC1[wt]P1P2,

ClpC1[F80I]P1P2, ClpC1[F80L]P1P2, ClpC1[F80V]P1P2, ClpC1[F80C]P1P2: complex formation of wild-type

and mutant ClpC1 with ClpP1 and ClpP2. ClpXP1P2: complex formation of wild-type ClpX with

ClpP1 and ClpP2. [*]: $P < 0.05$; [**]: $P < 0.001$; [****], $P < 0.00001$; ns, no significant difference, $P >$

0.05. Numbers after the drug names indicate the working concentrations (μg mL^{-1}). D-value of

fluorescence intensity represents the change in fluorescence intensity by subtracting the

fluorescence intensity of the first detection from the fluorescence intensity of each detection.

Extension of cell lengths and inhibition of cell division in ILE- and ILF-treated

Msm

After treating with 0.125 μg mL^{-1} of ILE or 5 μg mL^{-1} of ILF for 15 h, the cell

lengths of Msm (2.07 ± 0.48 μm) were extended to 5.95 ± 2.16 μm and 9.41 ± 2.40 μm,

respectively (Fig. 6). This morphological change indicates an inhibition of cell division,

as evidenced by the readily observable increase in division septa formation.

Fig. 6 | Cell morphology of Msm strains before or after treatment with ILE or ILF. a Cell morphology before treatment. **b** Cell morphology of 0.125 μg mL^{-1} ILE-treated Msm. **c** Cell morphology of 5 μg mL^{-1} ILF-treated Msm. White arrows indicate division septa.

Proteomic signature of the ILE-treated cells

Proteomic analysis was performed to provide insights into the adaptive responses

of Mtb during ILE treatment to better understand the mechanism of action of ILE.

Compared to untreated cells, ILE-treated Mtb displayed upregulation of proteins

involved in protein synthesis, including various 50S and 30S ribosomal proteins, except

for RplR and RpsC, which were downregulated. Regulatory sigma factors such as SigE,

SigB, and SigL, crucial for gene transcription and translation, were also affected.

Intriguingly, components of the toxin-antitoxin system, including anti-sigma factors

Rv3221A and Rv0516c, as well as toxins Rv0660c and Rv0748, were upregulated upon

ILE treatment. This observation aligns with reports that some toxin-antitoxin proteins

are targets of the ClpXP1P2 complex[14]. Correspondingly, ClpX was found to be

downregulated, potentially leading to increased levels of toxin-antitoxin proteins and

contributing to the altered cell morphology in ILE-treated Mtb.

Membrane protein expression was notably affected, with some increasing

expression while others decreasing. Proteins implicated in intermediary metabolism

and respiration, such as QcrA, QcrB, and CydD, were downregulated, hinting at the

impact of ILE on respiratory pathways. Alterations in lipid metabolism-related proteins,

including upregulation of FadE9, FadE29, and FadD24 and downregulation of FadD9,

FadE6, FadE12, and FadE24, were also observed. Additionally, many membrane

proteins (e.g., MmpL2, EccD5, EccE2) showed significant expression changes, while

others like MmpS3, SecF, and TatA were downregulated in the presence of ILE.

Discussion

This study describes ILE and ILF, two cyclic peptides with potent anti-mycobacterial properties, isolated from a deep South China Sea-derived strain, *Streptomyces atratus* SCSIO ZH16[15]. Both ILE and ILF exhibited significant activity against Msm, Mmr, and Mtb, with ILE also effective against various drug-resistant clinical Mtb isolates. This suggests that ILE targets molecular mechanisms distinct from those drugs of current TB therapeutics. Additionally, ILE showed robust antibacterial activity against Mab. In mouse macrophages, ILE also has potent bactericidal activity. Notably, neither ILE nor ILF affected certain non-mycobacterial pathogens, such as *Staphylococcus aureus* and *Acinetobacter baumannii*, indicating a mycobacteria-specific action. This specificity mirrors that of other cyclic peptides like rufomycin I, lassomycin, ecumicin, and cycolmycin A, which also selectively target mycobacteria over common Gram-negative and Gram-positive bacteria, including *Escherichia coli* (*E. coli*) and *Staphylococcus aureus*[12,27-29].

Combination studies of ILE and ILF with various anti-TB drugs, including RIF, INH, STR, EMB, and TB47 revealed synergistic or additive interactions, except for an indifferent response with ILF + RIF. These results suggest the potential of ILE and ILF to enhance the efficacy of existing anti-TB therapies. Interestingly, our proteomic analysis revealed changes in four proteins associated with the mechanism of INH after ILE treatment, potentially explaining the synergistic effect of ilamycins with INH. Specifically, SigL (fold change > 28, $P < 0.01$) and Ndh (fold change = 4.79, $P < 0.001$) were significantly upregulated, while FadE24 (fold change = 0.35, $P < 0.5$) and Glf (fold change = 0.33, $P < 0.01$) were downregulated. SigL has been reported to activate KatG, which is critical for the effect of INH[30,31], while Ndh influences the NADH/NAD$^+$ ratio, affecting INH-NAD formation[32]. FadE24 was reported to be upregulated in the prescence of INH, and mutations in it have also been found in INH-resistant Mtb[33]. Glf is responsible for catalyzing the conversion of UDP-galactopyranose to UDP-galactofuranose. Its upregulation was hypothesized to contribute to INH resistance in Mtb either by binding a modified form of INH or by

sequestering a factor such as NAD^+ required for INH activity[34]. Additionally, crucial

proteins in the electron transport chain of Mtb such as QcrA, QcrB, and CydD were all

downregulated after treatment of ILE. Given the mutual substitutability of the two

electron transfer chains , it was previously speculated that the inhibitory effect of the

QcrB inhibitor TB47 on Mtb would be enhanced when cyt-bds is inhibited[24]. Therefore,

the downregulation of these proteins may explain its synergistic effect with ilamycins.

WGS and Sanger sequencing of spontaneous mutants resistant to ILE or ILF

identified ClpC1 as the primary target in Mtb, Mab, Msm, and Mmr. The mutations in

ClpC1 associated with ILE or ILF resistance mirror those reported in mutants resistant

to other ClpC1-targeting compounds. For instance, rufomycin I-resistant mutants

exhibited V13F, H77R, F80I/V/L/S mutations[27]; lassomycin-resistant strains harbored

Q17R/H, R21S, P79T mutations[29]; cyclomarin A-resistant variants showed F2A/Y,

F80A/Y, E89A/Q alterations[12]; and ecumicin resistance was linked to L92S/F, L96P

mutations[28]. Interestingly, all of these mutations were located in the MtbClpC1-NTD

(residues 1–145). However, no mutation was observed in ClpX of rufomycin I-,

lassomycin-, cyclomarin-, and ecumicin-resistant mutants[27-29,35]. On the other hand,

ILE-resistant mutants conferred ClpX mutations and also exhibited cross-resistance to

ILF, suggesting that ilamycins may interact with multiple targets within the

mycobacterial proteolytic machinery.

Molecular docking simulations elucidated the interactions between ILE and

MtbClpC1-NTD, as well as MtbClpX-NTD. These simulations identified key residues

such as Q17, H77, F80, and L88 in MtbClpC1-NTD, and P30 in MtbClpX-NTD that

consistent with sites of mutation in resistant strains. Comparable docking studies with

cyclic peptides rufomycin, ecumicin, and cyclomarin A against MtbClpC1-NTD

corroborated that these crucial residues align with mutation sites previously identified[36].

This consistency reinforces the relevance of these residues in the binding and efficacy

of anti-mycobacterial cyclic peptides.

Comparative analysis of Ilamycin-resistant mutants and gene-edited strains

suggests that mutations at the F2 position of ClpC1 in both Mab and Msm confer

relatively low-level resistance to ilamycins. Contrastingly, P30 mutations in Mab ClpX

correlate with high-level resistance in Mab, signifying its critical role. Notably, the lack of impact of Msm $clpC1^{I197N}$ mutation on ilamycin resistance is attributed to its location outside the NTD of ClpC1, suggesting that resistance is primarily driven by alterations within the NTD. Additionally, the mutant strain harbored this mutation also harbored a mutation in F2 in ClpC1 which may mainly contribute to the resistance to ilamycins. Overexpression studies further revealed that AlMmr strains overexpressing $clpC1^{L88R}$ exhibited resistance to ILE and ILF, paralleling previous observations of mutant $clpC1$ gene overexpression conferring resistance to ClpC1-targeting compounds. For example, Msm strains expressing $clpC1^{F80Y}$ and $clpC1^{F89Y}$ mutants and *Mycobacterium bovis* BCG strains expressing $clpC1^{F2Y}$, $clpC1^{F80Y}$, and $clpC1^{E89A}$ mutants showed statistically significant resistance ($P < 0.05$) to cyclomarin A compared with parent strains[35]. Conversely, most overexpressed strains did not demonstrate drug resistance. As both proteins ClpX and ClpC1 are used to regulate protein degradation, we hypothesized that the proteins produced by overexpression are likely to be bound by some proteins within the bacteria, rendering them non-functional or directly degraded, so that most of the overexpression strains do not develop drug resistance. Moreover, knockdown of $clpX$ or $clpC1$ genes heightened sensitivity to ilamycins in Mab and Msm, also impacting strain growth, consistent with the essential nature of these genes for mycobacterial viability[25,26]. Additionally, although sgRNAs in our study intended to target the same site in $clpC1$ and $clpX$ in Mab and Msm, we observed that the effective dose of aTc in Mab is evidently higher than in Msm, which may be due to the intrinsic resistance of Mab towards tetracyclines by overexpression of a tetracycline-degrading enzyme MabTetX (MAB_1496c)[37]. Collectively, these gene-level investigations substantiate ClpX and ClpC1 as molecular targets of ILE and ILF, providing valuable insight into their mechanism of action.

DSF analyses confirmed that both ILE and ILF enhance the thermal stability of MtbClpC1-NTD with ILE showing a stronger affinity than ILF, which correlates with its lower MIC. Unlike ecumicin and lassomycin, which stimulate ClpC1 ATPase activity but impede substrate degradation, or cyclomarin A, which activates both ATPase and proteolytic functions[27-29], ILE does not alter the ATPase activities of ClpX

and ClpC1. Instead, ILE inhibits the proteolytic activity of the ClpC1P1P2 complex without affecting ATPase activities of ClpC1, suggesting a distinct mechanism that disrupts protein homeostasis and leads to bacterial cell death. Although ILE had no effect on the ATPase activity of ClpX and the hydrolytic activity of ClpXP1P2, it may have other ways of exerting its antimicrobial effects through ClpX or ClpXP1P2, which further need to be explored.

Scanning electron microscopy revealed that ILE and ILF treatments impede cell division in Msm, consistent with the phenotype of *clpX* gene silencing[26]. This is also consistent with the role of ClpX in regulating FtsZ assembly and Z-ring formation, essential for cell division in Mtb[38]. Proteomic data further indicated that ILE markedly increases toxin and antitoxin protein expression, known substrates of the ClpXP1P2 complex[14], hinting at a potential targeting of ClpX by ilamycins.

Future studies should focus on specifying the precise interactions between ilamycins, ClpX, and the ClpXP1P2 complex to fully understand the mechanism. The collective evidence from this and related studies[27-29,35] establishes ClpC1 and ClpX as promising targets for TB therapy.

Methods

Bacterial strains and media

Bacterial strains used in this study are listed in Supplementary Table 1. For Mtb, Mmr, Mab, and Msm, cells were grown in Middlebrook 7H9 broth supplemented with OADC (0.005% oleic acid, 0.5% bovine serum albumin, 0.2% dextrose, 0.085% catalase). For *E. coli*, *Enterococcus faecium*, *Staphylococcus aureus*, *Acinetobacter baumannii*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*, cells were grown in Luria-Bertani (LB) broth.

MICs against mycobacteria

The MICs of ILE and ILF were detected using a cost-effective *in vitro* assay as previously described against a selectable marker-free autoluminescent Mtb H$_{37}$Ra (AlRa), AlRv, AlMab, AlMsm and AlMmr[22-24,39]. In such cases, the MIC$_{lux}$ was defined as the lowest concentration that can inhibit > 90% relative light units (RLUs) compared with that from the untreated controls[22-24,39]. The MICs of the ILE were also determined by a well-established microplate alamar blue assay (MABA) against Mtb H$_{37}$Ra, H$_{37}$Rv and nine clinical Mtb strains isolated from pulmonary TB patients hospitalized in Guangzhou Chest Hospital, Guangzhou, China[20,31]. The MIC by MABA method was defined as the lowest concentration corresponding to the blue column on the microplate was observed by naked eye.

MICs against non-mycobacteria

The MICs against *Enterococcus faecium*, *Staphylococcus aureus*, *Acinetobacter baumannii*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* were determined by observing the turbidity after 16 h of incubation in LB broth at 37°C. The MIC was defined as the lowest concentration, resulting in clarity in turbidity relative to that of untreated control cultures[40,41].

Combined activity of ILE or ILF with other compounds *in vitro*

The combined activity of ILE or ILF with INH, RIF, EMB, STR and TB47 was tested by a checkerboard method. The fractional inhibitory concentration index (FICI)

369   was interpreted as follows: synergistic (FICI of ≤ 0.5), partially synergistic (0.5 < FICI

370   < 1.0), additive (FICI = 1.0), indifferent (1.0 < FICI ≤ 4.0), and antagonistic (FICI >

371   4.0)[42,43].

Activity in macrophage culture

The growth inhibition of AlRv in macrophage culture was assessed as previously

described[42]. The cell density of mouse macrophages RAW264.7 was adjusted to 5×10^5

cells mL^{-1}. 100 μL cells per well were added to 96-well plates and cultured overnight

for adherence. The AlRv was adjusted to 3×10^6 RLU mL^{-1}, and 100 μL AlRv was

added to each well. The plates were then incubated at 37°C for 6 h. The cells were

washed three times with DMEM to remove the AlRv that was not phagocytosed and

treated with DMEM containing amikacin (100 μg mL^{-1}) for 2 h to eliminate the

extracellular bacteria. The cells were washed twice with DMEM to remove any residual

amikacin. A total of 195 μL of DMEM containing 10% fetal bovine serum and 5 μL of

the two-fold dilutions of different ILE concentrations were added to the cells. The

mixture was incubated at 37°C with 5% CO_2 and the RLUs were measured on days 1-

5 or 7 of incubation. RIF (1 μg mL^{-1}) and dimethyl sulfoxide (DMSO) were used as

positive and negative controls, respectively.

ILE or ILF-resistant mutants

Broth cultures of AlRa, AlRv, AlMab, and AlMmr, having OD_{600} 0.6-1.3, were

plated on 7H11 plates containing 0.1, 0.5, 2.5, 5, 10 or 20 μg mL^{-1} of ILE. Broth cultures

of AlMsm (OD_{600} 0.6-1.3) were plated on 7H11 plates containing 10, 20 or 40 μg mL^{-1}

of ILF. The bacterial colonies grown on the ILE- or ILF-containing plates were picked

up to confirm the drug resistance phenotype by drug susceptibility testing in liquid

culture.

WGS analysis

The whole-genomes of the parent Mtb, Mab and a part of the confirmed ILE-

resistant mutants were sequenced by the Beijing Genomics Institute in China. The

resulting reads were aligned to the reference genome sequence and compared with that

of the parent strain. The mutations identified by WGS analysis in the drug-resistant

mutants were verified by amplifying and sequencing the *clpX* and *clpC1* genes by

Sanger sequencing. The *clpX* and *clpC1* genes of other ILE- or ILF-resistant mutants,

including Mtb, Mab, Msm, and Mmr, were amplified and detected by Sanger

sequencing (primers are listed in Supplementary Table 2).

Molecular docking analysis

To assess the binding affinities and interaction modes among ILE, MtbClpC1-

NTD, and MtbClpX-NTD, Autodock Vina 1.2.2, a computational protein–ligand

docking software, was utilized[44]. The structure of ILE was obtained from PubChem

Compound[45]. The 3D coordinates of MtbClpC1-NTD (PDB code, 6CN8; resolution,

1.4 Å) were retrieved from the Protein Data Bank (PDB), and the structure of MtbClpX-

NTD was predicted using AlphaFold2.

For the docking analysis, all protein and molecular files were converted to the

PDBQT format, excluding water molecules, and polar hydrogen atoms were added. The

grid box was centered to encompass the domain of each protein, allowing for free

molecular movement. The dimensions of the grid box were set to 30 Å × 30 Å × 30 Å,

with a grid point distance of 0.05 nm. Molecular docking studies were conducted using

Autodock Vina 1.2.2, available at http://autodock.scripps.edu/.

Overexpression of *clpX* or *clpC1* in Mab, Mmr or Msm

clpC1^{wt}, *clpC1*^{mt} (mutated *clpC1*), *clpX*^{wt} and *clpX*^{mt} from Mtb, Mab, Mmr and

Msm were amplified (primers are listed in Supplementary Table 2) and cloned into the

plasmid p60A or pMVA with high copies under the control of the *hsp60* promoter

(Supplemantary Fig. 1)[24]. The plasmids were transformed into AlMab, AlMmr, and

AlMsm. Furthermore, the *clpP1*, *clpP2* and *clpX* or *clpC1* of Mab were amplified from

the genome of Mab^{wt} and inserted into the plasmid pMVA under the control of the *hsp70*

and *hsp60* promoters for expressing ClpXP1P2 or ClpC1P1P2 simultaneously. The

MICs of the recombinant strains to ILE or ILF were determined as described above.

Gene silencing of *clpX* or *clpC1* genes in Mab or Msm

CRISPR interference was used to perform *clpX* or *clpC1* gene silencing[46]. 20-bp-

long oligonucleotides were designed as sgRNA to bind the sense strand of the coding

region of *clpX* or *clpC1*. The oligonucleotides were annealed and then ligated with linearized pLJR962[46]. Subsequently, the empty vector pLJR962 and the plasmids containing sgRNAs targeting *clpX* or *clpC1* (Supplemantary Table 3) separately were electroporated into receptive cells of Mab or Msm. The gene-silenced strains and the empty vector transformant were cultured and the gene-silencing effects were determined. 4 μL ILE was added to the 96-well microplate making the final concentration of 8 ~ 0.0313 μg mL^{-1}. The concentration of bacteria, whose OD$_{600}$ were adjusted to ~ 0.5, was 1000-fold diluted in 7H9 with or without aTc. The diluted bacteria were then divided into the experimental group (+ aTc 50 ng mL^{-1}) and the control group (- aTc). 196 μL of these two groups were then inoculated into the 96-well microplate separately. The microplates were incubated at 37°C for 3 days. The MIC was defined as the lowest concentration that prevented visible bacterial growth. The experiment was performed in triplicate and repeated twice.

Ten-fold serial dilutions of Mab^{wt} or Msm^{wt}, the empty vector transformant, and the gene-silenced strains were prepared after adjusting OD$_{600}$ to ~0.9. For Mab, 1 μL of each dilution was spotted on plates containing 0.25, 0.5, and 1 μg mL^{-1} ILE or 5, 10, and 20 μg mL^{-1} with different concentrations of aTc (0, 100, 200 and 400 ng mL^{-1}). For Msm, 1 μL of each dilution was spotted on plates containing 0.004, 0.008, and 0.01625 μg mL^{-1} ILE or 0.125, 0.25, and 0.5 μg mL^{-1} ILF with different concentrations of aTc (0, 12.5, 25 and 50 ng mL^{-1}). As a control, 1 μL of each dilution was also spotted on plates containing only aTc. The plates were incubated at 37°C for 3 days. The experiment was performed in triplicate and repeated twice.

CRISPR/Cpf1-mediated gene editing of *clpX* and *clpC1* genes in Mab and Msm

The CRISPR/Cpf1-associated recombination have been used for gene editing in Mab and Msm previously[20,21]. Firstly, the pJV53-Cpf1 was electroporated into Mab^{wt} and Msm^{wt} for the Mab::pJV53-Cpf1 and Msm::pJV53-Cpf1. Secondly, sgRNAs were designed to target 25 nucleotide sequences, which followed the sequence 5'-YTN-3' on the template strand of each target gene (Supplementary Table 3). All oligonucleotides designed for sgRNA expression were ligated into pCR-zeo. Subsequently, the vectors

were electroporated into receptive cells of induced Mab::pJV53-Cpf1 or Msm::pJV53-

Cpf1. For Cpf1 expression, aTc was added to 7H11 agar plates at a final concentration

of 200 ng mL^{-1}. These plates were cultured in 30℃ incubators for 3-5 days, and then

single colonies were selected to determine whether the site of the target gene had been

edited successfully by PCR and Sanger sequencing (primers are listed in

Supplementary Table 2). The MICs of the edited strains to ILE and ILF were tested as

described above.

Protein expression and purification

Mtb ClpP1 (residues 7-200) and ClpP2 (residues 13-214) with N-terminal His$_6$-

SUMO tags were expressed and purified. In short, ClpP1 and ClpP2 were individually

overexpressed in *E. coli* BL21 (λDE3) in LB broth at 16℃, following induction with

0.5 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) for approximately 16 h. Cells

were resuspended and lysed; His-tagged ClpP1/ClpP2 proteins were subsequently

purified via nickel affinity chromatography and eluted with the same buffer

supplemented with 250 mM imidazole. The His$_6$-SUMO tag of eluted proteins was

removed by ubiquitin-like-protease 1 at 4℃. Following the purification of the tag-free

proteins, peak fractions were collected and concentrated through anion exchange

chromatography (MonoQ; GE Healthcare). These fractions were then applied to

Superdex 200 16/600 (GE Healthcare) for final purification. ClpP1 and ClpP2 were

then concentrated in amicon centrifugal concentration devices (Millipore) to a

tetradecamer concentration of more than 80 μM.

clpX^{wt} and *clpX*^{P30H} were inserted into the pGEX-6P-1 vector, which had a His$_6$-

tag at the C-terminal and an HRV 3C protease cleavage site. After induction for 18 h at

16℃, the cells were collected by centrifugation. The pellet was then resuspended and

lysed. After centrifugation, the supernatant was loaded onto a GST column and eluted.

The eluent was digested at 4℃ for 26 h with enough HRV 3C protease. The fractions

containing the target protein were then pooled, concentrated to 1 mL, and purified with

a SuperoseTM 6 increase 10/300 GL column.

clpC1^{wt}, its mutants (F80I/V/L/C), and *gfp*-ssrA (AGKEKQNLAFAA) were

inserted into the pET-28a vector that had an N-terminal His_6-tag. The recombinant

strains were cultured in LB broth at 37°C until OD_{600} reached 0.6-0.8. After 18 h of

induction with 0.5 mM IPTG at 18°C, the pellet was suspended and lysed. Following

the loading of the supernatant onto a Ni-NTA column, the proteins were washed and

then eluted. The fractions were then pooled and loaded onto a SuperoseTM 6 increase

10/300 GL column.

The coding gene sequence of MtbClpC1-NTD (residues 1–145) and MtbClpX-

NTD (residues 1–112) were inserted into the pET-28a vector with a C-terminal His_6-

tag and an HRV 3C protease cleavage site. The expression and purification methods of

MtbClpC1-NTD and MtbClpX-NTD were consistent with the above methods.

DSF

DSF was used to detect ILE or ILF binding to the MtbClpC1-NTD and MtbClpX-

NTD based on the established methods[47][48]. 5 µM MtbClpX-NTD or MtbClpC1-NTD

was incubated with different concentration of ILE or ILF (range from 12.5 to 400 µM)

or DMSO for 0.5 h, then mixed with Sypro Orange fluorescent dye. The fluorescence

readings were observed using the CFX96 Real-time System (Bio-rad), in which the 8-

tube PCR Strips were smoothly heated from 25°C to 95°C at a constant heating rate of

0.4°C min^{-1}. The excitation and emission filters were set at $\lambda = 460$ nm and $\lambda = 510$ nm,

respectively. The data were collected with CFX Manager software and analyzed using

GraphPad Prism 8.0.2. All measurements were carried out in triplicate.

ClpC1 and ClpX ATPase activity assay

The ATPase activity of Mtb ClpC1 and ClpX was assessed via a free-phosphate

detection assay using the Biomol Green reagent. 5 µL of various concentrations of ILE

(0, 0.08, 0.16, 0.31, 0.625, 1.25, 2.5, 5 and 10 $\mu g\ mL^{-1}$) was added to each well of a

transparent 96-well plate. The ClpX or ClpC1 were diluted to 0.1 $\mu g\ mL^{-1}$ in the reaction

mixture. 90 µL ClpC1 and ClpX were added to a 96-well plate and incubated at 37°C

for 1 h. The reaction was initiated by adding 5 µL of 1 mM ATP to the reaction mixture

at 37°C for 1 h. The final volume of the reaction mixture was adjusted to 100 µL. After

the reaction, 100 µL of Biomol Green reagent was added to determine the amount of

free phosphate. After incubation at room temperature for 30 min, the resulting reaction product was measured at 620 nm using a Multimode Plate Reader (Envision, PerkinElmer, USA). The concentration of free phosphate released from ATP by ClpX and ClpC1 was calculated using a standard curve according to the concentration of free phosphate. Measurements were performed in triplicate.

Proteolytic activity assay of ClpC1P1P2 and ClpXP1P2

The proteolytic activity of ClpC1P1P2 and ClpXP1P2 were measured at 37°C in white 96-well plates using a Multimode Plate Reader (Envision, PerkinElmer, USA). ILE was added at 0, 0.1, 1, 10, and 20 μg mL^{-1} as the final concentrations in each well of a white 96-well plate. 100 μL reaction mixture was prepared as 2 μM ClpP1P2, 1 μM ClpC1 or ClpX, and 2.5 μM fluorescein isothiocyanate (FITC)-casein for ClpC1P1P2 or 2 μM GFP-SsrA for ClpXP1P2. The FITC-casein and GFP-SsrA hydrolysis was continuously monitored at 535 nm (excitation at 485 nm). Measurements were performed in triplicate.

Microscopy of ILE- or ILF-treated Msm

We treated exponentially growing Msm with 0.125 μg mL^{-1} ILE or 5 μg mL^{-1} ILF for 15 h. The bacteria immobilized by 4% glutaraldehyde were washed three times and resuspended with phosphate-buffered saline (PBS). 5 mm cell slides were soaked in bacterial suspension and placed in a 24-well plate for drying naturally. Subsequently, the cell slides were washed three times with PBS. Successive soaks with gradually more concentrated (70%, 80%, 90%, 100%) ethanol solutions were performed for 10 minutes each. Cell slides were dried using the critical point drying method and then coated with electrically-conducting material. The bacterial morphology was observed by a GeminiSEM 300 ultra-high resolution field emission cryo-scanning electron microscope.

Proteomic analysis

AlRa was treated for 5 days, in triplicate, with 0.0125 μg mL^{-1} ILE for 4D label-free phospho-proteomic analysis. The untreated cells were diluted on the same day, and the compounds were added to reach a comparable cell density between untreated and

ILE-treated cells before sample collection. The cells were resuspended and boiled for

20 min. The samples were then sent to APTBIO (Shanghai, China) for proteomics

testing and analysis.

To compare the abundance of phosphopeptides between the control and treatment

samples, label-free quantification was performed with a minimum fold change of 1.5

to determine the differentially expressed phosphopeptides. In addition, the student's *t-*

test was employed to identify significant differences between the control and treatment

samples. $P < 0.05$ were considered to indicate significance. Functional category

analysis was conducted according to Mycobrowser (https://mycobrowser.epfl.ch/).

Protein functional annotation was performed with DAVID

(https://david.ncifcrf.gov/summary.jsp).

Statistical analysis

Statistical analysis was performed with GraphPad Prism 8.0.2. The unpaired *t-test*

was applied to determine statistical significance. $P < 0.05$ was considered significant.

References

1. Bouzeyen R, Javid B. Therapeutic vaccines for tuberculosis: An overview. *Front Immunol* **13**, 878471 (2022).

2. Miotto P, Zhang Y, Cirillo DM, Yam WC. Drug resistance mechanisms and drug susceptibility testing for tuberculosis. *Respirology* **23**, 1098-1113 (2018).

3. Pfyffer G. Drug-resistant tuberculosis: resistance mechanisms and rapid susceptibility testing. *Schweiz med Wochenschr* **130**, 1909-1913 (2000).

4. Fedrizzi T, *et al.* Genomic characterization of nontuberculous mycobacteria. *Sci rep* **7**, 45258 (2017).

5. Henkle E, Winthrop KL. Nontuberculous mycobacteria infections in immunosuppressed hosts. *Clin chest med* **36**, 91-99 (2015).

6. Fraga H, *et al.* Development of high throughput screening methods for inhibitors of ClpC1P1P2 from *Mycobacteria tuberculosis*. *Anal Biochem* **567**, 30-37 (2019).

7. Lupoli TJ, Vaubourgeix J, Burns-Huang K, Gold B. Targeting the proteostasis network for mycobacterial drug discovery. *ACS infect dis* **4**, 478-498 (2018).

8. Lee H, Suh JW. Anti-tuberculosis lead molecules from natural products targeting *Mycobacterium tuberculosis* ClpC1. *J Ind Microbiol Biotechnol* **43**, 205-212 (2016).

9. d'Andrea FB, Poulton NC, Froom R, Tam K, Campbell EA, Rock JM. The essential *M. tuberculosis* Clp protease is functionally asymmetric *in vivo*. *Sci Adv* **8**, eabn7943 (2022).

10. Raju RM, *et al. Mycobacterium tuberculosis* ClpP1 and ClpP2 function together in protein degradation and are required for viability *in vitro* and during infection. *PLoS pathog* **8**, e1002511 (2012).

11. Akopian T, Kandror O, Raju RM, UnniKrishnan M, Rubin EJ, Goldberg AL. The active ClpP protease from *M. tuberculosis* is a complex composed of a heptameric ClpP1 and a ClpP2 ring. *EMBO J* **31**, 1529-1541 (2012).

12. Schmitt EK, *et al.* The natural product cyclomarin kills *Mycobacterium*

tuberculosis by targeting the ClpC1 subunit of the caseinolytic protease. *Angew*

Chem Int Ed Engl **50**, 5889-5891 (2011).

13. Sassetti CM, Boyd DH, Rubin EJ. Genes required for mycobacterial growth

defined by high density mutagenesis. *Mol Microbiol* **48**, 77-84 (2003).

14. Texier P*, et al.* ClpXP-mediated degradation of the TAC antitoxin is neutralized

by the SecB-like chaperone in *Mycobacterium tuberculosis*. *J Mol Biol* **433**,

166815 (2021).

15. Ma J*, et al.* Biosynthesis of ilamycins featuring unusual building blocks and

engineered production of enhanced anti-tuberculosis agents. *Nat Commun* **8**, 391

(2017).

16. Gutiérrez AV, Viljoen A, Ghigo E, Herrmann J-L, Kremer L. Glycopeptidolipids,

a double-edged sword of the *Mycobacterium abscessus* complex. *Front*

microbiol **9**, 1145 (2018).

17. Jankute M, Grover S, Birch HL, Besra GS. Genetics of mycobacterial

arabinogalactan and lipoarabinomannan assembly. *Molecular Genetics of*

Mycobacteria, 535-557 (2014).

18. Lu XY*, et al.* Pyrazolo[1,5-]pyridine inhibitor of the respiratory cytochrome

complex for the treatment of drug-resistant tuberculosis. *Acs Infect Dis* **5**, 239-

249 (2019).

19. Bergval I*, et al.* Pre-existing isoniazid resistance, but not the genotype of

Mycobacterium tuberculosis drives rifampicin resistance codon preference *in*

vitro. *PLoS One* **7**, e29108 (2012).

20. Wang S*, et al.* Amino acid 17 in QRDR of gyrase A plays a key role in

fluoroquinolones susceptibility in mycobacteria. *Microbiol Spectr* **11**, e0280923

(2023).

21. Yan M-Y, Yan H-Q, Ren G-X, Zhao J-P, Guo X-P, Sun Y-C. CRISPR-Cas12a-

Assisted Recombineering in Bacteria. *Appl Environ Microbiol* **83**, (2017).

22. Liu Y*, et al.* Assessment of clofazimine and TB47 combination activity against

Mycobacterium abscessus using a bioluminescent approach. *Antimicrob Agents*

Chemother **64**, (2020).

23. Yang F*, et al.* Engineering more stable, selectable marker-free autoluminescent

mycobacteria by one step. *PloS One* **10**, e0119341 (2015).

24. Liu Y*, et al.* The compound TB47 is highly bactericidal against *Mycobacterium*

ulcerans in a Buruli ulcer mouse model. *Nat Commun* **10**, 524 (2019).

25. Rifat D, Chen L, Kreiswirth BN, Nuermberger EL. Genome-wide essentiality

analysis of *Mycobacterium abscessus* by saturated transposon mutagenesis and

deep sequencing. *mBio* **12**, e0104921 (2021).

26. Bai J, Chi M, Hu Y, Hao M, Zhang X. Construction and biological characteristics

of ClpC and ClpX knock-down strains in *Mycobacterium smegmatis*. *Chin. J.*

Biotechnol **41**, 13-22 (2021).

27. Choules MP*, et al.* Rufomycin targets ClpC1 proteolysis in *Mycobacterium*

tuberculosis and *M. abscessus*. *Antimicrob Agents Chemother* **63**, e02204-18

(2019).

28. Gao W*, et al.* The cyclic peptide ecumicin targeting ClpC1 is active against

Mycobacterium tuberculosis in vivo. *Antimicrob Agents Chemother* **59**, 880-889

(2015).

29. Gavrish E*, et al.* Lassomycin, a ribosomally synthesized cyclic peptide, kills

Mycobacterium tuberculosis by targeting the ATP-dependent protease

ClpC1P1P2. *Chem biol* **21**, 509-518 (2014).

30. Sklar JG, Makinoshima H, Schneider JS, Glickman MS. *M. tuberculosis*

intramembrane protease Rip1 controls transcription through three anti-sigma

factor substrates. *Mol microbiol* **77**, 605-617 (2010).

31. Hameed HA*, et al.* Molecular targets related drug resistance mechanisms in

MDR-, XDR-, and TDR-*Mycobacterium tuberculosis* strains. *Front cell infect*

microbiol **8**, 114 (2018).

32. Unissa AN, Subbian S, Hanna LE, Selvakumar N. Overview on mechanisms of

isoniazid action and resistance in *Mycobacterium tuberculosis*. *Infect Genet Evol*

45, 474-492 (2016).

33. Ramaswamy SV*, et al.* Single nucleotide polymorphisms in genes associated

with isoniazid resistance in *Mycobacterium tuberculosis*. *Antimicrob Agents*

Chemother **47**, 1241-1250 (2003).

34.   Chen P, Bishai WR. Novel selection for isoniazid (INH) resistance genes

supports a role for NAD+-binding proteins in mycobacterial INH resistance.

Infect Immun **66**, 5099-5106 (1998).

35.   Vasudevan D, Rao SP, Noble CG. Structural basis of mycobacterial inhibition by

cyclomarin A. *J Biol Chem* **288**, 30883-30891 (2013).

36.   Wolf NM*, et al.* High-resolution structure of ClpC1-rufomycin and ligand

binding studies provide a framework to design and optimize anti-tuberculosis

leads. *ACS Infect Dis* **5**, 829-840 (2019).

37.   Rudra P, Hurst-Hess K, Lappierre P, Ghosh P. High levels of intrinsic tetracycline

resistance in *Mycobacterium abscessus* are conferred by a tetracycline-modifying

monooxygenase. *Antimicrob Agents Chemother* **62**, e00119-18 (2018).

38.   Dziedzic R*, et al. Mycobacterium tuberculosi*s ClpX interacts with FtsZ and

interferes with FtsZ assembly. *PloS One* **5**, e11058 (2010).

39.   Zhang T, Li S-Y, Nuermberger EL. Autoluminescent *Mycobacterium*

tuberculosis for rapid, real-time, non-invasive assessment of drug and vaccine

efficacy. *PloS One* **7**, e29774 (2012).

40.   Tian X*, et al.* Rapid visualized assessment of drug efficacy in live mice with a

selectable marker-free autoluminescent *Klebsiella pneumoniae. Biosens*

Bioelectron **177**, 112919 (2021).

41.   Wang S*, et al.* Rapid, serial, non-invasive quantification of *Pseudomonas*

aeruginosa in live mice with a selectable marker-free autoluminescent strain.

Biosens Bioelectron **165**, 112396 (2020).

42.   Yu W*, et al.* TB47 and clofazimine form a highly synergistic sterilizing block in

a second-line regimen for tuberculosis in mice. *Biomed pharmacother* **131**,

110782 (2020).

43.   Liu P*, et al.* Design and synthesis of novel pyrimidine derivatives as potent

antitubercular agents. *Eur J Med Chem* **163**, 169-182 (2019).

44.   Morris GM, Huey R, Olson AJ. Using autodock for ligand-receptor docking.

Curr protoc bioinformatics **24**, 8.14. 11-18.14. 40 (2008).

45. Wang Y*, et al.* Pubchem bioassay: 2017 update. *Nucleic Acids Res* **45**, D955-D963 (2017).

46. Rock JM*, et al.* Programmable transcriptional repression in mycobacteria using an orthogonal CRISPR interference platform. *Nat microbiol* **2**, 1-9 (2017).

47. Vivoli M, Novak HR, Littlechild JA, Harmer NJ. Determination of protein-ligand interactions using differential scanning fluorimetry. *J Vis Exp*, e51809 (2014).

48. Khadiullina R, Mirgayazova R, Davletshin D, Khusainova E, Chasov V, Bulatov E. Assessment of thermal stability of mutant p53 proteins via differential scanning fluorimetry. *Life (basel)* **13**, 31 (2022).

Acknowledgments

We acknowledge the Yicheng Sun group from the Institute of Pathogenic Biology, Chinese Academy of Medical Sciences for kindly sending the pJV53-Cpf1 and pCR-Zeo plasmids as the tool of gene editing. This work was supported by the National Key R&D Program of China (2021YFA1300900), the National Natural Science Foundation of China (81973372, 32300152, 82022067, 22037006), China Postdoctoral Science Foundation (2022M723164), Guangdong Provincial Basic and Applied Basic Research Fund (2022A1515110505, 2019B030302004), the grant of State Key Lab of Respiratory Disease (SKLRD-Z-202414, SKLRD-Z-202301), Open Program of Shen Zhen Bay Laboratory (SZBL2021080601006) and Nansha District Science and Technology Plan Project (NSJL202102). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contributions

T.Z., J.M, and X.X. supervised the project; T.Z., Y.G., and C.F. designed the whole project and most of the experiments; J.M. and C.S. prepared and characterized the ilamycin E and ilamycin F; Y.G. and C.F. performed the susceptibility testing and resistant mutant screening; B.Z. and J.J. contributed to molecular docking simulation; C.F., X.T. and J.H. constructed overexpression and gene-editing strains and peformed

704 the susceptibility testing; C.F. and X.H. performed the gene-scilencing experiments;

705 Y.G. and B.Z. contributed to protein expression and mode of action experiments; C.F.

706 and H.Z. contributed to the microscopy results; Y.G. performed the statistical analysis;

707 Y.G., F.C., and T.Z. wrote the manuscript; H.M.A., J.M., X.C., N.Z., X.X., J.J., and T.Z.

708 revised the manuscript.

Competing interests

710 The authors have no competing interests to declare that are relevant to the content of

711 this article.