Mutations in ClpC1 or ClpX subunit of the caseinolytic protease confer resistance to natural product ilamycins in mycobacteria

Yamin Gao[1,2,3#], Cuiting Fang[1,2,3#], Biao Zhou[1,4,5], H.M. Adnan Hameed[1,2,3], Changli Sun[3,6], Xirong Tian[1,2,3], Jing He[1,2,7], Xingli Han[1,2,3], Han Zhang[1,2,8], Jun Li[9], Jianhua Ju[3,6], Xinwen Chen[5], Nanshan Zhong[5], Junying Ma[3,6*], Xiaoli Xiong[1,2,3*], Tianyu Zhang[1,2,3*]

[1] State Key Laboratory of Respiratory Disease, Institute of Drug Discovery, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China
[2] Guangdong-HongKong-Macau Joint Laboratory of Infectious Respiratory Diseases, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China
[3] University of Chinese Academy of Sciences, Beijing 100049, China
[4] Guangzhou Medical University, Guangzhou 510180, China
[5] Guangzhou National Laboratory, Guangzhou 510005, China
[6] CAS Key Laboratory of Tropical Marine Bio-Resources and Ecology, Guangdong Key Laboratory of Marine Materia Medica, RNAM Center for Marine Microbiology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China
[7] Institute of Physical Science and Information Technology, Anhui University, Hefei 230601, China
[8] School of Life Sciences, University of Science and Technology of China, Hefei 230026, China
[9] Shanghai Institute for Advanced Immunochemical Studies and School of Life Science and Technology, ShanghaiTech University, Shanghai 201210, China

[#] These authors contributed equally to this work.
[*] Corresponding authors: Tianyu Zhang (Tel: +86-20-32015270, E-mail: zhang_tianyu@gibh.ac.cn); Xiaoli Xiong (Tel: +86-20-32015327, E-mail: xiong_xiaoli@gibh.ac.cn); Junying Ma (Tel: +86-20-34066449, E-mail: majunying@scsio.ac.cn).

Abstract

The mycobacterial caseinolytic protease (Clp) system has been recognized as a

promising therapeutic target. In this study, we identified two novel ilamycin analogs,

ilamycin E (ILE) and ilamycin F (ILF), both targeting the ClpC1 component of the

ClpC1P1P2 proteasome. ILE potently disrupts ClpC1P1P2-mediated proteolysis,

leading to delayed bactericidal activity, while ILF also binds ClpC1, albeit with lower

affinity. Notably, we discovered a unique mutation in *clpX* and a novel insertion in

clpC1, both conferring resistance to ILE and ILF in mycobacterium, validated by gene

editing. Furthermore, ILE can also inhibit the proteolytic activity of ClpXP1P2 in a

manner dependent on the substrate's tag sequence and adaptor. This first

demonstration of *clpX*- and *clpC1*-mediated ilamycin resistance underscores the

potential of ilamycins to target multiple components of the Clp protease system,

offering a novel dual-target strategy for combating mycobacterial infections.

Introduction

Tuberculosis (TB) caused by *Mycobactrium tuberculosis* (Mtb) continues to pose a significant global public health threat, with the situation being exacerbated by the emergence of drug-resistant Mtb strains[1,2]. Additionally, infections caused by nontuberculous mycobacteria (NTM), such as *Mycobacterium abscessus* (Mab), are on the rise due to intrinsic resistance to almost all the existing antibiotics. In several countries, the incidence of NTM infections has surpassed that of TB[3,4]. This highlights an urgent need for new drugs with novel mechanisms of action to manage drug-resistant mycobacterial diseases effectively.

The caseinolytic protease (Clp) proteolytic system has recently emerged as a promising therapeutic target in mycobacteria[5-8]. This system is pivotal not only for maintaining intracellular protein quality but also for modulating responses to environmental stressors and contributing to the pathogenicity of virulent strains[6,9]. It comprises multi-subunit protein complexes that facilitate intracellular protein degradation. In mycobacteria, the core complex of the hetero-tetradecamer Clp protease assumes a barrel-shaped structure, constructed from two rings, each composed of seven ClpP1 and seven ClpP2 peptidase subunits. The entry of substrates into the ClpP1P2 complex is stringently regulated and reliant on ATP-dependent AAA+ unfoldases (ClpX or ClpC1), which act as adaptors to facilitate substrate entry into the protease chamber through ATP hydrolysis[8,10,11]. Both the Clp protease genes and the ATPase adaptor genes *clpX* and *clpC1* have been found essential for mycobacterial survival[8,12].

Many non-ribosomal natural cyclic peptides, such as ilamycins (also known as rufomycins), cyclomarin A (CYMA), and ecumicin (ECU), have shown antitubercular activities by targeting the ClpC1 component of the ClpC1P1P2 proteasome[13]. However, aside from the mutations identified in the *clpC1* gene from spontaneous resistant mutants, robust genetic evidence is lacking. Although these compounds all disrupt protein metabolism through the Clp system, their mechanisms of action (MOAs) differ. For example, rufomycin □ (RUFI) inhibits the proteolytic activity of

the ClpP1P2 complexes by binding to ClpC1, interfering with its binding to ClpP1P2,

while CYMA stimulates ClpC1's ATPase and promotes proteolysis by the ClpC1P1P2

proteasome[14,15]. On the other hand, ECU stimulates the ATPase activity of ClpC1 but

inhibit the proteolysis by ClpC1ClpP1P2[16]. To the best of our knowledge, there are no

reported anti-Mtb compounds targeting ClpX.

We previously reported that ilamycin E (ILE) and ilamycin F (ILF), synthesized

by *Streptomyces atratus* SCSIO ZH16 isolated from the South China Sea, exhibited

activities against Mtb[17]. Particularly, ILE demonstrated superior activity against Mtb

compared to other ilamycin components and the aforementioned compounds[13,17,18].

However, their MOAs, activities against NTM and other non-mycobacterial

pathogenic bacteria, and potential drug interactions remain to be elucidated. In this

study, we discovered that ilamycins target ClpC1 and inhibit the proteolytic activity of

the ClpC1P1P2 protease. Additionally, ILE can also impair the proteolytic activity of

the ClpXP1P2 protease depending on the tag sequences of the substrate and adaptor-

mediated recognition. These findings highlight the potential of Clp complexes as

viable therapeutic targets, particularly when employing a dual-targeting approach.

Results

ILE and ILF exhibit selective and potent antimycobacterial activities *in vitro* and *ex vivo*

Mtb and 3 different NTM (Mab, *Mycobacterium marinum* (Mmr), *Mycobacterium smegmatis* (Msm) were selected to investigate the activities of ILE and ILF. These mycobacteria were chosen due to their clinical relevance and growth characteristics. Mab, a rapidly growing mycobacterium, is frequently associated with severe human infections and is known for its intrinsic antibiotic-resistant nature[19,20]. Mmr, a slowly growing mycobacterium, is closely related to Mtb but less virulent and serves as a model organism for Mtb studies[21,22]. Msm is also utilized widely as a model organism due to its rapid growth, non-virulence and the availability of genetic tools[22].

ILE and ILF demonstrated significant inhibitory activities against autoluminescent Mtb $H_{37}Ra$ (AlRa)[23], Msm (AlMsm)[23] and Mmr (AlMmr)[24] with varying potencies (Table 1). Notably, ILE exhibited potent activity against autoluminescent Mab (AlMab)[25] at a concentration of 1 µg mL^{-1}, whereas ILF only showed weak activity at 20 µg mL^{-1}, though they both showed highly bactericidal activities against Mtb. This disparity in efficacy may stem from ILE and ILF structural differences and the unique cell wall permeabilities of each mycobacterial species[26,27]. Compared to RIF, ILE and ILF exhibited delayed antimicrobial activities, initially displaying minor bacteriostatic effects on AlRa, with bactericidal activity emerging after 6 days of incubation (Fig. 1). Similar killing dynamics were observed for AlMsm, AlMab, and AlMmr (Supplementary Fig. 1). Consistent bactericidal activity of ILE was also confirmed in autoluminescent $H_{37}Rv$ (AlRv) (Supplementary Fig. 2), addressing concerns regarding the slower growth of $H_{37}Ra$[28]. The classical colony-forming unit (CFU) enumeration were also performed to evaluate the antimicrobial activities of ILE and ILF. Consistent with the luminescence inhibition data, ILE and ILF exhibited concentration- and time-dependent bactericidal activities against AlRa, as demonstrated by the substantial reduction in CFU counts after 15

days of incubation (Fig. 1). Specifically, based on CFU enumeration in AlRa, the minimal bactericidal concentration (MBC) was determined to be 0.05 µg mL^{-1} for ILE and 1.25 µg mL^{-1} for ILF.

The minimal inhibitory concentrations (MICs) of ILE against clinically isolated drug-resistant Mtb strains ranged from 0.01 to 0.05 µg mL^{-1}, which aligns with the laboratory standard Mtb H$_{37}$Rv strain (Table 1). Synergistic effects were observed when combining ILE with rifampicin (RIF) or the novel anti-TB drug TB47 or combining ILF with isoniazid (INH) or TB47[29]. Partial synergy was noted when ILE was paired with INH, ethambutol (EMB), clofazimine (CFZ), or bedaquiline (BDQ), and when ILF was combined with CFZ. Additive effects were observed when ILE was combined with streptomycin (STR), or ILF combined with EMB, STR or BDQ. No antagonism was detected between any of the selected drugs and either ILE or ILF. In mouse RAW264.7 macrophages, ILE effectively inhibited AlRv[30] with an MIC of 0.2 µg mL^{-1}, while ILF showed activity at 5 µg mL^{-1}. Cytotoxicity testing of ILE in RAW264.7 cells revealed a 50% toxic concentration (TC$_{50}$) of 28.24 µg mL^{-1}, suggesting a favorable selectivity index (TC$_{50}$/MIC = 141.2) relative to its anti-Mtb activity. However, neither ILE nor ILF exhibited activity against common non-mycobacterial clinical isolates, with MICs exceeding 128 µg mL^{-1} (Table 1).

Table 1 | Antibacterial spectrum of ILE and ILF

Strains or drug combinations	ILE MIC (µg mL^{-1})	ILF MIC (µg mL^{-1})	FICI
AlRa	0.05[†]	1.25[†]	-
AlRv	0.05[†]	ND	-
AlMsm	0.0625[†]	2.5[†]	-
AlMab	1[†]	20[†]	-
AlMmr	0.08[†]	0.625[†]	-
H$_{37}$Rv	0.025[‡]	2.5[‡]	-
K11[*]	0.025-0.05[‡]	ND	-
K12[*]	0.025[‡]	ND	-
K13[*]	0.01[‡]	ND	-
K14[*]	0.01[‡]	ND	-
K16[*]	0.05[‡]	ND	-
K17[*]	0.01[‡]	ND	-
K18[*]	0.025-0.05[‡]	ND	-
K19[*]	0.01[‡]	ND	-
K20[*]	0.05[‡]	ND	-

Staphylococcus aureus	> 128	> 128	-
Acinetobacter baumannii	> 128	> 128	-
Enterococcus faecium	> 128	> 128	-
Klebsiella pneumoniae	> 128	> 128	-
Pseudomonas aeruginosa	> 128	> 128	-
ILE + RIF	-	-	0.3125[†]
ILE + INH	-	-	0.625[†]
ILE + EMB	-	-	0.5625[†]
ILE + STR	-	-	1[†]
ILE + CFZ	-	-	1[†]
ILE + BDQ	-	-	0.62[†]
ILE + TB47	-	-	0.3125[†]
ILF + RIF	-	-	1.0625[†]
ILF + INH	-	-	0.3125[†]
ILF + EMB	-	-	1[†]
ILF + STR	-	-	1[†]
ILF + CFZ	-	-	0.63[†]
ILF + BDQ	-	-	1[†]
ILF + TB47	-	-	0.3125[†]

[*]Clinical resistant Mtb; [†]MICs obtained by detection of relative light units; [‡]MICs obtained by the microplate alamar blue assay; -Not applicable; ND, not detected ; FICI, fractional inhibitory concentration index.

Fig 1 | ILE and ILF exhibit potent activity against AlRa. Chemical structures, time-kill curves based on relative light units (RLUs), and CFU counts at day 15 against AlRa of ILE **(a)** and ILF **(b)**. DMSO, solvent control; RIF, positive control. Numbers following the drug names indicate the working concentrations (µg mL^{-1}).

114   The MIC_{lux} was defined as the lowest concentration that can inhibit > 90% RLUs

115   compared with that from the untreated controls. RLU and CFU data were obtained

116   from independent experiments. Statistical significance was determined using one-way

117   ANOVA. Data are presented as mean ± SD from three biological replicates (n = 3).

118   ****, $P < 0.0001$.

Mutations in *clpC1* and *clpX* were identified in ILE/ILF-resistant mutants

To identify target(s) of ILE and ILF, spontaneous mutants of different

mycobacterium species were screened. The spontaneous mutation rate for producing

ILE-resistant Mtb was approximately 2.5×10^{-9}, which is similar to that of RUFI and

lower than that of ECU and lassomycin, as well as RIF[15,16,18,31].

Whole genome sequencing (WGS) of 7 ILE-resistant Mtb strains from 4

independent screens revealed no mutations in ClpC1 or its flanking regions, but 5

strains carried a ClpX P30H mutation. Further Sanger sequencing of 17 ILE-resistant

Mtb strains identified ClpX (P30H) mutation in 8 strains (47%) and ClpC1 (F80V/C/I)

mutations in 11 strains (61%), with 2 strains harboring both mutations (ClpX P30H

and ClpC1 F80V/I) (Table 2). Similarly, WGS of 8 ILE-resistant Mab strains revealed

4 strains with ClpC1 mutations (Q17R, F2L, H77D), one with ClpX P30H, and three

without mutations in either gene. Sanger sequencing of the *clpX* and *clpC1* genes of

other ILE-resistant Mab strains revealed that there were also ClpC1 (M1-FERF-F2;

L88R) mutations in Mab strains (Table 2). Besides, a small subset of ILE-resistant

Mtb or Mab strains did not carry mutations in neither *clpX* nor *clpC1*, suggesting the

potential involvement of alternative resistance mechanisms. In these strains, WGS

identified sporadic mutations in genes unrelated to the Clp proteases, such as *Rv1572*,

MAB_2872c (*hisS*), or *MAB_2670c* (*hisD*) (Supplementary Table 1 and

Supplementary Table 2), though their relevance to resistance remains to be validated.

However, all examined ILE-resistant Mmr strains and ILF-resistant Msm strains

carried mutations exclusively in ClpC1 without any mutations in ClpX (Table 2).

Cross-resistance between ILE and ILF was observed in all ILE- or ILF-resistant NTM

mutants. These findings suggest that ILE and ILF may both target the ClpX and

ClpC1, especially in Mtb and Mab.

Table 2 | Summary of mutations of ClpC1 and ClpX in ILE- and ILF-resistant

mutants

Strains No.[a]	MICs (µg mL^{-1})		Mutation			
	ILE	ILF	ClpC1	*clpC1*	ClpX	*clpX*
Mtb^{R1*}	0.625	-	-	-	P30H	CCC→CAC
Mtb^{R2*}	20	-	F80I	TTT→ATT	P30H	CCC→CAC
Mtb^{R3*}	20	-	F80V	TTT→GTT	P30H	CCC→CAC
Mtb^{R4*}	0.1	-	F80I	TTT→ATT	-	-
Mtb^{R5*}	0.1	-	F80V	TTT→GTT	-	-
Mtb^{R6*}	0.1	-	F80L	TTT→CTT	-	-
Mtb^{R7*}	20	-	F80C	TTT→TGT	-	-
Mab^{R1*}	> 128	> 128	-	-	P30H	CCC→CAC
Mab^{R2*}	4	> 128	F2L	TTC→TCC	-	-
Mab^{R3*}	4	> 128	M1-FERF-F2	Ins: TTCGAGAGATTC	-	-
Mab^{R4*}	4	> 128	Q17R	CAA→CGA	-	-
Mab^{R5*}	> 128	> 128	H77D	CAC→GAC	-	-
Mab^{R6*}	> 128	> 128	L88R	CTG→CGG	-	-
Mmr^{R1*}	1	4	F2L	TTC→TCC	-	-
Mmr^{R2*}	10	40	H77D	CAC→GAC	-	-
Mmr^{R3*}	10	40	L88R	CTC→CGC	-	-
Msm^{R1†}	0.5	20	F2S	TTT→TCT	-	-
Msm^{R2†}	0.25	10	F2C	TTT→TGT	-	-
Msm^{R3†}	1	20	M1-L-F2	Ins: GTT	-	-
Msm^{R4†}	1	10	F5I	TTT→ATT	-	-
Msm^{R5†}	0.5	40	F80S	TTC→TCC	-	-
Msm^{R6†}	0.25	10	L88P	CTC→CCC	-	-
Msm^{R7†}	0.25	20	L88R	CTC→CGC	-	-
Msm^{R8†}	0.25	10	F2S; I197N	TTT→TCT; ATC→AAC	-	-

[a]Resistant strains were renumbered; *Resistant mutants selected from ILE-containing plates; †Resistant mutants selected from ILF-containing plates; Ins, insertion; Underlines indicate mutated bases; - showed no mutation.

Docking predicts ILE and ILF bind the N-terminal domain (NTD) of ClpC1 and

ClpX

To investigate the interactions of ILE and ILF with ClpC1 and ClpX, we

performed molecular docking simulations. Since all mutation sites of ILE and ILF-

resistant bacteria are located in ClpC1-NTD (residues 1–145) or ClpX-NTD (residues

1–112), we employed AutoDock to dock ILE into the reported structure of the

MtbClpC1-NTD (Fig. 2a)[32-34] and the AlphaFold2-predicted structure of the

MtbClpX-NTD (Fig. 2b), respectively. The docking results indicated that ILE binds to

MtbClpC1-NTD and MtbClpX-NTD with predicted binding energies of –7.9 kcal

mol^{-1} and –6.7 kcal mol^{-1}, and corresponding root mean square deviations of 3.5 Å

and 2.1 Å, respectively, confirming acceptable docking accuracy. Key residues

involved in ILE binding include the known mutation sites Q17, H77, F80, and L88 in

MtbClpC1-NTD, and P30 in MtbClpX-NTD (Table 2). Given the structural similarity

between ILE and ILF, along with the observed cross-resistance, it is likely that ILF

interacts with ClpC1 and ClpX at similar binding sites. Notably, H77 and F80 are also

critical for CYMA and RUFI binding, while RUFI additionally involves M1, F2, V13,

and E89, and ECU interacts with L92 and L96, indicating a partially conserved

binding interface across cyclopeptides[32].

Fig 2 | The graphical representation of crucial residues in the interaction between

ILE and MtbClpC1-NTD as well as MtbClpX-NTD. a Crucial residues in the

interaction between ILE and MtbClpC1-NTD. **b** Crucial residues in the interaction

between ILE and MtbClpX-NTD. MtbClpC1-NTD, N-terminal domain of MtbClpC1

(residues 1–145); MtbClpX-NTD, N-terminal domain of MtbClpX (residues 1–112).

Mutations in *clpC1* and *clpX* confer resistance to ILE and ILF in mycobacteria

To validate the significance of ClpC1 and ClpX in relation to ILE and ILF, we

initially constructed a series of overexpression strains. The wild-type (wt) genes

clpX^{wt} and *clpC1*^{wt}, along with their mutant (mt) variants *clpX*^{mt} and *clpC1*^{mt}, were

inserted into two extrachromosomal plasmids, p60A or pMVA (Supplementary Fig.

3)[35]. These genes were under the robust *hsp60* promoter to facilitate overexpression in

AlMab, AlMsm, and AlMmr[23-25]. Among these strains, only the AlMmr strain

overexpressing ClpC1^{L88R} demonstrated resistance to both ILE and ILF when

compared to the parent strain, exhibiting an 8-fold increase in MICs (Supplementary

Table 3). However, the overexpression of ClpC1$^{wt/mt}$ or ClpX$^{wt/mt}$, as well as the

concurrent overexpression of genes encoding the ClpC1P1P2 or ClpXP1P2 proteases

in AlMab and AlMsm, did not alter their susceptibility to ILE or ILF. These findings

are consistent with previous observations, suggesting that the overexpression of all

mutated *clpC1* does not necessarily confer resistance to ClpC1-targeting compounds,

and that efficacy varies across different mycobacterial species[36].

In order to further determine the roles of mutations in *clpC1* and *clpX* in ILE and

ILF resistance directly, we aimed to obtain gene-edited strains. Given the slower

growth rate of Mtb compared to rapidly-growing mycobacteria such as Msm and Mab,

we initially utilized these two NTM[37]. Utilizing the newly developed CRISPR/Cpf1-

mediated gene editing techniques, we introduced specific *clpC1* and *clpX* mutations

into Mab and Msm, resulting in two Mab mutants (ClpC1^{wt} to ClpC1^{F2L}; ClpX^{wt} to

ClpX^{P30H}) and three Msm mutants (ClpC1^{wt} to ClpC1^{F2C}, ClpC1^{wt} to ClpC1$^{M1-L-F2}$ and

ClpC1^{wt} to ClpC1^{I197N})[38,39]. These genetically modified strains displayed varying

levels of resistance as summarized in Table 3. Notably, the ClpX^{P30H} mutation in Mab

conferred the highest level of resistance, with MICs exceeding 128 µg mL^{-1} for both

ILE and ILF. The ClpC1^{F2L} mutation in Mab led to a more modest increase in

resistance, elevating the MIC for ILE by eight times relative to the parent strain. The

mutations ClpC1^{F2C} and ClpC1$^{M1-L-F2}$ in Msm both resulted in a greater than 4-fold

increase in the MICs of ILE and ILF, compared to the parent strain. A comparative

analysis of spontaneous mutants and gene-edited strains revealed that mutations at the

F2 of ClpC1 in both Mab and Msm confer relatively low-level resistance to ILE and

ILF, whereas the P30H mutation in Mab ClpX is associated with high-level resistance,

underscoring its crucial role. It is noteworthy that the lack of impact of the Msm

ClpC1^{I197N} mutation on ILE and ILF resistance may be attributed to its location

outside ClpC1-NTD, suggesting that resistance is primarily driven by alterations

within the NTD. Furthermore, the spontaneous mutant strain harboring $clpC1^{I197N}$ also

possessed $clpC1^{F2S}$ (Table 2) which may be the principal factor contributing to ILE

and ILF resistance.

Table 3 | MICs of ILE and ILF against gene edited strains and corresponding

spontaneous mutants

Strains	Amino acid change	MICs (µg mL^{-1}) / (Fold change)	
		ILE	ILF
Mab^{wt}	/	1	20
Mab^{R2*}	ClpC1^{wt} to ClpC1^{F2L}	4 / (4)	> 128 / (> 6)
Mab-C1^{F2L}	ClpC1^{wt} to ClpC1^{F2L}	8 / (8)	> 128 / (> 6)
Mab^{R1*}	ClpX^{wt} to ClpX^{P30H}	> 128 / (> 128)	> 128 / (> 6)
Mab-X^{P30H}	ClpX^{wt} to ClpX^{P30H}	> 128 / (> 128)	> 128 / (> 6)
Msm^{wt}	/	0.0625	2.5
Msm^{R2†}	ClpC1^{wt} to ClpC1^{F2C}	0.25 / (4)	10 / (4)
Msm-C1^{F2C}	ClpC1^{wt} to ClpC1^{F2C}	0.5 / (8)	20 / (8)
Msm^{R3†}	ClpC1^{wt} to ClpC1$^{M1-L-F2}$	1 / (16)	20 / (8)
Msm-C1$^{M1-L-F2}$	ClpC1^{wt} to ClpC1$^{M1-L-F2}$	0.25 / (4)	16 / (6.4)
Msm^{R8†}	ClpC1^{wt} to ClpC1^{F2S} ClpC1^{wt} to ClpC1^{I197N}	0.25 / (4)	10 / (4)
Msm-C1^{I197N}	ClpC1^{wt} to ClpC1^{I197N}	0.0625 / (1)	2.5 / (1)

*Resistant mutants selected from ILE-containing plates; †Resistant mutants selected from ILF-containing plates; -C1 or -X means ClpC1 or ClpX was edited.

ILE and ILF treatment alter Msm cell morphology and increases sensitivity in

clpC1 **and** ***clpX*** **knockdown strains**

To elucidate the bactericidal mechanisms of ILE and ILF, we detected the effect

of ILE or ILF treatment on the morphology of Msm cells. Msm was cultured with

0.125 µg mL^{-1} of ILE or 5 µg mL^{-1} of ILF for 15 hours (h). As a control, Msm was

also exposed to 1 µg mL^{-1} clarithromycin (CLR), a macrolide antibiotic that inhibits

bacterial protein synthesis by specifically targeting the 50S subunit of the ribosome.

The lengths of CLR-treated cells showed only a modest increase, from 2.07 ± 0.48 µm

to 2.64 ± 0.80 µm (Fig. 3a and Fig. 3b). Notably, the cell lengths of Msm increased to

5.95 ± 2.16 µm and 9.41 ± 2.40 µm following treatment with ILE and ILF,

respectively (Fig. 3c and Fig. 3d). In comparison to untreated and CLR-treated Msm,

ILE and ILF both induced a pronounced inhibition of cell division, as evidenced by

the formation of division septa (Fig. 3c and Fig. 3d). This observation is consistent

with the phenotype observed in Msm upon *clpX* gene silencing, and the established

role of ClpX in regulating FtsZ assembly and Z-ring formation which are crucial

processes for cell division in Mtb[40,41].

Fig 3 | The cell morphology of Msm strains before or after treatment. a The cell

morphology of Msm without any drug treatment. **b** The cell morphology of Msm

treated with a concentration of 1 µg mL^{-1} CLR. **c** The cell morphology of Msm

treated with a concentration of 0.125 µg mL^{-1} ILE. **d** The cell morphology of Msm

treated with a concentration of 5 µg mL^{-1} ILF. The white arrows highlight the division

septa.

Correspondingly, we performed knockdown of *clpX* and *clpC1* in Mab and Msm[42].

The knockdown of *clpX* or *clpC1* in Mab and Msm strains resulted in heightened

sensitivity to ILE and ILF (Fig. 4a and Fig. 4b). Upon induction with

anhydrotetracycline (aTc) in the absence of ILE or ILF, growth inhibition was

observed in both the *clpX* or *clpC1* knockdown strains of Msm and Mab, highlighting

the crucial role of these proteins in cellular proliferation. Additionally, we examined

ILE's intracellular activities against *clpC1* and *clpX* knockdown strains of Mab in

murine RAW264.7 macrophages (Fig. 4c). Upon infection and drug treatment, ILE

exhibited potent intracellular activity, with significantly enhanced efficacy against

clpX and *clpC1* knockdown strains compared to the vector control. Conclusively, ILE

and ILF may disrupt essential cellular processes regulated by ClpX and ClpC1,

leading to bactericidal effects.

Fig 4 | Impact of *clpX* or *clpC1* gene suppression on the susceptibility of Mab and Msm to ILE or ILF. a *In vitro* susceptibility of Mab to ILE or ILF upon aTc-induced gene suppression. **b** *In vitro* susceptibility of Msm to ILE or ILF upon aTc-induced gene suppression. The concentrations of ILE and ILF are in µg mL^{-1}. *, Mab strains contained varying plasmids: vector, a vector control with the empty plasmid pLJR962;

clpC1, pLJR962-clpC1 for inducible inhibition of the *clpC1* gene; *clpX*, pLJR962-

clpX for inducible inhibition of the *clpX* gene. [#], Similarly, Msm strains were

differentiated by the plasmids they harbored as that in Mab. **c** Intracellular

susceptibility of Mab to ILE upon gene suppression in RAW264.7 macrophages.

Survival rates were calculated as (CFU of ILE-treated group / CFU of untreated group)

× 100%. Numbers following the drug names indicate the working concentrations (μg

mL^{-1}). Statistical significance was determined by two-way ANOVA followed by

Tukey's multiple comparisons test. Data are presented as mean ± SD from three

biological replicates (n = 3). [**], $P < 0.01$; [****], $P < 0.0001$.

ILE and ILF selectively binds to ClpC1-NTD and is disrupted by resistance

mutations

Since all mutations occurred within ClpC1-NTD or ClpX-NTD across Mtb, Mab,

Msm, and Mmr, and their ClpC1-NTD or the first 64 amino acids of the ClpX-NTD

sequences are identical (Supplementary Fig. 4 and Supplementary Fig. 5), we

specifically expressed and purified the MtbClpC1-NTD$^{wt/mt}$ and MtbClpX-NTD$^{wt/mt}$.

Our goal was to investigate whether there is any interaction between these proteins

and ILE or ILF. The interactions between the MtbClpC1-NTD or MtbClpX-NTD with

ILE or ILF were examined by differential scanning fluorimetry (DSF)[43]. We

discovered that ILE increased the melting temperature (Tm) of MtbClpC1-NTD^{wt}

from 68.5□ to 77.5□ (Fig. 5a), signifying a significant enhancement in thermal

stability (ΔTm = 9□). Conversely, ILF marginally changed the thermal stability of the

MtbClpC1-NTD by merely 0.5□ (Fig. 5a) compared to ILE. The stronger affinity of

ILE over ILF correlates with its lower MIC. These findings suggest potential

interactions of ILE and ILF with MtbClpC1-NTD. To further test whether these

interactions are specific and mutation-dependent, we performed DSF assays using a

panel of MtbClpC1-NTD variants harboring the acquired mutations from Mtb

resistant strains (F80I/L/V/C). Notably, all four mutations markedly reduced the

thermal shift induced by ILE (Fig. 5b - Fig. 5e), indicating that these residues are

critical for compound binding. The loss of thermal stabilization strongly supports that

resistance arises via disruption of drug-target interaction. Notably, DSF assays of

MtbClpX-NTD failed to produce typical sigmoidal melting curves, likely due to the

small size and compact nature of the domain (residues 1-112), which may limit the

exposure of hydrophobic surfaces upon unfolding. As a result, we could not reliably

assess ligand binding to MtbClpX-NTD using this method.

Fig 5 | ILE and ILF bind to MtbClpC1-NTD directly. The melting curves of 5 μM

MtbClpC1-NTD^{wt} **(a)**, MtbClpC1-NTD^{F80I} **(b)**, MtbClpC1-NTD^{F80L} **(c)**, MtbClpC1-

NTD^{F80V} **(d)**, and MtbClpC1-NTD^{F80C} **(e)** in the presence of 10 μM ILE and ILF.

ILE significantly impedes the proteolytic function of the ClpC1P1P2 and

ClpXP1P2 proteases

Considering the stronger interactions observed between MtbClpC1-NTD and ILE

compared to ILF, we focused our subsequent biochemical assays on evaluating ILE.

We observed that ILE inhibited the proteolysis of the model substrate, fluorescein

isothiocyanate-casein (FITC-casein), by ClpC1^{wt}P1P2 complex compared to the

control group (Fig. 6a). However, the proteolytic activities of multiple mutant

ClpC1P1P2 proteases remain unaffected at 10 µg mL^{-1} ILE compared to the control,

with only partial inhibition noted at 20 µg mL^{-1} ILE (Fig. 6b - Fig. 6e).

For the ClpXP1P2 proteasome, we first used ssrA tagged green florescent protein

(GFP-ssrA) as the substrate. The results showed that even when the concentrations of

ILE or ILF reached as high as 2000 µg mL^{-1}, it had no effect on the proteolytic

activity of ClpXP1P2 (Supplementary Fig. 6a and Supplementary Fig. 6b). Previous

studies have demonstrated that the adaptor protein SspB from *E. coli* is required for

efficient degradation of proteins tagged with a mutated form of the ssrA tag DAS+4

by ClpXP1P2 in both *E. coli* and Msm[44,45]. Therefore, we expressed GFP tagged with

DAS+4 (GFP-DAS+4) as the substrate and SspB as the adaptor. Importantly, when

using DAS+4-tagged substrates, protein degradation could only be observed in the

presence of the adaptor SspB (Supplementary Fig. 6c)**.** It was observed a significant

inhibition of the proteolytic activity of the ClpX^{wt}P1P2 complex during the hydrolysis

of GFP-DAS+4 in the presence of SspB when ILE is added (Fig. 6f). However, we

discovered that the mutant ClpX^{P30H}P1P2 complex was incapable of degrading GFP-

DAS+4 (Supplementary Fig. 6d).

Additionally, we noted that ILE did not impact the ATP-dependent enzymatic

activities of ClpC1 and ClpX, as no significant differences in ATPase activities were

detected across varying ILE concentrations compared to controls (Supplementary Fig.

7).

Fig 6 | ILE significantly impedes the proteolytic function of both the ClpC1P1P2 proteases and the ClpXP1P2 proteases. Degradation of FITC-casein by **(a)** ClpC1^{wt}P1P2, **(b)** ClpC1^{F80I}P1P2, **(c)** ClpC1^{F80L}P1P2, **(d)** ClpC1^{F80V}P1P2, and **(e)** ClpC1^{F80C}P1P2 proteases in the presence or absence of ILE. Data are presented as

mean ± SD from three biological replicates (n = 3). **f** Degradation of GFP-DAS+4 by

ClpX^{wt}P1P2 proteases in the presence or absence of ILE. Panel (f) shows data from a

representative experiment without error bars. Numbers following the drug names

indicate the working concentrations (μg mL^{-1}). DMSO was added as control. The D-

value of fluorescence intensity represents the change in fluorescence intensity by

subtracting the initial fluorescence intensity from the fluorescence intensity of

subsequent detections. Statistical significance was determined using one-way

ANOVA. *, $P < 0.05$; ***, $P < 0.0001$; ****, $P < 0.00001$; ns, no significant difference,

where $P \geq 0.05$.

ILE disrupts protein homeostasis with accumulation of ribosomal components

and toxin-antitoxin (TA) systems

To elucidate the adaptive responses of Mtb to ILE and gain insight into its

mechanism of action, we performed quantitative proteomic analysis of Mtb cells

treated with ILE (Supplementary Table 4). Compared to untreated controls, a total of

773 proteins were upregulated and 280 proteins were downregulated (Fig. 7a),

indicating a global increase in protein abundance.

Fig 7 | Proteomic profiling reveals global protein expression changes of Mtb in

response to ILE treatment. a Bar graph summarizing the number of differentially

expressed proteins upon ILE treatment compared to the control group. Upregulated

and downregulated proteins are categorized by fold change (FC): FC 2-10, FC >10,

FC 0.1-0.5, and FC <0.1. **b** Volcano plot showing the distribution of differentially

expressed proteins.

Notably, ribosomal proteins—including components of the 50S and 30S subunits

(e.g., RpsG, RpsS)—were broadly upregulated, with 21 out of 23 ribosomal proteins

showing increased abundance. RplR and RpsC were the only ribosomal proteins

found to be downregulated.

All differentially expressed TA system proteins were upregulated, including

proteins from VapBC (27 proteins), MazEF (11 proteins), ParDE/RelBE (7 proteins),

HigBA (3 proteins) and mt-PemIK (1 protein) families. Additionally, the reported

ClpP1P2 substrates PanD (FC = 2.67, $P < 0.05$) and CarD (FC = 5.99, $P < 0.001$) was

also significantly upregulated[46,47]. Proteins associated with cell division, such as FtsZ

and FtsE, were upregulated approximately two-fold ($P < 0.001$), consistent with

scanning electron microscopy observations.

To visualize changes in proteins functionally associated with the Clp protease

system, key components including ClpX, ClpS, and ClgR were highlighted in the

volcano plot (Fig. 7b). ClpX was downregulated (FC = 0.41, $P < 0.05$), whereas ClpS

and ClgR were highly upregulated (FC = 187.26 and 8.53, respectively; both $P <$

0.01). Other components such as ClpC1, ClpC2, ClpP1, and ClpP2 showed no

significant changes and were therefore not highlighted.

Discussion

The study of natural cyclic peptides has emerged as a focal point in the current research on drugs against Mtb. In comparison to the reported cycloheptapeptides, ILE demonstrates superior anti-Mtb activity. Specifically, RUFI inhibited Mtb growth at a MIC of 60 nM by day 14 when tested with the autoluminescent Mtb strain, while ILE achieves an MIC of 48 nM (equivalent to 0.05 µg mL^{-1}) by day 15 (Fig. 1a)[48]. Moreover, the MIC of ILE is significantly lower than those of CYMA (100 nM), ECU (160 nM), and lassomycin (410 nM) reported[13,18]. While these compounds exhibit activity against a range of NTM, only RUFI and ILE exhibit efficacy against Mab, indicating their potential as therapeutic agents for diseases caused by Mab[15].

Previous research had not explored the combination effects of cycloheptapeptides with anti-TB drugs, leaving their potential as adjuvants to current TB therapies unresolved[14-16,18]. Notably, our findings on combinatorial activity suggest that ILE have the potential to enhance the efficacy of existing anti-TB drugs, including RIF, INH, EMB, and TB47. Proteomic analysis provided mechanistic insights into these interactions, revealing that ILE treatment modulated the expression of multiple proteins involved in pathways targeted by these drugs. As for RIF, ILE-induced downregulation of Rho and NusA, two key transcriptional regulators that interact with RNA polymerase. Given that Rho suppression causes widespread transcriptional disruption and rapid cell death, and that NusA modulates transcriptional pausing and termination, their downregulation may sensitize Mtb to RIF-induced RNA polymerase inhibition[49,50]. In the case of INH, four proteins—SigL, Ndh, FadE24, and Glf—were differentially expressed in response to ILE. Specifically, SigL and Ndh were significantly upregulated, whereas FadE24 and Glf were downregulated. SigL has been reported to regulate the expression of KatG, which is crucial for the activation of INH, and Ndh has been identified to influence the NADH/NAD+ ratio, affecting INH-NAD formation[51-53]. INH-resistant Mtb has been found to carry mutations in FadE24[54]. Upregulation of Glf was hypothesized to contribute to INH resistance in Mtb either by binding to a modified form of INH or by sequestering a factor such as

NAD+ required for INH activity[55]. For EMB, ILE downregulated UbiA, a key

enzyme in the decaprenylphosphoryl-D-arabinose biosynthesis pathway required for

cell wall assembly. Given that UbiA overexpression has been associated with EMB

resistance, its suppression by ILE may underlie the observed partial synergy[56].

Additionally, synergy with TB47—a cytochrome bc1 complex (QcrB) inhibitor—can

be explained by ILE-mediated suppression of several electron transport chain

components, including QcrA, QcrC, and CydD[24,57]. As the Qcr and Cyt-bd branches

of the electron transport chain are functionally redundant, simultaneous inhibition by

TB47 and ILE likely causes a collapse in respiratory function, resulting in pronounced

synergistic killing[24].

Although ILE and ILF have demonstrated potent activity against Mtb, their

MOAs remained elusive[17]. Previous studies had identified ClpC1 as the primary target

of several cyclic peptides[32]. Consistent with this, we found that mutations in ClpC1—

particularly in its NTD—confer resistance to ILE and ILF across multiple

mycobacterial species, with the exception of unique insertion mutations discovered

between M1 and F2 in both Mab and Msm[15,16,18,36]. Previous studies indirectly

validated these mutations via overexpression[36], but using advanced mycobacterial

gene-editing tools, we now directly confirm that specific mutations (e.g., Mab-F2L,

Msm-F2C, and a novel Msm insertion) confer ILE/ILF resistance. Additionally, WGS

of 2 ILE-resistant Mtb mutants revealed identical large insertions in Rv1572c, a

conserved hypothetical protein whose role remains to be explored. Moreover, 3 ILE-

resistant Mab mutants were found to carry mutations in *hisS* or *hisD*, which encode

histidyl-tRNA synthetase and histidinol dehydrogenase, respectively. Although

overexpression of mutated version of these genes did not restore drug sensitivity, gene

editing-based validation is planned to confirm their potential role in resistance.

Interestingly, no ClpX mutations had been reported in mutants resistant to these

compounds[15,16,18]. However, our findings indicate that ILE-resistant mutants

harboring ClpX mutations also exhibit cross-resistance to ILF, suggesting a shared

resistance mechanism. Moreover, gene editing confirmed that ClpX mutations can

directly confer high-level resistance to both ILE and ILF, revealing ClpX as a novel

contributor to cyclic peptide's resistance. These findings also shed light on possible

limitations of previous resistance screening approaches. Whether the ClpX mutation

can lead to resistance of mycobacteria to other ClpC1-targeting cyclic peptides—such

as RUFI, ECU, and lassomycin—remains to be determined. Given RUFI's close

structural similarity to ILE/ILF and strong activity against Mab, this is particularly

worth investigating[15,58]. Molecular docking and gene editing techniques provide

effective strategies to explore this possibility.

The MOA of ILE differs from ECU and lassomycin, which stimulate ClpC1

ATPase activity but hinder substrate degradation, as well as CYMA, which activates

both ATPase and proteolytic functions[16,18,36]. In this study, we demonstrated that ILE

and ILF inhibited the proteolytic activity of both ClpC1P1P2 and ClpXP1P2 proteases

without affecting ATPase activity, which is similar to RUFI's mechanism[15]. Notably,

a higher concentration of ILE is required to inhibit ClpXP1P2 than ClpC1P1P2,

aligning with the higher resistance levels associated with ClpX mutations. These

observations highlight ClpX mutation as a previously unrecognized resistance

mechanism in mycobacteria, warranting further investigation.

Proteomic analysis further revealed a broad upregulation of proteins in response

to ILE treatment, consistent with a disruption of proteostasis resulting from ClpC1

and ClpX inhibition. Impaired protease activity would prevent normal turnover of

these proteins, resulting in their accumulation. This may explain the observed increase

in ribosomal protein levels as a compensatory response to maintain translational

capacity. Notably, multiple TA systems—including VapBC, MazEF, ParDE/RelBE,

HigBA, and mt-PemIK families—were strongly upregulated, aligning with their

known degradation by ClpC1P1P2 or ClpXP1P2[9,59]. Given their roles in growth

modulation, dormancy, and virulence, such dysregulation could contribute to the

antibacterial effects of ILE treatment[60]. Additionally, accumulation of PanD and CarD,

two validated ClpP1P2 substrate, further supports protease inhibition[46,47]. ClpX was

downregulated, potentially relieving repression of FtsZ and explaining its

upregulation[40]. This is consistent with previous reports that ClpX knockdown leads to

filamentous cell morphology due to disrupted FtsZ regulation in mycobacteria[41], and

may explain the elongated cells we observed under SEM in ILE/ILF-treated strains.

ClpS was dramatically induced, likely to compensate for impaired substrate

recognition[59]; and the transcriptional activator ClgR was upregulated, consistent with

proteostasis stress responses observed under proteasome inhibition[61]. These findings

indicate a coordinated regulatory response aimed at restoring protein homeostasis

under Clp protease dysfunction.

However, we discovered that the mutant $ClpX^{P30H}P1P2$ complex was incapable of

degrading GFP-DAS+4. This may be attributed to the fact that the enhancement of

ClpXP1P2 degradation by SspB requires an intact zinc binding domain (ZBD) in

ClpX (residues 1-60)[62]. A mutation in P30, which is located in the hydrophobic

surfaces of ZBD, may suppress the binding of SspB to ClpX, thereby preventing the

delivery of the tagged protein to ClpXP1P2 complexes for degradation. Although

SspB homologs have not been identified in mycobacterium, it is possible that native

adaptor-like mechanisms exist, and disruption of such interactions may contribute to

ILE resistance. Notably, despite the presence of $ClpC1^{wt}$, the $ClpX^{P30H}$ mutation

confers resistance, suggesting that it may compensate for ClpC1 dysfunction by

altering substrate specificity or proteostasis regulation. Additionally, the roles of some

clpC1 mutation sites in ILE and ILF resistance remain unproven due to the inability to

acquire corresponding gene-edited strains. Therefore, further studies are needed to

identify native ClpX substrates or adaptors in mycobacteria, assess ClpX mutant

proteolytic activity in more physiological contexts, and attempt to obtain the

remaining *clpC1*-edited strains to confirm their roles in ILE and ILF resistance.

While ClpX is not currently considered the target of ilamycins in Mtb, its

presence in humans—unlike ClpC1—raises concerns about potential off-target effects.

Therefore, we performed sequence analysis shows low similarity (< 25% identity) and

no conservation at resistance-related residues between Mtb and human ClpX,

suggesting minimal risk. However, further validation is warranted to fully exclude

off-target interactions.

In summary, the potent and broad antimycobacterial activities of ILE and ILF

underscore their potential for further development as effective agents against

mycobacteria, especially drug-resistant strains. Our findings not only reinforce the

critical role of the ClpC1P1P2 protease as a validated target, but also uncover ClpX—

via mutation—as a previously unrecognized mechanism of resistance[15,16,18,36]. This

discovery significantly expands the current understanding of resistance pathways

beyond *clpC1*, and suggests that ClpXP1P2 itself represents a promising and

underexplored target for antimycobacterial therapy. A dual-targeting strategy that

affects both simultaneously is a rational approach, providing a new avenue for

developing treatments against mycobacterial infections.

Materials and Methods

Bacterial strains and culture conditions

The bacterial strains used in this study are listed in Supplementary Table 5. Mtb, Mab, Mmr, Msm, and derived strains were grown in Middlebrook 7H9 broth (BD, USA) or 7H10 agar (BD) or 7H11 agar (Acmec, China) supplemented with OADC (comprising 0.005% oleic acid, 0.5% bovine serum albumin, 0.2% dextrose, 0.085% catalase). Mtb, Msm, and Mab were grown at 37□. Mmr were grown at 30□. For *E. coli*, *Enterococcus faecium*, *Staphylococcus aureus*, *Acinetobacter baumannii*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa*, cells were grown in Luria-Bertani (LB) broth at 37□. Antibiotics were supplemented as required at the appropriate screening concentrations for distinct bacterial species. Specifically, for *E. coli* and Msm, the antibiotics added were kanamycin (KAN, 50 µg mL^{-1}, Solarbio, China), apramycin (APR, 50 µg mL^{-1}, MeilunBio, China), and Zeocin (ZEO, 30 µg mL^{-1}, InvivoGen, France). For Mmr, only APR (50 µg mL^{-1}) were utilized. Meanwhile, for Mab, KAN was used at a higher concentration of 100 µg mL^{-1}, along with ZEO (30 µg mL^{-1}) and an increased concentration of APR (230 µg mL^{-1}).

Determination of MICs and MBC against mycobacteria *in vitro* and *ex vivo*

The MICs of ILE and ILF were determined using a cost-effective *in vitro* assay as previously described against a selectable marker-free AlRa, AlRv, AlMab, AlMsm and AlMmr[23-25,30]. The MIC$_{lux}$ was defined as the lowest concentration that could inhibit > 90% relative light units (RLUs) compared with that from the untreated controls[23-25,30]. The MICs of the ILE were also determined by a well-established microplate alamar blue assay (MABA) against Mtb H$_{37}$Ra, H$_{37}$Rv and nine clinical Mtb strains isolated from pulmonary TB patients hospitalized at Guangzhou Chest Hospital, Guangzhou, China[29]. The MIC, as determined by the MABA method, is defined as the lowest concentration that prevents the color change from blue to pink, as observed with the naked eye.

MBC was defined as the minimum concentration resulting in ≥ 99.9% reduction in CFUs compared to the bacterial load before drug addition. For CFU enumeration,

mycobacterial cultures treated with different concentrations of compounds or control

were collected before or after incubation, with the incubation time varying depending

on the species. Bacterial suspensions were serially diluted in phosphate-buffered

saline (PBS, GENOM, China), plated onto 7H10 agar plates, and incubated at 37°C

until visible colonies appeared. The number of visible colonies was counted, and

CFUs were calculated to evaluate bacterial viability.

Intracellular survival assay was performed as previously reported[63]. RAW264.7

macrophages were seeded at $5 \times 10\square$ cells per well in 12-well plates and infected

with different Mab at an MOI of 10:1 for 4 h. After removing extracellular bacteria

with 250 µg mL^{-1} amikacin for 2 h, infected cells were cultured with or without

ilamycins as well as 200 ng mL^{-1} aTc. At 72 h post-infection, macrophages were lysed,

and intracellular bacteria were quantified by CFU enumeration on 7H10 agar. All

experiments were performed in triplicate.

MICs to non-mycobacteria determination

The MICs of ILE or ILF against *Enterococcus faecium*, *Staphylococcus aureus*,

Acinetobacter baumannii, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* were

determined by observing the turbidity after 16 h of incubation in LB broth at 37□.

The MIC was defined as the lowest concentration, resulting in clarity in turbidity

relative to that of untreated control cultures[64-66].

Combined activity of ILE or ILF with other compounds *in vitro*

The combined activity of ILE or ILF with other compounds, such as INH

(MeilunBio), RIF (MeilunBio), EMB (MeilunBio), STR (MeilunBio), CFZ

(MeilunBio), or BDQ (biochempartner, China), and TB47 (BojiMed, China), was

evaluated *in vitro* using the checkerboard method. The FICI was interpreted as

follows: synergistic (FICI of ≤ 0.5), partially synergistic ($0.5 < \text{FICI} < 1.0$), additive

(FICI = 1.0), indifferent ($1.0 < \text{FICI} \leq 4.0$), and antagonistic (FICI > 4.0)[67,68].

Determination of MIC and TC$_{50}$ in macrophage culture

The cell density of mouse macrophages RAW264.7 was adjusted to 5×10^5 cells

mL^{-1}. 100 µL of cells per well were added to 96-well plates and cultured overnight for

adherence. The AlRv concentration was adjusted to 3×10^6 RLU mL^{-1}, and 100 μL of

AlRv was added to each well. The plates were then incubated at 37□ for 6 h. The

cells were washed three times with DMEM (Gibco, USA) to remove non-

phagocytosed AlRv and treated with DMEM containing amikacin (100□μg mL^{-1},

MeilunBio, China) for 2□h to eliminate the extracellular bacteria. The cells were

washed twice with DMEM to remove residual amikacin. A total of 195 μL of DMEM

containing 10% fetal bovine serum and 5 μL of the two-fold dilutions of various ILE

concentrations were added to the cells. The mixture was incubated at 37□ with 5%

CO_2, and the RLUs were measured on days 1-5 or 7 of incubation. RIF (1 μg mL^{-1})

and dimethyl sulfoxide (DMSO, Xilong Chemical, China) were used as positive and

negative controls, respectively.

The TC$_{50}$ of ILE was assessed in mouse RAW264.7 macrophages using the Cell

Counting Kit-8 (CCK-8, MeilunBio, China). Cells were seeded in 96-well plates at a

density of 1□×□10□ cells per well and allowed to adhere overnight. Serial dilutions

of ILE were added and incubated for 24□h at 37□ with 5% CO_2. CCK-8 reagent was

then added according to the manufacturer's instructions, and absorbance was

measured at 450□nm using a microplate reader. The TC□□ value was calculated

from dose–response curves using GraphPad Prism.

ILE- or ILF-resistant mutants

Broth cultures of AlRa, AlRv, AlMab, and AlMmr, OD$_{600}$ reaching 0.6-1.3, were

plated on 7H11 plates containing ILE at 0.1, 0.5, 2.5, 5, 10 or 20 μg mL^{-1}. Broth

cultures of AlMsm (OD$_{600}$ 0.6-1.3) were plated on 7H11 plates containing ILF at 10,

20 or 40 μg mL^{-1}. The bacterial colonies grown on the ILE- or ILF-containing plates

were selected and grown in liquid culture for drug susceptibility testing to confirm the

drug resistance phenotype.

WGS analysis

WGS analysis was conducted on the parent strains of Mtb, Mab, and a subset of

the confirmed ILE-resistant mutants. The sequencing was performed by the Beijing

Genomics Institute in China. Subsequently, the obtained reads were aligned with the

reference genome sequence and compared to that of the parent strain. Mutations

identified through WGS in the drug-resistant mutants were validated by amplifying

and sequencing the *clpX* and *clpC1* genes using Sanger sequencing. Additionally, the

clpX and *clpC1* genes from other ILE- or ILF-resistant mutants, including Mtb, Mab,

Msm, and Mmr, were amplified and sequenced using Sanger sequencing (primer

sequences are provided in Supplementary Table 6).

Molecular docking analysis

To evaluate the binding affinities and interaction modes among ILE, MtbClpC1-

NTD, and MtbClpX-NTD, we employed Autodock Vina 1.2.2, a computational

protein-ligand docking software previously reported[33]. The structure of ILE was

obtained from PubChem Compound[34]. The 3D coordinates of MtbClpC1-NTD were

retrieved from the Protein Data Bank (PDB) with PDB code 6CN8 and a resolution of

1.4 Å, while the structure of MtbClpX-NTD was predicted using AlphaFold2.

For the docking analysis, all protein and ligand files were converted to the

PDBQT format; water molecules were excluded, and polar hydrogen atoms were

supplemented. A grid box was positioned to envelop the domain of each protein,

facilitating unrestricted molecular movement. The dimensions of the grid box were set

to 30 Å $\times$ 30 Å $\times$ 30 Å, with a grid point spacing of 0.05 nm. The molecular docking

experiments were executed using Autodock Vina 1.2.2, which is accessible at

http://autodock.scripps.edu/.

Overexpression of *clpX* or *clpC1* in Mab, Mmr or Msm

The *clpC1*^{wt}, *clpC1*^{mt}, *clpX*^{wt} and *clpX*^{mt} from Mtb, Mab, Mmr and Msm were

amplified using primers detailed in Supplementary Table 3 and subsequently cloned

into the plasmid p60A or pMVA with multi-copy under the control of the *hsp60*

promoter as depicted in Supplementary Fig. 3. The plasmids were then transformed

into ΔMab, ΔMmr, and ΔMsm. Additionally, the *clpP1*, *clpP2* and *clpX* or *clpC1* of

Mab were amplified from the genome of Mab^{wt} and inserted into the plasmid pMVA

under the control of the *hsp70* or *hsp60* promoters to enable the simultaneous

expression of ClpXP1P2 or ClpC1P1P2. The MICs of the recombinant strains to ILE

or ILF were determined using the aforementioned protocols.

Silencing *clpX* or *clpC1* genes in Mab or Msm

The CRISPR-Cas9 system was employed for targeted gene silencing of *clpX* and

clpC1[42]. Small guide RNAs (sgRNAs) of 20 base pairs were designed to target the

coding regions of *clpX* and *clpC1* on their sense strands. The oligonucleotides were

annealed and then ligated with linearized pLJR962[42]. Plasmids harboring sgRNAs

against *clpX* or *clpC1*, along with the empty vector pLJR962 as a control, were

introduced into Mab and Msm cells via electroporation (Supplementary Table 7).

Tenfold serial dilutions of Mab^{wt} or Msm^{wt}, the control strain carrying the empty

vector, and the gene-silenced strains were prepared after reaching an OD$_{600}$ of

approximately 0.9. For Mab, 1 µL of each dilution was spotted on plates containing

either 0.25, 0.5, and 1 µg mL^{-1} ILE or 5, 10, and 20 µg mL^{-1} ILF with varying

concentrations of aTc (0, 100, 200 and 400 ng mL^{-1}). For Msm, 1 µL of each dilution

was spotted on plates containing either 0.004, 0.008, and 0.01625 µg mL^{-1} ILE or

0.125, 0.25, and 0.5 µg mL^{-1} ILF with varying concentrations of aTc (0, 12.5, 25 and

50 ng mL^{-1}). Control plates received 1 µL of each dilution with aTc alone. The plates

were incubated at 37□ for 3 days. The experiment was performed in triplicate and

repeated twice.

CRISPR/Cpf1-mediated gene editing of *clpX* and *clpC1* genes in Mab and Msm

The CRISPR/Cpf1-associated recombination have been previously employed for

gene editing in both Mab and Msm recently[38,39]. Initially, the pJV53-Cpf1 plasmid

was electroporated into Mab^{wt} and Msm^{wt} strains, resulting in the Mab::pJV53-Cpf1

and Msm::pJV53-Cpf1, respectively. Subsequently, crRNAs were designed to target

specific 25 nucleotide sequences adjacent to the 5'-YTN-3' motif on the template

strand of the target genes (Supplementary Table 7). Oligonucleotides for crRNA

expression were ligated into the pCR-zeo vector. These constructed vectors were then

electroporated into the respective acetamide-induced cells of Mab::pJV53-Cpf1 or

Msm::pJV53-Cpf1. For the induction of Cpf1 expression, aTc was added to 7H11

agar plates to achieve a final concentration of 200 ng mL^{-1}. These plates were cultured

at 30□ for 3-5 days. Individual colony was picked and screened to confirm successful

editing at the target gene site via PCR and Sanger sequencing (primers listed in

Supplementary Table 6). The MICs of the edited strains to ILE and ILF were

determined using the methods described earlier.

Expression and purification of proteins

The Mtb ClpP1 (residues 7-200) with a C-terminal Strep-II tag and ClpP2

(residues 13-214) with a C-terminal His$_6$-tag were expressed and purified. Briefly,

clpP1 and *clpP2* genes were inserted into the pETDute-1 vector and expressed in *E.*

coli BL21 (λDE3) in LB broth at 25□, following induction with 0.5 mM isopropyl β-

D-1-thiogalactopyranoside (IPTG, Macklin, China) for approximately 16 h. Cells

were resuspended and lysed; The target protein was subsequently purified via nickel

affinity chromatography and eluted with the same buffer supplemented with 200 mM

imidazole. Eluted fractions containing the target protein were loaded onto a Strep-II

column. The target protein was eluted with the strep buffer containing 50 mM

HEPES-KOH (pH 7.5), 150 mM KCl, 10 mM MgCl$_2$, 50 mM d-Desthiobiotin and 10%

glycerol (v/v), before a final gel filtration using on a SuperoseTM 6 increase 10/300

GL column (GE Healthcare) in SEC buffer (25 mM HEPES-KOH (pH 7.5), 150 mM

KCl, 10 mM MgCl$_2$).

The sequence of *clpX*^{wt} or *clpX*^{P30H} was inserted into the pGEX-6P-1 vector,

which had a His$_6$-tag at the C-terminal and an HRV 3C protease cleavage site. After

induction for 18 h at 16□, the cells were collected by centrifugation. The pellet was

then resuspended and lysed. After centrifugation, the supernatant was loaded onto a

GST column and eluted. The eluent underwent digestion at 4□ for 26 h with enough

HRV 3C protease. The fractions containing the target protein were then pooled,

concentrated to 1 mL, and purified with a SuperoseTM 6 increase 10/300 GL column.

The sequence of *clpC1*^{wt}, its mutants (F80I/V/L/C), the gene encoding *gfp-ssrA*

(AADSHQRDYALAA), *gfp*-DAS+4 (AANDENYSENYADAS) or *E. coli sspB* was

inserted into the pET-28a vector that had an N-terminal His$_6$-tag. The recombinant

strains were cultured in LB broth at 37□ until the OD$_{600}$ reached 0.6-0.8. After 18 h of

induction with 0.5 mM IPTG at 18□, the pellet was suspended and lysed. Following

the loading of the supernatant onto a Ni-NTA column, the proteins were washed and

then eluted. The fractions were then pooled and loaded onto a SuperoseTM 6 increase

10/300 GL column.

The gene encoding MtbClpC1-NTD$^{wt/mt}$ or MtbClpX-NTD$^{wt/mt}$ was inserted into

the pET-28a vector with a C-terminal His$_6$-tag and an HRV 3C protease cleavage site.

The expression and purification methods of MtbClpC1-NTD and MtbClpX-NTD

were consistent with the methods described above.

DSF

The binding of ILE or ILF to the MtbClpC1-NTD and MtbClpX-NTD was

detected using DSF, following established protocols[43,69]. A 5 µM solution of either

MtbClpX$^{wt/mt}$-NTD or MtbClpC1$^{wt/mt}$-NTD was incubated with varying

concentrations of ILE or ILF (ranging from 10 to 400 µM), as well as DMSO for 30

minutes, subsequently mixed with Sypro Orange fluorescent dye (Invitrogen, USA).

Fluorescence measurements were recorded using the CFX96 Real-Time System (Bio-

rad, USA), which involved heating 8-tube PCR Strips from 25□ to 95□ at a constant

rate of 0.4□ per minute. The excitation and emission filters were adjusted to

wavelengths of 460 nm and 510 nm, respectively. Data collection was facilitated by

the CFX Manager software, with analysis performed using GraphPad Prism version

8.3.0. To ensure accuracy, all measurements were performed in triplicate.

Proteolytic activity assays of ClpC1P1P2 and ClpXP1P2

The proteolytic activities of ClpC1$^{wt/mt}$P1P2 and ClpX$^{wt/mt}$P1P2 were measured at

37□ in black 96-well plates using a Multimode Plate Reader. For the ClpC1P1P2

proteolytic activity experiments, ILE was added at final concentrations of 0, 0.1, 1, 10,

and 20 µg mL^{-1} in each well. A 100 µL reaction mixture was formulated with 2 µM

ClpP1P2, 1 µM ClpC1$^{wt/mt}$, 2 mM ATP, and 2.5 µM FITC-casein (AAT Bioquest,

USA).

For the ClpXP1P2 proteolytic activity experiments, two different ssrA-tagged

GFP substrates were used: GFP-ssrA and the mutated ssrA-tagged GFP (GFP-

DAS+4). For the GFP-ssrA substrate, the reaction mixture (100 μL) contained 0.2 μM

ClpP1P2, 0.7 μM ClpX, 2 mM ATP, 1 μM GFP-ssrA, and 7 μM bortezomib (AbMole,

USA) as an activator[70]. For the DAS+4-tagged substrate, GFP-ssrA was replaced with

1 μM GFP-DAS+4, and 2 μM SspB adaptor protein was added.

The hydrolysis of FITC-casein, GFP-ssrA and GFP-DAS+4 was continuously

monitored at 535 nm (with excitation at 485 nm). Measurements were performed in

triplicate.

ATPase activity assays of ClpC1 and ClpX

The ATPase activities of Mtb ClpC1 and ClpX were assessed using a free-

phosphate detection assay with the Biomol Green reagent (Enzo Life Sciences, USA).

To each well of a transparent 96-well plate, 5 μL of various concentrations of ILE (0,

0.08, 0.16, 0.31, 0.625, 1.25, 2.5, 5 and 10 μg mL^{-1}) was added. Subsequently, either

ClpX or ClpC1 was diluted to a concentration of 0.1 μg/mL in the reaction mixture.

90 μL ClpC1 and ClpX were then added to a 96-well plate and incubated at 37□ for 1

h. The reaction was initiated by adding 5 μL of 1 mM ATP to the reaction mixture,

which was maintained at 37□ for 1 h. The final volume of the reaction mixture was

adjusted to 100 μL. Post reaction, 100 μL of Biomol Green reagent was introduced to

determine the amount of free phosphate. After a 30-minute incubation at room

temperature, the resulting reaction product was measured at 620 nm using a

Multimode Plate Reader. The concentration of free phosphate liberated from ATP by

ClpX and ClpC1 was calculated using a standard curve based on the known

concentration of free phosphate. Measurements were performed in triplicate.

Microscopy of drug-treated Msm

Treatment of exponentially growing Msm was conducted using 0.125 μg mL^{-1}

ILE, 5 μg mL^{-1} L ILF, or 1 μg mL^{-1} L CLR (Meilunbio, China) for a duration of 15 h.

The bacteria were immobilized with a 4% glutaraldehyde solution, followed by three

washes and resuspension in PBS. Bacterial suspensions were then applied to 5 mm

cell slides, which were placed in a 24-well plate to dry naturally. Afterward, the cell

slides underwent three additional PBS washes. The cell slides were subsequently

soaked in ascending concentrations of ethanol solutions (70%, 80%, 90%, 100%) for

10 minutes each. Finally, the cell slides were dried using the critical point drying

method and coated with an electrically-conducting material. Observations of bacterial

morphology were made using a GeminiSEM 300 ultra-high resolution field emission

cryo-scanning electron microscope.

Proteomic analysis

The AlRa cells were treated with 0.0125 μg mL^{-1} of ILE for 5 days in triplicate

for 4D label-free phosphoproteomic analysis. To maintain comparable cell density

between the untreated and ILE-treated groups, the untreated cells were diluted on the

same day as the compound addition, prior to sample collection. The cells were

resuspended and boiled for 20 minutes. Subsequently, the samples were sent to

APTBIO (Shanghai, China) for proteomic testing and analysis.

To compare the abundance of phosphopeptides between the control and treatment

samples, label-free quantification was performed with a minimum FC of 2 to

determine the differentially expressed phosphopeptides. Additionally, the student's *t*-

test was employed to detect significant differences between the control and treatment

samples, with $P < 0.05$ indicating significance. rotein functional annotation was

conducted using NCBI and Mycobrowser (https://mycobrowser.epfl.ch/). All

differentially expressed proteins were listed in Supplementary Table 4.

Data availability

The Sequence Read Archive (SRA) for a subset of ILE-resistant Mab strains

have been deposited in the National Center for Biotechnology Information (NCBI)

under BioProject Name PRJNA1262689. For the remaining samples, only variant

analysis results provided by the sequencing company are available and are included in

Supplementary Table 2 and 3. All other data are available from the corresponding

author on reasonable request.

References

1. Miotto, P., Zhang, Y., Cirillo, D. M. & Yam, W. C. Drug resistance mechanisms and drug susceptibility testing for tuberculosis. *Respirology* **23**, 1098-1113 (2018).

2. Dheda, K. *et al.* Multidrug-resistant tuberculosis. *Nature Reviews. Disease Primers* **10**, 22 (2024).

3. Fedrizzi, T. *et al.* Genomic characterization of nontuberculous mycobacteria. *Sci. Rep.* **7**, 45258 (2017).

4. Wu, M.-L., Aziz, D. B., Dartois, V. & Dick, T. NTM drug discovery: status, gaps and the way forward. *Drug Discov. Today* **23**, 1502-1519 (2018).

5. Lee, H. & Suh, J. W. Anti-tuberculosis lead molecules from natural products targeting *Mycobacterium tuberculosis* ClpC1. *J. Ind. Microbiol. Biotechnol.* **43**, 205-212 (2016).

6. Lupoli, T. J., Vaubourgeix, J., Burns-Huang, K. & Gold, B. Targeting the proteostasis network for mycobacterial drug discovery. *ACS Infect. Dis.* **4**, 478-498 (2018).

7. Fraga, H. *et al.* Development of high throughput screening methods for inhibitors of ClpC1P1P2 from *Mycobacteria tuberculosis*. *Anal. Biochem.* **567**, 30-37 (2019).

8. d'Andrea, F. B. *et al.* The essential *M. tuberculosis* Clp protease is functionally asymmetric *in vivo*. *Sci. Adv.* **8**, eabn7943 (2022).

9. Texier, P. *et al.* ClpXP-mediated degradation of the TAC antitoxin is neutralized by the SecB-like chaperone in *Mycobacterium tuberculosis*. *J. Mol. Biol.* **433**, 166815 (2021).

10. Akopian, T. *et al.* The active ClpP protease from *M. tuberculosis* is a complex composed of a heptameric ClpP1 and a ClpP2 ring. *EMBO J.* **31**, 1529-1541 (2012).

11. Raju, R. M. *et al. Mycobacterium tuberculosis* ClpP1 and ClpP2 function together in protein degradation and are required for viability *in vitro* and during infection. *PLoS Pathog.* **8**, e1002511 (2012).

12. Sassetti, C. M., Boyd, D. H. & Rubin, E. J. Genes required for mycobacterial growth defined by high density mutagenesis. *Mol. Microbiol.* **48**, 77-84 (2003).

13. Boshoff, H. I., Malhotra, N., Barry III, C. E. & Oh, S. The antitubercular activities of natural products with fused-nitrogen-containing heterocycles. *Pharmaceuticals (Basel)* **17**, 211 (2024).

14. Schmitt, E. K. *et al.* The natural product cyclomarin kills *Mycobacterium tuberculosis* by targeting the ClpC1 subunit of the caseinolytic protease. *Angew. Chem., Int. Ed. Engl.* **50**, 5889-5891 (2011).

15. Choules, M. P. *et al.* Rufomycin targets ClpC1 proteolysis in *Mycobacterium tuberculosis* and *M. abscessus*. *Antimicrob. Agents Chemother.* **63**, e02204-02218 (2019).

16. Gao, W. *et al.* The cyclic peptide ecumicin targeting ClpC1 is active against *Mycobacterium tuberculosis in vivo*. *Antimicrob. Agents Chemother.* **59**, 880-889

(2015).

17   Ma, J. *et al.* Biosynthesis of ilamycins featuring unusual building blocks and
engineered production of enhanced anti-tuberculosis agents. *Nat. Commun.* **8**,
391 (2017).

18   Gavrish, E. *et al.* Lassomycin, a ribosomally synthesized cyclic peptide, kills
Mycobacterium tuberculosis by targeting the ATP-dependent protease
ClpC1P1P2. *Chem. Biol.* **21**, 509-518 (2014).

19   Victoria, L., Gupta, A., Gómez, J. L. & Robledo, J. *Mycobacterium abscessus*
complex: a review of recent developments in an emerging pathogen. *Front. Cell.*
Infect. Microbiol. **11**, 659997 (2021).

20   Luthra, S., Rominski, A. & Sander, P. The role of antibiotic-target-modifying and
antibiotic-modifying enzymes in *Mycobacterium abscessus* drug resistance.
Front. Microbiol. **9**, 2179 (2018).

21   Stinear, T. P. *et al.* Insights from the complete genome sequence of
Mycobacterium marinum on the evolution of *Mycobacterium tuberculosis.*
Genome Res. **18**, 729-741 (2008).

22   Shiloh, M. U. & Champion, P. A. D. To catch a killer. What can mycobacterial
models teach us about *Mycobacterium tuberculosis* pathogenesis? *Curr. Opin.*
Microbiol. **13**, 86-92 (2010).

23   Yang, F. *et al.* Engineering more stable, selectable marker-free autoluminescent
mycobacteria by one step. *PLoS ONE* **10**, e0119341 (2015).

24   Liu, Y. *et al.* The compound TB47 is highly bactericidal against *Mycobacterium*
ulcerans in a Buruli ulcer mouse model. *Nat. Commun.* **10**, 524 (2019).

25   Liu, Y. *et al.* Assessment of clofazimine and TB47 combination activity against
Mycobacterium abscessus using a bioluminescent approach. *Antimicrob. Agents*
Chemother. **64**, e01881-01819 (2020).

26   Gutiérrez, A. V., Viljoen, A., Ghigo, E., Herrmann, J.-L. & Kremer, L.
Glycopeptidolipids, a double-edged sword of the *Mycobacterium abscessus*
complex. *Front. Microbiol.* **9**, 1145 (2018).

27   Jankute, M., Grover, S., Birch, H. L. & Besra, G. S. Genetics of mycobacterial
arabinogalactan and lipoarabinomannan assembly. *Microbiol Spectr* **2**, MGM2-
2013 (2014).

28   Zhang, M., Gong, J., Lin, Y. & Barnes, P. F. Growth of virulent and avirulent
Mycobacterium tuberculosis strains in human macrophages. *Infect. Immun.* **66**,
794-799 (1998).

29   Lu, X. Y. *et al.* Pyrazolo[1,5-]pyridine inhibitor of the respiratory cytochrome
complex for the treatment of drug-resistant tuberculosis. *ACS Infect. Dis.* **5**, 239-
249 (2019).

30   Zhang, T., Li, S. Y. & Nuermberger, E. L. Autoluminescent *Mycobacterium*
tuberculosis for rapid, real-time, non-invasive assessment of drug and vaccine
efficacy. *PLoS ONE* **7**, e29774 (2012).

31   Bergval, I. *et al.* Pre-existing isoniazid resistance, but not the genotype of
Mycobacterium tuberculosis drives rifampicin resistance codon preference *in*
vitro. PLoS ONE **7**, e29108 (2012).

32   Wolf, N. M. *et al.* High-resolution structure of ClpC1-rufomycin and ligand
binding studies provide a framework to design and optimize anti-tuberculosis
leads. *ACS Infect. Dis.* **5**, 829-840 (2019).

33   Eberhardt, J., Santos-Martins, D., Tillack, A. F. & Forli, S. AutoDock Vina 1.2.0:
New docking methods, expanded force field, and python bindings. *J. Chem. Inf.*
Model. **61**, 3891-3898 (2021).

34   Wang, Y. *et al.* Pubchem bioassay: 2017 update. *Nucleic Acids Res.* **45**, D955-
D963 (2017).

35   Makafe, G. G. *et al.* Role of the Cys154Arg substitution in ribosomal protein L3
in Oxazolidinone resistance in *Mycobacterium tuberculosis*. *Antimicrob. Agents*
Chemother. **60**, 3202-3206 (2016).

36   Vasudevan, D., Rao, S. P. & Noble, C. G. Structural basis of mycobacterial
inhibition by cyclomarin A. *J. Biol. Chem.* **288**, 30883-30891 (2013).

37   Chhotaray, C. *et al.* Advances in the development of molecular genetic tools for
Mycobacterium tuberculosis. *J Genet Genomics* (2018).

38   Yan, M. *et al.* CRISPR-Cas12a-assisted recombineering in bacteria. *Appl.*
Environ. Microbiol. **83**, e00947-00917 (2017).

39   Wang, S. *et al.* Amino acid 17 in QRDR of Gyrase A plays a key role in
fluoroquinolones susceptibility in mycobacteria. *Microbiol Spectr* **11**, e0280923
(2023).

40   Dziedzic, R. *et al. Mycobacterium tuberculosi*s ClpX interacts with FtsZ and
interferes with FtsZ assembly. *PLoS ONE* **5**, e11058 (2010).

41   Bai, J., Chi, M., Hu, Y., Hao, M. & Zhang, X. Construction and biological
characteristics of ClpC and ClpX knock-down strains in *Mycobacterium*
smegmatis. *China Biotechnology* **41**, 13-22 (2021).

42   Rock, J. M. *et al.* Programmable transcriptional repression in mycobacteria using
an orthogonal CRISPR interference platform. *Nat. Microbiol.* **2**, 1-9 (2017).

43   Vivoli, M., Novak, H. R., Littlechild, J. A. & Harmer, N. J. Determination of
protein-ligand interactions using differential scanning fluorimetry. *J. Vis. Exp.*,
e51809 (2014).

44   Kim, J. H. *et al.* Protein inactivation in mycobacteria by controlled proteolysis
and its application to deplete the beta subunit of RNA polymerase. *Nucleic Acids*
Res. **39**, 2210-2220 (2011).

45   McGinness, K. E., Baker, T. A. & Sauer, R. T. Engineering controllable protein
degradation. *Mol. Cell* **22**, 701-707 (2006).

46   Gopal, P. *et al.* Pyrazinamide triggers degradation of its target aspartate
decarboxylase. *Nat. Commun.* **11**, 1661 (2020).

47   Raju, R. M. *et al.* Post-translational regulation via Clp protease is critical for
survival of *Mycobacterium tuberculosis*. *PLoS Pathog.* **10**, e1003994 (2014).

48   Shetye, G. S., Choi, K. B., Kim, C. Y., Franzblau, S. G. & Cho, S. *In vitro*
profiling of antitubercular compounds by rapid, efficient, and nondestructive
assays using autoluminescent *Mycobacterium tuberculosis*. *Antimicrob. Agents*
Chemother. **65**, e0028221 (2021).

49   Botella, L., Vaubourgeix, J., Livny, J. & Schnappinger, D. Depleting

Mycobacterium tuberculosis of the transcription termination factor Rho causes
pervasive transcription and rapid death. *Nat. Commun.* **8**, 14731 (2017).

50   Jin, D. J. *et al.* Effects of rifampicin resistant *rpoB* mutations on antitermination
and interaction with *nusA* in *Escherichia coli*. *J. Mol. Biol.* **204**, 247-261 (1988).

51   Sklar, J. G., Makinoshima, H., Schneider, J. S. & Glickman, M. S. *M.*
tuberculosis intramembrane protease Rip1 controls transcription through three
anti☐sigma factor substrates. *Mol. Microbiol.* **77**, 605-617 (2010).

52   Unissa, A. N., Subbian, S., Hanna, L. E. & Selvakumar, N. Overview on
mechanisms of isoniazid action and resistance in *Mycobacterium tuberculosis*.
Infect. Genet. Evol. **45**, 474-492 (2016).

53   Hameed, H. A. *et al.* Molecular targets related drug resistance mechanisms in
MDR-, XDR-, and TDR-*Mycobacterium tuberculosis* strains. *Front. Cell. Infect.*
Microbiol. **8**, 114 (2018).

54   Ramaswamy, S. V. *et al.* Single nucleotide polymorphisms in genes associated
with isoniazid resistance in *Mycobacterium tuberculosis*. *Antimicrob. Agents*
Chemother. **47**, 1241-1250 (2003).

55   Chen, P. & Bishai, W. R. Novel selection for isoniazid (INH) resistance genes
supports a role for NAD^+-binding proteins in mycobacterial INH resistance.
Infect. Immun. **66**, 5099-5106 (1998).

56   He, L. *et al. ubiA* (*Rv3806c*) encoding DPPR synthase involved in cell wall
synthesis is associated with ethambutol resistance in *Mycobacterium tuberculosis*.
Tuberculosis (Edinb) **95**, 149-154 (2015).

57   Gao, Y. *et al.* Ultra-short-course and intermittent TB47-containing oral regimens
produce stable cure against Buruli ulcer in a murine model and prevent the
emergence of resistance for *Mycobacterium ulcerans*. *Acta Pharm. Sin. B* **11**,
738-749 (2021).

58   Zhou, B. *et al.* Rufomycins or Ilamycins: naming clarifications and definitive
structural assignments. *J. Nat. Prod.* **84**, 2644-2663 (2021).

59   Ziemski, M., Leodolter, J., Taylor, G., Kerschenmeyer, A. & Weber-Ban, E.
Genome-wide interaction screen for *Mycobacterium tuberculosis* ClpCP protease
reveals toxin-antitoxin systems as a major substrate class. *FEBS J.* **288**, 111-126
(2021).

60   Bordes, P. & Genevaux, P. Control of toxin-antitoxin systems by proteases in
Mycobacterium tuberculosis. *Front Mol Biosci* **8**, 691399 (2021).

61   Yamada, Y. & Dick, T. Mycobacterial caseinolytic protease gene regulator ClgR
is a substrate of caseinolytic protease. *mSphere* **2**, e00338-00316 (2017).

62   Thibault, G. *et al.* Specificity in substrate and cofactor recognition by the N-
terminal domain of the chaperone ClpX. *Proc Natl Acad Sci U S A* **103**, 17724-
17729 (2006).

63   Ju, Y. *et al.* The gene *MAB_2362* is responsible for intrinsic resistance to various
drugs and virulence in *Mycobacterium abscessus* by regulating cell division.
Antimicrob. Agents Chemother. **69**, e0043324 (2025).

64   Jiang, H. *et al.* One-step engineering of a stable, selectable marker-free
autoluminescent *Acinetobacter baumannii* for rapid continuous assessment of

drug activity. *J. Microbiol. Biotechnol.* **29**, 1488-1493 (2019).

65   Tian, X. *et al.* Rapid visualized assessment of drug efficacy in live mice with a
selectable marker-free autoluminescent *Klebsiella pneumoniae*. *Biosens.*
Bioelectron. **177**, 112919 (2021).

66   Wang, S. *et al.* Rapid, serial, non-invasive quantification of *Pseudomonas*
aeruginosa in live mice with a selectable marker-free autoluminescent strain.
Biosens. Bioelectron. **165**, 112396 (2020).

67   Yu, W. *et al.* TB47 and clofazimine form a highly synergistic sterilizing block in
a second-line regimen for tuberculosis in mice. *Biomed. Pharmacother.* **131**,
110782 (2020).

68   Liu, P. *et al.* Design and synthesis of novel pyrimidine derivatives as potent
antitubercular agents. *Eur. J. Med. Chem.* **163**, 169-182 (2019).

69   Khadiullina, R. *et al.* Assessment of thermal stability of mutant p53 proteins via
differential scanning fluorimetry. *Life (Basel)* **13**, 31 (2022).

70   Zhou, B. *et al.* Structural insights into bortezomib-induced activation of the
caseinolytic chaperone-protease system in *Mycobacterium tuberculosis*. *Nat.*
Commun. **16**, 3466 (2025).

Acknowledgments

We acknowledge the team of professor Yicheng Sun from the Institute of
Pathogenic Biology, Chinese Academy of Medical Sciences for kindly sending the
pJV53-Cpf1 and pCR-Zeo plasmids as the tool of gene editing. This work was
supported by fundings from National Key R&D Program of China
(2021YFA1300900), National Natural Science Foundation of China (81973372,
32300152, 82022067, 22037006), China Postdoctoral Science Foundation
(2022M723164), Guangdong Provincial Basic and Applied Basic Research Fund
(2022A1515110505, 2019B030302004), Grant of State Key Lab of Respiratory
Disease (SKLRD-Z-202414, SKLRD-Z-202301), Open Program of Shen Zhen Bay
Laboratory (SZBL2021080601006) and Nansha District Science and Technology Plan
Project (NSJL202102).The funders had no role in study design, data collection and
analysis, decision to publish, or preparation of the manuscript.

Author contributions

T.Z., J.J., and X.X. supervised the project; T.Z., Y.G., and C.F. designed the
whole project and most of the experiments; J.M. and C.S. prepared and characterized
the ilamycin E and ilamycin F; Y.G. and C.F. performed the susceptibility testing and
resistant mutant screening; B.Z. and J.J. contributed to molecular docking simulation;

C.F., X.T., and J.H. constructed overexpression and gene-editing strains and peformed

the susceptibility testing; C.F. and X.H. performed the gene-scilencing experiments;

Y.G. and B.Z. contributed to protein expression and mode of action experiments; C.F.

and H.Z. contributed to the microscopy results; Y.G. performed the statistical analysis;

Y.G., F.C., and T.Z. wrote the manuscript; H.M.A., J.L., J.M., X.C., N.Z., X.X., J.J.,

and T.Z. revised the manuscript.

Competing interests

The authors have no competing interests to declare that are relevant to the

content of this article.

968