Wake EEG oscillation dynamics reflect both sleep pressure and brain maturation across childhood and adolescence

Sophia Snipes,[1,2] Valeria Jaramillo,[1,3,4,5] Elena Krugliakova,[1,6] Carina Volk,[1,7] Melanie Furrer,[1] Mirjam Studler,[1,8] Monique LeBourgeois,[9,10,11] Salome Kurth,[1,12,13] Oskar G. Jenni,[1] Reto Huber[1,14]

[1] *Child Development Center, University Children's Hospital Zurich, University of Zurich, Zurich, Switzerland*
[2] *Paris Brain Institute, Sorbonne Université, Inserm-CNRS, Paris, France* [3] *School of Psychology, University of Surrey, Guildford, UK*
[4] *Surrey Sleep Research Centre, Faculty of Health and Medical Sciences, University of Surrey, Guildford, United Kingdom*
[5] *UK Dementia Research Institute Care Research and Technology Centre, Imperial College London and the University of Surrey, Guildford, UK*
[6] *Donders Institute, Radboud University Medical Center, Nijmegen, the Netherlands*
[7] *Center for MR Research, University Children's Hospital Zurich, Switzerland*
[8] *Department of Social Neuroscience and Social Psychology, Institute of Psychology, University of Bern, Bern, Switzerland*
[9] *University of Colorado at Boulder, Department of Integrative Physiology, Boulder, Colorado, USA*
[10] *The Warren Alpert Medical School of Brown University, Department of Psychiatry and Human Behavior, Providence, Rhode Island, USA*
[11] *In memoriam*
[12] *Department of Psychology, University of Fribourg, Fribourg, Switzerland*
[13] *Department of Pulmonology, University Hospital Zurich, Zurich, Switzerland*
[14] *Department of Child and Adolescent Psychiatry and Psychotherapy, Psychiatric Hospital, University of Zurich, Switzerland*

Corresponding author: Sophia Snipes (publishing@sophia.science)

1 ABSTRACT

An objective measure of brain maturation is highly insightful for monitoring both typical and atypical development. Slow wave activity, recorded in the sleep electroencephalogram (EEG), reliably indexes age-related changes in sleep pressure as well as deficits related to developmental disorders such as attention-deficit hyperactivity disorder (ADHD). We aimed to determine whether wake EEG measured before and after sleep could index the same developmental changes in sleep pressure, using data collected from 163 participants 3-25 years old. We analyzed age- and sleep-dependent changes in two measures of oscillatory activity, amplitudes and density, as well as two measures of aperiodic activity, offsets and exponents. We then compared these wake measures to sleep slow wave amplitudes and slopes. Finally, we compared wake EEG in children with ADHD (N=58) to neurotypical controls. Of the four wake measures, only

oscillation amplitudes consistently exhibited the same changes as sleep slow waves. Wake amplitudes decreased with age, decreased after sleep, and this overnight decrease decreased with age. Furthermore, wake amplitudes were significantly related to both sleep slow wave amplitudes and slopes. Wake oscillation densities decreased overnight in children but increased overnight in adolescents and adults. Aperiodic offsets decreased linearly with age, decreased after sleep, and were significantly related to sleep slow wave amplitudes. Aperiodic exponents also decreased with age, but increased after sleep. No wake measure showed significant effects of ADHD. Overall, our results indicate that wake oscillation amplitudes, and to some extent aperiodic offsets, behave like sleep slow waves across sleep and development. At the same time, overnight changes in oscillation densities independently reflect some yet-unknown shift in neural activity around puberty.

Keywords: development, EEG, sleep, synaptic homeostasis hypothesis, aperiodic activity, oscillation bursts

1 INTRODUCTION

The EEG is one of few tools available to study the human developing brain already from birth (Korotchikova et al., 2009). It is non-invasive, relatively cheap, and provides a real-time readout of neuronal activity. It is an incredibly rich signal, with the potential as a prognostic and diagnostic tool for both typical development and disease. Sleep EEG, and in particular slow wave activity (0.5-4 Hz) during NREM sleep, has proven especially sensitive to brain maturation (Campbell & Feinberg, 2009) and developmental disorders such as ADHD (Furrer et al., 2019). This is because slow wave activity reflects the overall synchronicity of the brain, which decreases with age following decreasing synaptic density across adolescence (Campbell & Feinberg, 2009; Huttenlocher, 1979; Jenni & Carskadon, 2004), and may be lower in ADHD due to reduced cortical thickness (Shaw et al., 2006). Furthermore, slow wave activity is greater in occipital regions in younger children, and greater in frontal regions in adolescents and adults (Kurth et al., 2010), possibly reflecting the slower maturation of higher order association areas (Shaw et al., 2008).

Introduction

In addition to these developmental phenomena, sleep slow waves reflect the buildup
and dissipation of homeostatic sleep pressure, increasing following wake and decreas-
ing during sleep (Borbély, 1982). These hourly changes in sleep pressure detected
through slow wave activity are hypothesized to reflect synaptic *plasticity*: synaptic
strength increases with wake and daytime learning, and decreases during sleep (Tononi
& Cirelli, 2003, 2014). Generally, plasticity decreases with brain maturation across
childhood and adolescence and this is reflected in decreases in the overnight *changes*
in slow waves with age (Jaramillo et al., 2020). Thus, both absolute slow wave activity
and changes in slow wave activity are markers of brain development, reflecting average
synaptic density and synaptic plasticity respectively. While these predictions have been
repeatedly tested and validated in sleep, their consequences on the wake EEG have not
been as systematically investigated. Previous research has found correlations between
wake and sleep EEG power (Finelli et al., 2000), however, greater insights can be de-
rived from more specific analyses.

The EEG is made up of both *periodic* activity and *aperiodic* activity (Figure 1A, Do-
noghue et al., 2020). Periodic activity refers to oscillations, which appear as quasi-
gaussian bumps in the power spectrum at their corresponding frequency. Instead, ape-
riodic activity is a form of background "noise," producing the characteristic 1/f curve in
the EEG power spectrum. Aperiodic activity is defined by its *offset* (reflecting the overall
aperiodic power) and *exponent* (the steepness of the curve). Changes in exponent in
particular are hypothesized to reflect alterations in excitatory/inhibitory balance of neu-
ronal activity (Brake et al., 2024; Gao et al., 2017), and this could explain the multitude
of conditions which affect aperiodic signals.

Aperiodic exponents reflect levels of consciousness, becoming progressively steeper
with increasing sleep depth (Lendner et al., 2020, 2023; Schneider et al., 2022), anes-
thesia (Colombo et al., 2019), and disorders of consciousness (Colombo et al., 2023;
Maschke et al., 2023). Additionally, exponents become progressively shallower during
sleep, reflecting the dissipation of sleep pressure (Bódizs et al., 2024; Horváth et al.,
2022). Brain maturation also strongly affects exponents and offsets, becoming shal-
lower and lower from childhood to adulthood (Cellier et al., 2021; Hill et al., 2022; Trön-
dle et al., 2022), with age and sleep depth interacting (Favaro et al., 2023). Lastly, dif-

ferences in aperiodic exponents and offsets have been found in children with ADHD

compared to neurotypical controls (Ostlund et al., 2021; Robertson et al., 2019).

Periodic activity, like aperiodic activity, can be estimated from spectral power by simply

subtracting the aperiodic activity from overall power, giving *periodic power* (Figure 1A).

However this misses two important independent changes that can happen to oscilla-

tions: they can change in *amplitude*, and they can change in *density* (Figure 1B). The

amplitudes of oscillations reflect the synchronicity of the oscillating neuronal popula-

tion, and that synchronicity is determined both by the number of neurons in phase with

each other and the strength of their synaptic connections. Whether an oscillation oc-

curs at all (i.e., density) will instead depend on the activity of "pacemaker" interneurons

which entrain a population of neurons to the same rhythm (Le Bon-Jego & Yuste, 2007;

Perkel et al., 1964), and this pattern of rhythmic firing will be in service of some underly-

ing function that will come and go as needed. In short, amplitudes reflect synchronicity

and densities reflect activity.

Since oscillation amplitudes reflect synchronicity, this means they should reflect the

same information as slow waves measured during sleep. Sleep slow wave activity in-

creases along a saturating exponential function relative to the prior duration of wake-

fulness, increasing rapidly after only a few hours awake and increasing less and less

with further hours awake (Borbély, 1982; Dijk et al., 1987). We found that wake oscilla-

tion amplitudes follow the same trajectory across 24 hours awake (Snipes et al., 2023).

Instead, alpha oscillation densities (8-12 Hz) decreased, supporting the specificity of

the effect to amplitudes and masking the effect when measuring spectral power.

Given these results, we hypothesized that wake oscillation amplitudes should behave

like sleep slow waves also across development: absolute amplitudes should decrease

with age, and overnight changes in amplitude should decrease with age. Likewise,

changes in amplitude should also manifest an anterior-posterior gradient with age:

larger amplitudes in occipital regions in children and larger in frontal regions in adults.

Furthermore, given that children with ADHD have lower sleep slow wave activity, they

should likewise have lower wake amplitudes. In short, if both wake amplitudes and

sleep slow waves are supposed to reflect brain-wide synchronicity, then wake ampli-

tudes should also reflect sleep pressure, brain development, and the pathophysiology

of ADHD. We further hypothesized that all these effects would be specific to ampli-

tudes, with other wake EEG parameters changing independently with sleep and age.

Finally, we expected that wake amplitudes, more than any other wake EEG measure,

should directly correlate with specific sleep markers of synaptic plasticity: slow wave

amplitudes and slopes (the steepness of the waves) measured at the beginning and end

of the night (Esser et al., 2007; Jaramillo et al., 2020; Riedner et al., 2007; Vyazovskiy et

al., 2009). Slopes in particular are thought to reflect changes in synaptic strength. The

steepness of a slow wave is determined by the rate at which neuronal populations syn-

chronize and desynchronize firing, and increased synaptic strength will increase the

speed of such synchronization (Vyazovskiy et al., 2009). Therefore, any wake measure

of sleep pressure and synaptic plasticity should correlate with both sleep slow wave

amplitudes and especially sleep slow wave slopes.

To answer these questions, we analyzed data collected from previous studies at the

University Children's Hospital of Zurich, with high-density wake EEG recordings meas-

ured the evening before and morning after a night of sleep. The final dataset included

105 neurotypical participants from the ages of 3.5 to 25, and 58 children with ADHD.

From 44 neurotypical participants, the wake EEG was further compared to their sleep

EEG.

Figure 1: Wake EEG measures. A: Measures based on spectral power. Given the nature of power, it is traditionally
analyzed log-transformed to have more normally distributed values. These are then aggregated into bands. Here, we
focus on classical wake EEG bands: theta (4-7 Hz, yellow), alpha (8-11 Hz, orange) and low beta (12-16 Hz, red), with
gaps between bands to avoid overlapping information. The example comes from a 15-year-old male participant,
used for the entire figure. The EEG signal is composed of aperiodic "background activity" (gray signal in B) and oscil-

latory activity (colored signal in B). When plotting the power spectrum on a log-log scale, a line can be fitted to the aperiodic component of the signal, which can be subtracted from the whole spectrum, leaving behind only periodic power. The aperiodic line can then be quantified by its offset (where it intersects 0 on the log-log scale, i.e. the log power at 1 Hz), and its exponent (how tilted it is). Thus, the power spectrum provides four measures: log-transformed power, periodic power, exponent, and offset of the aperiodic signal. **B**: Cycle-by-cycle analysis is used to detect bursts of oscillations by identifying sections of the EEG signal that show periodic activity (colored), relative to the aperiodic background activity (gray). Once bursts are detected, there are two main parameters to quantify them: density (how much of the signal in time contains an oscillation) and amplitude (average peak-to-peak voltage of each oscillation). Densities are expressed in percentage, and when pooling bursts detected in all channels, they can easily exceed 100%, as the same burst will appear in multiple channels and multiple bursts can co-occur. Amplitudes are in microvolts. The EEG trace was stitched together for illustrative purposes. N.B. Beta periodic power is lower than alpha periodic power (in panel A), but their densities (in B) are roughly the same; this is because periodic power is also influenced by the lower amplitudes of low beta compared to alpha. The densities of other participants are provided in Suppl. Figure 1-1.

2 METHODS

2.1 Datasets

The data for this manuscript was assembled from previous studies conducted between 2008 and 2021. The participant demographics of each dataset are in Table 1. In total, we included 163 participants between the ages 3.5 and 24.7, 38% female, 7% left-handed. Of these, 36% were diagnosed with ADHD at the department for Child and Adolescent Psychiatry at the University of Zurich, the outpatient clinic of the Child Development Center, and at private children's clinics in Zurich Oerlikon. Patients were not excluded based on medication status, and therefore were a mixture of medicated, previously medicated, and unmedicated (see Table 2, and (Furrer et al., 2019; Ringli et al., 2013)). Otherwise, all participants were screened by telephone such that they all were completely healthy, took no (other) medication, had no (other) comorbidities, and were good sleepers. All participants were recruited from canton Zurich, Switzerland, and recorded at the University Children's Hospital of Zurich, except for the children from 3.5 to 8 (Dataset2009, N=11), who were recruited in Providence, RI, USA, and recorded at home. Sleep time was determined by their individual preferred sleep and wakeup time, which they had to maintain the week prior to each measurement. Wake measurements were done just before going to sleep, and ~30 minutes after waking up. 115 participants had 2 sessions, spaced at least 1 week apart, both included in these analyses. Depending on the dataset, different paradigms were used involving different wake tasks (described below) and there could be 1-4 tasks at each time point (see Table 1). Every set of tasks for each dataset was repeated in the evening and in the morning. In total, 1243 recordings were included in these analyses lasting on average 4.2 min (task time + buff-

er − artefacts), with a standard deviation of 2.5 min. The data (averaged across chan-

nels) for all participants is provided in Suppl. Data 1.

Informed consent was obtained from all adult participants, and from the legal guardi-

ans of all children below 14, as well as from adolescent participants 14-18. All studies

were approved by the local ethics committees and performed according to the declara-

tion of Helsinki.

| | N | Female | Left-handed | ADHD | Age range | Mean age | Paradigm | Sessions | Recordings |
	#	%	%	%	(years)	(years)		#	#
Dataset2008	38	32	0	0	8.7-23.4	14.1 (3.8)	Adaptation	2	7.3/8
Dataset2009	11	73	0	0	3.5-8.0	5.6 (1.5)	Oddball	1	1.8/2
Dataset2010	28	21	14	100	9.7-16.3	12.7 (1.9)	Adaptation	1	2.8/4
Dataset2016	18	44	0	0	18.4-24.7	21.6 (2.1)	Oddball	2	3.9/4
Dataset2017	42	43	17	36	8.1-17.6	12.2 (2.7)	Attention	2	12.7/16
Dataset2019	26	38	0	58	8.8-16.8	11.4 (2.0)	Attention	2	10.1/12
All	163	38	7	36	3.5-24.7	13.2 (4.4)			

Table 1: All participants' demographics, split by dataset. The year for each dataset indicates the beginning of data
collection. *N* indicates the number of participants. After the mean age of each dataset, the standard deviation is
provided in parentheses. *Paradigm* indicates which set of wake tasks were recorded. *Sessions* indicate the number
of sessions expected for each dataset, although in practice due to dropouts, some participants only completed 1.
Recordings indicate the average number of recordings per participant compared to the total number of recordings
expected by the experimental paradigm. Recordings were missing either because they contained too many artefacts
or were omitted entirely during data collection.

| | N | Female | Left-handed | Age range | Mean age | Oddball |
	#	%	%	(years)	(years)	%
Medicated in the past	6	17	0	9.7-14.8	12.5 (2.2)	33
Unmedicated	14	21	14	9.5-15.3	11.3 (1.6)	64
Medication the day before	22	23	14	8.7-16.3	12.3 (2.2)	45
Medication the day of	13	15	0	9.0-16.1	12.3 (2.0)	54
All patients	55*	20	9	8.7-16.3	12.1 (2.0)	51
Controls	105	47	6	3.5-24.7	13.8 (5.2)	58

14 **Table 2: ADHD demographics, split by patient status.** For each patient group, the *Oddball* column indicates how
15 many came from a dataset performing the oddball task (rather than the attention tasks). *For 3 patients, medication
16 status was missing, therefore the true total is 58.

2.1.1 Oddball & motor adaptation paradigms

95 participants (Dataset2008, Dataset2009, Dataset2010, Dataset2016) performed an auditory oddball task during their wake EEG. The task lasted 4 minutes and was performed in the evening just before going to bed and in the morning ~30 minutes after waking up. The task involved 300 tones at ~80 dB, with an interstimulus interval of 0.8 s. A random 10% of stimuli were targets to which the participant had to push a button in response. For the youngest children (Dataset2009), the 4-minute task was split into 2 segments, and for the adults (Dataset2016) the task was extended to 6 minutes.

66 of these participants (Dataset2008, Dataset2010) also performed a half-hour visuo-motor adaptation task (Ghilardi et al., 2000) followed by a second oddball. One dataset (Dataset2008) also included a second session with a control visuomotor task (no adaptation), counterbalanced with the motor adaptation task. The motor tasks were not included in this analysis, because they further differed from evening to morning. For more details on the adaptation task see Wilhelm et al. (2014). The youngest (Dataset2009) and oldest (Dataset2016) participants only conducted one oddball and no motor task. Dataset2016 also had two sessions, one night with phase-targeted auditory stimulation during NREM sleep and the other sham. The experimental conditions did not affect the wake EEG (data not shown).

The sleep data from these participants has been published (Buchmann et al., 2011; Furrer et al., 2019, 2020; Jaramillo et al., 2020; Kurth et al., 2010; Ringli et al., 2013; Volk et al., 2018; Wilhelm et al., 2014), as has a subset of the wake EEG data (Fattinger et al., 2017).

2.1.2 Attention paradigm

participants (Dataset2017, Dataset2019) performed three tasks with a focus on attention. These were studies investigating the relationship between slow waves, behavior, and MR spectroscopy (Jaramillo et al., 2020; Volk et al., 2019). This included two sessions to compare the effects of phase targeted auditory stimulation on slow waves in sleep (sham and stimulation; data currently unpublished). The wake tasks were part of the Test Battery for Attentional Performance (TAP) (Zimmermann & Fimm, 2012), which included 2 minutes of a visual Go/No-Go task (respond to 1 stimulus, withhold

response to another), 5.5 minutes of a visual and acoustic Alertness task, and then 2

1.5-minute Fixation recordings. For one dataset (Dataset2019) the Go/No-Go task was

adapted using cartoon images (extended to 12 minutes), but only 1 Fixation recording

was measured (lasting 2 minutes).

2.2   EEG recordings and preprocessing

All datasets were measured using 128 channel EGI Geodesic Sensor nets and EGI am-

plifiers (Electrical Geodesics Inc., Eugene, OR, USA). Wake recordings were done with

Cz reference, 1000 Hz sampling rate, and impedances kept below 50 kOhm. All anal-

yses were performed in MATLAB 2023b, with the EEGLAB toolbox v2023.1 (Delorme &

Makeig, 2004), the FOOOF/specparam toolbox v1.1.0 (Donoghue et al., 2020), and cus-

tom scripts.

EEG data was first mean-centered, then lowpass filtered at 40 Hz (EEGLAB's

pop_eegfiltnew function) and notch-filtered at either 50 or 60 Hz (Dataset2009) along

with subsequent harmonics. The data was downsampled to 250 Hz, then highpass-

filtered over 0.5 Hz (Kaiser filter, stopband = 0.25 Hz, stopband attenuation = 60, pass-

band ripple = 0.05).

Artifacts were removed with a fully automated procedure. Movement and other large

artifacts were detected in data filtered between 1 and 40 Hz, in 3 s segments. A seg-

ment was labeled a "major artifact" if it exceeded 500 μV, or a "minor artifact" if the

correlation with neighboring channels was below .3. Major artifacts were always re-

moved, either by removing all data in all channels during those 3 s, or removing the en-

tire channel with such an artifact, depending on which (channel or segment) removed

the least amount of clean data. Minor artifacts were removed in a similar way, removing

iteratively either the channel with the most artifactual segments, or the segments with

the most artifactual channels, until all channels and all segments had at most 30% of

the data containing a minor artifact. Flat channels were removed using EEGLAB's

clean_artifacts function. The missing Cz channel was added (as a vector of zeros), then

the data was average referenced. Physiological artifacts (blinks, eye movements, mus-

cle tone, heartbeat) were removed with independent component analysis (ICA), with

components calculated separately as described in the next section. After these were

removed, a second pass was conducted using EEGLAB's *clean_windows* function

(MaxBadChannels = .3, PowerTolerances = [-inf, 12]), then bad segments/channels still

containing amplitudes over 140 μV were removed, and finally EEGLAB's

clean_channels_nolocs was applied (MinCorrelation = .5, IgnoredQuantile = .1, Max-

BrokenTime = .5). Recordings for which more than 25 channels were removed, or which

had less than 1 minute of data, were excluded from analysis. In a last step, EEG chan-

nels were interpolated, for a total of 123 channels, excluding the external electrodes

(49 56, 107, 113) and the face electrodes (126, 127).

2.2.1 Automatic detection and removal of physiological artefacts using ICA

For ICA, a separate copy of the EEG data was preprocessed as previously described,

however the data was filtered between 2.5 and 100 Hz, and downsampled to 500 Hz.

Automatically detected bad channels and bad time windows were removed, an empty

Cz channel added, and then the data was re-referenced to the average of all channels.

EEGLAB's *runica* function was run with principal component analysis (PCA) rank reduc-

tion. Then, components were automatically classified with EEGLAB's *iclabel*, as either

brain, muscle, eye, heart, line, channel noise, or other. This function provides a proba-

bility score for each label from 0 to 1, so the label with the largest score for each com-

ponent was taken. Components classified as muscle, eye, or heart were removed. Of

the remaining noise classifications (line, channel, other), due to poor classification ac-

curacy, an additional step was implemented. Spectral power was calculated for each

component (*pwelch*, 4 s Hanning windows, 50% overlap), then smoothed over 5 Hz

(*lowess*) to facilitate model fitting. Unlike for the main analyses, the periodic signal was

not of interest, therefore stronger smoothing was possible. The *specparam* algorithm

(Donoghue et al., 2020; Ostlund et al., 2022) was applied to the power spectrum be-

tween 8 and 30 Hz to extract aperiodic exponents. Components for which the spectral

exponent was shallower than 0.5 (so almost flat or even tilted positive), were consid-

ered noise and therefore excluded, as they reflected either muscle activity or other non-

physiological signals. Using the manually labeled components in an independent adult

dataset (Snipes et al., 2022), we confirmed that this procedure was sufficiently compa-

rable to human detection of artifactual components. We further confirmed that the

outcome matched human component classification in a small subset of the children's

data. However, considering the trend towards 0.5 exponents observed in Figure 2, for

future datasets with older participants we would recommend a lower threshold.

For the Dataset2009 cohort of <8-year-olds, given how little data there was and how

many more movement artifacts, we chose to apply to same manual artifact rejection as

in Snipes et al. (2022) to preserve as much data as possible.

2.3   Burst detection

Bursts of oscillations were detected using cycle-by-cycle analysis (Cole & Voytek, 2019)

implemented in MATLAB (Snipes et al., 2023). Bursts were detected with the same

thresholds as in Snipes et al. (2024). Briefly, EEG was narrow-band-pass filtered in over-

lapping ranges (2-6 Hz, 4-8 Hz...), from which zero-crossings were detected. Then, in

the broadband filtered data (0.5-40 Hz), peaks were identified between the zero-

crossings, and a cycle was considered an oscillation from positive-to-positive peak. A

minimum number of consecutive cycles must meet a set of criteria (monotonicity, peri-

od consistency, amplitude consistency, shape consistency, etc.) for this to be consid-

ered a burst. Importantly, amplitude itself is never used as a threshold, as this would

create a greater dependency between amplitude and density (such that a decrease in

an amplitude threshold would result in an automatic increase in density).

Three sets of criteria were used. The first aimed to detect bursts relying on many low-

threshold criteria (frequency in range of narrowband filter; PeriodConsistency = .5; Am-

plitudeConsistency = .4; FlankConsistency = .5; ShapeConsistency = .2; Monotonic-

tyInTime = .4; MonotonicityInAmplitude = .4; ReversalRatio = .6; MinCycles = 4). The

second had fewer criteria with intermediate thresholds but a higher minimum number

of cycles (PeriodConsistency = .6; AmplitudeConsistency = .6; MonotonicityInAmplitde

= .6; FlankConsistency = .6; MinCycles = 5). The third set had fewer criteria but stricter

monotonicity thresholds (frequency in range of narrowband filter; PeriodConsistency =

.7; FlankConsistency = .3; MontonocityInAmplitude = .9; MinCycles = 3). These criteria

were chosen a-priori based on manual tuning of the burst detection on an independent

dataset of wake EEG in adults during sleep deprivation.

After bursts were detected in each channel separately, they were grouped into clusters

when they occurred simultaneously in multiple channels with roughly the same fre-

quency. The frequency of bursts was calculated as the inverse of the average distance

between negative peaks (1/period). Bursts for which the shorter one overlapped at least

50% of the longer one, and were within 1 Hz of each other, were considered part of the

same burst cluster. Bursts identified separately in each channel were used for all the

topographies, otherwise burst clusters were used to reduce the effect of burst globality

(spread across the scalp) on measures of density.

2.3.1   Oscillation measures

Oscillation amplitudes were calculated as the average negative-to-positive peak volt-

age difference for all cycles involved in all bursts, with units in microvolts (µV). *Oscilla-*

tion densities were calculated as the percentage of the recording occupied by bursts

(sum of all the bursts' durations divided by the duration of the recording). This accounts

for both the duration and overall quantity of bursts in the recording. The average dura-

tion and number of bursts can be affected by the background aperiodic signal which

can break up sustained oscillations into smaller bursts (Tal et al., 2020), therefore burst

density was preferable. When calculating across multiple channels (e.g. Figure 2), os-

cillation density could easily exceed 100%, as burst clusters in different frequency

ranges often co-occur. When combining densities across multiple frequency bands,

bursts were pooled rather than averaged (sum of the durations of all the bursts of any

frequency, divided by the duration of the recording). In our previous publication (Snipes

et al., 2023), we referred to oscillation densities as "quantities", however this term did

not properly account for the normalization in time.

2.3.2   The choice of frequency bands

Only bursts between 2 and 16 Hz were detected. Below 4 Hz very few bursts could be

identified, therefore only bursts above 4 Hz were included in the analysis. Bursts over

16 Hz could be detected, but with higher false-positive rates, as determined by visual

inspection. The choice of cutoff at 16 Hz was done arbitrarily a-priori to capture alpha

(8-12 Hz) with generous padding. The division of bands for Figure 6 and Figure 7 was

done using conventional bands (Pernet et al., 2020) with 1 Hz gaps to reduce overlap-

ping information due to the drift in peak frequencies across individuals and ages: theta

(4-7 Hz), alpha (8-11 Hz) and low beta (12-16 Hz). The inclusion of low beta was done a-

posteriori based on results observed in Figure 5B.

Many researchers advocate for the use of an individual alpha frequency (IAF) to define

frequency ranges, especially when analyzing data across development (Bazanova &

Vernon, 2014; Klimesch, 1999; Ostlund et al., 2022; Tröndle et al., 2022). This is done by

selecting the peak alpha frequency separately for each individual and defining the band

around this peak. The shift in IAF with age makes a strong case for such an approach

(Freschl et al., 2022). The main problem with using IAF is that it assumes the largest

oscillation will be functionally the same for all participants. Given that our participants

displayed large heterogeneity in the number and amplitude of frequency peaks within

the 2-16 Hz range (Suppl. Figure 1-1), and the vast majority of our participants were old

enough that the peak alpha frequency was larger than 8 Hz (Freschl et al., 2022), we

preferred to use fixed bands with gaps. The completely distinct topographies across the

three bands for all age groups support this decision for this dataset (Figure 6, Figure 7).

2.4   Spectral power analysis

Spectral power was calculated using MATLAB's *pwelch* function, with 4 s Hanning win-

dows and 50% overlap. When average power across channels was calculated, edge

channels were excluded (total count: 98). To dissociate periodic and aperiodic spectral

power, we used the MATLAB extension of specparam (formerly known as FOOOF (Do-

noghue et al., 2020)). Spectra were smoothed over 2 Hz, and the aperiodic signal was

fitted between 2 and 35 Hz (frequencies sufficiently separated from the 0.5-40 Hz filter

range of the preprocessed data). Otherwise, the default settings were used (peak width:

[0.5 12], max number of peaks: inf; minimum peak height: 0; peak threshold: 2; aperiod-

ic mode: fixed).

2.4.1   Spectral and aperiodic measures

Power was calculated by averaging the log-transformed power values between 4 and 16

Hz. Aperiodic *offsets* were provided by specparam as the log power value at 1 Hz of the

aperiodic signal, and *exponents* as the x value in the $1/f^x$ model that describes the

steepness of the aperiodic signal. The values are such that positive exponents refer to a

downward descending aperiodic signal, and the larger the value the steeper the de-

scent. *Periodic power* was calculated as the log-transformed power from which the

aperiodic signal was subtracted. Fitting model parameters *r-squared* and *mean abso-*

lute error (MAE) were similarly analyzed to evaluate whether the model fit could ac-

count for the results (Ostlund et al., 2022), with the results provided in Suppl. Figure 2-

1. Overall, the specparam model fit was very good, with average MAE of 0.028 (inter-

quartile range: 0.019, 0.035) and r-squared values of 0.998 (0.997, 0.999).

2.5 Sleep EEG

A subset of wake data was compared to previously analyzed sleep data (Jaramillo et al.,

2020), comprising the *sleep-wake dataset*. This included only neurotypical participants

from Dataset2016 and Dataset2017, resulting in 44 participants, 50% female, with

mean age of 16.3 (8.1-24.8). Only one night per participant (experimental baseline) was

included. The corresponding wake data was the auditory oddball for Dataset2016 (>18-

year-olds), and the TAP Alertness task for Dataset2017 (<18-year-olds). The channel-

averaged data is provided in Suppl. Data 2.

EEG sleep data was recorded at 500 Hz with Cz reference. Sleep stages were scored

according to standard AASM scoring guidelines (Iber, 2007). Epochs containing arte-

facts were rejected after visual inspection using a semi-automatic approach (Huber et

al., 2000). Channels with poor signal quality were removed and interpolated, and outer

edge channels were removed (total count: 98).

Traditionally, sleep pressure is quantified with average spectral power ("slow wave ac-

tivity" is power between 0.5-4 Hz). However, directly measuring individual slow wave

parameters is considered a more precise marker of sleep pressure and synaptic plas-

ticity (Esser et al., 2007; Riedner et al., 2007; Vyazovskiy et al., 2009). This is why both

here and in our previous study (Jaramillo et al., 2020) we analyze sleep data focusing on

slow wave amplitudes and especially slopes.

Slow wave detection was performed similarly to Riedner et al. (2007). The EEG was

bandpass filtered from 0.5 to 4 Hz (stopband 0.1 and 10 Hz, Chebyshev Type II filter)

and rereferenced to the average. Negative deflections between zero-crossings were

identified as slow waves if they were separated by 0.25-1 s. *Slow wave amplitude* con-

sisted of the most negative amplitude between zero-crossings of the signal. The de-

scending *slow wave slope* was the amplitude of the negative peak divided by the time

from the positive-to-negative zero-crossing to the negative peak. The negative peak was

used because this corresponded to silent periods of neuronal spiking, and the de-

scending slope is thought to reflect the synchronization of this silent period across neu-

rons (Nir et al., 2011; Vyazovskiy et al., 2009).

Slow waves were detected in the first and last hour of artifact-free NREM sleep. Given

that slopes are directly calculated from amplitudes, slow waves from the first to last

hour were first matched by amplitude (see Jaramillo et al. (2020) for details), and only

the slopes of matched slow waves were included in the analysis (~60% of total waves).

Thus, the decrease in slow wave slopes across the night is independent from the de-

crease in amplitudes.

2.6   Statistics

Statistics were performed using the MATLAB Statistics and Machine Learning Toolbox.

For all analyses, statistical significance was determined for p-values < .05. Given the

heterogeneous datasets pooled together for this analysis, we chose to conduct linear

mixed effects models to model the relationship between age, sleep, ADHD, and EEG

measures. This was done with the function *fitlme*.

For each fixed factor of the model, β estimates, t-values, p-values, and degrees of free-

dom (df) are reported in the text. β estimates of continuous variables (e.g., age) indicate

by how much the EEG measure (e.g. density) changes for each unit of the continuous

variable (e.g., 1 year) when all other variables are 0. Similarly, the β estimates of cate-

gorical variables (e.g. group) indicate how much the EEG measure changes from that

category (e.g., ADHD) to the baseline category (e.g., controls), for all other factors set to

0. Lastly, t-values allow a comparison of the magnitude of the effect of each factor

when comparing models with different measuring units. To determine the topography of

the effects, we plotted β estimates and their associated statistical significance, cor-

rected for multiple comparisons across channels using false discovery rates (FDR;

(Benjamini & Yekutieli, 2001)).

We ran models with fixed factors Task, Time, Age, Group, and Sex, the interaction be-

tween Time and Age, and nested mixed factors Participant and Session. *Task* compared

the levels *Oddball* vs. *go/no-go*, *alertness*, and *fixation*. *Time* compared the time of re-

cording; evening vs. morning for wake, and first vs. last hour of NREM sleep. *Group*

compared neurotypical participants vs. those with ADHD. *Sex* compared females vs.

males. Depending on the analysis and subset of the data, different factors were includ-

ed or not, and so the exact model is specified before each analysis and in each figure

caption.

As a simpler analysis and sanity check, we conducted Pearson's correlations between

age and each wake measure, including only auditory oddball recordings, averaging ses-

sions (Figure 2). In each figure, r values are provided as effect sizes. Pearson's correla-

tions were also conducted between all measures in the sleep-wake dataset (Suppl.

Figure 2-2).

3   RESULTS

3.1   Effects of age, sleep, sex, and ADHD on wake EEG measures

The following linear mixed effects model was applied to each wake EEG measure, aver-

aged across channels: $Measure \sim Task + Time * Age + Group + Sex +$

$(1|Participant) + (1|Participant{:}Session)$. The full outputs of the model are provided

in Suppl. Data 3.

Oscillation amplitudes significantly decreased with age (β = -0.783, t = -11.33, p < .001,

df = 1234), significantly decreased the morning following sleep (β = -3.977, t = -15.90, p

< .001, df = 1234), with a significant positive interaction (β = 0.143, t = 7.74, p < .001, df =

1234), such that amplitudes decreased less overnight with increasing age (Figure 2,

bottom row). Amplitudes were significantly lower in males than females (β = -1.476, t = -

2.38, p = .017, df = 1234), and were not significantly different in participants with ADHD

(β = -0.574, t = -0.89, p = .373, df = 1234). As can be seen in Figure 2, the relationship

between age and amplitudes was quite robust, both as absolute values (r_{eve} = -.66, r_{mor} =

-.63) and overnight changes (r = .56). Overall, amplitudes changed in the expected di-

rections for both development and sleep pressure, except for the lack of an effect of

ADHD.

Oscillation densities significantly decreased with age (β = -3.540, t = -2.11, p = .035, df =

1234), and the morning after sleep (β = -77.506, t = -10.39, p < .001, df = 1234), with a

significant positive interaction between age and time of recording (β = 4.878, t = 8.85, p

< .001, df = 1234). Unlike amplitudes, the correlation between age and density was

weak (r_{eve} = -.28, r_{mor} = .01). Instead, the correlation between age and overnight change

in density was quite strong (r = .57), such that oscillation densities decreased overnight

in children under 15 and increased in young adults (Figure 2). There was no effect of

ADHD (eta = -0.154, t = -0.01, p = .992, df = 1234) or sex (β = -16.446, t = -1.10, p = .271,

df = 1234). Overall, oscillation densities behaved independently from amplitudes, es-

pecially in the direction of overnight changes in adolescents and adults.

Aperiodic exponents became significantly shallower with age (β = -0.027, t = -9.23, p <

.001, df = 1234) but significantly steeper overnight (β = 0.099, t = 5.51, p < .001, df =

1234), with a trending negative interaction between time of recording and age (β = -

0.003, t = -1.91, p = .056, df = 1234). The increased overnight steepness was driven pri-

marily by a decrease in higher frequency power (Suppl. Figure 5-1), whereas decreasing

steepness with age is driven by decreases in low frequency power (Favaro et al., 2023).

The correlations between exponents and age were as robust as for oscillation ampli-

tudes (r_{eve} = -.60, r_{mor} = -.66), but the correlation with overnight change was weak (r = -

.15, not significant). There was no significant effect of ADHD (β = 0.028, t = 1.07, p =

.284, df = 1234) or sex (β = 0.041, t = 1.63, p = .102, df = 1234). This means that aperiodic

exponents also change independently from oscillation amplitudes, and do not reflect

the direction of changes expected for sleep pressure.

Aperiodic offsets also significantly decreased with age (β = -0.067, t = -16.59, p < .001,

df = 1234), but with no significant effect of time (β = 0.029, t = 1.65, p = .100, df = 1234),

and a significant negative interaction (β = -0.004, t = -3.11, p = .002, df = 1234), such

that offsets decreased more overnight with age. The correlations between age and off-

sets were the strongest of all measures (r_{eve} = -.81, r_{mor} = -.83), however, the correlation

between age and overnight change in offset was negligible (r = -.05). Again, there was no

effect of ADHD (β = 0.021, t = 0.55, p = .579, df = 1234) or sex (β = -0.003, t = -0.07, p =

.944, df = 1234). Overall, offsets correlated with age in the same direction as ampli-

tudes, changed overnight in the same direction, but the overnight change was not larger

in children compared with adults, counter to what would be expected for sleep pres-

sure.

Curiously, fitting parameters of the specparam algorithm showed both significant age

and time of recording effects, with the model fit improving in the morning and worsen-

ing with age (Suppl. Figure 2-1). Therefore, differences in fit quality may partially con-

tribute to the effects observed for aperiodic measures. However, the changes in model

fitting go in the opposite direction one would expect of data quality (i.e. worse in the

morning and better in adults). This suggests that some other aspect of the EEG signal

that changes with age and time can affect the fit quality of the specparam algorithm.

Figure 2: Wake EEG measures correlated with age. Only auditory oddball recordings were included, pooling both
neurotypical and ADHD participants. Each dot represents the value for a single participant. For participants with
multiple sessions, values across sessions were first averaged. Pearson's correlations were done for each figure, with
rho values provided in the corner. If the p-value was less than .05, a correlation line was drawn (without correcting for
multiple comparisons). Amplitudes and densities of oscillations were obtained from burst clusters, pooling all fre-
quencies (4-16 Hz). Exponent values are such that larger values indicate a steeper aperiodic signal; units are "arbi-
trary" (a.u.), in that exponents are scale-free measures. Power spectra were calculated for each channel, then aver-
aged across all channels, excluding the outermost ring of channels. Power and periodic power were then calculated
by averaging values from 4 to 16 Hz. The same figure for the specparam fitting estimates is provided as Suppl. Figure
2-1. The correlations between each measure with the other is provided in Suppl. Figure 2-2.

In summary, of the four EEG measures, only amplitudes followed the same trajectories

expected for both development and sleep pressure. The absolute values of all four

measures had a negative correlation with age, and differed primarily in the overnight

response and the relationship between age and overnight response. Oscillation densi-

ties in particular showed a strong effect of age on overnight changes, reversing direction

between childhood and adolescence. No measure showed any relationship with ADHD,

and only amplitudes were affected by sex. Therefore, in later analyses we did not in-

clude these factors, and pooled patients and controls for greater statistical power.

3.1.1 Comparison of oscillation and aperiodic measures to spectral measures

To quantify the extent to which wake spectral measures were influenced by oscillation

and aperiodic measures, we directly correlated each measure to the other in the sleep-

wake dataset (Suppl. Figure 2-2), and then controlled for time of the recording, task,

and age using linear mixed effects models in the complete wake dataset:

$Measure1 \sim Measure2 + Time * Age + Task + (1|Participant) +$

$(1|Participant: Session)$ (Suppl. Table 2-1).

Without correcting for anything, all measures significantly correlated with each other,

except periodic power which did not correlate with either aperiodic exponents or off-

sets. Power was most related to amplitudes, also in the mixed effects model (correla-

tion r = .93, mixed effects t-value = 39.3). Power was also significantly related to densi-

ties (r = .82, t = 28.2) and offsets (r = .80, t = 17.5), but less so with exponents (r = .54, t =

4.7). Periodic power was most correlated with densities (r = .80, t = 46.9) and somewhat

with amplitudes (r = .53, t = 20.6).

3.1.2 Comparisons between wake and sleep EEG measures

Using the sleep-wake dataset, we directly compared wake measures with sleep slow

wave amplitudes and slopes, first through simple correlations (Suppl. Figure 2-2) and

then with the linear mixed effects model: $SleepMeasure \sim WakeMeasure + Time *$

$Age + Task + (1|Participant)$. The full models are provided in Suppl. Data 4.

All wake measures were significantly correlated with both slow wave amplitudes (wake

amplitude r = .61; density r = .27; exponent r = .46; offset r = .66) and slopes (amplitude r

= .70; density r = .38; exponent r = .60; offset r = .80). When controlling for the time of

the recording, task, and age, wake amplitudes were significantly related to both slow

wave slopes (β = 4.21, t = 2.8, p = .006, df = 82) as well as slow wave amplitudes (β =

0.78, t = 2.7, p = .007, df = 82). The same was true for wake power (slow wave slopes: β

= 26.36, t = 2.3, p = .023, df = 82; slow wave amplitudes: 5.84, t = 2.6, p = .011, df = 82).

While aperiodic offsets had shown the strongest correlations to sleep measures, with

the mixed effects model they were only significantly related to amplitudes (β = 14.69, t =

2.8, p = .006, df = 82) but not slopes (β = 40.26, t = 1.6, p = .111, df = 82). No other wake

measure was related to either slopes (density: β = 0.05, t = 1.1, p = .289, df = 82; expo-

nent: β = 7.78, t = 0.3, p = .785, df = 82; periodic power: β = 44.92, t = 1.2, p = .247, df =

82) or amplitudes (density: β = 0.02, t = 1.2, p = .235, df = 82; exponent: β = 5.99, t = 0.8,

p = .453, df = 82; periodic power: β = 7.69, t = 0.8, p = .412, df = 82).

Overall, wake amplitudes were most related to sleep slow waves, with significant rela-

tionships to both amplitudes and slopes. This association was reflected in wake power.

Wake offsets were also significantly correlated to sleep measures, however the associ-

ation between offsets and sleep slopes did not survive the correction for age and time

in the mixed effects model.

3.2   Topography of wake EEG measures by age

Figure 3 provides the average topographical maps of each measure for five age bins,

averaging (or pooling for densities) all frequencies from 4-16 Hz, from the oddball task.

Amplitudes, densities, exponents, and offsets all showed unique topographies from

each other. Across ages, for each measure there were primarily changes in magnitude

more so than major regional differences. However, oscillation amplitudes in the young-

est cohort began as a single midline occipital spot, which spread bilaterally in the 7–10-

year-olds. Prominent central bilateral peaks also appeared in the 7–10-year-olds. Oscil-

lation densities similarly started as a single midline occipital spot, but these became

more lateral-parietal in the 14–18-year-olds. Like amplitudes, two small bilateral cen-

tral peaks emerged in the densities of 7–10-year-olds, which merged with the primary

occipital-parietal cluster in the 14–18-year-olds. Furthermore, a frontal peak gradually

emerged with age. Exponents were steepest in midline channels, whereas offsets

showed both a frontal midline and occipital peak. As with the correlations between

measures, power topographies most resemble amplitudes, and periodic power resem-

bles densities.

Figure 3: Topography averages of wake EEG measures. Each plot is a schematic of the EEG viewed from above, with the nose on top. Lighter colors indicate greater magnitude over a given location for that measure (rows). Only neurotypical participants and oddball recordings were included, and participants were grouped into age bins (columns). Multiple oddball recordings from different sessions and times of day were first averaged for each participant. The number of participants included is indicated in the top right corner of each plot. Acronyms: y.o., years-old; a.u., arbitrary units.

To determine the topography of overnight changes in EEG measures, we performed linear mixed effects models for each channel, with the model Measure ~ Time + Task + (1|Participant) + (1|Participant: Session), dividing participants into 4 age bins (the

youngest 3-7 were excluded as they were too few, with too few recordings). Figure 4

plots the β estimates for the effect of Time.

Amplitudes showed widespread overnight decreases across all age groups, however

the decrease was largest in occipital channels for the youngest group, and slightly more

fronto-temporal in young adults. The overnight density topographies resembled the av-

erage density topographies from Figure 3, in terms of location of the effects. The young-

est group showed the largest overnight decrease in the same midline-occipital spot

where there were the most oscillations (Figure 3), and adults showed the largest in-

crease in the same bilateral occipital-parietal areas where they had the largest densi-

ties.

The overnight increase in the steepness of exponents was widespread, but peaked in an

occipital spot in all age groups, with additional bilateral frontal spots in <18-year-olds.

These topographies do not correspond to the average topography of exponents from

Figure 3. Offsets revealed widespread decreases, with localized increases in the same

occipital locations for which exponents increased the most. This suggests that aperiod-

ic offsets generally decrease, although the increase in exponent steepness contrasts

this effect.

Power and periodic power again showed similarities to amplitudes and densities re-

spectively, however while densities increased in the 14–18-year-olds, periodic power

decreased. Likewise, the overnight increase in periodic power for >18-year-olds was

more occipital and lateral than the increase in densities, and the larger decrease in

amplitudes in occipital regions of 7–10-year-olds was less evident in power than for

amplitudes.

Figure 4: Topographies of wake EEG overnight changes. A linear mixed effects model was run for each measure, each age group, and each channel: Measure ~ Time + Task + (1|Participant) + (1|Participant:Session). Color reflects β estimates for the fixed effect *Time*, such that red indicates an overnight increase in that measure. The factor *Task* was not included for the 18-25 y.o. group, as these participants only performed oddballs. White dots indicate channels for which the β estimate was statistically significant, corrected for multiple comparisons with FDR (false discovery rate). Black dots indicate remaining channels. Data includes both patients and neurotypical controls. Degrees of freedom (df) are provided for each plot. For each column, the sample sizes were N = 29, 67, 36, & 23.

3.3 Frequency-specific wake EEG changes with age and sleep pressure

In the previous analyses, we had pooled all frequencies between 4 and 16 Hz. Subsequently, we explored how oscillation and spectral EEG measures changed for each frequency, this time averaging channels. Average evening values for each age and frequency are plotted in Figure 5A, and the overnight differences are plotted in Figure 5B.

Average oscillation amplitudes and spectral power showed gradual gradients, with highest values in the youngest participants and lowest frequencies, and lowest values in the oldest participants and highest frequencies. Like average amplitudes, overnight changes in amplitude showed largely gradual decreases with age and frequency.

Average density and periodic power instead had distinct peak values between 8 and 11 Hz, with the peak shifting upwards with age, a well-known property of alpha oscillations during development (Freschl et al., 2022; Smith, 1938; Tröndle et al., 2022). Overnight densities (and periodic power) showed decreases in higher frequencies (>11 Hz, i.e. low beta) and increases in alpha. These increases only began between 8-10 years of age, they were strongest in adults, and the range shifted to higher frequencies with age.

Figure 5: Average spectrograms across ages. From the oddball task, pooling controls and ADHD participants. Average spectra from the other tasks are provided in Suppl. Figure 5-1. **A**: Average values, such that lighter colors indicate greater magnitude for a given frequency and age. For spectra extending to older ages and sleep stages, see Sun et al. (2023). **B**: Difference values between morning and evening recordings, such that red indicate a greater magnitude in the morning. The measurement unit of each figure is the same as that of A.

Given the dissociation between alpha and beta for density, we explored the topographic changes in density, split by both age and frequency. In Figure 6, we plot average values. Theta oscillations were the overall rarest. They were most prevalent in the youngest children, and with age, the peak in theta density gradually shifted upward and decreased in magnitude. Alpha density instead started as a midline occipital cluster in the 3–7-year-olds. Alpha densities in the occipital spot decreased in the 7-10 cohort, with bilateral central spots instead becoming more pronounced. With age, these three peaks morphed into a continuous occipital-parietal cluster, while in the remaining channels overall density of alpha increased. Lastly, low beta oscillations showed yet another topography. They were almost completely absent in the youngest group, and started to appear in the 7–10-year-olds as lateral occipital peaks and a frontal midline

spot. Gradually, the lateral peaks converge towards the midline, and in adults became

partially overlapping with alpha, on average right-lateralized.

Figure 6: Average topographies of oscillation densities, split by frequency band. Recordings are the same as in
Figure 3. Theta is 4-7 Hz, alpha is 8-11 Hz, and low beta is 12-16 Hz. Each band has a different color scale range.

The overnight changes in density split by frequency are in Figure 7, based on the model

$\text{Density} \sim \text{Time} + \text{Task} + (1|\text{Participant}) + (1|\text{Participant:Session})$. Like in Figure 4,

these reflect the β estimates for the fixed effect of Time.

The overnight changes in theta density were small (around 1%), however quite variable

by region and age. In the 7–10-year-olds, theta density generally increased overnight

except in a central spot, exactly where the largest theta densities were seen in Figure 6,

which instead decreased. This continued for the 10–14-year-olds. In the 14–18-year-

olds, there were no significant effects, however the midline theta spot, now more

frontal, showed a decrease. In adults, only some scattered theta increases were ob-

served.

For alpha, the main occipital spot in the 7–10-year-olds decreased overnight. Already in

the 10–14-year-olds a bilateral central alpha rhythm started to increase overnight, with

still some slight decreases in the occipital spot. The overnight increases spread to the

entire scalp in adolescents and adults, peaking in occipital parietal areas, especially

1 right lateralized. For low beta, across all ages there were decreases, with the peak shift-

2 ing across age bins.

Figure 7: Topographies of overnight density changes, split by frequency band. Participants and plot information are the same as in Figure 4, with color indicating the β estimates for the linear mixed effects model, such that red indicates an overnight increase in density. The model was Density ~ Time + Task + (1|Participant) + (1|Participant:Session).

3.4 Effect of ADHD on the wake EEG topography

In our initial mixed effects models pooling channels and frequencies, we found no sig-

nificant effects for ADHD, which is why in subsequent analyses and figures we no long-

er included a Group effect. However, we nevertheless conducted mixed effects models

to determine the effect of ADHD for each channel: Measure ~ Task + Time ∗ Age +

Group + (1|Participant) + (1|Participant: Session). We found no significant effects

when correcting for multiple comparisons (Figure 8). However, amplitudes were on av-

erage lower in participants with ADHD and frontal exponents were steeper compared to

controls.

Figure 8: Effects of ADHD on EEG measures. Red indicates larger values in patients compared to controls. The scale for each topography is the same as for Figure 4. The model was Measure ~ Time*Age + Task + Group + (1|Participant) + (1|Participant:Session). White dots would have indicated statistically significant channels, following FDR correction for multiple comparisons.

4 DISCUSSION

In this study, we compared four distinct wake EEG measures and their relationship to development, sleep pressure, and ADHD. Specifically, we hypothesized that oscillation amplitudes would behave like sleep slow waves, because both measures likely reflect neuronal synchronization due to synaptic density and plasticity. Our predictions were met on all accounts except for the sensitivity of wake amplitudes to ADHD. Of the four measures, only amplitudes decreased overnight in all ages *and* the overnight decrease was largest in younger children (Figure 2). Wake amplitudes were significantly correlated to sleep slow wave measures, also when correcting for age and participant. Aperiodic offsets showed similar development and overnight changes, as well as strong correlations to sleep measures, but they did not display larger overnight decreases in younger children, a key indicator of higher neural plasticity.

While oscillation amplitudes were the only wake measure that followed all the patterns expected for sleep pressure, every EEG measure reflected development. Average amplitudes, exponents, and offsets were all strongly anticorrelated with age, with offsets showing the strongest relationship (Figure 2). The correlation between average densities and age was weak, but we found distinctive regional patterns across age groups (Figure 6). Overnight changes in both oscillation amplitudes and density were robustly correlated with age (Figure 2, Figure 7), whereas overnight changes in exponents and offsets were not as affected by age, as evidenced by the low correlations in Figure 2.

1   These results are summarized in Figure 9. Finally, no measure showed significant ef-

2   fects of ADHD (Figure 8).

		Sleep pressure	Development
Oscillation	**Amplitude**	n.a.	✓
	Overnight amplitude	✓	✓
	Density	n.a.	✓
	Overnight density	✗	✓
Aperiodic	**Exponent**	n.a.	✓
	Overnight exponent	✗	n.s.
	Offset	n.a.	✓
	Overnight offset	mixed	mixed

4   **Figure 9: Summary of which measure reflects sleep pressure or development.** Sleep pressure was determined by
5   whether a measure decreased overnight and whether the decrease was larger in younger children. Development was
6   determined by whether there were strong effects of age. Overnight offsets had "mixed" results across analyses.
7   Acronyms: n.a., not applicable; n.s., not significant.

4.1 Oscillation amplitudes

Like for slow waves during sleep, the decrease in wake oscillation amplitudes with age

could be explained by decreasing synaptic density in the cortex across adolescence

(Huttenlocher, 1979). The overnight decrease in wake amplitudes could be explained by

sleep's role in reducing net synaptic strength (Cirelli & Tononi, 2022; Tononi & Cirelli,

2014), and such plastic changes are more pronounced in children than adults (Jaramil-

lo et al., 2020). The larger decrease in occipital regions in children compared to adults

may be because primary sensory and motor areas obtain peak cortical thickness earlier

in children, followed by adjacent secondary areas and finally frontal association areas

(Shaw et al., 2008), resulting in an overall posterior-anterior maturation trajectory (Kurth

et al., 2010). Therefore, younger children show larger overnight decreases in amplitudes

in occipital areas because these areas undergo higher plastic changes at that matura-

tional stage. Finally, even the significant sex difference was comparable to sleep slow

wave activity (Mourtazaev et al., 1995), with wake oscillation amplitudes higher in fe-

males than males (although the effect was relatively small).

Higher slow wave activity in females has been hypothesized to reflect their smaller

heads and thinner skulls (Dijk et al., 1989). It is then also possible that head and skull

size affect the age-related changes we observed for wake amplitudes. However, Scha-

woronkow and Voytek (2021) did not find any changes in wake oscillation amplitudes

from 1-7 month-old infants, despite this period corresponding to the fastest growth in

head circumference (Roche et al., 1986). Therefore, it is unlikely that the more modest

increases in head size from childhood to adolescence can entirely explain the large de-

crease in amplitudes we observed here.

While wake oscillation amplitudes behave largely like sleep slow waves, the effects are

reduced. In adults the overnight decrease in wake amplitudes was near 0 µV (Figure 2),

which is not the case for either slow wave slopes or amplitudes during sleep (Jaramillo

et al., 2020; Riedner et al., 2007). One possible explanation for this is that the evening

wake recordings fell within the wake maintenance zone. This is a circadian time window

just before bedtime characterized by increased alertness (Shekleton et al., 2013;

Strogatz et al., 1987). We had found in our previous study that oscillation amplitudes

are significantly reduced in this window, counteracting the otherwise monotonic

buildup in amplitudes that occurred throughout the day (Snipes et al., 2023). In adults

the contrasting effect of the wake maintenance zone may be sufficient to equalize

evening and morning amplitudes. Overall, wake oscillation amplitudes may reflect the

same information as sleep slow waves but may be more affected by circadian or other

factors. Therefore, wake amplitudes may be affected by the same underlying neuronal

changes as sleep slow wave activity, but they are not sufficiently specific nor sensitive

to replace this gold standard measure of sleep pressure.

4.2 Oscillation densities

As could be expected, the densities of oscillations were the most complex EEG meas-

ure, with effects differing depending on age, time, topography, and oscillation frequen-

cy.

Theta oscillations were most prevalent in early childhood, and decreased progressively

with age, supporting previous results measuring relative theta power (Somsen et al.,

1997). We further found that the peak in theta densities steadily drifted more frontally

across childhood and adolescence. This frontal theta in adults is known to originate

from the midline prefrontal cortex, and to be anti-correlated to the default mode net-

work (Ishii et al., 2014; Michels et al., 2010; Scheeringa et al., 2008). Therefore, this drift

in theta may reflect the steady maturation of both frontal cortices and the default mode

network (Fan et al., 2021).

There is still unresolved contradictory evidence on the role of theta in adults (Snipes et

al., 2022), without including the question of theta during development. On the one

hand, theta is often associated with cognitive effort (Buzsáki, 2005; Cavanagh & Frank,

2014; Meyer et al., 2019; Mitchell et al., 2008). On the other, it is also associated with

sleepiness (Finelli et al., 2000; Smith, 1938; Snipes et al., 2022) and fatigue (Arnau et

al., 2021; Tran et al., 2020; Wascher et al., 2014). A possible resolution to this paradox

is that there are distinct oscillations that originate from different circuits with different

functions and just happen to occur at the same frequency. Our results in Figure 7 would

support this, as the peak source of theta shows overnight *decreases*, whereas theta

from the rest of the cortex instead shows overnight increases, even as the theta peak

drifts more frontally. Alternatively, theta could reflect a general form of "idling rhythm"

(Snipes et al., 2022, 2024), originating from disengaged cortical areas, and what chang-

es from evening to morning is which circuits tend to idle. Theta as an idling rhythm is

supported by simultaneous EEG-fMRI studies that find theta activity anti-correlating

with brain metabolism (Scheeringa et al., 2008). This would make theta functionally

comparable to alpha (Laufs et al., 2003), differing only by source and frequency.

Alpha begins in young childhood as a midline spot (Figure 6). Two lateral central peaks

become more defined at 7-10 years of age. These likely reflect sensorimotor mu

rhythms which appear when motor activity is absent or even suppressed (Pfurtscheller

et al., 2006; Pineda, 2005), and is already present in infants (Berchicci et al., 2011). We

find that with age, they become topographically indistinguishable from occipital alpha,

at least when recorded during an oddball task. These lateral central peaks are the first

to show overnight increases in 10-14 year-olds, while the overnight decrease in the oc-

cipital midline spot becomes less prominent. The overnight increase in alpha densities

then spreads over bilateral parietal and occipital areas across adolescence and adult-

hood. This dissociation between overnight decreases in childhood and increases in

adulthood, as well as the slight differences in topography, could suggest that occipital

alpha is in fact qualitatively distinct in children and adults. However, these rhythms

were previously considered functionally equivalent because also in infants alpha power

increases with eyes closed compared to eyes open (Stroganova et al., 1999). More re-

search is needed on the sources of these oscillations.

It is also possible that this dissociation in overnight density changes is driven by some

other difference with age, such as a longer window of sleep inertia in young children,

longer sleep duration, or a shifted circadian rhythm compared to adults. Melatonin in

the morning is elevated in children under 10, whereas older children and adolescents

have morning melatonin levels comparable with the rest of the day (Attanasio et al.,

1985). In adults, alpha power fluctuates with circadian rhythm and is therefore syn-

chronized to melatonin levels (Cajochen et al., 2002). Therefore, it is possible that the

dissociation of decreasing/increasing alpha densities originates from children and

adults being at different phases of their alpha circadian rhythm in the morning. More

research is needed into the circadian effects on the EEG across development.

Finally, we found overnight decreases in low beta densities across ages. There is ample

literature on "beta bursts" from sensorimotor areas (Feingold et al., 2015; Jones et al.,

2009; Little et al., 2019; Lundqvist et al., 2024; Shin et al., 2017; Wessel, 2020), however

these are likely qualitatively distinct from the beta oscillations we observed. First, the

topography of sensorimotor beta rhythms was found to be either bilateral-central in

infants or frontal in adults (Rayson et al., 2023), unlike the bilateral occipital topogra-

phies in Figure 6. Second, sensorimotor beta was found to occur in "bursts" of power

less than 150 ms long (Sherman et al., 2016) and could thus correspond to only 1-3 cy-

cles of beta (Jones, 2016; van Ede et al., 2018). Supporting this, lagged coherence de-

creases rapidly after 2 cycles (Rayson et al., 2023). We detected bursts that were *at*

least 3 cycles long, only up to 16 Hz, and therefore always longer than 187 ms. There-

fore, the low-beta bursts we captured with cycle-by-cycle analysis are likely distinct

from the previously investigated sensorimotor beta activity. It remains an open ques-

tion what is the functional role of these low-beta oscillations, and why they may be less

common the morning after sleep.

4.3 Aperiodic offsets and exponents

Our results on offsets and exponents replicate previous findings: they decrease linearly with age (Cellier et al., 2021; Hill et al., 2022; Tröndle et al., 2022) and originate from broad, primarily midline sources (Favaro et al., 2023). As in Lendner et al. (2023), we found exponents becoming increasingly steeper after sleep, extending this finding across ages. We further found, unusually, that exponents and offsets change in opposite directions, with offsets decreasing after a night of sleep. For aperiodic activity measured *during* sleep, both exponents and offsets decrease across the night, and the decrease decreases with age (Horváth et al., 2022). This indicates that changes in aperiodic activity related to sleep-wake history do not reflect the same information when measured during wake or during sleep. It is possible that during sleep the decrease in exponents reflects the decrease in sleep pressure, whereas during wake the increase in exponents could reflect something like lingering sleep inertia in the morning. Notably, the change in aperiodic exponents with age and sleep depth affect primarily lower frequencies (Favaro et al., 2023), whereas the overnight wake EEG changes affect primarily higher frequencies (Suppl. Figure 5-1). This means that the pivot point (the frequency at which the aperiodic signal rotates) of the exponent differs, and this likely has biological significance. Future studies should, as in Bódizs et al. (2021), specifically identify such pivot points, also to better dissociate changes in offset and exponent.

Second to oscillation amplitudes, aperiodic offsets showed the closest resemblance to sleep slow wave activity, decreasing overnight and showing strong correlations with slow wave measures. While aperiodic offsets were not significantly related to slow wave slopes when controlling for age and participant, this does not imply a significant difference from wake oscillation amplitudes. However, offsets did not show the characteristic decrease in overnight changes with age that was observed for slow waves and wake amplitudes (Figure 2). This could suggest that offsets also reflect changes in synaptic strength and plasticity, but may be less sensitive or specific than wake amplitudes, especially to overnight changes.

4.4   Spectral power vs oscillation burst detection

As explained by previous papers, changes in spectral power do not differentiate between changes in oscillation amplitudes or densities (Donoghue et al., 2022; Quinn et al., 2019; Snipes et al., 2023; Tal et al., 2020; Zich et al., 2020). With this study, we again demonstrate that amplitudes and densities meaningfully change independently, both across sleep and development. However, we also found that the changes in amplitude were largely captured by changes in average power, whereas changes in density were largely captured by changes in periodic power. This suggests that power and periodic power can be used as proxies for amplitudes and densities. Previous studies of development have likewise found differences between power and *relative power* (similar to periodic power) across development (Somsen et al., 1997). These results may be reinterpreted as the dissociation between changes in amplitude and density.

Arguing against this approach, however, is the topography of overnight changes in 14–18-year-olds: oscillation densities increased overnight, whereas periodic power decreased, likely reflecting the greater influence of the overnight decrease in amplitudes. Similarly, the overnight increase in densities is more central than the overnight increase in periodic power. Therefore, differences between power and periodic power may suggest different influences of oscillation amplitudes and densities, but to know for certain, oscillations should be measured directly.

4.5   ADHD

Despite a relatively largesample size (N = 58), we did not observe any significant effects of ADHD on our EEG measures. We did find exponents to be steeper on average in patients, supporting the results of Robertson et al. (2019), and contrasting those of Ostlund et al. (2021). One explanation could be that our participants were performing tasks for most recordings, and the differences between patients and controls may mostly emerge in resting EEG; patients may have developed compensation mechanisms masking potential differences during the tasks. It's also possible our analysis did not reach significance because our participants were a combination of both medicated and unmedicated patients, and it is known that medication will reduce the effects of ADHD

on the EEG (Furrer et al., 2019; Karalunas et al., 2022). Additionally, our participants

were screened for good sleep quality (to have a chance of falling asleep in the laborato-

ry with an EEG net, and to have similar levels of sleep pressure as controls). However,

around 40-55% of children with ADHD report sleep deficits (Becker et al., 2019; Corkum

et al., 1998; Holmberg & Hjern, 2006; Konofal et al., 2010), so it is possible that poor

sleep quality in patients contributes to differences in the wake EEG observed in prior

studies (Clarke et al., 2020). Finally, it's possible that only subtypes of patients with

ADHD have a specific relationship to any of the measured EEG markers; ADHD is highly

heterogeneous with varying symptoms among individuals.

Regardless of the reason, given that we do not see any systematic differences between

patients and controls, none of the wake EEG measures we tested make for a reliable

intrinsic marker of ADHD which could potentially be used to aid diagnosis. Instead, re-

search investigating such markers in these and other patient populations should take

special care to control for sleep/wake history and sleep quality, as these may have a

greater impact on the EEG. Instead, within the same participants we observed signifi-

cant differences in sleep slow wave activity from controls (Furrer et al., 2019), suggest-

ing sleep is more sensitive to developmental deficits.

4.6 Limitations

An important limitation of this study is the scarcity of datasets under 8. It is known that

in young children there is a switch from primarily synaptic growth to primarily synaptic

pruning, peaking in different cell populations and regions at different ages (Cao et al.,

2020; Petanjek et al., 2011; Shaw et al., 2008), which is also reflected in sleep slow

wave activity peaking around this age (Feinberg & Campbell, 2013). Wilkinson et al.

(2024) found an increase with age in both offsets and exponents across infants 0-3

years old, and McSweeney et al. (2023) found a quadratic relationship between age and

offsets/exponents, such that they peaked at 5-7 years old. This would suggest more

complex relationships between age and EEG measures than the linear trends observed

here.

A further limitation of the data is its lack of uniformity, pooling multiple experimental

paradigms and tasks, and medicated and unmedicated patients. Our primary findings

remain significant also for more uniform subsets of the data (e.g. Figure 2 is only odd-

ball, Suppl. Figure 2-2 only neurotypical controls), but it's possible that different tasks

would generate different topographies, and significant effects of ADHD would have

emerged from entirely unmedicated patients.

This paper poses two potential weaknesses in interpretability. First, if one assumes that

aperiodic and oscillatory signals are a linear sum of each other in the time-domain of

the EEG signal, it is possible that the aperiodic changes explain part of the changes in

oscillation amplitudes. A better understanding of the relationship between oscillations

and the aperiodic signal are needed to validate such an assumption, and then precise

simulations would be needed to determine the magnitude of this effect. However, in

practice we have demonstrated that oscillation amplitudes can behave differently from

aperiodic measures, and therefore it is at least useful to measure them independently.

Second, overnight changes do not dissociate between homeostatic (related to sleep

pressure) and circadian (related to clock time) effects. To do so would require substan-

tially more intensive protocols, such as sleep deprivation, sleep restriction, or shifting

sleep windows over several days. However, collecting more wake recordings through-

out the day would already provide an indication as to when an effect is circadian or ho-

meostatic.

Finally, our data is limited to EEG. Future studies and analyses would greatly benefit by

comparing these measures to structural and functional brain changes observable with

MRI, as well as cognitive and behavioral outcome measures related to development.

This would bridge the gap between a purely basic research finding to practical applica-

tions.

4.7 Conclusions

We have found that overnight changes in wake oscillations provide distinct markers of

brain maturation. Both absolute amplitudes and overnight changes in amplitudes de-

crease linearly with age, the effect more occipital in younger children. Overnight chang-

es in oscillation density, especially of alpha oscillations, dissociate children from ado-

lescents and adults by switching from an overnight decrease to an increase in density.

Understanding the reason behind this effect would likely provide important information

on brain development around puberty and adolescence. More generally, we have
shown that there are a multitude of changes in the EEG with development that go be-
yond simple spectral power, each with their own functional significance. Moving for-
ward, researchers should analyze independently oscillatory amplitudes and densities
as well as aperiodic exponents and offsets, as they offer different insights into neuronal
activity and structure.

5   DATA AND CODE AVAILABILITY

The data tables used for the linear mixed effects models are provided as supplementary
material (Suppl. Data 1, Suppl. Data 2), and the output of the linear models for wake
and sleep data are likewise provided (Suppl. Data 3, 4). The raw and preprocessed EEG
is available upon request, requiring permission from the local ethics committee.

The burst detection can be conducted either with the original python toolbox bycycle
(https://github.com/bycycle-tools/bycycle), or with our MATLAB implementation
(https://github.com/HuberSleepLab/Matcycle). The preprocessing and analysis code is
likewise open source (https://github.com/snipeso/children-wake/).

6   AUTHOR CONTRIBUTIONS

17 Methodology, SS, EK, VJ, CV, ML, SK, RH; Software, SS; Formal Analysis, SS, VJ; Investi-
18 gation, EK, VJ, CV, MF, MS, ML, SK, RH; Resources, ML, OGJ, RH; Writing–Original draft,
19 SS; Writing-Review & Editing, SS, EK, VJ, MF, MS, SK, OGJ, RH; Supervision, RH; Funding
20 acquisition EK, ML, OGJ, RH

7   DECLARATION OF COMPETING INTERESTS

22 None of the authors have competing interests.

8   ACKNOWLEDGEMENTS & FUNDING

24 This work was supported by the Swiss National Science Foundation (PP00A-114923,
25 PP00P3-135438, CRSII3_160803, 320030_153387, 320030_179443, PCEFP1_181279),
26 the University Research Priority Program of the University of Zurich, Forschungszentrum
27 für das Kind (FZK) of the University Children's Hospital Zurich, the Clinical Research

Priority Program "Sleep and Health" of the University of Zurich, the HMZ Flagship Project "SleepLoop" of University Medicine Zurich, the Borbély-Hess Foundation, Gottfried und Julia Bangerter-Rhyner-Stiftung, the Waterloo Foundation (2462-4978), and the National Institute of Health (K01MH074643). We thank everyone involved in data collection, and preprocessing, especially Maya Ringli, Renato Merki, Michelle Suppiger, Léa Geiger, Martina Liechti, Stefano Maurizio, Dr. Renate Drechsler, Stephanie Cares, Margot Schaerer, Jessica Bühler, Natalie Birnbaum, Demy Marcella Gutjahr, and Soraya Souissi. We thank the children and parents for their participation, and Giulio Tononi for the loan of technical material.

9 REFERENCES

Arnau, S., Brümmer, T., Liegel, N., & Wascher, E. (2021). Inverse effects of time-on-task in task-related and task-unrelated theta activity. *Psychophysiology*, *58*(6), e13805. https://doi.org/10.1111/psyp.13805

Attanasio, A., Borrelli, P., & Gupta, D. (1985). Circadian rhythms in serum melatonin from infancy to adolescence. *The Journal of Clinical Endocrinology and Metabolism*, *61*(2), 388–390. https://doi.org/10.1210/jcem-61-2-388

Bazanova, O. M., & Vernon, D. (2014). Interpreting EEG alpha activity. *Neuroscience & Biobehavioral Reviews*, *44*, 94–110. https://doi.org/10.1016/j.neubiorev.2013.05.007

Becker, S. P., Langberg, J. M., Eadeh, H.-M., Isaacson, P. A., & Bourchtein, E. (2019). Sleep and daytime sleepiness in adolescents with and without ADHD: Differences across ratings, daily diary, and actigraphy. *Journal of Child Psychology and Psychiatry*, *60*(9), 1021–1031. https://doi.org/10.1111/jcpp.13061

Benjamini, Y., & Yekutieli, D. (2001). The Control of the False Discovery Rate in Multiple Testing under Dependency. *The Annals of Statistics*, *29*(4), 1165–1188.

Berchicci, M., Zhang, T., Romero, L., Peters, A., Annett, R., Teuscher, U., Bertollo, M., Okada, Y., Stephen, J., & Comani, S. (2011). Development of Mu Rhythm in Infants and Preschool Children. *Developmental Neuroscience*, *33*(2), 130–143. https://doi.org/10.1159/000329095

Bódizs, R., Schneider, B., Ujma, P. P., Horváth, C. G., Dresler, M., & Rosenblum, Y. (2024). Fundamentals of sleep regulation: Model and benchmark values for fractal and oscillatory neurodynamics. *Progress in Neurobiology*, *234*, 102589. https://doi.org/10.1016/j.pneurobio.2024.102589

Bódizs, R., Szalárdy, O., Horváth, C., Ujma, P. P., Gombos, F., Simor, P., Pótári, A., Zeising, M., Steiger, A., & Dresler, M. (2021). A set of composite, non-redundant EEG measures of NREM sleep based on the power law scaling of the Fourier spectrum. *Scientific Reports*, *11*(1), 2041. https://doi.org/10.1038/s41598-021-81230-7

Borbély, A. A. (1982). A two process model of sleep regulation. *Human Neurobiology*, *1*(3), 195–204.

Brake, N., Duc, F., Rokos, A., Arseneau, F., Shahiri, S., Khadra, A., & Plourde, G. (2024). A neurophysiological basis for aperiodic EEG and the background spectral trend. *Nature Communications*, *15*(1), 1514. https://doi.org/10.1038/s41467-024-45922-8

Buchmann, A., Ringli, M., Kurth, S., Schaerer, M., Geiger, A., Jenni, O. G., & Huber, R. (2011). EEG Sleep Slow-Wave Activity as a Mirror of Cortical Maturation. *Cerebral Cortex*, *21*(3), 607–615. https://doi.org/10.1093/cercor/bhq129

Buzsáki, G. (2005). Theta rhythm of navigation: Link between path integration and landmark navigation, episodic and semantic memory. *Hippocampus*, *15*(7), 827–840. https://doi.org/10.1002/hipo.20113

References

Cajochen, C., Wyatt, J. K., Czeisler, C. A., & Dijk, D. J. (2002). Separation of circadian and wake duration-dependent modulation of EEG activation during wakefulness. *Neuroscience*, *114*(4), 1047–1060. https://doi.org/10.1016/S0306-4522(02)00209-9

Campbell, I. G., & Feinberg, I. (2009). Longitudinal trajectories of non-rapid eye movement delta and theta EEG as indicators of adolescent brain maturation. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(13). https://doi.org/10.1073/pnas.0812947106

Cao, J., Herman, A. B., West, G. B., Poe, G., & Savage, V. M. (2020). Unraveling why we sleep: Quantitative analysis reveals abrupt transition from neural reorganization to repair in early development. *Science Advances*, *6*(38), eaba0398. https://doi.org/10.1126/sciadv.aba0398

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive Sciences*, *18*(8), 414–421. https://doi.org/10.1016/j.tics.2014.04.012

Cellier, D., Riddle, J., Petersen, I., & Hwang, K. (2021). The development of theta and alpha neural oscillations from ages 3 to 24 years. *Developmental Cognitive Neuroscience*, *50*, 100969. https://doi.org/10.1016/j.dcn.2021.100969

Cirelli, C., & Tononi, G. (2022). The why and how of sleep-dependent synaptic down-selection. *Seminars in Cell & Developmental Biology*, *125*, 91–100. https://doi.org/10.1016/j.semcdb.2021.02.007

Clarke, A. R., Barry, R. J., & Johnstone, S. (2020). Resting state EEG power research in Attention-Deficit/Hyperactivity Disorder: A review update. *Clinical Neurophysiology*, *131*(7), 1463–1479. https://doi.org/10.1016/j.clinph.2020.03.029

Cole, S., & Voytek, B. (2019). Cycle-by-cycle analysis of neural oscillations. *Journal of Neurophysiology*, *122*(2). https://doi.org/10.1152/JN.00273.2019

Colombo, M. A., Comanducci, A., Casarotto, S., Derchi, C.-C., Annen, J., Viganò, A., Mazza, A., Trimarchi, P. D., Boly, M., Fecchio, M., Bodart, O., Navarro, J., Laureys, S., Gosseries, O., Massimini, M., Sarasso, S., & Rosanova, M. (2023). Beyond alpha power: EEG spatial and spectral gradients robustly stratify disorders of consciousness. *Cerebral Cortex*, *33*(11), 7193–7210. https://doi.org/10.1093/cercor/bhad031

Colombo, M. A., Napolitani, M., Boly, M., Gosseries, O., Casarotto, S., Rosanova, M., Brichant, J.-F., Boveroux, P., Rex, S., Laureys, S., Massimini, M., Chieregato, A., & Sarasso, S. (2019). The spectral exponent of the resting EEG indexes the presence of consciousness during unresponsiveness induced by propofol, xenon, and ketamine. *NeuroImage*, *189*, 631–644. https://doi.org/10.1016/j.neuroimage.2019.01.024

Corkum, P., Tannock, R., & Moldofsky, H. (1998). Sleep Disturbances in Children With Attention-Deficit/Hyperactivity Disorder. *Journal of the American Academy of Child & Adolescent Psychiatry*, *37*(6), 637–646. https://doi.org/10.1097/00004583-199806000-00014

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dijk, D. J., Beersma, D. G. M., & Bloem, G. M. (1989). Sex Differences in the Sleep EEG of Young Adults: Visual Scoring and Spectral Analysis. *Sleep*, *12*(6), 500–507. https://doi.org/10.1093/sleep/12.6.500

Dijk, D. J., Beersma, D. G. M., & Daan, S. (1987). EEG Power Density during Nap Sleep: Reflection of an Hourglass Measuring the Duration of Prior Wakefulness. *Journal of Biological Rhythms*, *2*(3), 207–219. https://doi.org/10.1177/074873048700200304

Donoghue, T., Haller, M., Peterson, E. J., Varma, P., Sebastian, P., Gao, R., Noto, T., Lara, A. H., Wallis, J. D., Knight, R. T., Shestyuk, A., & Voytek, B. (2020). Parameterizing neural power spectra into periodic and aperiodic components. *Nature Neuroscience*, *23*(12), Article 12. https://doi.org/10.1038/s41593-020-00744-x

Donoghue, T., Schaworonkow, N., & Voytek, B. (2022). Methodological considerations for studying neural oscillations. *European Journal of Neuroscience*, *55*(11–12), 3502–3527. https://doi.org/10.1111/ejn.15361

Esser, S. K., Hill, S. L., & Tononi, G. (2007). Sleep Homeostasis and Cortical Synchronization: I. Modeling the Effects of Synaptic Strength on Sleep Slow Waves. *Sleep*, *30*(12), 1617–1630. https://doi.org/10.1093/sleep/30.12.1617

Fan, F., Liao, X., Lei, T., Zhao, T., Xia, M., Men, W., Wang, Y., Hu, M., Liu, J., Qin, S., Tan, S., Gao, J.-H., Dong, Q., Tao, S., & He, Y. (2021). Development of the default-mode network during childhood and adolescence: A longitudinal resting-state fMRI study. *NeuroImage*, *226*, 117581. https://doi.org/10.1016/j.neuroimage.2020.117581

Fattinger, S., Kurth, S., Ringli, M., Jenni, O. G., & Huber, R. (2017). Theta waves in children's waking electroencepha-
logram resemble local aspects of sleep during wakefulness. *Scientific Reports*, *7*(1), Article 1.
https://doi.org/10.1038/s41598-017-11577-3

Favaro, J., Colombo, M. A., Mikulan, E., Sartori, S., Nosadini, M., Pelizza, M. F., Rosanova, M., Sarasso, S., Massimini,
M., & Toldo, I. (2023). The maturation of aperiodic EEG activity across development reveals a progressive differentia-
tion of wakefulness from sleep. *NeuroImage*, *277*, 120264. https://doi.org/10.1016/j.neuroimage.2023.120264

Feinberg, I., & Campbell, I. G. (2013). Longitudinal sleep EEG trajectories indicate complex patterns of adolescent
brain maturation. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, *304*(4), R296-
303. https://doi.org/10.1152/ajpregu.00422.2012

Feingold, J., Gibson, D. J., DePasquale, B., & Graybiel, A. M. (2015). Bursts of beta oscillation differentiate postper-
formance activity in the striatum and motor cortex of monkeys performing movement tasks. *Proceedings of the Na-
tional Academy of Sciences*, *112*(44), 13687–13692. https://doi.org/10.1073/pnas.1517629112

Finelli, L. A., Baumann, H., Borbély, A. A., & Achermann, P. (2000). Dual electroencephalogram markers of human
sleep homeostasis: Correlation between theta activity in waking and slow-wave activity in sleep. *Neuroscience*,
101(3), 523–529. https://doi.org/10.1016/S0306-4522(00)00409-7

Freschl, J., Azizi, L. A., Balboa, L., Kaldy, Z., & Blaser, E. (2022). The development of peak alpha frequency from infan-
cy to adolescence and its role in visual temporal processing: A meta-analysis. *Developmental Cognitive Neurosci-
ence*, *57*, 101146. https://doi.org/10.1016/j.dcn.2022.101146

Furrer, M., Jaramillo, V., Volk, C., Ringli, M., Aellen, R., Wehrle, F. M., Pugin, F., Kurth, S., Brandeis, D., Schmid, M.,
Jenni, O. G., & Huber, R. (2019). Sleep EEG slow-wave activity in medicated and unmedicated children and adoles-
cents with attention-deficit/hyperactivity disorder. *Translational Psychiatry*, *9*(1). https://doi.org/10.1038/s41398-
019-0659-3

Furrer, M., Ringli, M., Kurth, S., Brandeis, D., Jenni, O. G., & Huber, R. (2020). The experience-dependent increase in
deep sleep activity is reduced in children with attention-deficit/hyperactivity disorder. *Sleep Medicine*, *75*.
https://doi.org/10.1016/j.sleep.2019.09.018

Gao, R., Peterson, E. J., & Voytek, B. (2017). Inferring synaptic excitation/inhibition balance from field potentials.
NeuroImage, *158*, 70–78. https://doi.org/10.1016/j.neuroimage.2017.06.078

Ghilardi, M.-F., Ghez, C., Dhawan, V., Moeller, J., Mentis, M., Nakamura, T., Antonini, A., & Eidelberg, D. (2000). Pat-
terns of regional brain activation associated with different forms of motor learning. *Brain Research*, *871*(1), 127–145.
https://doi.org/10.1016/S0006-8993(00)02365-9

Hill, A. T., Clark, G. M., Bigelow, F. J., Lum, J. A. G., & Enticott, P. G. (2022). Periodic and aperiodic neural activity
displays age-dependent changes across early-to-middle childhood. *Developmental Cognitive Neuroscience*, *54*,
101076. https://doi.org/10.1016/j.dcn.2022.101076

Holmberg, K., & Hjern, A. (2006). Health complaints in children with attention-deficit/hyperactivity disorder. *Acta
Paediatrica, International Journal of Paediatrics*, *95*(6). https://doi.org/10.1080/08035250600717121

Horváth, C. G., Szalárdy, O., Ujma, P. P., Simor, P., Gombos, F., Kovács, I., Dresler, M., & Bódizs, R. (2022). Overnight
dynamics in scale-free and oscillatory spectral parameters of NREM sleep EEG. *Scientific Reports*, *12*(1), Article 1.
https://doi.org/10.1038/s41598-022-23033-y

Huber, R., Graf, T., Cote, K. A., Wittmann, L., Gallmann, E., Matter, D., Schuderer, J., Kuster, N., Borbély, A. A., &
Achermann, P. (2000). Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep
EEG. *NeuroReport*, *11*(15), 3321.

Huttenlocher, P. R. (1979). Synaptic density in human frontal cortex—Developmental changes and effects of aging.
Brain Research, *163*(2), 195–205. https://doi.org/10.1016/0006-8993(79)90349-4

Iber, C. (2007). The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology, and Technical
Specification. *(No Title)*. https://cir.nii.ac.jp/crid/1370004237604151044

Ishii, R., Canuet, L., Ishihara, T., Aoki, Y., Ikeda, S., Hata, M., Katsimichas, T., Gunji, A., Takahashi, H., Nakahachi, T.,
Iwase, M., & Takeda, M. (2014). Frontal midline theta rhythm and gamma power changes during focused attention on
mental calculation: An MEG beamformer analysis. *Frontiers in Human Neuroscience*, *8*(JUNE).
https://doi.org/10.3389/fnhum.2014.00406

References

Jaramillo, V., Volk, C., Maric, A., Furrer, M., Fattinger, S., Kurth, S., Lustenberger, C., & Huber, R. (2020). Character-
ization of overnight slow-wave slope changes across development in an age-, amplitude-, and region-dependent
manner. *Sleep*, *43*(9). https://doi.org/10.1093/sleep/zsaa038

Jenni, O. G., & Carskadon, M. A. (2004). Spectral Analysis of the Sleep Electroencephalogram During Adolescence.
Sleep, *27*(4), 774–783. https://doi.org/10.1093/sleep/27.4.774

Jones, S. R. (2016). When brain rhythms aren't "rhythmic": Implication for their mechanisms and meaning. *Current
Opinion in Neurobiology*, *40*, 72–80. https://doi.org/10.1016/j.conb.2016.06.010

Jones, S. R., Pritchett, D. L., Sikora, M. A., Stufflebeam, S. M., Hämäläinen, M., & Moore, C. I. (2009). Quantitative
Analysis and Biophysically Realistic Neural Modeling of the MEG Mu Rhythm: Rhythmogenesis and Modulation of
Sensory-Evoked Responses. *Journal of Neurophysiology*, *102*(6), 3554–3572. https://doi.org/10.1152/jn.00535.2009

Karalunas, S. L., Ostlund, B. D., Alperin, B. R., Figuracion, M., Gustafsson, H. C., Deming, E. M., Foti, D., Antovich, D.,
Dude, J., Nigg, J., & Sullivan, E. (2022). Electroencephalogram aperiodic power spectral slope can be reliably meas-
ured and predicts ADHD risk in early development. *Developmental Psychobiology*, *64*(3), e22228.
https://doi.org/10.1002/dev.22228

Klimesch, W. (1999). EEG alpha and theta oscillations reflect cognitive and memory performance: A review and anal-
ysis. *Brain Research. Brain Research Reviews*, *29*(2–3), 169–195. https://doi.org/10.1016/s0165-0173(98)00056-3

Konofal, E., Lecendreux, M., & Cortese, S. (2010). Sleep and ADHD. *Sleep Medicine*, *11*(7).
https://doi.org/10.1016/j.sleep.2010.02.012

Korotchikova, I., Connolly, S., Ryan, C. A., Murray, D. M., Temko, A., Greene, B. R., & Boylan, G. B. (2009). EEG in the
healthy term newborn within 12hours of birth. *Clinical Neurophysiology*, *120*(6), 1046–1053.
https://doi.org/10.1016/j.clinph.2009.03.015

Kurth, S., Ringli, M., Geiger, A., LeBourgeois, M., Jenni, O. G., & Huber, R. (2010). Mapping of cortical activity in the
first two decades of life: A high-density sleep electroencephalogram study. *Journal of Neuroscience*, *30*(40).
https://doi.org/10.1523/JNEUROSCI.2532-10.2010

Laufs, H., Kleinschmidt, A., Beyerle, A., Eger, E., Salek-Haddadi, A., Preibisch, C., & Krakow, K. (2003). EEG-
correlated fMRI of human alpha activity. *NeuroImage*, *19*(4), 1463–1476. https://doi.org/10.1016/S1053-
8119(03)00286-6

Le Bon-Jego, M., & Yuste, R. (2007). Persistently active, pacemaker-like neurons in neocortex. *Frontiers in Neurosci-
ence*, *1*. https://www.frontiersin.org/articles/10.3389/neuro.01.1.1.009.2007

Lendner, J. D., Helfrich, R. F., Mander, B. A., Romundstad, L., Lin, J. J., Walker, M. P., Larsson, P. G., & Knight, R. T.
(2020). An electrophysiological marker of arousal level in humans. *eLife*, *9*, e55092.
https://doi.org/10.7554/eLife.55092

Lendner, J. D., Niethard, N., Mander, B. A., van Schalkwijk, F. J., Schuh-Hofer, S., Schmidt, H., Knight, R. T., Born, J.,
Walker, M. P., Lin, J. J., & Helfrich, R. F. (2023). Human REM sleep recalibrates neural activity in support of memory
formation. *Science Advances*, *9*(34), eadj1895. https://doi.org/10.1126/sciadv.adj1895

Little, S., Bonaiuto, J., Barnes, G., & Bestmann, S. (2019). Human motor cortical beta bursts relate to movement
planning and response errors. *PLOS Biology*, *17*(10), e3000479. https://doi.org/10.1371/journal.pbio.3000479

Lundqvist, M., Miller, E. K., Nordmark, J., Liljefors, J., & Herman, P. (2024). Beta: Bursts of cognition. *Trends in Cogni-
tive Sciences*, *0*(0). https://doi.org/10.1016/j.tics.2024.03.010

Maschke, C., Duclos, C., Owen, A. M., Jerbi, K., & Blain-Moraes, S. (2023). Aperiodic brain activity and response to
anesthesia vary in disorders of consciousness. *NeuroImage*, *275*, 120154.
https://doi.org/10.1016/j.neuroimage.2023.120154

McSweeney, M., Morales, S., Valadez, E. A., Buzzell, G. A., Yoder, L., Fifer, W. P., Pini, N., Shuffrey, L. C., Elliott, A. J.,
Isler, J. R., & Fox, N. A. (2023). Age-related trends in aperiodic EEG activity and alpha oscillations during early- to
middle-childhood. *NeuroImage*, *269*, 119925. https://doi.org/10.1016/j.neuroimage.2023.119925

Meyer, M., Endedijk, H. M., van Ede, F., & Hunnius, S. (2019). Theta oscillations in 4-year-olds are sensitive to task
engagement and task demands. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-019-42615-x

Michels, L., Bucher, K., Lüchinger, R., Klaver, P., Martin, E., Jeanmonod, D., & Brandeis, D. (2010). Simultaneous
EEG-fMRI during a working memory task: Modulations in low and high frequency bands. *PLoS ONE*, *5*(4).
https://doi.org/10.1371/journal.pone.0010298

References

Mitchell, D. J., McNaughton, N., Flanagan, D., & Kirk, I. J. (2008). Frontal-midline theta from the perspective of hippocampal 'theta'. *Progress in Neurobiology*, *86*(3), 156–185. https://doi.org/10.1016/j.pneurobio.2008.09.005

Mourtazaev, M. S., Kemp, B., Zwinderman, A. H., & Kamphuisen, H. A. C. (1995). Age and Gender Affect Different Characteristics of Slow Waves in the Sleep EEG. *Sleep*, *18*(7), 557–564. https://doi.org/10.1093/sleep/18.7.557

Nir, Y., Staba, R. J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I., & Tononi, G. (2011). Regional Slow Waves and Spindles in Human Sleep. *Neuron*, *70*(1), 153–169. https://doi.org/10.1016/j.neuron.2011.02.043

Ostlund, B. D., Alperin, B. R., Drew, T., & Karalunas, S. L. (2021). Behavioral and cognitive correlates of the aperiodic (1/f-like) exponent of the EEG power spectrum in adolescents with and without ADHD. *Developmental Cognitive Neuroscience*, *48*, 100931. https://doi.org/10.1016/j.dcn.2021.100931

Ostlund, B. D., Donoghue, T., Anaya, B., Gunther, K. E., Karalunas, S. L., Voytek, B., & Pérez-Edgar, K. E. (2022). Spectral parameterization for studying neurodevelopment: How and why. *Developmental Cognitive Neuroscience*, *54*, 101073. https://doi.org/10.1016/j.dcn.2022.101073

Perkel, D. H., Schulman, J. H., Bullock, T. H., Moore, G. P., & Segundo, J. P. (1964). Pacemaker Neurons: Effects of Regularly Spaced Synaptic Input. *Science*, *145*(3627), 61–63. https://doi.org/10.1126/science.145.3627.61

Pernet, C., Garrido, M. I., Gramfort, A., Maurits, N., Michel, C. M., Pang, E., Salmelin, R., Schoffelen, J. M., Valdes-Sosa, P. A., & Puce, A. (2020). Issues and recommendations from the OHBM COBIDAS MEEG committee for reproducible EEG and MEG research. *Nature Neuroscience*, *23*(12), 1473–1483. https://doi.org/10.1038/s41593-020-00709-0

Petanjek, Z., Judaš, M., Šimić, G., Rašin, M. R., Uylings, H. B. M., Rakic, P., & Kostović, I. (2011). Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proceedings of the National Academy of Sciences*, *108*(32), 13281–13286. https://doi.org/10.1073/pnas.1105108108

Pfurtscheller, G., Brunner, C., Schlögl, A., & Lopes da Silva, F. H. (2006). Mu rhythm (de)synchronization and EEG single-trial classification of different motor imagery tasks. *NeuroImage*, *31*(1), 153–159. https://doi.org/10.1016/j.neuroimage.2005.12.003

Pineda, J. A. (2005). The functional significance of mu rhythms: Translating "seeing" and "hearing" into "doing". *Brain Research Reviews*, *50*(1), 57–68. https://doi.org/10.1016/j.brainresrev.2005.04.005

Quinn, A. J., van Ede, F., Brookes, M. J., Heideman, S. G., Nowak, M., Seedat, Z. A., Vidaurre, D., Zich, C., Nobre, A. C., & Woolrich, M. W. (2019). Unpacking Transient Event Dynamics in Electrophysiological Power Spectra. *Brain Topography*, *32*(6), 1020–1034. https://doi.org/10.1007/s10548-019-00745-5

Rayson, H., Szul, M. J., El-Khoueiry, P., Debnath, R., Gautier-Martins, M., Ferrari, P. F., Fox, N., & Bonaiuto, J. J. (2023). Bursting with Potential: How Sensorimotor Beta Bursts Develop from Infancy to Adulthood. *Journal of Neuroscience*, *43*(49), 8487–8503. https://doi.org/10.1523/JNEUROSCI.0886-23.2023

Riedner, B. A., Vyazovskiy, V. V., Huber, R., Massimini, M., Esser, S., Murphy, M., & Tononi, G. (2007). Sleep homeostasis and cortical synchronization: III. A high-density EEG study of sleep slow waves in humans. *Sleep*, *30*(12), 1643–1657. https://doi.org/10.1093/sleep/30.12.1643

Ringli, M., Souissi, S., Kurth, S., Brandeis, D., Jenni, O. G., & Huber, R. (2013). Topography of sleep slow wave activity in children with attention-deficit/hyperactivity disorder. *Cortex*, *49*(1), 340–347. https://doi.org/10.1016/j.cortex.2012.07.007

Robertson, M. M., Furlong, S., Voytek, B., Donoghue, T., Boettiger, C. A., & Sheridan, M. A. (2019). EEG power spectral slope differs by ADHD status and stimulant medication exposure in early childhood. *Journal of Neurophysiology*, *122*(6), 2427–2437. https://doi.org/10.1152/jn.00388.2019

Roche, A. F., Mukherjee, D., & Guo, S. (1986). Head Circumference Growth Patterns: Birth to 18 Years. *Human Biology*, *58*(6), 893–906.

Schaworonkow, N., & Voytek, B. (2021). Longitudinal changes in aperiodic and periodic activity in electrophysiological recordings in the first seven months of life. *Developmental Cognitive Neuroscience*, *47*, 100895. https://doi.org/10.1016/j.dcn.2020.100895

Scheeringa, R., Bastiaansen, M. C. M., Petersson, K. M., Oostenveld, R., Norris, D. G., & Hagoort, P. (2008). Frontal theta EEG activity correlates negatively with the default mode network in resting state. *International Journal of Psychophysiology*, *67*(3), 242–251. https://doi.org/10.1016/j.ijpsycho.2007.05.017

References

Schneider, B., Szalárdy, O., Ujma, P. P., Simor, P., Gombos, F., Kovács, I., Dresler, M., & Bódizs, R. (2022). Scale-free and oscillatory spectral measures of sleep stages in humans. *Frontiers in Neuroinformatics*, *16*. https://www.frontiersin.org/articles/10.3389/fninf.2022.989262

Shaw, P., Kabani, N. J., Lerch, J. P., Eckstrand, K., Lenroot, R., Gogtay, N., Greenstein, D., Clasen, L., Evans, A., Rapoport, J. L., Giedd, J. N., & Wise, S. P. (2008). Neurodevelopmental trajectories of the human cerebral cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *28*(14), 3586–3594. https://doi.org/10.1523/JNEUROSCI.5309-07.2008

Shaw, P., Lerch, J., Greenstein, D., Sharp, W., Clasen, L., Evans, A., Giedd, J., Castellanos, F. X., & Rapoport, J. (2006). Longitudinal Mapping of Cortical Thickness and Clinical Outcome in Children and Adolescents With Attention-Deficit/Hyperactivity Disorder. *Archives of General Psychiatry*, *63*(5), 540–549. https://doi.org/10.1001/archpsyc.63.5.540

Shekleton, J. A., Rajaratnam, S. M. W., Gooley, J. J., Van, R. E., Czeisler, C. A., & Lockley, S. W. (2013). Improved Neurobehavioral Performance during the Wake Maintenance Zone. *Journal of Clinical Sleep Medicine*, *09*(04), 353–362. https://doi.org/10.5664/jcsm.2588

Sherman, M. A., Lee, S., Law, R., Haegens, S., Thorn, C. A., Hämäläinen, M. S., Moore, C. I., & Jones, S. R. (2016). Neural mechanisms of transient neocortical beta rhythms: Converging evidence from humans, computational modeling, monkeys, and mice. *Proceedings of the National Academy of Sciences*, *113*(33), E4885–E4894. https://doi.org/10.1073/pnas.1604135113

Shin, H., Law, R., Tsutsui, S., Moore, C. I., & Jones, S. R. (2017). The rate of transient beta frequency events predicts behavior across tasks and species. *eLife*, *6*, e29086. https://doi.org/10.7554/eLife.29086

Smith, J. R. (1938). The Electroencephalogram During Normal Infancy and Childhood: I. Rhythmic Activities Present in the Neonate and Their Subsequent Development. *The Pedagogical Seminary and Journal of Genetic Psychology*, *53*(2), 431–453. https://doi.org/10.1080/08856559.1938.10533820

Snipes, S., Krugliakova, E., Meier, E., & Huber, R. (2022). The theta paradox: 4-8 Hz EEG oscillations reflect both sleep pressure and cognitive control. *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.1063-22.2022

Snipes, S., Meier, E., Accascina, S., & Huber, R. (2024). Extended wakefulness alters the relationship between EEG oscillations and performance in a sustained attention task. *Journal of Sleep Research*, *n/a*(n/a), e14230. https://doi.org/10.1111/jsr.14230

Snipes, S., Meier, E., Meissner, S. N., Landolt, H.-P., & Huber, R. (2023). How and when EEG reflects changes in neuronal connectivity due to time awake. *iScience*, *26*(7). https://doi.org/10.1016/j.isci.2023.107138

Somsen, R. J. M., van't Klooster, B. J., van der Molen, M. W., van Leeuwen, H. M. P., & Licht, R. (1997). Growth spurts in brain maturation during middle childhood as indexed by EEG power spectra. *Biological Psychology*, *44*(3), 187–209. https://doi.org/10.1016/S0301-0511(96)05218-0

Stroganova, T. A., Orekhova, E. V., & Posikera, I. N. (1999). EEG alpha rhythm in infants. *Clinical Neurophysiology*, *110*(6), 997–1012. https://doi.org/10.1016/S1388-2457(98)00009-1

Strogatz, S. H., Kronauer, R. E., & Czeisler, C. A. (1987). Circadian pacemaker interferes with sleep onset at specific times each day: Role in insomnia. *American Journal of Physiology - Regulatory Integrative and Comparative Physiology*, *253*(1 (22/1)). https://doi.org/10.1152/ajpregu.1987.253.1.r172

Sun, H., Ye, E., Paixao, L., Ganglberger, W., Chu, C. J., Zhang, C., Rosand, J., Mignot, E., Cash, S. S., Gozal, D., Thomas, R. J., & Westover, M. B. (2023). The sleep and wake electroencephalogram over the lifespan. *Neurobiology of Aging*, *124*, 60–70. https://doi.org/10.1016/j.neurobiolaging.2023.01.006

Tal, I., Neymotin, S., Bickel, S., Lakatos, P., & Schroeder, C. E. (2020). Oscillatory Bursting as a Mechanism for Temporal Coupling and Information Coding. *Frontiers in Computational Neuroscience*, *14*. https://doi.org/10.3389/fncom.2020.00082

Tononi, G., & Cirelli, C. (2003). Sleep and synaptic homeostasis: A hypothesis. *Brain Research Bulletin*, *62*(2), 143–150. https://doi.org/10.1016/j.brainresbull.2003.09.004

Tononi, G., & Cirelli, C. (2014). Sleep and the Price of Plasticity: From Synaptic and Cellular Homeostasis to Memory Consolidation and Integration. *Neuron*, *81*(1), 12–34. https://doi.org/10.1016/j.neuron.2013.12.025

Tran, Y., Craig, A., Craig, R., Chai, R., & Nguyen, H. (2020). The influence of mental fatigue on brain activity: Evidence from a systematic review with meta-analyses. *Psychophysiology*, *57*(5), e13554. https://doi.org/10.1111/psyp.13554

References

Tröndle, M., Popov, T., Dziemian, S., & Langer, N. (2022). Decomposing the role of alpha oscillations during brain maturation. *eLife*, *11*, e77571. https://doi.org/10.7554/eLife.77571

van Ede, F., Quinn, A. J., Woolrich, M. W., & Nobre, A. C. (2018). Neural Oscillations: Sustained Rhythms or Transient Burst-Events? *Trends in Neurosciences*, *41*(7), 415–417. https://doi.org/10.1016/j.tins.2018.04.004

Volk, C., Jaramillo, V., Merki, R., O'Gorman Tuura, R., & Huber, R. (2018). Diurnal changes in glutamate + glutamine levels of healthy young adults assessed by proton magnetic resonance spectroscopy. *Human Brain Mapping*, *39*(10), 3984–3992. https://doi.org/10.1002/hbm.24225

Volk, C., Jaramillo, V., Studler, M., Furrer, M., O'Gorman Tuura, R. L., & Huber, R. (2019). Diurnal changes in human brain glutamate + glutamine levels in the course of development and their relationship to sleep. *NeuroImage*, *196*, 269–275. https://doi.org/10.1016/j.neuroimage.2019.04.040

Vyazovskiy, V. V., Olcese, U., Lazimy, Y. M., Faraguna, U., Esser, S. K., Williams, J. C., Cirelli, C., & Tononi, G. (2009). Cortical Firing and Sleep Homeostasis. *Neuron*, *63*(6), 865–878. https://doi.org/10.1016/j.neuron.2009.08.024

Wascher, E., Rasch, B., Sänger, J., Hoffmann, S., Schneider, D., Rinkenauer, G., Heuer, H., & Gutberlet, I. (2014). Frontal theta activity reflects distinct aspects of mental fatigue. *Biological Psychology*, *96*, 57–65. https://doi.org/10.1016/j.biopsycho.2013.11.010

Wessel, J. R. (2020). β-Bursts Reveal the Trial-to-Trial Dynamics of Movement Initiation and Cancellation. *Journal of Neuroscience*, *40*(2), 411–423. https://doi.org/10.1523/JNEUROSCI.1887-19.2019

Wilhelm, I., Kurth, S., Ringli, M., Mouthon, A.-L., Buchmann, A., Geiger, A., Jenni, O. G., & Huber, R. (2014). Sleep Slow-Wave Activity Reveals Developmental Changes in Experience-Dependent Plasticity. *Journal of Neuroscience*, *34*(37), 12568–12575. https://doi.org/10.1523/JNEUROSCI.0962-14.2014

Wilkinson, C. L., Yankowitz, L. D., Chao, J. Y., Gutiérrez, R., Rhoades, J. L., Shinnar, S., Purdon, P. L., & Nelson, C. A. (2024). Developmental trajectories of EEG aperiodic and periodic components in children 2–44 months of age. *Nature Communications*, *15*(1), 5788. https://doi.org/10.1038/s41467-024-50204-4

Zich, C., Quinn, A. J., Mardell, L. C., Ward, N. S., & Bestmann, S. (2020). Dissecting Transient Burst Events. *Trends in Cognitive Sciences*, *24*(10), 784–788. https://doi.org/10.1016/j.tics.2020.07.004

Zimmermann, P., & Fimm, B. (2012). *Testbatterie zur Erfassung von Aufmerksamkeitsstörungen (TAP 2.3.)*. Psytest.