Stimulation of saliva affects the release of aroma in wine; a study of microbiota,

biochemistry and participant origin

Xinwei Ruan[1], Yipeng Chen[1], Aafreen Chauhan[1] and Kate S. Howell[1*]

[1]School of Agriculture, Food and Ecosystem Sciences, Faculty of Science, University of Melbourne,

Parkville 3010, Australia

Xinwei Ruan: ORCID: 0000-0002-1963-3807

Yipeng Chen: ORCID: 0009-0002-7511-374X

Aafreen Chauhan: aschauhan@student.unimelb.edu.au

* Corresponding author: Kate S. Howell

Email: khowell@unimelb.edu.au; **ORCID:** 0000-0001-6498-0472

Abstract

Saliva influences the release of aroma in the oral cavity. The composition of human saliva varies

depending on stimulation and host's origin; however, the compositional differences of saliva and their

influences on aroma release have not been fully evaluated. In this study, we recruited 30 healthy adults

(15 Australians and 15 Chinese) and collected saliva samples at three stages: before, during, and after

stimulation. Salivary samples were characterised by flow rate, total protein concentration, esterase

activity, microbiome composition by full-length 16S rRNA gene sequencing, and the ability to release

aroma from wine by headspace solid-phase microextraction gas chromatography–mass spectrometry

(HS-SPME-GC-MS). Differences in salivary composition and specific wine volatiles were found

between Australian and Chinese participants, and amongst the three stimulation stages. Significant

correlations between the relative abundance of 3 bacterial species and 10 wine volatiles were observed.

Our results confirm the influence of participant's geographic origin and stimulation on salivary

composition, highlighting the role of salivary components, especially salivary bacteria, on the release

of aroma from wine.

Keywords: Microbiota, Salivary microbiome, Geographic variation, Aroma perception,

Preference

1   Introduction

Saliva plays an important role in aroma release from foods and beverages [1-3]. When mixing with saliva in oral cavity, volatile compounds are released from food matrix and then travel to olfactory receptors via a retronasal route. Physiochemically, saliva has a dilution effect on aroma due to continuous salivation [4]. Saliva contains salt, sugar and the proteinaceous fraction of mucin, which may influence how volatiles partition in a solution [5]. Further biochemical effects by degrading odorants or releasing aroma from aroma precursors have been reported [6-8]. Two critical components of saliva, proteins and microbiota, are implicated with release or formation of aroma compounds and thus affect aroma release from food

Proteins are able to bind, trap and interact with volatile compounds, thereby influencing the aroma release [9]. Two abundant proteins in whole human saliva, mucin and α-amylase, modify release of aroma compounds through hydrophobic interactions [10]. Pérez-Jiménez, Rocha-Alcubilla and Pozo-Bayón [11] demonstrated the efficacy of esterase in human saliva to degrade esters by comparing whole saliva with enzymatically inhibited saliva. Large inter-individual variations have been observed in the total salivary esterase activity [12]. The chemical structure of esters also influences the extent of enzymatic degradation [6]. In addition, the effect of salivary enzymatic reactions on aldehydes [7] and thiols [6] has also been demonstrated by comparing the aroma release ability of thermal-treated saliva with untreated saliva.

In contrast to salivary proteins, the role of oral bacteria in aroma release is not well characterised. The oral microbiota contribute to the ability of human saliva to hydrolyse odourless grape glycosides [8, 13]. A previous study [14] identified the differences in salivary microbial profile between obese and normal-weight individuals. However, no association was established between the differences of these two groups in aroma release of wine and the microbial profiles. In recent years, the advances in culture-independent methods for oral microbiome identification have deepened our understanding of microorganisms in the oral cavity [15] and reveal the salivary microbial composition with a stable core bacterial community shared by geographically diverse populations [16]. Increased accuracy of taxonomic identification would facilitate to study the role of salivary bacteria in aroma release.

Salivary protein and microbiota are likely to be dynamic during oral processing. Saliva production

switches between stimulated and unstimulated depending on the presence of stimuli. During stimulation

evoked by taste or mastication, the saliva secretion volume increases [17]. More than 90% of saliva comes

from three pairs of major salivary glands, including the submandibular, sublingual, and parotid glands.

The saliva produced by these three glands differs in composition and function because of the cells that

constitute the glands [18]. For example, the parotid glands secrete watery fluids since they exclusively

consist of serous acinar cells, while the sublingual glands produce viscous saliva rich in mucins and

other glycoproteins due to the high proportion of mucous acinar cells [19]. Therefore, the salivary

composition varies depending on the contribution of different glands with or without stimulation [20]. It

has been documented that parotid glands have a higher secretory rate under stimulation than in

unstimulated conditions. Meanwhile, the submandibular and sublingual glands produce more saliva

under unstimulated conditions [21].

Recently, the published literature has offered contradictory findings on whether the microbial diversity

in saliva is different between unstimulated and stimulated status [22-24]. Specifically, the alpha- and beta-

diversity of salivary microbiome were reported as significantly different by Gomar-Vercher, Simón-

Soro, Montiel-Company, Almerich-Silla and Mira [22]. In contrast, the other two studies observed no

significance [23, 24]. Significant differences have also been observed by previous studies between saliva

collected before and during stimulation in various salivary biochemical parameters, including protein

concentration [25], antioxidant activity [25], esterase activity [26]. Meanwhile, the composition of saliva

collected after stimulation by food or other stimuli have not been characterised. Since the after-odour

and after-taste generated after the swallowing or expectoration of food is also considered an essential

sensory parameter [27, 28], studying the salivary composition after stimulation may give insight into

understanding this process.

Meanwhile, the microbiota and type of protein in saliva varies among populations and is also shaped

by the host characteristics, such as age [29], gender [30], body mass index (BMI) [31] and other lifestyle factors

[32]. The correlation of a host's ethnicity on the bacterial salivary composition has received considerable

critical attention. Mason, Nagaraja, Camerlengo, Joshi and Kumar [33] investigated the oral microbial

profile of 192 participants belonging to Chinese, Latinos, non-Hispanic whites, and non-Hispanic

blacks, finding the lower alpha diversity of non-Hispanic blacks than the other three groups. Yang,

Zheng, Cai, Shrubsole, Pei, Brucker, Steinwandel, Bordenstein, Li and Blot [34] observed the differences

in overall microbial composition between African Americans and European Americans. A recent study

documented different microbial alpha and beta diversity of supragingival plaque microbiome among 96

children from four ethnic groups, including African American, Burmese, Caucasian and Hispanic [35]. A

study conducted by our group demonstrated the differences between Australian and Chinese wine

experts' salivary protein profile and their sensory responses [36]. The mode of variation between

Australian and Chinese saliva has not been established, but our preliminary data suggest that dietary

differences might be correlated with these differences (data not shown).

Food preference is strongly influenced by a consumer's cultural, geographic and personal background

[37, 38]. The preferences of Australian and Chinese consumers for 14 red wines have been compared by

Williamson, Robichaud and Francis [37], finding variance in some attributes for liking, such as "alcohol",

"sweetness" and "dark fruit". Aroma perception is a prominent factor that influences the acceptability

and preference of customers for different wines [39]. However, much uncertainty exists about the

relationships between host factors, salivary composition, and aroma. We ask if the functional properties

of diverse salivary microbiome communities can provide insight into the differences in sensory

perception amongst wine consumers.

To enhance our understanding of saliva composition and aroma perception, we analysed samples from

30 participants from Australia and China and mixed the saliva samples with wine to observe the ability

of human saliva to release aroma compounds from wine. Each participant provided three types of saliva

samples collected before, during and after mechanical stimulation to extend our knowledge of reactions

that occur in the oral cavity during stages of appreciating wine. Enzymatic activity tests and full-length

16S rRNA sequencing highlighted the differences of saliva type and origin of participants on salivary

composition. Combining these findings with the wine aroma profile detected by headspace solid-phase

microextraction gas chromatography-mass spectrometry (HS-SPME-GC-MS), we demonstrate

differences in salivary proteins and microbiome due to the mode of salivary stimulation and how this

varies based on geographic origin of the participant.

2   Material and Methods

2.1   Materials and reagents

All reagents and chemicals used in this study were of analytical grade. Methanol, citric acid, 4-

Nitrophenyl acetate, sodium chloride, sodium phosphate, sodium bicarbonate, potassium chloride,

potassium phosphate dibasic trihydrate, calcium chloride dihydrate, mucin from porcine stomach,

phosphate-buffered saline, Tris-buffered saline, 4-octanol, amylase activity assay kit, and standards for

volatile compounds detected in GC-MS analysis was obtained from Sigma-Aldrich (Castle Hill NSW,

Australia). Micro bicinchoninic acid (BCA) protein assay kit was acquired from Thermo Fisher

Scientific (Waltham, MA). Pure water was obtained from a Milli-Q system (Millipore Australia,

Bayswater, Victoria, Australia). A Shiraz wine (Ladies Who Shoot Their Lunch; vintage 2019; 14.9%

v/v alcohol) produced in Victoria, Australia, was purchased from a local wine producer, and used in all

GC-MS assays with wine (section 2.6). This wine was selected because different aroma perception were

observed between Chinese and Australian panellists in a previous sensory evaluation [36].

2.2   Study cohort

We recruited 30 healthy adults to meet the following criteria: (i) self-reported good systemic health (i.e.,

no diagnosed diabetes, cardiovascular diseases, endocrine disorders, or hypertension); (ii) no clinically

diagnosed oral diseases; (iii) not receiving antibiotic or steroid hormone use within the previous three

months; (iv) no smoking habits (less than several times a month); (v) not a regular drinker (less than

several times a week); (vi) have a BMI index no higher than 24 kg/m^2. The sample collection was

performed at the University of Melbourne, Australia. An online questionnaire was provided to all

volunteers to ensure they met the inclusion criteria. Demographics were also acquired using self-

reported questionnaires before the collection. Fifteen Chinese and fifteen Australian participants (aged

18-28) joined the study. All participants resided in Melbourne, Australia. The participants were grouped

according to their country of birth and residence history. The Australian group includes participants

born in Australia and have spent most of their lives there. The Chinese group contains participants born

in China and have lived in Australia for less than five years. The ethnic origin of the participant's parents

or grandparents was not collected nor considered in this study. Participants self-identified as 'Chinese'

or 'Australian'. The gender and BMI index of participants were balanced in each group **(Table S1)**.

This study was approved by the University of Melbourne Human Research Ethics Committees (Review

Reference Number: 2022-24220-29889-3).

2.3   Sample collection

Before collection, sampling procedures were explained in detail to all participants. Participants were

asked to refrain from eating and drinking any beverage except water for an hour prior to providing

saliva samples. Three types of saliva were collected before (Phase A), during (Phase B), and after (Phase

C) stimulation. The collection procedure is illustrated in **Figure S1**. For Phase A, participants naturally

expectorated the unstimulated saliva into a 50 mL sterile collection tube until the volume reached 5 mL.

For Phase B, participants were instructed to chew a piece of paraffin wax and expectorated the

stimulated saliva into another collection tube in 5 minutes. Finally, after removing paraffin wax from

the oral cavity, the participant repeated the saliva collection procedure described in Phase A. The saliva

samples collected after stimulation were named as Phase C. Time was recorded in Phases A and C to

calculate the flow rate as volume collected per minute (mL/min). For the same participant, Phase A, B,

and C collections must be completed consecutively on the same day. Immediately after the collection,

saliva samples were separated in smaller aliquots and kept frozen at -80 $^\circ$C until analysis.

2.4   Biochemical analyses

Salivary total protein content, and enzymatic activities of 90 samples were tested individually to find

the differences among different types of saliva. All analyses were performed in triplicates. Protein

concentration of whole saliva samples was tested by Micro BCA™ Protein Assay Kit (Thermo

Scientific™) using bovine serum albumin as standard for calibration.

For the enzymatic activity measurements, saliva samples were centrifuged at 15,000 ×g for 15 min at

4 °C to remove insoluble matter. All enzymatic activity assays were performed in triplicate. Total

salivary esterase activity was conducted on individual samples accordng to the method from Pérez-

Jiménez, Muñoz-González and Pozo-Bayón [26]. α-Amylase activity assay was performed following the

user manual of an Amylase Activity Assay Kit (Sigma-Aldrich), we pooled the samples for this assay

because of the limited material for some participants.

2.5   Full length 16S rRNA gene sequencing

Bacterial genomic DNA was extracted from 90 whole human saliva samples using the QIAGEN®

DNeasy® PowerSoil® Kit. The concentration of genomic DNA was measured using a Qubit HS DNA

kit (Thermo Scientific™) and normalised to 1 ng/μL prior to Polymerase Chain Reaction (PCR).

Following PacBio's recommended PCR protocol, full-length 16S amplicons were generated using the

Platinum™ SuperFi II High-fidelity PCR enzyme (Thermo Scientific™), with 2ng of input DNA. The

amplicons generated were visualised on a 2% agarose gel and pooled. SMRTbell™ cleanup

beads (PacBio) were used to purify the pooled amplicons according to the PacBio protocol. The

amplicon library was contracted with the SMRTbell® prep kit 3.0 library prep kit (PacBio) following

the protocol. The library was then polymerase-bounded using the Revio™ Polymerase Kit (PacBio) and

loaded on a single-molecule real-time (SMRT) Cell 25M (PacBio). 16S rRNA full-length sequencing

was performed for 30 hours on the PacBio Revio instrument.

2.6   Gas chromatography-mass spectrometry (GC-MS) analysis for whole human saliva

Different types of saliva samples were pooled together for the headspace solid-phase microextraction

GC-MS analyses (HS-SPEM-GC-MS) to overcome the inter-individual variation in saliva composition

and the availability of saliva from each participant. Saliva was pooled based on stimulation phase, by

participant origin, resulting in 6 groups of saliva samples: Australia-Phase A, Australia-Phase B,

Australia-Phase C, Chinese-Phase A, Chinese-Phase B and Chinese-Phase C. Each sample type was

measured in triplicate.

Three types of controls were used: the mixture of wine and water, the mixture of wine and artificial

saliva, and the mixture of six pooled saliva samples and model wine. The mixture of wine and water

were used to control for the dilution effect of saliva in the GC analysis matrix [4, 40]. An artificial saliva

solution was spiked with wine to control for the effect of mucin and salts in aroma release. It was

prepared according to the method of Genovese, Piombino, Gambuti and Moio [1]. The artificial saliva

samples were freshly prepared by dissolving mucin (2.16 g/L), $NaHCO_3$ (5.208g/L), $K_2HPO_4 \cdot 3H_2O$

(1.369 g/L), NaCl (0.877 g/L), KCl (0.477 g/L) and $CaCl_2 \cdot 2H_2O$ (0.441 g/L) in Milli-Q water. The

mixture of the model wine and six types of saliva were used to control for the confounder effect of the

volatiles that may present in saliva itself. The model wine used in this study was prepared by saturating

12% ethanol (v/v) with potassium hydrogen tartrate, adjusting to pH 3.0 by 40% (w/v) tartaric acid.

Each mixture was measured in triplicate.

The static headspace volatile extraction method was conducted following the method of Muñoz-

González, Feron, Guichard, Rodríguez-Bencomo, Martín-Álvarez, Moreno-Arribas and Pozo-Bayón [41]

with some modifications. The 20 mL GC vial was incubated at 37°C for 20 min to stabilise the

temperature of the reaction environment. Then 5 mL of wine and 1 mL of pooled saliva were added to

the vial with 10 μL of internal standard (100 mg/L 4-octanol). The wine-saliva mixture was incubated

at 37 °C for 12 min. The headspace volatile extraction from the GC vial was performed by a 65 μL

polydimethylsiloxane/divinylbenzene (PDMS/DVB) SPME fibre with 1 cm fibre length (Supelco,

Bellefonte, PA) at 37 °C for 5 min.

GC-MS analysis was carried out with an Agilent 7890A GC system (Agilent Technologies, Mulgrave,

VIC, Australia) coupled with an Agilent 5975C MS (Agilent Technologies). The fibre was desorbed at

220 °C in splitless mode at 8.7 psi for 30 s. Chromatographic separation of volatile compounds were

performed with a DB-Wax polar capillary column (Agilent Technologies; 30 m × 0.25 mm × 0.25 μm

film thickness). Helium was used as the carrier gas at a flow rate of 0.7 mL/min. After initially holding

the oven temperature at 40°C for 4 mins, it then ramped at 3°C/min to 220 °C, and held at 220 °C for

10 min. For the MS system, electron ionisation mode at 70 eV was used to record the electron impact

mass spectra. The scanning mass range was from m/z 35 to 350. The temperature of the MS interface,

MS source, MS quadrupole was 250, 230 and 150 °C.

The volatile compounds identification was based on the retention indices (RIs) with the facility of NIST

library (version 11.0). RIs were obtained by calculating the retention times of alkane standards (C_5–C_{30})

on the same column. Peaks were semi-quantified using the internal standard, 4-octanol. The

concentrations of volatiles were expressed as μg/L 4-octanol equivalence. Among the 27 compounds

identified, sixteen compounds have also been quantified using external standards to validate the results

of the semi-quantified method, facilitated by the calibration curve of standard compounds as:

217
$$Concentration \ (g/mL) = \left(\frac{peak \ area \ of \ identified \ compound}{peak \ area \ of \ internal \ standard} \times slope \right) + intercept$$

2.7   Statistical analyses

The data analyses of salivary biochemical parameters and the volatile compounds generated from the

wine-saliva mixture were conducted in R (version 4.1.0). Student t-tests were used to examinate the

significant ($p < 0.05$) differences between samples from Chinese and Australian participants. Two-way

analysis of variance (ANOVA) with Tukey's comparison were performed to test the significant ($p <$

0.05) differences among saliva samples collected before, during and after stimulation. The Spearman's

correlations were calculated with *rcorr* function in the R package *Hmisc* [42].

For bioinformatic analyses, the demultiplexed PacBio HiFi full-length 16S rRNA gene sequences are

quality filtered using QIIME2 [43]. DADA2 [44] was used to denoise the sequences into amplicon single

variants (ASVs) using the function "dada2 denoise-ccs". A phylogenetic tree was generated from the

mafft alignment. The naïve Bayesian machine learning based classification was performed using the

expanded Human Oral Microbiome Database (eHOMD) version 15.23 [45]. If a species-level match was

not found, it was further classified based on three other databases that successively refer to the next one.

The three databases are GTDB r207, Silva v138 [46] and the NCBI RefSeq 16S rRNA database

supplemented by the Ribosomal Database Project (RDP). The resulting objects were imported to R

(version 4.1.0) for further analyses. Alpha diversity was calculated with Simpson's diversity index.

Wilcoxon rank-sum tests were used to determine the statistically significant ($p < 0.05$) differences. Beta

diversity was assessed using Bray-Curtis dissimilarly. Permutational multivariate analysis of variance

(PERMANOVA) with 999 permutations was conducted to examine the statistically significant ($p <$

0.05) difference in beta-diversity using the *adonis2* function of *vegan* package [47]. The differential taxa

were defined from phylum level to species level by *metacoder* package [48] with false discovery rate-

adjusted Wilcoxon rank-sum test. ANCOM-BC2 [49] was performed to identify the differential species

with the default parameters.

3   Results

3.1   Simulated saliva (Phase B) has a distinguishable protein and biochemical composition

3.1.1   Differences in saliva composition amongst the three phases

To determine whether human saliva produced before, during and after stimulation is different, the

biochemical properties and microbial composition of human saliva samples were tested. The salivary

flow rates were calculated as volume collected per minute, reflecting the speed of participants to

produce saliva. *Ex vivo* measurements of α-amylase and esterase were conducted to assess enzymatic

activity of the centrifuged saliva. The flow rate observed in Phase B and total salivary esterase activity

were significantly higher than the other two phases, while Phase A and C saliva were not significantly

different **(Figure 1A, E)**. For the total protein concentration of saliva samples, the value of Phase A

was significantly higher than that of the other two phases **(Figure 1C)**. No significant differences were

found among the three groups in the amylase activity **(Table S2)**.

Full-length 16S rRNA gene sequencing was used to detect the differences in the salivary microbiome.

In contrast to the protein concentration and enzymatic activities, no significant differences were

detected among Phases A, B and C regarding the alpha-diversity and beta-diversity of the microbial

communities ($p > 0.05$). Although several species have been identified as differential taxa **(Figure**

S2). Most differential species had the lowest relative abundance in Phase B **(Figure S2A)**. The

differential species identified in Phase A and C were clustered together in the heatmap **(Figure S2B)**,

suggesting the microbial profile is distinct in Phase B at species level.

As the volatiles may also have originated from the human oral cavity, the volatiles in the saliva samples

were assessed by mixing each pooled saliva sample with model wine. Among the detected volatiles, (–)

– clovene was found in the mixture of saliva and wine and the mixture of saliva and model wine.

Therefore, this compound was excluded from further analysis.

3.1.2   Effect of stimulation on the release of aroma from wine

To evaluate the ability of different saliva samples to release aroma compounds from wine, HS-SPME-

GC-MS analyses were performed on the same wine sample spiked with saliva samples. Saliva samples

were pooled into six groups: Australia-Phase A, Australia-Phase B, Australia-Phase C, Chinese-Phase

A, Chinese-Phase B and Chinese-Phase C. After mixing the six pooled saliva samples with wine,

twenty-seven volatile compounds were identified, consisting of 12 esters, 6 alcohols, 3 acids, 2

benzenoids and 4 terpenes **(Table 1)**. The wine sample was also mixed with water and artificial saliva

as the control groups **(Figure S3A)**. For many compounds, especially esters, the addition of artificial

saliva induced a significant decrease in their concentration compared to the wine-water mixture.

Depending on the type of pooled saliva added, the decrease caused by artificial saliva could either be

counteracted or further decreased **(Figure S3B)**. It suggests the whole human saliva induces other

reactions with the wine matrix other than the effect of mucin.

For the 26 included volatiles, twenty-one volatiles were constantly detected across all six groups, while

the other 5 volatiles were only occasionally found **(Table 1)**. Methyl hexanoate is the only ester that

was not constantly detected in all groups. The other occasionally detected compounds are all benzenoids

and terpenes. These compounds were mostly absent in Phases A and C samples except limonene, which

was not detected in the Australian-Phase B group. The one-way ANOVA conducted with the

concentration of 16 volatiles quantified using external standards is compatible with the results

calculated using semi-quantification method **(Table 1; Table S3)**.

The dataset was divided into two subgroups, Australian **(Figure 2A)** and Chinese **(Figure 2B)**, to show

the changing trend for 21 constant-detected compounds among Phases A, B and C. The statistical

analyses suggest that the Australian and Chinese groups individually have 9 and 8 compounds with

significantly higher concentrations in Phase B than in the other two phases ($p < 0.05$). Most of the

distinguishing compounds are esters, while octanoic acid and 1,3-ditertiary butyl benzene show the

same trend in either the Australian or both groups. In addition, several compounds demonstrated

significant differences between Phase A and C, including diethyl succinate, 3-methylbutyl acetate, 3-

ethyl-2-pentanol and acetic acid. 3-Ethyl-2-pentanol and acetic acid only showed a significant

difference among phases in the Australian group, with significantly lower concentration in Phase A (p

< 0.05). Further, a significant difference was observed between the Australian and Chinese groups of

these two compounds in Phase A **(Figure 2C)**.

To better understand the relationship between three types of saliva and the aroma released from wine,

a principal component analysis (PCA) was performed on the 21 detected compounds shared amongst

all samples **(Figure 2D)**. The internal standard equivalent concentration of the aroma compounds,

(expressed as μg/L 4-octanol equivalents), distinguished the saliva samples from different phases. The

first two principal components explained 68.57% of the variance. The first principal component (PC1)

separated the samples of Phase B from those of Phase A and C. The samples of Phase B were highly

associated with most esters, 1,3−ditertiarybutylbenzene and 3−ethyl−2−pentanol, positioned in the

space defined by negative PC1. On the contrary, 1-butanol was positively related to PC1. The second

principal component (PC2) was negatively associated with most alcohols, including 2-methyl-1-

propanol, 3-methyl-1-butanol, 1-hexanol and 2,3-butanediol. Phase A and C samples were separated on

PC2 with some overlap retained, suggesting the interaction between the two investigated covariates,

phase, and subject origin **(Table S2)**. In addition, PCAs were conducted using a total 27 compounds

and 16 compounds quantified using external standards, separately **(Figure S4)**. These results agree with

the PCA constructed by the semi-quantified compounds.

3.2 The composition of saliva from Australian and Chinese participants is different

3.2.1 Differences in salivary proteins between two groups

The flow rate, total protein concentration, and activities of alpha-amylase and esterase of the saliva

samples collected from Australian and Chinese participants were compared to investigate the influence

of the host's cultural background on salivary composition. Although no differences were found between

the flow rate of Australian and Chinese participants' saliva **(Figure 1B)**, the saliva of Australian

participants demonstrated higher total protein concentration **(Figure 1D)**. For the enzymatic activities,

the two groups demonstrated no statistical differences in either amylase or esterase activity **(Table S2;**

Figure 1F).

3.2.2 Differences in salivary microbiome profiles

For the comparison of the salivary microbial profile of participants from Australia and China, the full-

length 16rRNA gene was sequenced. After denoising by DADA2, the samples were rarefied to 5,232,

resulting in 11,052 amplicon sequence variants (ASVs). The Australian and Chinese groups differ in

the alpha-diversity characterised by Simpson Index **(Figure 4A)**. Beta-diversity analysis conducted

using Bray-Curtis dissimilarity also demonstrated the significant differences between these two groups

(PERMANOVA, $p = 0.001$). The principal coordinate analysis (PCoA) result showed the separation

between the microbial composition of samples from two groups **(Figure 4B)**.

When investigating the taxa that discriminate saliva samples from Australian and Chinese participants,

we first compared the relative abundance of the two groups at phyla **(Figure 3A)** and genus

levels **(Figure 3B)**. Firmicutes, the phylum with the highest relative abundance in both groups, showed

higher relative abundance in the saliva samples from Australian participants. A higher relative

abundance of *Veillonella* in Australian saliva was observed. We performed a Wilcoxon Rank Sum test

with a false discovery rate correction to each taxon, spanning all taxonomic levels **(Figure**

3C). Genus *Neisseria*, *Lautropia*, *Porphyromonas*, *Capnocytophaga*, *Cardiobacterium*, *Filifactor* and

Parvimonas were higher in Chinese participants' saliva, with several species of these genre. All taxa

belong to phyla Synergistetes and Spirochaetes, except the species were more prevalent in the Chinese

group. In contrast, the saliva microbiota of Australian participants was distinguished mainly by taxa under the order *Veillonellales*, especially the genus *Veillonella* and *Megasphasera*. The genus with high abundances, such as *Streptococcus*, *Prevotella* and *Haemonphilus*, contain both species identified as biomarkers of Chinese and Australian participants.

We also used ANCOM-BC2 to identify the differential species between Australian and Chinese saliva to accomplish more precise results at the species level. We found 8 differential species for comparing two groups, mostly under the phylum *Bateroidetes* and *Firmicutes* **(Figure 4C)**. Among them, *Porphyromonas pasteri* and *Neisseria subflava* have higher relative abundance in the Chinese group, which agrees with the results of the Wilcoxon Rank Sum tests **(Figure 3C)**. Similarly, the six differential species enriched in Australian samples identified by ANCOM-BC2 are also found in the Wilcoxon Rank Sum tests. In the Australian group, the differential species of Phase A and C were clustered together, suggesting they are closer to each other than Phase B. Meanwhile, the differential species in the Chinese Phase A and B samples were clustered.

3.2.3   Differences in saliva composition between groups are associated with aroma release

To investigate the aroma consequences of compositional differences in the saliva of saliva samples collected from Australian and Chinese participants, the volatile profile was collected by HS-SPME-GC-MS and compared. The result of two-way ANOVA showed that 11 compounds were released differently after mixing with saliva samples from two groups. Different concentrations due to the saliva samples collected at different phases were observed **(Figure 4E)**. All differential compounds showed higher concentration in the Chinese group, especially in unstimulated Phases A and C. Meanwhile, ethyl heptanoate released after mixing with Chinese saliva samples demonstrated higher concentration in Phases A and B.

We suspected that the salivary bacteria contributed to aroma release, and so we calculated the Spearman's rank correlation between the relative abundance of the species in pooled saliva samples and the aroma compounds released after mixing the wine samples with those saliva samples. We found three species significantly correlated to wine aroma profile, including *Streptococcus sp.*

HMT006, *Prevotella shahii* and an unclassified species belongs to *Selenomonas* **(Figure 4D)**. The

concentration of three differential esters, ethyl propanoate, 3-methylbutyl acetate and ethyl hexanoate,

are positively associated with the relative abundance of *Streptococcus sp. HMT006*, while negatively

associated with the relative abundance of the unclassified species of *Selenomonas* **(Figure 4F)**. The

higher concentration of 3-ethyl-2-pentanol and several esters in the Chinese group has been positively

correlated with the increased relative abundance of *Streptococcus sp. HMT006* in saliva samples from

Chinese participants in Phase A. As the only benzenoid identified, the concentration of 1,3-

ditertiarybutylbenzene showed a negative correlation with the unclassified species

of *Selenomonas.* Although two-way ANOVA concluded that the concentration of 2,3-butanediol was

not different between the Chinese and Australian groups, it was positively associated with the higher

relative abundance of *Prevotella shahii* in the Chinese saliva

4   Discussion

In the current study, we observed the differences in salivary composition between saliva samples collect

at different stimulation stages and from participants with different geographic origin. This study further

demonstrates the association between host factors, salivary composition, and aroma release from wine.

For the two parameters controlled in this study, the phase of saliva production and the host's ethnic

background, we simultaneously observed the differences in salivary composition and the aroma profiles

of wine after mixing with saliva samples. The salivary stimulation stages and subject origin both

influence aroma release ability as the composition of saliva is different. These findings shed light on

the role of human saliva in immediate and prolonged retronasal perception of aroma and has introduced

unstimulated saliva as a control in *ex vivo* experiments.

Overall, saliva samples collected before, during and after stimulation were different in composition. We

observed a higher total protein concentration in the saliva samples produced in Phase A. The lower

protein concentration in Phase B and C may be due to the dilution of the substantial water content

secreted in Phase B [50]. Interestingly, even though the flow rate dropped significantly after removing the

stimulus from the oral cavity, the total protein concentration in saliva did not return to the level when

the stimulus was not received. Further analyses, such as proteomics should be applied to assess the

decreasing rate of individual proteins. By observing whether the decreasing rate of different proteins

are even, it is possible to infer if dilution is the main reason for the lower protein concentration in Phase

C. In contrast, a higher total salivary esterase activity was found in Phase B, while the esterase activity

of Phases A and C had no significant difference. The esterase in saliva is an important enzyme for

hydrolysing esters into carboxylic acids and alcohols [6]. The higher esterase activity in stimulated saliva

than in unstimulated saliva has also been observed by Pérez-Jiménez, Muñoz-González and Pozo-

Bayón [26]. Esterase did not follow a consistent trend with total protein concentration, suggesting the

mechanisms beyond dilution that may be responsible for modulating the concentration of different

proteins after stimulation. Meanwhile, no significant differences were observed in the current study

between the microbial diversities of saliva samples collected at different stimulation stages. The

association between oral microbiota and the type of saliva has been widely investigated by previous

studies but remains controversial. A previous study suggested a significant difference between

unstimulated and stimulated saliva in both alpha and beta diversity [22]. On the contrary, another two

studies [23, 24] reported no significant differences between the Shannon diversity index of bacteria in

unstimulated and stimulated saliva samples. Several species have been identified as differential among

the mouth-rinse water, unstimulated saliva, and stimulated saliva by Jo, Nishimoto, Umezawa, Yama,

Aita, Ichiba, Murakami, Kakizawa, Kumagai, Yamada and Fukuda [24], which agrees with our results.

However, no intersection has been found between the differential species defined by the previous work

and our study. The reason could be the different statistical tools used. It is possible that the differences

could be too subtle to detect. Further study with higher resolution, such as metagenomic sequencing,

need to be applied to monitor the sub-species level or community-level differences.

The relationship between the stimulation stages and aroma release ability was found to be focused on

the esters in wine and the esterase activity in human saliva. Esters are an essential group of aromatic

compounds to contribute fruitiness to wine [55]. Most esters detected in this study have the odour activity

values greater than 1 **(Table S3)**, meaning that these compounds are likely to have a sensory impact.

Saliva samples produced in Phase B demonstrated the highest esterase activity and induced the highest

concentration of esters released from wine samples. However, the presence of esterase was previously

associated with the hydrolysis of esters [3] and the generation of carboxylic acids [11]. Esters are relatively

less persistent in the oral cavity after swallowing, probably due to the modification by oral microbiota

directly or with enzymatic action [56]. This rather contradictory result may be explained by the presence

of reactions other than enzymatic hydrolysis of the esters. In addition, the total salivary esterase activity

could be attributed to different types of enzymes, such as carboxylesterases, carbonic anhydrases,

trypsin, and cholesterol esterases [26]. The catalytic effect of specific enzymes could be accompanied by

concurrent synthesis of esters. Interestingly, the same phenomenon was reported in our recent study [36],

which strengthens our assumption that perhaps more than one type of reaction involving esterases

occurs when human saliva mixes with esters in wine.

Notably, several volatiles showed differences between Phase A and Phase C in the Australia group,

including 3-methylbutyl acetate, 3-ethyl-2-pentanol and acetic acid. 3-Methylbutyl acetate is an ester

with fruity, banana-like aroma (**Figure S3**). In contrast, 3-ethyl-2-pentanol and acetic acid may provide

an unpleasant flavour to wine [57]. Those volatiles may play a role in the after-stimulation stage. During

retronasal perception of wine, two distinct types of aroma release are perceived by consumers: the

immediate impression produced when swallowing and the prolonged perception after swallowing [58].

By comparing the saliva samples collected during and after stimulation with the control, our study

demonstrates that it is important to specify the type of saliva collected and used in *ex vivo* experiments.

Stimulated saliva needs to be specified for the research targeting the reactions with stimulus. These

findings could also shed light on understanding the differences between immediate and prolonged

perception of wine aroma. Meanwhile, although we demonstrated the different aroma release capacities

of saliva samples collected during and after the mechanical stimulation, more complex mechanisms are

introduced in aroma perception occurs during the wine tasting experience of human beings. Mixing the

saliva samples collected after the mechanical stimulation with wine could not completely reflect the

complexity and dynamics of "after-odour" occurring in human oral cavity. We set the stopping point

for the collection of Phase C as the time that volume reaches 5 mL to allow enough time for saliva to

change from "stimulated" to "unstimulated". However, after swallowing the wine, various amounts of

odorants will be captured by oral mucosa and keep working as "stimulus" in mouth [28]. Therefore, the

changes of saliva from "stimulated" to "unstimulated" could not be as abruptly as what we mimicked

using paraffin chewing. The proportion of wine and saliva used in our static headspace SPME method

does not reflect all the consequences of oral processing to explain aroma production in the 'after-taste'

period. Further analytic approaches to simulate the relationships between post-stimulated saliva and

wine, include modelling the oral mucosa [60], and should be used to further understand the role of saliva

in the after-taste phase.

Moreover, the chemical composition of wine also influences the aroma release of wine aroma volatiles.

For example, the phenolic compounds impact the aroma release depending on the chemical structure [61,

62]. Specifically, phenolic acids facilitate the intra-oral release of β-phenyl ethanol, linalool and β-ionone,

while flavonoids in wine decrease the release of some esters [61]. Meanwhile, ethanol in wine may also

affect the salivary proteins [63]. Future research is required to explore the influences of host factors on the

aroma release ability of saliva on other food matrices, such as non-alcoholic beverages and solid foods.

The influence of the host's origin on salivary composition, especially the oral microbiome, has also

been observed in the current study. To control the confounder effect of other host factors, we applied a

series of selection criteria to ensure the two groups of participants have balanced gender and BMI.

Although significant difference was identified in age of two groups, all participants in this study were

aged between 18 to 28. The differences induced by host origin have been attributed to genetic or external

factors, such as diet [51] and altitude [52]. Differences in cultural background and lifestyle habits between

Asian countries and Western countries have led to interest in understanding the differences in salivary

composition between different countries [53]. Our previous study collected salivary microbiome data from

47 publicly available datasets, reporting the differences in beta diversity of saliva samples from three

geographic locations, including North America, Europe, and China [16]. Notably, the variance explained

in the previous study could be confounded by external factors other than the differences among three

populations, such as climate and collection methods. To better understand the differences between two

participant groups, the current study effectively eliminated the variance caused by geographic location

by collecting all the saliva samples from the same location within a short period. *Veillonella* spp.,

including *Veillonella atypica* and *Veillonella dispar*, were identified as the differential taxa that were

higher in Australian samples, which was consistent with the result of our previous study. Among

them, *Veillonella atypica* has also been identified as a biomarker to differentiate people with obesity

from normal-weight individuals [54]，underlying a potential association among diet, body weight and

ethnicity.

Although the differences between the Chinese and Australia group have been observed in our study,

association between ethnicity and salivary protein composition is far from conclusive. A previous study

[29] compared protein concentration, flow rate, amylolytic activity, and lipolytic activity of unstimulated

saliva samples from 15 Chinese and 15 Caucasian participants living in the Netherlands. Like the

current study, they focused on the difference in saliva samples between Chinese and residents of a

country defined as "Western countries". However, they observed higher total salivary protein

concentrations in Chinese than in Caucasian participants. This discrepancy between two studies could

be attributed to the differences between the Western countries selected. Australian and Dutch residents

could have specific differences in diets and lifestyles, leading to the different biochemical parameters

of saliva. Meanwhile, no significant differences in flow rate and amylolytic activity was detected by

Mosca et al. [29] , which is consistent with our findings. Further proteomic analyses on samples collected

in the current study would facilitate the observation of the differences on specific protein concentration;

we suspect that differences in published studies might also be due to the collection of stimulated or

unstimulated saliva (or merging of these two classes).

Our *ex vivo* results show that Australian and Chinese have different aroma compounds released in the

oral cavity. Our previous study reported that people from Australia and China have different sensory

responses to the same wine [36] and reported increased levels of esters in the mixture of Chinese saliva

and wine. According to the results of current study, the prevalence of *Streptococcus* sp. HMT 066 has

been associated with the concentration of esters released. It has been reported that *Streptococcus* spp.

have the ability to release aroma from odourless precursors [8]. 2,3-butanediol is the second most

abundant by-product generated during the alcoholic fermentation of wine, contributing to its bitterness

[64]. The higher 2,3- butanediol concentration found in our study using wine and saliva from the Chinese

group positively correlates with the relative abundance of *Prevotella shahii* **(Figure 4F)**. *Prevotella* spp.

has been described as a potential contributor to aroma perception by enzymatic activity [13]. Nevertheless,

within genera like *Streptococcus* and *Prevotella*, the function of different species varies. Functional

annotation and taxonomic profiling with higher resolution at special or even stain level could be done

to differentiate the bacteria belongs to these two genera. Further experiments, such as strain isolation,

are required to verify the function of specific species. *Veillonella* spp. have been found different in

relative abundance between hosts from China and Western countries in both a previous study [16] and the

current study. Although no correlation was found between *Veillonella* spp. and any aroma compounds

in this study, more research on *Veillonella* could help understand the differences in aroma perception

between Chinese and Western people.

In summary, this study demonstrated that saliva secreted before, during and after stimulation differs in

composition and the ability of the three types of saliva to release aroma compounds from wine varies.

Although Phase A and C have more in common than Phase B, there are still some differences in their

composition and the role in aroma release. The enzymatic activity of esterase may explain the variation

in the ability of human saliva to release aroma compounds attributed to stimulation. Another important

finding of this study was the difference in salivary composition, especially salivary microbiota, between

Australian and Chinese participants. Several differential species were identified by full-length 16S

rRNA sequencing. We took samples from 30 people in this study, which is a relatively small number,

yet our findings mirror data collected in other studies. For example, our findings of a differential relative

abundance of *Veillonella* spp. contributes additional evidence to support this genus as a biomarker to

differentiate people originating in different geographic areas. The volatiles released from the wine-

saliva mixture of these two groups highlight the associations between salivary microbiota and aroma

release. Our findings serve as a basis for future studies targeting the relationship between salivary

composition, aroma perception, and wine preference. A better understanding of the variations in wine

preference gives insights into designing products for specific consumer groups and should be

complemented in the future by sensory tests to verify the findings in the human oral cavity.

     Figure Legends

     Figure 1. The differences of salivary composition amongst three phases of collection and two

     participant groups. Boxplots showing the flow rate of (**A**) Phase A, B, C and (**B**) Australian and

     Chinese participants; the total protein concentration of (**C**) Phase A, B, C and (**D**) Australian and

     Chinese participants; the total salivary esterase activity of (**E**) Phase A, B, C and (**F**) Australian and

     Chinese participants. Significant differences ($p< 0.05$, Student's t-test; $p< 0.05$, one-way ANOVA)

     were tested and are marked with "*"; with NS denoting no significant differences observed.

     Figure 2. The aroma released from wine with saliva samples from three collection phases are

     different. Point plots with lines showing the change of each volatile released from (**A**) wine-Australia

     saliva mixture and (**B**) wine-Chinese saliva mixture. The green lines represent the compounds showing

     highest concentration in Phase B and have no significant differences between Phase A and C (one-way

     ANOVA, $p < 0.05$). Among them, light green lines represent compounds only significantly higher than

     Phase A, while dark green lines represent compounds significantly higher than both Phase A and C.

     The pink lines represent the compounds that also demonstrate significant differences between Phase A

     and C. The dark pink lines showing the compounds have significant differences among all three

     comparisons. The light pink lines showing the compounds have differential concentration between

     Phase A and B, and Phase A and C. The volatiles with significantly changes are indicate by letters: a)

     Ethyl acetate; b) Diethyl succinate; c) Ethyl hexanoate; d) 3-Methylbutyl acetate; e) Ethyl butanoate; f)

     Ethyl 3-methylbutanoate; g) Ethyl propanoate; h) Ethyl 2-methylbutanoate; i) Hexyl acetate; j) 3-Ethyl-

     2-pentanol; k) Acetic acid; l) 1,3-Ditertiarybutylbenzene; m) Octanoic acid. (**C**) Bar plots showing the

     concentration of 3-Ethyl-2-pentanol and Acetic acid in six subgroups. Significant differences ($p< 0.05$,

     Student's t-test) between Australian and Chinese group in Phase A are marked with "*". (**D**) Principal

     component analysis (PCA) plot demonstrating the wine aroma release after spiking with samples from

     three groups.

Figure 3. The microbial profiles of Australian and Chinese participants' saliva samples are

different. Comparison of the salivary microbial composition between two groups in stacked bar plots

at (**A**) phylum level and (**B**) genus level. (**C**) Taxonomic hierarchies showing the enrichment of taxa

from two groups from phylum to species level. The taxa with significant differences (false discovery

rate-adjusted Wilcoxon rank-sum $q < 0.05$) were coloured with green (higher in Australian group) and

pink (higher in Chinese group).

Figure 4. The differences in salivary microbial composition between Australian and Chinese

participants associate with the differences in aroma released after mixing with wine. (**A**)

Comparison of the alpha-diversity between two groups, representing by Simpson's Diversity Index

(Wilcoxon rank-sum test; $p = 0.029$). (**B**) Principal coordinate analysis (PCoA) with Bray-Curtis

dissimilarity showing the difference between two groups (PERMANOVA; $p = 0.001$). (**C**) and (**D**)

Heatmap represents the mean relative abundance of (**C**) eight differential species identified by

ANCOM-BC2 and (**D**) three species that significantly correlated ($p < 0.01$, FDR-adjusted Spearman's

rank correlation) with aroma compounds in six pooled saliva groups. (**E**) Bar plots showing the log-

transformed concentration (μg/L 4-octanol equivalence) of aroma compounds detected in two groups.

The coloured squares on the bottom indicate the result of two-way ANOVA. (**F**) Spearman's correlation

heatmap between aroma compounds and relative abundance of species. Significant correlated pairs

($p < 0.01$, FDR-adjusted Spearman's rank correlation) are labelled with "*".

Funding

This study was funded by the Faculty of Science at the University of Melbourne. XR and YC are

supported by a Melbourne Research Scholarship administered by the University of Melbourne.

Acknowledgements

The authors would like to acknowledge all participants. The authors would also like to acknowledge

the assistance of Zijian Liang to the sample collection process. The work of Dr Pangzhen Zhang to set

up methodologies and verify volatile analysis is gratefully acknowledged.

Ethics approval and consent to participate

The study is under the approval of the University of Melbourne Human Research Ethics Committees

(Review Reference Number: 2022-24220-29889-3). All participants provided written informed consent.

Data availability

The sequencing data generated and analysed in the present study are available in NCBI BioProject

database, PRJNA1077481(https://www.ncbi.nlm.nih.gov/bioproject/1077481).

Declaration of Competing Interest

The authors declare that they have no competing financial interest.

Supporting Information

Supplementary Table 1 Background information of participants from Chinese and Australia

Supplementary Table 2 The influence of two factors, phase and ethnicity on the amylase activity and

aroma compounds detected by HS-SPME-GC-MS from the mixture of six types of pooled saliva

samples and wine, evaluated by two-way ANOVA and Tukey's pairwise comparison.

Supplementary Table 3 The concentration and odour activity values (OAV) of aroma compounds

detected by headspace solid-phase microextraction gas chromatography mass spectrometry (HS-SPME-

GC-MS) from the mixture of six types of pooled saliva samples and wine, quantified using external

standards.

Supplementary Figure 1 Flow chart showing the sample collection procedure of three types of saliva

from each participant: before, during and after stimulation.

Supplementary Figure 2 Thirteen species identified as differential among three phases by

ANCOM-BC. A) Point plots with lines showing the log-transformed relative abundance of differential

species in three phases. The green lines indicate the species with the lowest relative abundance in Phase

A, and the pink lines indicate the species with the lowest relative abundance in Phase B; **B)** Heatmap

showing the log-transformed mean relative abundance of differential species.

Supplementary Figure 3 The aroma released from wine after mixing with water, artificial saliva,

and pooled saliva samples. A) The table showing the concentration of volatiles released from wine-

water and wine-artificial saliva mixture. Superscripts indicate significant differences ($p < 0.05$;

Student's t test) between the two groups. **B)** The coloured squares indicate the comparison among wine-

water, wine-artificial saliva, and wine-saliva mixtures. The green squares represent the higher

concentration of released aroma after mixing with the pooled saliva than mixing with water or artificial

saliva. The pink squares represent the lower concentration of released aroma after mixing with the

pooled saliva than mixing with water or artificial saliva. The grey squares represent the volatiles that

have not been detected. The significant differences ($p < 0.05$; one-way ANOVA) were indicated by "*".

Supplementary Figure 4 Principal component analysis (PCA) plots of A) 27 semi-quantified
compounds and B) 17 compounds quantified using external standards showing the similar results with
the PCA plots of 21 semi-quantified compounds detected constantly in all subgroups.

References

(1) Genovese, A.; Piombino, P.; Gambuti, A.; Moio, L. Simulation of retronasal aroma of white and red wine in a model mouth system. Investigating the influence of saliva on volatile compound concentrations. *Food Chem.* **2009**, *114* (1), 100-107. DOI: 10.1016/j.foodchem.2008.09.022.

(2) Feron, G.; Ayed, C.; Qannari, E. M.; Courcoux, P.; Laboure, H.; Guichard, E. Understanding Aroma Release from Model Cheeses by a Statistical Multiblock Approach on Oral Processing. *PLoS One* **2014**, *9* (4), e93113. DOI: 10.1371/journal.pone.0093113.

(3) Ployon, S.; Morzel, M.; Canon, F. The role of saliva in aroma release and perception. *Food Chem.* **2017**, *226*, 212-220. DOI: 10.1016/j.foodchem.2017.01.055.

(4) Taylor, A. J.; Roozen, J. P. Volatile flavor release from foods during eating. *Crit. Rev. Food Sci. Nutr.* **1996**, *36* (8), 765-784. DOI: 10.1080/10408399609527749.

(5) Friel, E. N.; Taylor, A. J. Effect of Salivary Components on Volatile Partitioning from Solutions. *J. Agric. Food. Chem.* **2001**, *49* (8), 3898-3905. DOI: 10.1021/jf010371e.

(6) Buettner, A. Influence of Human Salivary Enzymes on Odorant Concentration Changes Occurring in Vivo. 1. Esters and Thiols. *J. Agric. Food. Chem.* **2002**, *50* (11), 3283-3289. DOI: 10.1021/jf011586r.

(7) Buettner, A. Influence of Human Saliva on Odorant Concentrations. 2. Aldehydes, Alcohols, 3-Alkyl-2-methoxypyrazines, Methoxyphenols, and 3-Hydroxy-4,5-dimethyl-2(5H)-furanone. *J. Agric. Food. Chem.* **2002**, *50* (24), 7105-7110. DOI: 10.1021/jf020714o.

(8) Muñoz-González, C.; Cueva, C.; Ángeles Pozo-Bayón, M.; Victoria Moreno-Arribas, M. Ability of human oral microbiota to produce wine odorant aglycones from odourless grape glycosidic aroma precursors. *Food Chem.* **2015**, *187*, 112-119. DOI: 10.1016/j.foodchem.2015.04.068.

(9) Guichard, E. Flavour retention and release from protein solutions. *Biotechnol. Adv.* **2006**, *24* (2), 226-229. DOI: 10.1016/j.biotechadv.2005.11.003.

(10) Pagès-Hélary, S.; Andriot, I.; Guichard, E.; Canon, F. Retention effect of human saliva on aroma release and respective contribution of salivary mucin and α-amylase. *Food Res. Int.* **2014**, *64*, 424-431. DOI: 10.1016/j.foodres.2014.07.013.

(11) Pérez-Jiménez, M.; Rocha-Alcubilla, N.; Pozo-Bayón, M. Á. Effect of saliva esterase activity on ester solutions and possible consequences for the in-mouth ester release during wine intake. *J. Texture Stud.* **2019**, *50* (1), 62-70. DOI: 10.1111/jtxs.12371.

(12) María, P.-J.; Carolina, M.-G.; María Ángeles, P.-B. Understanding human salivary esterase activity and its variation under wine consumption conditions. *RSC Advances* **2020**, *10* (41), 24352-24361, 10.1039/D0RA04624H. DOI: 10.1039/D0RA04624H.

(13) Parker, M.; Onetto, C.; Hixson, J.; Bilogrevic, E.; Schueth, L.; Pisaniello, L.; Borneman, A.; Herderich, M.; de Barros Lopes, M.; Francis, L. Factors Contributing to Interindividual Variation in Retronasal Odor Perception from Aroma Glycosides: The Role of Odorant Sensory Detection Threshold, Oral Microbiota, and Hydrolysis in Saliva. *J. Agric. Food. Chem.* **2020**, *68* (38), 10299-10309. DOI: 10.1021/acs.jafc.9b05450.

(14) Piombino, P.; Genovese, A.; Esposito, S.; Moio, L.; Cutolo, P. P.; Chambery, A.; Severino, V.; Moneta, E.; Smith, D. P.; Owens, S. M.; et al. Saliva from Obese Individuals Suppresses the Release of Aroma Compounds from Wine. *PLoS One* **2014**, *9* (1), e85611. DOI: 10.1371/journal.pone.0085611.

(15) Wade, W. G. The oral microbiome in health and disease. *Pharmacol. Res.* **2013**, *69* (1), 137-143. DOI: 10.1016/j.phrs.2012.11.006.

(16) Ruan, X.; Luo, J.; Zhang, P.; Howell, K. The salivary microbiome shows a high prevalence of core bacterial members yet variability across human populations. *NPJ Biofilms Microbi.* **2022**, *8* (1), 85. DOI: 10.1038/s41522-022-00343-7.

(17) Emmelin, N. Control of Salivary Glands. In *Oral Physiology*, Emmelin, N., Zotterman, Y. Eds.; Pergamon, 1972; pp 1-16.

(18) Humphrey, S. P.; Williamson, R. T. A review of saliva: Normal composition, flow, and function. *The Journal of Prosthetic Dentistry* **2001**, *85* (2), 162-169. DOI: https://doi.org/10.1067/mpr.2001.113778.

(19) Jasim, H.; Olausson, P.; Hedenberg-Magnusson, B.; Ernberg, M.; Ghafouri, B. The proteomic profile of whole and glandular saliva in healthy pain-free subjects. *Sci. Rep.* **2016**, *6* (1), 39073. DOI: 10.1038/srep39073.

(20) Proctor, G. B.; Carpenter, G. H. Regulation of salivary gland function by autonomic nerves. *Auton.*
Neurosci. **2007**, *133* (1), 3-18. DOI: 10.1016/j.autneu.2006.10.006.
(21) Shannon, I. L.; Suddick, R. P.; Chauncey, H. H. Effect of atropine-induced flow rate depression
on the composition of unstimulated human parotid fluid. *Arch. Oral Biol.* **1969**, *14* (7), 761-770. DOI:
10.1016/0003-9969(69)90167-8.
(22) Gomar-Vercher, S.; Simón-Soro, A.; Montiel-Company, J. M.; Almerich-Silla, J. M.; Mira, A.
Stimulated and unstimulated saliva samples have significantly different bacterial profiles. *PLoS One*
2018, *13* (6), e0198021. DOI: 10.1371/journal.pone.0198021.
(23) Belstrøm, D.; Holmstrup, P.; Bardow, A.; Kokaras, A.; Fiehn, N. E.; Paster, B. J. Comparative
analysis of bacterial profiles in unstimulated and stimulated saliva samples. *J. Oral Microbiol.* **2016**, *8*,
30112. DOI: 10.3402/jom.v8.30112.
(24) Jo, R.; Nishimoto, Y.; Umezawa, K.; Yama, K.; Aita, Y.; Ichiba, Y.; Murakami, S.; Kakizawa, Y.;
Kumagai, T.; Yamada, T.; et al. Comparison of oral microbiome profiles in stimulated and unstimulated
saliva, tongue, and mouth-rinsed water. *Sci. Rep.* **2019**, *9* (1), 16124. DOI: 10.1038/s41598-019-52445-
6.
(25) Neyraud, E.; Palicki, O.; Schwartz, C.; Nicklaus, S.; Feron, G. Variability of human saliva
composition: Possible relationships with fat perception and liking. *Arch. Oral Biol.* **2012**, *57* (5), 556-
566. DOI: 10.1016/j.archoralbio.2011.09.016.
(26) Pérez-Jiménez, M.; Muñoz-González, C.; Pozo-Bayón, M. Á. Specificity of Saliva Esterases by
Wine Carboxylic Esters and Inhibition by Wine Phenolic Compounds Under Simulated Oral Conditions.
Front. Nutr. **2021**, *8*, Original Research. DOI: 10.3389/fnut.2021.761830.
(27) Carvalho, F. M.; Moksunova, V.; Spence, C. Cup texture influences taste and tactile judgments in
the evaluation of specialty coffee. *Food Qual.* **2020**, *81*, 103841. DOI:
https://doi.org/10.1016/j.foodqual.2019.103841.
(28) Esteban-Fernández, A.; Rocha-Alcubilla, N.; Muñoz-González, C.; Moreno-Arribas, M. V.; Pozo-
Bayón, M. Á. Intra-oral adsorption and release of aroma compounds following in-mouth wine exposure.
Food Chem. **2016**, *205*, 280-288. DOI: 10.1016/j.foodchem.2016.03.030.
(29) Mosca, A. C.; Stieger, M.; Neyraud, E.; Brignot, H.; van de Wiel, A.; Chen, J. How are
macronutrient intake, BMI, ethnicity, age, and gender related to the composition of unstimulated saliva?
A case study. *J. Texture Stud.* **2019**, *50* (1), 53-61. DOI: 10.1111/jtxs.12362.
(30) Bhuptani, D.; Kumar, S.; Vats, M.; Sagav, R. Age and gender related changes of salivary total
protein levels for forensic application. *J. Forensic Odontostomatol.* **2018**, *36* (1), 26-33.
(31) Burcham, Z. M.; Garneau, N. L.; Comstock, S. S.; Tucker, R. M.; Knight, R.; Metcalf, J. L.;
Miranda, A.; Reinhart, B.; Meyers, D.; Woltkamp, D.; et al. Patterns of Oral Microbiota Diversity in
Adults and Children: A Crowdsourced Population Study. *Sci. Rep.* **2020**, *10* (1), 2133. DOI:
10.1038/s41598-020-59016-0.
(32) Morita, E.; Narikiyo, M.; Yokoyama, A.; Yano, A.; Kamoi, K.; Yoshikawa, E.; Yamaguchi, T.;
Igaki, H.; Tachimori, Y.; Kato, H.; et al. Predominant presence of Streptococcus anginosus in the saliva
of alcoholics. *Oral Microbiol. Immunol.* **2005**, *20* (6), 362-365. DOI: 10.1111/j.1399-
302X.2005.00242.x.
(33) Mason, M. R.; Nagaraja, H. N.; Camerlengo, T.; Joshi, V.; Kumar, P. S. Deep Sequencing
Identifies Ethnicity-Specific Bacterial Signatures in the Oral Microbiome. *PLoS One* **2013**, *8* (10),
e77287. DOI: 10.1371/journal.pone.0077287.
(34) Yang, Y.; Zheng, W.; Cai, Q.; Shrubsole, M. J.; Pei, Z.; Brucker, R.; Steinwandel, M.; Bordenstein,
S. R.; Li, Z.; Blot, W. J. Racial differences in the oral microbiome: data from low-income populations
of African ancestry and European ancestry. *mSystems* **2019**, *4* (6). DOI: 10.1128/msystems. 00639-19.
(35) Premaraj, T. S.; Vella, R.; Chung, J.; Lin, Q.; Hunter, P.; Underwood, K.; Premaraj, S.; Zhou, Y.
Ethnic variation of oral microbiota in children. *Sci. Rep.* **2020**, *10* (1), 14788. DOI: 10.1038/s41598-
020-71422-y.
(36) Luo, J.; Ruan, X.; Ang, C.-S.; Nolvachai, Y.; Marriott, P. J.; Zhang, P.; Howell, K. Variation of
wine preference amongst consumers is influenced by the composition of salivary proteins. *NPJ Sci.*
Food **2023**, *7* (1), 51. DOI: 10.1038/s41538-023-00222-1.
(37) Williamson, P. O.; Robichaud, J.; Francis, I. L. Comparison of Chinese and Australian consumers'
liking responses for red wines. *Aust. J. Grape Wine Res.* **2012**, *18* (3), 256-267. DOI: 10.1111/j.1755-
0238.2012.00201.x.

(38) Kim, S.; Petard, N.; Hong, J. What is lost in translation: A cross-cultural study to compare the concept of nuttiness and its perception in soymilk among Korean, Chinese, and Western groups. *Food Res. Int.* **2018**, *105*, 970-981. DOI: 10.1016/j.foodres.2017.12.002.

(39) Sáenz-Navajas, M.-P.; Avizcuri, J. M.; Echávarri, J. F.; Ferreira, V.; Fernández-Zurbano, P.; Valentin, D. Understanding quality judgements of red wines by experts: Effect of evaluation condition. *Food Qual.* **2016**, *48*, 216-227. DOI: 10.1016/j.foodqual.2015.10.001.

(40) Benjamin, O.; Silcock, P.; Beauchamp, J.; Buettner, A.; Everett, D. W. Tongue Pressure and Oral Conditions Affect Volatile Release from Liquid Systems in a Model Mouth. *J. Agric. Food. Chem.* **2012**, *60* (39), 9918-9927. DOI: 10.1021/jf3028232.

(41) Muñoz-González, C.; Feron, G.; Guichard, E.; Rodríguez-Bencomo, J. J.; Martín-Álvarez, P. J.; Moreno-Arribas, M. V.; Pozo-Bayón, M. Á. Understanding the Role of Saliva in Aroma Release from Wine by Using Static and Dynamic Headspace Conditions. *J. Agric. Food. Chem.* **2014**, *62* (33), 8274-8288. DOI: 10.1021/jf503503b.

(42) Harrell, F. E. J. *Hmisc: Harrell Miscellaneous*. 2024. https://CRAN.R-project.org/package=Hmisc (accessed 2024-06-05).

(43) Bolyen, E.; Rideout, J. R.; Dillon, M. R.; Bokulich, N. A.; Abnet, C. C.; Al-Ghalith, G. A.; Alexander, H.; Alm, E. J.; Arumugam, M.; Asnicar, F.; et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat. Biotechnol.* **2019**, *37* (8), 852-857. DOI: 10.1038/s41587-019-0209-9.

(44) Callahan, B. J.; McMurdie, P. J.; Rosen, M. J.; Han, A. W.; Johnson, A. J. A.; Holmes, S. P. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **2016**, *13* (7), 581-583. DOI: 10.1038/nmeth.3869.

(45) Escapa, I. F.; Chen, T.; Huang, Y.; Gajare, P.; Dewhirst, F. E.; Lemon, K. P. New Insights into Human Nostril Microbiome from the Expanded Human Oral Microbiome Database (eHOMD): a Resource for the Microbiome of the Human Aerodigestive Tract. *mSystems* **2018**, *3* (6), 10.1128/msystems.00187-00118. DOI: 10.1128/msystems.00187-18.

(46) Quast, C.; Pruesse, E.; Yilmaz, P.; Gerken, J.; Schweer, T.; Yarza, P.; Peplies, J.; Glöckner, F. O. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **2013**, *41*, D590-596. DOI: 10.1093/nar/gks1219.

(47) Oksanen, J.; Blanchet, F. G.; Friendly, M.; Kindt, R.; Legendre, P.; McGlinn, D.; Minchin, P. R.; O'Hara, R. B.; Simpson, G. L.; Solymos, P.; et al. *vegan: Community Ecology Package. R package version 2.5-7.* 2020. https://CRAN.R-project.org/package=vegan (accessed 2024-06-05).

(48) Foster, Z. S. L.; Sharpton, T. J.; Grünwald, N. J. Metacoder: An R package for visualization and manipulation of community taxonomic diversity data. *PLoS Comp. Biol.* **2017**, *13* (2), e1005404. DOI: 10.1371/journal.pcbi.1005404.

(49) Lin, H.; Peddada, S. D. Multigroup analysis of compositions of microbiomes with covariate adjustments and repeated measures. *Nat. Methods* **2024**, *21* (1), 83-91. DOI: 10.1038/s41592-023-02092-7.

(50) Lee, V.; Linden, R. An olfactory-submandibular salivary reflex in humans. *Exp. Physiol.* **1992**, *77* (1), 221-224. DOI: 10.1113/expphysiol.1992.sp003578.

(51) Liu, K.; Chen, S.; Huang, J.; Ren, F.; Yang, T.; Long, D.; Li, H.; Huang, X. Oral Microbiota of Children Is Conserved across Han, Tibetan and Hui Groups and Is Correlated with Diet and Gut Microbiota. *Microorganisms* **2021**, *9* (5), 1030. DOI: 10.3390/microorganisms9051030.

(52) Hao, Y.; Tang, C.; Du, Q.; Zhou, X.; Peng, X.; Cheng, L. Comparative analysis of oral microbiome from Zang and Han populations living at different altitudes. *Arch. Oral Biol.* **2021**, *121*, 104986. DOI: 10.1016/j.archoralbio.2020.104986.

(53) Kim, T. S.; Kang, N. W.; Lee, S.-B.; Eickholz, P.; Pretzl, B.; Kim, C. Differences in subgingival microflora of Korean and German periodontal patients. *Arch. Oral Biol.* **2009**, *54* (3), 223-229. DOI: 10.1016/j.archoralbio.2008.10.005.

(54) López-Dávalos, P. C.; Requena, T.; Pozo-Bayón, M. Á.; Muñoz-González, C. Decreased retronasal olfaction and taste perception in obesity are related to saliva biochemical and microbiota composition. *Food Res. Int.* **2023**, *167*, 112660. DOI: 10.1016/j.foodres.2023.112660.

(55) Ferreira, V.; López, R.; Escudero, A.; Cacho, J. F. The aroma of Grenache red wine: hierarchy and nature of its main odorants. *J. Sci. Food Agric.* **1998**, *77* (2), 259-267. DOI: 10.1002/(SICI)1097-0010(199806)77:2<259::AID-JSFA36>3.0.CO;2-Q.

(56) Goodstein, E. S.; Bohlscheid, J. C.; Evans, M.; Ross, C. F. Perception of flavor finish in model
white wine: A time-intensity study. *Food Qual.* **2014**, *36*, 50-60. DOI: 10.1016/j.foodqual.2014.02.012.
Muñoz-González, C.; Pérez-Jiménez, M.; Pozo-Bayón, M. Á. Oral persistence of esters is affected by
wine matrix composition. *Food Res. Int.* **2020**, *135*, 109286. DOI: 10.1016/j.foodres.2020.109286.
(57) Guo, C.; Deng, H.; Li, E. Removal of acetic acid in mulberry wine by co-inoculating
Saccharomyces cerevisiae with indigenous non-Saccharomyces yeast. *Food Bioscience* **2024**, *58*,
103658. DOI: https://doi.org/10.1016/j.fbio.2024.103658.
(58) Buettner, A. Investigation of Potent Odorants and Afterodor Development in Two Chardonnay
Wines Using the Buccal Odor Screening System (BOSS). *J. Agric. Food. Chem.* **2004**, *52* (8), 2339-
2346. DOI: 10.1021/jf035322b.
(59) Mayr, C. M.; Parker, M.; Baldock, G. A.; Black, C. A.; Pardon, K. H.; Williamson, P. O.; Herderich,
M. J.; Francis, I. L. Determination of the Importance of In-Mouth Release of Volatile Phenol
Glycoconjugates to the Flavor of Smoke-Tainted Wines. *J. Agric. Food. Chem.* **2014**, *62* (11), 2327-
2336. DOI: 10.1021/jf405327s.
(60) Ployon, S.; Brulé, M.; Andriot, I.; Morzel, M.; Canon, F. Understanding retention and
metabolization of aroma compounds using an in vitro model of oral mucosa. *Food Chem.* **2020**, *318*,
126468. DOI: 10.1016/j.foodchem.2020.126468.
(61) Esteban-Fernández, A.; Muñoz-González, C.; Jiménez-Girón, A.; Pérez-Jiménez, M.; Pozo-Bayón,
M. Á. Aroma release in the oral cavity after wine intake is influenced by wine matrix composition.
Food Chem. **2018**, *243*, 125-133. DOI: https://doi.org/10.1016/j.foodchem.2017.09.101.
(62) Perez-Jiménez, M.; Esteban-Fernández, A.; Muñoz-González, C.; Pozo-Bayón, M. A. Interactions
among Odorants, Phenolic Compounds, and Oral Components and Their Effects on Wine Aroma
Volatility. *Molecules* **2020**, *25* (7), 1701.
(63) Waszkiewicz, N.; Szajda, S. D.; Jankowska, A.; Zwierz, P.; Czernikiewicz, A.; Szulc, A.; Zwierz,
K. The Effect of Acute Ethanol Intoxication on Salivary Proteins of Innate and Adaptive Immunity.
Alcoholism: Clinical and Experimental Research **2008**, *32* (4), 652-656. DOI:
https://doi.org/10.1111/j.1530-0277.2007.00613.x.
(64) Romano, P. Metabolic characteristics of wine strains during spontaneous and inoculated
fermentation. In *Food Technology and Biotechnology*, 1997; Vol. 35, pp 255-260.

Table 1 Aroma compounds detected by headspace solid-phase microextraction gas chromatography mass spectrometry (HS-SPME-GC-MS) from the mixture of six types of pooled saliva samples and wine.

No.	Aroma Compound	RI-NIST*	RI-Actual	Internal standard (4-octanol) equivalence (µg/L)					
				Phase A		Phase B		Phase C	
				Australian	Chinese	Australian	Chinese	Australian	Chinese
	Esters								
1	Ethyl acetate	888	887	4646.99 ± 691.59^c	7449.38 ± 35.97^a	8822.61 ± 158.22^a	8606 ± 759.79^a	5935.32 ± 842.17^{bc}	7368.83 ± 103.25^{ab}
2	Ethyl propanoate	960	958	60.23 ± 4.97^{bc}	74.47 ± 2.46^{ab}	89.82 ± 2.16^a	87.58 ± 10.23^a	56.58 ± 7.30^c	71.78 ± 2.21^{bc}
3	Ethyl isobutyrate	962	959	153.31 ± 37.03^a	171.05 ± 54.71^a	120.65 ± 38.81^a	128.48 ± 38.89^a	123.29 ± 34.79^a	70.75 ± 61.53^a
4	Ethyl butanoate	1039	1035	49.91 ± 43.35^d	121.16 ± 3.14^b	151.59 ± 5.06^a	149.92 ± 17.20^a	93.75 ± 11.98^{cd}	120.11 ± 4.30^{bc}
5	Ethyl 2-methylbutanoate	1055	1050	41.49 ± 6.72^c	79.85 ± 2.41^b	102.68 ± 4.39^a	102.34 ± 11.04^a	56.79 ± 8.78^c	76.28 ± 2.18^b
6	Ethyl 3-methylbutanoate	1072	1067	48.24 ± 9.48^d	107.18 ± 5.42^b	137.85 ± 2.7^a	134.99 ± 11.07^a	72.17 ± 13.35^c	101.04 ± 4.98^b
7	3-Methylbutyl acetate	1126	1122	545.84 ± 89.42^d	1084.34 ± 9.03^{bc}	1302.7 ± 21.06^{ab}	1347.1 ± 143.51^a	765.84 ± 108.53^d	1066.07 ± 59.63^c
8	Methyl hexanoate	1190	1184	n.d.	2.88 ± 0.57^{bc}	3.58 ± 0.37^{ab}	4.06 ± 0.39^a	0.96 ± 0.32^d	2.36 ± 0.25^c
9	Ethyl hexanoate	1235	1233	851.85 ± 223.93^d	1977.03 ± 53.02^{bc}	2391.76 ± 66.67^{ab}	2478.48 ± 252.05^a	1259.94 ± 253.43^d	1830.77 ± 80.92^c
10	Hexyl acetate	1235	1232	7.02 ± 6.64^c	15.17 ± 3.00^{abc}	20.62 ± 0.86^a	21.65 ± 1.20^a	10.06 ± 2.16^{bc}	16.05 ± 1.80^{ab}
11	Ethyl heptanoate	1334	1332	0.09 ± 0.01^b	4.29 ± 3.68^{ab}	2.7 ± 4.18^{ab}	8.01 ± 2.27^a	0.2 ± 0.11^b	0.21 ± 0.16^b
12	Diethyl succinate	1681	1681	2752.88 ± 136.55^a	2256.51 ± 215.43^b	2471.91 ± 151.36^{ab}	2314.02 ± 68.01^{ab}	2557.77 ± 315.12^{ab}	2695.89 ± 66.46^{ab}
	Alcohols								
13	2-Methyl-1-propanol	1098	1096	493.94 ± 24.95^a	468.86 ± 20.24^a	521.16 ± 20.25^a	492.19 ± 50.54^a	491.19 ± 35.89^a	515.29 ± 17.78^a

#	Compound	RI	RI-NIST					
14	1-Butanol	1144	1141	12.36 ± 0.42^a	7.74 ± 6.67^a	8.97 ± 7.73^a	12.93 ± 1.23^a	12.3 ± 0.37^a	12.59 ± 0.86^a
15	3-Methyl-1-butanol	1204	1205	10562.24 ± 491.79^a	9444.18 ± 259.00^a	10666.84 ± 311.22^a	10051.84 ± 1014.17^a	10159.14 ± 712.50^a	10557.02 ± 315.42^a
16	1-Hexanol	1348	1346	319.34 ± 15.53^a	305.42 ± 12.75^a	346.57 ± 13.21^a	331.82 ± 31.04^a	315.04 ± 19.65^a	335.08 ± 13.11^a
17	3-Ethyl-2-pentanol	N/A	1531	19.18 ± 1.58^b	32.62 ± 2.77^a	40.21 ± 6.14^a	39.85 ± 5.86^a	29.76 ± 4.30^{ab}	36.01 ± 4.12^a
18	2,3-Butanediol	1542	1540	43.59 ± 1.18^a	43.61 ± 13.28^a	37.71 ± 5.61^a	47.51 ± 6.63^a	41.67 ± 6.48^a	51.4 ± 14.36^a
	Acids								
19	Acetic acid	1461	1454	94.23 ± 21.35^c	165.63 ± 34.94^b	221.29 ± 12.79^{ab}	205.82 ± 25.96^{ab}	228.81 ± 13.09^{ab}	230.26 ± 23.85^a
20	Pentanoic acid	1744	1732	8.81 ± 0.10^a	7.26 ± 0.85^a	8.37 ± 0.78^a	8.39 ± 0.49^a	7.94 ± 0.63^a	8.4 ± 1.55^a
21	Octanoic acid	2087	2079	3.91 ± 0.82^a	5.9 ± 2.16^a	7.74 ± 0.66^a	8.52 ± 2.42^a	4.63 ± 1.53^a	7.33 ± 3.15^a
	Benzenoid								
22	Styrene	1256	1254	n.d.	2.56 ± 0.23^b	4.47 ± 0.42^a	4.79 ± 0.77^a	n.d.	1.19 ± 1.03^b
23	1,3-Ditertiarybutylbenzene	N/A	1424	17.57 ± 1.55^{bc}	32.37 ± 3.73^b	56.56 ± 4.08^a	69.43 ± 4.61^a	14.76 ± 9.41^c	29.05 ± 7.84^{bc}
	Terpenes								
24	D-Limonene	1199	1196	4.5 ± 3.55^a	n.d.	n.d.	0.92 ± 0.70^a	n.d.	n.d.
25	p-&m-Cymene	1269	1272	n.d.	n.d.	5.16 ± 0.65^a	6.96 ± 1.21^a	n.d.	n.d.
26	(-)-Clovene	N/A	1304	190.44 ± 38.62^a	40.73 ± 9.73^b	45.69 ± 19.52^b	65.47 ± 12.48^b	16.35 ± 0.60^b	27.51 ± 3.92^b
27	1,1,6-Trimethyl-1,2-dihydronaphthalene (TDN)	1712	1704	22.88 ± 5.95^a	15.8 ± 1.76^{ab}	15.54 ± 1.19^{ab}	22.88 ± 5.95^a	n.d.	9.23 ± 1.57^b

Superscripts letters for the same compound indicate significant differences ($p < 0.05$; Tukey test) on the aroma release from wine after spiking with different saliva samples.

RI = Retention index.

*RI-NIST was obtained from NIST Webbook (2018).