Nevertheless sex persisted: facultative sex persists despite high costs in a sexually polymorphic aphid parasitoid

Rebecca A Boulton[*1,2]

[1] Biological and Environmental Sciences, University of Stirling, FK9 4NF, United Kingdom
[2] Laboratory of Genetics, Plant Sciences Group, Wageningen University and Research, 6708 8 PB, Netherlands

* Corresponding author: Rebecca.boulton@stir.ac.uk

Abstract

The prevalence of sexual reproduction has long puzzled evolutionary biologists. This is because of the 'twofold cost of sex', or, more accurately, 'the twofold cost of males'; an individual that reproduces sexually is expected to invest half its resources in producing male offspring, but an equivalent asexual individual will produce all daughters and so should have twice as many grandoffspring. To understand how sex coexists with asexual reproduction despite the twofold cost I assayed behaviour and offspring production of sexual and asexual females of the sexually polymorphic aphid parasitoid *Lysiphlebus fabarum.* To fully measure the costs of sex (including the costs of mating and exposure to males) I exposed half of these females to sexual males. I found that when males were available, asexual females are often facultatively sexual; they readily mate with males and producee offspring with paternal genetic contributions. Sexual females did not appear to be limited in their fitness by the 'twofold cost of males'; although they produced sons, they also produced the same number of daughters as asexual females because their overall offspring production was higher. Instead, the main cost of sex was increased reproductive failure, particularly when sex was facultative. Stable coexistence of both reproductive modes in *L. fabarum* may occur because this main cost of sex is paid by both asexual (under facultative sex) and sexual females. Coexistence and interbreeding with male-producing sexuals may be the very thing that has allowed asexual *L. fabarum* lineages to persist in the long-term; ensuring sufficient genetic variation for adaptation through occasional facultative sex, as well as reducing the costs, such as low fecundity, that accumulate under obligate asexual reproduction.

1. Introduction

Understanding how sex became the dominant reproductive strategy in eukaryotes has long been a major theme in evolutionary biology [1–3]. The initial interest in this question arose from the observation that asexually reproducing individuals are expected to have twice the number of grand offspring (and four times the number of great-grand offspring etc) as a sexual individual. This became known as the 'twofold cost of sex' [1] and arises because sexual individuals must 'waste' resources on producing males, which do not efficiently convert resources to offspring [4]. When the sex ratio is 50:50, sexuals produce half as many daughters as asexuals, so the twofold cost of sex is sometimes known as the 'twofold cost of males' [1–4]. How then, did sex become more prevalent than asexual reproduction in eukaryotes after it evolved over two billion years ago? And more recently, given that asexual reproduction readily re-evolves in eukaryotes [2] why does sex still dominate?

Historically, two broad models which encompass different types of benefits have been proposed to explain why sex evolved and persisted: mutational models and ecological models. Mutational models emphasize the importance of sex in exposing mutations to selection, so that deleterious mutations can be purged, and beneficial mutations can be fixed more effectively [3,5–8]. Ecological models on the other hand emphasize the benefits of sex for adaptive potential; by creating new gene combinations sex facilitates adaptation under environmental change [9]. Mutational and ecological benefits of sex are not mutually exclusive and may interact in complex ways, sometimes synergistically.

The complex and context-dependent ways in which the different benefits of sex interact likely explains the complex, labile and dynamic ways that reproductive mode evolves in the eukaryotes [2–4,10,11]. For instance, there is often substantial plasticity in reproductive mode, as many eukaryotes reproduce asexually only some of the time, either as an obligate part of their lifecycle (as in cyclical parthenogens like many aphids and *Daphnia*; [12]Allen & Lynch, 2008)facultatively when a suitable mate can't be found [11], or sometimes spontaneously (so-called *Tychoparthenogenesis* [2,13]. Asexual reproduction has also proved not to be the evolutionary 'dead end' predicted by theory. 'Ancient' asexual lineages exist despite the widespread view that sex is required for long-term persistence, speciation and adaptation to changing environments due to the genetic variation it generates [3,14,15]. It even seems as though sex may be a case where Dollo's law, which states that complex ancestral states once lost cannot be reacquired, is broken; Domes et al [16] found that sexual reproduction has reevolved in a group of parthenogenetic mites.

Whilst the results of these broad scale phylogenetic and population-level studies have been enlightening, there is much more that we could learn from them about the evolutionary origin and maintenance of sex. Measuring the performance of sexual and asexual individuals (and populations) in a range of ecologically relevant contexts can help us to understand these broad scale patterns, and more accurately develop theoretical predictions [4]. While such studies can be empirically challenging, they have revealed that the 'twofold cost of males' often deviates far from the theoretical twofold prediction. Finding a mate, avoiding sexually transmitted diseases, evading persistent and undesirable mates, and, finding sufficient resources for reproduction when males deplete them are just some of the additional costs that can further reduce the relative fitness of sexuals compared to asexuals [4]. Importantly, some of these costs of males also affect asexual females when they overlap in time and space with their sexual relatives [4,] potentially reducing the fitness of asexuals and supressing the spread of asexual reproduction.

The sexual-asexual aphid parasitoid *Lysiphlebus fabarum* offers an ideal system to explore how the costs of sex influence the success and proliferation of asexual reproduction. We already have a comprehensive understanding of the molecular basis and population genetics of reproductive mode in *L. fabarum* [17–19]. Asexual reproduction seems to have initially arisen 0.5 million years ago in *L. fabarum,* and it is determined by the inheritance of a single recessive allele which results in parthenogenesis by central-fusion automixis (the haploid products of meiosis I fuse to form diploid gametes which develop without fertilisation; [19,20]). Whilst this form of parthenogenesis does maintain heterozygosity to some extent, because recombination still occurs, asexual lineages can only become more homozygous, eventually resulting in fitness costs[2]. However, parthenogenetic populations of *L. fabarum*

exhibit high levels of heterozygosity and high allelic variation [17,18]. This genetic diversity is
thought to underlie the success of asexual reproduction in this species, resulting in
parthenogenetic populations dominating across Europe [17,18]. Sexual populations are less
commonly encountered in the field, but sex does persist in *L. fabarum*, and even occurs in
sympatry with asexuality in some regions [17,18].

This study aims to elucidate the mechanisms underlying the population-level and
phylogenetic patterns of reproductive mode observed in *L. fabarum* by measuring the
individual fitness of sexual and asexual females. I tested whether sexual females experience
reduced fitness compared to asexuals due to the need to produce male offspring. I also
tested whether exposure to males further reduces female fitness (for instance due to costs of
mating or resisting mating; [21]) and whether these costs are equivalent for sexual and
asexual *L. fabarum* females. To do this I gave female *L. fabarum* from 7 asexual lines and
one outbred sexual population aphid hosts to parasitise after (i) being exposed to males
(from the sexual population) or (ii) being kept in isolation (due to haplodiploidy sexual
females can produce unfertilised eggs that develop as male offspring). I found that asexual
females readily copulated when they were exposed to males and so in a follow-up
experiment, I genotyped their G1 offspring to test whether mating reflects a behavioural
vestige of sexual reproduction without fertilisation or whether these females can still
reproduce sexually. I also assayed the reproductive success of daughters produced by
asexual females which either mated or remained virgin. This allowed me to test whether
facultative sex allows females to reap the short-term demographic benefits of asexual
reproduction (not producing males) and also gain the long-term genetic benefits of sex
(which only appear in hybrid offspring).

2. Methods

(a) Study system

In the aphid parasitoid *Lysiphlebus fabarum* both sexual and asexual populations exist,
separately as well as in sympatry in some regions [17,18]. Asexual reproduction is inherited
via a single recessive allele which appeared approximately 0.5 million years ago [17].
Inheritance of two copies of the asexual allele results in parthenogenetic asexual
reproduction through central-fusion automixis, where the products of meiosis 1 fuse and
recombine, restoring diploidy without integrating paternal DNA [19]. A panel of microsatellite
markers has been developed to study the population genetics of *L. fabarum,* one of which,
Lysi07, shows perfect linkage to the recessive genetic factor associated with asexual
reproduction [22]

(b) Lab rearing

This study used females from 7 asexual lines and males from an outbred sexual population
of the aphid parasitoid *Lysiphlebus fabarum*. These lines were provided by Prof Christoph
Vorburger (EAWAG, Zürich, Switzerland) who has been maintaining them since they were
collected between 2006 and 2017 (see [19] for details). The wasps have been cultured at the
University of Stirling under equivalent conditions since March 2023. The wasps are given
black bean aphid (*Aphis fabae*) colonies to parasitise; they are able to parasitise all instars of
aphid but prefer 1[st] instar nymphs when given a choice [23]. Female wasps lay a single egg

inside each aphid (they are solitary endoparasitoids) and the host continues to grow as the
wasp larva develops (they are koinobionts; [23]). In the lab aphids are maintained on broad
bean seedlings (*Vicia faba*). All insects are kept in climate-controlled chambers at 20°C, 70%
RH on a 16:8 hour light:dark cycle. Under these conditions the generation time of *L. fabarum*
is 12-14 days. Several microsatellite markers are available for genotyping these lines (see
[22]), which are regularly genotyped to check their genetic identity has been maintained.
150
151 (c) Experimental design
152
To measure the benefits and costs of sex in *L. fabarum* I assayed offspring production by
sexual and asexual *L. fabarum* females (referred to as G0) which were exposed to a male
(from the obligately sexual population), M, or remained virgin, V (fig 1a). All females,
including asexuals, in the M treatment readily copulated with the male, so I refer to these
females as 'mated'. To test for longer-term costs of mating (and potentially facultative sex) I
repeated the same fitness assays with the unmated daughters of V and M asexual females
(fig 1c; V_2 and M_2; referred to as G1 females). The fitness measures I used were: (i) the
number of aphid mummies (parasitised dead aphids) produced and (ii) the number of
mummies from which adult wasps successfully emerged (fig 1b and 1d). Emerging wasps
(1b and 1d) were also sexed in order to test whether asexual females avoid the 'twofold' cost
of male production regardless of their mated status. To test whether mating by asexual
females does equate to facultative sex I genotyped full broods of emerged wasps (fig 1b and
1d) to test for the presence of sexual alleles the offspring of mated asexual wasps.
166
To assess the 'twofold' cost of sex (male production) and determine whether it applies to
facultative and obligately sexual females I looked at the number of sons and daughters
produced by V, M and V_2 and M_2 females. To estimate this cost of sex in the case of obligate
sex I compared son and daughter production between G0 obligately sexual mated M
females and all (V and M) G0 asexual females (fig 1, grey boxes). To test whether the cost of
male production also applies to facultatively sexual females (asexual M females) I compared
son and daughter production of (i) G0 asexual V and M females (fig 1, dashed outlined
boxes) and (ii) G1 asexual V_2 and M_2 females (fig 1 grey ovals). I predicted that the 'twofold'
cost of sex (producing males) would only be paid by obligately sexual females (sexual M);
who would produce ~50% male offspring and as a consequence would produce around half
as many daughters as asexual females (asexual M and V). I predicted male production
would be negligible for all asexual females, including those which mated with a sexual male
(M) or whose mother had mated (M_2). Section 2 (f) below outlines the statistical models I
applied to make these comparisons.
181

Figure 1 Schematic of the experimental design showing (a) the 2 treatments (virgin, V;
mated, M) that sexual and asexual females were subjected to initially (G0). On day 0 mated
females were paired with a male (produced by a virgin sexual female) and observed until
copulation was recorded. Virgin females were kept in isolation. All wasps were then given
aphid nymphs on a broad bean leaf disc to parasitise for 3 days after which the parasitised
aphids were moved onto a broad bean seedling in a cellophane bag. After 14 days
parasitised aphid mummies and live emerged wasps were collected (G1), (b) sexed and
counted, and preserved in ethanol for DNA extraction. I retained 1 live G1 asexual female
per replicate from (b) to test for effects of mothers mated status on reproductive success
(virgin mothers, V_2 or mated mothers, M_2). These G1 females were given aphids to parasitise
on the day they emerged, which as with G0 were moved onto a broad bean seedling after 3
days. When the next generation (G2) wasps emerged (d) they, and all mummies, were
counted and sexed. Boxes around focal wasps (in a and c) indicate treatments which were
compared to each other to estimate the costs of obligate (grey boxes) and facultative
(dashed boxes; grey ovals) sex.

(d) Experimental set up

Generation -1: standardised rearing

To standardise rearing conditions for females to be used in experiments, recently emerged
females were removed from stock for all lines (7 asexual and 1 sexual) and 20 females per

line were transferred onto broad beans seedlings infested with *A. fabae* in cellulose bags
(Celloclair, Liestal, Switzerland). To generate enough males for the mating treatments 20
virgin sexual *Lysiphlebus* females were allowed to parasitise an aphid colony. Virgin sexual
females lay only unfertilised eggs that develop as haploid males. To ensure virginity
parasitised aphid 'mummies' were removed from infested plants in the sexual stock cage
before the wasps emerged and were isolated in cellulose pill capsules. When the wasps
emerged 20 females were transferred onto broad bean seedlings infested with *A. fabae* in a
cellulose bag. This process was repeated for each asexual line but emerged females were
removed from stock, and not isolated as mummies. This process was repeated with freshly
emerged wasps every day for 3 days to stagger the emergence of focal wasps to increase
the sample size by conducting the experiment across 3 blocks.
217
218 *Generation 0: mating treatments*
219
Ten to 12 days after aphids were exposed to the G-1 wasps I checked the seedlings for the
presence of parasitised aphid 'mummies' and isolated any that I found into cellulose pill
capsules. These were labelled with the line ID, whether the mother was mated or virgin (for
sexuals) and the date the mummy was isolated. Mummies were checked regularly for
emergence. On the peak emergence days (day 13) I selected focal females and males and
assigned them to mating treatments. I set up between 3 and 11 females per treatment for
each asexual line and 18 and 22 sexual females for the virgin and mated treatments
respectively resulting in a total sample size of N = 126 replicates.
228
Asexual female wasps from 7 lines and sexual females were assigned to either mated (M) or
virgin (V) treatments. For the M treatments a male produced by a sexual virgin female was
introduced into the pill capsule with the female, the pair was observed continuously for 30
minutes or until copulation occurred. In *Lysiphlebus fabarum* it is possible to determine
successful mating visually, the female becomes quiescent and lowers her antenna and the
male is able to insert his aedeagus into her vagina. Females often reject male mating
attempts in this species by curling their abdomen under and moving rapidly. For all M
females I recorded whether males attempted copulation and if so whether females rejected
or accepted the mating attempt. For M females that accepted copulation the male was
removed from the pill capsule and the female was retained (I did not assay the fitness of M
females which did not copulate in this experiment). For the V treatments no male was added
to the females pill capsule, to control for potential mechanical costs of mating, V females
were gently moved around in their pill capsule using a paintbrush for 5-10 seconds.
242
On each day that wasps were put into M and V treatments (over 3 days) once all females
had been subjected to their mating treatment, they were transferred onto broad bean leaves
cut into discs and embedded in 6mL 1% agarose gel in an insect rearing dish with a mesh
ventilation hole (diameter, 15mm depth; SPL Life Sciences, Pocheon-si, South Korea). Leaf
discs had been infested with 5 adult aphids 5 days previously. On the day of the experiments
the adult aphids were removed from the leaf disc and the nymphs that they produced were
counted and numbers recorded. Wasps on leaf discs were checked every day between 9:00-
10:00, 12:00-13:00 and 17:00-18:00 for 3 days after they were set up and any dead females
were recovered, and their time and date of death recorded. After 3 days the leaf disc was
removed from the agar and transferred onto a 2-week-old broad bean seedling in a cellulose

bag. If the female wasp was still alive, she was transferred onto the seedling along with the
aphids.
255
Twelve days after the aphids had been exposed to the wasps, I cut the stem of the seedling
and closed the bag (still containing the seedling) with a bulldog clip. Cutting the plant
resulted in any unparasitized aphids dropping off the plant before wasps started to emerge.
The bags were checked each morning from day 12 to day 16 and any emerging wasps were
collected. I collected one female from each replicate (for asexual females only; sexual virgin
females do not produce daughters) to start generation G1 (see below). The rest of the G1
brood were collected into 99% ethanol for genotyping. Bags containing mummies were
retained and the mummies counted after all parasitoids had emerged.
264
265 *Generation 1: daughters of asexual mated and virgin females*
266
To test whether mothers mated status influences the fitness of asexuals, I assayed the
fitness of the daughters produced by G0 asexual females in the M and V treatments.
Generation G1 females which had emerged into cellulose bags were collected subjected to
the same protocol as G0 females, but G1 focal females all remained virgin; they were the
daughters of asexual females that had either mated (labelled M_2) or remained virgin (labelled
V_2). Females were placed onto leaf discs for 3 days, which were then moved onto seedlings.
The number of aphid nymphs and wasp survival on leaf discs was also recorded in for these
G1 females. I counted the number of mummies and emerged wasps produced by these G1
females. For each combination of line by treatment I was able to set up between 3 and 11
focal females, resulting in a total sample size of N = 80 females.
277
278 (d) Assaying sperm use of asexual females
279
To test whether asexual females used sperm from sexual males after they mated, I
genotyped 41 whole broods of offspring (8 broods from asexual virgin females, 29 broods
from mated asexual females, and 4 broods from sexual females) using a single microsatellite
marker (*Lysi07*; Sandrock et al 2007) which exhibits complete linkage with the asexuality-
inducing allele (Sandrock and Vorburger 2011). DNA was extracted in a pooled reaction (up
to 10 wasps from one brood per reaction) using the Qiagen DNAeasy blood & tissue kit
(QIAGEN Inc, Hilden, Germany). After extraction DNA was amplified in a multiplex PCR
reaction using fluorescently labelled *Lysi07* primers. PCR reactions were carried out using a
Qiagen Multiplex PCR kit (QIAGEN Inc. Hilden, Germany) in 11 µl reactions and contained:
1 µL DNA template, 10 µL PCR mastermix (made up 1x QIAGEN Multiplex PCR mix and
0.1µM of each locus-specific labelled (forward) and unlabelled (reverse) primer). PCR
conditions were as follows: initial denaturation at 95°C for 15 min, followed by 30 cycles of
94°C for 30 s, 58°C for 90 s, 72°C for 60 s, and final extension at 60°C for 30 min. PCR
amplification was performed using a DNA Engine Tetrad ® Thermal Cycler (MJ Research,
Bio-Rad, Hemel Hempstead, Herts, UK). PCR products were then sent to MRC PPU DNA
sequencing and services at the University of Dundee for fragment analysis. This involves
capillary electrophoresis which provides a series of different coloured peaks which vary in
location on an X-axis based on the size of the microsatellite fragment. I scored these peaks
using the software Geneious™. For *Lysi07,* peaks scored at 184 are found exclusively in
asexual wasps and 192 in sexual wasps. When a sample had scores at both 184 and 192 it

indicated the presence of both sexual and asexual alleles suggesting that an asexual wasp
had mated and used sperm.
302
303 (e) Statistical analysis
304 *Parasitism failure, mummy production and wasp emergence*
305
To test whether mated status (and reproductive mode) were related to reproductive success
I ran 3 sets of generalised linear mixed models (GLMMs) with the following response
variables: (i) parasitism failure (binomial, scored as 1 if no mummies were recovered; using
the glmer function in the lme4 package in R; [24]), (ii) number of mummies (negative
binomial, count of the number of mummies recovered from each replicate; using *glmmTMB*
in R; [25]), (iii) proportion of mummies that emerged as adult wasps (binomial, modelled
using the cbind function in R and using *glmmTMB*).
313

The G0 models included reproductive mode (sexual or asexual), mated status (virgin or
mated) and their interaction effect as fixed factors. Covariates were also fitted; these were
the number of aphid nymphs on the leaf disc and female wasp survival (in hours; 72 hours
for all wasps which survived until transfer to seedling). I also included a random effect of
Line ID nested within reproductive mode. Block was not included in any model; treatments
were replicated evenly across daily blocks and in no case did I find a significant effect of
block on any outcome variable, nor did inclusion of block in any models influence the results.
Broods that were recorded from two G0 virgin sexual females were removed from the
analyses as female offspring were recovered (virgin sexual females only produce unfertilized
eggs which develop as males) suggesting mislabelling.

The G1 models were similar but the data was only from asexual females, so no fixed effect
of reproductive mode was fitted. The only fixed effects were mother's mated status, (mated
or virgin), survival of the focal female and number of nymphs (as covariates), and Line ID
was included as a random effect. The package *DHARMa* [26] was used to check that all
models were appropriately specified.
329
330 *Daughter production and the cost of males*
331
To compare the traditional 'twofold' cost of sex (male production), I tested whether obligately
sexual females produced fewer daughters than all asexual females (grey boxes in fig. 1). To
do this I ran a model with the number of female offspring produced as the outcome variable
(negative binomial in *glmmTMB*). Data from virgin sexual females was not used in the G0
models (as virgin sexual females produce only male offspring). The G0 model included a
fixed effect of reproductive mode. The number of male offspring produced was also included
as a covariate, and as an interaction with reproductive mode, to test for a trade-off between
son and daughter production.
340
To test whether facultative sex (by normally asexual females) imposes a cost of male
production I ran two additional models. These models had the same structure but included
data either from only G0 or only G1 asexual females and included only a fixed effect of
mated status (or mother's mated status). The G0 model tested whether asexual females that
mated produced fewer daughters than females which remained virgin. The G1 model tested

whether females with mated mothers produced the same number of daughters as females
with virgin mothers. As with the other models, mothers' survival and number of nymphs were
included as covariates and random effects were Line ID nested within reproductive mode or
and Line ID only (if only data from asexual females used).

3. Results

(a) Summary

In the first generation (G0) 103 females were offered a male to mate with (M) and 66 were
not (they remained virgin, V). Of the 103 females assigned to the M treatment, 70 mated
Thirty-three M females did not mate either because males failed to initiate courtship (N =
10), or the female did not signal receptivity (N = 23). The 70 M and 66 V female wasps were
given aphids to parasitise. Table 1 shows the numbers of females which remained virgin (V),
mated (M), or did not mate (due to rejection or no attempt) by line.

Table 1. Number of first generation G0 females that remained virgin (V), accepted a mating
(M: successful matings), did not accept a mating because the male did not attempt
courtship, or because the female rejected the male. Also shown are the proportion of male-
female introductions that resulted in a successful mating by line.

Line	Collection year	Remained virgin	Attempted matings	Successful matings	No attempt	Rejected	proportion attempts successful
CHC17-66	2017	5	5	5	0	0	1.00
CV17-84	2017	4	6	5	0	1	0.83
IL06-658	2006	3	5	5	0	0	1.00
IL07-64	2007	11	25	10	4	11	0.40
IL09-348	2009	10	14	8	1	5	0.57
IL09-402	2009	10	20	10	4	6	0.50
IL09-554	2009	5	5	5	0	0	1.00
Sexual	Various	18	23	22	1	0	0.96

366
367

Of the 136 G0 females that were given aphids to parasitise 58 died before they were
transferred onto the seedlings and 70 survived. In the second generation (G1), I took the
daughters of 40 M and 40 V G0 asexual mothers and put them on leaf discs. Of the 80
females transferred onto leaf discs 57 survived and 23 died before being transferred to
seedlings 72 hours later (note that there was no effect of mating treatment on wasp survival
in the first 72 hours in either the G0 or G1 generations: G0 (effects of reproductive mode and
mated status): Log-rank X^2= 1.4, df = 3, p = 0.70 and G1 (mated status only): Log-rank X^2 =
2.8, df = 1, p = 0.33).

(b) Sperm use

Forty-one broods were genotyped in total (see table S1 for a summary of brood genotypes
broken down by treatment and line). All broods produced by sexual females were
homozygous for the sexual allele (peak 192) at *Lysi07*. Broods produced by asexual females
were significantly more likely to test positive for sexual alleles if the female had mated (M)
than if she was virgin (V) (X^2 =4.56, df = 1, p = 0.03); 9/30 broods produced by mated M
asexual females were homozygous for the asexual allele (peak 184), 19/30 were

heterozygous for sexual and asexual alleles (184/192) and the 2 were homozygous for the
sexual allele (192; most likely due to a PCR failure/scoring error or misnumbering of a
replicate as there is no evidence for introgression of sexual alleles in the stock lines these
females were derived from). Only 1 brood produced by a virgin (V) asexual female was
scored as positive (heterozygous) for the sexual allele (suggesting either mislabelling or
contamination between samples); the other 7 broods were homozygous for the asexual
allele.
392
(c) Parasitism success

In many cases no mummies were recovered from aphids that were exposed to *L. fabarum*
females, suggesting that these females either failed to oviposit, or their eggs failed to
hatch/larvae to develop resulting in complete reproductive failure. In G0 100 out of 126
females successfully parasitised aphids and produced adult offspring. A few individuals
successfully produced a small number of mummies but no adult offspring (12), these
females were excluded from the analyses below as they may have produced diapause
larvae. In the first generation G0, sexual females were more likely to fail to parasitise aphids
than asexual females (= 4.49, df = 1, 124, p = 0.03). Mating had no effect on G0 females
ability to parasitise aphids and this was true for both sexual and asexual females (= *1.54,*
df = 1,124, p = 0.21. Interaction effect between reproductive mode and mated status: =
0.42, df = 1, 124, p = 0.52; fig. 2a).

In G1 54/80 daughters of asexual G0 females successfully parasitised aphids and produced
adult offspring, 6 females successfully produced a small number of mummies but no adult
offspring (again, females which produced mummies but not adult offspring were removed
from the analysis below). In G1 there was a large negative effect of mothers mated status on
parasitism success. The G1 daughters of mated M G0 asexual females were 25% more
likely to fail to parasitise aphids than the daughters of virgin V females (= *7.43, df = 1,79,*
p = 0.0.006; fig. 2b)*.*

Figure 2. Bar plot showing the effect of mated status on % parasitism success in **(a)** G0
across 7 asexual lines and one outbred sexual line of *L. fabarum*. In G0 sexual females were
more likely to fail to parasitise than asexuals but there was no effect of mated status and **(b)**

G1 in asexual daughters of G0 females; daughters of mated M females were significantly
more likely to fail to parasitise than daughters of virgin V females.

(d) Mummy production and emergence

Figure 3 Number of mummies produced (**a & b**) and % mummies adult wasps emerged
from (**c & d**) in G0 (**a & c**) and G1 (**b & d**). G0 sexual females produced more mummies than
asexual females (**a**) which were more likely to emerge as adult wasps (**c**). Mummies
produced by virgin mothers (sexual and asexual) were more likely to emerge as adult wasps
(**b**). For asexual G1 females there was no effect of mothers mated status on mummy
production (**c**) or the proportion of mummies that successfully emerged (**d**). Error bars
represent 95% confidence intervals.

Although sexual females in G0 were more likely to fail to parasitise any aphids, when I
considered only the females that did parasitise, sexual females were at an advantage
producing more mummies ($X^2 = 5.04, df = 1, 98, p = 0.03$; fig. 3a) more of which produced a
wasp ($X^2 = 4.41, df = 1, 98, p = 0.03$; fig. 3c) than their asexual conspecifics. Whilst sex had
an effect on mummy production in G0 there was no effect of mating; virgin V and mated M
females produced the same numbers of mummies ($X^2 = 0.30, df = 1, 98, p = 0.59$)
regardless whether they were sexual or asexual (interaction: $X^2 = 0.29, df = 1, 98, p = 0.59$;
fig. 3a). There was an effect of mating on adult offspring production though; virgin V G0

mothers were more successful ($X^2 = 3.85, df = 1, 98, p = 0.05$; fig. 3c) for sexual and
asexual mothers (interaction: $X^2 = 0.78, df = 1, 98, p = 0.38$).

These results were not replicated in the second generation, there was no effect of G0
mother's mated status on the number of mummies their G1 daughters produced ($X^2 = 2.41,$
$df = 1, 52, p = 0.121$; fig. 3b) or the proportion of these mummies that emerged ($X^2 = 0.21,$
$df = 1, 52, p = 0.65$; fig. 3d).

Daughter production and the cost of producing males

To test whether the classic 'twofold' cost of sex applies in *L. fabarum* I looked at daughter
production by focal females. In G0, obligately sexual females did not produce fewer
daughters than asexual females (M or V; $X^2 = 0.02, df = 1, 98, p = 0.90$). Contrary to
expectations females that produced more males did not produce fewer daughters (number of
sons: $X^2 = 0.68, df = 1, 98, p = 0.40$), a pattern that was the same for sexual and asexual
females (reproductive mode*number male offspring: $X^2 = 3.23, df = 1, 98, p = 0.07$). I ran a
second model with sexual females removed to test whether facultative sex by G0 mated
asexual females imposes a cost in terms of male production, but again M and V females
produced the same number of daughters ($X^2 = 0.49, df = 1, 71, p = 0.48$). Son production
was very low for M and V asexual females in G0 (< 5 per mother) and there was no
evidence that females which produced more males produced fewer daughters (mated status:
$X^2 = 0.49, df = 1, 71, p = 0.48$; male production; $= 3.22, df = 1, 71, p = 0.07$; mated
status*number male offspring $X^2 = 0.36, df = 1, 71, p = 0.55$).

I found the same pattern in the second generation (G1); male production was again very low
and had no effect on daughter production ($X^2 = 1.88, df = 1, 52, p = 0.17$) regardless of
mothers mated status (mothers mated status*number male offspring: $X^2 = 0.65, df = 1, 52, p$
$= 0.42$). Importantly the G1 daughters of mated M G0 females (some of which are
facultatively sexual) produced the same number of daughters as females produced
asexually by virgin V G0 mothers ($X^2 = 2.24, df = 1, 52, p = 0.14$). Taken together these
results from G0 and G1 suggest that the twofold cost of male production does not constrain
the fitness of obligately or facultatively sexual females in *L. fabarum*.

4. Discussion

Populations of the aphid parasitoid *Lysiphlebus fabarum* vary in their reproductive mode:
asexual (parthenogenetic) reproduction dominates across Europe, but sexual populations
persist in sympatry with asexuals in some regions [17,18]. In this study I show for the first
time that asexual *L. fabarum* females are sexually functional and can use sperm to fertilise
eggs and produce daughters. I found that some females from all 7 asexual *L. fabarum* lines
assayed were facultatively sexual; they mated with sexual males and produced offspring
with genetic contributions from both parents. However, asexual females that reproduced
sexually did not get the best of both worlds, in terms of short-term benefits of
parthenogenesis (higher daughter production) or genetic benefits of sex (increased fecundity
or offspring emergence). In fact, the pattern was quite the opposite. The main short-term
benefit of asexual reproduction is that it eliminates the 'twofold cost of males'. In *L. fabarum*

asexuals do occasionally produce male offspring, Sandrock and Vorburger [19] found that around 1 in every 3000 asexual wasps is male in the lab. In the current study I also found low rates of male production by *all* asexual females, but crucially asexual females did not produce more daughters than sexual females. Obligately sexual females had higher fecundity than females from asexual lines, such that despite producing many more sons they produced the same number of daughters as asexuals, suggesting a long-term genetic benefit of sex. Females from asexual lines which engaged in facultative sex did not gain this benefit of sex though, the fecundity of asexual females was not influenced by their mated status or their mothers mated status. There was a significant fitness cost to sex though, it increased the risk of reproductive failure by 12% compared to parthenogenesis. This cost was even higher for asexual females that mated and reproduced sexually; when asexual G0 females mated and could reproduce sexually, 40% of their G1 daughters failed to reproduce compared to only 10% of the G1 daughters of asexual G0 females that did not mate.

Asexual reproduction is thought to have first evolved approximately 0.5 million years ago in *L. fabarum,* arising through the inheritance of a single recessive allele which results in the haploid products of meiosis I fusing to form diploid gametes which develop without fertilisation (central-fusion automixis; [19]). Although asexuality may have arisen 0.5 million years ago it is unlikely that the asexual lineages I tested have existed for so long without sex and recombination (see [19]). The asexual lines used in this experiment have, however been reared in the lab for 7-18 years without sex. Despite the lack of sex (potentially for up to 400 generations) some females from all seven asexual *L. fabarum* populations exhibited sexual behaviour and were able to integrate paternal DNA into the female offspring they produced. The maintenance of sexual traits in *L. fabarum* makes them unique compared to many other asexual eukaryotes, where sexual traits such as mate location, attraction, allowing copulation and storing sperm decay rapidly due to high costs of maintenance [27]. In other parasitoid species that have reverted to asexual reproduction due to PI (Parthenogenesis Inducing) *Wolbachia* infection, females no longer produce costly volatile chemoattractants which render them attractive to males (i.e. [28–30]). In these cases sex, fertilisation and associated traits are rapidly lost due to selection on the bacteria and the wasps. In terms of the former, parthenogenesis increases the spread of PI-*Wolbachia* (which is maternally inherited through the cytoplasm). Selection on loss of fertilisation in the wasps occurs because increasingly female biased population sex ratios under PI-*Wolbachia* means that females who leave more eggs unfertilised to develop as haploid males will have higher fitness. As such selection on parasitoid and endosymbiont appears to result in rapid conversion to parthenogenesis as well as the loss of sexual traits which would allow facultative sex (see [28]). In *L. fabarum* parthenogenesis is inherited through the nuclear genome, not an endosymbiont, which may be one reason why I found that males find and court asexual females from some lines as readily as sexual females (see Table 1), suggesting that both sexual and asexual females invest in olfactory signals to attract males.

What remains unclear is why sexual traits have persisted in asexual *L. fabarum* despite being costly in terms of reproductive success. Whilst obligate sexual females do produce, on average, more offspring, both obligate and facultative sex are associated with a higher risk of failure to produce any offspring in *L. fabarum*. One explanation for this, which applies to facultative and obligate sex is 'genetic slippage', which occurs when combinations of genes that normally work well together are broken up by sex, creating new genotypes which have low fitness [3–8,12]. Sex and recombination can produce novel multilocus genotypes that

perform better than the parental genotypes in some environments, but genetic slippage can
mean that they perform much worse [12]. In this study it seems that the new genotypes
created by sex were costly for *L. fabarum,* and this cost may have been more severe under
facultative sex because because normally asexual females have gone without sex for so
long (giving beneficial allele combinations more time to build up than when sex is regular).
Another, non-mutually exclusive explanation for the high reproductive failure of G1 daughters
of facultatively sexual females (G0 M females) is that some them may have been triploid with
low fertility. Triploid females may represent a proportion of the females in G1 that failed to
produce any offspring, a possibility that is supported by field evidence; Sandrock et al [18]
found around 2% of females in all-female broods were triploid. However, triploidy is not a
given under central fusion automixis as eggs are laid when they are in metaphase I, but
diploidy is not restored until anaphase II[20]. Follow up studies which replicate this
experiment but genotype individual mothers and offspring are needed to shed light on the
extent to which triploidy vs genetic slippage leads to reproductive failure in the daughters of
mated asexual females.
540
The mating systems and population dynamics characteristic of aphid parasitoids like
Lysiphlebus might explain why facultative sex has persisted in *L. fabarum* despite the costs.
Parasitoid abundance increases over the season as hosts become more numerous, until
populations crash later in the summer [31]. At the end of the season the abundance of *L.*
fabarum females in parasitised aphid colonies becomes very high which could have a
number of effects on the rate of facultative sex and its consequences. First, high population
densities are expected to increase the likelihood that normally asexual females will
encounter and mate with rare asexual males (or sexual males in regions of sympatry).
Second, mating between kin is common in parasitoids because many females parasitise
aphids in the same colony that they emerged from, which may counter the detrimental
effects of genetic slippage (i.e. they are less likely to produce low fitness recombinants with
closely related mating partners: [32]). Genetic slippage is essentially a form of severe
outbreeding depression [33] , and so its costs may be reduced under inbreeding. Third, the
main cost of sex (increased reproductive failure), may be relaxed at certain times in the
season. For instance, during the 'mid-season crash' aphid populations become locally
extinct over just a few days, in part due to high natural enemy pressure [34]. At such times
the cost of sex may be less significant if the costs of host-finding and host competition
constrain parasitoid fitness to a greater extent.
559
In addition to relaxed costs, facultative sex could also be an important 'bet-hedging' strategy
for *L. fabarum* late in the season if it reduces the variation in fitness across environments
compared to parthenogenesis. Bet-hedging has been proposed to maintain traits that are
costly for individual fitness, like sex (and polyandry [35,36]) ensuring the genes underlying
these traits persist even if the traits themselves can be costly to the bearers. Williams [37]
used the analogy of reproduction as a lottery to explain sex as bet-hedging: in this analogy
asexual reproduction is like buying many lottery tickets with the same number, if you win,
you win big. Reproducing sexually, you don't buy so many tickets, but they have different
numbers, increasing the chances you win, but reducing the amount you win. Sex and
recombination result in more genetically diverse offspring, which differ from each other and
their parents. By producing sexual offspring with more diverse genotypes, some of them will
have low fitness in the same conditions as their parents. The genotypes that perform poorly
in the parental environment may be better adapted to different conditions, they may thrive

when the environment changes [8,37]. Asexual reproduction on the other hand tends to
result in offspring that are genetically very similar, if not identical, to their parents, and so do
very well in the same conditions their parents experienced, but if the environment changes
they are more likely to fail to reproduce entirely. Under bet-hedging, sex is a risk-averse
strategy associated with moderate fitness under a range of environments, while asexual
reproduction could provide big benefits under a stable environment but is risky if conditions
change. The scenario that aphid parasitoids like *L. fabarum* face at the end of the season
may increase the importance of facultative sex as a bet-hedging strategy because some
offspring will need to survive overwinter as diapause pre-pupae and reproduce the following
spring, in drastically different environmental conditions from their parents.
583
When viewed in the light of the natural history of aphid parasitoids like *L. fabarum* the results
of this study suggest that relatively frequent sex, potentially between closely related males
and parthenogenetic females, may be responsible for maintaining high levels of genetic
variation and heterozygosity in asexual *L. fabarum.* If sex occurs mainly late in the season it
is also likely to be fairly synchronous, mirroring the reproductive phenology of their cyclically
parthenogenetic aphid hosts. When normally parthenogenetic individuals all reproduce
sexually at the same time in this way, Kokko [11] has shown that facultative sex is more
stable and robust to the invasion of obligate asexual reproduction. Synchronous seasonal
sex between siblings may explain not only the how heterozygosity and allelic diversity is
maintained in asexual *L. fabarum* [17,18] but also why sex and its associated traits have
persisted. Studies that measure the frequency of sex in the field, and its associated costs will
reveal the full extent of facultative parthenogenesis and its evolutionary significance in *L.*
fabarum, so that we can establish whether facultative sex has in this case made asexual
reproduction more successful.
598
599 **Data accessibility**
600
601 All data and code can be found on github https://github.com/DrBecky-
602 B/Asexual_mating_Lysi
603
604 **References**
605

1. Maynard-Smith J. The evolution of sex. Cambridge: Cambridge
University Press; 1978.
2. Simon J-C, Delmotte F, Rispe C, Crease T. Intraclonal genetic variation:
ecological and evolutionary aspects. Biol J Linn Soc. 2003;79: 151–163.
3. Schwander T, Crespi BJ. Twigs on the tree of life? Neutral and selective
models for integrating macroevolutionary patterns with microevolutionary
processes in the analysis of asexuality. Mol Ecol. 2009;18: 28–42.
doi:10.1111/j.1365-294X.2008.03992.x
4. Lehtonen J, Jennions MD, Kokko H. The many costs of sex. Trends Ecol
Evol. 2012;27: 172–178. doi:10.1016/j.tree.2011.09.016
5. Fisher RA. The genetical theory of natural selection. Oxford: Clarendon;
1930.
6. Muller HJ. Some genetic aspects of sex . Am Nat. 1932;66: 118–138.

7. Muller HJ. The relation of recombination to mutational advance. Mutat
Res. 1964;1: 2–9.
8. Hartfield M, Keightley PD. Current hypotheses for the evolution of sex
and recombination. Integr Zool. 2012;7: 192–209. doi:10.1111/j.1749-
4877.2012.00284.x
9. West SA, Lively CM, Read AF. A pluralist approach to sex and
recombination. J Evol Biol. 1999;12: 1003–1012. doi:10.1046/j.1420-
9101.1999.00119.x
10. Aanen D, Beekman M, Kokko H. Weird sex: The underappreciated
diversity of sexual reproduction. Phil Trans Roy Soc B Biol Sci.
2016;371. doi:10.1098/rstb.2016.0262
11. Kokko H. When synchrony makes the best of both worlds even better:
How well do we really understand facultative sex? Am Nat. 2020;195:
380–392. doi:10.1086/706812
12. Allen DE, Lynch M. Both costs and benefits of sex correlate with relative
frequency of asexual reproduction in cyclically parthenogenic Daphnia
pulicaria populations. Genetics. 2008;179: 1497–1502.
doi:10.1534/genetics.107.082479
13. Schwander T, Vuilleumier S, Dubman J, Crespi BJ. Positive feedback in
the transition from sexual reproduction to parthenogenesis. Proc R Soc
Lond B Biol Sci. 2010;277: 1435–1442. doi:10.1098/rspb.2009.2113
14. Judson OP, Normark BB. Judson Normark 1996 ancient asexual
scandals. Trends Ecol Evol. 1996;11: 41–46. doi:10.1016/0169-
5347(96)81040-8
15. Schwander T. Evolution: The end of an ancient asexual scandal. Curr
Biol. 2016;26: R233–R235. doi:10.1016/j.cub.2016.01.034
16. Domes K, Norton RA, Maraun M, Scheu S. Reevolution of sexuality
breaks Dollo's law. Proc Nat Acad Sci. 2007;104:7139-7144.
doi:10.1073/pnas.0700034104
17. Belshaw R, Quicke DLJ, Volkl W, Charles AH, Godfrayl J. Molecular
markers indicate rare sex in a predominantly asexual parasitoid wasp.
Evolution. 1999;53: 1189–1199. doi:10.1111/j.1558-5646.1999.tb04532.x
18. Sandrock C, Schirrmeister BE, Vorburger C. Evolution of reproductive
mode variation and host associations in a sexual-asexual complex of
aphid parasitoids. BMC Evol Biol. 2011;11. doi:10.1186/1471-2148-11-
348
19. Sandrock C, Vorburger C. Single-Locus recessive inheritance of asexual
reproduction in a parasitoid wasp. Curr Biol. 2011;21: 433–437.
doi:10.1016/j.cub.2011.01.070
20. Belshaw R, Quicke DLJ. The cytogenetics of thelytoky in a
predominantly asexual parasitoid wasp with covert sex. Genome.
2003;46: 170–173. doi:10.1139/g02-112

21.   Boulton RA, Zuk M, Shuker DM. An Inconvenient Truth: The
662   unconsidered benefits of convenience polyandry. Trends Ecol Evol.
663   2018;33: 904–915. doi:10.1016/j.tree.2018.10.002

664   22.   Sandrock C, Frauenfelder N, Von Burg S, Vorburger C. Microsatellite
665   DNA markers for the aphid parasitoid *Lysiphlebus fabarum* and their
666   applicability to related species. Mol Ecol Notes. 2007;7: 1080–1083.
667   doi:10.1111/j.1471-8286.2007.01783.x

668   23.   Najafpour P, Rasekh A, Esfandiari M. The effect of choice and no-choice
669   access on host (Aphis fabae) instars preference, in different ages and
670   sizes of Lysiphlebus fabarum (Hymenoptera: Braconidae: Aphidiinae). J
671   Entomol Soc Iran. 2016;36: 99–109.

672   24.   Bates D, Mächler M, Bolker B, Walker S. Fitting linear mixed-effects
673   models using lme4. J Stat Softw. 2015;67. doi:10.18637/jss.v067.i01

674   25.   Brooks ME, Kristensen K, Benthem KJ, van, Magnusson A, Berg CW,
675   Nielsen A, et al. glmmTMB Balances speed and flexibility among
676   packages for zero-inflated generalized linear mixed modeling. R J.
677   2017;9: 378. doi:10.32614/RJ-2017-066

678   26.   Hartig F. DHARMa: residual diagnostics for hierarchical (multi-
679   level/mixed) regression models. R package version 020; 2018.

680   27.   van der Kooi CJ, Schwander T. On the fate of sexual traits under
681   asexuality. Biol Rev. 2014;89: 805–819. doi:10.1111/brv.12078

682   28.   Pijls JWAM, Van Steenbergen HJ, Van Alphen JJM. Asexuality cured: the
683   relations and differences between sexual and asexual Apoanagyrus
684   diversicornis. Heredity. 1996;76: 506–513. doi:10.1038/hdy.1996.73

685   29.   Pannebakker BA, Beukeboom LW, Van Alphen JJM, Brakefield PM,
686   Zwaan BJ. The genetic basis of male fertility in relation to haplodiploid
687   reproduction in *Leptopilina clavipes* (Hymenoptera: Figitidae). Genetics.
688   2004;168: 341–349. doi:10.1534/genetics.104.027680

689   30.   Stouthamer R, Russell JE, Vavre F, Nunney L. Intragenomic conflict in
690   populations infected by Parthenogenesis Inducing *Wolbachia* ends with
691   irreversible loss of sexual reproduction. BMC Evol Biol. 2010. doi:
692   10.1186/1471-2148-10-229

693   31.   Jervis MA, Kidd NAC, Mills NJ, van Nouhuys S, Singh A, Yazdani M.
694   Population Dynamics. In: Hardy I, Wajnberg E, editors. Jervis's Insects
695   as Natural Enemies: Practical Perspectives. Cham: Springer
696   International Publishing; 2023. pp. 591–667. doi:10.1007/978-3-031-
697   23880-2_7

698   32.   Lynch M, Deng H-W. Genetic slippage in response to sex. Am Nat.
699   1994;144: 242–261. doi:10.1086/285673

700   33.   Lynch M. The genetic interpretation of inbreeding depression and
701   outbreeding depression. Evolution. 1991;45: 622–629.
702   doi:10.1111/j.1558-5646.1991.tb04333.x

703   34.   Karley AJ, Parker WE, Pitchford JW, Douglas AE. The mid-season crash
704   in aphid populations: Why and how does it occur? Ecol Entomol.
705   2004;29: 383–388. doi:10.1111/j.0307-6946.2004.00624.x
706   35.   Yasui Y, Garcia-Gonzalez F. Bet-hedging as a mechanism for the
707   evolution of polyandry, revisited. Evolution. 2016;70: 385–397.
708   doi:10.1111/evo.12847
709   36.   Li X-Y, Lehtonen J, Kokko H. Sexual reproduction as bet-hedging. In:
710   Apaloo J, Viscolani B, editors. Advances in Dynamic and Mean Field
711   Games Theory, Applications, and Numerical Methods. Cham,
712   Switzerland: Birkhäuser; 2017. pp. 218–235.
713   doi:https://doi.org/10.1007/978- 3- 319- 70619- 1
714   37.   Williams GC. Sex and evolution. Princeton, New Jersey: Princeton
715   University Press; 1975.
716