Corticosterone release in very young siblicidal seabird chicks (*Rissa tridactyla*) is sensitive to environmental variability and responds rapidly and robustly to external challenges

Z.M. Benowitz-Fredericks[1], A.P. Will[2,3], S.N. Pete[1], S. Whelan[4] & AS Kitaysky[3]*

* corresponding author (zmbf001@bucknell.edu)
1 Bucknell University, Department of Biology 1 Dent Drive, Lewisburg PA
2 World Wildlife Fund, US Arctic Program
3 University of Alaska Fairbanks, Department of Biology and Wildlife, Institute of Arctic Biology
4 Institute for Seabird Research and Conservation

Keywords: development, facultative siblicide, food availability, glucocorticoids, HPA-axis, ontogeny

<u>Abstract.</u>

In birds, patterns of development of the adrenocortical response to stressors vary among individuals, types of stressors, and species. Since there are benefits and costs of exposure to elevated glucocorticoids, this variation is presumably a product of selection such that animals modulate glucocorticoid secretion in contexts where doing so increases their fitness. In this study, we evaluated hypothalamic-pituitary-adrenal (HPA) activity in first-hatched free-living seabird nestlings that engage in intense sibling competition and facultative siblicide (black-legged kittiwakes, *Rissa tridactlya*). We sampled 5 day old chicks (of the ~45 day nestling period), a critical early age when food availability drives establishment of important parent-offspring and intra-brood dynamics. We experimentally supplemented parents with food ("supplemented") and measured chick baseline corticosterone secretion and capacity to rapidly increase corticosterone in response to an acute challenge (handling and a 15 min restraint in a bag). We also used topical administration of corticosterone to evaluate the ability of chicks to downregulate physiologically relevant corticosterone levels on a short (minutes) time scale. We found that 5 day old chicks are not hypo-responsive but release corticosterone in proportion to the magnitude of the challenge, showing differences in baseline between parental feeding treatments (supplemented vs non-supplemented), moderate increases in response to handling, and a larger response to restraint (comparable to adults) that also differed between chicks from supplemented and control nests. Topical application of exogenous corticosterone increased circulating levels nearly to restraint-induced levels and induced downregulation of HPA responsiveness to the acute challenge of handling. Parental supplemental feeding did not affect absorbance/clearance or negative feedback. Thus, while endogenous secretion of corticosterone in young chicks is sensitive to environmental context, other aspects of the HPA function, such as rapid negative feedback and/or the ability to clear acute elevations in corticosterone, are not. We conclude that 5 day old kittiwake chicks are capable of robust adrenocortical responses to novel challenges, and are sensitive to parental food availability, which may be transduced behaviorally, nutritionally, or via maternal effects. Questions remain about the function of such rapid, large acute stress-induced increases in corticosterone in very young chicks.

1. Introduction

The activity of the vertebrate hypothalamo-pituitary adrenal (HPA) axis is often considered in terms of costs and benefits. At a basal level, activity of the axis and the consequent release of glucocorticoid hormones is mandatory for survival, normal development (Wada, 2008), and modulation of metabolism (Jimeno and Verhulst, 2023), while the ability to rapidly increase glucocorticoids allows animals to maintain or reestablish homeostasis in the face of internal or external challenges (Sapolsky et al., 2000). While developmental exposure to glucocorticoids can increase fitness, presumably by better-preparing offspring for environmental conditions (Crino and Breuner, 2015), prolonged exposure to glucocorticoids during development (whether chronic or repeated acute elevations) can be associated with costs such as cognitive impairment, and reduced growth, immune function, and neural development (Kitaysky et al., 2003; Sapolsky and Meaney, 1986; Schoech et al., 2011). Thus, as with any process presumably shaped by selection, developing animals should only increase glucocorticoids when doing so increases fitness (i.e., benefits outweigh costs).

In birds, there is variation among species in the developmental stage at which chicks begin to respond to environmental challenges by activating the HPA axis and increasing circulating corticosterone (e.g. Adams et al., 2008; Bebus et al., 2020; Wada et al., 2007), as well as variation in which stimuli can elicit an HPA axis response from nestlings (e.g. Lynn and Kern, 2018, 2014). Many altricial and some semi-altricial species have a post-hatch "hypo-responsive" period (the "stress non-responsive period" in mammals) of relatively low HPA axis responsiveness (Bebus et al., 2020; Blas et al., 2006; Fiske et al., 2013, Jones et al., 2021; Sims and Holberton, 2000; Tilgar et al. 2009; Wada et al., 2009, 2007). This developmental pattern is thought to protect young chicks who are both vulnerable to excessive glucocorticoid exposure and too early in development to benefit from elevated glucocorticoids because they have life-histories that leave them unable to change physiology or behavior in sufficiently rapid or appropriate ways to increase fitness (Fiske et al., 2013; Sims and Holberton, 2000; Torres☐Medina et al., 2019; Wada, 2008; Wada et al., 2007). However, some species have life-histories that involve complex early-life social dynamics that may be influenced by glucocorticoids (reviewed in Smiseth et al., 2011) and would therefore be expected to activate the HPA axis early in life.

For nest bound chicks that compete with siblings for limited parental resources via begging,

aggression, and, in some species, siblicide, corticosterone may be a relevant mediator of social

behavior (Loiseau et al., 2008; Vallarino et al., 2006). While glucocorticoids in adults are

thought to help mediate the tradeoffs associated with allocating resources and behavior between

self-maintenance and reproduction to maximize fitness (Bókony et al., 2009), chicks should be

selected to maximize survival and optimize somatic growth such that they can successfully

recruit to the breeding population. Food availability may be one of the most important factors

that affects chicks' growth and survival, behavior, and adrenocortical activity (Kitaysky et al.,

2003). In some species, presumably those with few options available to increase food intake,

food availability has minimal impacts on corticosterone release (Kitaysky et al., 2005).

However, in other species, survival, and optimal growth in the face of limited food availability

can evoke intense intra-brood competition at a very young age (reviewed in Morandi and Ferrer,

2014, Osorno and Drummond, 2003). These species may be selected for HPA activation early in

postnatal life, despite potential costs of elevating glucocorticoids during development.

Food availability affects sibling competition in semi-precocial black-legged kittiwake (*Rissa*

tridactyla; hereafter "kittiwake") chicks (White et al., 2010). Kittiwakes reside in cliffside nests

with 0 - 2 siblings and rely on parental food provisioning for ~40-45 days before fledging

(Hatch et al., 2020). Both parents forage at sea and deliver regurgitated meals to chicks multiple

times a day. Chicks engage in facultative siblicide, where the first-hatched chick (hereafter "A-

chick") often out-competes the second-hatched chick (hereafter "B-chick") and monopolizes

parental food resources. A-chicks accomplish this by interrupting B-chick begging and feeding

bouts, driving them out of the nest with repeated attacks, or killing them outright (Braun and

Hunt, 1983; reviewed in Dickins, 2021). Most siblicide occurs during the first 10 days post-

hatch, with chicks capable of intense aggression just a few days after hatching (Braun and Hunt,

1983). While the proximate cause of death is rarely clear (often the B-chick is simply missing

from the nest), in 2022, hour-long observations at 28 nests documented aggression in 13 of 28

nests and found that only 5 of those 28 B-chicks survived to fledge (ZMBF unpubl.) While there

are dozens of hypotheses about the evolution of brood reduction, most of them center on

maximizing fitness when resources are limited (Morandini and Ferrer, 2015). Thus, compared to

more "helpless" altricial nestlings, the complex behavioral repertoire of young kittiwake chicks

is likely to be highly sensitive to energy balance and resource availability, and therefore may be

modulated by glucocorticoids.

Black-legged kittiwake physiology has been studied intensively, including several investigations

of HPA activity (both baseline and restraint-induced elevations of circulating corticosterone) in

captive and free-living chicks (Table 1). These studies confirm that food availability is likely to

be an important driver of variation in kittiwake chick HPA activity. Captive studies demonstrated

clear consequences of food intake (both quality and quantity) on HPA activity of older chicks

(29-31 days) subject to several weeks of experimental diets (Kitaysky et al., 1999), with low

quality/quantity diet treatments dramatically increasing both baseline and restraint-induced

corticosterone, but there was no change in baseline corticosterone with age in captive chicks fed

ad libitum (Table 1). However, these captive chicks were raised in isolation, without any of the

natural, complicated social interactions with parents and siblings which may influence HPA

activity. The interannual and intercolony differences observed in free-living 12–15-day old

chicks (Brewer et al., 2010, 2008) was hypothesized to have been driven by differences in food

availability. While experimental food supplementation to free-living kittiwake nests at a different

colony did not generate large differences in corticosterone in at least some breeding seasons

(Schultner et al., 2013; Young et al., 2017), the lack of treatment effect was attributed to the high

food availability naturally available to control (non-supplemented) adults in those years.

Thus, overall, corticosterone release in kittiwake chicks appears to be responsive to

environmental and social context at a variety of ages. However, all but one of these studies

(Merkling et al., 2014; baseline corticosterone only) measured corticosterone in chicks that had

already passed the most critical developmental window in terms of competition and resource

monopolization, which occurs between age 3- and 10-days post hatch (Merkling et al. 2014).

Intrabrood aggression drops to very low levels by 10 days post hatch (Leclaire et al., 2011;

White et al. 2010), and 62-80% of B-chicks who died during the years of this study did so when

both chicks were 10 days of age or younger (Whelan and Hatch, unpubl.). Thus, this early stage

of development is critical for fitness, yet information about adrenocortical physiology at this age

is lacking, a gap we sought to fill with this study. Because kittiwake life-history involves

complex early-life social dynamics that may be influenced by glucocorticoids, we tested the

hypothesis that the HPA axis of 5 day old black-legged kittiwakes is not "hypo-responsive" at

very early developmental stages but is instead already sensitive to environmental context

(interannual variability, experimental food supplementation) and physiological state (genetic sex,

body condition). We measured baseline and restraint-induced corticosterone, as well as

corticosterone uptake, negative feedback and clearance following exogenous corticosterone

administration. We predicted that very young kittiwake chicks have well-developed

adrenocortical activity, which is dampened by the experimental food supplementation of their

parents.

Table 1. Review of studies measuring corticosterone in first-hatched black-legged kittiwake
(*Rissa tridactyla)* chicks. Superscript letters indicate whether corticosterone
radioimmunoassays were conducted using the same antibody (a = Esoterix, b = Biogenesis)
and/or in the same lab (a = Wingfield, UW, b = O'Reilly, UP, c = Chastel, CEBC, d =
Kitaysky, UAF). Note that comparison of absolute levels across studies may only be
appropriate if hormone assays were conducted in the same lab using the same antibody, given
the variation generated by inter-laboratory differences in extraction and assay methods, as well
as specificity of antibodies used (Fanson et al. 2007). "~" indicates that age is presented as
approximate in some studies, or that hormone levels were estimated from figures.
147

Citation	Age (days)	Location	Baseline corticosterone (ng/mL)	Restraint-induced max corticosterone (ng/mL)	Sources of variation in corticosterone	No significant effect of …
[a/a]Kitaysky, Piatt et al. 1999	29-31	Captivity (*source - Alaska: Cook Inlet*)	4-5; food ad libitum	30; high food	(-) food intake; (-) body composition (lipid content)	—
			16; food restricted	100; food restricted		
[a/a]Kitaysky et al. 2001	~16	Alaska: Cook Inlet	~7	—	—	—
[a/a]Kitaysky et al. 2003 (*including unpubl. data*)	14-56 (weekly samples); 75	Captivity (*source - Alaska: Cook Inlet*)	2.9; food ad libitum	31 (@ 75 days only)	—	age
[a/b]Brewer et al. 2008; 2010	12-15	Alaska: Shoup Bay	3.3 - 7.4	~30-35 (peak: 10 min)	Year (highest in year of lowest productivity); colony	brood size; hatching order
[b/c]Merkling et al. 2014	5	Alaska: Middleton Island	~3-4	—	hatching order x experimental hatching asynchrony interaction; (-) growth rate	sex; body condition
[a/d]Schultner et al. 2013	25	Pacific - Alaska, Middleton Island	1.6; control	14.8; control	population	supplemental feeding, parental corticosterone manipulation
			1.4; food supplemented	10.3; food supplemented		
		Atlantic - Norway: Svalbard	3.4; control	26.2; control		
[a/d]Young et al. 2017	10	Alaska: Middleton Island	2.5	—	(-) age; hatching order; brood swapping	sex; food supplementation; brood enlargement
	25		1.5			

2. Methods

2.1 Study site

This research was conducted in two years (2021 and 2022) at a free-living colony of black-legged kittiwakes nesting on a modified radar tower on Middleton Island, in the Gulf of Alaska (59°25'41.2"N 146°20'10.3"W). Details of the study site are described in Gill et al. (2002) and Gill and Hatch (2002), but in short: semi-partitioned nest sites on the outside face of the tower are each equipped with a sliding one-way mirrored glass window. The sliding glass provides access to individual nests from inside the tower for monitoring and manipulation with minimal disturbance to neighboring nests. All nests are monitored twice daily for events including egg-laying, hatching, chick death, and fledging. Chicks are not banded until day 25 post-hatch, so younger chicks are identified by hatching order using colored pen marks, applied on hatch day and refreshed every 5 days afterwards. Thus ages are known to the day. Tower nest sites are inaccessible to both terrestrial and aerial predators.

2.2 Food supplementation

The tower is also the site of a long-term food-supplementation experiment in which a subset of nests (hereafter "supplemented") are provided with whole fish (either capelin (*Mallotus mallotus)* or herring *(Clupea spp.);* both preferred prey items) ad libitum, 3 times per day for the entirety of the breeding season (Gill et al. 2002), beginning several weeks before egg-laying. Adults readily consume the supplemental food; chicks consume the food supplement once they are large enough to consume whole fish (never observed prior to 7 days post-hatch, thus chicks in this study were not yet directly consuming the supplement). Adults at supplemented nests continue to forage at sea, but benefit from the supplementation in terms of reproductive success in most years (e.g. Kahane-Rapport et al., 2022). Occupants of control (hereafter "non-supplemented") nests do not receive any supplemental food and rely solely on foraging at sea.

2.3 Blood sampling

In both years and regardless of treatment, all blood sampling occurred between 0600 and 0845, prior to the first daily supplemental feeding for supplemented nests at 0900. All chicks had been briefly handled only once before: on hatch day for measurements and marking. To prevent parents from startling abruptly and dislodging chicks, we started a timer then gently tapped on

windows to alert parents and induce them to stand before retrieving a chick to take a blood

sample. All blood samples were taken within 3 minutes of this initial tapping disturbance (2021:

range = 38 - 180 sec, mean. = 82 sec, SD = 32 sec; in 2022: range = 45 - 172 sec, mean = 89 sec,

SD = 25 sec). At each blood sampling event (hereafter "bleed"), the alar vein was punctured

using a sterile 26G needle and 100-200 µL of whole blood was collected in heparinized

microhematocrit tubes. Samples were refrigerated for up to 3 hours. They were then centrifuged

to separate erythrocytes from plasma, both of which were frozen at -20°C for up to three weeks,

then transported off the island and shipped on dry ice to the lab where they were stored at -80°C

until use.

2.4 Experiment 1: Sources of variation in baseline corticosterone and response to restraint

(Fig 1)

In 2021, between 24 Jun and 7 Jul, chicks from supplemented and non-supplemented nests

containing two chicks were bled for baseline corticosterone ("undisturbed"; Table 2). Nests

were sampled when the first-hatched A-chick was 5 days old (second-hatched B-chick ages

ranged from 2 to 5 days old). For each nest, both A- and B- chicks were removed simultaneously

and the A-chick was bled in < 3 min from initial disturbance. Chicks were then weighed and

measured, B-chicks were returned to the nest and A-chicks were placed in a breathable cloth bag

in proximity to a space heater, since 5 day old chicks do not have full thermoregulatory capacity.

Fifteen min after initial disturbance, A-chicks were bled a second time to evaluate short-term

adrenocortical response to restraint, then returned to their nest. Previous serial samples of

restrained three-week old kittiwake chicks at this site took blood samples at < 3, 10, 30 and 50

min and found that corticosterone was highest at 10 min in some individuals, and at 30 min in

others (Kitaysky, unpubl.). Because it is not possible to identify the timing of the true peak, we

selected 15 min as a time point likely to capture variation in rates of increase, if not the peak

increase. In 2022, baseline blood samples were collected from A-chicks between 29 June and 4

July for interannual comparisons.

2.5 Experiments 2 and 3: Responses to exogenous corticosterone

To bypass some of the challenges and constraints associated with the use of subcutaneous

corticosterone implants (Torres-Medina et al, 2018) and minimize changes in HPA activity

associated with handling, we used topical administration of corticosterone in dimethyl sulfoxide

(DMSO) to rapidly elevate corticosterone with minimal disturbance. This manipulation allowed

us to evaluate differences in glucocorticoid uptake and/or clearance, as well as negative

feedback. DMSO is a polar solvent that has been used in veterinary and human medicine as a

delivery vehicle for topical application of lipophilic compounds for decades (Kumar and Darreh-

Shori, 2017; Patil, 2013). It is considered non-toxic, though it can have off-target cellular effects

such as inhibition of acetylcholinesterase, and is used by itself for a variety of conditions to

reduce inflammation (Santos et al., 2003). In avian endocrinology, it has been used successfully

for topical application of corticosterone (Busch et al., 2008; Pegan et al., 2019; Vitousek et al.,

2018).

2.5.1 Preparation of exogenous corticosterone treatments

Topical treatments were made following Pegan et al. (2019), though with a higher ratio of liquid

DMSO to facilitate pipetting and surface area of contact on skin. In short, for the "CORT"

treatment, 0.05 g corticosterone powder (Sigma #27840) was dissolved in 4 mL liquid DMSO,

which was mixed with 6 mL veterinary grade DMSO gel to achieve a corticosterone

concentration of 5 μg/μL. A control DMSO gel solution was made using the same ratio of liquid

to gel DMSO, but with no corticosterone.

We used average body masses from 5 day old chicks measured in 2019 (90g) to anticipate that a

20 μL dose would deliver approximately 1 microgram of corticosterone per gram of body mass.

This dose was selected based on previous experiments with topical administration of

corticosterone in DMSO in birds (Busch et al, 2008, Pegan et al. 2019) in which it elevated

corticosterone within a naturally occuring range. We used food coloring to visually confirm that

vortexing rapidly and thoroughly dispersed liquid DMSO throughout the gel (though did not

include dye in the final solutions). In the years of our study, average body mass of 5 day old

chicks was lower than in 2019 (73g in 2021; 75g in 2022), such that 20μL of gel would have

been a higher dose than intended. However, while 20μL of gel was drawn into each pipette tip,

the viscosity of the gel meant that several drops were always left behind on the inside of the

pipette tip. Repeated measurements in the lab using a microbalance to weigh pipetted DMSO gel

mix (n = 20; compared to the weight of 20µL dispensed using a displacement syringe) indicated

that an average of 12.4 ± 0.0019 µL of gel was delivered to each chick (57% of the dose

originally calculated for 90g chicks). Thus, the average dose delivered to a 74g chick was 0.62

µg corticosterone/g body mass. We confirmed that this dose elevated corticosterone in kittiwake

chicks to levels comparable to an endogenous response to handling in this study (see Fig. 4B)

2.5.2 Administration of topical treatments

In 2021 and 2022, 5-day old - chicks from either one- or two-chick nests were treated with a

drop of either corticosterone in DMSO gel (hereafter "CORT" treatment) or DMSO gel

(hereafter "DMSO" treatment; see section 2.5.1). CORT and DMSO gels were kept in physically

different locations (and away from where chicks were bled) with dedicated pipettes, tip boxes,

and glove boxes, to minimize the risk of corticosterone contamination. Chicks were treated after

tapping on the nest window to alert the parent and opening the window part way (causing the

brooding parent to stand or briefly flush). We then reached into the nest with a pipettor set to 20

µL, and gently touched the tip to the skin on the back of the A-chick between the scapulae and

behind the neck, dispensing the gel there (see supplemental material for video footage). We

administered the gel to this location, where it is tucked behind the head when a chick is sitting, to

minimize the likelihood that the gel was dislodged or transferred to brooding parents.

Application of the treatment usually took ~15 seconds from initial disturbance of the nest site to

completion (closing of window), unless the brooding parent was particularly aggressive, in

which case it could take up to 30 seconds to work around the adult. Parents resumed brooding

within seconds of window closure. Chicks were not weighed until the final blood sample was

taken; chicks were excluded from further analyses *a priori* if they weighed < 50 g, a threshold

below which chicks have dramatically reduced survival.

2.6 Experiment 2: Effects of exogenous corticosterone on endogenous production (Fig. 1)

We used A-chicks from food supplemented and non-supplemented nests containing one or two

chicks (Table 2). We applied the topical gel treatment, took a blood sample 15 minutes after gel

application and immediately returned the chick to the nest. Fifteen min after the first sample (i.e.

30 min after gel treatment), we removed the chick to take a second blood sample to evaluate the

effect of exogenous corticosterone on subsequent endogenous corticosterone production in

response to the handling that occurred during the first bleed. Finally, though DMSO is estimated

to take at least 5 minutes to penetrate the skin and enter the bloodstream (Patil 2013), nothing is

known about the acute time course of corticosterone uptake from topical DMSO application in

young seabird chicks. Therefore, we checked for very rapid changes in corticosterone by treating

a small subset of chicks with topical CORT or DMSO and taking a single blood sample 5

minutes after treatment administration (Table 2).

2.7 Experiment 3: Time course of exogenous corticosterone (Fig. 1)

In 2022, we repeated the topical CORT experiment (mixing fresh batches of gel solutions using

the same protocol), but only used chicks from non-supplemented nests containing two chicks.

Chicks were distributed across 4 treatment groups. Half of the topical CORT-treated chicks and

half of the topical DMSO-treated chicks were bled for the first time 15 min after topical

application to confirm rapid increase in corticosterone in CORT group, and the other half of the

topically treated chicks were bled for the first time 60 min after application to evaluate chicks'

ability to clear exogenous corticosterone from circulation. Corticosterone levels in these four

groups were compared to baseline levels in the same year (see Experiment 1, section 2.4).

2.8 Genetic sexing

All chicks were genetically sexed from DNA extracted from erythrocytes (Qiagen DNeasy Blood

and Tissue kits, Cat # 69556). We used primers and protocols for PCR and gel electrophoresis

described in Merkling et al. (2012) to differentiate the sexes by number of bands.

2.9 Corticosterone assays

Corticosterone was measured blind to treatment using radioimmunoassay, following the protocol

described in Kitaysky et al. (2007). In short, plasma was equilibrated with tritiated

corticosterone, extracted using dichloromethane, and assayed in duplicate. 2021 and 2022

samples were run in separate assays (three and one assays in 2021 and 2022, respectively), but

all samples from the same chick were run in the same assay, with treatments distributed across

assays. Each assay included a standard treated identically to samples to calculate interassay

variability (interassay CV = 2.7%). Final corticosterone values were corrected for the percent of

tritiated corticosterone lost during extraction (average recovery for 2021 = 96% +/- 3.18%, for

2022 = 92% +/- 2.96%).

2.10 Data analysis

To meet assumptions for parametric statistical tests, all corticosterone values were log-

transformed, which normalized their distribution. Corticosterone concentrations among food, and

CORT treatment groups had equal variance in all but one case (see below for modeling

approach).

We first determined whether brood size (one or two chicks) affected baseline corticosterone

concentrations using a Welch's two-sample t-test and tested for differences in body condition

between years and feeding treatments (ANOVA) to account for potential confounding effects

among explanatory variables on the measured physiological response (i.e. corticosterone

concentration).

2.10.1 Experiment 1: Sources of variation in baseline corticosterone and response to

restraint

To test which ecological and biological factors affect 5-day old chicks' baseline corticosterone

and adrenocortical response to 15 min of handling and restraint (including whether there is a

response) we used a model selection approach. We built an *a priori* candidate models of interest

(Table 3) using a small set of variables that have been previously shown to affect adrenocortical

response in birds: body condition, sex, feeding treatment, and in baseline adrenocortical function.

We used Akaike's Information Criterion (Burnham and Anderson, 2004) to identify the top

performing model(s), defined as all models with AIC scores with a difference of < 2 from the top

performing model. We calculated the effect size for the variables from each best fit model and

considered any variable with an effect size whose 95% confidence interval does not overlap zero

as statistically significant. We calculated AIC scores in R (v. 4.2.1, R Core Team 2022) using

packages *AICcmodavg* (Mazarolle 2023). We calculated effect size using the package *effectsize*

(Ben-Shachar et al., 2020), and tested variables in the top models for multicollinearity using the

vif function in the *multcomp* package (Hothorn et al., 2008).

2.10.2 Experiment 2: Effects of exogenous corticosterone on endogenous production

To test the effects of topical application of corticosterone via DMSO on chick's subsequent adrenocortical response we ran a single mixed effects ANOVA testing whether corticosterone concentrations (both from the first and second bleed) differed between treatment groups. We included body condition as a covariate, and individual chick as a random effect. The variance in corticosterone levels at the second bleed was not equal among treatments, so we also accounted for unequal variance in the model. To test for the effect of feeding treatment on the response of chicks we ran two ANCOVA, one for corticosterone at the first bleed, and the other for corticosterone at the second bleed. For both models, feeding treatment was the predictor and body condition was a covariate.

2.10.3 Experiment 3: Time course of exogenous corticosterone

We used the same mixed effects ANOVA model described above to test for differences in response over time to the exogenous treatments. To test for differences in the magnitude of the response between the first and second bleed in treatment groups we ran two difference ANCOVAs with the change in corticosterone concentration (i.e. delta corticosterone) as the response, corticosterone treatment (CORT or DMSO) and feeding treatment (supplemented or non-supplemented) as predictors and body condition as a covariate. The first was to examine change in corticosterone at comparable time intervals among the two exogenous treatments only. The second model included the delta corticosterone values for the control group (the difference between baseline and post-bag restraint).

To test for differences in clearance of corticosterone over time among the treatment groups, for non-supplemented chicks only, we ran an ANCOVA with corticosterone concentration as the response, exogenous treatment as a predictor, and body condition as a covariate.

2.10.4 Post-hoc analyses

We followed up all ANOVA and ANCOVA models with post-hoc comparisons and report p-values and F statistics from a Type III ANOVA using Satterthwaite's method for modeled variables, and p-values from post-hoc Tukey contrasts. We analyzed data in R (v. 4.2.1, R Core Team 2022) packages *nlme* (for mixed effects ANOVA, Pinheiro and Bates 2023) and *multcomp*

364   (for ANCOVA and post-hoc contrasts, Hothorn et al., 2008). We considered variables and

365   contrasts with p-values < 0.05 as significant.

368   # 3. Results

Table 2. Circulating corticosterone levels (mean ± sd; ng/mL) in 5 day old, first-hatched black-legged kittiwake (*Rissa tridactyla)* chicks by treatment and time. All chicks sampled on Middleton Island, Alaska. Parents at supplemented nests were provided with whole fish ad libitum 3x/day; non-supplemented relied only on natural food avilability. DMSO = treated topically with DMSO (control) and CORT = treated topically with corticosterone dissolved in DMSO. 15 min restraint values are from the same chicks bled for baseline samples, and 30 min DMSO and CORT bleeds in 2021 are from the same chicks in the 15 min DMSO and CORT groups. NA = no samples collected.

	Year			
	2021		**2022**	
	Parental food supplementation			
Treatment/time	**Non- supplemented**	**Supplemented**	**Non- supplemented**	**Supplemented**
Baseline (1st bleed)	15.4 ± 10.9 (n=19)	3.4 ± 1.9 (n=17)	14.5 ± 15.6 (n=10)	10.8 ± 7.5 (n=12)
15 min restraint (2nd bleed)	54.7 ± 10.7 (n=19)	51.0 ± 17.9 (n=17)	NA	NA
5 min DMSO (1st bleed)	14.4 ± 12.6 (n=4)	NA	NA	NA
5 min CORT (1st bleed)	24.4 ± 20.5 (n=4)	NA	NA	NA
15 min DMSO (1st bleed)	11.0 ±7.7 (n=16)	11.6 ±14.6 (n=11)	13.7 ± 10.9 (n=11)	NA
15 min CORT (1st bleed)	41.0 ±22.8 (n=17)	25.5 ± 19.2 (n=9)	34.7 ± 16.7 (n=11)	NA
30 min DMSO (2nd bleed)	31.8 ± 17.0 (n=16)	23.8 ±13.5 (n=11)	NA	NA
30 min CORT (2nd bleed)	57.2 ±20.9 (n=17)	45.1 ±20.4 (n=9)	NA	NA
60 min DMSO (1st bleed)	NA	NA	6.4 ± 3.8 (n=14)	NA
60 min CORT (1st bleed)	NA	NA	27.7 ± 16.1 (n=14)	NA

3.1 Experiment 1: Sources of variation in HPA activity

3.1.1 Sources of variation in baseline corticosterone

There was no difference in A-chick baseline corticosterone between one- and two-chick nests

($t_{18.16}= -0.06$, p = 0.95), so brood size was omitted from subsequent analyses. Chick body

condition did not differ between years ($F_{1,55} = 1.69$, p = 0.2), or feeding treatments ($F_{1,54} = 1.77$,

p = 0.28).

Model comparison indicated strong support for an effect of feeding treatment that differed

between years, with both top models including the year*treatment interaction (Table 3). Post-hoc

analyses of the top model indicated significant effects of year, treatment and their interaction -

baseline corticosterone was higher in non-supplemented nests (effect size ± 95% confidence

interval from top model: 0.99 ± 0.58, 1.41) but more so in 2021 than 2022 (-0.73 ± -1.16, -0.31),

and was higher in chicks from both supplemented and non-supplemented nests in 2022 (0.57 ±

0.27, 0.87; Table 2, 3). The next best performing model included body condition; however, it

was not an informative predictor on its own (-0.0042 ± -0.32, 0.31). Body condition did explain

some of the variability with respect to feeding treatment (Fig 2) and so was included as a

covariate in subsequent analyses.

Table 3. Model selection results of variables affecting baseline corticosterone secretion of 5-day

old chicks and *a priori* justification for model inclusion. Treatment = food supplementation to

parents (food supplemented or non-supplemented). Top performing models are italicized.

Model	K	AICc	Delta AICc	AICc Wt	*a priori* justification
*Treatment*Year*	*5*	*38.93*	*0*	*0.56*	*Effects of parental nutritional status depend on interannual environmental variability*
*Year*Treatment*Body Condition*	*9*	*39.66*	*0.73*	*0.39*	*The effect of parental nutritional status on the relationship between chick energy stores and corticosterone varies interannually*
Full Two-Way Interaction Model*	12	44.31	5.38	0.04	
Year+Treatment*Body Condition	6	48.37	9.45	0	Interannual environmental variation affects corticosterone, but parental nutritional status also modulates the impact of chick energy stores on corticosterone
Treatment	3	49.35	10.43	0	Parental nutritional status affects corticosterone
Treatment*Body Condition	5	49.36	10.44	0	Parental nutritional status modulates the impact of chick energy stores on corticosterone
Year+Treatment+Body Condition	5	49.83	10.91	0	Corticosterone is affected by interannual environmental variability, parental nutritional status, and chick energy stores
Treatment+Sex	4	50.25	11.32	0	Corticosterone is affected by parental nutritional status and chick sex.
Treatment+Body Condition	4	51.27	12.35	0	Both chick energy balance and parental nutritional status affect chick corticosterone

Year*Treatment+Body Condition+Sex	6	51.36	12.44	0	Corticosterone is affected by interannual environmental variability, parental nutritional status, chick energy balance and sex
Treatment*Sex	5	52.39	13.46	0	Parental nutritional status affects corticosterone in male and female chicks differently
Null (intercept only) Model	2	62.82	23.9	0	None of these variables explain variation in corticosterone
Year	3	63.14	24.21	0	Corticosterone reflects interannual environmental variability
Year+Body Condition	4	63.5	24.58	0	Corticosterone reflects both energy balance and interannual environmental variability
Body Condition	3	63.82	24.9	0	Corticosterone reflects energy balance
Year*Body Condition	5	64.08	25.16	0	The effect of chick energy balance on corticosterone is modulated by interannual environmental variability
Sex	3	64.1	25.18	0	Corticosterone reflects sex-specific differences
Body Condition*Sex	5	64.14	25.22	0	Corticosterone reflects energy balance in sex-specific ways
Year+Sex	4	64.55	25.63	0	Corticosterone reflects both sex and interannual environmental variability
Body Condition+Sex	4	65.55	26.62	0	Corticosterone reflects both energy balance and sex
Year*Sex	5	66.76	27.83	0	Male and female chicks respond differently to impacts of interannual environmental variability on corticosterone

392 *BCI+Sex+Treatment+Year+BCI*Sex+BCI*Treatment+BCI*Year+Sex*Treatment+Sex*Year+Treatment*Year
393
3.1.2 Endogenous corticosterone secretion in response to restraint. After 15 minutes of

handling and restraint in a bag, 5-day old chicks increased corticosterone substantially (Table 2,

Fig. 3, 4). The top model for corticosterone after 15 min of restraint included the interaction

between baseline corticosterone and feeding treatment (effect size: 1.55 ± 0.65, 2.44; Table 4). In

nests where parents were supplementally fed, higher baseline corticosterone was associated with

lower HPA response - this was not the case in non-supplemented nests. The next best performing

model included all two-way interaction terms and indicated that body condition was negatively

correlated with chicks' HPA response (effect size: -0.57 ± -1.05, -0.08).

Table 4. Model selection results of variables affecting corticosterone response to 15 minutes of
handling and restraint in 5-day old chicks. Baseline corticosterone is log-transformed. Treatment
= food supplementation to parents (supplemented or non-supplemented). *A priori* justification is
the same as in Table 3 with the addition of baseline corticosterone: a chick's baseline
physiological state affects their response to a stressor, and that response may exist in parallel or
be modified by the other study variables. Since chick sex was not a significant predictor of
baseline corticosterone but there were significant interactions among the other variables of
interest, we included a full two-way interaction without sex in the model set. The top performing
model is italicized.

Model	K	AICc	Delta AICc	AICcWt
*Treatment*Baseline corticosterone*	5	-49.38	0	0.66
Two-Way Interaction Model without Sex*	8	-47.29	2.09	0.23
Body Condition	3	-42.28	7.1	0.02
Treatment+Body Condition+Baseline corticosterone	5	-41.24	8.14	0.01
Null (intercept only) Model	2	-41.23	8.15	0.01
Treatment+Body Condition	4	-40.84	8.54	0.01
Treatment+Baseline corticosterone	4	-40.27	9.11	0.01
Treatment	3	-40.16	9.22	0.01
Body Condition+Baseline corticosterone	4	-39.86	9.52	0.01
Sex+Body Condition	4	-39.83	9.55	0.01
Sex	3	-39.28	10.1	0
Treatment*Body Condition	5	-38.96	10.41	0
Baseline CORT	3	-38.87	10.51	0
Treatment+Sex+Body Condition+Baseline corticosterone	6	-38.86	10.52	0
Treatment*Body Condition+Baseline corticosterone	6	-38.81	10.57	0
Treatment+Sex	4	-38.16	11.22	0

Treatment+Sex+Body Condition+Baseline corticosterone	5	-37.67	11.7	0
Treatment*Sex	5	-37.26	12.12	0
Sex*Body Condition	5	-37.15	12.23	0
Sex*Baseline corticosterone	5	-36.96	12.42	0
Sex+Baseline corticosterone	4	-36.8	12.58	0
Full Two-Way Interaction Model including Sex**	12	-32.27	17.1	0

*Treatment+Sex+BC+BaselineCorticosterone+Treatment*Sex+Treatment*BC+Treatment*BaselineCorticosterone
+Sex*BC+Sex*BaselineCorticosterone+BC*BaselineCorticosterone
**Treatment+BC+BaselineCorticosterone+Treatment*BC+Treatment*BaselineCorticosterone+BC*BaselineCortic
osterone

3.3 Experiment 2: Exogenous corticosterone and negative feedback

3.3.1 Response to exogenous corticosterone treatment

Topical application of corticosterone significantly increased plasma corticosterone ($F_{5, 84} =$

61.08, p < 0.0001, controlled for effect of body condition $F_{1, 88} = 5.76$, p = 0.018, with individual

as a random effect). Post-hoc tests showed that the CORT treatment elevated circulating

corticosterone compared to undisturbed baseline (p < 0.001) and to the DMSO treatment (p <

0.001), but the identical application of DMSO alone did not elevate corticosterone compared to

undisturbed baseline (p = 0.998), confirming that neither DMSO nor the topical administration

procedure evoked significant changes in HPA activity (Table 2; Fig 4). The topical CORT

treatment increased circulating corticosterone to levels that were slightly (but significantly; p =

0.001) lower than levels secreted endogenously in response to 15 minutes of restraint (Fig 4A/B).

Parental feeding treatment had a strong effect on corticosterone levels 15 min after exogenous

corticosterone application ($F_{1,50} = 4.56$, p = 0.037, controlling for body condition $F_{1,50} = 0.34$, p =

0.56) -- chicks from supplemented nests had much lower circulating corticosterone 15 min after

the topical hormone application than chicks from non-supplemented nests (Table 2; Fig 4). This

difference between supplemented and non-supplemented groups mirrored the difference between

them in the baseline group: i.e. corticosterone in non-supplemented chicks was 12 ng/mL higher

than in supplemented chicks at baseline. A separate small subset of chicks bled 5 min after gel

application showed no significant difference between the DMSO and the CORT treatments ($t_5 =$

-1.07, p = 0.33; Table 2).

3.3.2 Effects of exogenous treatment on endogenous response to handling.

Both the CORT and DMSO-treated chicks increased circulating corticosterone at the second

bleed (30 min after topical gel application and 15 min after the first bleed) compared to their

respective first bleeds (post-hoc comparison from mixed-effects ANCOVA, DMSO p < 0.001,

CORT p = 0.003; Table 2; Fig 4C). Although the corticosterone in the CORT treated group

remained higher at the second bleed compared to the DMSO group (p < 0.001; Fig 4B), the

adrenocortical response (magnitude of change) was larger in the DMSO treatment ($F_{1,49}$ =

8.6626, p = 0.005; Fig 4C).

In addition, DMSO and CORT treated chicks from supplemented nests had lower corticosterone

at both the first ($F_{1,50}$ = 5.17, p = 0.03) and second bleed ($F_{1,50}$ = 4.56, p = 0.04) compared to

those from non-supplemented nests (in both cases controlling for body condition, though body

condition was not significant; bleed 1: $F_{1,50}$ = 0.12, p = 0.45; bleed 2: $F_{1,50}$ = 0.02, p = 0.56; Fig

4A/B). However, feeding treatment did not affect negative feedback. The differences in

corticosterone levels between parental feeding treatments at the second bleed were of the same

magnitude as the difference between feeding treatments at the 1st bleed (15 min after gel

application; $F_{1,49}$ = 1.2688, p = 0.26). Consequently, there were no differences between feeding

treatments in the magnitude of change in corticosterone within CORT or DMSO treatments

(bleed 2 corticosterone - bleed 1 corticosterone; Fig 4C). Finally, the increase in endogenous

corticosterone after the brief disturbance of handling followed by return to the nest for ~ 15 min

(delta corticosterone for the DMSO treated group) was much lower than the endogenous increase

in corticosterone after 15 min of continuous bag restraint ($F_{2.82}$ = 21.29, p < 0.0001; Fig 4C).

3.4 Experiment 3. Clearance of exogenous corticosterone

In non-supplemented nests in 2022, circulating corticosterone in DMSO-treated chicks bled for

the first time 15 or 60 min after treatment did not differ from undisturbed baseline corticosterone

(Tukey Contrasts, p = 1 and 0.63 respectively; Fig 5, Table 2). Corticosterone in the CORT

treatment was higher than baseline and DMSO at both 15 (p = 0.01 and 0.007) and 60 min post-

treatment (p = 0.02 and <0.001) but was not different between the 15- and 60-min CORT groups

(p = 0.97; Fig 5).

4. Discussion

In this study, we demonstrate that kittiwake chicks do not exhibit a post-hatch HPA-axis

hypo-responsive period. Instead, they have HPA axes that a) can rapidly secrete a large amount

of corticosterone and b) are sensitive to environmental and physiological context. At five days

old, just 10-15% of the way through the nestling stage, chicks mounted corticosterone responses

proportional to the intensity of the challenge, with levels 15 min after a moderate challenge (brief

handling to take a blood sample followed by return to the nest) that were higher than the

undisturbed baseline, but lower than levels evoked by a more intense challenge (15 min of

continuous restraint). Chick corticosterone levels after the 15 min restraint challenge were not

only much higher than their baseline levels, but higher than any of the previously reported

maximum values for kittiwake chicks with the exception of much older, severely food-restricted

captive chicks (though all previous studies that measured restraint-induced levels used chicks

older than 5 days; Table 1). Restraint-induced levels of corticosterone in chicks were comparable

to (with some individuals higher than) maximal levels secreted by adult kittiwakes at this colony

during 50 min of restraint (12 - 63 ng/mL; Kitaysky et al., 2010). In addition, nutritional status

of parents interacted with interannual variability and body condition to affect baseline HPA axis

activity, though neither chick sex nor number of siblings did. Our study design minimized

potential sources of variation in HPA activity by using kittiwake chicks of known age, hatching

order and sex, and sampling within a narrow hatch date range (< 2 weeks) and time of day (< 3

hours).

4.1 Food availability to parents and year effects

Despite the fact that 5-day old chicks are very early in development and parents should be able to

meet their relatively low food needs at this stage, there were strong effects of year and parental

supplemental feeding treatment on chick HPA activity. Baseline corticosterone was significantly

lower in chicks from food supplemented nests in one of two years. An obvious explanation for

this difference might be that the current nutritional status of parents is transmitted to chicks via

differences in provisioning, which affects chick nutritional status and thereby their HPA activity.

However, two lines of evidence indicate that this was a contributing factor but not the sole

driver. First, parental food treatment had a strong effect in 2021 even when chick body condition

was controlled for statistically. Second, there were no significant differences in chick body

condition between years or feeding treatments, yet their baseline circulating corticosterone levels

differed between years. In fact, although climate indices and measures of reproductive success

indicated that 2021 and 2022 were both years of intermediate reproductive success for kittiwakes

at this site, 2022 was a better year than 2021 (S. Hatch, unpublished data). Yet baseline

corticosterone was higher in 2022 than in 2021, and in 2022 chicks from supplemented nests had

corticosterone levels that were similar to non-supplemented. Though reproductive success at the

end of the season is temporally distant from conditions at 5 days post hatch, the mismatch

between annual breeding success and young chick endocrine physiology suggest that the

differences in chick baseline corticosterone are affected by food availability but also modulated

by additional factors that we were unable to measure in this study.

There are several potential modulators of the relationship between chick corticosterone and

current food availability. First, it could be a result of prenatal effects. Supplemental feeding at

experimental nests began before egg-laying, potentially interacting with early season or even

overwinter environmental conditions in each year to influence egg composition (Benowitz-

Fredericks et al., 2013). Variation in egg composition can affect chick HPA activity (Ahmed et

al., 2014; Love and Williams, 2008; Schwabl and Partecke, 2020; Sockman and Schwabl 2001)

and may be a mechanism for intergenerational transmission of information that prepares chicks

for the environment into which they are born (reviewed in Engqvist and Reinhold, 2016). In

addition, adults show high nest site fidelity at this colony, thus most parents at supplemented and

non-supplemented nests have likely experienced the same feeding treatments in past years. This

combination of early indicators of natural food availability during the pre-lay period (Whelan et

al., 2021) and anticipated changes in food availability during chick-rearing based on prior

experience may shape investment in reproduction (Shultz et al., 2009), and thus the

programming of chicks' HPA axes. Second, the differences in HPA activity could be driven by

post-natal behavioral cues from parents or siblings. While chick body condition indicated that

parents seemed to be meeting the nutritional needs of their chicks at this early age regardless of

the supplemental food treatment, differences in competition from siblings or responsiveness of

parents to begging may impact corticosterone secretion (Braasch et al., 2014; Dreiss et al.,

2016). Finally, other aspects of interannual environmental variability could modulate the

relationship between food availability and corticosterone by imposing additional energetic

challenges. For example, weather can affect chick corticosterone (Crino et al., 2020; De Bruijn

and Romero, 2018), as can ectoparasites (Dudek et al., 2021; Kitaysky, 2001; Quillfeldt et al.,

2004).

4.2 Body condition

Control chicks (from non-supplemented nests) in higher body condition had lower baseline

corticosterone levels, and body condition was generally negatively correlated with restraint-

induced corticosterone levels. While negative relationships between body condition or body

mass and corticosterone have been frequently reported in birds, including adult black legged

kittiwakes (Kitaysky et al. 1999) and red-legged kittiwake chicks (Kitaysky et al. 2001), a

phylogenetically controlled meta-analysis found no consistent relationship between baseline

corticosterone and body condition in seabirds (Sorenson et al., 2017). In adult kittiwakes, low

body condition can be associated with either low or high levels of corticosterone (Schultner et

al., 2013) - the first because animals with a dependable food supply and low corticosterone can

afford to avoid the costs of carrying extra weight (Rogers, 2015), and the second because severe

food shortages elevate corticosterone, which mobilize and deplete reserves. Growing chicks are

under different selection pressures than adults and may be less likely to benefit from maintaining

a "lean" phenotype, though the relationships between growth and fitness in kittiwakes are

complex and context-dependent (Vincenzi et al., 2013; Vincenzi and Mangel, 2013). Indeed, in

this experiment the relationship between body condition and baseline corticosterone was

eliminated by supplemental feeding. While some models predict that the relationships between

glucocorticoids and phenotype will be obscured by high resource availability (Breuner and Berk,

2019), there were no differences in body condition between treatments in the young chicks

included in this study. This indicates that at least at this age, factors in addition to internal

indicators of energy reserves may modify chick HPA activity. For example, other aspects of

interannual variability, such as weather, may be buffered to different extents by parents in the

different supplemental feeding treatments. The negative relationship with response to restraint

indicates that chicks in better condition may be able to buffer themselves from potential costs of

elevated glucocorticoids exposure by downregulating their sensitivity to challenges. However, it

is also worth noting that food intake can change tissue composition of chicks including increased

water retention (Kitaysky et al., 1999) and, potentially, differences in rate of yolk absorption
(Noy and Sklan, 2001), such that measures of body condition in young chicks may not always
provide an accurate proxy of nutritional status.
565
4.3 Genetic sex
There are relatively few studies examining sex differences in chick HPA activity. While sexual
dimorphism in other traits might be hypothesized to predict sexual dimorphism in physiology,
there is variation among species in whether chick HPA axis activity varies by sex, with some
showing sex differences in HPA responsiveness before dimorphism emerges (e.g. Lynn et al.
2022) and others showing no differences despite sexual dimorphism (e.g. Sockman and Schwabl
2001). Though young kittiwake chicks are initially sexually monomorphic, males grow faster
and are generally considered more costly for parents to rear (Merkling et al., 2012; Young et al.,
2017). While the dimorphism manifests when chicks are older than 5 days (Merkling et al.
2012), the physiological substrates for these differences may be established earlier. One study
measured baseline corticosterone in 5-day old kittiwakes and, similar to this study, found no
effect of sex. However, none of the additional aspects of HPA-function that we measured in this
study (response to restraint, negative feedback) varied by sex either. Whether a sex difference
manifests later in development as dimorphism increases, or whether the effects of elevated
corticosterone on physiology and behavior differ by sex even when HPA activity does not,
remains to be determined.
582
4.4 Sources of variation in restraint-induced corticosterone.
The magnitude of the corticosterone response to restraint in chicks has been proposed to reflect
consistent phenotypic variation in adrenocortical responsiveness with fitness implications,
predicting subsequent recruitment and survival in at least one species (Blas et al. 2007). What
drives variation in restraint-induced corticosterone levels in chicks? Though we did not measure
it here, HPA axis activity has a heritable component in many birds, with higher heritability of
restraint induced levels than baseline levels (Beziers et al. 2019, Jenkins et al. 2014). While body
condition may also play a role (see section 4.2), the strongest evidence in this study was for a
negative relationship between baseline corticosterone and corticosterone after restraint, but only
among chicks in the food-supplemented treatment. Chicks from supplemented nests secreted

more corticosterone to reach similar levels as their counterparts from the non-supplemented nests

after 15 min of restraint, despite starting at much lower baseline levels. While these results

suggest that lower baseline corticosterone levels in chicks from supplemented nests was not a

consequence of a difference in adrenal capacity, they do not identify sources of interindividual

variation within treatments.

Variation in adrenal capacity may be explained in part by the "facilitation" phenomenon, in

which restraint-induced corticosterone levels reflect adrenal capacity as shaped by HPA activity

in the recent past (Akana et al., 1992; Dallman et al., 2004, 1992; Kitaysky et al. 2007). In a

study of adult kittiwakes, maximum restraint-induced corticosterone was correlated with food

availability two weeks prior to sampling (Kitaysky et al., 2010). Though the adrenal capacity of

very young chicks has had limited opportunity to be shaped by post-hatch experience, the HPA

axis of avian embryos is active and capable of responding to stimuli (Jenkins and Porter, 2004)

and its activity in chicks may be shaped by both prenatal and postnatal exposure to

corticosterone (Benowitz-Fredericks et al. 2015; Marasco et al., 2016), as well as environmental

context (Groothuis et al., 2020). Thus, it is possible that prenatal experience plays a role (see

section 4.1) and the HPA responsiveness of chicks to restraint is determined by a combination of

ontogenetic changes and embryonic/early post-hatch exposures to external cues.

4.5 Effects of exogenous glucocorticoids.

We used topical corticosterone administration to evaluate rapid uptake, clearance, and negative

feedback of glucocorticoids. The ability to clear circulating glucocorticoids via a combination of

negative feedback and metabolism is an important aspect of HPA function that likely has fitness

consequences, as it allows animals to buffer themselves from costs of extended glucocorticoid

exposure (Lattin and Kelly, 2020; Romero, 2004; Romero et al., 2009; Taff et al., 2018). Though

the effects were not apparent after 5 minutes, topical corticosterone treatment elevated

circulating corticosterone within a physiologically relevant range after 15 min. This elevation

persisted for at least 60 min. In contrast, administration of the vehicle control (DMSO alone) did

not change corticosterone levels, compared to undisturbed baseline levels, at any point.

We did not find evidence that chicks from supplemented and non-supplemented nests processed

exogenous corticosterone differently. The difference between parental feeding treatments in

levels of corticosterone 15 min after administration of exogenous corticosterone was similar to

the difference between them at baseline. However, the second bleed for the DMSO and CORT

treated chicks provided evidence that a) exogenous corticosterone evoked rapid negative

feedback on endogenous production (the magnitude of the handling-induced increase was higher

in the DMSO control treatment compared to the CORT treatment) and b) the levels of negative

feedback did not differ by parental feeding treatment (the magnitude of difference in

corticosterone between CORT treated chicks in supplemented and non-supplemented nests at the

2nd bleed were the same magnitude of difference as in the first bleed). Thus, while endogenous

secretion of corticosterone in 5-day old chicks is sensitive to environmental context, rapid

negative feedback and/or the ability to clear acute exogenous elevations in corticosterone are not.

Though altered sensitivity to negative feedback has been proposed as a potential mechanism to

explain ontogenetic changes in HPA activity in chicks, this hypothesis was not supported in

white storks (Torres☐Medina et al., 2019). Similarly, our study suggests that it does not explain

variation in HPA activity between young kittiwake chicks from supplemented and non-

supplemented nests, though age-related changes have not yet been tested.

4.6 Conclusion.

The developmental trajectory of avian HPA axes has been hypothesized to be driven by a

species' position on the altricial-precocial continuum (Blas et al., 2006), latitude (Jones et al.,

2021), or by other aspects of life-history including provisioning strategies, but no consistent

patterns have emerged (Sagar et al., 2019). The HPA axes of black-legged kittiwake chicks are

sensitive to internal and external cues and are capable of mounting a large glucocorticoid

response to an acute challenge (restraint) during a developmental window of intense social

interaction with both siblings and parents. While this is consistent with the hypothesis that

glucocorticoids are important for species with intra-brood competition, the ability of young

chicks to mount an adult-like HPA response to restraint has been found in a number of other

seabird species that have one-egg clutches, including three species of petrels (Adams et al., 2008;

Angelier et al., 2022; Sagar et al., 2019), three species of alcids (Benowitz-Fredericks et al.,

2015; Kitaysky et al., 2005; Will et al., 2015), and a congener of the black-legged kittiwakes

studied here: the red-legged kittiwake (Kitaysky et al., 2001; though red-legged kittiwakes

historically had larger clutch sizes). Thus, sibling competition cannot be the sole selective

pressure on the development of HPA axis activity. A phylogenetically controlled meta-analysis

characterizing the ontogeny of HPA activity across species with varied ecologies and life-

histories would be valuable for understanding the pressures and constraints on HPA axis

development in birds. However, for species like kittiwakes with very early activation of the axis,

the next steps are to understand why young chicks are capable of such robust responses to acute

challenges by characterizing the rapid metabolic and behavioral effects of elevated

corticosterone in chicks.

Acknowledgements

We are grateful for help with the following: establishment, maintenance, and long-term

funding/logistics for the tower colony: S. Hatch and M. Hatch; field logistics and assistance - F.

Tremblay and E.C. Kelly. Field crews: 2021 - K. Bacile, A. Colligan, J. Dunoyer, E. Fernandes-

McDade, J. Morales Valenzuela, S. Morrow. 2022 - G. Dennis, H. Gee, L. Jackson, D. Jadhon,

L. Marcouillier, S. Solasz, A. Turmaine; impeccable hormone assays - E. Kitaiskaia; genetic

sexing - A. Jackson, A. Le, S. Lin, and S. Walsh. We also thank two anonymous reviewers for

their thoughtful feedback. Financial support came from a Dean's Fellowship from Bucknell

University (ZMBF), a Gates Millennium Scholarship and an American Indigenous Business

Leaders internship (SNP), Gulf Watch Alaska and Alaska Ocean Observing System

(SW/Institute for Seabird Research and Conservation), and the Institute of Arctic Biology,

University of Alaska Fairbanks (ASK and AW). All research was conducted under permits from

the U.S. Fish and Wildlife Service (#MB33779D-1), Alaska Department of Fish and Game

(#21-089; #22-080) and with approval from McGill University's Animal Use Committee.

Figure captions

Figure 1. Experimental design. Timeline for blood sampling and topical hormone treatments of
5 day old black-legged kittiwake A-chicks.

Figure 2. Body condition and baseline corticosterone. Body condition of 5 day old first-
hatched kittiwake chicks was negatively related to baseline corticosterone levels, but this
relationship was eliminated in nests where parents were provided with supplemental food
("supplemented"). Solid line = significant, dashed = non-significant.

Figure 3. Baseline and restraint-induced corticosterone. Baseline corticosterone of 5-day old
kittiwake chicks was negatively correlated with corticosterone levels after 15 min bag restraint,
but only in chicks from nests where parents received supplemental feeding ("supplemented").
Chicks from supplemented nests had lower baseline corticosterone to begin with (see figure
4A). Solid line = significant, dashed = non-significant.

Figure 4. Hormone treatments and repeated blood samples. Circulating corticosterone levels
in 5-day old kittiwake chicks by parental feeding status and chick treatment. A) Bleed 1 is an
undisturbed baseline bleed for the control group (< 3 min) and 15 min after topical treatments for
the DMSO (DMSO gel) and CORT (corticosterone dissolved in DMSO gel) groups. B) Bleed 2
is after 15 min of continuous bag restraint for the control group, and 15 min after the first bleed
for the DMSO and CORT groups (treated chicks were returned to nests between bleed 1 and
bleed 2). C) Change in corticosterone levels between the first and second bleed. All statistical
analyses were performed using log-transformed data, but raw data are shown for panels A and B.

Figure 5. Time course for topical corticosterone. Corticosterone levels in 5-day old chicks
from nests without parental food supplementation (non-supplemented) at undisturbed baseline ($<$
3 min), 15 and 60 minutes after application of topical gel treatments. No chick was bled more
than once. Statistical analyses were performed using log-transformed data but raw data are
shown here.

FIGURE 1

Experiment 1 (2021 & 2022; supplemented & non-supplemented)

Experiment 2 (2021; supplemented & non-supplemented)

Experiment 3 (2022; non-supplemented only)

710   **FIGURE 2**

711 **FIGURE 3**

712 **FIGURE 4**

A. Bleed 1

B. Bleed 2

C. Change in Corticosterone (Bleed 2 – Bleed 1)

714   **FIGURE 5**

References:

Adams, N.J., Cockrem, J.F., Candy, E.J., Taylor, G.A., 2008. Non-precocial grey-faced petrel chicks (*Pterodroma macroptera gouldi*) show no age-related variation in corticosterone responses to capture and handling. Gen. Comp. Endocrinol. 157, 86–90. https://doi.org/10.1016/j.ygcen.2008.02.007

Ahmed, A.A., Ma, W., Ni, Y., Zhou, Q., Zhao, R., 2014. Embryonic exposure to corticosterone modifies aggressive behavior through alterations of the hypothalamic pituitary adrenal axis and the serotonergic system in the chicken. Horm. Behav. 65, 97–105. https://doi.org/10.1016/j.yhbeh.2013.12.002

Akana, S.F., Dallman, M.F., Bradbury, M.J., Scribner, K.A., Strack, A.M., Walker, C.D., 1992. Feedback and facilitation in the adrenocortical system: unmasking facilitation by partial inhibition of the glucocorticoid response to prior stress. Endocrinology 131, 57–68. https://doi.org/10.1210/endo.131.1.1319329

Angelier, F., Wingfield, J.C., Barbraud, C., Parenteau, C., Trouvé, C., Chastel, O., 2022. Ontogeny and individual heterogeneity of the corticosterone stress response in a wild altricial seabird, the snow petrel (*Pagodroma nivea*). Front. Ecol. Evol. 10, 902202. https://doi.org/10.3389/fevo.2022.902202

Béziers, P., San-Jose, L.M., Almasi, B., Jenni, L., Roulin, A., 2019. Baseline and stress-induced corticosterone levels are heritable and genetically correlated in a barn owl population. Heredity 123, 337–348. https://doi.org/10.1038/s41437-019-0203-5

Bebus, S.E., Jones, B.C., Anderson, R.C., 2020. Development of the corticosterone stress response differs among passerine species. Gen. Comp. Endocrinol. 291, 113417. https://doi.org/10.1016/j.ygcen.2020.113417

Benowitz-Fredericks, Z.M., Kitaysky, A.S., Welcker, J., Hatch, S.A., 2013. Effects of Food Availability on Yolk Androgen Deposition in the Black-Legged Kittiwake (*Rissa tridactyla*), a Seabird with Facultative Brood Reduction. PLOS ONE 8, e62949. https://doi.org/10.1371/journal.pone.0062949

Benowitz-Fredericks, Z.M., Schultner, J., Kitaysky, A.S., 2015. Effects of prenatal environment on phenotype are revealed by postnatal challenges: Embryonic hormone exposure, adrenocortical function, and food in seabird chicks. Physiol. Biochem. Zool. 88, 607–623. https://doi.org/10.1086/683259

Ben-Shachar, M., Lüdecke, D., Makowski, D., 2020. effectsize: Estimation of Effect Size Indices and Standardized Parameters. J. Open Source Softw. 5, 2815. https://doi.org/10.21105/joss.02815

Blas, J., Baos, R., Bortolotti, G.R., Marchant, T.A., Hiraldo, F., 2006. Age-related variation in the adrenocortical response to stress in nestling white storks (*Ciconia cico*nia) supports the developmental hypothesis. Gen. Comp. Endocrinol. 148, 172–180. https://doi.org/10.1016/j.ygcen.2006.02.011

Blas, J., Bortolotti, G.R., Tella, J.L., Baos, R., Marchant, T.A., 2007. Stress response during development predicts fitness in a wild, long lived vertebrate. Proc. Natl. Acad. Sci. U.S.A. 104, 8880–8884. https://doi.org/10.1073/pnas.0700232104

Bókony, V., Lendvai, Á.Z., Liker, A., Angelier, F., Wingfield, J.C., Chastel, O., 2009. Stress response and the value of reproduction: Are birds prudent parents? Am. Nat. 173, 589–598. https://doi.org/10.1086/597610

Braasch, A., Becker, P.H., Groothuis, T.G.G., 2014. Response of testosterone and corticosterone plasma levels to the challenge of sibling competition: A study in common terns. Gen. Comp. Endocrinol. 204, 95–103. https://doi.org/10.1016/j.ygcen.2014.05.007

Braun, B.M., Hunt, G.L., 1983. Brood reduction in black-legged kittiwakes. The Auk 100, 469–476. https://doi.org/10.1093/auk/100.2.469

Breuner, C.W., Berk, S.A., 2019. Using the van Noordwijk and de Jong Resource Framework to Evaluate Glucocorticoid-Fitness Hypotheses. Integr. Comp. Biol. 59, 243–250. https://doi.org/10.1093/icb/icz088

Brewer, J.H., O'Reilly, K.M., Buck, C.L., 2010. Effect of nestling status and brood size on concentration of corticosterone of free-living kittiwake chicks. Gen. Comp. Endocrinol. 166, 19–24. https://doi.org/10.1016/j.ygcen.2009.08.005

Brewer, J.H., O'Reilly, K.M., Kildaw, S.D., Buck, C.L., 2008. Interannual variation in the adrenal responsiveness of black-legged kittiwake chicks (*Rissa tridactyla*). Gen. Comp. Endocrinol. 156, 361–368. https://doi.org/10.1016/j.ygcen.2008.01.010

Burnham, K.P., Anderson, D.R., 2004. Multimodel Inference: Understanding AIC and BIC in Model Selection. Sociol. Methods Res. 33, 261–304. https://doi.org/10.1177/0049124104268644

Busch, D.S., Sperry, T.S., Wingfield, J.C., Boyd, E.H., 2008. Effects of repeated, short-term, corticosterone administration on the hypothalamo–pituitary–adrenal axis of the white-crowned sparrow (*Zonotrichia leucophrys gambelii*). Gen. Comp. Endocrinol. 158, 211–223. https://doi.org/10.1016/j.ygcen.2008.06.004

Crino, O.L., Breuner, C.W., 2015. Developmental stress: evidence for positive phenotypic and fitness effects in birds. J. Ornithol. 156, 389–398. https://doi.org/10.1007/s10336-015-1236-z

Crino, O.L., Driscoll, S.C., Brandl, H.B., Buchanan, K.L., Griffith, S.C., 2020. Under the weather: Corticosterone levels in wild nestlings are associated with ambient temperature and wind. Gen. Comp. Endocrinol. 285, 113247. https://doi.org/10.1016/j.ygcen.2019.113247

Dallman, M.F., Akana, S.F., Scribner, K.A., Bradbury, M.J., Walker, C. D., Strack, A.M., Cascio, C.S., 1992. Stress, Feedback and Facilitation in the Hypothalamo Pituitary Adrenal Axis. J. Neuroendocrinol. 4, 517–526. https://doi.org/10.1111/j.1365-2826.1992.tb00200.x

Dallman, M.F., Akana, S.F., Strack, A.M., Scribner, K.S., Pecoraro, N., La Fleur, S.E., Houshyar, H., Gomez, F., 2004. Chronic Stress Induced Effects of Corticosterone on Brain: Direct and Indirect. Ann. N. Y. Acad. Sci. 1018, 141–150. https://doi.org/10.1196/annals.1296.017

De Bruijn, R., Romero, L.M., 2018. The role of glucocorticoids in the vertebrate response to weather. Gen. Comp. Endocrinol. 269, 11–32. https://doi.org/10.1016/j.ygcen.2018.07.007

Dickins, D.W., 2021. "Punishing" pecks and "siblicidal" pecks in kittiwake (*tridactyla*) chicks. Learn. Motiv. 73, 101695. https://doi.org/10.1016/j.lmot.2020.101695

Dreiss, A.N., Gaime, F., Delarbre, A., Moroni, L., Lenarth, M., Roulin, A., 2016. Vocal communication regulates sibling competition over food stock. Behav. Ecol. Sociobiol. 70, 927–937. https://doi.org/10.1007/s00265-016-2114-2

Dudek, B.M., Henderson, M.T., Hudon, S.F., Hayden, E.J., Heath, J.A., 2021. Haematophagous ectoparasites lower survival of and have detrimental physiological effects on golden eagle nestlings. Conserv. Physiol. 9, coab060. https://doi.org/10.1093/conphys/coab060

Engqvist, L., Reinhold, K., 2016. Adaptive trans☐generational phenotypic plasticity and the
lack of an experimental control in reciprocal match/mismatch experiments. Methods Ecol.
Evol. 7, 1482–1488. https://doi.org/10.1111/2041-210X.12618
Fanson, K.V., Németh, Z., Ramenofsky, M., Wingfield, J.C., Buchanan, K.L., 2017.
Inter☐laboratory variation in corticosterone measurement: Implications for comparative
ecological and evolutionary studies. Methods Ecol Evol 8, 1745–1754.
https://doi.org/10.1111/2041-210X.12821
Fiske, J.A., Gannon, D., Newman, A.E.M., 2013. Effects of repeated investigator handling of
Leach's Storm-Petrel chicks on growth rates and the acute stress response: Repeated
Handling and Petrel Chick Development. J. Field Ornithol. 84, 425–432.
https://doi.org/10.1111/jofo.12041
Gill, V.A., Hatch, S.A., 2002. Components of productivity in black-legged kittiwakes *Rissa*
tridactyla☐: response to supplemental feeding. J. Avian Biol. 33, 113–126.
https://doi.org/10.1034/j.1600-048X.2002.330201.x
Gill, V.A., Hatch, S.A., Lanctot, R.B., 2002. Sensitivity of breeding parameters to food supply
in Black-legged kittiwakes *Rissa tridactyla*: Sensitivity of breeding kittiwakes to food. Ibis
144, 268–283. https://doi.org/10.1046/j.1474-919X.2002.00043.x
Groothuis, T.G., Kumar, N., Hsu, B.-Y., 2020. Explaining discrepancies in the study of
maternal effects: the role of context and embryo. Curr. Opin. Behav. Sci. 36, 185–192.
https://doi.org/10.1016/j.cobeha.2020.10.006
Hatch, S.A., Robertson, G.J., Baird, P.H., 2020. Black-legged Kittiwake (*Rissa tridactyla*), in:
Billerman, S.M., Keeney, B.K., Rodewald, P.G., Schulenberg, T.S. (Eds.), Birds of the
World. Cornell Lab of Ornithology. https://doi.org/10.2173/bow.bklkit.01
Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous Inference in General Parametric
Models. Biom. J. 50, 346–363. https://doi.org/10.1002/bimj.200810425
Jenkins, S.A., Porter, T.E., 2004. Ontogeny of the hypothalamo–pituitary–adrenocortical axis in
the chicken embryo: a review. Domest. Anim. Endocrinol. 26, 267–275.
https://doi.org/10.1016/j.domaniend.2004.01.001
Jenkins, B.R., Vitousek, M.N., Hubbard, J.K., Safran, R.J., 2014. An experimental analysis of
the heritability of variation in glucocorticoid concentrations in a wild avian population. Proc.
R. Soc. B. 281, 20141302. https://doi.org/10.1098/rspb.2014.1302
Jimeno, B., Verhulst, S., 2023. Meta-analysis reveals glucocorticoid levels reflect variation in
metabolic rate, not 'stress.' eLife 12, RP88205. https://doi.org/10.7554/eLife.88205.3
Jones, B.C., Nguyen, L.T., DuVal, E.H., 2021. Testing the developmental hypothesis of the
HPA axis in a tropical passerine: Dampened corticosterone response and faster negative
feedback in nestling lance-tailed manakins (*Chiroxiphia lanceolata*). Gen. Comp.
Endocrinol. 300, 113639. https://doi.org/10.1016/j.ygcen.2020.113639
Kahane-Rapport, S.R., Whelan, S., Ammendolia, J., Hatch, S.A., Elliott, K.H., Jacobs, S., 2022.
Food supply and individual quality influence seabird energy expenditure and reproductive
success. Oecologia 199, 367–376. https://doi.org/10.1007/s00442-022-05191-y
Kitaysky, A., Kitaiskaia, E., Wingfield, J., Piatt, J., 2001. Dietary restriction causes chronic
elevation of corticosterone and enhances stress response in red-legged kittiwake chicks. J.
Comp. Physiol. [B] 171, 701–709. https://doi.org/10.1007/s003600100230
Kitaysky, A.S., 2001. Corticosterone facilitates begging and affects resource allocation in the
black-legged kittiwake. Behav. Ecol. 12, 619–625. https://doi.org/10.1093/beheco/12.5.619

Kitaysky, A.S., Kitaiskaia, E.V., Piatt, J.F., Wingfield, J.C., 2003. Benefits and costs of
increased levels of corticosterone in seabird chicks. Horm. Behav. 43, 140–149.
https://doi.org/10.1016/S0018-506X(02)00030-2

Kitaysky, A.S., Romano, M.D., Piatt, J.F., Wingfield, J.C., Kikuchi, M., 2005. The
adrenocortical response of tufted puffin chicks to nutritional deficits. Hormones and
Behavior 47, 606–619. https://doi.org/10.1016/j.yhbeh.2005.01.005
Kitaysky, A., Piatt, J., Wingfield, J., 2007. Stress hormones link food availability and
population processes in seabirds. Mar. Ecol. Prog. Ser. 352, 245–258.
https://doi.org/10.3354/meps07074
Kitaysky, A.S., Piatt, J.F., Hatch, S.A., Kitaiskaia, E.V., Benowitz-Fredericks, Z.M., Shultz,
M.T., Wingfield, J.C., 2010. Food availability and population processes: severity of
nutritional stress during reproduction predicts survival of long-lived seabirds: Nutritional
stress and population processes. Funct. Ecol. 24, 625–637. https://doi.org/10.1111/j.1365-
2435.2009.01679.x
Kitaysky, A.S., Piatt, J.F., Wingfield, J.C., Romano, M., 1999. The adrenocortical stress-
response of Black-legged Kittiwake chicks in relation to dietary restrictions. J. Comp.
Physiol. [B] 169, 303–310. https://doi.org/10.1007/s003600050225
Kitaysky, A.S., Romano, M.D., Piatt, J.F., Wingfield, J.C., Kikuchi, M., 2005. The
adrenocortical response of tufted puffin chicks to nutritional deficits. Horm. Behav. 47, 606–
619. https://doi.org/10.1016/j.yhbeh.2005.01.005
Kumar, A., Darreh-Shori, T., 2017. DMSO: A mixed-competitive inhibitor of human
acetylcholinesterase. ACS Chem. Neurosci. 8, 2618–2625.
https://doi.org/10.1021/acschemneuro.7b00344
Kuznetsova, A, Brockhoff, PB, Christensen, RHB. 2017. "lmerTest Package: Tests in Linear
Mixed Effects Models." J. Stat. Software, **82**(13), 1–26. doi:10.18637/jss.v082.i13.
Lattin, C.R., Kelly, T.R., 2020. Glucocorticoid negative feedback as a potential mediator of
trade-offs between reproduction and survival. Gen. Comp. Endocrinol. 286, 113301.
https://doi.org/10.1016/j.ygcen.2019.113301
Leclaire, S., Bourret, V., Wagner, R.H., Hatch, S.A., Helfenstein, F., Chastel, O., Danchin, É.,
2011. Behavioral and physiological responses to male handicap in chick-rearing black-legged
kittiwakes. Behav. Ecol. 22, 1156–1165. https://doi.org/10.1093/beheco/arr149
Loiseau, C., Sorci, G., Dano, S., Chastel, O., 2008. Effects of experimental increase of
corticosterone levels on begging behavior, immunity and parental provisioning rate in house
sparrows. Gen. Comp. Endocrinol. 155, 101–108.
https://doi.org/10.1016/j.ygcen.2007.03.004
Love, O., Williams, T., 2008. Plasticity in the adrenocortical response of a free-living
vertebrate: The role of pre- and post-natal developmental stress. Horm. Behav. 54, 496–505.
https://doi.org/10.1016/j.yhbeh.2008.01.006
Lynn, S.E., Kern, M.D., 2018. Interactions of body temperature and nutritional status on
hypothalamo-pituitary-adrenal axis activity in pre-thermoregulatory eastern bluebird chicks
(*Sialia sialis*). Gen. Comp. Endocrinol. 267, 82–89.
https://doi.org/10.1016/j.ygcen.2018.06.003

Lynn, S.E., Kern, M.D., 2014. Environmentally relevant bouts of cooling stimulate
corticosterone secretion in free-living eastern bluebird (*Sialia sialis*) nestlings: Potential links
between maternal behavior and corticosterone exposure in offspring. Gen. Comp.
Endocrinol. 196, 1–7. https://doi.org/10.1016/j.ygcen.2013.11.011
Lynn, S.E., Kern, M.D., Fitzgerald, K., Will, A., Kitaysky, A., 2022. Cooling increases
corticosterone deposition in feathers of eastern bluebird chicks. Gen. Comp. Endocrinol. 320,
114001. https://doi.org/10.1016/j.ygcen.2022.114001
Marasco, V., Herzyk, P., Robinson, J., Spencer, K.A., 2016. Pre￿ and post￿natal stress
programming: Developmental exposure to glucocorticoids causes long￿term brain￿region
specific changes to transcriptome in the precocial Japanese quail. J. Neuroendocrinol. 28,
jne.12387. https://doi.org/10.1111/jne.12387
Mazerolle, M.J. 2023. *AICcmodavg: Model selection and multimodel inference based on*
(Q)AIC(c). R package version 2.3.2, https://cran.r-project.org/package=AICcmodavg.
Merkling, T., Chastel, O., Blanchard, P., Trouvé, C., Hatch, S.A., Danchin, E., 2014.
Physiological and fitness correlates of experimentally altered hatching asynchrony magnitude
in chicks of a wild seabird. Gen. Comp. Endocrinol. 198, 32–38.
https://doi.org/10.1016/j.ygcen.2013.12.011
Merkling, T., Leclaire, S., Danchin, E., Lhuillier, E., Wagner, R.H., White, J., Hatch, S.A.,
Blanchard, P., 2012. Food availability and offspring sex in a monogamous seabird: insights
from an experimental approach. Behav. Ecol. 23, 751–758.
https://doi.org/10.1093/beheco/ars023
Morandini, V., Ferrer, M., 2015. Sibling aggression and brood reduction: a review. Ethol. Ecol.
Evol. 27, 2–16. https://doi.org/10.1080/03949370.2014.880161
Noy, Y., Sklan, D., 2001. Yolk and exogenous feed utilization in the posthatch chick. Poult. Sci.
80, 1490–1495. https://doi.org/10.1093/ps/80.10.1490
Osorno, J.L., Drummond, H., 2003. Is obligate siblicidal aggression food sensitive? Behavioral
Ecology and Sociobiology 54, 547–554. https://doi.org/10.1007/s00265-003-0667-3
Patil, M., 2013. Pharmacology and clinical use of dimethyl sulfoxide (DMSO): A review. Int. J.
Mol. Vet. Res. https://doi.org/10.5376/ijmvr.2013.03.0006
Pegan, T.M., Winkler, D.W., Haussmann, M.F., Vitousek, M.N., 2019. Brief increases in
corticosterone affect morphology, stress responses, and telomere length but not post fledging
movements in a wild songbird. Physiol. Biochem. Zool. 92, 274–285.
https://doi.org/10.1086/702827
Pinheiro J, Bates D, R Core Team (2023). nlme: Linear and Nonlinear Mixed Effects Models. R
package version 3.1-164, https://CRAN.R-project.org/package=nlme
Quillfeldt, P., Masello, J.F., Möstl, E., 2004. Blood chemistry in relation to nutrition and
ectoparasite load in Wilson's storm-petrels *Oceanites oceanicus*. Polar Biol. 27, 168–176.
https://doi.org/10.1007/s00300-003-0572-2
R Core Team (2022). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/
Rogers, C.M., 2015. Testing optimal body mass theory: Evidence for cost of fat in wintering
birds. Ecosphere 6, 1–12. https://doi.org/10.1890/ES14-00317.1
Romero, L.M., 2004. Physiological stress in ecology: lessons from biomedical research. Trends
Ecol. Evol. 19, 249–255. https://doi.org/10.1016/j.tree.2004.03.008

Romero, L.M., Dickens, M.J., Cyr, N.E., 2009. The reactive scope model — A new model
integrating homeostasis, allostasis, and stress. Horm. Behav. 55, 375–389.
https://doi.org/10.1016/j.yhbeh.2008.12.009
Sagar, R.L., Cockrem, J., Rayner, M.J., Stanley, M.C., Welch, J., Dunphy, B.J., 2019. Regular
handling reduces corticosterone stress responses to handling but not condition of semi-
precocial mottled petrel (*Pterodroma inexpectata*) chicks. Gen. Comp. Endocrinol. 272, 1–8.
https://doi.org/10.1016/j.ygcen.2018.11.004
Santos, N.C., Figueira-Coelho, J., Martins-Silva, J., Saldanha, C., 2003. Multidisciplinary
utilization of dimethyl sulfoxide: pharmacological, cellular, and molecular aspects. Biochem.
Pharmacol. 65, 1035–1041. https://doi.org/10.1016/S0006-2952(03)00002-9
Sapolsky, R.M., Meaney, M.J., 1986. Maturation of the adrenocortical stress response:
Neuroendocrine control mechanisms and the stress hyporesponsive period. Brain Res. Rev.
11, 65–76. https://doi.org/10.1016/0165-0173(86)90010-X
Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence stress
responses? Integrating permissive, suppressive, stimulatory, and preparative actions. Endocr.
Rev. 21, 55–89. https://doi.org/10.1210/edrv.21.1.0389
Schoech, S.J., Rensel, M.A., Heiss, R.S., 2011. Short- and long-term effects of developmental
corticosterone exposure on avian physiology, behavioral phenotype, cognition, and fitness: A
review. Curr. Zool. 57, 514–530. https://doi.org/10.1093/czoolo/57.4.514
Schultner, J., Kitaysky, A.S., Gabrielsen, G.W., Hatch, S.A., Bech, C., 2013. Differential
reproductive responses to stress reveal the role of life-history strategies within a species.
Proc. R. Soc. B Biol. Sci. 280, 20132090–20132090. https://doi.org/10.1098/rspb.2013.2090
Schultner, J, Kitaysky, A.S., Welcker, J., Hatch, S., 2013. Fat or lean: adjustment of
endogenous energy stores to predictable and unpredictable changes in allostatic load. Funct.
Ecol. 27, 45–55. https://doi.org/10.1111/j.1365-2435.2012.02058.x
Schwabl, H., Partecke, J., 2020. Developmental programming of the adrenocortical stress
response by yolk testosterone depends on sex and life history stage. J. Exp. Biol. jeb.220152.
https://doi.org/10.1242/jeb.220152
Shultz, M., Piatt, J., Harding, A., Kettle, A., Van Pelt, T., 2009. Timing of breeding and
reproductive performance in murres and kittiwakes reflect mismatched seasonal prey
dynamics. Mar. Ecol. Prog. Ser. 393, 247–258. https://doi.org/10.3354/meps08136
Sims, C.G., Holberton, R.L., 2000. Development of the Corticosterone Stress Response in
Young Northern Mockingbirds (*Mimus polyglottos*). Gen. Comp. Endocrinol. 119, 193–201.
https://doi.org/10.1006/gcen.2000.7506
Smiseth, P.T., Scott, M.P., Andrews, C., 2011. Hormonal regulation of offspring begging and
mediation of parent–offspring conflict. Anim. Behav. 81, 507–517.
https://doi.org/10.1016/j.anbehav.2010.11.029
Sockman, K.W., Schwabl, H., 2001. Plasma corticosterone in nestling American kestrels:
Effects of age, handling stress, yolk androgens, and body condition. General and
Comparative Endocrinology 122, 205–212. https://doi.org/10.1006/gcen.2001.7626
Sorenson, G.H., Dey, C.J., Madliger, C.L., Love, O.P., 2017. Effectiveness of baseline
corticosterone as a monitoring tool for fitness: a meta-analysis in seabirds. Oecologia 183,
353–365. https://doi.org/10.1007/s00442-016-3774-3
Taff, C.C., Zimmer, C., Vitousek, M.N., 2018. Efficacy of negative feedback in the HPA axis
predicts recovery from acute challenges. Biol. Lett. 14, 20180131.
https://doi.org/10.1098/rsbl.2018.0131

Tilgar, V., Saag, P., Moks, K., 2009. Development of stress response in nestling pied flycatchers. J Comp Physiol A 195, 799–803. https://doi.org/10.1007/s00359-009-0452-5

Torres☐Medina, F., Cabezas, S., Marchant, T.A., Blas, J., 2019. Dexamethasone treatment supports age☐related maturation of the stress response in altricial nestling birds. J. Avian Biol. 50, jav.02091. https://doi.org/10.1111/jav.02091

Vallarino, A., Wingfield, J.C., Drummond, H., 2006. Does extra corticosterone elicit increased begging and submissiveness in subordinate booby (*Sula nebouxii*) chicks? Gen. Comp. Endocrinol. 147, 297–303. https://doi.org/10.1016/j.ygcen.2006.01.016

Vincenzi, S., Hatch, S., Mangel, M., Kitaysky, A., 2013. Food availability affects onset of reproduction in a long-lived seabird. Proc. R. Soc. B Biol. Sci. 280, 20130554. https://doi.org/10.1098/rspb.2013.0554

Vincenzi, S., Mangel, M., 2013. Linking food availability, body growth and survival in the black-legged kittiwake Rissa tridactyla. Deep Sea Res. Part II Top. Stud. Oceanogr. 94, 192–200. https://doi.org/10.1016/j.dsr2.2013.03.029

Vitousek, M.N., Taff, C.C., Ardia, D.R., Stedman, J.M., Zimmer, C., Salzman, T.C., Winkler, D.W., 2018. The lingering impact of stress: brief acute glucocorticoid exposure has sustained, dose-dependent effects on reproduction. Proc. R. Soc. B Biol. Sci. 285, 20180722. https://doi.org/10.1098/rspb.2018.0722

Wada, H., 2008. Glucocorticoids: Mediators of vertebrate ontogenetic transitions. Gen. Comp. Endocrinol. 156, 441–453. https://doi.org/10.1016/j.ygcen.2008.02.004

Wada, H., Hahn, T.P., Breuner, C.W., 2007. Development of stress reactivity in white-crowned sparrow nestlings: Total corticosterone response increases with age, while free corticosterone response remains low. Gen. Comp. Endocrinol. 150, 405–413. https://doi.org/10.1016/j.ygcen.2006.10.002

Wada, H., Salvante, K.G., Wagner, E., Williams, T.D., Breuner, C.W., 2009. Ontogeny and Individual Variation in the Adrenocortical Response of Zebra Finch (*Taeniopygia guttata*) Nestlings. Physiol. Biochem. Zool. 82, 325–331. https://doi.org/10.1086/599320

Whelan, S., Hatch, S.A., Benowitz-Fredericks, Z.M., Parenteau, C., Chastel, O., Elliott, K.H., 2021. The effects of food supply on reproductive hormones and timing of reproduction in an income-breeding seabird. Horm. Behav. 127, 104874. https://doi.org/10.1016/j.yhbeh.2020.104874

White, J., Leclaire, S., Kriloff, M., Mulard, H., Hatch, S.A., Danchin, E., 2010. Sustained increase in food supplies reduces broodmate aggression in black-legged kittiwakes. Anim. Behav. 79, 1095–1100. https://doi.org/10.1016/j.anbehav.2010.02.003

Will, A., Watanuki, Y., Kikuchi, D.M., Sato, N., Ito, M., Callahan, M., Wynne-Edwards, K., Hatch, S., Elliott, K., Slater, L., Takahashi, A., Kitaysky, A., 2015. Feather corticosterone reveals stress associated with dietary changes in a breeding seabird. Ecol. Evol. 5, 4221–4232. https://doi.org/10.1002/ece3.1694

Young, R.C., Welcker, J., Barger, C.P., Hatch, S.A., Merkling, T., Kitaiskaia, E.V., Haussmann, M.F., Kitaysky, A.S., 2017. Effects of developmental conditions on growth, stress and telomeres in black-legged kittiwake chicks. Mol. Ecol. 26, 3572–3584. https://doi.org/10.1111/mec.14121