Full Title: Meiotic DSB repair DNA synthesis tracts in *Arabidopsis thaliana*

Short Title: SPO11-dependent meiotic DNA synthesis.

Authors: Miguel Hernández Sánchez-Rebato[1&], Veit Schubert[2], Charles I. White[1]*

Affiliations:

[1] Institut de Génétique, Reproduction et Développement, CNRS UMR 6293, INSERM U1103,

Université Clermont Auvergne, 28, place Henri Dunant,

63001 Clermont-Ferrand, France.

[2] Leibniz Institute of Plant Genetics and Crop Plant Research (IPK) Gatersleben, D-06466 Seeland,

Germany

[&]Current address: Centro de Biología Molecular Severo Ochoa,

Calle Nicolás Cabrera 1, 28049 Madrid, Spain

* Corresponding author.

E-mail: charles.white@cnrs.fr

Keywords: Meiosis, Recombination, DNA synthesis, DSB repair, SPO11

Abstract

We report here the successful labelling of meiotic prophase I DNA synthesis in the flowering plant, *Arabidopsis thaliana*. Incorporation of the thymidine analogue, EdU, enables visualisation of the footprints of recombinational repair of programmed meiotic DNA double-strand breaks (DSB), with ~400 discrete, SPO11-dependent, EdU-labelled chromosomal foci clearly visible at pachytene and later stages of meiosis. This number equates well with previous estimations of 200-300 DNA double-strand breaks per meiosis in Arabidopsis, confirming the power of this approach to detect the repair of most or all SPO11-dependent meiotic DSB repair recombination. The chromosomal distribution of these DNA-synthesis foci accords with that of early recombination markers and MLH1, which marks Class I crossover sites, colocalises with the EdU foci.

It is currently estimated that ~10 cross-overs (CO) and an equivalent number of non-cross-overs (NCO) occur in each Arabidopsis male meiosis. Thus, at least 90% of meiotic recombination events, and very probably more, have not previously been accessible for analysis. Visual examination of the patterns of the foci on the synapsed pachytene chromosomes corresponds well with expectations from the different mechanisms of meiotic recombination and notably, no evidence for long Break-Induced Replication DNA synthesis tracts was found. Labelling of meiotic prophase I, SPO11-dependent DNA synthesis holds great promise for further understanding of the molecular mechanisms of meiotic recombination, at the heart of reproduction and evolution of eukaryotes.

Author Summary

Author Summary

Sexual reproduction involves the fusion of two cells, one from each parent. To maintain a stable

chromosome complement across generations, these specialized reproductive cells must be

produced through a specialized cell division called meiosis. Meiosis halves the chromosome

complement of gametes and recombines the parental genetic contributions in each gamete,

generating the genetic variation that drives evolution. The complex mechanisms of meiotic

recombination have been intensely studied for many years and we now know that it involves the

repair of programmed chromosomal breaks through recombination with intact template DNA

sequences on another chromatid. At the molecular level, this is known to involve new DNA

synthesis at the sites of repair/recombination and we report here the successful identification

and characterisation of this DNA neo-synthesis during meiosis in the flowering plant,

Arabidopsis. Both the characteristics and numbers of these DNA synthesis tracts accord with

expectations from theory and earlier studies. Potentially applicable to studies in many

organisms, this approach provides indelible footprints in the chromosomes and has the great

advantage of freeing researchers from dependence on indirect methods involving detection of

proteins involved in these dynamic processes.

58
59

Introduction

Meiosis is the specialised cell division that halves chromosome numbers during the

production of gametes, essential for the maintenance of stable ploidy across generations in

sexually-reproducing organisms. In contrast to mitosis, in which duplication of the genome is

followed by a single cellular division, meiosis involves one round of DNA replication followed by

two rounds of division. Separating sister chromatids to opposite poles, the second meiotic

division is, in broad terms, analogous to mitosis. Balanced chromosomal segregation during the

first division however requires that homologous chromosomes recognise each other, pair and

orient themselves on the spindle so that the two members of each pair move to opposite poles

during anaphase. This is ensured by the programmed induction of homologous recombination

through the induction and repair of DNA double-strand breaks (DSB) during the first meiotic

prophase.

This programmed, highly regulated process of DNA damage and repair is essential for

proper meiotic chromosome alignment and segregation and thus fertility in eukaryotes, with

some known exceptions, notably *Drosophila* and *C. elegans* (see review by [1]). Following the

programmed induction of DSB by the SPO11 complex, the cleaved DNA ends are resected to

produce 3'-ssDNA overhangs, onto which the recombinases DMC1 and RAD51 are loaded to

form presynaptic nucleoprotein filaments. These presynaptic nucleofilaments are the active

molecular species in the search for, and invasion of an intact homologous DNA template, the first

steps of DNA repair via homologous recombination. DNA synthesis then extends the invading 3'

end across the template and the joint-molecule intermediate structures are further processed to

yield the two main products of HR-mediated DSB repair: gene conversions associated with

reciprocal exchange of their flanking sequences (crossovers, CO), or not (non-crossovers, NCO).

DNA synthesis is thus inherent to DSB repair via homologous recombination. In the 1960s,

studies based on incorporation of H3-thymidine and C14-thymidine into newly synthesised DNA

gave evidence for meiotic prophase I DNA synthesis in the plants *Lilium longiflorum* and *Trillium*

erectum, the amphibian *Triturum viridescens*, mice and humans [2–6]. These early studies have

since been extended to show association of DNA synthesis with pachytene recombination

nodules in *Drosophila* female meiosis and with the synaptonemal complex in mouse

spermatocytes [7,8].

Labelling with 5-bromo-2′-deoxyuridine (BrdU) in budding yeast has been used to confirm

the presence of SPO11-dependent meiotic prophase DNA synthesis [9]. The authors

characterized both single-molecule BrdU tracts at a recombination hotspot and the timing of

their appearance in relation to other meiotic features. Observation of SPO11-dependent, BrdU

labelling pattern has also been described in *Tetrahymena* [10] but, to our knowledge, no further

data has been published in other organisms. In Arabidopsis, incorporation of BrdU or 5-ethynyl-

2′- deoxyuridine (EdU) during pre-meiotic S-phase has been used to establish the meiotic

timeline of meiosis in Arabidopsis pollen mother cells (PMC) [11–13]. There are thus far no

reports of labelling non-S-Phase DNA synthesis during Arabidopsis meiosis.

The induction of synchronous meiosis in populations of cells has been key to a number of

major advances in understanding of the molecular processes that make up the meiotic division

[14–18]. However, such direct approaches have not been available in other organisms and this

has resulted in significant gaps in understanding. A clear example of this is seen in the flowering

plant, *A. thaliana*. Estimations of numbers of SPO11-induced DSB in Arabidopsis pollen mother

cells (PMC, male meiosis) are ~200-300 per meiosis. Of these, ~10 yield inter-homologue CO and

another 10 have been shown to give inter-homologue NCO [19,20]. Thus, the repair outcome of

at least 90% of meiotic DSB in Arabidopsis is not known. The situation is strikingly different in

budding yeast with 66 NCO plus 90.5 CO estimated per meiosis, the sum of which is very close to

the estimated 140-170 meiotic DSB. Similarly in mouse, 273 NCO plus 27 CO sum close to the

estimated 200-400 meiotic DSB [21–24].

Detection of recombination-associated DNA synthesis tracts offers an attractive path to

understand the mechanisms underlying these "invisible" meiotic recombination events in

Arabidopsis and potentially in other organisms. We present here the successful labelling of

SPO11-dependent, meiotic prophase I DNA synthesis by EdU incorporation in Arabidopsis PMC.

Results

EdU labelling of DSB repair-associated DNA synthesis

In Arabidopsis, while each inflorescence contains flower buds bearing PMC at different pre-

meiotic and meiotic stages, the PMC of the anthers of a given flower bud are nearly synchronous

[25,26]. Along with the sequence of cytologically distinguishable meiotic stages in the buds of a

given inflorescence, in practice it is relatively straightforward to classify PMC subpopulations

within the meiotic timeline.

Timelines of Arabidopsis male meiosis concord that pre-meiotic G2 phase lasts 7-10 hours

and that pollen mother cells advance from the beginning of leptotene to the end of pachytene in

19 to 21 hours [11–13,27](summarized in Figure 1). Subsequent stages of meiosis are

considerably faster, with post-pachytene meiosis estimated to take from 2.7 to 9 hours. Given

that meiotic DSB formation begins at the onset of prophase I (early leptotene) and continues for

a few hours [28], any meiotic prophase I DSB-repair DNA synthesis should occur at least 7-10

hours after the end of pre-meiotic S-phase. We thus decided to incubate the inflorescences with

the nucleoside analogue for a 24-hour pulse to entirely cover the prophase I DSB repair-

associated DNA synthesis window. Given the relative synchrony of progression through meiosis

of the PMC from a given flower bud, the easily identifiable appearance of S-phase labelling and

the possibility to easily situate subpopulations of meiotic cells in the temporal meiotic sequence,

it seemed reasonable to expect that we would be able to exclude S-phase labelled meioses, and

thus focus our analyses on those labelled during Prophase I.

Notwithstanding the current absence of direct evidence of the timing of occurrence of

meiotic recombination intermediates in Arabidopsis, DAPI staining and immunolocalisation of

meiotic proteins implicated in early invasion intermediates (RAD51/DMC1), joint molecules

(MSH4, early HEI10) and resolution (MLH1, late HEI10) give a clear picture of the progression of

key events through meiotic prophase I. In pachytene, when synapsis is complete, mean numbers

of MLH1 foci correspond to the number of Class I crossovers (~90% of total CO), measured by

counting of chiasmata at metaphase I. In analogy to organisms for which recombination

intermediates can be tracked directly, it is reasonable to assume that most recombination events

are either resolved or close to resolution at pachytene in Arabidopsis.

As summarised in Figure 1 , inflorescences were thus incubated with EdU for 24 hours to

ensure coverage of the window of DSB repair-associated DNA synthesis through prophase I up to

the end of pachytene. Pachytene cells observed at the end of the 24-hour pulse would have

traversed prophase I up to that point in the presence of the analogue. As mentioned above,

nuclei with pre-meiotic S-phase labelling were easily identified and excluded from the analysis.

Non-replicative DNA synthesis tracts are present from pachytene until the end of meiosis

Early prophase I meiocytes, including cells in leptotene and zygotene, showed complete or

extensive EdU labelling of their chromosomes (Figure 2a). This replicative labelling confirms, as

expected, that they had passed through pre-meiotic S-Phase during the EdU pulse [11–13,29,30].

Furthermore, nuclei with partial replicative labelling of DAPI-dense, late replicating

heterochromatic regions were observed, as expected for cells that were in mid/late S-phase at

the beginning of the EdU pulse (second row of Figure 2a) [30–33].

In striking contrast, pachytene nuclei showed faint, discrete foci of a considerably smaller

size (Figure 2a Pachytene, see also Figure 4a and Figure S1). These foci are very faint through the

microscope eyepiece, but are clearly visible when the images are processed. This difference in

labelling intensity between S-phase and prophase EdU incorporation is not unexpected,

considering that the latter would correspond to DNA synthesis tracts of tens to a few thousands

of base pairs, while replicative DNA synthesis would range from megabase-long late-replicating

regions to the whole genome (4C ≈ 520 Mb in these prophase I nuclei) [34]. We note also the

presence of labelled mitochondria in these images, confirming their replication during pre-

meiotic G2 and/or meiotic prophase I cells. These labelled mitochondria, clearly visible from

pachytene on, are also present at earlier stages, but are not readily visible due to the very strong

fluorescence of the S-phase replicative nuclear labelling (Figure 2).

The EdU foci observed in pachytene cells appear well-distributed across the chromosomes,

but are less present in DAPI-dense, heterochromatic regions (Figure 2a Pachytene, Figure 4a).

Furthermore, given the tighter-packed DNA in these regions, the observed lesser density of EdU

foci is presumably an underestimation. This pattern of foci concords with genetic crossover

mapping, studies of meiotic DSB (SPO11-oligo sequencing) and immunofluorescence studies of

the distributions of early recombination proteins, all of which show meiotic recombination to

follow a relatively even distribution across the Arabidopsis genome, with the exception of

(heterochromatic) centromeric regions which have significantly lower densities of SPO11-oligos

and crossovers [34–36].

As summarised in Figure 1, based on published timelines [11–13,27] and our own

measurements, a meiocyte observed at the telophase II/tetrad stage would have been at

leptotene/zygotene at the beginning of the 24-hour EdU pulse. We had thus good reason to

expect that prophase I EdU foci would be observable in all meiotic stages up to the end of

meiosis. This was indeed the case, with EdU foci visible from pachytene to the end of meiosis

(telophase II/tetrad Figure 2). Meiotic chromosomes condense progressively as they traverse

prophase I, evolving from a mass of hardly-distinguishable thin fibres at early leptotene to thick

synapsed fibres in pachytene and to individualised dense bivalents with a more polygonal shape

at metaphase I. This condensation produces significant shortening of the chromosomes and

results in changes of the visual character of the EdU signals. While most EdU foci at pachytene

appear as discrete spots, they merge into larger patches at diplotene and this continues through

diakinesis, up to the greatest condensation at metaphase I (Figure 2, diplotene-metaphase I).

Furthermore, heterochromatic, centromeric regions become more easily identifiable when

bivalents individualize at diakinesis and the relatively low density of EdU signals in these regions

becomes clearly visible. This is even more pronounced at metaphase I, with clear visual

differentiation between the centromeric and pericentromeric regions of the two homologous

chromosomes of each bivalent. Finally, the EdU signals are observed throughout the second

meiotic division, changing their appearance with evolving chromosome condensation and shape

in a way similar to that seen during the first meiotic division (Figure 2, prophase II-tetrad).

Prophase I DNA synthesis is SPO11-dependent

Meiotic recombination is initiated by the induction of DNA DSB by the SPO11 complex at the

beginning of meiotic prophase I. To verify that the discrete non-replicative DNA synthesis tracts

observed at pachytene and subsequent stages originate from SPO11-dependent meiotic DSB

repair, we carried out this analysis in parallel in *spo11-1* mutant plants [37] (Figure 3).

Pachytene images were analysed with the IMARIS software package (Bitplane AG, Oxford

Instruments, UK) *Spots* function. EdU foci were selected based on diameter and mean intensity

in the EdU channel, and filtered to retain only those colocalising with chromosome fibres (mean

intensity cutoff in the DAPI channel). A high-cut filter of mean intensity in the EdU channel was

also applied to filter out the much-brighter organellar foci.

WT pachytenes show 409.5 ± 7.96 EdU foci per nucleus (mean ± SEM; n = 45 nuclei; Figure

2, Figure S1), with counts ranging from 323 to 510 (plus one outlier with 263). Ten of the 45

image stacks analysed were acquired via 3D-SIM, for comparison with the Zeiss LSM

800/Airyscan confocal image stacks and to confirm that we were not missing foci. WT cells

imaged with 3D-SIM had a mean 400.6 ± 14.74 EdU foci (n = 10) while those imaged with the

confocal microscope had a mean of 412.1 ± 9.38 EdU foci (n = 35). Hence, no significant

difference was observed between both imaging systems (unpaired t-test; p = 0.55) (Figure 4).

In *spo11-1* plants, 39.2 ± 2.83 EdU foci per cell (mean ± SEM; n = 35 cells; Figures 3, 4) were

detected in zygotene/pachytene nuclei. Thus, some EdU foci were detected in *spo11-1* meioses,

but in significantly lower numbers and, in general, considerably smaller and weaker than the foci

in the WT. A similar signature could be observed in *spo11-1* cells at subsequent meiotic stages

(Figure 3, diakinesis to Telophase II), with some EdU foci detected but in much lower numbers.

This ~10-fold reduction in mean numbers of foci with respect to the WT is highly significant

(unpaired t-test; p < 0.0001) and confirms the SPO11-dependence of the great majority of the

observed pachytene EdU foci. That the expected S-phase EdU labelling was observed in early

prophase I *spo11-1* PMC nuclei (Figure 3, early PI, mid PI, second row) and organellar EdU

labelling was comparable in *spo11-1* and WT meioses (Figures 2, 3), confirms that the EdU

labelling pulse worked as expected in both WT and *spo11-1* plants.

In interpreting these numbers it is important to note that although Arabidopsis *spo11*

meiosis does complete and produce (mostly inviable) pollen, chromosome pairing and synapsis

are very severely reduced in *spo11* meiosis and the zygotene/pachytene stages are only

identifiable by their characteristic chromosome condensation and their temporal positioning

[37]. Thus, *spo11-1* PMC at leptotene can be differentiated from the zygotene-like state by the

condensation of the chromosomes (thickness of the fibres) and the clustering of DAPI-dense

regions. Also, *spo11-1* zygotene/pachytene-like cells that do not show S-phase EdU labelling are

in a more advanced state that those that do show it, as they would have completed S-phase at

the beginning of the pulse (Figure 3a, mid PI, third row vs second row). Identification of later

meiotic stages in *spo11-1* meioses presents no difficulties (Figure 3).

Prophase I DNA synthesis patterns mirror meiotic recombination features

Meiotic recombination models predict different patterns of DNA synthesis tracts depending

on how recombination intermediates are formed and resolved. Non-crossover events generated

by either synthesis-dependent strand annealing (SDSA) or dHJ dissolution recombination, both

expected to produce DNA synthesis tracts in only one of the two implicated chromatids and thus

in only one of the homologous chromosomes in interhomologue recombination. On the other

hand, dHJ resolution resulting in either crossover or non-crossover products would generate

DNA synthesis tracts in both participating chromatids - and both chromosomes in

interhomologue events.

The expected patterns of resulting EdU foci are shown schematically in Figure 5a and

examples of segments of the synaptonemal complex (SC) from SIM pachytene micrographs are

shown in Figure 5b. The paired, homologous chromosomes are clearly distinguishable in these

images, permitting a first analysis of different patterns of EdU foci. No obvious bias is visible in

the disposition of Individual EdU foci in these images. Foci are seen at both the internal (facing

the homologous chromosome) and external faces of the paired chromosomes in the SC fibre, as

well as in the middle of the SC (Figure 5).

Differences can however be seen when EdU foci are analysed with respect to adjacent foci,

with three clear configurations of pairs of foci, examples of which are circled on the images

shown in Figure 5b:

Class (1) isolated individual foci located on one of the two homologues (one SC lateral

axis; Fig. 5b, white circle).

Class (2) "pairs" of foci on one of the two homologues (one SC lateral axis; Fig. 5b, green

circle).

Class (3) "pairs" of foci on the two homologues (both SC lateral axes; Fig. 5b, yellow

circle).

Further examples of these patterns can be found in the pachytene images of Figure S2.

Finally, we note the absence of the long labelling tracts expected from BIR recombination,

implying that such events are either rare or "short", or both, in Arabidopsis meiosis.

Prophase I DNA synthesis colocalize with class I crossovers

As the SPO11-dependent, meiotic prophase I DNA synthesis foci result from recombinational

repair of meiotic DSB, genetic crossovers should co-localize with a subset of these EdU foci. Class

I CO are detectable cytologically in Arabidopsis by the immunolocalization of proteins such as

MLH1.

Combining labelling of EdU tracts with immunolocalization of MLH1 (Figure 6) gave 7.86 ±

0.24 (mean ± SEM; n = 29) MLH1 foci per WT pachytene, as expected [38]. 95.6% (n=228) of all

MLH1 foci colocalized with an EdU focus and the remaining 4.4% were adjacent to one. Although

this conclusion must be tempered due to the high number of EdU foci in these images, Arabidopsis

Class I CO are thus physically associated with the prophase I DNA synthesis tracts at this level of

resolution.

Discussion

Identification of meiotic DNA repair-associated DNA synthesis tracts

We present here the labelling and characterisation of SPO11-dependent, meiotic prophase I

DNA synthesis tracts through incorporation of the thymidine analogue EdU in pollen mother cells

of Arabidopsis plants. By labelling DNA synthesis associated with the repair of meiotic DSB, this

approach holds great promise for the future understanding of recombination patterns and

mechanisms, notably for studies of NCO and sister-chromatid exchange (SCE) recombination. For

practical reasons, molecular and genetic studies of meiotic recombination have essentially been

focussed on events implicating the transfer of DNA sequence polymorphisms between the two

recombining DNA molecules. The necessary presence of such polymorphism has, for instance,

complicated significantly the analysis of the roles of meiotic SCE [39–46] and while EdU labelling

has recently been applied to analysis of meiotic sister-chromatid CO in *C. elegans* (Almanzar *et*

al, 2021; Toraason *et al*, 2021; Billmyre & Hughes, 2021), this approach does not appear adapted

to the study of NCO.

Detection of NCO events in Arabidopsis has been elusive. Mapping NCO genome-wide by

sequencing Arabidopsis tetrads has estimated NCO numbers to be of the same order as CO, ~10

events per meiosis, with gene conversion tracts ranging between a few tens to a few hundreds

of bp [19,20,50,51]. Notwithstanding this (surprisingly) low number of NCO, numbers of CO are

in agreement with chiasmata counting and immunolocalisation of CO markers. Thus, given the

200-300 DSB per meiosis in the these cells, the repair of at least 90% (and very possibly >98%

[52] of meiotic DSB in Arabidopsis is not accessible to analysis with current methods. In this

work, we present the development of meiotic prophase DNA synthesis labelling as a means to

fully characterise meiotic DSB-repair in Arabidopsis meiocytes.

Initial tests showed that incubating inflorescence stems for 24 hours in EdU solution

resulted in two clear sub-populations of meiocytes, distinguishable by their meiotic stage and

pattern of EdU labelling. The first consists of late G2 + early prophase I cells (leptotene,

zygotene) whose chromosomes show very strong EdU signals, either genome-wide or in large

patches. These patterns are consistent with those expected from pre-meiotic S-phase replicative

EdU incorporation:

I) This extensive EdU labelling is observed in meioses of both wild-type and *spo11* mutant

plants.

II) The extensive and intense fluorescent signals are similar to those observed in previous

reports of EdU/BrdU incorporation during pre-meiotic replication [11–13,29].

III) Extrapolating from published meiotic timelines and our own studies, PMC in pre-meiotic

S-phase at the beginning of the pulse are expected to advance at most to leptotene-

zygotene during the 24 hour EdU pulse (see Figure 1).

IV) The signal in partially labelled early prophase I cells corresponds principally to late-

replicating, DAPI-dense, heterochromatic regions, in accordance to expectations for cells in

late S-phase at the beginning of the EdU pulse.

The second population of EdU-labelled meiocytes includes cells at meiotic stages from

pachytene to the end of meiosis (tetrads), whose chromosomes show much fainter, discrete EdU

labelling. Clearly resolving into individual foci at the stages in which the chromosomes are less

condensed, these foci group into "patches" as chromosome condensation proceeds post-

prophase I. These foci have a number of characteristics which permit their identification with

confidence as meiotic DSB repair-associated DNA synthesis tracts:

I) These discrete EdU-label foci are SPO11-dependent, analogously to earlier observations in

yeast [9], confirming with high confidence that they result from EdU incorporation at DNA

synthesis tracts associated to meiotic DSB repair. Some EdU foci are observed in *spo11-1*

meioses, although at considerably lower numbers than in the wild type (39 versus 410 per

meiosis respectively). While some of these may come from mistaken assignation of organellar

DNA foci, we note that some crossing-over is detected in Arabidopsis *spo11* plants, RAD51 foci

are detected in Arabidopsis *spo11* meiosis and DMC1 foci in rice meioses [37,53–55].

Furthermore, given that both meiotic Arabidopsis SPO11 proteins are needed for normal meiotic

DSB induction, some leakage may occur in plants lacking one or the two SPO11 proteins [54,56–

59]. Alternatively, they may correspond to SPO11-independent breakage of mechanical,

chemical or enzymatic origin. If so, that they do produce DNA neo-synthesis tracts argues for

their repair by HR rather than End Joining.

II) The SPO11-dependent foci are seen at later meiotic stages than those with replicative

labelling and these nuclei had thus completed S-phase prior to the beginning of the pulse. Given

the known timing of meiosis in these cells, meiotic DSB repair DNA synthesis will have occurred

during the EdU pulse in these cells [9,13,14,38,60]. We note that while present in considerably

lower numbers than in the wild type (39 versus 410 per meiosis, respectively), there are some

EdU foci in *spo11-1* meioses.

III) The discrete EdU foci observed at pachytene and later stages are both much less intense

and smaller in size than the generalised labelling in leptotene-zygotene nuclei. This is expected

given the predicted size of DNA synthesis tracts associated to recombination events (few bases

to some kilobases of DNA), with respect to replicative labelling (megabases to the full genome).

IV) The numbers of EdU foci measured at pachytene concord with published estimations of

numbers of meiotic DSB and HR intermediates in Arabidopsis. A mean of 409.5 ± 7.96 (mean ±

SEM; n = 45 meioses) DNA repair-associated DNA synthesis foci were observed in WT

pachytenes. This number is a little higher than estimations of meiotic DSB numbers through

immunolocalisation of early recombination proteins in Arabidopsis (RAD51 in most cases).

Reported numbers of RAD51 foci per Arabidopsis meiosis range from 90 to 250 per meiosis, and

compare well with a recent report of 239 ± 30 (mean ± SD) prophase I SPO11-1 foci [56]. Given

the limitations of immunodetection and that the recruitment/release of these proteins at

DSB/recombination sites is dynamic, these numbers presumably under-estimate the true values.

Notably, a recent report using a STED super-resolution microscope found more than 1000 RAD51

foci per PMC meiosis [52]. Thus, the mean of 409.5 DNA repair-associated synthesis tracts per

meiosis reported here is probably closer to the true value than that of protein-immunodetection

approaches. It is of course necessary to take into account the fact that resolution of dHJ

recombination intermediates is expected to result in patches of neo-synthesized DNA on both

implicated chromatids, resulting in two foci from a single recombination event. Such events are

however estimated to represent ~5% or less of meiotic DSB repair events in Arabidopsis and so

are not expected to impact significantly the estimations of total events.

V) DAPI-dense heterochromatic, centromeric regions have a visibly lower density of SPO11-

dependent EdU foci in pachytene/diplotene and beyond (particularly noticeable at

diakinesis/MI). This concords with SPO11-oligo mapping of meiotic DSB patterns in Arabidopsis

PMC, which shows that centromeric regions have a much lower density of DSB than the rest of

the chromosome [34,36] .

VI) Combining labelling of EdU tracts with immunolocalization of MLH1 confirms that, as

expected, Class I CO are physically associated with the prophase I DSB repair DNA synthesis tracts

(Figure 6).

Meiotic DSB repair-associated DNA synthesis tracts mirror meiotic recombination features

As presented in Figure 7, different mechanisms of recombinational repair of meiotic DSB will

result in different patterns of DNA synthesis tracts associated to an HR event. The most obvious

difference is that DNA synthesis is expected to be found on only the recipient chromatid, which

carried the initiating DSB (Figure 7f), or on both the recipient and donor chromatids (Figure 7g,

h). SDSA and dHJ dissolution mechanisms would lea d to the former and dHJ resolution to the

latter. A special case is the BIR mechanism, which is expected to result in a long DNA synthesis

tract, potentially extending all the way to the end of the recipient chromatid (Figure 7c).

379   Visual inspection of the pachytene images clearly shows different configurations of the EdU

380   foci, which we have arbitrarily divided into 3 classes suggestive of the DNA synthesis patterns

381   expected at sites of recombination:

382   Class (1) isolated individual foci located on one of the two homologues (one lateral axis) (Fig.

383   5b, Fig. S2). Only the initially broken chromatid (recipient) will bear (EdU-substituted)

384   neosynthesized DNA tracts in repair via SDSA (Fig. 7a, b, f) and dHJ dissolution (Fig. 7a, d, e, f)

385   recombination.

386   Resolution of double Holiday junction (dHJ) intermediates (Fig7d) can give rise to gene

387   conversions, accompanied (Fig. 7g) or not (Fig. 7h) by a CO. In both cases, this can lead to the

388   presence of neo-synthesized DNA on both the donor and recipient chromatids. These events

389   would thus be expected to result in paired DNA synthesis tracts on the two participating

390   chromatids: either on one chromosome (Class (2) - one lateral axis) for sister chromatid events,

391   or on both homologues (Class (3) - both lateral axes).

392   Class (2) are thus "pairs" of foci on one of the two homologues (one SC lateral axis), which

393   could result from recombination between sister chromatids through the dHJ resolution pathway.

394   Class (3) are "pairs" of foci on the two homologues (both lateral axes), which could result

from recombination between homologues through the dHJ resolution pathway.

We find no evidence for long DNA synthesis tracts indicative of BIR recombination in our

images. Thus, either BIR replication tracts are not distinguishably longer than other DSB repair-

DNA synthesis, or the BIR mechanism is a minor (or non-) contributor to DSB repair in

Arabidopsis meiosis. In this context, we note that an earlier effort to detect mitotic BIR repair in

bleomycin-treated *Vicia faba* root tips described asymetric interstitial G2-phase EdU labelling of

interstitial tandem-repeat regions, although no extended labelling tracts out to the end of the

chromatid were observed [61].

Inference of the recombination pathways underlying the 3 classes of foci of course remains

speculative in the absence of molecular analyses of the EdU tracts themselves. FISH on extended

DNA fibres was used in an earlier study in yeast to determine the lengths and positioning of

BrdU-substituted DNA synthesis tracts with respect to DSB hotspots [9]. They observed two main

classes of tracts: tracts that extended from the DSB towards only one side of the DSB and tracts

that extended on both sides. These two classes correspond to the tracts expected in single DNA

molecules that took part in CO events in the first case and NCO events via SDSA in the second,

which were further validated studying their dynamics in recombination mutants. We have

previously carried out DNA combing plus fibre FISH in Arabidopsis [62], but the lack of strong

meiotic DSB hotspots with a precisely localised DSB site complicates the possible use of this

approach in Arabidopsis meiosis. Rather, we are actively developing an approach based on the

direct detection of EdU-substituted DNA tracts by Nanopore sequencing. It is hoped that this

approach will permit unambiguous characterisation of the principal meiotic recombination

products and underlying mechanisms in Arabidopsis, as well as of more complex events.

Materials and Methods

Plant material and growing conditions

Standard conditions were used for growth of wild-type *A. thaliana Col-0* and *spo11*-1-2 [37] plants. Seeds were sown on soil substrate (Klasmann-Deilmann GmbH TS3 FIN 416GF, Geeste, Germany), stratified for 2-4 days at 4°C and grown in climate chambers at 23°C and 60% relative humidity, under a daily cycle of 16 hours of light (110-140 µmol m^{-2} s^{-1}) and 8 hours of darkness.

Meiotic chromosome preparation by spreading

Meiosis was analysed following the chromosome spreading method of Ross et al. [63], with minor modifications. Inflorescences were collected in 3:1 ethanol:glacial acetic acid fixative solution and left overnight at room temperature. The fixative was then substituted with fresh fixative solution and the tubes were transferred to 4°C. This step was repeated the following 2-3 days. At this point, fixed inflorescences can be stored at 4°C for further use or prepared for microscopy.

A number of fixed inflorescences were selected, deposited in a glass dissection well with 3:1 fixative solution and dissected under a stereomicroscope into individual flower buds, discarding

open flowers and buds with pollen (yellow anthers). The flower buds were then washed 3 x 2

minutes in citrate buffer and the buffer replaced with 500 µl of digestion enzyme mix (0.3% w/v

cellulase from *Trichoderma* sp. (Sigma-Aldrich), 0.3% w/v pectolyase from *Aspergillus japonicus*

(Sigma) and 0.3% w/v cytohelicase from *Helix pomatia* (Duchefa Biochemie) in citrate buffer).

The flower buds were incubated in the digestion mix for 2 hours at 37°C and the digestion

stopped by adding a greater volume of ice-cold citrate buffer. The well was placed under the

stereomicroscope and a single flower bud with a small volume of liquid was transferred to a slide

with a Pasteur pipette. The flower bud was macerated with a needle, a 10µl drop of ice-cold 60%

glacial acetic acid was added and the slide incubated for one minute on a hot plate at 45°C while

gently moving the drop with a flat needle. After the incubation, an extra 10 µl of ice-cold 60%

glacial acetic acid were added, followed by 100 µl of 3:1 fixative - first circling and flooding the

drop and then mixing with the drop by vigorously pipetting the last microliters. The liquid was

drained by tilting the slide, which was then further washed with another 100 µl of 3:1 fixative,

drained and left to air dry at room temperature.

To visualize the chromosomes, the slides were stained in Vectashield antifade mounting

medium with 1.5 µg/ml DAPI (# H-1200, Vector Laboratories, Burlingame, CA, USA), by putting a

drop on a coverslip which was then placed on a slide and gently squashed by thumb pressure.

Spreads were analysed with a Zeiss AxioImager Z1 microscope with a 100×/1.40 Oil Plan-

Apochromat objective and the Zeiss ZEN2Blue software (Carl Zeiss GmbH)in the channel for blue

fluorescence (Zeiss #49 HE filter).

EdU labelling and cytological detection

Among the different thymidine analogues commonly used to label *in vivo* DNA synthesis, 5-

Ethynyl-2'-deoxyuridine (EdU) has a number of advantages and has been preferred by the

Arabidopsis community in recent studies labelling DNA synthesis [12,29,30,64–67]. EdU can be

detected in intact double-strand DNA and this antibody-free technique simplifies the protocol

and offers better reproducibility.

EdU (#Ab146186, Abcam, UK) working solutions were prepared at 10mM in 1×PBS buffer.

Arabidopsis stems with apical inflorescences were submerged in water to avoid the introduction

of air bubbles (submerging only the stem segment to be cut, without wetting the inflorescences)

and cut obliquely 4-5cm below the inflorescence. Care was taken to make the cleanest cut

possible in order to avoid crushing vascular tissues. Open flowers and big leaves (if present) were

removed to avoid crowding the tube and minimize the risk of exhausting the solution due to

excessive transpiration. The inflorescences were incubated in 800µl of 10mM EdU in a 2ml

eppendorf tube for 24 hours, under the same growing conditions as the plants. To avoid tight

packing, a maximum of four or five inflorescence stems were incubated in one tube. After 24

hours the inflorescences were removed from the EdU solution and fixed in 3:1 ethanol:glacial

acetic acid solution.

Chromosome spreads were prepared as previously described and visualised under the

microscope to identify those slides with meiocytes at the desired meiotic stages. The slides were

first incubated in 4T (4×SSC, 0.5% v/v Tween 20) in a Coplin jar until the coverslips detached,

followed by 30' without coverslips in fresh 4T. The Click-iT reaction mix (Invitrogen Click-iT plus

Alexa-fluor 647 kit, Thermo-Fisher Scientific) to label EdU with the fluorochrome was prepared

following the manufacturer's instructions. Before labelling, the slides were washed for 5 minutes

with 1×PBS. A 30µl drop of Click-iT reagent was placed on a 32x24mm piece of parafilm and the

inverted slides were placed on top of the mix on the parafilm. The slides were incubated with the

mix for 30 minutes at 37°C in a dark moist chamber, washed for 5 minutes with 1×PBS in a Coplin

jar in the dark and mounted in Vectashield + DAPI for microscopy.

Pictures were taken using a Zeiss LSM800 confocal microscope applying the Airyscan

module and selecting the presets for each specific dye. EdU foci were quantified semi-

automatically using the IMARIS software *Spots* tool (Bitplane AG, Oxford Instruments, UK),

adding intensity filters for both channels to specifically selected visually detectable EdU foci that

co-localised with the DAPI fluorescence of the chromosomes. To analyze the chromatin and EdU

signal ultrastructure, we applied super-resolution spatial structured illumination microscopy (3D-

SIM) using a 63×/1.40 Oil Plan-Apochromat objective of an Elyra PS.1 microscope system and the

ZENBlack software (Carl Zeiss GmbH) [68]. Maximum intensity projections were calculated from

3D-SIM image stacks. Zoom-in sections were presented as single slices to indicate the subnuclear

chromatin structures at the super-resolution level.

Detection of MLH1 and EdU

For the detection of MLH1 and neo-synthesized DNA tracts with incorporated EdU, the

protocol for the immunolocalization of MLH1 [69] was adapted to include the Click-iT EdU

labelling. Chromosome spreads were prepared as previously described and visualized under the

microscope to identify those slides with meiocytes at the desired stages. The slides were

incubated in a Coplin jar with 100% ethanol to detach the coverslip and washed with 1×PBST

(0.1% Triton X-100 in 1×PBS. Another glass staining jar was filled with tri-sodium citrate buffer

pH7 and heated to boiling point in a microwave. The slides were placed in the heated tri-sodium

citrate solution for 45 seconds, after which they were returned to room temperature 1×PBST.

504   The rabbit anti-MLH1 primary antibodies (kindly provided by Mathilde Grelon, INRAe

Versailles [38]) was prepared at 1/200 in 1×PBST + 1% BSA. A 45µl drop was placed on a

32x24mm piece of parafilm and the inverted slide place on this. The slides were incubated for 2

days at 4°C in a dark moist chamber, after which they were washed 3 x 5 minutes with 1×PBST.

The secondary antibodies (goat anti-rabbit Alexa fluor 488) was prepared at 1/100 1×in PBST +

1% BSA, and 45µl were used per slide. Slides were incubated 30 minutes at 37°C in a dark moist

chamber, after which they were washed 2 x 5 minutes in 1×PBST, followed by one wash with

1×PBS. The Click-iT reaction was then performed as described above and the slide was mounted

in Vectashield + DAPI mounting medium for microscopy. Counting of MLH1 foci and co-

localization with EdU foci was carried out by visual inspection of images taken using the Zeiss

LSM800 confocal microscope plus the Airyscan module.

Acknowledgements

We thank the members of the recombination group and especially Olivier Da Ines, Jérémy

Verbeke and Floriane Chéron for their help and suggestions. Mathilde Grelon kindly sent us the

anti-MLH1 antiserum.

This work was supported by a European Marie Sklodowska Curie Actions Innovative Training

Network grant (EU H2020-MSCA-ITN-2017-765212) to CIW, the Centre National de Recherche

Scientifique (CNRS), the Institut National de la Santé et de la Recherche Médicale (Inserm) and

the Université Clermont Auvergne. The funders had no role in study design, data collection and

analysis, decision to publish, or preparation of the manuscript.

References

1.   Arter M, Keeney S. Divergence and conservation of the meiotic recombination machinery.

Nat Rev Genet. 2023 [cited 6 Feb 2024]. doi:10.1038/s41576-023-00669-8

2.   Hotta Y, Stern H. Transient phosphorylation of deoxyribosides and regulation of

deoxyribonucleic acid synthesis. J Biophys Biochem Cytol. 1961/11/01 ed. 1961;11: 311–9.

doi:10.1083/jcb.11.2.311

3.   Lima-De-Faria A, German J, Ghatnekar M, McGovern J, Anderson L. In vitro labelling of

human meiotic chromosomes with H3-thymidine. Hereditas. 1968;60: 249–261.

doi:10.1111/j.1601-5223.1968.tb02205.x

4.   Mukherjee AB, Cohen MM. DNA Synthesis during Meiotic Prophase in Male Mice. Nature.

1968;219: 489–490. doi:10.1038/219489a0

5.   Prensky W. Uptake of thymine-methy6H3 by pachytene chromosomes of Lilium

longiflorum. Genetics. 1962;47: 977.

6.   Wimber DE, Prensky W. Autoradiography with Meiotic Chromosomes of the Male Newt

(Triturus Viridescens) Using H3-Thymidine. Genetics. 1963/12/01 ed. 1963;48: 1731–8.

doi:10.1093/genetics/48.12.1731

7.   Carpenter AT. EM autoradiographic evidence that DNA synthesis occurs at recombination

nodules during meiosis in Drosophila melanogaster females. Chromosoma. 1981/01/01 ed.

1981;83: 59–80. doi:10.1007/BF00286016

8.   Moses MJ, Dresser ME, Poorman PA. Composition and role of the synaptonemal complex.

550   Symp Soc Exp Biol. 1984/01/01 ed. 1984;38: 245–70.

551   9.   Terasawa M, Ogawa H, Tsukamoto Y, Shinohara M, Shirahige K, Kleckner N, et al. Meiotic

552   recombination-related DNA synthesis and its implications for cross-over and non-cross-

553   over recombinant formation. Proc Natl Acad Sci U S A. 2007/03/27 ed. 2007;104: 5965–70.

554   doi:10.1073/pnas.0611490104

555   10. Loidl J, Lukaszewicz A, Howard-Till RA, Koestler T. The Tetrahymena meiotic chromosome

556   bouquet is organized by centromeres and promotes interhomolog recombination. J Cell

557   Sci. 2012/09/15 ed. 2012;125: 5873–80. doi:10.1242/jcs.112664

558   11. Armstrong SJ, Franklin FCH, Jones GH. A meiotic time-course for Arabidopsis thaliana. Sex

559   Plant Reprod. 2003;16: 141–149. doi:10.1007/s00497-003-0186-4

560   12. Stronghill PE, Azimi W, Hasenkampf CA. A novel method to follow meiotic progression in

561   Arabidopsis using confocal microscopy and 5-ethynyl-2'-deoxyuridine labeling. Plant

562   Methods. 2014;10: 33. doi:10.1186/1746-4811-10-33

563   13. Sanchez-Moran E, Santos J-L, Jones GH, Franklin FCH. ASY1 mediates AtDMC1-dependent

564   interhomolog recombination during meiosis in Arabidopsis. Gene Dev. 2007;21: 2220–

565   2233. doi:10.1101/gad.439007

566   14. Allers T, Lichten M. Differential Timing and Control of Noncrossover and Crossover

567   Recombination during Meiosis. Cell. 2001;106: 47–57. doi:10.1016/s0092-8674(01)00416-

568   0

15. Börner GV, Hochwagen A, MacQueen AJ. Meiosis in budding yeast. Rothstein R, editor.

GENETICS. 2023;225: iyad125. doi:10.1093/genetics/iyad125

16. Collins I, Newlon CS. Meiosis-specific formation of joint DNA molecules containing

sequences from homologous chromosomes. Cell. 1994;76: 65–75. doi:10.1016/0092-

8674(94)90173-2

17. Hunter N, Kleckner N. The Single-End Invasion: An Asymmetric Intermediate at the Double-

Strand Break to Double-Holliday Junction Transition of Meiotic Recombination. Cell.

2001;106: 59–70. doi:10.1016/S0092-8674(01)00430-5

18. Schwacha A, Kleckner N. Identification of joint molecules that form frequently between

homologs but rarely between sister chromatids during yeast meiosis. Cell. 1994;76: 51–63.

doi:10.1016/0092-8674(94)90172-4

19. Wijnker E, James GV, Ding J, Becker F, Klasen JR, Rawat V, et al. The genomic landscape of

meiotic crossovers and gene conversions in Arabidopsis thaliana. eLife. 2013;2: e01426.

doi:10.7554/elife.01426

20. Lu P, Han X, Qi J, Yang J, Wijeratne AJ, Li T, et al. Analysis of Arabidopsis genome-wide

variations before and after meiosis and meiotic recombination by resequencing Landsberg

erecta and all four products of a single meiosis. 2012;22: 508–518.

doi:10.1101/gr.127522.111

21. Li R, Bitoun E, Altemose N, Davies RW, Davies B, Myers SR. A high-resolution map of non-

crossover events reveals impacts of genetic diversity on mammalian meiotic

recombination. Nat Commun. 2019;10: 3900. doi:10.1038/s41467-019-11675-y

22. Serrentino M-EM, Borde VV. The spatial regulation of meiotic recombination hotspots: are

all DSB hotspots crossover hotspots? Experimental cell research. 2012;318: 1347–1352.

doi:10.1016/j.yexcr.2012.03.025

23. Chakraborty P, Pankajam AV, Lin G, Dutta A, Krishnaprasad GN, Tekkedil MM, et al.

Modulating Crossover Frequency and Interference for Obligate Crossovers in

Saccharomyces cerevisiae Meiosis. G3 (Bethesda). 2017;7: 1511–1524.

doi:10.1534/g3.117.040071

24. Mancera E, Bourgon R, Brozzi A, Huber W, Steinmetz LM. High-resolution mapping of

meiotic crossovers and non-crossovers in yeast. Nature. 2008;454: 479–485.

doi:10.1038/nature07135

25. Valuchova S, Mikulkova P, Pecinkova J, Klimova J, Krumnikl M, Bainar P, et al. Imaging plant

germline differentiation within Arabidopsis flowers by light sheet microscopy. eLife.

2020;9: 3645–19. doi:10.7554/elife.52546

26. Rossig C, Lievre LL, Pilkington SM, Brownfield L. A simple and rapid method for imaging

male meiotic cells in anthers of model and non-model plant species. Plant Reprod.

2021;34: 37–46. doi:10.1007/s00497-021-00404-5

27. Prusicki MA, Keizer EM, Rosmalen RP van, Komaki S, Seifert F, Müller K, et al. Live cell

imaging of meiosis in Arabidopsis thaliana. eLife. 2019;8: e42834. doi:10.7554/elife.42834

28. Kurzbauer M-T, Uanschou C, Chen D, Schlögelhofer P. The recombinases DMC1 and RAD51

are functionally and spatially separated during meiosis in Arabidopsis. Plant Cell. 2012;24:

2058–2070. doi:10.1105/tpc.112.098459

29. Singh G, Da Ines O, Gallego ME, White CI. Analysis of the impact of the absence of RAD51

612  strand exchange activity in Arabidopsis meiosis. PLoS ONE. 2017;12: e0183006-16.

613  doi:10.1371/journal.pone.0183006

614 30. Varas J, Graumann K, Osman K, Pradillo M, Evans DE, Santos JL, et al. Absence of SUN1 and

615  SUN2 proteins in Arabidopsis thaliana leads to a delay in meiotic progression and defects in

616  synapsis and recombination. Plant J. 2015;81: 329–346. doi:10.1111/tpj.12730

617 31. Concia L, Brooks AM, Wheeler E, Zynda GJ, Wear EE, LeBlanc C, et al. Genome-Wide

618  Analysis of the Arabidopsis Replication Timing Program. PLANT PHYSIOLOGY. 2018;176:

619  2166–2185. doi:10.1104/pp.17.01537

620 32. Da Ines O, White CI. Centromere Associations in Meiotic Chromosome Pairing. Annual

621  review of genetics. 2015;49: 95–114. doi:10.1146/annurev-genet-112414-055107

622 33. Koornneef M, Fransz P, Jong H de. Cytogenetic tools for Arabidopsis thaliana. Chromosom

623  Res. 2003;11: 183–194. doi:10.1023/a:1022827624082

624 34. Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmücker A, et al. The genetic

625  and epigenetic landscape of the Arabidopsis centromeres. Science. 2021;374: eabi7489.

626  doi:10.1126/science.abi7489

627 35. Rowan BA, Heavens D, Feuerborn TR, Tock AJ, Henderson IR, Weigel D. An Ultra High-

628  Density Arabidopsis thaliana Crossover Map That Refines the Influences of Structural

629  Variation and Epigenetic Features. Genetics. 2019;213: genetics.302406.2019.

630  doi:10.1534/genetics.119.302406

631 36. Choi K, Zhao X, Tock AJ, Lambing C, Underwood CJ, Hardcastle TJ, et al. Nucleosomes and

632  DNA methylation shape meiotic DSB frequency in Arabidopsis thaliana transposons and

gene regulatory regions. Genome research. 2018;28: 532–546. doi:10.1101/gr.225599.117

37. Grelon M, Vezon D, Gendrot G, Pelletier G. AtSPO11-1 is necessary for efficient meiotic

recombination in plants. EMBO J. 2001;20: 589–600. doi:10.1093/emboj/20.3.589

38. Chelysheva L, Grandont L, Vrielynck N, le Guin S, Mercier R, Grelon M. An Easy Protocol for

Studying Chromatin and Recombination Protein Dynamics during Arabidopsisthaliana

Meiosis: Immunodetection of Cohesins, Histones and MLH1. Cytogenetic and Genome

Research. 2010;129: 143–153. doi:10.1159/000314096

39. Schwacha A, Kleckner N. Interhomolog Bias during Meiotic Recombination: Meiotic

Functions Promote a Highly Differentiated Interhomolog-Only Pathway. Cell. 1997;90:

1123–1135. doi:10.1016/S0092-8674(00)80378-5

40. Goldfarb T, Lichten M. Frequent and efficient use of the sister chromatid for DNA double-

strand break repair during budding yeast meiosis. PLoS biology. 2010;8: e1000520.

doi:10.1371/journal.pbio.1000520

41. Oh SD, Lao JP, Hwang PY-H, Taylor AF, Smith GR, Hunter N. BLM Ortholog, Sgs1, Prevents

Aberrant Crossing-over by Suppressing Formation of Multichromatid Joint Molecules. Cell.

2007;130: 259–272. doi:10.1016/j.cell.2007.05.035

42. Oh SD, Lao JP, Taylor AF, Smith GR, Hunter N. RecQ Helicase, Sgs1, and XPF Family

Endonuclease, Mus81-Mms4, Resolve Aberrant Joint Molecules during Meiotic

Recombination. Molecular Cell. 2008;31: 324–336. doi:10.1016/j.molcel.2008.07.006

43. Jackson JA, Fink GR. Meiotic recombination between duplicated genetic elements in

Saccharomyces cerevisiae. Genetics. 1985;109: 303–332. doi:10.1093/genetics/109.2.303

44. Haber JE, Thorburn PC, Rogers D. Meiotic and mitotic behaviour of dicentric chromosomes

in Saccharomyces cerevisiae. Genetics. 1984;106: 185–205.

doi:10.1093/genetics/106.2.185

45. Webber HA, Howard L, Bickel SE. The cohesion protein ORD is required for homologue bias

during meiotic recombination. The Journal of Cell Biology. 2004;164: 819–829.

doi:10.1083/jcb.200310077

46. Cole F, Baudat F, Grey C, Keeney S, De Massy B, Jasin M. Mouse tetrad analysis provides

insights into recombination mechanisms and hotspot evolutionary dynamics. Nat Genet.

2014;46: 1072–1080. doi:10.1038/ng.3068

47. Almanzar DE, Gordon SG, Rog O. Meiotic sister chromatid exchanges are rare in C. elegans.

Current Biology. 2021;31: 1499-1507.e3. doi:10.1016/j.cub.2020.11.018

48. Toraason E, Horacek A, Clark C, Glover ML, Adler VL, Premkumar T, et al. Meiotic DNA

break repair can utilize homolog-independent chromatid templates in C. elegans. Current

Biology. 2021;31: 1508-1514.e5. doi:10.1016/j.cub.2021.03.008

49. Billmyre KK, Hughes SE. Meiosis: The elusive sister chromatid repair. Current Biology.

2021;31: R454–R456. doi:10.1016/j.cub.2021.03.093

50. Yang S, Yuan Y, Wang L, Li J, Wang W, Liu H, et al. Great majority of recombination events

in Arabidopsis are gene conversion events. Proceedings of the National Academy of

Sciences of the United States of America. 2012;109: 20992–20997.

doi:10.1073/pnas.1211827110

51. Drouaud J, Khademian H, Giraut L, Zanni V, Bellalou S, Henderson IR, et al. Contrasted

patterns of crossover and non-crossover at Arabidopsis thaliana meiotic recombination

hotspots. Hong Ma, editor. PLoS Genetics. 2013;9: e1003922.

doi:10.1371/journal.pgen.1003922

52. Sims J, Schlögelhofer P, Kurzbauer M-T. From Microscopy to Nanoscopy: Defining an

Arabidopsis thaliana Meiotic Atlas at the Nanometer Scale. Front Plant Sci. 2021;12:

672914. doi:10.3389/fpls.2021.672914

53. Fayos I, Meunier AC, Vernet A, Navarro-Sanz S, Portefaix M, Lartaud M, et al. Assessment

of the roles of SPO11-2 and SPO11-4 in meiosis in rice using CRISPR/Cas9 mutagenesis.

Journal of Experimental Botany. 2020;71: 7046–7058. doi:10.1093/jxb/eraa391

54. Stacey NJ, Kuromori T, Azumi Y, Roberts G, Breuer C, Wada T, et al. Arabidopsis SPO11-2

functions with SPO11-1 in meiotic recombination. The Plant Journal. 2006;48: 206–216.

doi:10.1111/j.1365-313X.2006.02867.x

55. Sims J, Copenhaver GP, Schlögelhofer P. Meiotic DNA Repair in the Nucleolus Employs a

Nonhomologous End-Joining Mechanism. The Plant Cell. 2019;31: 2259–2275.

doi:10.1105/tpc.19.00367

56. Vrielynck N, Schneider K, Rodriguez M, Sims J, Chambon A, Hurel A, et al. Conservation and

divergence of meiotic DNA double strand break forming mechanisms in Arabidopsis

thaliana. Nucleic Acids Res. 2021. doi:10.1093/nar/gkab715

57. Sprink T, Hartung F. Heterologous Complementation of SPO11-1 and -2 Depends on the

Splicing Pattern. IJMS. 2021;22: 9346. doi:10.3390/ijms22179346

58. Vrielynck N, Chambon A, Vezon D, Pereira L, Chelysheva L, Muyt AD, et al. A DNA

topoisomerase VI–like complex initiates meiotic recombination. Science (New York, NY).

2016;351: 939–943. doi:10.1126/science.aad5196

59. Hartung F, Wurz-Wildersinn R, Fuchs J, Schubert I, Suer S, Puchta H. The catalytically active

tyrosine residues of both SPO11-1 and SPO11-2 are required for meiotic double-strand

break induction in Arabidopsis. The Plant cell. 2007;19: 3090–3099.

doi:10.1105/tpc.107.054817

60. Alonso-Ramos P, Álvarez-Melo D, Strouhalova K, Pascual-Silva C, Garside GB, Arter M, et al.

The Cdc14 Phosphatase Controls Resolution of Recombination Intermediates and

Crossover Formation during Meiosis. Int J Mol Sci. 2021;22: 9811.

doi:10.3390/ijms22189811

61. Schubert I, Schubert V, Fuchs J. No Evidence for "Break-Induced Replication" in a Higher

Plant – But Break-Induced Conversion May Occur. Front Plant Sci. 2011;2: 8.

doi:10.3389/fpls.2011.00008

62. Vannier J-B, Depeiges A, White C, Gallego ME. Two roles for Rad50 in telomere

maintenance. The EMBO Journal. 2006;25: 4577–4585. doi:10.1038/sj.emboj.7601345

63. Ross KJ, Fransz P, Jones GH. A light microscopic atlas of meiosis in Arabidopsis thaliana.

Chromosom Res : Int J Mol, Supramol Evol Asp Chromosom Biol. 1996;4: 507–16.

doi:10.1007/bf02261778

64. Dvořáčková M, Raposo B, Matula P, Fuchs J, Schubert V, Peška V, et al. Replication of

ribosomal DNA in Arabidopsis occurs both inside and outside the nucleolus during S phase

progression. J Cell Sci. 2017;131: jcs202416. doi:10.1242/jcs.202416

65. Lambing C, Osman K, Nuntasoontorn K, West A, Higgins JD, Copenhaver GP, et al.

Arabidopsis PCH2 Mediates Meiotic Chromosome Remodeling and Maturation of

Crossovers. PLoS Genet. 2015;11: e1005372. doi:10.1371/journal.pgen.1005372

66. Vanstraelen M, Baloban M, Ines OD, Cultrone A, Lammens T, Boudolf V, et al. APC/C-

CCS52A complexes control meristem maintenance in the Arabidopsis root. Proceedings of

the National Academy of Sciences of the United States of America. 2009;106: 11806–

11811. doi:10.1073/pnas.0901193106

67. Amiard S, Charbonnel C, Allain E, Depeiges A, White CI, Gallego ME. Distinct roles of the

ATR kinase and the Mre11-Rad50-Nbs1 complex in the maintenance of chromosomal

stability in Arabidopsis. THE PLANT CELL. 2010;22: 3020–3033.

doi:10.1105/tpc.110.078527

68. Weisshart K, Fuchs J, Schubert V. Structured Illumination Microscopy (SIM) and

Photoactivated Localization Microscopy (PALM) to Analyze the Abundance and Distribution

of RNA Polymerase II Molecules on Flow-sorted Arabidopsis Nuclei. BIO-PROTOCOL.

2016;6. doi:10.21769/BioProtoc.1725

69. Chelysheva LA, Grandont L, Grelon M. Immunolocalization of Meiotic Proteins in

Brassicaceae: Method 1. In: Pawlowski WP, Grelon M, Armstrong S, editors. Plant Meiosis:

Methods and Protocols. Totowa, NJ: Humana Press; 2013. pp. 93–101. doi:10.1007/978-1-

62703-333-6_9

Figure Legends

Figure 1. EdU labelling strategy and timeline of Arabidopsis **male meiosis.** Pollen mother cell nuclei

were fixed and isolated at the end of a 24-hour pulse of EdU incorporporation. Cells which traversed all or

part of the pre-meiotic S-phase during the pulse will show replicative EdU incorporation and be at

G2/leptotene/zygotene stages at the end of the pulse, with some overlap into early pachytene. Fully

pachytene nuclei (mid-late pachytene) will not show replicative labeling, but will have traversed the

leptotene and zygotene stages in the presence of EdU. Any EdU incorporation in these (or later) nuclei

will be due to non-S-Phase DNA synthesis.

Figure 2. DSB repair-associated DNA synthesis tracts are detected from pachytene up to the end of

meiosis. a) Meiosis I and b) meiosis II stages in *A. thaliana* wildtype. S-phase, replicative DNA labelling is

seen at early G2/letotene and zygotene stages, while clear, discrete foci of EdU substituted DNA are

visible in the chromosomes at later stages from pachytene onwards. Rectangles indicate the enlarged

regions presented in the zooms. Non-chromosomal EdU foci correspond to labelling of mitochondrial

genome replication. 3μm scale bars are shown at the bottom left of each image (1μm for the zooms).

Images were taken with the confocal microscope + Airyscan module.

Figure 3. DSB repair-associated DNA synthesis is SPO11- dependent.

a) Meiosis I and b) meiosis II stages in *A. thaliana* spo11-1 mutants. As for WT meiosis (Figure 2),

replicative S-phase labelling is seen at early stages, corresponding to G2/leptotene and zygotene. As

expected, a striking reduction in numbers of EdU foci (compared to WT) is observed in spo11-1 mutants

at stages from mid-prophase I onwards (corresponding to pachytene and later stages). Rectangles

indicate the enlarged regions presented in the zooms. Images taken with the confocal microscope with

Airyscan module. 3µm scale bars are shown at the bottom left of each image (1µm for the zooms).

Figure 4. Concordance of numbers of SPO11-dependent, meiotic DNA synthesis foci and estimated

DSB numbers.

a) Representative images of wild-type pachytene and *spo11-1* pachytene-like meiotic stages. The merged

DAPI (blue) plus EdU (orange) images are accompanied by the computer-generated images of the scored

EdU foci (white, right panels). EdU foci colocalising with chromosome fibres were retained for counting

and mean intensity used to exclude the much-brighter organellar foci (see text). The Images taken with

the confocal microscope with Airyscan module. 3µm scale bars are shown at the bottom left of each

image. b) Numbers of EdU foci on the chromosomes quantified with Imaris software show clearly the

SPO11-dependence of these DNA synthesis tracts and that their numbers correspond to expectations

from DSB counting in PMC. The mean number of foci/nucleus and number of nuclei counted are shown

above the graph.

Figure 5. Ultrastructure of SPO11-dependent, meiotic DNA synthesis foci on the synaptonemal

complex (SC) during pachytene.

a) Expected configurations of DNA synthesis tracts resulting from different mechanisms of homologous

recombinational repair of meiotic DSB: Synthesis-Dependent Strand Annealing (SDSA), double-Holliday

Junction (dHJ), Gene Conversion ± Crossover (GC±CO) and Break-Induced Replication (BIR). b) Examples

of segments of pachytene SC. Isolated individual foci (one SC lateral axis, white circle) and pairs of foci on

one homologue (one SC lateral axis, green circle) or the two homologues (both SC lateral axes, yellow

circle) are highlighted. The images represent a single slice of 3D-SIM image stacks. 0.5µm scale bars are

shown at the bottom left of each image.

Figure 6. Class I crossovers (CO) colocalize with prophase I DNA synthesis tracts.

a) Wild-type pachytene cell showing DAPI staining (blue), prophase I DNA synthesis tracts (EdU, orange)

and Class 1 CO (MLH1, green). **b)** Examples of colocalising and adjacent MLH1 and EdU foci. The

quantification (right) shows that 95.6% of MLH1 foci co-localize with EdU tracts, and the remainder are

adjacent to one. a) is a confocal image and b) a single slice of a 3D-SIM image stack. Scale bars are 1μm in

(a) and 0.1μm in (b).

Figure 7. Recombination mechanisms and DNA synthesis tracts.

Following resection and invasion of the donor chromatid (a), DNA neo-synthesis tracts resulting from

repair of DSB are expected to be found only on the recipient chromatid in gene conversions (f) resulting

from SDSA (b) or dHJ dissolution (e) pathways. A special case of this is the BIR pathway (c), which will

result in a long synthesis tract, potentially extending all the way to the end of the recipient chromatid.

DNA synthesis tracts are expected to be found on both the recipient and donor chromatids in gene

conversions, associated (g) or not (h) to a CO, resulting from resolution of the dHJ intermediate (d).

Supplemental Figure Legends

Figure S1. Discrete foci of EdU substituted DNA are visible in pachytene chromosomes of Arabidopsis

PMC.

Examples of prophase I DNA labelling in pachtene nuclei of wildtype PMC. Each row is the same nucleus,

imaged with DAPI fluorescence (white, left), EdU (orange, middle) and the merged image (right). Images

taken with the confocal microscope with Airyscan module. 2μm scale bars are included at the bottom left

of each image.

Figure S2. Distribution of SPO11-dependent, meiotic DNA synthesis foci on paired pachytene

chromosomes from two different cells.

Zoomed regions from SIM (a) and Confocal (b) images of meiotic pachytenes, showing the distribution of

EdU-labelled meiotic prophase I DNA synthesis foci (a)red, b) orange) on the DAPI-stained chromosome

fibres (white) of the synaptonemal complexes. 0.5µm scale bars are included at the bottom left of each

image.

Figure 1

Figure 1. EdU labelling strategy and timeline of *Arabidopsis* male meiosis. Pollen mother cell nuclei were fixed and isolated at the end of a 24-hour pulse of EdU incorporporation. Cells which traversed all or part of the pre-meiotic S-phase during the pulse will show replicative EdU incorporation and be at G2/leptotene/zygotene stages at the end of the pulse, with some overlap into early pachytene. Fully pachytene nuclei (mid-late pachytene) will not show replicative labeling, but will have traversed the leptotene and zygotene stages in the presence of EdU. Any EdU incorporation in these (or later) nuclei will be due to non-S-Phase DNA synthesis.

Figure 2

a)

Figure 2

b)

Figure 2. DSB repair-associated DNA synthesis tracts are detected from pachytene up to the end of meiosis.
a) Meiosis I and b) meiosis II stages in *A. thaliana* wildtype. S-phase, replicative DNA labelling is seen at early G2/letotene and zygotene stages, while clear, discrete foci of EdU substituted DNA are visible in the chromosomes at later stages from pachytene onwards. Rectangles indicate the enlarged regions presented in the zooms. Non-chromosomal EdU foci correspond to labelling of mitochondrial genome replication. 3µm scale bars are shown at the bottom left of each image (1µm for the zooms). Images were taken with the confocal microscope + Airyscan module.

Figure 3

a)

Figure 3

Figure 3. DSB repair-associated DNA synthesis is SPO11-dependent.
a) Meiosis I and b) meiosis II stages in *A. thaliana* spo11-1 mutants. As for WT meiosis (Figure 2), replicative S-phase labelling is seen at early stages, corresponding to G2/leptotene and zygotene. As expected, a striking reduction in numbers of EdU foci (compared to WT) is observed in spo11-1 mutants at stages from mid-prophase I onwards (corresponding to pachytene and later stages). Rectangles indicate the enlarged regions presented in the zooms. Images taken with the confocal microscope with Airyscan module. 3μm scale bars are shown at the bottom left of each image (1μm for the zooms).

Figure 4

Figure 4. Concordance of numbers of SPO11-dependent, meiotic DNA synthesis foci and estimated DSB numbers.
a) Representative images of wild-type pachytene and spo11-1 pachytene-like meiotic stages. The merged DAPI (blue) plus EdU (orange) images are accompanied by the computer-generated images of the scored EdU foci (white, right panels). EdU foci colocalising with chromosome fibres were retained for counting and mean intensity used to exclude the much-brighter organellar foci (see text). The Images taken with the confocal microscope with Airyscan module. 3µm scale bars are shown at the bottom left of each image. b) Numbers of EdU foci on the chromosomes quantified with Imaris software show clearly the SPO11-dependence of these DNA synthesis tracts and that their numbers correspond to expectations from DSB counting in PMC. The mean number of foci/nucleus and number of nuclei counted are shown above the graph.

Figure 5

a)

b)

Figure 5. Ultrastructure of SPO11-dependent, meiotic DNA synthesis foci on the synaptonemal complex (SC) during pachytene. a) Expected configurations of DNA synthesis tracts resulting from different mechanisms of homologous recombinational repair of meiotic DSB: Synthesis-Dependent Strand Annealing (SDSA), double-Holliday Junction (dHJ), Gene Conversion ± Crossover (GC±CO) and Break-Induced Replication (BIR). b) Examples of segments of pachytene SC. Isolated individual foci (one SC lateral axis, white circle) and pairs of foci on one homologue (one SC lateral axis, green circle) or the two homologues (both SC lateral axes, yellow circle) are highlighted. The images represent a single slice of 3D-SIM image stacks. 0.5µm scale bars are shown at the bottom left of each image.

Figure 6

a)

Figure 6. Class I crossovers (CO) colocalize with prophase I DNA synthesis tracts.
a) Wild-type pachytene cell showing DAPI staining (blue), prophase I DNA synthesis tracts (EdU, orange) and Class 1 CO (MLH1, green). b) Examples of colocalising and adjacent MLH1 and EdU foci. The quantification (right) shows that 95.6% of MLH1 foci co-localize with EdU tracts, and the remainder are adjacent to one. a) is a confocal image and b) a single slice of a 3D-SIM image stack. Scale bars are 1µm in (a) and 0.1µm in (b).

Figure 7

Figure 7. Recombination mechanisms and DNA synthesis tracts.
Following resection and invasion of the donor chromatid (a), DNA neo-synthesis tracts resulting from repair of DSB are expected to be found only on the recipient chromatid in gene conversions (f) resulting from SDSA (b) or dHJ dissolution (e) pathways. A special case of this is the BIR pathway (c), which will result in a long synthesis tract, potentially extending all the way to the end of the recipient chromatid. DNA synthesis tracts are expected to be found on both the recipient and donor chromatids in gene conversions, associated (g) or not (h) to a CO, resulting from resolution of the dHJ intermediate (d).

Figure S1

Supplemental Figure S1. Discrete foci of EdU substituted DNA are visible in pachytene chromosomes of Arabidopsis PMC.
Examples of prophase I DNA labelling in pachytene nuclei of wildtype PMC. Each row is the same nucleus, imaged with DAPI fluorescence (white, left), EdU (orange, middle) and the merged image (right). Images taken with the confocal microscope with Airyscan module. 2μm scale bars are included at the bottom left of each image.

Figure S2

Figure S2

Supplemental Figure S2. Distribution of SPO11-dependent, meiotic DNA synthesis foci on paired pachytene chromosomes from two different cells.
Zoomed regions from SIM (a) and Confocal (b) images of meiotic pachytenes, showing the distribution of EdU-labelled meiotic prophase I DNA synthesis foci (a)red, b) orange) on the DAPI-stained chromosome fibres (white) of the synaptonemal complexes. $0.5\mu m$ scale bars are included at the bottom left of each image.