Components of Mnemonic Metacognition Constitutionally Supported by High Gamma Activity Between the Precuneus and Hippocampus

Kai Pan*[1,2,3], Xinxia Guo*[1,2,3], Zhe Zheng[1,2,3], Zhoule Zhu[1,2,3], Hemmings Wu[1,2,3], Junming Zhu[1,2,3], Hongyi Ye[1], Yudian Cai[8], Diogo Santos-Pata[5], Jianmin Zhang[1,2,3], Sze Chai Kwok[4,5,6,7], Hongjie Jiang[1,2,3]

[1] Department of Neurosurgery, the Second Affiliated Hospital of Zhejiang University School of Medicine, Hangzhou, Zhejiang, China

[2] Clinical Research Center of Neurological Disease of Zhejiang Province, Hangzhou, Zhejiang, China

[3] Zhejiang Key Laboratory of Research and Transformation for Major Neurosurgical Diseases, Hangzhou, Zhejiang, China

[4] Phylo-Cognition Laboratory, Division of Natural and Applied Sciences, Data Science Research Center, Duke Kunshan University, Duke Institute for Brain Sciences, Kunshan, Jiangsu, China

[5] Shanghai Key Laboratory of Brain Functional Genomics, Key Laboratory of Brain Functional Genomics (Ministry of Education), Shanghai Key Laboratory of Magnetic Resonance, Affiliated Mental Health Center (ECNU), School of Psychology and Cognitive Science, East China Normal University, Shanghai, China

[6] Shanghai Key Laboratory of Magnetic Resonance, East China Normal University, Shanghai, China

[7] Shanghai Changning Mental Health Center, Shanghai, China

[8] Psychological Counseling Center, East China University of Science and Technology, Shanghai 200237, China

Correspondence

(1) Hongjie Jiang, Department of Neurosurgery, Second Affiliated Hospital, School of Medicine, Zhejiang University, 88 Jiefang Road, Hangzhou, Zhejiang 310009, China. Email: insjhj@zju.edu.cn

(2) Sze Chai Kwok, Duke Kunshan University No. 8 Duke Avenue, Kunshan, Jiangsu, 215316 China. Email: sze-chai.kwok@st-hughs.oxon.org

(3) Jianmin Zhang, Department of Neurosurgery, Second Affiliated Hospital, School of Medicine, Zhejiang University, 88 Jiefang Road, Hangzhou, Zhejiang 310009, China. Email: zjm135@zju.edu.cn

Kai Pan and Xinxia Guo authors contributed equally to this work.

ABSTRACT

Metacognition, the ability to introspectively monitor one's own performance, remains an area with poorly understood neural mechanisms, particularly regarding the intercommunication among various brain regions. Recent studies have identified the precuneus as a key region in mnemonic metacognitive tasks, while the hippocampus is recognized for its role in memory function. This study aimed to investigate the role of the precuneus and hippocampus in mnemonic metacognition by utilizing intracranial electrode recordings from patients with intractable epilepsy to analyze the correlational dynamics between these regions. Our findings revealed that high gamma activity serve as a key feature in both the precuneus and hippocampus for confidence judgment. Based on signal detection theory (SDT), we discovered that high gamma activity in the precuneus is significantly linked to type 2 sensitivity (meta-d'), while hippocampal high gamma activity was primarily linked to type 1 sensitivity (d'). Additionally, the correlation between high gamma activity in the hippocampus and precuneus was exclusively related to type 2 sensitivity (meta-d'). Temporal analysis indicated that the hippocampus is initially engaged for the memory process, followed by its joint engagement together with the precuneus for confidence generation. These findings elucidate the distinct electrophysiological roles of the hippocampus and precuneus in mnemonic metacognition, providing deeper insights into the neural mechanisms underlying this cognitive process.

Keywords: metacognition, precuneus, hippocampus, high gamma, regional communication

INTRODUCTION

Metacognition, defined as the ability to introspectively monitor and regulate cognitive processes, involves reevaluating cognition and objectively assessing confidence in decision-making (Flavell, 1979; Fleming et al., 2024; Miyoshi et al., 2024). Numerous studies have elucidated the neural underpinnings of various domains of metacognition, particularly in relation to perception and memory. These investigations have highlighted that the prefrontal cortex's association with perception-related metacognition, and that the precuneus is more intricately linked to memory-related metacognition (Fleming et al., 2010; McCurdy et al., 2013; Baird et al., 2013). Despite advances in identifying the critical roles of the prefrontal (Hebart et al., 2016; Shekhar and Rahnev, 2018; Cai et al., 2022) and parietal cortices (Morales et al., 2018; Pereira et al., 2021) in metacognitive processes, the interregional communication underlying the computation for metacognition remain elusive.

The precuneus has been implicated not only in metacognitive processes related to memory but also in broader memory functions. Studies have demonstrated its involvement in maintaining object-location within working memory (Bicanski & Burgess, 2018), facilitating the retrieval of autobiographical memories (Hebscher et al., 2020), and reconstructing contextual information to aid temporal order memory tasks (Foudil et al., 2020). Given the precuneus's integral role in memory-related cognitive processes and its anatomical connection with the hippocampus (Tanglay et al., 2022), we hypothesized the existence of a neural circuit linking the precuneus and the hippocampus. Empirical evidence supports this hypothesis, documenting hippocampal-precuneus connections that can predict memory retrieval confidence levels (Ren et al., 2018) and highlighting the importance of hippocampal-posterior parietal cortex information flow in episodic memory tasks (Das and Menon, 2023). The pathway between the hippocampus and the precuneus has also been implicated in reasoning decision processes (Pudhiyidath et al., 2022) and in self- and goal-oriented processing (Zheng et al., 2021). The involvement of hippocampal-parietal circuits in these related

processes suggests that communication between the hippocampus and precuneus is also

crucial for memory-related metacognition.

Previous research, which has predominantly utilized magnetic resonance imaging,

animal neurophysiology, or cognitive alterations under pathological states, has yet to

ascertain alterations in neural electrical activity within the human brain during

mnemonic metacognitive processes. Our study addressed this lacuna by analyzing

intracranial electroencephalogram signals from patients with intractable epilepsy,

uncovering the correlation and potential neural pathways linking the precuneus and

hippocampus within mnemonic metacognition. To gain a better understanding of

metacognition and the roles of the precuneus and hippocampus within this performance,

our study employs methodologies based on the signal detection theory (SDT)

framework (Maniscalco and Lau, 2012). By integrating behavioral and

electrophysiological signal analyses, our study elucidated the unique neural

mechanisms of metacognition, thereby contributing to a deeper understanding of this

cognitive process.

RESULT

A   total   of   eleven   patients   with   intractable   epilepsy   underwent

stereoelectroencephalography (sEEG) surgery, during which their brain electrical

signals were recorded during metacognitive behavioral tasks. The demographic and

clinical characteristics of eleven subjects are provided in Supplementary Table S1.

Postoperative metacognitive behavioral testing was conducted in the ward, where

patients were first shown a video and subsequently asked to determine the temporal

order of two displayed pictures. For each trial, after the memory judgment, they were

required to make a confidence judgment (Fig. 1a; see Methods section for details).

Schematic diagram of electrodes was illustrated in Fig.1b. To ensure accurate electrode

placement, we calculated the Montreal Neurological Institute (MNI) coordinates and

visualized them using the AAL3 atlas (Fig. 1c).

Behavioral results

We calculated the mean levels of confidence (3.4148 ± 0.1206 mean ± SEM) and

accuracy (0.7738 ± 0.0288 mean ± SEM) for these eleven participants during the

behavioral experiment (Fig. 1d). After classifying the trials according to the level of

confidence, we found that trials with high confidence had significantly higher accuracy

compared to trials with low confidence (Fig. 1e, $p < 0.001$, paired t-test). Reaction times

during the confidence stage were significantly shorter in correct trials than in incorrect

trials (Fig. 1g, $p < 0.01$, paired t-test), although no significant difference was observed

between high confidence and low confidence trials (Fig. 1f, $p > 0.05$, paired t-test).

There was a significant positive correlation between accuracy and confidence level

across sessions (Fig. 1h, $r = 0.3508$, $p = 0.0075$, Pearson correlation).

Precuneus and hippocampus involvement in metacognition through low gamma and

high gamma frequency band

We first determined the frequency bands in the hippocampus and precuneus that are

most critical for and involved in metacognitive processes. We conducted a time-

frequency analysis, comparing the fixation stage with the confidence stage to calculate

t-values for these regions (Fig. 2a, 2e). The results revealed significant increased

activation in the precuneus and the hippocampus in both low gamma γ (35-70 Hz) and

high gamma Hγ (70-150 Hz) frequency bands. To assess the high frequency band

impact of the precuneus and hippocampus on confidence ratings, we employed

generalized linear models (GLMs) to regress the γ and H γ bands against the confidence

level. We identified significant differences in the -1.0 to -0.5 s interval for both the

hippocampus and precuneus (Fig. 2b, 2f, cluster-based permutation, $p < 0.05$).

Additionally, to verify the contribution of each frequency band to confidence, we

compared GLMs that included only high-gamma and gamma bands with all possible

GLMs that included these high-frequency bands plus any combination of low-

frequency bands. Bayesian model selection (see Methods) indicated that the high-

frequency bands (both low and high gamma) GLM provided the best account of

subjective value (exceedance probability (Xp) = 1). Regression coefficients for each

band showed significant differences for high gamma compared to other bands (Fig. 2c,

2g, one-sample, two-sided t-test, $p < 0.05$). To further explore the influence of other

task-related factors on high-frequency activity, we performed a regression analysis on

the γ and $H\gamma$ bands. The results indicated that, compared to other factors, confidence

level and confidence reaction time had a more pronounced impact. (Fig. 2d, 2h, one-

sample, two-sided t-test, $p < 0.05$). These results suggest that both the hippocampus and

precuneus contribute to confidence rating within the high gamma band, displaying

similar time windows of activity.

High gamma activity reflects distinct cognitive processes in the precuneus and

hippocampus

Given the involvement of high gamma band activity in the confidence rating process,

we investigated the cognitive processes represented by high gamma activity in the

precuneus and hippocampus. Metacognition is often measured by calculating d' and

meta-d' based on signal detection theory (SDT). To explore this, we computed an

accuracy index with the neural data ($H\gamma_{Correct} - H\gamma_{Incorrect}$) which corresponds to the

behavioral index d'. We also calculated an interaction index of confidence and accuracy

using the neural data [($H\gamma_{High\ confidence} - H\gamma_{Low\ confidence}$) - ($H\gamma_{Correct} - H\gamma_{Incorrect}$)] to

correspond to the behavioral index meta-d', which refers to one's efficacy with which

confidence ratings discriminate between correct and incorrect judgments.

Using a 0.1s window, we assessed changes in power spectral density of high gamma

activity during the confidence rating stage, standardizing the data using z-scores. In the

precuneus, high gamma activity began to change between approximately -0.3 s and 0 s,

for either comparing high vs. low confidence or comparing high vs. low accuracy trials

(Fig. 3a, left and right panels). In contrast, in the hippocampus, changes occurred

between -0.8 s and -0.5 s, also showing consistency for both confidence and accuracy

(Fig. 4a, left and right panels).

We then averaged the accuracy index and interaction index of high gamma within these

two windows across all subjects and conducted Pearson correlation analyses with d' and

meta-d', respectively. Pearson correlation analyses revealed that high gamma activity

in the precuneus was significantly correlated with meta-d', but not with d' (Fig. 3b).

Conversely, high gamma activity in the hippocampus was significantly correlated with

d', but not with meta-d' (Fig. 4b). Further analysis showed no significant correlation

between precuneus high gamma activity during the hippocampus's significant time

window and d' or meta-d' (Supplementary Fig. S1a). However, hippocampal high

gamma activity during the precuneus's significant time window showed a trend towards

correlation with meta-d' (Supplementary Fig. S1b). These findings suggest that the

hippocampus and precuneus both play a role in confidence judgment, with the

precuneus being more involved in metacognitive processes (reflected by meta-d') and

the hippocampus more engaged in memory processes (reflected by d'). The different

time windows emphasize their distinct contributions. Given the observed trend of

correlation between high gamma activity in the hippocampus and meta-d' within the

same time window, our next step is to investigate whether the activities between the

hippocampus and precuneus might be correlated or synchronized differently for these

distinct cognitive processes.

Correlation between high gamma activity in the precuneus and hippocampus related

with metacognition

Having established the relationship between the high gamma activity in the precuneus

and metacognition, we next examined how hippocampal gamma activity might

correlate with that of the precuneus. We applied a correlation analysis method using a

0.02 s window and 0.01s steps to compute Pearson correlation coefficient between the

hippocampus and precuneus signals. Amplitude segments with significant correlations

lasting over 0.04 s were counted (Fig. 5a). Additionally, segments lasting over 0.06 s

were calculated, showing results consistent with those obtained with using 0.04 s

segments (Supplementary Fig. S2). We ran correlation analyses for each pair of

electrode sites (82 pairs in total) from the 11 subjects. Overall, we found that 60 out of

these 82 electrode pairs between the precuneus and hippocampus showed significantly

higher correlation values compared to shuffled data. Taking one example participant,

the original data exhibited a higher probability density in greater counts than the

shuffled data (Fig. 5b, $p < 0.001$, permutation test, $n = 1000$). We then ran this analysis

on each participant individually and found the same significant differences

(Supplementary Fig. S3). The correlation count for the original data was significantly

higher than that for the shuffled data (Fig. 5c, Wilcoxon matched pairs signed-rank test,

$p < 0.0001$). This elevated correlation count indicates a stronger relationship between

high gamma activity in the hippocampus and precuneus. Moreover, we applied this

index to examine its relationship with the metacognitive indices. When categorized by

confidence level, high confidence trials had a significantly higher correlation count than

low confidence trials (Fig. 5d, Wilcoxon matched-pairs signed-rank test, $p < 0.05$).

Incorrect trials exhibited a significantly higher correlation count than correct trials (Fig.

5e, Wilcoxon matched-pairs signed-rank test, $p < 0.0001$). In alignment with the power

analysis, we also calculated the interaction index in correlation count [(Count $_{High}$

$_{confidence}$ − Count $_{Low\ confidence}$) - (Count $_{Correct}$ − Count $_{Incorrect}$)], which was significantly

greater than zero (Fig. 5g, one-sample two-sided t-test, $p < 0.0001$). For completeness,

we ran Pearson correlation analyses and revealed that the interaction index showed a

trend towards correlation with meta-d' (Supplementary Fig. S4b) but not between the

accuracy index and d' (Fig. S4a).

Given the strong correlation identified between the precuneus and hippocampus during

the confidence rating stage, we then examined the temporal dynamics of this

relationship. We calculated the Pearson correlation coefficient between the amplitudes

of the hippocampus and the precuneus using a 0.02s window, averaging it according to

the 0.1s window used in power analysis. We called this correlation coefficient as H-P

correlation. The H-P correlations were significant at approximately -0.3 s and -0.9 s

(Fig. 6a), aligning with previously identified changes in the power analysis (Fig. 3a,

4a). Additionally, we conducted Pearson correlation analyses between the H-P

correlation and either d' or meta-d'. The H-P correlation coefficients at approximately -

0.9s showed no significant correlation with either meta-d' or d' (Fig. 6b). In contrast,

the H-P correlation coefficients around -0.3s exhibited a significant correlation with

meta-d' but were not with d' (Fig. 6c). These results demonstrated that the precuneus

and hippocampus might jointly participate in the process of confidence rating, with

their interaction being specifically related to type 2 performance rather than type 1

performance, highlighting the significance of their connection in metacognitive

processes.

Discussion

In this study, we employed intracranial electrode recordings to investigate changes in

brain electrical activity during mnemonic metacognitive processes. Specifically, we

analyzed first-order and second-order performance in metacognition by examining the

distinct characteristics and correlations of specific frequency bands in the precuneus

and hippocampus. Our findings underscore the critical role of the temporal-to-parietal

pathway in the genesis of metacognition, emphasizing the significance of neural circuits

over isolated brain regions in metacognitive processes.

Our study is the first to identify the involvement of high gamma band activity in both

the precuneus and hippocampus during the confidence judgment stage. The role of the

precuneus in metacognition, including mnemonic metacognition, has been extensively

documented in previous research (McCurdy et al., 2013; Baird et al., 2013; Ye et al.,

2018; Ye et al., 2019), as well as in non-memory domains (Fleming et al., 2010;

Heereman et al., 2015; Morales et al., 2018). The literature suggests that high gamma

activity in the precuneus is associated with memory retrieval processes (Hebscher et al.,

2019), whereas our findings provide evidence that the precuneus contributes to

metacognitive processes via high-frequency bands (Manippa et al., 2024). This may

explain the cognitive improvements observed following high-frequency stimulation of

the precuneus in dementia patients (Morris and Hannesdottir, 2004; Mograbi et al.,

2012; Koch et al., 2022).

Metacognition, as a complex and advanced cognitive function, should not be

anatomically attributed to a single brain region. Indeed, like in memory, mnemonic

processes are multifaceted and can be neurally dissociated into precision, vividness,

and accuracy (Richter et al., 2016). Beyond that, our ability to monitor some of these

memory-related processes is also segregated neurally (Zou and Kwok, 2022). Previous

studies have suggested the role of functional connectivity between the precuneus and

the hippocampus in metacognitive processes. The two regions are linked by the superior

longitudinal fasciculus (SLF), emphasizing the relevance of both white and grey matter

pathways (Zheng et al., 2021). Building on that, this study focuses on the functional

connection between the precuneus and the hippocampus. Several studies have

identified significant interactions between the hippocampus and precuneus in various

cognitive functions, including memory (Ren et al., 2018), reasoning and decision-

making (Pudhiyidath et al., 2023), and self- and goal-oriented processing (Zheng et al.,

2021). Regarding the neural mechanisms underlying these interactions, phase-

amplitude coupling has been observed between the medial temporal lobe and precuneus

during memory retrieval (Hebscher et al., 2019). Our study supports the involvement

of both the hippocampus and the precuneus in metacognitive processes at high-

frequency bands, highlighting potential differences in the neural mechanisms

underlying memory retrieval and metacognition. As a control, we conducted a phase-

amplitude coupling (PAC) analysis between the hippocampus and precuneus (Voytek

et al., 2009; Jensen et al., 2016), which showed no significant difference in coupling

between the retrieval and confidence stages (Supplementary Fig. S5). Although this

finding provides no direct evidence for confidence-related PAC, we propose that this

PAC phenomenon likely originates from memory retrieval processes (Richter et al.,

2016).

Pereira et al. (2021) reported a strong correlation between evidence accumulation and

meta-perception in the posterior parietal cortex by single-neuron activity of a

microelectrode. However, the limitations of electrocorticography (ECoG) in their study

left unresolved whether deep brain structures, such as the hippocampus, are implicated

during metacognition. There is evidence suggesting the neural circuitry linking the

hippocampus to the cortex is important for metacognition. Some work reported that the

structural integrity of the fornix, as the output tract of the hippocampus, is related to

metacognition driven cognitive offloading (Zheng et al., 2024). This suggests that the

fornix is not only important for mnemonic functions (Kwok et al., 2015; Kwok &

Buckley, 2010) but also provides a pathway to deliver memory related signals for the

frontal-parietal cortice (Boldt & Gilbert, 2022; Kwok et al., 2019) s to generate

confidence and metacognitive computation. These are in line with reports that the

hippocampus plays a role in memory-related metacognition (Allen et al., 2017;

Miyamoto et al., 2017, 2018). This possibility is corroborated with findings showing

resting-state functional connectivity (rs-fcMRI) between the precuneus and

hippocampus is positively correlated with metamemory ability (Ye et al., 2019). This

is in line with anatomical evidence for the interaction between the two regions as they

are connected by the fiber bundle between the precuneus to the parahippocampal gyrus

(Tanglay et al. 2022). Our findings thus help characterize the interaction via these

cortical-subcortical regions in metacognitive processes in a memory task.

In the hippocampus, many studies have demonstrated the role of high gamma bands in

memory-related cognitive activities (Burke et al., 2014; Staresina et al., 2016; Mart,

2021). In addition to their involvement in memory processes, memory-selective

neurons in the hippocampus exhibit firing patterns associated with confidence

judgments (Rutishauser et al., 2015). Gray matter myelination in the hippocampus has

been linked to individuals' metacognitive abilities (Allen et al., 2017). Our findings

suggest that high gamma activity in the hippocampus also plays a role in metacognitive

processes. Given the joint involvement of the precuneus and hippocampus in
confidence reporting, the correlation between high gamma activity in these two brain
regions is particularly noteworthy. Indeed, our study confirms a significant correlation
between high gamma activity in the precuneus and the hippocampus.

Importantly, our results showed first-order and second-order performance in
metacognition could be dissociated through a high gamma band communication
between the hippocampus and precuneus. In memory tasks, first-order performance
refers to the memory performance. We identified a significant correlation between first-
order performance and high gamma band power in the hippocampus, with this activity
occurring approximately 0.8 to 0.5 s before the confidence reporting. Additionally,
elevated gamma band activity in the hippocampus during confidence reporting was
observed for trials with incorrect and high confidence responses. The high gamma band
activity was negatively correlated with type 1 sensitivity (d') across participants,
suggesting that the hippocampus may generate an "error signal" during first-order
performance, consistent with previous findings (Haque et al., 2020; Desender et al.,
2021). Regarding confidence, or second-order performance, we found that it to be
primarily correlated with high gamma power in the precuneus, occurring approximately
0.3 to 0 s before the confidence rating. A similar time window, spanning 0.2 to 0.5 s
after stimulus onset, had been observed in meta-perception (Xue et al., 2023).
Importantly, the correlation between the precuneus and hippocampus was significantly
associated with second-order performance but not with first-order performance. These
findings confirm that the hippocampus and precuneus cross talk to each other and
jointly contribute to metacognition. While we established the synergistic role of the
precuneus and hippocampus in metacognitive processes, we furthermore observed
some functional differences between the two regions. Neither the precuneus alone nor
its interaction with the hippocampus was associated with first-order performance; only
the hippocampus showed involvement in these first-order processes. We also observed
significant correlation around the first-order performance period of hippocampus,
which may reflect functional connectivity between the precuneus and hippocampus

during working memory processes (Ren et al., 2018; Hebscher et al., 2019). These findings underscore the distinct roles of the hippocampus and precuneus in metacognition process: the hippocampus contributes to both memory and metacognitive process, while the precuneus appears specifically engaged in metacognitive processes. These results also highlight a temporal distinction between first-order performance (d') and metacognitive performance (meta-d'). It would be important to further verify whether this temporal distinction is domain-specific or generalizable across different types of metacognition in future research.

In conclusion, our study demonstrates that the precuneus and hippocampus are both implicated in mnemonic metacognition. Through high gamma band activity communication, the precuneus and hippocampus together contribute to second-order performance, whereas first-order performance is primarily mediated by the hippocampus alone. Our findings provide electrophysiological evidence of a parietal-hippocampal circuit (see also the role of fornix in Zheng et al., 2024) (see also the role of fornix in Zheng et al., 2024) underpinning mnemonic metacognition in humans.

REFERENCES

1. Allen, M., Glen, J. C., Müllensiefen, D., Schwarzkopf, D. S., Fardo, F., Frank, D., Callaghan, M. F., & Rees, G. (2017). Metacognitive ability correlates with hippocampal and prefrontal microstructure. *NeuroImage*, *149*, 415–423. https://doi.org/10.1016/j.neuroimage.2017.02.008

2. Baird, B., Smallwood, J., Gorgolewski, K. J., & Margulies, D. S. (2013). Medial and lateral networks in anterior prefrontal cortex support metacognitive ability for memory and perception. *Journal of Neuroscience*, *33*(42), 16657–16665. https://doi.org/10.1523/JNEUROSCI.0786-13.2013

3. Bicanski, A., & Burgess, N. (2018). A neural-level model of spatial memory and imagery. *ELife*, *7*. https://doi.org/10.7554/eLife.33752

4. Boldt, A., & Gilbert, S. J. (2022). Partially Overlapping Neural Correlates of Metacognitive Monitoring and Metacognitive Control. Journal of Neuroscience, 42(17), 3622–3635. https://doi.org/10.1523/JNEUROSCI.1326-21.2022

5. Burke, J. F., Long, N. M., Zaghloul, K. A., Sharan, A. D., Sperling, M. R., & Kahana, M. J. (2014). NeuroImage Human intracranial high-frequency activity maps episodic memory formation in space and time. *NeuroImage*, *85*, 834–843. https://doi.org/10.1016/j.neuroimage.2013.06.067

6. Cai, Y., Jin, Z., Zhai, C., Wang, H., Wang, J., Tang, Y., & Kwok, S. C. (2022). Time-sensitive prefrontal involvement in associating confidence with task performance illustrates metacognitive introspection in monkeys. *Communications Biology*, *5*(1). https://doi.org/10.1038/s42003-022-03762-6

7. Das, A., & Menon, V. (2023). *Hippocampal-parietal cortex causal information flow in human episodic memory : replication across three experiments.*

8. Desender, K., Ridderinkhof, K. R., & Murphy, P. R. (2021). Understanding neural signals of post-decisional performance monitoring: An integrative review. *ELife*, *10*, 1–21. https://doi.org/10.7554/eLife.67556

9. Flavell, J. H. (1979). Metacognition and cognitive monitoring: A new area of cognitive–developmental inquiry. American Psychologist, 34(10), 906–911.

10. Fleming, Stephen M; Weil, Rimona S; Nagy, Zoltan; Dolan, Raymond J; & Rees, G. (2010). Tackling Attack Detection and Incident Response. *Science*, *329*(April), 1541–1543. https://doi.org/10.1126/science.1191883.Relating

11. Fleming, S. M. (2024). Metacognition and Confidence: A Review and Synthesis. *Annual Review of Psychology*, *75*(1), 1–28. https://doi.org/10.1146/annurev-psych-022423-032425

12. Foudil, S. A., Kwok, S. C., & Macaluso, E. (2020). Context-dependent coding of temporal distance between cinematic events in the human precuneus. *Journal of Neuroscience*, *40*(10). https://doi.org/10.1523/JNEUROSCI.2296-19.2020

13. Futing Zou, Sze Chai Kwok; Distinct Generation of Subjective Vividness and Confidence during Naturalistic Memory Retrieval in Angular Gyrus. J Cogn Neurosci 2022; 34 (6): 988–1000. doi: https://doi.org/10.1162/jocn_a_01838

14. Green, D.M., Swets, J.A., 1966. Signal Detection Theory and Psychophysics. John Wiley, Oxford.

15. Haque, R. U., Inati, S. K., Levey, A. I., & Zaghloul, K. A. (n.d.). *episodic memory retrieval. 2020*, 1–14. https://doi.org/10.1038/s41467-020-19828-0

16. Hebart, M. N., Schriever, Y., Donner, T. H., & Haynes, J. D. (2016). The Relationship between

Perceptual Decision Variables and Confidence in the Human Brain. *Cerebral Cortex*, *26*(1), 118–130. https://doi.org/10.1093/cercor/bhu181

17. Hebscher, M., Ibrahim, C., & Gilboa, A. (2020). Precuneus stimulation alters the neural dynamics of autobiographical memory retrieval. *NeuroImage*, *210*(June 2019), 116575. https://doi.org/10.1016/j.neuroimage.2020.116575

18. Hebscher, M., Meltzer, J. A., & Gilboa, A. (2019). A causal role for the precuneus in network-wide theta and gamma oscillatory activity during complex memory retrieval. *ELife*, *8*, 1–20. https://doi.org/10.7554/eLife.43114

19. Heereman, J., Walter, H., & Heekeren, H. R. (2015). A task-independent neural representation of subjective certainty in visual perception. *Frontiers in Human Neuroscience*, *9*(OCT). https://doi.org/10.3389/fnhum.2015.00551

20. Koch G, Casula EP, Bonnì S, Borghi I, Assogna M, Minei M, Pellicciari MC, Motta C, D'Acunto A, Porrazzini F, Maiella M, Ferrari C, Caltagirone C, Santarnecchi E, Bozzali M, Martorana A. Precuneus magnetic stimulation for Alzheimer's disease: a randomized, sham-controlled trial. Brain. 2022 Nov 21;145(11):3776-3786. doi: 10.1093/brain/awac285. PMID: 36281767; PMCID: PMC9679166.

21. Kwok, S. C., & Buckley, M. J. (2010). Fornix transection selectively impairs fast learning of conditional visuospatial discriminations. Hippocampus, 20(3), 413–422. https://doi.org/10.1002/hipo.20643

22. Kwok, S. C., Cai, Y., & Buckley, M. J. (2019). Mnemonic introspection in macaques is dependent on superior dorsolateral prefrontal cortex but not orbitofrontal cortex. Journal of Neuroscience, 39(30), 5922–5934. https://doi.org/10.1523/JNEUROSCI.0330-19.2019

23. Kwok, S. C., Mitchell, A. S., & Buckley, M. J. (2015). Adaptability to changes in temporal structure is fornix-dependent. Learning and Memory, 22(8), 354–359. https://doi.org/10.1101/lm.038851.115

24. Lopez-Persem, A., Bastin, J., Petton, M., Abitbol, R., Lehongre, K., Adam, C., Navarro, V., Rheims, S., Kahane, P., Domenech, P., & Pessiglione, M. (2020). Four core properties of the human brain valuation system demonstrated in intracranial signals. Nature Neuroscience, 23(5), 664–675. https://doi.org/10.1038/s41593-020-0615-9

25. Maniscalco, B., & Lau, H. (2012). A signal detection theoretic approach for estimating metacognitive sensitivity from confidence ratings. *Consciousness and Cognition*, *21*(1), 422–430. https://doi.org/10.1016/j.concog.2011.09.021

26. Manippa, V., Palmisano, A., Nitsche, M. A., Filardi, M., Vilella, D., Logroscino, G., & Rivolta, D. (2024). Cognitive and Neuropathophysiological Outcomes of Gamma-tACS in Dementia: A Systematic Review. Neuropsychology Review, 34(1), 338–361. https://doi.org/10.1007/s11065-023-09589-0

27. Mart, C. (2021). *The hippocampus as the switchboard between perception and memory*. *118*(50). https://doi.org/10.1073/pnas.2114171118/-/DCSupplemental.Published

28. McCurdy, L. Y., Maniscalco, B., Metcalfe, J., Liu, K. Y., de Lange, F. P., & Lau, H. (2013). Anatomical coupling between distinct metacognitive systems for memory and visual perception. *Journal of Neuroscience*, *33*(5), 1897–1906. https://doi.org/10.1523/JNEUROSCI.1890-12.2013

29. Miyamoto K, Osada T, Setsuie R, Takeda M, Tamura K, Adachi Y, Miyashita Y. Causal neural network of metamemory for retrospection in primates. Science. 2017 Jan 13;355(6321):188-193. doi: 10.1126/science.aal0162. PMID: 28082592.

30. Miyamoto, K., Setsuie, R., Osada, T., & Miyashita, Y. (2018). Reversible Silencing of the
Frontopolar Cortex Selectively Impairs Metacognitive Judgment on Non-experience in Primates.
Neuron, 97(4), 980-989.e6. https://doi.org/10.1016/j.neuron.2017.12.040

31. Miyoshi, K., Webb, T., Rahnev, D., & Lau, H. (2024). Confidence and metacognition. In *Reference*
Module in Neuroscience and Biobehavioral Psychology. Elsevier. https://doi.org/10.1016/b978-0-
12-820480-1.00049-8

32. Mograbi DC, Ferri CP, Sosa AL, Stewart R, Laks J, Brown R, Morris RG. Unawareness of memory
impairment in dementia: a population-based study. Int Psychogeriatr. 2012 Jun;24(6):931-9. doi:
10.1017/S1041610211002730. Epub 2012 Jan 17. PMID: 22251835.

33. Morales, J., Lau, H., & Fleming, S. M. (2018). Domain-general and domain-specific patterns of
activity supporting metacognition in human prefrontal cortex. *Journal of Neuroscience*, *38*(14),
3534–3546. https://doi.org/10.1523/JNEUROSCI.2360-17.2018

34. Morris, Robin G, and Kristin Hannesdottir, 'Loss of 'Awareness' in Alzheimer's Disease', i
n Robin G Morris, and James T Becker (eds), Cognitive Neuropsychology of Alzheimer's
Disease (Oxford, 2004; online edn, Oxford Academic, 31 Oct. 2023), https://doi.org/10.1093/o
so/9780198508304.003.0017, accessed 18 Aug. 2024.

35. Pereira, M., Megevand, P., Tan, M. X., Chang, W., Wang, S., Rezai, A., Seeck, M., Corniola, M.,
Momjian, S., Bernasconi, F., Blanke, O., & Faivre, N. (2021). Evidence accumulation relates to
perceptual   consciousness   and   monitoring.   *Nature   Communications*,   *12*(1),   1–11.
https://doi.org/10.1038/s41467-021-23540-y

36. Pudhiyidath, A., Morton, N. W., Duran, R. V., Schapiro, A. C., Momennejad, I., Hinojosa-Rowland,
D. M., Molitor, R. J., & Preston, A. R. (2022). Representations of Temporal Community Structure
in Hippocampus and Precuneus Predict Inductive Reasoning Decisions. *Journal of Cognitive*
Neuroscience, *34*(10), 1736–1760. https://doi.org/10.1162/jocn_a_01864

37. Ren, Y., Nguyen, V. T., Sonkusare, S., Lv, J., Pang, T., Guo, L., Eickhoff, S. B., Breakspear, M.,
& Guo, C. C. (2018). Effective connectivity of the anterior hippocampus predicts recollection
confidence   during   natural   memory   retrieval.   *Nature   Communications*,   *9*(1).
https://doi.org/10.1038/s41467-018-07325-4

38. Rigoux, L., Stephan, K. E., Friston, K. J., & Daunizeau, J. (2014). Bayesian model selection for
group   studies   -   Revisited.   NeuroImage,   84,   971   –   985.
https://doi.org/10.1016/j.neuroimage.2013.08.065

39. Richter, F. R., Cooper, R. A., Bays, P. M., & Simons, J. S. (2016). Distinct neural mechanisms
underlie the success, precision, and vividness of episodic memory. ELife, 5(OCTOBER2016), 1–
18. https://doi.org/10.7554/eLife.18260

40. Rutishauser, U., Ye, S., Koroma, M., Tudusciuc, O., Ross, I. B., Chung, J. M., & Mamelak, A. N.
(2015). Representation of retrieval confidence by single neurons in the human medial temporal lobe.
Nature Neuroscience, *18*(7), 1041–1050. https://doi.org/10.1038/nn.4041

41. Shekhar, M., & Rahnev, D. (2018). Distinguishing the roles of dorsolateral and anterior PFC in
visual   metacognition.   *Journal   of   Neuroscience*,   *38*(22),   5078–5087.
https://doi.org/10.1523/JNEUROSCI.3484-17.2018

42. Staresina, B. P., Michelmann, S., Bonnefond, M., Jensen, O., Axmacher, N., & Fell, J. (2016).
Hippocampal pattern completion is linked to gamma power increases and alpha power decreases
during recollection. 1–18. https://doi.org/10.7554/eLife.17397

43. Stephan, K. E., Penny, W. D., Daunizeau, J., Moran, R. J., & Friston, K. J. (2009). Bayesian model

508   selection   for   group   studies.   NeuroImage,   46(4),   1004   –   1017.

509   https://doi.org/10.1016/j.neuroimage.2009.03.025

44.   Tanglay, O., Young, I. M., Dadario, N. B., Briggs, R. G., Fonseka, R. D., Dhanaraj, V., Hormovas,

J., Lin, Y. H., & Sughrue, M. E. (2022). Anatomy and white-matter connections of the precuneus.

In *Brain Imaging and Behavior* (Vol. 16, Issue 2, pp. 574–586). https://doi.org/10.1007/s11682-

021-00529-1

45.   Voytek B, D'Esposito M, Crone N, Knight RT. A method for event-related phase/amplitude

coupling. *Neuroimage*. 2013;64:416-424. doi:10.1016/j.neuroimage.2012.09.023

46.   Xue, K., Zheng, Y., Rafiei, F., & Rahnev, D. (2023). The timing of confidence computations in

human prefrontal cortex. *Cortex*, 1–28. https://doi.org/10.1016/j.cortex.2023.08.009

47.   Ye, Q., Zou, F., Dayan, M., Lau, H., Hu, Y., & Kwok, S. C. (2019). Individual susceptibility to

TMS affirms the precuneal role in meta-memory upon recollection. *Brain Structure and Function*,

224(7), 2407–2419. https://doi.org/10.1007/s00429-019-01909-6

48.   Ye, Q., Zou, F., Lau, H., Hu, Y., & Kwok, S. C. (2018). Causal evidence for mnemonic

metacognition in human precuneus. *Journal of Neuroscience*, *38*(28), 6379–6387.

https://doi.org/10.1523/JNEUROSCI.0660-18.2018

49.   Zheng, Y., Bo, B., Wang, D., Liu, Y., Gilbert, S. J., Li, Y., & Kwok, S. C. (2024). Fornix and

Uncinate Fasciculus Support Metacognition-Driven Cognitive Offloading. 21, 1–26.

50.   Zheng, Y., Wang, D., Ye, Q., Zou, F., Li, Y., & Kwok, S. C. (2021). Diffusion property and

functional connectivity of superior longitudinal fasciculus underpin human metacognition.

Neuropsychologia,156   (October   2020),   107847.

https://doi.org/10.1016/j.neuropsychologia.2021.107847

51.   Zheng, A., Montez, D. F., Marek, S., Gilmore, A. W., Newbold, D. J., Laumann, T. O., Kay, B. P.,

Seider, N. A., Van, A. N., Hampton, J. M., Alexopoulos, D., Schlaggar, B. L., Sylvester, C. M.,

Greene, D. J., Shimony, J. S., Nelson, S. M., Wig, G. S., Gratton, C., McDermott, K. B., …

Dosenbach, N. U. F. (2021). Parallel hippocampal-parietal circuits for self- And goal-oriented

processing. *Proceedings of the National Academy of Sciences of the United States of America*,

118(34). https://doi.org/10.1073/pnas.2101743118

52.   Zou, Futing, and Sze Chai Kwok. "Distinct Generation of Subjective Vividness and Confidence

during Naturalistic Memory Retrieval in Angular Gyrus." *Journal of cognitive neuroscience* vol.

34,6 (2022): 988-1000. doi:10.1162/jocn_a_01838

53.   Zuo, S., Wang, L., Shin, J. H., Cai, Y., Lee, S. W., Appiah, K., Zhou, Y. Di, & Kwok, S. C. (2020).

Behavioral evidence for memory replay of video episodes in the macaque. *ELife*, *9*, 1–23.

https://doi.org/10.7554/eLife.54519

Figure 1. Experimental design, electrode's location and behavioral details. (a) Experimental
design. Participants performed a temporal order judgment task, by choosing the image that happened
earlier in the video. After this, participants needed to report confidence level from 1-4 points. **(b)**
Schematic diagram of stereoelectroencephalography electrodes used in this study. **(c)** A summary
of the recording locations from 11 participants. Each colored dot represents the location of a
recording site shown on a standard brain template (AAL Atlas). The two colors represent the brain
areas. Blue: hippocampus; Red: precuneus. **(d)** Bar plot shows average confidence level (3.40±0.41)
and average accuracy (77%±9.5) across participants. **(e)** Accuracy plotted in high confidence and
low confidence trials across participants, with a significant difference (paired t-test, p = 0.0078). **(f)**
Reaction time in confidence rating stage of high confidence trials compared with low confidence
trials across participants (paired t-test, p > 0.05). **(g)** Reaction time in confidence rating stage of
correct trials is significantly shorter than incorrect trials (paired t-test, p = 0.0058). **(h)** Correlation
between the confidence level and accuracy across sessions collapsed over participants（r = 0.3508,
p = 0.0075）. Each dot denotes a session.

Figure 2. Gamma activity in the precuneus and hippocampus associated with confidence. (a)
Time-frequency analysis (T-value map) during the confidence rating stage, compared with the
fixation stage, averaged across all electrodes in the precuneus. Horizontal dashed lines indicate
boundaries between frequency bands that are included in the analysis in (c). **(b)** Time course of
regression estimates for the gamma and high-gamma frequency bands in the precuneus. The solid
line (shaded areas) represents the SEM across the precuneual electrode sites (n=31). The horizontal
black rectangle on the x-axis represents significant time points (cluster-based permutation, p < 0.05).
(c) Regression estimates of power against confidence rating in the precuneus, averaged over the -
1.0 to -0.25s window, for each frequency band defined in (a). Asterisks indicate significance of
regression estimates (one-sample, two-sided t-test, p < 0.05). **(d)** Regression estimates of related
factors against gamma and high gamma in the precuneus, averaged over the -1.0 to -0.25s window.
Asterisks indicate significance of regression estimates (one-sample, two-sided t-test). **(e)** Time-
frequency analysis (T-value map) during the confidence rating stage, compared with the fixation
stage, averaged across all electrodes in the hippocampus. Horizontal dashed lines indicate
boundaries between frequency bands that are included in the analysis in (e). **(f)** Time course of
regression estimates for the gamma and high-gamma frequency bands in the hippocampus. The solid
line (shaded areas) represents the SEM across the hippocampal electrode sites (n=24). The
horizontal black rectangle on the x-axis represents significant time points (cluster-based permutation,
p < 0.05). **(g)** Regression estimates of power against confidence rating in the hippocampus, averaged
over the -1.0 to -0.25s window, for each frequency band defined in (e). Asterisks indicate
significance of regression estimates (one-sample, two-sided t-test, γ: p < 0.01, Hγ: p < 0.001). **(h)**
Regression estimates of related factors against gamma and high gamma in the hippocampus,
averaged over the -1.0 to -0.25s window. Asterisks indicate significance of regression estimates
(one-sample, two-sided t-test, p < 0.05).

Figure 3. High gamma power in the precuneus is associated with meta-d' but not with d'. (a)
Left panel: The time course of the difference in high gamma power (z-score) of the precuneus
between high confidence trials and low confidence trials. Right panel: The time course of the
difference in high gamma power (z-score) of the precuneus between correct trials and incorrect trials.
The solid line represents the mean with SEM (shaded areas) across electrode sites (n=31). The
horizontal black rectangle on the x-axis represents significant time points (cluster-based permutation,
$p < 0.05$). The vertical red shaded areas denote the time window of interest used in **(b)**. **(b)** The left
panel illustrates the correlation between the neural accuracy index (Hγ $_{Correct}$ − Hγ $_{Incorrect}$) and
behavioral type 1 sensitivity (d') (Pearson correlation, r = -0.2814, p = 0.4019), whereas the right
figure depicts the correlation between the neural interaction index [(Hγ $_{High\ confidence}$ − Hγ $_{Low\ confidence}$)
- (Hγ $_{Correct}$ − Hγ $_{Incorrect}$)] and behavioral type 2 sensitivity (meta-d') (Pearson correlation, r = 0.8733,
$p < 0.001$).

Figure 4. High gamma power in the hippocampus is associated with d' but not with meta-d'.
(a) Left panel: The time course of the difference in high gamma power (z-score) of the hippocampus between high confidence trials and low confidence trials. Right panel: The time course of the difference in high gamma power (z-score) of the hippocampus between correct trials and incorrect trials. The solid line represents the mean, with SEM (shaded areas) across electrode sites (n=24). The horizontal black rectangle on the x-axis represents significant time points (cluster-based permutation, p < 0.05). The vertical shaded areas highlight the time window of interest used in **(b)**.
(b) The left panel illustrates the correlation between the neural accuracy index ($H\gamma_{Correct} - H\gamma_{Incorrect}$) and behavioral type 1 sensitivity (d') (Pearson correlation, r = -0.6194, p = 0.0421), whereas the right panel depicts the correlation between the neural interaction index [($H\gamma_{High\ confidence} - H\gamma_{Low\ confidence}$) - ($H\gamma_{Correct} - H\gamma_{Incorrect}$)] and behavioral type 2 sensitivity (meta-d') (Pearson correlation, r = -0.4056, p = 0.2158).

636
Figure 5. Correlation analysis of high gamma activity between the precuneus and the
hippocampus and its relationship with metacognitive indices. (a) Schematic diagram of
correlation analysis procedure. We defined a 0.02s length window, step by 0.01s, and calculated
Person correlation in each window. Segments in which 4 or more consecutive windows (0.04s, at
least 2 cycles of high gamma) showing significant correlation are considered high correlation. We
calculated the count of highly correlated occurrences between the hippocampus and the precuneus.
(b) The probability density of one representative participant comparing shuffled and original data
is presented ($p < 0.001$, permutation test, n=1000), see also Figure S2. **(c)** Count of correlation of
original and shuffled data (Wilcoxon matched pairs signed rank test, $p < 0.0001$) taking 82 electrode
pairs electrodes from all participants into account. **(d)** Count correlation in high confidence trials
compared with low confidence trials (Wilcoxon matched pairs signed rank test, $p < 0.05$). **(e)** Count
correlation in correct trials compared with incorrect trials (Wilcoxon matched pairs signed rank test,
$p < 0.0001$). **(f)** Neural Interaction index [(Count $_{High\ confidence}$ − Count $_{Low\ confidence}$) - (Count $_{Correct}$ −
Count $_{Incorrect}$)] of high correlation count across all electrodes against zero (one-sample t-test, $p <$
0.0001). SEM is across all electrode pairs.

Figure 6. The time course of correlation coefficient of high gamma between the precuneus and the hippocampus towards confidence level reporting. (a) The time course of correlation coefficient before confidence rating in all trials. The solid line represents the mean with SEM (shaded areas) across all subjects (n=11). The horizontal black rectangle on the x-axis represents significant time points (cluster-based permutation, $p < 0.05$). The yellow and green vertical shaded areas represent the time window of interest used in **(b)** and **(c)** respectively. **(b)** The upper panel illustrates the correlation between behavioral type 1 sensitivity (d') and neural accuracy index (H-P $_{Correct}$ − H-P $_{Incorrect}$) of H-P correlation coefficient (Pearson correlation, $r = 0.4415$, $p = 0.1741$), whereas the lower panel depicts the correlation between behavioral type 2 sensitivity (meta-d') and neural interaction index $[(H\text{-}P_{\text{High confidence}} − H\text{-}P_{\text{Low confidence}}) - (H\text{-}P_{\text{Correct}} − H\text{-}P_{\text{Incorrect}})]$ of H-P correlation coefficient (Pearson correlation, $r = 0.5283$, $p = 0.0948$). **(c)** The upper panel illustrates the correlation between behavioral type 1 sensitivity (d') and neural accuracy index (H-P $_{Correct}$ − H-P $_{Incorrect}$) of correlation coefficient (Pearson correlation, $r = 0.2143$, $p = 0.5269$), whereas the lower

panel depicts the correlation between behavioral type 2 sensitivity (meta-d') and neural interaction index $[(H\text{-}P_{\text{High confidence}} - H\text{-}P_{\text{Low confidence}}) - (H\text{-}P_{\text{Correct}} - H\text{-}P_{\text{Incorrect}})]$ of H-P correlation coefficient (Pearson correlation, r = -0.6154, p = 0.0439).

METHODS AND MATERIALS

Participants

Eleven patients diagnosed with intractable epilepsy underwent implantation of subdural intracranial electrodes for epilepsy treatment at The Second Affiliated Hospital, Zhejiang University School of Medicine (SAHZU). The patients' ages ranged from 12 to 39 (mean=24.91 years, SD=9.63). All patients demonstrated a good understanding of the experimental procedure and had at least 1 electrode recording signals in each of the target brain regions (hippocampus and precuneus). No clinical seizures were observed during the course of the experiment. This study was approved by the SAHZU committee on Human Research.

Experimental task

Each trial began with a video of animals, which lasted for 5-6 s. These videos were validated in previous studies (e.g., Zuo et al., 2020). Subsequently, at the memory test, the participants had to determine which image extracted from the video was presented earlier within a 10-second timeframe (temporal order judgment, TOJ). Following the memory test, a fixation cross was presented for 5 s. In the final phase, participants reported their confidence level on a scale ranging from 1 to 4. The number of trials completed by each participant varied, with a range of 120-240 trials (4-6 sessions, 30-40 trials per session). Initiation of each trial was self-paced and the participants were allowed to take breaks between sessions. To exclude epileptic activity, we only included patients who did not experience seizures or pre-seizure activity within 24 hours.

Intracranial recordings and electrode localization

Recording sites in depth electrodes implanted in the hippocampus and precuneus were 2 mm platinum cylinders with 1.5 mm intercontact spacing and a diameter of 0.8 mm (Fig.1b). Before electrode implantation, T1-weighted Magnetic Resonance Imaging

(MRI) using a 3.0-T scanner and computed tomography were obtained for each patient.

The locations of the electrodes were determined based on clinical standards, utilizing

the SINOVATION software. We analyzed only electrodes localized to the

hippocampus and precuneus in this study. Electrode localization was calculated using

the Montreal Neurological Institute (MNI) coordinates, with the AAL3 atlas applied for

further visualization. The number of electrodes for each patient is shown in the

Supplementary table S1.

Preprocessing

All data analyses were performed in MATLAB 2020a using EEGLAB, procedure from

GitHub and Self-defining script. The sample rate of signal was at 2000 Hz. First, we

extracted markers of each event, excluding trials with missing or wrong markers from

further analysis. We selected electrodes from the same region based on MNI

coordinates before bipolar referencing (either the hippocampus or precuneus, see

Supplementary Table1). Then bipolar reference was applied to raw data of

hippocampus and precuneus by measuring the difference between adjacent recording

sites (i.e., 1.5 mm apart due to the electrode design). We used FIR band-stop filters to

remove 50-Hz and 100-Hz power line interference. We focused on three events in the

experimental task: 1. TOJ test (10s, retrieval); 2. Rest/Fixation (5s); 3. Report

confidence level (4s, confidence rating).

Data analysis

Generalized Linear Model (GLM).

To determine whether the neural signals are associated with metacognition, the time-

series power (P) was regressed against several relevant factors (R). These factors

included confidence level, confidence reaction time, response accuracy

(correct/incorrect), retrieval reaction time, touch side (left/right), and video category:

$$P = \alpha + \beta_1 R_1 + \beta_2 R_2 + \beta_3 R_3 + \beta_4 R_4 + \beta_5 R_5 + \beta_6 R_6$$

To assess the contribution of the different frequency bands to the confidence rating,

confidence levels(R) were regressed separately across trials against normalized

envelope of power P (Lopez-Persem et al., 2020) for each frequency band (delta: 1–4Hz, theta: 4–8Hz, alpha: 8–14 Hz, beta: 14–35 Hz, low-gamma: 35-70 Hz, high-gamma: 70-150 Hz), averaged over time between 0 and 1.5 s before reporting confidence level:

$$R=\alpha+\beta P$$

with α corresponding to the intercept. The significance of the regression estimates β was assessed across recording sites using two-sided, one-sample, Student's t-tests.

To further determine whether other frequency bands provided additional information, we compared the following 7 GLMs:

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma)$$

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma) +\beta_{\theta}P(\theta)$$

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma)+ \beta_{\alpha}P(\alpha)$$

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma) +\beta_{\beta}P(\beta)$$

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma) +\beta_{\theta}P(\theta) + \beta_{\alpha}P(\alpha)$$

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma) +\beta_{\theta}P(\theta) + \beta_{\beta}P(\beta)$$

$$R=\beta_{H\gamma}P(H\gamma) + \beta_{\gamma}P(\gamma) + \beta_{\alpha}P(\alpha) + \beta_{\beta}P(\beta)$$

The model comparison was conducted using the Variational Bayesian Analysis (VBA) toolbox. Log-model evidence obtained in each recording site was taken to a group-level, random-effect, Bayesian model selection (RFX-BMS) procedure (Rigoux et al., 2014; Stephan et al., 2009). The RFX-BMS provided an exceedance probability (Xp) that measures the likelihood of a given model being more frequently implemented relative to all the others considered in the model space in the population from which samples are drawn.

Correlation Analysis. We used Pearson correlation to examine the relationship of high gamma activity between the hippocampus and precuneus. For correlation count, we firstly extracted the high-gamma (70-150Hz) band from both the hippocampus and precuneus. We then defined a window length of 0.02s, step by 0.01s (50% overlapping), and calculated the Pearson correlation coefficient for each window. To ensure the sensitivity and specificity of the results, we considered the signals to be relevant only when all 4 windows (equivalent to 0.04 s, at least 2 cycles of high gamma) met the

minimum criteria of Pearson correlation (r > 0.8, p < 0.05). The occurrence of highly

relevant signals during confidence rating was counted. We quantified the count of high

correlation bands within the whole confidence rating stage. As a control, we shuffled

the trials to obtain a baseline for each participant for statistical inference. The shuffling

criteria were as follows: 1) random matching of signals from different trials, 2) a

minimum interval of 10 trials between shuffled signals. Finally, we applied permutation

test for origin data and shuffled data to detect significance (n=1000). As for the H-P

correlation coefficient within time series, a window width of 0.02 s was also utilized to

compute the Pearson correlation coefficient of high gamma frequency bands at each

time window, averaging it according to the 0.1s window.

Measurement of metacognitive indices. We computed type 1 sensitivity (d') and type 2

sensitivity (meta-d'), wherein meta-d' indicates metacognitive sensitivity (the ability to

discriminate between correct and incorrect judgments) and d' represents memory

performance according to signal detection theory (Green and Swets, 1966; Maniscalco

and Lau, 2012). We divided within-subjects trials by their confidence levels into two

categories for analysis: High confidence (3-4), Low confidence (1-2). In neural data,

we computed an interaction index [(Neural data High confidence − Neural data Low

confidence) - (Neural data Correct − Neural data Incorrect)] to reflect metacognitive

process and an accuracy index (Neural data Correct − Neural data Incorrect) to reflect

memory process.

Power Spectral Density (PSD). During the confidence rating stages, we calculated the

power spectral density (PSD) using the Welch periodogram method (pwelch function

in MATLAB) for the 1 s interval preceding the confidence rating. A Hanning window

of 0.1 s with 50% overlap was applied to minimize edge effects. Subsequently, we

calculated the z-score for each electrode, followed by the computation of the accuracy

index ($H\gamma_{Correct} - H\gamma_{Incorrect}$) and interaction index [($H\gamma_{High\ confidence} - H\gamma_{Low\ confidence}$) -

($H\gamma_{Correct} - H\gamma_{Incorrect}$)].

Time–frequency analysis. Time–frequency analyses were carried out using the Morlet wavelet transform method (cwt function in MATLAB) with default scale. We then computed the corresponding T values for each electrode.

ACKNOWLEDGEMENT

We extend our gratitude to the patients for their cooperation, as well as to the medical staff involved in the clinical treatment and the technicians in the EEG laboratories for their valuable contributions.

AUTHOR CONTRIBUTIONS

H.J.J and S.C.K. conceived the study and designed the experiment. K.P, X.X.G and H.Y.Y analyzed the data. J.M.Z supervised the surgeries and experiments. K.P and Z.L.Z. ran the experiments. Z.Z., J.M.Z, and H.J.J performed the surgeries and contributed to electrode localization. K.P and H.J.J wrote the paper. H.W, Y.C, D.S-P, and S.C.K further interpreted the data and contributed to the writing by reviewing and editing the manuscript.

COMPETING INTERESTS

The authors declare no competing financial interests.

FUNDING

S.C.K. received support from the National Natural Science Foundation of China (grant number: 32071060) and Jiangsu Provincial Department of Science and Technology (grant number: BK20221267).