Ontogenetic variation in the marine foraging of Atlantic salmon functionally links genomic diversity with a major life history polymorphism

Running title: Functional studies of the genotype-phenotype map of Atlantic salmon

Author names and affiliations:

Tutku Aykanat[1], Jan Arge Jacobsen[2], Kjetil Hindar[3]

1. Organismal and Evolutionary Biology Research Program, Faculty of Biological and Environmental Sciences, PO Box 56, FI-00014 University of Helsinki, Finland.
2. Faroe Marine Research Institute, Nóatún 1, FO-100 Tórshavn, Faroe Islands.
3. Norwegian Institute for Nature Research (NINA), NO-7485 Trondheim, Norway.

Keywords: Feeding strategy, Atlantic salmon, *vgll3*, *six6,* genomic diversity, life-history evolution, polymorphism

Abstract

The ecological role of heritable phenotypic variation in free-living populations remains largely unknown. Knowledge of the genetic basis of functional ecological processes can link genomic and phenotypic diversity, providing insight into polymorphism evolution and how populations respond to environmental change. By quantifying the marine diet, of sub-adult Atlantic salmon, we assessed how foraging behavior changes along the ontogeny, and in relation to genetic variation in two loci with major effect on age-at-maturity (*six6* and *vgll3*). We used a two-component, zero-inflated negative binomial model to simultaneously quantify foraging frequency (zero-inflation components) and foraging outcome (count component), separately for fish and crustaceans in the diet. We found that older salmon forage for both prey types more actively (as evidenced by increased foraging frequency), but with a decreased efficiency (as evidenced by fewer prey items in the diet), suggesting an age-dependent shift in foraging dynamics. The *vgll3* locus was linked to age-dependent changes in foraging behavior: younger salmon with $vgll3^{LL}$ (the genotype associated with late maturation) tend to forage crustaceans more often than those with $vgll3^{EE}$ (the genotype associated with early maturation), while the pattern was reversed in older salmon. $Vgll3^{LL}$ genotype was also linked to marginal increase in fish acquisition especially in younger salmon, while *six6* was not a factor explaining the diet variation. Our results suggest a functional role for variation in marine feeding behavior linking genomic diversity at *vgll3* with age-at-maturity among salmon, with potential age-dependent trade-offs maintaining the genetic variation. A shared genetic basis between dietary ecology and age-at-maturity likely subjects Atlantic salmon populations to evolution induced by bottom-up changes in marine productivity.

Introduction

Phenotypic polymorphisms are ubiquitous, discrete trait morphs (e.g., (Bataillon et al., 2022; Funk et al., 2021; Janssen, Bustnes, & Mundy, 2021; Johnston et al., 2013), the genetic basis of which are maintained at higher frequencies than what would be expected under mutation-selection balance (Ford, 1945). Much of the research on phenotypic polymorphism focuses on determining which selective landscapes generate and maintain heritable polymorphisms, such as negative frequency-dependence (Chouteau, Arias, & Joron, 2016; Goldberg et al., 2020; Gross, 1985; Svensson, Willink, Duryea, & Lancaster, 2020), antagonistic pleiotropy (Charmantier, Perrins, McCleery, & Sheldon, 2006; Marden, Langford, Robertson, & Fescemyer, 2021), fluctuating selection (Bonnet & Postma, 2018; Grant & Grant, 2002) and, heterozygote advantage (Coulson et al., 2011; Hedrick, 2012). Furthermore, by the advent of next-generation sequencing, an increasing number of studies report associations between discrete chromosomal elements (large-effect loci, inversions, supergenes) and phenotypic polymorphisms (Barson et al., 2015; Funk et al., 2021; Johnston et al., 2013; Matschiner et al., 2022). Such an intense research focus is unsurprising given that studying polymorphisms provides insight into speciation (Jamie & Meier, 2020), one of the cornerstones of biology since Darwin. Despite the increasing number of known genotype-phenotype associations, how such genomic diversity is functionally translated into phenotypic diversity within wild populations is mostly unresolved. This is likely due to the experimental studies and sampling designs required for monitoring wild individuals being financially and logistically unfeasible without applying various simplifying assumptions (Mousseau, Sinervo, & Endler, 2000; Reznick, 2016). As such, our understanding of how genomic

variation influences individual-level differences in ecology, and how such ecological differences

might covary with polymorphic trait states, is incomplete.

One well studied genotype-polymorphism association is that between age-at-maturity and the

vgll3 and *six6* haploblocks in Atlantic salmon, *Salmo salar* (Ayllon et al., 2015; Barson et al.,

2015). Atlantic salmon most often display an anadromous life history. They hatch in rivers where

they can spend several years growing before migrating to the ocean. The marine environment is

much more productive than fresh water (Gross, Coleman, & McDowall, 1988) and juvenile

salmon undertake this costly migration in order to maximize growth and lipid storage. This, in

turn, supports future reproductive efforts when they return to fresh water to spawn (Esteve,

2005; Ferguson, Reed, Cross, McGinnity, & Prodohl, 2019). Atlantic salmon most commonly stay

at sea for one to three years prior to return to fresh water for reproduction, which is termed as

sea-age at maturity. Atlantic salmon exhibit a high degree of divergence in this polymorphic

phenotype within and among populations, the genetic variation of which is largely explained by

two genomic regions, so called *vgll3* and *six6*, after the most prominent genes within the

haploblock (Barson et al., 2015; Sinclair-Waters et al., 2022). The *vgll3* gene encodes for a

functionally-conserved transcription co-factor that regulates adiposity in mice (Halperin, Pan,

Lusis, & Tontonoz, 2013) and influences pubertal timing in vertebrates (Cousminer et al., 2013).

The functional relationship between adiposity and maturation via *vgll3* haploblock likely

operates also in Atlantic salmon since condition factor has been shown to drive *vgll3* dependent

maturation response in the species (Debes et al., 2021). Salmon with the 'early' *EE* genotype are

more likely to mature after one year at sea (one-sea-winter salmon, 1SW) than individuals with

the 'late' *LL* genotype, which are more likely to spend multiple years at sea before maturing

(multi-sea-winter, MSW, predominantly composed of 2SW and 3SW salmon). The second major

haploblock, *six6*, displays substantial population-level effects on sea-age at maturity among wild

salmon populations (Barson et al., 2015), and within aquaculture strains (Sinclair-Waters et al.,

2022). The *six6* haploblock contains genes with known functional roles in sensory and digestive

system development (Lee et al., 2012; Moustakas-Verho et al., 2020). The molecular functions

associated with *vgll3* (adiposity regulation) and *six6* (sensory and digestive system development)

suggest that dietary ecology may underpin processes by which genomic variation at the *vgll3* and

six6 haploblocks is expressed as phenotypic variation in sea-age at maturity.

Variation in how food is acquired by organisms and how derived metabolic energy is invested

into different biological processes can be conceptualized using the resource acquisition-

allocation framework (Enberg et al., 2012). Briefly, individuals acquire a given amount of a

resource from which they service their basal metabolism. Any surplus resources are then

allocated to behavior, structural (skeletal/muscle) growth, or maturation/gonadal growth

(Enberg et al., 2012). A supposition arising from the resource acquisition-allocation framework is

that individuals vary in how much/which resources they acquire, and this may alter their

maturation trajectories. As such, variation in feeding opportunities and strategies may play a role

in survival and maturation of the organisms.

Atlantic salmon is recognized as an opportunistic feeder with extensive ontogenetic and

spatiotemporal variation in diet breadth (Rikardsen & Dempson, 2010). Salmon feed increasingly

on higher trophic levels as they age, which is strongly correlated to their size (Aykanat et al.,

2020; Jacobsen & Hansen, 2001; Salminen, Erkamo, & Salmi, 2001; Salminen, Kuikka, & Erkamo,

1995). The variation in available prey items, as well as preference for particular prey items, can

affect the net energy gain from foraging activities, and this may subsequently alter the sea-age at

which they mature (Czorlich, Aykanat, Erkinaro, Orell, & Primmer, 2022; Hvidsten et al., 2009;

Vollset et al., 2022). For example, Vollset et al. (2022) showed that the reduced marine growth of

Atlantic salmon (and subsequent decrease in maturation age) was linked to reduction in the

Arctic cold water flux to the Northeast Atlantic Ocean and the subsequent decrease in the

plankton biomass in the feeding grounds in the Northeast Atlantic Ocean. Similarly, using a 40

year time series from large salmon population, Czorlich et al. (2022) linked the evolutionary

response in the *vgll3* genotype towards early maturation to changes in Capelin abundance in the

Barents Sea, a major prey for Atlantic salmon in the region. On this background, one can posit

that genetic variation in ontogeny dependent foraging behavior can be maintained in the

populations via life-history trade-offs. For example, salmon carrying alleles associated with a

foraging behavior that provides higher net-energy gain at early stages at sea will be more likely

to acquire enough resources to initiate maturation, and mature at an earlier age (Chaput &

Benoit, 2012; Thorpe, Mangel, Metcalfe, & Huntingford, 1998). However, demonstrating a genetic

basis to foraging behavior, and how this might covary with sea-age at maturity, remains highly

challenging in salmon due to a lack of adequate samples in the wild. As such, whether foraging

behavior varies as a function of genotype, and the evolutionary consequences that this variation

could have on the sea-age at maturity phenotype, remains largely unknown. One previous study

has demonstrated a relationship between the L allele of *six6* and increased stomach content (that

is composed of fish) among maturing Atlantic salmon sampled on their return migration along

the Norwegian coast (Aykanat et al., 2020). However, a shared genetic basis between sea-age and

foraging among sub-adult, immature fish (i.e. fish that are still in the oceanic feeding phase,

before the onset of maturation) would be an indirect evidence for dietary ecology acting as a

potential process that functionally links genomic diversity at the *vgll3* and *six6* loci with sea-age

at maturity in salmon.

An extensive dataset of the diet of wild Atlantic salmon sub-adults (i.e., the marine stage prior to

maturation) sampled in the oceanic feeding ground (Jacobsen & Hansen, 2001) provided us with

the material to test if ontogenetic variation in foraging behavior is explained by sea-age at

maturity loci (*vgll3 – six6*) in Atlantic salmon. The Atlantic salmon in the study almost exclusively

feed on crustaceans and fish, the former of which was more numerous (∼95% of diet items were

crustaceans) but comprised only 30% of the content by weight. We propose that a genetic basis

in feeding behavior results in variation in the net-energy gain, which subsequently alters

individuals' probability of reaching to maturation threshold. The salmon diet in our data has a

right-skewed distribution with a substantial zero inflation, which we presumed to be shaped by

two independent biological processes. The first process explains the "structural-zeros" in the

data (i.e. zero inflation), which is used as a proxy to measure the active foraging frequency for

particular diet types. The second process is the "count" component, and is used as a proxy to

measure the foraging outcome providing that salmon actively pursue for a prey (see Materials

and Methods for details). Thus, as a first step, using the a zero-inflation model to account for

these two processes, we tested if foraging behavior changes along the ontogeny, and secondly, if

genomic regions associated with sea-age at maturity in salmon, in part, explain the variation in

foraging behavior, across age groups. Within the same framework, we also estimated if the

variation in diet is also determined by age-dependent sex effects, since Atlantic salmon also

exhibited sex dependent maturation pattern (Barson et al., 2015; Fleming & Einum, 2010).

Methods

Sampling procedure

Sampled fish used in this study are originally described in Jacobsen and Hansen (2001). Briefly, Atlantic salmon were sampled from known marine feeding grounds around the Faroe Islands, inhabited by all sea-age groups but predominately by MSW fish (Jacobsen & Hansen, 2001; O'Sullivan et al., 2022). Fish were sampled between November and March (excluding January due to rough sea conditions) during the 1992-'93 and 1993-'94 fishing periods. Salmon sampled from November-December were considered to have been caught in the autumn and those sampled from February-March in the winter, which exhibited marked differences in prey diversity, sea surface temperature and stomach content (Jacobsen & Hansen, 2001). The locations of the longlines used to sample fish were also recorded (n = 55, see Fig. S1 in O'Sullivan et al., 2022). Individual salmon had their fork lengths measured, were gutted, and gutted fish were weighed (in kilograms). Gut contents were also removed, identified, and enumerated. Fish scales were taken from each individual salmon for determination of sea-age via scale reading, and for later, DNA extraction for genotyping. The fish sampled in the study are immature, sub-adults with no phenotypic indication of maturation. However, it is reasonable to assume that virtually all 3SW fish will mature in the following autumn, since 4SW and above sea-ages are rare in salmon populations.

Dietary quantification and somatic measurements

The stomach content data used in this study is a subset of the data that was quantified by Jacobsen and Hansen (2001). We did not include data from the 1994-'95 fishing period in Jacobsen and Hansen (2001), since material for genotyping was not available. The most common

diet items among crustaceans were amphipods (e.g., *Themisto* spp.), shrimps (e.g., *Hymenodora*

glacialis) and euphausiids (e.g., *Meganyctiphanes norvegica*), and among fishes were pearlsides

(e.g., *Maurolicus muelleri*), lanternfishes (e.g., *Benthosema glaciale* and Myctophidae),

barracudinas (e.g., *Notolepis rissoi* and Paralepidae), blue whiting (*Micromesistius poutassou*), and

capelin (*Mallotus villosus*) (Supplementary table 1). Crustaceans were substantially more

numerous than fishes (~95% of diet items were crustaceans) but represented only 30% of the

content by weight (Supplementary table 1). For example, the average size of crustacean species

ranged from 0.027 g for *Themisto compressa* to 0.766 for *Sergestes arcticus*, and > 99% were

species that were smaller than 0.20 g. The average size of fish subcategories ranged from 0.26 g

for fry (mostly consisting of capelin) to 72.0 g for herring, *Clupea harengus* (Supplementary table

1). For the purposes of this study, and to reduce the complexity of data structure and statistical

analyses, the stomach contents were primarily categorized as crustaceans and fishes.

Apart from than crustaceans and fishes, the diet also contained remains from fish and

crustaceans, and organic remains, which were mostly not possible to further identify or

enumerate, (Supplementary table 1). In addition, a few stomachs contained bird remains (n=1)

and squid (n=11). Fish and crustacean remains were included in the count-data analysis if

numerated (n=12 and n=16, respectively). Squids were grouped together with other fishes due to

their size similarity (excluding them from the analysis did not change any results). One fish with

bird remains was excluded from the analysis.

Genotyping and data filtering

DNA was extracted from the sampled fish scales (O'Sullivan et al., 2022). Individuals were

genotyped using a GTSeq approach (Campbell, Harmon, & Narum, 2015) with a panel of 167

SNPs as outlined in (Aykanat et al., 2020). This panel allowed individuals to be assigned to their

reporting group of origin (O'Sullivan et al., 2022), as well as to score the genotypes at focal SNPs

within our target loci. The focal SNPs were $vgll3_{TOP}$ and $six6_{TOP.LD}$ (hereafter, $vgll3$ and $six6$).

$Vgll3_{TOP}$ exhibits the strongest association with sea-age within the haploblock (Barson et al.,

2015), while $six6_{TOP.LD}$ exhibits the second strongest association overall with sea-age prior to

population structure correction (Barson et al., 2015) . If an individual was missing a genotype for

$vgll3_{TOP}$, the missing genotype was inferred from the genotypes of adjacent SNP markers in the

same haploblock, which display strong linkage disequilibrium due to close physical proximity to

$vgll3_{TOP}$ (The loci with the strongest linkage with $vgll3_{TOP}$ was prioritized when imputing the

missing genotype, with an order of $vgll3_{missense1}$, $vgll3_{missens2}$, $akap11$, see, Aykanat, Lindqvist,

Pritchard, & Primmer, 2016). Sex was determined genetically by targeting the sex determination

locus (sdy, Yano et al., 2013), as outlined in Aykanat et al. (2016). There was 82.3 % concordance

between phenotypic and genetic sex, of which, we prioritize genetic sexing over phenotypic

sexing since sub-adults at sea do not yet develop secondary sexual characteristics, hence

phenotypic sex may be prone to misidentification. For a small subset (n = 39), the genetic sex of

an individual could not be determined, and phenotypic sex was used instead. Individuals lacking

genotypic information for either the $vgll3$, $six6$ loci, or sex, were excluded from downstream

analyses (n = 77). A small number of fish with sea-age greater than 3SW were also excluded from

the data (n=15). Reporting group was included as a three-level categorical variable, in which an

individual is assigned to a reporting group with the highest assignment probability, where levels

corresponded to large-scale geographical regions based on the genetic stock identification in

O'Sullivan et al. (2022). These reporting groups were Ireland and the UK (n = 473), Southern

Norway (n = 832), and the Northern group (n = 271), the latter which consisting of fish from

Northern Norway, including Teno River, Eastern Finnmark/Northern coast of Kola, and the

Barents/White Sea reporting groups. In total, 1536 individual salmon consisting of 168 1SW, 950

2SW and, 418 3SW individuals were used in the downstream analyses.

Statistical analyses

We modelled the total number of crustaceans and the total number of fishes in the stomachs of

individual salmon as a pair of separate two-component zero-inflation models, for which 46.1 and

79.9 % of stomachs had no crustaceans and fishes, respectively (Supplementary Figure 1,

Supplementary Table 2). Zero-inflation models simultaneously estimated two independent

processes that shapes the distribution of the response variable (diet in the stomach). The first

process was modelled with the zero inflation component (using binomial error distribution), and

quantified for the "structural-zeros" (i.e., inflated zero counts) in the data that were due to

individuals not actively foraging for the specific prey type. We used this statistic as a proxy to

measure the active foraging frequency. The second process is modelled with the count

component (using a negative binomial error distribution that includes the "random zeroes" in the

data) that were due to sampling variation when salmon actively forage for prey. We used this

statistic as a proxy to measure the outcome of the active feeding, i.e. foraging outcome.

The model structure (for both components) is as follows:

$Y = \mu + Season + fishing\ period + Reporting\ Group + Sea\text{-}Age \times (Sex + vgll3 + six6 + L_{age.std}) +$
$\varepsilon_{location} + \varepsilon_{error}$

whereby, for both for crustacean and fish models, season (autumn, winter), fishing period (1992-

'93, 1993-'94), and reporting group were included as categorical fixed effects. The *vgll3* and *six6*

genotypes, as well as sex and age-standardized length ($L_{age.std}$) were modelled interacting with

sea-age, since we quantify a sea-age-dependent diet response associated with these variables. For

the crustacean diet model, sea-age was parameterized as a continuous variable, which was more

parsimonious than models that parameterize sea-age as a categorical variable (see results). For

the fish diet model, sea-age was also parameterized as a continuous variable, but comparing

continuous vs categorical parameterization was not possible due to convergence problems with

the latter (see results). For both crustaceans and diet models, the *vgll3* and *six6* genotypes were

modelled additively (EE, EL, and LL genotypes were coded numerically by 1, 2, and 3) which

were more parsimonious than non-additive models (i.e., when genotype values were

categorically modelled. See results.). All covariates interacting with sea-age (*vgll3*, *six6, sex, and*

$L_{age.std}$) were scaled by their respective means and standard deviations to improve convergence,

and interpretability of the coefficient tables. Spatial variation in sampling effort (sampling

location, N=55) was included in the model as a random effect.

In order to evaluate if any taxonomic groups within crustaceans drive the relation between

genotypes and diet, we further broke down the response variable into lower order taxonomic

groups when numbers within specific taxa were sufficiently large. For that, we reanalyzed the

data using Hyperiid amphipods (*Themisto* spp.) and Euphausiids as the response variable, which

comprised the majority of the crustacean diet in the stomach (Supplementary Table 1). The

representative sizes in the diet between these two groups were also different, with the latter

being substantially larger than the former (Supplementary Table 1). We did not implement the

same approach in fishes which had restrictively small number of prey items per species.

All models were run using the *glmmTMB* package (version 1.1.5, Brooks et al., 2017) in R version

4.1.0 (R Core Team, 2019), using a negative binomial error structure with a logit link function for

the count component, and with a binomial error structure with a logit link function for the zero-

inflation (zi) component, using default optimizer settings. Marginal means for genotype effects

and their confidence intervals were calculated using the *emmeans* package (version, 1.8.2, Lenth,

2023). Model diagnostics were carried out using *DHARMa* package in R (version 0.4.5, Hartig

2022), whereby randomized quantile residuals were simulated for both models in order to

determine if any explanatory covariate displayed heteroscedasticity (Hartig 2022). Using the

aforementioned two-component models, we primarily asked whether sub-adult foraging

behavior varies as a function of life history genotype and age. We also tested if sex had an age-

dependent effect on adult foraging behavior.

271 Results

272 *Crustaceans model*

Sea-age: Based on the zero-inflation models, older sea-age groups more likely forage for

crustaceans (zi component: z = -3.196, *P* = 0.001), but younger sea-age groups had substantially

higher number of crustaceans in the stomach, indicating a higher foraging outcome (count

component: z= -4.478, *P* < 0.001, Supplementary Table 3, Figure 1).

Vgll3 and six6: Genetic variation in the *vgll3* locus was strongly associated with an age-dependent

change in the crustacean foraging frequency (zi component: z = 2.690, $P_{\text{sea-age x } vgll3}$ = 0.007, Figure

1a, Supplementary Table 3). As such, *vgll3*^{LL} genotype was insignificantly associated with

increased foraging in younger sea-age groups (Odds ratio = 2.33, 95% CI = 0.78 – 7.04, *P* = 0.132,

Figure 1d), but in the older sea-age groups, the alternative, *vgll3*^{EE} genotype, was significantly

associated with increased foraging for crustaceans (for 2SW, odds ratio = 2.04, 95% CI = 1.04–

3.98, *P* = 0.038; for 3SW, odds ratio = 9.67, 95% CI = 2.15– 43.57, *P* = 0.003, Figure 1a). The

difference appeared to be driven by the *vgll3*^{E} allele, which exerts a significant age-dependent

response to crustacean's foraging frequency in the diet (z = -4.081, $P_{\text{sea-age}|vgll3EE}$ < 0.001), while

the $vgll3^L$ allele had no age-dependent effect (z = - 1.112, $P_{\text{sea-age}|vgll3EE}$ = 0.507, Figure 1a).

However, $vgll3$ genotypes did not alter the foraging outcome, as there was no difference in in the

abundance of crustaceans in salmon diet in relation to $vgll3$ genotype (Figure 1d).

Figure 1: Crustaceans acquisition patterns of sub-adult Atlantic salmon as a function of sea-age-dependent genotype

($vgll3$ and $six6$) and sex. Upper and lower panels indicate marginal means for foraging frequency (zero-inflation

component), and foraging outcome (count component), respectively. Error bars are 95% confidence intervals.

Crustacean diet as a function of $six6$ locus was insignificant for both foraging frequency (zero-

inflation components), and foraging outcome (zero-inflation components, Figure 1b,e,

Supplementary Table 3).

Sex: Sex exhibited a strong age-dependent association with foraging frequency (zi component: z =

-2.822, _P_ = 0.005, Figure 1c), but did not alter the foraging outcome (count component: z = -0.703,

_P_ = 0.482, Figure 1f). Female salmon forages for crustacean more often than males in the 1SW

group (Odds ratio = 3.27, 95% CI = 1.46 – 7.31, _P_ = 0.0042), whereas the pattern reversed in the

older sea-age groups and the 3SW males forage crustaceans more often than the 3SW females

(Odds ratio = 3.55, 95% CI = 1.13 – 11.21, _P_ = 0.031, Figure 1c). The difference appeared to be

driven by greater age-dependent response by males (z = -4.297, $P_{\text{sea-age x females}}$ < 0.001), compared

to females (z = -1.489, $P_{\text{sea-age x males}}$ = 0.296, Figure 1c).

Season, year, reporting group and length effects: Season was an important determinant for

crustacean diet. Salmon more likely forage for crustaceans in the winter (zi component: _z_ = -

2.439, _P_ = 0.015), but had overall lower abundance of prey in the diet (count component: _z_ = -

5.107, _P_ < 0.001, Supplementary Table 3). Fishing period did not have a significant effect on

foraging frequency yet the number of crustaceans in the diet was significantly different between

the two sampling years (count component: _z_ = 5.512, _P_ < 0.001, Supplementary Table 3).

Reporting group generally did not affect variation in the crustacean diet, but fish assigned to the

Northern Norway reporting group had marginally fewer crustaceans in the diet than the fish

assigned to Ireland and the UK reporting group (_z_ = -2.504, _P_ = 0.037, Supplementary Table 3).

Age standardized length as a function of sea-age ($L_{\text{age.std}}$) was not a significant factor for the

crustaceans diet (Supplementary Table 3). Sequential exclusion of these variables from the model

(Season, fishing period, reporting groups, and $L_{\text{age.std}}$) did not meaningfully influence the

genotype or sex effects observed in the crustacean diet (data not shown).

Alternative parameterization of variables: When crustaceans diet was subdivided into lower

taxonomic orders, two major groups (Hyperiid amphipods and Euphausiids) appeared to exhibit

similar effects and effect sizes (and similar to the overall crustaceans model) in the *vgll3* locus,

albeit not always significant due to lower sample sizes, (Supplementary Table 4 and 5),

suggesting neither a particular order nor size drives the age-dependent genetic or sex relation.

In the implemented framework, we modelled the life-history loci and sea-age as a continuous

variable, assuming additivity for the latter, and linear relation of the response variable with the

former. This model structure was substantially more parsimonious than models that assume

non-additivity (where genotypes were modelled as categorical variables), and/or non-linear age-

dependent response (where sea-age was modelled as a categorical variable, Supplementary

Table 6). Similarly, a model that accounts for sex and *vgll3* interaction, which was previously

suggested as a mechanism for a sex dependent genetic effect on sea-age at maturity (Barson et al.,

2015), was not supported any better (data not shown).

Convergence and diagnostics: Convergence was achieved in all models enabling us to compare

different parameterizations of sea-age and genotype. Overall, the diagnostics for the crustaceans

model suggested a good model fit (Supplementary Figure 2a).

Fish model

Sea-age: Similar to the crustaceans model, older sea-age groups more likely forage for fish (zi

component: $z = -3.704$, $P < 0.001$), and younger sea-age groups exhibited higher foraging

outcome (count component: $z = -4.204$, $P < 0.001$, Supplementary Table 7, Figure 2).

Vgll3 and six6: Frequency of foraging for fish was generally insignificant for *vgll3* and *six6* loci.

The foraging outcome, however, was significantly higher for *vgll3*^{L} allele (count component: $z =$

2.088, $P = 0.037$, Supplementary Table 7), whereby the effect was diminished in older age-groups

due to a marginal age-dependent effect (count component: $z = -1.796$, $P_{\text{sea-age x } vgll3} = 0.073$,

Supplementary Table 7). Salmon with *vgll3*[LL] genotype had more fish in the stomach than salmon

with *vgll3*[EE] genotype in 1SW and 2SW sea-age groups (for 1SW, odds ratio = 3.36, 95% CI = 1.15

– 9.81, *P* = 0.027; for 2SW, odds ratio = 1.65, 95% CI = 1.01 – 2.71, *P* = 0.047), but the difference

was negligible in the 3SW age group (odds ratio = 0.81, 95% CI = 0.39 – 1.70, *P* = 0.582, Figure

2d). *Six6* was not a significant factor for foraging outcome for fish diet.

Figure 2: Fish acquisition patterns of immature Atlantic salmon as a function of sea-age-dependent genotype (*vgll3*

and *six6*) and sex. Upper and lower panels indicate marginal means for foraging frequency (zero-inflation

component), and foraging outcome (count component), respectively. Error bars are 95% confidence intervals.

Sex: Sex exhibited a marginal effect for foraging frequency and outcome for fish diet. Frequency of

foraging for fish increases in females as salmon gets older (zi component: z = 2.127$P_{\text{sea-age x }sex}$ =

0.033, Figure 2c, Supplementary Table 7), and females forage for fish more often than males in

the 3SW age group (odds ratio = 26.506, 95% CI = 1.45 – 483.96, P = 0.027, Figure 2c). In

contrast, the number for fish in the diet was higher in females in 1SW age groups (odds ratio =

2.64, 95% CI = 1.14 – 6.14, P = 0.024, Figure 2f, Supplementary Table 5), and the pattern was

reversed in 3SW, and males had marginally higher fish diet in the 3SW age group (odds ratio =

1.80, 95% CI = 0.99 – 3.29, P = 0.055, Figure 2f) suggesting an age specific sex pattern in foraging

outcome (count component: z = 2.484, $P_{\text{sea-age x sex}}$ = 0.013) . For both foraging frequency and

outcome the age-dependent effects were driven by age-dependent response of females, rather

than males (Figure 2c, f).

Season, year, reporting group and length effects: Season was an important determinant of fish diet.

Salmon forage for fish more often in the winter (zi component: z = -2.014, P = 0.044), and acquire

more fish (count component: z = 4.413, P < 0.001, Supplementary Table 7). Fishing period,

reporting group, or age standardized length did not have a significant effect on fish diet

(Supplementary Table 7).

The estimated effects of season or reporting group did not meaningfully influence the genotype

or sex effects observed, since the sequential exclusion of these parameters did not meaningfully

change the effect size and/or significance (data not shown). However, the model did not converge

when fishing period was excluded from the model, so their effect on the focal variables are not

assessed. On the other hand, the marginal *vgll3* effect to foraging outcome appeared to be

improved and became significant upon excluding $L_{\text{age.std}}$ from the model, suggesting possible

covariation between the two (Supplementary Figure 3). As such, salmon with *vgll3*[LL] genotype

had more fish in the stomach than salmon with *vgll3*[EE] genotype in 1SW and 2SW sea-age groups

(for 1SW, odds ratio = 4.64, 95% CI = 1.52 – 14.16, P = 0.007; for 2SW, odds ratio = 2.02, 95% CI

= 1.22 – 3.35, *P* = 0.006), but the difference was negligible in the 3SW age group (odds ratio =

0.88, 95% CI = 0.41 – 1.90, *P* = 0.746, Supplementary Figure 3).

Convergence and diagnostics: As evidenced from diagnostics plot from obtained from the

DHARMa package, the fish model had suboptimal fitting (Supplementary Figure 2b). Yet, this

shortcoming may have little impact to main conclusions. For example, when additional

dispersion parameters were added to account for variance heterogeneity for fishing season,

reporting group, season and sex, the homogeneity of variance (or within group deviation from

uniformity) was achieved for all parameters except for fishing season, and Kolmogorov–Smirnov

deviation test was not significant anymore (Supplementary Figure 2c). The addition of these

dispersion parameters did not alter the conclusions associated with genetic or sex effects

(Supplementary Table 8). The convergence was not achieved when testing for alternative

parameterization of model covariates, such as when sea-age and genotype effects was modelled

as a categorical variables, or when testing the effect of non-focal variables to main conclusions

(by sequentially excluding them from the model). Therefore, the model results from continuous

parametrization of sea-age should be cautiously approached.

Discussion

We demonstrated an age-dependent genetic pattern in marine diet variation in wild, sub-adult

Atlantic salmon. By simultaneously capturing the effects of two biological processes that shape

the diet variation, namely active foraging frequency (via the zero-inflated component) and

foraging outcome (via the count component), we provided insights to the foraging dynamics of

Atlantic salmon at sea.

Intriguingly, the two processes had remarkably different sea-age-dependent patterns: younger

individuals contain more crustaceans in the stomach than older age groups suggesting a higher

foraging outcome, despite lower foraging frequencies. This pattern cannot be explained by higher

assimilation rate of older salmon (i.e. which have larger guts) since higher assimilation rates

should also correlate with higher frequencies of empty stomach. Results from these two

biological process combined, suggest older sub-adult salmon adopts a continuous feeding

strategy, albeit a less efficient one, perhaps to fulfill the energetic demands of a larger body size

(Jonsson & Jonsson, 2011; Nina Jonsson & Jonsson, 2003). On the other hand, these results are in

contrast with the previous findings by (Aykanat et al., 2020), whereby an opposite pattern was

shown in mature Atlantic salmon that were sampled in coastal waters, on their migration to

freshwaters for reproduction. Their study suggested that older age groups adopted a feast and

famine strategy as a foraging adaptation to maintain high energy balance. It is unclear what

underlies these opposing patterns in our study. However, it might be possible that a combination

of physiological and behavioral differences associated with differences in life histories (immature

sub-adults vs mature adults), habitat (oceanic feeding grounds vs coastal vs oceanic feeding), or

diet composition could be the underlying reason for the contrast. For example, the diet was

primarily composed of four fish species in Aykanat et al. (2020), a substantial portion of which

was larger than three grams while only a small number of fish species in the diet had on average

weighed more than three grams, suggesting a striking difference in diet breadth. All above factors

may contribute to change in feeding strategies as a function of age.

Our results indicate that older sea-age groups feed more frequently, despite that they lack the

foraging outcome of younger age groups. This is especially intriguing in relation to the

crustaceans diet, since it is contrary to the expected patterns of ontogenetic diet shifts in salmon

(Rikardsen & Dempson, 2010). The larger body of older sea-age groups are energetically more

demanding (Jonsson & Jonsson, 2011; Jonsson & Jonsson, 2003), and it might be that older

salmon are in need of constant searching for food to support the energetic demands of larger

body, relative to younger age groups, especially when prey size is small, a strategy of which may

come in the expense of efficiency of foraging. It could be also speculated that the salmon partly

use filter feeding using their gill-rakers as sieve during high density crustacean prey

distributions. Small salmon (parr and smolts) uses filter feeding to catch small prey (e.g. Keeley &

Grant, 1997). If ocean feeding post-smolts and small salmon also uses this technique,

acknowledging that gill-raker spacing is proportional to fish length (Wankowski 1979), 1SW

salmon might benefit from their smaller gill-raker spacing when targeting the relatively small

crustacean prey compared to larger MSW salmon.

We also demonstrate a shared genetic basis between a major life-history polymorphism (sea-

age) and feeding strategy in Atlantic salmon with a substantial sea-age-dependence. Specifically,

the age-dependent patterns associated with the *vgll3* locus, predominantly in crustacean foraging

rates, provides evidence that feeding strategy is a functional ecological process by which

differences between sea-age phenotypes might be realized. This is a rare example of genetically-

controlled differences in resource acquisition of a highly migratory fish species in the wild and

demonstrates that salmon could be subject to selection if oceanic food webs change (Czorlich et

al., 2022; Strøm, Ugedal, Rikardsen, & Thorstad, 2023; Vollset et al., 2022). Crustaceans are a

primary source of dietary material for Atlantic salmon in the Northeastern Atlantic (Jacobsen &

Hansen, 2001; Rikardsen & Dempson, 2010). A reduction in the water flux from the colder Arctic

region into the northern Atlantic around 2005 was associated with a decrease in biotic and

abiotic factors including plankton abundance, as well as specific growth rate of salmon, the latter

which eventually resulted in delay in maturation timing (Vollset et al., 2022). While our results

do not support that *vgll3* is linked to a change in feeding efficacy (as quantified by the count

component of the model), foraging for crustaceans (as quantified by the zi component) appeared

to be regulated by the *vgll3* locus. These results suggest that an ecological shift in the northeast

Atlantic might exert an evolutionary change in the *vgll3* genomic region.

The crustacean acquisition patterns in relation to *vgll3* might provide some clues on how the

variation in sea-age at maturity may be mediated via a genetic polymorphism in resource

acquisition. To illustrate, 1SW salmon with the *vgll3*^{*EE} genotype appear to obtain as many

crustaceans as 1SW salmon with the the *vgll3*^{*LL} genotype (Figure 1d) despite foraging for

crustaceans less frequently (Figure 1a). In other words, 1SW individuals with *vgll3*^{EE} genotype

gain a similar amount of food (estimated from count model) with less active foraging. Such

increase in acquisition efficiency may be one of the factors that allows salmon with the *vgll3*^{EE}

genotype to mature earlier as 1SW fish, i.e. by allocating surplus resources (saved through

efficient foraging) towards fat deposition and reproductive investment. Furthermore, we also

observed a parallelism in age-dependent crustaceans foraging activity between *vgll3* genotypes

and sex, and in relation to their effect to maturation timing. For example, females, which matures

later than males (Barson et al., 2015; Fleming & Einum, 2010), forage for crustaceans more often

than males (and with similar acquisition efficiencies), a pattern that is similar between *vgll3*^{LL}

and *vgll3*^{EE} genotypes. Similar to salmon with *vgll3*^{LL} locus, perhaps, females' foraging behavior is

not sufficient to acquire energy to trigger early maturation. It might be that an optimal foraging

strategy for earlier maturation is to decrease active foraging time and to increase the foraging

efficiency. On the other hand, it is not clear if age-dependent changes in diet patterns associated

with early maturation in *vgll3*^{EE} genotype and males indicate an age-dependent trade-off

associated with early maturation. In fact, the age-dependent changes in foraging time were

mainly driven by age-dependent changes in the *vgll3*[EE] genotype and males, which are associated

with early maturing. Based on the foraging patterns, early maturation genotype and males are

becoming less efficient in acquiring crustaceans in older sea-ages (i.e., 3SW). Thus, we propose

that early maturation genotype and males have evolved to have higher foraging efficiencies for

crustaceans in early ages, potentially linked to earlier maturation, but this may come with an

efficiency trade-off in later years at sea. In turn, a seemingly maladapted foraging strategy in

younger ages might be selected for in salmon populations by eventual delay in the maturation

timing, resulting to mature at a larger size.

The crustacean acquisition model provides a functional ecological process that explains genetic

(via the *vgll3* locus) and between sexes basis of maturation timing. However, it is not clear if this

effect may be manifested via condition factor, a primary phenotypic trait associated with

maturation timing (Thorpe et al., 1998), and one that is linked to *vgll3* during precocious male

maturation (Debes et al., 2021). To explore if *vgll3* or sex was linked to age-dependent

differences in condition factor, and consistently so with the expected maturation patterns, we

implemented a linear mixed model with a Gaussian error structure using condition factor as

response variable, and using a similar covariate structure as above (except that sea-age was

modeled categorically which was more parsimonious than a numeric model). There was no

evidence of age-dependent association between condition factor and *vgll3* structure

(Supplementary Table 9). Males, on the other hand exhibited a higher condition factor in 1SW age

group (z = 1.982, P = 0.0476), and a lower condition factor in 2SW (z = -4.776, *P* < 0.001)

compared to females, but not in 3SW sea-age groups (t = -0.282, P = 0.7779). This is indeed

consistent with expected maturation patterns in Atlantic salmon, where 1SW is more

predominately males, compared with 2SW more females (Fleming & Einum, 2010), suggesting

that the sex specific diet differences might drive maturation via condition factor effects. However,

the condition factor used here was calculated using the gutted weight, hence did not include any

variation in condition that might be related to guts and other visceral tissue, both of which are

likely important ecological traits in the processes of maturation (Mobley et al., 2021).

Unlike crustaceans, the fish acquisition was generally similar between genotypes across sea-age

groups. However, foraging outcome was meaningfully different within *vgll3* genotypes and sexes

across sea-ages, whereby *vgll3*^{LL} genotype and females consumed higher number of fish in the

1SW age group with marginal and significant age-dependent changes for *vgll3* and sex,

respectively. This pattern further highlight resource acquisition as a potential ecological function

that links maturation timing to genetic polymorphisms in the *vgll3* locus, and sex. However,

caution is required when interpreting the results for the fish model, the fit of which had some

deviations from linear model assumption and since convergence for alternative parameterization

options was not achieved.

Caveats

Studying the evolutionary ecology of Atlantic salmon in the marine phase provides

unprecedented possibilities to understand the ecological and evolutionary basis of life history

variation in the wild, but is inherently difficult. This is due to their extensive oceanic migration

with substantial spatio-temporal variation both within (Gilbey et al., 2021) and between age-

classes. Such stochasticity (due to the marine ecology of salmon as well as the way fish were

sampled) results in structurally complicated data. It was impossible for us to quantify and

compare all possible model structures to account for this complexity. Likewise, the complex

model structure resulted in convergence issues, especially with alternative parameterization of

the fish model, which limits exploration of potential confounding effects. However, especially for

the crustaceans model, we were able to assess the effect of alternative parameterizations at a

great extent, which appears to not affect the main conclusions.

1SW salmon utilize the feeding grounds around the Faroe Islands relatively less often than older

age groups, suggesting that the 1SW fish have spatially distinct feeding ranges in the ocean.

Therefore, it is unclear if the patterns in this study are also observed on feeding grounds that are

predominantly used by 1SW salmon. This encourages further work to verify potential acquisition

patterns on such feeding grounds, and to better understand acquisition-dependent life history

variation associated with the *vgll3* haploblock. Similarly, the low numbers in 1SW are associated

with large 95% prediction intervals around 1SW. This limits the inferential strength of the

relevant results, and predictive power of the analyses.

Given the complex data structure and associated challenges with the interpretations, we limited

our analyses to asking broad questions concerning the eco-evolutionary dynamics of marine

feeding strategy and sea-age phenotype. As such, we primarily grouped diet into two major

categories (crustaceans and fish), and did not address more detailed or refined hypotheses, due

to lacking the statistical power to do so robustly. However, two major crustacean taxa (Hyperiid

amphipods and Euphausiids) had sufficient data to achieve model convergence, by which we

concluded that the general patterns observed in crustaceans were similarly reflected in these

taxa, also irrespective of their size distribution, suggesting prey size may not be important in

shaping age-dependent genetic and sex diversity on diet acquisition.

Conclusions

An important consequence of anthropogenic change has been the large-scale disturbance of

ecosystems globally (Sydeman, Poloczanska, Reed, & Thompson, 2015; Vollset et al., 2022). In

particular, climate-mediated changes are predicted to continue altering marine food webs, with

ecological, genetic, and economic consequences (Daufresne, Lengfellner, & Sommer, 2009;

Pershing et al., 2015). Therefore, linking community-level processes to eco-evolutionary

processes at the individual-level (such as polymorphic expression) is a prerequisite for

understanding how taxa will respond to these large-scale changes. Such an understanding is

essential for effective conservation and ecologically sensitive fisheries management.

This study demonstrates that individual-level variation in prey acquisition both within and

between sea-age phenotypes, which also displays a shared genetic basis (to varying degrees)

with the *vgll3* locus. Furthermore, we show an apparent decrease in crustacean feeding efficiency

for MSW salmon with the *vgll3*^{EE} genotype relative to 1SW salmon with the same genotype, but

not for the *vgll3*^{LL} genotype. Taken together, these results are indicative of variation in feeding

strategy acting as a functional ecological process that links *vgll3* genotype and sea-age in Atlantic

salmon. Our results could help to provide a mechanism to explain the observed changes in the

frequencies of 1SW and MSW phenotypes in some Atlantic salmon populations (Czorlich,

Aykanat, Erkinaro, Orell, & Primmer, 2018; Utne et al., 2021; Vollset et al., 2022). Changes in

marine food webs have long been expected to negatively impact the energetic status of salmon

(Friedland, Chaput, & MacLean, 2005; B. Jonsson, Jonsson, & Albretsen, 2016; N. Jonsson &

Jonsson, 2004; Strøm et al., 2023). The shared genetic basis between prey acquisition and sea-age

that we demonstrate indicates that Atlantic salmon sea-age is susceptible to evolution from

human-induced changes to marine ecosystems.

Data accessibility

555 The compiled datasets and R-codes to reproduce the results are available from the dryad

556 repository https://doi.org/XXX

Acknowledgements

558 This study was funded by the Academy of Finland (grants no. 328860, 353388 and 325964 to TA)

559 and the Research Council of Norway (project number 280308 to KH). We acknowledge DNA

560 Sequencing and Genomics Laboratory (supported by HiLIFE and Biocenter Finland funding),

561 Institute of Biotechnology, University of Helsinki for sequencing. We also thank Katja Enberg for

562 providing comments to an earlier version of this manuscript.

Author contributions

564 TA and KH conceptualized the paper. KH organized sample retrieval from scale archives at NINA.

565 JAJ coordinated the Faroese fieldwork and collected the diet data. TA conducted the statistical

566 analyses, genotyped the samples, and drafted the MS. All co-authors contributed to subsequent

567 drafts.

References

Aykanat, T., Lindqvist, M., Pritchard, V. L., & Primmer, C. R. (2016). From population genomics to
conservation and management: a workflow for targeted analysis of markers identified
using genome-wide approaches in Atlantic salmon *Salmo salar*. *Journal of Fish Biology,*
89(6), 2658-2679. doi:10.1111/jfb.13149
Aykanat, T., Rasmussen, M., Ozerov, M., Niemela, E., Paulin, L., Vaha, J. P., . . . Primmer, C. R. (2020).
Life-history genomic regions explain differences in Atlantic salmon marine diet
specialization. *Journal of Animal Ecology, 89*(11), 2677-2691. doi:10.1111/1365-
2656.13324

Ayllon, F., Kjaerner-Semb, E., Furmanek, T., Wennevik, V., Solberg, M. F., Dahle, G., . . . Wargelius, A. (2015). The vgll3 Locus Controls Age at Maturity in Wild and Domesticated Atlantic Salmon (*Salmo salar* L.) Males. *PLoS Genetics, 11*(11), e1005628. doi:10.1371/journal.pgen.1005628

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., . . . Primmer, C. R. (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature, 528*(7582), 405-408. doi:10.1038/nature16062

Bataillon, T., Gauthier, P., Villesen, P., Santoni, S., Thompson, J. D., & Ehlers, B. K. (2022). From genotype to phenotype: Genetic redundancy and the maintenance of an adaptive polymorphism in the context of high gene flow. *Evolution Letters, 6*(2), 189-202. doi:10.1002/evl3.277

Bonnet, T., & Postma, E. (2018). Fluctuating selection and its (elusive) evolutionary consequences in a wild rodent population. *Journal of Evolutionary Biology, 31*(4), 572-586. doi:10.1111/jeb.13246

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., . . . Bolker, B. M. (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *R Journal, 9*(2), 378-400. doi:Doi 10.32614/Rj-2017-066

Campbell, N. R., Harmon, S. A., & Narum, S. R. (2015). Genotyping-in-Thousands by sequencing (GT-seq): A cost effective SNP genotyping method based on custom amplicon sequencing. *Molecular Ecology Resources, 15*(4), 855-867. doi:10.1111/1755-0998.12357

Chaput, G., & Benoit, H. P. (2012). Evidence for bottom-up trophic effects on return rates to a second spawning for Atlantic salmon (Salmo salar) from the Miramichi River, Canada. *ICES Journal of Marine Science, 69*(9), 1656-1667. doi:10.1093/icesjms/fss055

Charmantier, A., Perrins, C., McCleery, R. H., & Sheldon, B. C. (2006). Quantitative genetics of age at reproduction in wild swans: Support for antagonistic pleiotropy models of senescence. *Proceedings of National Academy of Sciences U S A, 103*(17), 6587-6592. doi:10.1073/pnas.0511123103

Chouteau, M., Arias, M., & Joron, M. (2016). Warning signals are under positive frequency-dependent selection in nature. *Proceedings of National Academy of Sciences U S A, 113*(8), 2164-2169. doi:10.1073/pnas.1519216113

Coulson, T., MacNulty, D. R., Stahler, D. R., vonHoldt, B., Wayne, R. K., & Smith, D. W. (2011). Modeling effects of environmental change on wolf population dynamics, trait evolution, and life history. *Science, 334*(6060), 1275-1278. doi:10.1126/science.1209441

Cousminer, D. L., Berry, D. J., Timpson, N. J., Ang, W., Thiering, E., Byrne, E. M., . . . Early Growth Genetics, C. (2013). Genome-wide association and longitudinal analyses reveal genetic loci linking pubertal height growth, pubertal timing and childhood adiposity. *Human Molecular Genetics, 22*(13), 2735-2747. doi:10.1093/hmg/ddt104

Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P., & Primmer, C. R. (2018). Rapid sex-specific evolution of age at maturity is shaped by genetic architecture in Atlantic salmon. *Nature Ecology and Evolution, 2*(11), 1800-1807. doi:10.1038/s41559-018-0681-5

Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P., & Primmer, C. R. (2022). Rapid evolution in salmon life history induced by direct and indirect effects of fishing. *Science, 376*(6591), 420-423. doi:10.1126/science.abg5980

Daufresne, M., Lengfellner, K., & Sommer, U. (2009). Global warming benefits the small in aquatic
ecosystems. *Proceedings of National Academy of Sciences U S A, 106*(31), 12788-12793.
doi:10.1073/pnas.0902080106
Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E., Parre, N., . . .
Primmer, C. R. (2021). Polygenic and major-locus contributions to sexual maturation
timing in Atlantic salmon. *Molecular Ecology, 30*(18), 4505-4519. doi:10.1111/mec.16062
Enberg, K., Jørgensen, C., Dunlop, E. S., Varpe, Ø., Boukal, D. S., Baulier, L., . . . Heino, M. (2012).
Fishing-induced evolution of growth: concepts, mechanisms and the empirical evidence.
Marine Ecology, 33(1), 1-25. doi:10.1111/j.1439-0485.2011.00460.x
Esteve, M. (2005). Observations of Spawning Behaviour in Salmoninae: *Salmo*, *Oncorhynchus* and
Salvelinus. Reviews in Fish Biology and Fisheries, 15(1-2), 1-21. doi:10.1007/s11160-005-
7434-7
Ferguson, A., Reed, T. E., Cross, T. F., McGinnity, P., & Prodohl, P. A. (2019). Anadromy,
potamodromy and residency in brown trout Salmo trutta: the role of genes and the
environment. *Journal of Fish Biology, 95*(3), 692-718. doi:10.1111/jfb.14005
Fleming, I. A., & Einum, S. (2010). Reproductive Ecology: A Tale of Two Sexes. In *Atlantic Salmon
Ecology* (pp. 33-65): Wiley-Blackwell.
Ford, E. B. (1945). Polymorphism. *Biological Reviews, 20*(2), 45-88. doi:10.1111/j.1469-
185X.1945.tb00315.x
Friedland, K. D., Chaput, G., & MacLean, J. C. (2005). The emerging role of climate in post-smolt
growth of Atlantic salmon. *ICES Journal of Marine Science, 62*(7), 1338-1349.
doi:10.1016/j.icejms.2005.04.013
Funk, E. R., Mason, N. A., Palsson, S., Albrecht, T., Johnson, J. A., & Taylor, S. A. (2021). A supergene
underlies linked variation in color and morphology in a Holarctic songbird. *Nature
Communications, 12*(1), 6833. doi:10.1038/s41467-021-27173-z
Gilbey, J., Utne, K. R., Wennevik, V., Beck, A. C., Kausrud, K., Hindar, K., . . . Verspoor, E. (2021). The
early marine distribution of Atlantic salmon in the North-east Atlantic: A genetically
informed stock-specific synthesis. *Fish and Fisheries, 22*(6), 1274-1306.
doi:10.1111/faf.12587
Goldberg, J. K., Lively, C. M., Sternlieb, S. R., Pintel, G., Hare, J. D., Morrissey, M. B., & Delph, L. F.
(2020). Herbivore-mediated negative frequency-dependent selection underlies a trichome
dimorphism in nature. *Evolution Letters, 4*(1), 83-90. doi:10.1002/evl3.157
Grant, P. R., & Grant, B. R. (2002). Unpredictable evolution in a 30-year study of Darwin's finches.
Science, 296, 707-711.
Gross, M. R. (1985). Disruptive Selection for Alternative Life Histories in Salmon. *Nature,
313*(5997), 47-48. doi:Doi 10.1038/313047a0
Gross, M. R., Coleman, R. M., & McDowall, R. M. (1988). Aquatic productivity and the evolution of
diadromous fish migration. *Science, 239*(4845), 1291-1293.
doi:10.1126/science.239.4845.1291
Halperin, D. S., Pan, C., Lusis, A. J., & Tontonoz, P. (2013). Vestigial-like 3 is an inhibitor of
adipocyte differentiation. *Journal of Lipid Research, 54*(2), 473-481.
doi:10.1194/jlr.M032755
Hartig F (2022). DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. R package version 0.4.6.
Hedrick, P. W. (2012). What is the evidence for heterozygote advantage selection? *Trends in
Ecology & Evolution, 27*(12), 698-704. doi:10.1016/j.tree.2012.08.012

Hvidsten, N. A., Jensen, A. J., Rikardsen, A. H., Finstad, B., Aure, J., Stefansson, S., . . . Johnsen, B. O. (2009). Influence of sea temperature and initial marine feeding on survival of Atlantic salmon Salmo salar post-smolts from the Rivers Orkla and Hals, Norway. *Journal of Fish Biology, 74*(7), 1532-1548. doi:10.1111/j.1095-8649.2009.02219.x

Jacobsen, J. A., & Hansen, L. P. (2001). Feeding habits of wild and escaped farmed Atlantic salmon, *Salmo salar* L., in the Northeast Atlantic. *ICES Journal of Marine Science, 58*(4), 916-933. doi:10.1006/jmsc.2001.1084

Jamie, G. A., & Meier, J. I. (2020). The Persistence of Polymorphisms across Species Radiations. *Trends in Ecology & Evolution, 35*(9), 795-808. doi:10.1016/j.tree.2020.04.007

Janssen, K., Bustnes, J. O., & Mundy, N. I. (2021). Variation in Genetic Mechanisms for Plumage Polymorphism in Skuas (*Stercorarius*). *Journal of Heredity, 112*(5), 430-435. doi:10.1093/jhered/esab038

Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J. M., & Slate, J. (2013). Life history trade-offs at a single locus maintain sexually selected genetic variation. *Nature, 502*(7469), 93-95. doi:10.1038/nature12489

Jonsson, B., & Jonsson, N. (2011). *Ecology of Atlantic salmon and brown trout : habitat as a template for life histories*. Dordrecht: Springer.

Jonsson, B., Jonsson, N., & Albretsen, J. (2016). Environmental change influences the life history of salmon Salmo salar in the North Atlantic Ocean. *Journal of Fish Biology, 88*(2), 618-637. doi:10.1111/jfb.12854

Jonsson, N., & Jonsson, B. (2003). Energy allocation among developmental stages, age groups, and types of Atlantic salmon (Salmo salar) spawners. *Canadian Journal of Fisheries and Aquatic Sciences, 60*(5), 506-516. doi:10.1139/f03-042

Jonsson, N., & Jonsson, B. (2004). Size and age of maturity of Atlantic salmon correlate with the North Atlantic Oscillation Index (NAOI). *Journal of Fish Biology, 64*(1), 241-247. doi:10.1046/j.1095-8649.2004.00269.x

Keeley, E. R., & Grant, J. W. A. (1997). Allometry of diet selectivity in juvenile Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, 54(8), 1894-1902. doi:10.1139/f97-096

Lee, B., Rizzoti, K., Kwon, D. S., Kim, S. Y., Oh, S., Epstein, D. J., . . . Jeong, Y. (2012). Direct transcriptional regulation of Six6 is controlled by SoxB1 binding to a remote forebrain enhancer. *Developmental Biology, 366*(2), 393-403. doi:10.1016/j.ydbio.2012.04.023

Lenth R (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. R package version 1.8.5.

Marden, J. H., Langford, E. A., Robertson, M. A., & Fescemyer, H. W. (2021). Alleles in metabolic and oxygen-sensing genes are associated with antagonistic pleiotropic effects on life history traits and population fitness in an ecological model insect. *Evolution, 75*(1), 116-129. doi:10.1111/evo.14095

Matschiner, M., Barth, J. M. I., Torresen, O. K., Star, B., Baalsrud, H. T., Brieuc, M. S. O., . . . Jentoft, S. (2022). Supergene origin and maintenance in Atlantic cod. *Nature Ecology Evolution, 6*(4), 469-481. doi:10.1038/s41559-022-01661-x

Mobley, K. B., Aykanat, T., Czorlich, Y., House, A., Kurko, J., Miettinen, A., . . . Primmer, C. R. (2021). Maturation in Atlantic salmon (*Salmo salar*, Salmonidae): a synthesis of ecological, genetic, and molecular processes. *Reviews in Fish Biology and Fisheries, 31*(3), 523-571. doi:10.1007/s11160-021-09656-w

Mousseau, T. A., Sinervo, B., & Endler, J. A. (2000). *Adaptive genetic variation in the wild*. New
York: Oxford University Press.
Moustakas-Verho, J. E., Kurko, J., House, A. H., Erkinaro, J., Debes, P., & Primmer, C. R. (2020).
Developmental expression patterns of six6: A gene linked with spawning ecotypes in
Atlantic salmon. *Gene Expr Patterns, 38*, 119149. doi:10.1016/j.gep.2020.119149
O'Sullivan, R. J., Ozerov, M., Bolstad, G. H., Gilbey, J., Jacobsen, J. A., Erkinaro, J., . . . Grant, W. S.
(2022). Genetic stock identification reveals greater use of an oceanic feeding ground
around the Faroe Islands by multi-sea winter Atlantic salmon, with variation in use across
reporting groups. *ICES Journal of Marine Science, 79*(9), 2442-2452.
doi:10.1093/icesjms/fsac182
Pershing, A. J., Alexander, M. A., Hernandez, C. M., Kerr, L. A., Le Bris, A., Mills, K. E., . . . Thomas, A.
C. (2015). Slow adaptation in the face of rapid warming leads to collapse of the Gulf of
Maine cod fishery. *Science, 350*(6262), 809-812. doi:10.1126/science.aac9819
R Core Team. (2019). R: A language and environment for statistical computing. In: R Foundation
for Statistical Computing, Vienna, Austria.
Reznick, D. (2016). Hard and Soft Selection Revisited: How Evolution by Natural Selection Works
in the Real World. *Journal of Heredity, 107*(1), 3-14. doi:10.1093/jhered/esv076
Rikardsen, A. H., & Dempson, J. B. (2010). Dietary Life-Support: The Food and Feeding of Atlantic
Salmon at Sea. In *Atlantic Salmon Ecology* (pp. 115-143): Wiley-Blackwell.
Salminen, M., Erkamo, E., & Salmi, J. (2001). Diet of post-smolt and one-sea-winter Atlantic
salmon in the Bothnian Sea, Northern Baltic. *Journal of Fish Biology, 58*(1), 16-35. doi:DOI
10.1006/jfbi.2000.1426
Salminen, M., Kuikka, S., & Erkamo, E. (1995). Annual variability in survival of sea-ranched Baltic
salmon,Salmo salarL: significance of smolt size and marine conditions. *Fisheries
Management and Ecology, 2*(3), 171-184. doi:10.1111/j.1365-2400.1995.tb00110.x
Sinclair-Waters, M., Piavchenko, N., Ruokolainen, A., Aykanat, T., Erkinaro, J., & Primmer, C. R.
(2022). Refining the genomic location of single nucleotide polymorphism variation
affecting Atlantic salmon maturation timing at a key large-effect locus. *Molecular Ecology,
31*(2), 562-570. doi:10.1111/mec.16256
Strøm, J. F., Ugedal, O., Rikardsen, A. H., & Thorstad, E. B. (2023). Marine food consumption by
adult Atlantic salmon and energetic impacts of increased ocean temperatures caused by
climate change. *Hydrobiologia, 850*(14), 3077-3089. doi:10.1007/s10750-023-05234-2
Svensson, E. I., Willink, B., Duryea, M. C., & Lancaster, L. T. (2020). Temperature drives pre-
reproductive selection and shapes the biogeography of a female polymorphism. *Ecology
Letters, 23*(1), 149-159. doi:10.1111/ele.13417
Sydeman, W. J., Poloczanska, E., Reed, T. E., & Thompson, S. A. (2015). Climate change and marine
vertebrates. *Science, 350*(6262), 772-777. doi:10.1126/science.aac9874
Thorpe, J. E., Mangel, M., Metcalfe, N. B., & Huntingford, F. A. (1998). Modelling the proximate
basis of salmonid life-history variation, with application to Atlantic salmon, *Salmo salar* L.
Evolutionary Ecology, 12(5), 581-599. doi:Doi 10.1023/A:1022351814644
Utne, K. R., Pauli, B. D., Haugland, M., Jacobsen, J. A., Maoileidigh, N., Melle, W., . . . Secor, D. (2021).
Poor feeding opportunities and reduced condition factor for salmon post-smolts in the
Northeast Atlantic Ocean. *ICES Journal of Marine Science, 78*(8), 2844-2857.
doi:10.1093/icesjms/fsab163
Vollset, K. W., Urdal, K., Utne, K., Thorstad, E. B., Saegrov, H., Raunsgard, A., . . . Fiske, P. (2022).
Ecological regime shift in the Northeast Atlantic Ocean revealed from the unprecedented

reduction in marine growth of Atlantic salmon. *Science Advances, 8*(9), eabk2542.
doi:10.1126/sciadv.abk2542
Wańkowski, J. W. J., & Thorpe, J. E. (2006). Spatial distribution and feeding in atlantic salmon,
Salmo salar L. juveniles. *Journal of Fish Biology*, 14(3), 239-247. doi:10.1111/j.1095-
8649.1979.tb03515.x