1 **The transcriptional response of cortical neurons to concussion reveals divergent fates**
2 **after injury**
3
4

5 Mor R. Alkaslasi[1,2], Eliza Y. H. Lloyd[1], Austin S. Gable[1], Hanna Silberberg[1], Hector E. Yarur[3],
6 Valerie S. Tsai[3], Mira Sohn [4], Gennady Margolin [4], Hugo A. Tejeda[3], Claire E. Le Pichon*[,1]
7
8
[1]Unit on the Development of Neurodegeneration, *Eunice Kennedy Shriver* National Institute of Child Health
and Human Development, National Institutes of Health, Bethesda, MD, USA

[2]Department of Neuroscience, Brown University, Providence, RI, USA

[3]Unit on Neuromodulation and Synaptic Integration, National Institute of Mental Health, National Institutes
of Health, Bethesda, MD, USA.

[4]Bioinformatics and Scientific Programming Core, *Eunice Kennedy Shriver* National Institute of Child Health
and Human Development, National Institutes of Health, Bethesda, MD, USA

* Correspondence: claire.lepichon@nih.gov

Abstract

Traumatic brain injury (TBI) is a risk factor for neurodegeneration, however little is known about how different neuron types respond to this kind of injury. In this study, we follow neuronal populations over several months after a single mild TBI (mTBI) to assess long ranging consequences of injury at the level of single, transcriptionally defined neuronal classes. We find that the stress responsive Activating Transcription Factor 3 (ATF3) defines a population of cortical neurons after mTBI. We show that neurons that activate ATF3 upregulate stress-related genes while repressing many genes, including commonly used markers for these cell types. Using an inducible reporter linked to ATF3, we genetically mark damaged cells to track them over time. Notably, we find that a population in layer V undergoes cell death acutely after injury, while another in layer II/III survives long term and retains the ability to fire action potentials. To investigate the mechanism controlling layer V neuron death, we genetically silenced candidate stress response pathways. We found that the axon injury responsive kinase MAP3K12, also known as dual leucine zipper kinase (DLK), is required for the layer V neuron death. This work provides a rationale for targeting the DLK signaling pathway as a therapeutic intervention for traumatic brain injury. Beyond this, our novel approach to track neurons after a mild, subclinical injury can inform our understanding of neuronal susceptibility to repeated impacts.

Introduction

Close to half the population is expected to experience a mild traumatic brain injury (mTBI) at some point in their life[1]. A common form of mTBI is concussion, a brain injury caused by mechanical force and resulting in temporary neurological dysfunction. Although most people seemingly recover, for some the impact can lead to long term damage. There is increasing evidence that repeated mTBI can cause chronic traumatic encephalopathy (CTE)[2] and is a potential risk factor for other neurodegenerative disorders[3–5]. The primary insult of mTBI triggers a cascade of damage termed 'secondary injury' that involves multiple brain cell types and unfolds during the days and weeks following the impact[1]. Even if symptoms during this time can be relatively mild, it is during this chronic phase that neurons are thought to become more vulnerable to repeated injuries. Despite this, the typical treatment for mild TBI is limited to pain management and rest. It is therefore likely that neuroprotective treatments would be beneficial to stave off risk of permanent damage.

To develop effective treatments, we must first better understand the pathways initiated in particular neurons. Many studies to date have lacked the resolution to discern cell type-specific responses[6–8]. More recent work has examined the effect of injury on particular neuron types[9,10]. We reasoned that a detailed look at a model of single mild TBI would provide important insight into the nature and extent of neuronal injury immediately following a concussion. We previously generated a mouse line to track neurons that are transcriptionally responsive to peripheral nerve injury[11] and wondered if it could be used to investigate mTBI.

Peripheral nerve injuries cause a transcriptional response in sensory neurons that is regulated by *Atf3* (activating transcription factor 3) and is essential for functional recovery[11,12]. In sensory neurons, *Atf3* is responsible for upregulating select regeneration-associated genes while repressing many other genes during the recovery process. We wondered whether a similar transcriptional response might occur in the brain after mTBI. Previous studies have observed neuronal *Atf3* activation following injury to the central nervous system. *Atf3* was activated in cortical neurons by TBI and in corticospinal neurons by axon transection depending on the proximity of the injury to the soma[13,14]. Studies in *Atf3*-deficient mice found worse outcomes following TBI and ischemia[15,16], suggesting a protective role for *Atf3*, but did not distinguish between neuronal and glial activation of Atf3. We therefore hypothesized Atf3 would be activated in neurons after mTBI, but wondered how these neurons would compare to peripheral neurons in their ability to exhibit plasticity and recover.

Using genetic reporter mice, single nucleus RNA-sequencing, and slice electrophysiology, we performed a detailed characterization of the neurons that transcriptionally activate *Atf3* after mTBI. We demonstrate that several subclasses of cortical neurons engage the *Atf3* response, but that these undergo divergent fates (death vs survival) that are linked to their identities. We probe the role of multiple candidate pathways for their contribution to cortical neuron death after mTBI and find that dual leucine zipper kinase (DLK), an upstream regulator of *Atf3*, drives neuron death in layer V, highlighting it as a potential therapeutic target for mTBI. These results underscore a

79 differential vulnerability of cortical neurons to mTBI and emphasize the importance of studying
80 injury-induced pathology at the level of individual neuronal subtypes.
81
82 **Results**
83
84 **Characterizing neurodegeneration in a closed skull model of mild TBI**
85
To study mild TBI, we characterized a unilateral closed-skull injury model[17] that provides a
clinically relevant view of concussion injury and allowed us to accurately dissect the resulting
pathological cascade. We used a controlled cortical impact injury wherein an impact was delivered
directly to the surface of the skull at a specified depth and velocity. The impact was provided by
a 3 mm diameter tip positioned over the mouse's left sensorimotor cortex (Fig. 1a). Following this
injury, mice presented with no overt long-term symptomology and no tissue loss, but did exhibit
reproducible cortical astrogliosis in an area approximately 2 mm in diameter and confined to the
ipsilateral cortex (Fig. 1b). This model resulted in a loss of righting reflex concordant with mild TBI
(righting time < 15 minutes[18,19], Fig. 1c), as well as a small yet consistent increase in the serum
biomarker of neuron degeneration, neurofilament light[20] (NfL, average 3-fold higher than baseline
between 1-14 dpi, Fig. 1d).
97
We started by examining pathology in the Thy1-YFP-H mouse, which sparsely expresses a
fluorescent protein in layer V cortical neurons, highlighting their morphology[21,22]. We histologically
confirmed cortical neurodegeneration in these mice (Fig. 1e). At 7 days post injury (dpi), we
observed hallmarks of degenerating dendrites, cell bodies, and axons specifically in the ipsilateral
cortex but not the side contralateral to injury (Fig. 1e-k, Supplementary Fig. 1a). Cortical dendrite
fragmentation in the region above layer V was quantified using a degeneration index calculation[23]
and revealed significant degeneration of YFP+ dendrites[24] only in the ipsilateral cortex when
compared to the contralateral side or the ipsilateral cortex of sham injury controls (Fig. 1f,i). Below
layer V there was a significant increase in YFP-positive structures that did not correspond to cell
bodies but rather to pathological enlargements of axons (area = 10-250 μm^2, mean = 20 μm^2, Fig.
1h,j). Some of these axonal swellings were of comparable size to cell bodies but none contained
DAPI-positive nuclei (Supplementary Fig. 1b). The swellings likely correspond to disruption in
axon transport leading to organellar and protein accumulations, also called diffuse axonal
injury[9,25–27]. We also observed beading of axons (fragments < 10 μm^2) representing axon
degeneration. We found that both axon beading and swelling were increased only in the ipsilateral
cortex at 7 dpi (Fig. 1h,j).
114
Because the Thy1 reporter is stochastically expressed, and because we had specifically observed
inflammatory, dendritic, and axonal pathology only on the side ipsilateral to injury, to quantify any
potential cell loss, we compared the number of YFP+ neurons in the cortex ipsilateral to the injury,
and normalized them to the contralateral cortex of each section. We measured a 15.3% ± 1.8 loss
of cell bodies at 7 dpi, and 26.3% ± 7.9 loss at 14 dpi (Fig. 1e,k). Thus, we find that a single
unilateral closed head impact over the sensorimotor cortex reproducibly leads to degeneration of
layer V projection neurons ipsilateral to the injury and across neuronal compartments.
122

123
mTBI produces an Atf3 response in a subset of cortical neurons
125

Transcription factors play major roles in neuronal responses to injury, and their activation can determine whether a neuron degenerates or regenerates. The transcription factor ATF3, in particular, is a master regulator of the transcriptional response to neuronal injury, and is responsible for driving a transcriptional shift toward an injured cell state[11,12]. We looked for ATF3 expression in brains following mTBI and observed that at 7 dpi, ATF3 immunolabeling localized specifically to the injured side of the cortex (Fig. 2a). We found 10% ± 2.3 of YFP-expressing neurons expressing ATF3 at this timepoint[13] (Fig. 2b). Some ATF3-positive cells were also present in layer II/III (Fig 2a).

Our initial attempts to identify the ATF3-positive neurons in layer V using markers of projection neurons such as CTIP2[28] were unsuccessful. We observed no double labeled cells upon staining for ATF3 and CTIP2 (Fig. 2d-f). We reasoned that ATF3 might be repressing marker genes in the cortex after TBI as has been observed in peripheral sensory neurons after axon injury[11,12]. We therefore performed single nucleus RNA sequencing of these neurons to obtain a more comprehensive picture of their repertoire of RNA expression.

Transcriptional profiling of cells that activate Atf3 in the injured cortex

We employed targeted snRNAseq of *Atf3*-expressing neurons using an inducible Atf3-IRES-CreER mouse line[12] crossed to the INTACT nuclear envelope protein reporter[29]. The resulting animals express GFP tethered to the nuclear envelope in cells that are expressing Atf3 at the time of tamoxifen treatment. Tamoxifen was administered at 4 and 5 dpi to induce expression of the nuclear GFP reporter and cortical tissue at the injury site was collected at 7 dpi, enabling the isolation of nuclei for single nucleus sequencing from cells expressing *Atf3* during this acute phase (Fig. 2c).

We sequenced 8,065 GFP+ nuclei and found a significant number of microglia (4274, 53.0%), excitatory (2057, 25.5%) and inhibitory (744, %) neurons, and small populations of astrocytes (107, 1.3%), oligodendrocytes (75, 0.9%), and other cells (808, 10%) (Supplementary Fig. 2a-e). In this study we focus on the role of ATF3 as an injury marker in neurons, but we note its role in microglial function as an interesting avenue for future investigation[30,31].

Specific subclasses of excitatory and inhibitory cortical neurons activate Atf3-associated injury pathways

We mapped the neuronal nuclei from this experiment onto a highly annotated mouse motor cortex reference atlas[32] to assign cellular subclasses based on the nuclear transcriptome. Of the 2,801 neuronal nuclei sequenced using the Atf3-CreER approach, we identified excitatory neurons across cortical layers, parvalbumin (Pvalb) and somatostatin (Sst) interneurons, and small

numbers of Lamp5, Vip, and Sncg interneurons (Fig. 2d,f, Supplementary Fig. 2a,b,
Supplementary Table 1). Using multiplex *in situ* hybridization with markers from the data, we
validated the presence of Gfp+ excitatory and inhibitory neurons (Supplementary Fig. 3a-c).
Interestingly, all interneuron subclasses expressed the inhibitory neuron marker *Gad2*, but lacked
their subclass markers, including *Pvalb* or *Sst* (Fig. 2f). Similarly, neurons assigned to excitatory
cortical layer identities lacked the typical expression of *Slc17a7* (VGLUT1), but expressed some
layer-specific markers, such as *Cux2*, *Rorb*, and *Foxp2* (Fig. 2f). Thus, in the cortex - similar to
the peripheral sensory nervous system[11,12] - Atf3 expression leads to the downregulation of
multiple marker genes.
176
In our experiment, we captured neurons that had expressed *Atf3* at 4 and 5 dpi, and sequenced
them at 7 dpi. Some of these still expressed *Atf3*, but many only had low or undetectable *Atf3*
expression (Fig. 2g, Supplementary Fig. 3a). In contrast, the damage-induced neuronal
endopeptidase *Ecel1*, whose expression is directly downstream of *Atf3*[33], was highly expressed
in both excitatory and inhibitory neurons (Fig. 2g, Supplementary Fig. 3a). We investigated the
expression of a panel of injury-induced genes across neuronal subclasses and discovered that
not all neuron subclasses that activated *Atf3* underwent the same subsequent transcriptional
programs (Fig. 2g). For example, pro-apoptotic and endoplasmic reticulum (ER) stress genes
such as *Ddit3*[34] were most highly expressed in layer V neurons and low in layer II/III (validated by
in situ hybridization, Supplementary Fig. 3d), while *Atf3* and axon growth genes were most highly
expressed in Pvalb and other interneuron subclasses. Thus although many neurons upregulate
Atf3, their overall transcriptional changes differ according to cell type. Our data thus highlight
heterogeneous transcriptional programs and fates among the Atf3-captured neurons.
190
191
192 **Genetic labeling of Atf3-expressing neurons highlights layer-specific vulnerability**
193
To visualize and map the neurons that express *Atf3* after mTBI, we generated a neuron-specific
Atf3 reporter mouse (Atf3-GFP) in which GFP is permanently expressed only in neurons once
Atf3 is upregulated. This mouse results from a cross between Atf3-IRES-Cre and a Cre-
dependent reporter line expressing GFP under control of the neuron-specific Snap25 promoter
(Jax 021879). In control mice, sparse GFP labeling is observed in the cortex and in some
hippocampal neurons, likely due to developmental Atf3 expression (Supplementary Fig. 4a). We
assessed the extent of Atf3-GFP labeling in the cortex and in other brain regions, observing injury-
induced GFP primarily on the ipsilateral side of the cortex, as well as in the ipsilateral anterior
thalamic nuclei (Supplementary Fig. 4b). The anterior thalamic neurons project into the cortex
around the site of injury, and thus their axons may be damaged in this injury model[35,36].
204
Because this Cre-dependent system results in permanent labeling of neurons in which *Atf3* is
induced, we concluded that loss of GFP-expressing neurons together with signs of
neurodegenerative pathology would indicate cell death. Longitudinal quantification up to 70 dpi of
GFP-expressing neurons in the ipsilateral cortex revealed that a prominent group of layer V
cortical neurons expressed Atf3-dependent GFP between 5 and 10 dpi and subsequently
disappeared by 14 dpi, while a population of layer II/III neurons persisted at 70 dpi (Fig. 3a,b).

The loss of layer V Atf3-GFP neurons by 14 dpi echoes the layer V neuron loss observed in the Thy1-YFP mouse (Fig. 1e,f). Parallel quantification of ATF3 protein revealed a comparative delay in GFP expression and extensive activation of ATF3 in non-neuronal cells, which was consistent with our snRNAseq data (Supplementary Fig. 5a). Interestingly, amplification of GFP signal with immunolabeling revealed that some layer II/III neurons initially exhibit lower expression of GFP (Fig. 3b).

Layer V Atf3-expressing neurons undergo cell death and are phagocytosed following mTBI

Atf3 activation promotes axon regeneration in peripheral injuries, but here we found its expression in neurons that are unlikely to regenerate. Because GFP+ neurons at 7 dpi were present only on the ipsilateral side of the cortex and primarily in layer V (Supplementary Fig. 5c), we were able to use the stereotyped projection patterns of layer V neurons to inspect the dendrites and axons of these neurons. Similarly to the Thy1-YFP+ layer V neurons (Fig 1e-k), we discovered that the GFP+ dendrites were severely fragmented, indicating dendrite degeneration, while GFP+ axons exhibited axonal swellings typical of diffuse axonal injury (Supplementary Fig. 5a). Morphologically unhealthy neurons that exhibited cell body vacuolization and loss of nuclear DAPI signal were also observed in layer V (Supplementary Fig. 5b, d).

Neuron death in layer V was confirmed by examining the expression of several types of apoptotic markers in Atf3-GFP tissue. The pro-apoptotic gene *Ddit3*[34] was significantly increased in GFP+ neurons compared to GFP-negative neurons either ipsilateral or contralateral to injury. In GFP+ neurons with high *Ddit3* expression, we observed high DAPI intensity reflecting chromatin condensation during apoptosis[37] (Supplementary Fig. 5d,e). These apoptotic cells had lower GFP expression and appeared morphologically misshapen (shriveled/deformed, Supplementary Fig. 5c). Additionally, the DNA damage marker phospho-H2AX, which is phosphorylated during apoptosis[38], was elevated in GFP+ neurons at 10 dpi but not in GFP- neurons in the ipsilateral or contralateral cortex (Supplementary Fig. 5f,g). The specificity of phospho-H2AX upregulation to GFP+ neurons and not their GFP- neighbors highlights that this mechanism of cell death is specific to neurons undergoing *Atf3*-associated injury responses at this timepoint. Thus, we conclude that layer V neurons that activate *Atf3* undergo apoptosis in the weeks following injury.

Related to this neuron death, we found that microglia exhibited increased phagocytic activity in the ipsilateral cortex and engulfed debris from dead GFP+ neurons. A significant proportion of CD68+ microglial lysosomes contained GFP+ debris (10% ± 4.2 of at 7 dpi) which increased by 14 dpi (15% ± 3.5), and coincided with the maximal loss of layer V neurons (Supplementary Fig. 5h,i). Microgliosis occurred specifically in the ipsilateral cortex where it peaked around 10 dpi and returned to baseline by 42 dpi (Supplementary Fig. 6a,b). Astrogliosis occurred in a delayed yet prolonged peak of GFAP expression which remained elevated at 70 dpi (Supplementary Fig. 6a,c).

These findings demonstrate mTBI leads to activation of *Atf3*-associated pathways in layer V cortical neurons, to their degeneration and death within two weeks after injury, and to cortical glial responses.

Layer II/III Atf3-expressing neurons survive and remain electrophysiologically active

The analysis of Atf3-GFP neurons across cortical layers over time (Fig. 3a) revealed the loss of most GFP-positive layer V neurons, and highlighted those in layer II/III as the main surviving GFP-positive population at 70 dpi. To evaluate whether this resulted from long-term survival of *Atf3*-marked neurons, we used the inducible Atf3-CreER reporter line. By injecting tamoxifen at 4 and 5 days post injury, we could permanently label cells expressing *Atf3* at this time point and evaluate their localization over time (Fig. 3c). At 7 dpi, most labeled neurons were located in layer V, while those remaining at 21 dpi were primarily found in layer II/III. These layer II/III *Atf3*-marked neurons persisted until at least 42 dpi (Supplementary Fig. 7a), and likely represent a resilient population of neurons that activate this stress response pathway soon after injury and survive. Additionally, layer II/III neurons at 21 dpi appeared morphologically healthy, unlike the degenerative profiles observed in some layer V neurons at 7 dpi (Supplementary Fig. 7b,c).

Next, we determined whether Atf3-GFP neurons displayed injury-induced alterations in excitability and basic membrane properties. We hypothesized the neurons would traverse a cellular state with altered electrophysiological properties reflecting the Atf3-response, and that this would differ between neurons from layer V versus layer II/III. We performed whole-cell patch clamp recordings of excitatory neurons from the neuron-specific Atf3-GFP mice at an acute (5-7 dpi) and late (21 dpi) timepoint. At the acute timepoint, layer II/III neurons expressed low levels of endogenous GFP and were thus too sparse and poorly defined to record from (Fig. 3b). Therefore, at this timepoint, we only recorded from GFP+ neurons in Layer V.

Layer V neurons at the acute stage were sufficiently healthy to record from, therefore not yet undergoing apoptosis, but they had clear alterations in electrophysiological properties compared to GFP-negative control neurons in the ipsilateral or contralateral cortex. They exhibited reduced intrinsic excitability: they were unable to sustain repetitive regular firing (Fig. 4a,b), had a higher rheobase (the minimal current required to initiate an action potential), and a significant decrease in the maximum number of spikes produced (Fig. 4c,d). They were also more depolarized (-54 mV vs -68 mV in controls, Fig. 4e). However, their lowered excitability did not stem from alterations in passive membrane properties since these were no different from controls (input resistance, Supplementary Fig. 7f, and capacitance, which reflects cell size and/or arborization complexity, Fig 4f).

GFP-negative neurons in the ipsilateral cortex were indistinguishable from contralateral uninjured neurons by nearly all measures, suggesting the decline in intrinsic excitability observed in GFP+ layer V neurons is a specific consequence in Atf3-positive neurons. Consistent with this, in our snRNAseq profiling of Atf3-captured neurons, we noted a pattern of downregulation of ion channels involved in action potential firing and maintenance of membrane potential (Fig. 4g). We confirmed the downregulation of *Scn1a* and *Kcnq5* in GFP+ layer V neurons in the tissue (Fig.

297   4h). The downregulation of these genes provides an explanation for the changes in excitability
298   we observed in the Atf3-marked neurons.
299
At the later time point, 21 dpi, Atf3-GFP layer II/III neurons were able to maintain sustained firing,
similar to GFP-negative controls (Fig. 4i, j, l). However, they exhibited increased tonic firing
(Supplementary Fig. 7h), decreased rheobase threshold (Fig. 4k), increased input resistance
(Supplementary Fig. 7j), and depolarized membrane potential (-53 mV vs –66 mV in controls, Fig.
4m) - all measures that reflect increased excitability. GFP+ neurons also exhibited a reduced
capacitance, suggesting that they might be more compact and/or less complex than GFP- controls
(Fig. 4n). Interestingly, amplified hyperpolarization-activated (Ih) current, an inward current that is
important in regulating action potential firing frequency, may contribute to enhanced tonic firing in
GFP+ neurons in layer II/III[39,40] (Supplementary Fig. 7k, i). Together, these findings suggest that
surviving layer II/III Atf3-GFP neurons adapt passive membrane properties to maintain sustained
firing.
311
312
313   **Layer II/III neurons undergo axon initial segment reorganization following mTBI**
314
Because we found that GFP+ neurons in layer II/III survive but are hyperexcitable, we wondered
what the consequences would be on their output. We therefore examined whether their axon
initial segment (AIS) underwent alterations. The AIS is a specialized structure at the base of the
axon that is essential for generating action potentials[41,42]. Other neuron types have been reported
to transiently lose their AIS during regeneration[43–45]. Immunolabeling for Ankyrin-G, a master
scaffolding protein of the AIS, suggested that the AIS in layer II/III neurons was lost at 7 dpi but
regained by 14 dpi (Supplementary Fig. 8a-c). This transient loss of AIS markers was confirmed
with staining for another AIS protein, β4-spectrin (Supplementary Fig. 8d). The hyperexcitability
observed at 21 dpi in layer II/III neurons may be linked to a reorganization of the AIS after mTBI.
By contrast, layer V neurons did not lose their AIS (Supplementary Fig. 8a-d) — their reduced
excitability may be due to lack of necessary machinery for ion flux caused by ion channel
dysregulation (Figure 4f). Previous studies of closed skull TBI have described an early loss of
activity followed by a stage of hyperactivity[46]. Our observation of the transient disappearance of
the AIS in layer II/III neurons may suggest that the observed changes in excitability are inherent
to the recovery process. Interestingly, the AIS is also a site of neuronal polarization in
development and regeneration[47].
331
332
333   **Atf3 is not required for mTBI-induced layer V neuron degeneration or death, but is required**
334   **for downregulation of ion channels**
335
336   Although *Atf3* is required for regeneration of sensory neurons following peripheral nerve injury[12,48],
337   our finding that Atf3-GFP cortical neurons in layer V die following mTBI suggested *Atf3* may play
338   a pro-degenerative role in the central nervous system. To investigate this, we deleted *Atf3* in layer
339   V neurons using an Rbp4-Cre driver (Rbp4-Cre::Atf3[fl/fl], Atf3 cKO) and quantified degenerative
340   pathology. We confirmed that layer V ATF3 expression was effectively reduced in Atf3 cKO mice

at 7 dpi (Supplementary Fig. 9a,b), with any remaining ATF3+ nuclei likely representing non-neuronal cells. Layer V deletion of *Atf3* did not affect dendrite degeneration[24] or presence of axonal swellings at 7 dpi, nor did it prevent mTBI-induced cell death or microgliosis (Supplementary Fig. 9d-i, 12c,d).

Because ATF3 drives downregulation of marker genes[11,12], we wondered if it is required for the loss of ion channel genes following mTBI (Fig. 4g,h). *In situ* hybridization of *Scn1a* in Atf3 cKO tissue (Rbp4-Cre::Atf3[fl/fl]::Sun1-GFP) revealed that the injury-induced loss of *Scn1a* in layer V neurons at 7 dpi is prevented by Atf3 cKO (Supplementary Fig. 10). This demonstration that downregulation of *Scn1a* is *Atf3*-dependent suggests that *Atf3* could also drive transcriptional repression of additional ion channels leading to the observed functional deficits in Atf3-expressing neurons.

The integrated stress response and SARM1 pathway are not required for mTBI-induced degeneration

Several interconnected signaling pathways influence neurodegeneration (Supplementary Fig. 11a). The integrated stress response (ISR), which halts translation in response to multiple neuronal stressors, and the SARM1 pathway, which controls Wallerian degeneration have both been shown to play neurotoxic roles in mTBI[6,49–53]. We wondered if we could protect layer V neurons by targeting these pathways. We use three genetic models to manipulate key players: a phospho-dead mutant of eIF2α (knock-in point mutation, serine to alanine substitution, eIF2a^{S51A}) that results in approximately 50% reduction of eIF2α and thus reduces ISR function[54], a conditional knockout of the pro-apoptotic effector of the ISR *Ddit3* (Rbp4-Cre::Ddit3[fl/fl], Ddit3 cKO) in layer V neurons[34], and a global knockout of the executor of Wallerian degeneration Sarm1[55] (Sarm1 KO). We find that targeting each of these pathways on its own is not sufficient to prevent dendrite degeneration, axon beading or swelling, or cell death (Supplementary Fig. 11). Therefore, we conclude that either none of these pathways is important for neuron death following mTBI, or that their collective action is required.

Layer V deletion of Dlk prevents degeneration and death

We had previously noted the upregulation of multiple stress responsive genes in deeper layer cortical neurons (Fig 2g), so it was perhaps unsurprising that deletion of individual pathway effectors was not sufficient to prevent death of layer V neurons. We observed, however, that phosphorylated c-Jun (p-c-Jun), a known binding partner of ATF3, was distributed in a similar pattern as ATF3 in the ipsilateral cortex (Fig. 5a). These transcription factors, ATF3 and p-cJun, are known to be activated by the axon damage sensing protein DLK[56–58] (dual leucine zipper kinase; Supplementary Fig. 11a). DLK can drive ISR activity through phosphorylation of an ISR kinase, PERK[59,60] (Supplementary Fig. 11a, i-k), and promote Sarm1 activation through inhibition of its regulator, NMNAT[61–63] (Supplementary Fig. 11). We thus reasoned that targeting DLK might be protective as a node sufficiently upstream of multiple neuronal stress responses. Furthermore, while the marker of DLK pathway activation p-c-Jun was detected in both layer V and layer II/III

385 neurons, there was significantly higher expression in layer V neurons, correlating with their
386 differential vulnerability (Fig. 5a,b). We thus tested if deleting DLK would promote survival of Atf3-
387 neurons in layer V.
388
We conditionally deleted Dlk in layer V neurons using the Rbp4-Cre driver line (Rbp4-Cre::Dlk[fl/fl],
Dlk cKO). We validated that *Dlk* transcript was selectively reduced in layer V DLK cKO neurons
(Supplementary Fig. 12b,c). DLK deletion completely prevented layer V neuron death, rescuing
the ~15% loss of these neurons in the ipsilateral cortex (Fig. 5c,d). This rescue was maintained
in Dlk cKO animals at 42 dpi (Supplementary Fig. 12d). By crossing the DLK cKO line with the
Thy1-YFP reporter, we also observed that DLK cKO prevented mTBI-induced dendrite
degeneration, and significantly reduced (but did not completely suppress) axon beading and
swelling (Figure 5e-g).
397
DLK signaling is essential in sensory neurons for recruitment of microglia and other inflammatory
cells to sites of injury[64–66]. We investigated whether DLK cKO locally reduced microgliosis in the
cortex after mTBI. We found that microgliosis was selectively reduced in layer V, where Dlk was
depleted (Supplementary Fig. 13a, b). Interestingly, although DLK is required for *Csf1*
upregulation in sensory neurons to recruit microglia following peripheral nerve injury, we found
that CSF1 does not play a role in mTBI-induced cortical microgliosis (Supplementary Fig. 13e,f).
Thus, mTBI-induced layer V microgliosis is not initiated through a neuronal injury response that
actively recruits microglia via Csf1. Instead, microgliosis may occur as a response to factors
released by apoptotic neurons, such as ATP, and prevention of neuron death by DLK deletion is
therefore sufficient to prevent microglial recruitment.
408
409 We thus find that DLK activation is required for the degeneration of layer V neurons following
410 mTBI, and that its differential activation in layer V and layer II/III neurons may be responsible for
411 the differential vulnerability.
412
413
414 **Discussion**
415
In this study, we develop and characterize a model of mild TBI in which we follow the cellular and
molecular sequelae of a single impact to the skull on specific cortical neuron types during several
weeks following injury. We used a closed-skull injury in order to model a clinically-relevant mild
trauma to the cortex. We find that despite its mild nature, this single impact injury produces wide
ranging consequences to neurons within the cortex, from cell death to survival, with specific
neuron types undergoing specific reproducible fates. We used the neuronal injury marker *Atf3* as
a reporter to gain genetic access to a subset of neurons that transcriptionally responds to the
injury. This allowed us to molecularly and spatially identify these neurons, describe their location
and morphology, record their electrophysiological properties, and determine how their cellular
states evolve over time. We discovered that the Atf3-responsive population of neurons falls into
two categories: one located in layer V that undergoes neuronal death within 2 weeks after injury,
and one in layer II/III that survives at least 2 months after injury. We found that the DLK signaling

pathway is responsible for the death of the Atf3-responsive neurons in layer V, highlighting its
value as a potential target for prevention treatments for TBI.
430
One challenge in studying models of mTBI is the ability to accurately quantify events or cell
numbers using markers that are transient or altered by the injury itself. For example, commonly
used markers to assess apoptosis are quite transient and may be missed. In this study, we
demonstrate that many marker genes of cortical neuron types are lost after mTBI, making it
impossible to accurately label or quantify the neuronal cell types in which they are normally
expressed. This is consistent with other work showing that stress response mechanisms adopted
by injured neurons to regain homeostasis after injury often result in the loss of expression of
markers genes and proteins[11,12,67]. To overcome this, we relied on genetic labeling to track and
record from our neurons of interest. This strategy also facilitated the enrichment of this relatively
rare population of cells within the cortex, allowing us to perform single nucleus transcriptomics to
molecularly identify them using their transcriptome instead of individual marker genes.
Additionally, this novel approach to study injured neurons is applicable to investigating the
neuronal Atf3 response in a range of neurodegenerative conditions, beyond TBI.
444
The Atf3-reporter strategy highlighted the existence of two main populations of neurons after
mTBI, and a striking differential vulnerability between them. The neurons in layer V mostly died
within 2 weeks of the injury while those in layer II/III survived. An intriguing aspect of the Atf3-
GFP neurons in layer II/III is that they initially express the GFP reporter much more weakly than
those in layer V. The initially faint reporter labeling may reflect a stronger translational repression
in these neurons than in layer V that is reversed as they regain homeostasis. A question naturally
arises: what is it about layer V neurons that makes them more vulnerable to degeneration? Layer
V neurons have been highlighted as a vulnerable population within the cortex in multiple diseases
and experimental models[4,68–72]. Using machine learning to classify the regeneration potential of
neurons, a recent study identified layer V neurons as the least regenerative within the cortex[73]. It
may be that their large somas, long axons, and metabolic demands require a molecular make-up
that confers an increased vulnerability[74]. Selective vulnerability is seen across neurodegenerative
diseases, necessitating further research to identify causative elements.
458
We found that DLK is essential for the death of layer V neurons, consistent with the known role
of DLK signaling in promoting neuron death[56,57,75–80]. It is reasonable to imagine that the mild
concussive TBI in our model produces axotomy of layer V projection neurons, and thus DLK-
dependent death. While some layer II/III neurons have corticocortical projections, others project
locally; thus, it is less clear whether they activate the Atf3-stress response pathway as a result of
axotomy or another mechanism. Regardless, the majority of these layer II/III neurons do not
degenerate within 70 days of the injury. Our snRNAseq data clearly demonstrate that, despite
sharing the expression of *Atf3* and *Ecel1*, the neurons from each layer express differential
transcriptional programs, consistent with their divergent fates.
468
Recently, DLK inhibitors have been developed to treat neurodegenerative conditions[57,64,75,81–83],
and understanding the role of the DLK pathway in mTBI will be critical for determining if it may be
a viable therapeutic target. Recently, a Phase I clinical trial of a DLK inhibitor in ALS patients was

halted after weeks of treatment due to observations of adverse effects including low platelet count,
ocular toxicity, and altered touch sensation[84]. However, mTBI may be a more appropriate
indication for trial as the injury timing can be known or even anticipated and dosing could be acute
rather than chronic to reduce undesirable side-effects. Alternate methods to target DLK signaling
may also be developed in future.
477
This study deepens our understanding of how cortical neurons respond to mTBI, and reveals the
heterogeneous nature of their responses. By uncovering the differential vulnerability of distinct
neuronal populations and their diverging engagement of multiple stress response pathways, we
pave the way for targeted therapeutic interventions. It is possible that the initiation of neuronal
stress responses after a single acute injury, as described in this study, can make surviving
neurons susceptible to further injury, highlighting the need for therapeutic approaches for
populations prone to recurrent injury, such as athletes and military personnel. The identification
of the DLK pathway as a potential target for neuroprotection opens new avenues for treatment
strategies. Ultimately, these findings offer crucial insights into the complex landscape of neuronal
injury responses, providing a foundation for future investigations and potential clinical
interventions aimed at mitigating the long-term impact of mTBI on cortical neurons.
489
Limitations
One key limitation of this study is the reliance on a Cre-dependent labeling strategy, which
requires functional translational machinery and sufficient time for reporter expression. Using the
Atf3-Cre line, we may have missed Atf3-positive neurons that are unable to produce the reporter
and/or that die too rapidly. Our analyses using the inducible Atf3CreER mouse are limited to the
specific times of tamoxifen delivery. The use of the Atf3-CreER line for snRNA sequencing of
injured neurons also meant that we lacked the appropriate set of control nuclei, so we relied on a
reference atlas to analyze the data. Finally, our conditional deletion experiments using Rbp4-Cre
result in late embryonic deletion of genes which may have affected layer V neuron development.
499
500 **Methods**
501
Mice
All animal care and experimental procedures were performed in accordance with animal study
proposals approved by the *Eunice Kennedy Shriver* National Institute of Child Health and Human
Disease Animal Care and Use Committee. Adult (>7 weeks of age) male and female mice were
used for all experiments, but were not analyzed separately. Thy1-YFP mice were acquired from
The Jackson Laboratory (B6.Cg-Tg(Thy1-YFP)HJrs/J, Jax Stock No. 003782) Atf3-Cre mice, as
previously described[11], were generated via knockin of an IRES-Cre sequence after the stop codon
of Atf3 at the endogenous locus, such that endogenous Atf3 would remain intact. Atf3-CreER and
Atf3 fl/fl mice were obtained from Dr. Clifford Woolf. Dlk fl/fl mice were obtained from Dr. Aaron
DiAntonio. For sequencing studies, Atf3-CreER mice were crossed to Sun1-sfGFP (B6;129-
Gt(ROSA)26Sortm5(CAG-Sun1/sfGFP)Nat/J, Jax Stock No. 021039) and bred to heterozygosity
for both alleles. For visualization of Atf3-expressing neurons, Atf3-Cre or Atf3-CreER mice were
crossed to Snap25-LSL-eGFP (B6.Cg-Snap25tm1.1Hze/J, Jax Stock No. 021879) or Ai14
(B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J, Jax Stock No. 007914) and bred to

heterozygosity for both alleles. For all conditional knockout experiments, Rbp4-Cre mice (Rbp4-
Cre (B6.FVB(Cg)-Tg(Rbp4-cre)KL100Gsat/Mmucd, MMRRC Stock No. 037128-UCD) were
crossed to the flox line such that the floxed allele would be homozygous and Cre-negative
littermates could be used as controls (this was for Dlk-fl, Atf3-fl and Csf1-fl lines). Chop fl mice
(B6.Cg-Ddit3tm1.1Irt/J, Jax Stock No. 030816), Csf1 fl mice (shared by Dr. Sherry Werner[85]) and
Sarm1 KO (C57BL/6J-Sarm1em1Agsa/J, Jax Sock No. 034399) were used for knockouts. For
Sarm1 KO, heterozygote littermates were used as controls, as heterozygous KO is not sufficient
to prevent Wallerian degeneration. For ISR manipulation, we used Eif2-S51A mice (B6;129-
Eif2s1tm1Rjk/J, Jax Stock No. 017601). Because homozygosity in this mutation is lethal,
heterozygotes were used for experiments, and WT littermates were used as controls. For
inducible Cre experiments, mice were dosed intraperitoneally with 75 mg/kg of a 20 mg/mL
solution of tamoxifen mixed in corn oil at 4 and 5 dpi.
528

Mild traumatic brain injury
Closed-skull mild traumatic brain injury was administered using the Leica Impact One (Leica
Biosystems, Cat. No. 39463920) controlled cortical impact (CCI) device. Mice receiving injury
were anesthetized with 2-2.5% isofluorane and positioned in a nose cone on foam pad. No
stereotaxic restraint was used, however neonatal ear bars were used to loosely stabilize the head
to enhance consistency while maintaining movement upon impact. Mice were shaved and
depilated around bregma. The 3 mm piston tip, mounted on the stereotax at an angle of 10
degrees from the vertical plane, was centered roughly at bregma and moved 2 mm lateral to the
midline. The impactor was driven at a velocity of 5 m/s, depth of 1.5 mm, and dwell time of 200
ms. Animals receiving sham injuries were shaved, depilated, and anesthetized for the same
amount of time as those receiving TBI, but were not administered the injury. Animals were given
5 mg/kg Meloxicam subcutaneously for analgesia immediately after injury and monitored after
removal of anesthesia to evaluate righting reflex. Mice exhibiting tissue deformation following
injury were excluded.
543

Serum collection and Neurofilament Light Simoa Assay
Animals were lightly anesthetized using isofluorane until response to painful stimuli was lost.
Blood was collected at baseline, 1, 9, and 14 dpi. The retro-orbital sinus of one eye was
penetrated with a sterile unfiltered P1000 pipette tip. Blood was collected into BD Microtainer
Capillary Blood Collector tubes (Cat. No. 365967) and allowed to clot at room temperature for 10
min. No more than 10% of the animal's body weight by volume was collected per session. Tubes
were spun down at 6500 rpm for 10 minutes at 4 °C and the supernatant was aliquoted for storage
at -80 °C. The Quanterix Neurology 4-plex A (Item 102153) assay was run following manufacturer
instructions. Briefly, standards were plated in triplicate and test samples were plated in duplicate.
Only neurofilament-light measurements were detectable and consistent between replicates. The
average of the two replicates were reported as the final sample NfL measurement.
555

Single nucleus RNA isolation and sequencing
Mice were anesthetized using 2.5% avertin and decapitated, and the brain was rapidly dissected.
Ipsilateral cortical regions, roughly 4-5 mm diameter and centered around the injury site (as shown
in Fig 1a), were collected and rapidly frozen in pre-chilled tubes on dry ice, then stored at -80 °C.

For the dissection of the desired cortical region, the ipsilateral hemibrain was rapidly isolated from
the skull. Brain tissue anterior, posterior, and lateral to the area of interest (as visualized in Figs
1b and Supp Fig 4b) was removed, as well as any tissue below the corpus callosum. Nuclei
isolation and sequencing was performed as previously described[86]. Ipsilateral cortical regions
from 4 animals were pooled for each sequencing run to remove individual variability. Two datasets
were integrated for the study, one collected from male animals and one from female animals. Sex
differences were not observed. Samples were homogenized in a dounce homogenizer (Kimble
Chase 2 ml Tissue Grinder) containing 1 ml freshly prepared ice-cold lysis buffer. The
homogenate was filtered through a 40 µm cell strainer (FisherScientific #08-771-1), transferred to
a DNA low bind microfuge tube (Eppendorf, #022431048), and centrifuged at 300×g for 5 min at
4 °C. The washing step was repeated, and the nuclei were resuspended in 1× PBS with 1% BSA
and 0.2 U/µl SUPERaseIn RNase Inhibitor (ThermoFisher, #AM2696) and loaded on top of a
1.8 M Sucrose Cushion Solution (Sigma, NUC-201). The sucrose gradient was centrifuged at
13,000×g for 45 min at 4 °C for extra cleanup. The supernatant was discarded, the nuclei were
resuspended in 1× PBS with 1% BSA, 0.2 U/µl SUPERaseIn RNase Inhibitor, and filtered through
a 35 µm cell strainer (Falcon #352235). Before FACS sorting, 5 mM DRAQ5 (ThermoFisher
#62251) was added to label nuclei.
577
GFP+/DRAQ5+ nuclei were sorted and collected on a Sony SH800 Cell Sorter with a 100 mm
sorting chip, and 10k GFP+ nuclei were loaded for sequencing. Using a Chromium Single Cell 3′
Library and Gel Bead Kit v3 (10X Genomics), GFP + nuclei were immediately loaded onto a
Chromium Single Cell Processor (10X Genomics) for barcoding of RNA from single nuclei.
Sequencing libraries were constructed according to the manufacturer's instructions and resulting
cDNA samples were run on an Agilent Bioanalyzer using the High Sensitivity DNA Chip as quality
control and to determine cDNA concentrations. The samples were combined and run on an
Illumina HiSeq2500. There were a total of 370 million reads passing the filter between the two
experiments (replicate 1 = 187,823,841, replicate 2 = 183,968,050). Reads were aligned and
assigned to Ensembl GRm38 transcript definitions using the CellRanger v7.0.1 pipeline (10X
Genomics). The transcript reference was prepared as a pre-mRNA reference as described in the
Cell Ranger documentation.
590
Single nucleus RNA sequencing data analysis
Following the CellRanger pipeline, filtered sequencing data were analyzed using the R package
Seurat version 4.1.3 following standard procedures. Outliers were identified based on the number
of expressed genes (nFeature > 6000) and mitochondrial proportions (percent.mt > 5) and
removed from the data. The data were normalized and scaled with the SCTransform function,
dimensional reduction was performed on scaled data, significant principal components (PCs)
were identified, and 30 significant PCs were used for downstream clustering. Clustering was
performed using the Seurat functions FindNeighbors and FindClusters (resolution = 0.6). Clusters
were then visualized with t-SNE or UMAP. Datasets were integrated with the IntegrateData
function, and integrated data were then processed by the same methods. Data was visualized
with the SCT assay or the RNA assay for dot plots, and plots were generated using Seurat
functions. To assign cell types in an unbiased manner, sequenced nuclei were mapped onto a
published and annotated mouse motor cortex snRNAseq reference dataset[32,87]using the Seurat

MapQuery function. Clusters containing under 10 nuclei were removed from subsequent
analyses. Comparisons to the nuclear reference dataset were made by merging it with our
sequencing dataset and visualizing the RNA assay.
607

Quantification of cell subtype similarity across the datasets

The similarity across cell subtypes between the reference and Atf3-CreER datasets was assessed
using MetaNeighbor[88] v1.22.0 in R v4.3.3. After annotating the cells using reference mapping,
cell subtypes and genes absent in one or both of datasets were removed. This pre-filtering
process resulted in an input count matrix consisting of 17,642 genes and 7,079 cells.
MetaNeighbor is designed to quantify the degree to which cell subtypes replicate across the
datasets based on the expression profiling of highly variable genes (HVGs). The HVGs were
computed using the variableGenes function provided in MetaNeighbor with default argument
setting. Briefly, a gene was selected as an HVG if it was in the top quartile of variable genes and
the top decile of expression bins for each dataset.
618

MetaNeighbor scored subtype-to-subtype similarity using the area under the receiver operator
characteristic curve (AUROC). This computation was performed using the MetaNeighborUS
function provided in MetaNeighbor with the fast_version and node_degree_normalization
arguments set to TRUE. Technically, the function builds a cell network based on the Spearman
correlation computed using the raw counts of HVGs between all pairs of cells in both datasets.
Here, a node represents a cell, and an edge represents the strength of the correlations between
nodes. For each cell (node), the cell subtype is predicted by accounting for the connectivity to
neighboring cells using neighbor-voting algorithm. This algorithm creates a weighted matrix of
predicted cell subtypes by performing matrix multiplication between the network and the binary
vector (0,1) indicating cell subtype membership. Afterwards, this matrix is divided by the node
degree, which returns a score for each cell equal to the fraction of its neighbors. The classification
of cell subtype is performed subsequently by computing the AUROC scores for each cell. This
score is interpreted as the probability that the classifier correctly predicts that a cell subtype in
membership ranked higher than that one not in membership. The AUROC ranges from 0 to 1,
where 1 represents perfect classification and 0.5 represents a prediction as poor as random
guessing. The output AUROC values are returned by averaging all pairs within a subtype.

Fixed tissue harvest and immunostaining
Mice were anesthetized with 2.5% avertin and transcardially perfused with saline, followed by 4%
paraformaldehyde. Tissue was post-fixed overnight and cryopreserved in 30% sucrose prior to
sectioning. Thirty micrometer thick coronal slices were collected free-floating using a Leica
CM3050 S Research Cryostat and stored in antigen preservation solution at 4 °C. For
immunostaining, tissue was washed and permeabilized in 0.1% Triton-X100 in 1× PBS (PBSTx),
then blocked in 5% normal donkey serum in 0.1% PBSTx. Primary antibodies were diluted in
0.5% normal donkey serum in 0.1% PBSTx and tissue was incubated overnight at 4 °C. Tissue
was washed in 0.1% PBSTx and incubated in secondary antibody (ThermoFisher) diluted in 0.1%
PBSTx for 1 h, washed in 1× PBS, mounted on positively charged slides, and coverslipped with
Prolong Diamond (ThermoFisher #P36961). NeuroTrace (1:500, Life Tech. N21483) was applied

following washes for 30 min. Primary antibodies: guinea pig anti-Ankyrin-G (1:500, Synaptic
Systems, 386-005), rabbit anti-ATF3 (1:500, Novus Biologicals, NBP1-85816), rat anti-CD68
(1:500, Bio-Rad, MCA1957), rat anti-CTIP2 (1:500, abcam, ab18465), mouse anti-GFAP (1:500,
Sigma-Aldrich, G3893), chicken anti-GFP (1:500, Invitrogen, A10262), chicken anti-IBA1 (1:500,
Synaptic Systems, 234-006), rabbit anti-phospho-cJun Ser63 (1:300, Cell Signaling Tech., 9261),
rabbit anti-phospho-H2AX (1:400, Cell Signaling Tech., 2577), rabbit anti-Olig2 (1:500, Millipore,
AB9610). A custom made rabbit anti-β4-Spectrin was shared by Dr. Damaris Lorenzo.
654

Multiplexed in situ hybridization
Tissue was sectioned coronally at 16 μm onto positively charged slides using a Leica CM3050 S
Research Cryostat. Slides were dried in the cryostat, then stored at −80 °C. Multiplexed in situ
hybridization was performed according to the manufacturer's instructions for PFA fixed sections
(ACD v2 kit). Probe targets were visualized using Opal dyes 520, 570, 690, or 780 (Akoya). Each
in situ analysis was performed in at least n = 2 mice. Probes: Atf3 (Cat. No. 426891), Atf4 (Cat.
No. 405101), Ddit3 (Cat. No. 317661), Dlk (Cat. No. 458151), Ecel1 (Cat. No. 475331), Gad2
(Cat. No. 439371), Kcnq5 (Cat. No. 511131), Satb2 (Cat. No. 413261), Scn1a (Cat. No. 556181),
Tubb3 (Cat. No. 423391).
664

Electrophysiology brain slice preparation
Mice were anesthetized using Pentobarbital Sodium (NIH Veterinarian Services) and
subsequently decapitated. Brains were swiftly removed and placed in an ice-cold cutting solution
containing (in mM): 92 NMDG, 20 HEPES, 25 glucose, 30 NaHCO3, 2.5 KCl, 1.2 NaPO4
saturated, 10 Mg-sulfate, and 0.5 CaCl2 with 95% O2/5% CO2. The solution had an osmolarity
of 303-306 mOsm (Wescorp). The extracted brain was promptly blocked, dried on filter paper,
and affixed to a platform immersed in ice-cold NMDG-based cutting solution within a chamber of
a Leica VT1200 Vibratome. Coronal slices (300 μm thick) encompassing the somatosensory
cortex, were cut at a speed of 0.07 mm/s. Post-slicing, sections were incubated in an NMDG-
based cutting solution in a chamber for 5-10 min at 34°C. Slices were then transferred to a
chamber filled with a modified holding aCSF saturated with 95% O2/5% CO2. The solution
contained (in mM): 92 NaCl, 20 HEPES, 25 glucose, 30 NaHCO3, 2.5 KCl, 1.2 NaPO4, 1 mM
Mg-sulfate, and 2 mM CaCl2, with an osmolarity of 303-306 mOsm, at room temperature for a
minimum of 1 hr. Slices were kept in the holding solution until being transferred to the recording
chamber.
680

Ex-vivo Whole-Cell Electrophysiology
Whole-cell patch-clamp electrophysiology studies were conducted following the methodology
previously described[89]. Cells were visualized using infrared-differential interference contrast (IR-
DIC) optics on an inverted Olympus BX5iWI microscope. The recording chamber was perfused
at a flow rate of 1.5-2.0 ml per minute with artificial cerebrospinal fluid (aCSF) comprising (in mM):
126 NaCl, 2.5 KCl, 1.4 NaH2PO4, 1.2 MgCl2, 2.4 CaCl2, 25 NaHCO3, and 11 glucose (303-305
mOsm), using a pump from World Precision Instruments. For whole-cell recordings of intrinsic
excitability, glass microelectrodes (3-5 MΩ) were employed, containing (in mM): 135 K-gluconate,
10 HEPES, 4 KCl, 4 Mg-ATP, and 0.3 Na-GTP. GFP-positive and GFP-negative cells were

identified based on the presence or absence of GFP fluorescence in the DIC. Data were filtered
at 10 kHz and digitized at 20 kHz using a 1440A Digidata Digitizer (Molecular Devices). Series
resistance (<20 MΩ) was monitored with a -5 mV voltage step. Cells exhibiting >20% change in
series resistance were excluded from further analysis. For intrinsic excitability, following
membrane rupture in voltage clamp, cells were transitioned to the current clamp configuration
without holding current injection. Intrinsic excitability was evaluated by applying hyperpolarizing
and depolarizing current steps (25 pA steps: 1-sec duration), and changes in voltage and action
potential firing were measured. Whole-cell recordings were conducted using a potassium
gluconate-based internal solution. For all experiments, cells experiencing a 20% or higher
increase in access resistance or high max spike rate consistent with interneurons were excluded
from analysis.
701
Imaging and quantifications
Images were collected using either a Zeiss slide scanner, Zeiss Axiocam 506, or Zeiss confocal
LSM800. Images were quantified using FIJI. For cell counts, slide scanner images were cropped
to equivalent contralateral and ipsilateral area for 3-4 sections and cells were counted by a blinded
observer. For dendrite degeneration quantifications, 63X confocal images were collected from 3-
4 sections (3 regions of interest (ROIs) per section per side). A FIJI macro was created to turn
each image to binary and use the 'Analyze Particles' feature to collect circular area (circularity >=
0.2) and total area. For axon pathology area measurements, the magic wand tool in the Arivis
Vision 4D software was used to manually select all axon blebs (YFP-high, DAPI-negative) for 3-
4 sections per animal. Objects with a 3-10 μm^2 area were defined as axon beading, and those
with area > 10 μm^2 were defined as axon swellings. For intensity quantifications, Z-planes were
summed and ROIs were drawn around cells of interest based on either GFP expression, DAPI
expression (specifically large nuclei > 40μm^2 to select for neurons), or *Tubb3* expression for
RNAscope, depending on the analysis. For normalized mean intensity, mean intensities were
collected for background ROIs (negative for any signal), and cell mean intensity values were
normalized to average background intensity. For percent area quantifications for IBA1 and GFAP,
ROIs were drawn around the ipsilateral or contralateral cortex, an automated threshold (Triangle
method) was set and images were turned to binary, then percent area was calculated using the
'Analyze Particles' feature. For all layer-specific quantifications, either DAPI or NeuroTrace was
used to determine layers. Layer V was defined as the cortical layer with larger and more dispersed
cells. Layer II/III was defined as everything above layer V, and layer VI is everything below.
723
724
Statistical analysis
Wherever possible, quantification of microscopy images was performed blinded. Statistical
analyses were performed using GraphPad Prism 9, except in the case of mixed models analyses.
Normality tests were performed and non-normal data were analyzed using non-parametric tests,
as reported in figure legends. For supplementary figures 10 and 11, in which our comparison of
interest was *ipsi vs contra* in WT *vs cKO* (interaction), we used R package *nlme* and its function
lme for the mixed effect modeling. The fixed effects were *Genotype*Side*, with a random effect
being Side of each animal (i.e., *Side|animal*). We considered models with fixed variances among
all the groups as well as models allowing for separate variances for either each Side and

Genotype, or each Side and animal. Model selection criteria (AIC, BIC and LRT) all pointed to
models allowing for different variances as the best (given that there are typically hundreds of
values per each side of each animal). The comparison of interest was the interaction: change
between cKO and wt genotypes in intensity differences between the sides, (ipsi - contra). For
supplementary figure 5, we were looking at differences between ipsi_pos & contra, and between
ipsi_pos & ipsi_neg (GFP+ neurons and GFP- neurons of the ipsilateral and contralateral cortices;
there is no contra+/-, just contra). To assess the differences between ipsi_pos & contra, and
between ipsi_pos & ipsi_neg, we used R package *nlme* and its function *lme* for the mixed effect
modeling. The fixed effects were the side and GFP status (i.e., ipsi_pos, ipsi_neg and contra),
with each one having a random component for each animal. Each data group (data for a given
side and GFP status of each animal) was allowed to have its own variance. Reported are Holm-
adjusted p-values of Tukey contrasts for multiple comparisons of means.
746

Acknowledgments
We would like to thank Niamh Crowley for his help with the Quanterix assay, Dr. Vincent Schram
and the NICHD Microscopy Core, Dr. Ryan Dale and the NICHD Molecular Genomics Core, Dr.
Clifford Woolf, Dr. Aaron DiAntonio, and Dr. Sherry Werner for sharing mice, Dr. Damaris Lorenzo
for sharing a custom-made antibody, and Drs. Alex Chesler, Mark Cookson, Mark Hoon, Ariel
Levine, Tim Petros, and Nick Ryba for helpful discussions about the manuscript. Biorender.com
was used to make figure graphics.
754

Author Contributions
MRA and CLP designed the experiments and wrote the manuscript. MRA performed
computational analyses. MRA, EYHL, ASG, and HS performed data collection and image
analysis. MRA, HAT, and HEY designed whole-cell patch clamp experiments. HEY and VST
collected and analyzed electrophysiological recordings. MS and GM performed computational
and statistical analyses.
761

762 **Funding**
763 CLP is intramurally funded by the National Institutes of Health, ZIA-HD008966 (NICHD). HAT is
764 intramurally funded by ZIA MH002970-04 (NIMH), a NARSAD Young Investigator Award from the
765 Brain and Behavior Research Foundation, and the BRAIN Initiative (R01). HEY is funded by an
766 NIH Center for Compulsive Behaviors Fellowship.
767

768 **Data availability**
769 The datasets generated in the current study are available from the corresponding author on
770 reasonable request. Data will be deposited to GEO and accession codes will be available before
771 publication.
772

773 **Code availability**
774 Code used in analyzing nuclear sequencing data will be uploaded to GitHub before publication.
775

References

1. Maas, A. I. R. *et al.* Traumatic brain injury: integrated approaches to improve prevention,
clinical care, and research. *Lancet Neurol.* **16**, 987–1048 (2017).
2. McKee, A. C. et al. Chronic Traumatic Encephalopathy in Athletes: Progressive
Tauopathy After Repetitive Head Injury. *J Neuropathology Exp Neurology* **68**, 709–735
(2009).
3. Kahriman, A. et al. Repeated mild traumatic brain injury triggers pathology in
asymptomatic C9ORF72 transgenic mice. *Brain* (2023) doi:10.1093/brain/awad264.
4. Alkaslasi, M. R. et al. Poor Corticospinal Motor *Neuron* Health Is Associated with
Increased Symptom Severity in the Acute Phase Following Repetitive Mild TBI and
Predicts Early ALS Onset in Genetically Predisposed Rodents. *Brain Sci* **11**, 160 (2021).
5. Chiasseu, M., Fesharaki-Zadeh, A., Saito, T., Saido, T. C. & Strittmatter, S. M. Gene-
environment interaction promotes Alzheimer's risk as revealed by synergy of repeated
mild traumatic brain injury and mouse App knock-in. *Neurobiol. Dis.* **145**, 105059 (2020).
6. Chou, A. et al. Inhibition of the integrated stress response reverses cognitive deficits
after traumatic brain injury. *P Natl Acad Sci Usa* **114**, E6420–E6426 (2017).
7. Sarkar, C. et al. Impaired autophagy flux is associated with neuronal cell death after
traumatic brain injury. *Autophagy* **10**, 2208–22 (2014).
8. Mierzwa, A. J., Marion, C. M., Sullivan, G. M., McDaniel, D. P. & Armstrong, R. C.
Components of Myelin Damage and Repair in the Progression of White Matter
Pathology After Mild Traumatic Brain Injury. J Neuropathology Exp Neurology **74**, 218–
232 (2015).
9. Greer, J. E., McGinn, M. J. & Povlishock, J. T. Diffuse Traumatic Axonal Injury in the
Mouse Induces Atrophy, c-Jun Activation, and Axonal Outgrowth in the Axotomized
Neuronal Population. *J Neurosci* **31**, 5089–5105 (2011).
10. Frankowski, J. C. et al. Brain-wide reconstruction of inhibitory circuits after traumatic
brain injury. *Nat Commun* **13**, 3417 (2022).
11. Nguyen, M. Q., Pichon, C. E. L. & Ryba, N. Stereotyped transcriptomic transformation of
somatosensory neurons in response to injury. *Elife* **8**, e49**67**9 (2019).
12. Renthal, W. et al. Transcriptional Reprogramming of Distinct Peripheral Sensory Neuron
Subtypes after Axonal Injury. Neuron (2020) doi:10.1016/j.neuron.2020.07.026.
13. Witcher, K. G. et al. Traumatic brain injury-induced neuronal damage in the
somatosensory cortex causes formation of rod-shaped microglia that promote
astrogliosis and persistent neuroinflammation. *Glia* **66**, 2719–2736 (2018).
14. Mason, M. R. J., Lieberman, A. R. & Anderson, P. N. Corticospinal neurons up-regulate
a range of growth-associated genes following intracortical, but not spinal, axotomy. *Eur J
Neurosci* **18**, 789–802 (2003).
15. Förstner, P. et al. Neuroinflammation after Traumatic Brain Injury Is Enhanced in
Activating Transcription Factor 3 Mutant Mice. *J Neurotraum* **35**, 2317–2329 (2018).
16. Ma, N., Li, G. & Fu, X. Protective role of activating transcription factor 3 against neuronal
damage in rats with cerebral ischemia. *Brain Behav* **12**, e2522 (2022).
17. Xiong, Y., Mahmood, A. & Chopp, M. Animal models of traumatic brain injury. *Nat Rev
Neurosci* **14**, 1**28**–1**42** (2013).
18. Namjoshi, D. R. et al. Merging pathology with biomechanics using CHIMERA (Closed-
Head Impact Model of Engineered Rotational Acceleration): a novel, surgery-free model

of traumatic brain injury. *Mol Neurodegener* **9**, 55 (20**14**).

19. DeWitt, D. S., Perez-Polo, R., Hulsebosch, C. E., Dash, P. K. & Robertson, C. S.
Challenges in the Development of Rodent Models of Mild Traumatic Brain Injury. *J.*
Neurotrauma **30**, 688–701 (2013).

20. Khalil, M. et al. Neurofilaments as biomarkers in neurological disorders. *Nat. Rev.*
Neurol. **14**, 577–589 (2018).

21. Porrero, C., Rubio-Garrido, P., Avendaño, C. & Clascá, F. Mapping of fluorescent
protein-expressing neurons and axon pathways in adult and developing Thy1-eYFP-H
transgenic mice. *Brain Res* **1345**, 59–72 (2010).

22. Feng, G. et al. Imaging Neuronal Subsets in Transgenic Mice Expressing Multiple
Spectral Variants of GFP. Neuron **28**, 41–51 (2000).

23. Gerdts, J., Sasaki, Y., Vohra, B., Marasa, J. & Milbrandt, J. Image-based Screening
Identifies Novel Roles for IκB Kinase and Glycogen Synthase Kinase 3 in Axonal
Degeneration. *J Biol Chem* **286**, 28011–28018 (2011).

24. Ahlgren, H. et al. The Nuclear Calcium Signaling Target, Activating Transcription Factor
3 (ATF3), Protects against Dendrotoxicity and Facilitates the Recovery of Synaptic
Transmission after an Excitotoxic Insult. J Biol Chem **289**, 9970–9982 (2014).

25. Greer, J. E., Hånell, A., McGinn, M. J. & Povlishock, J. T. Mild traumatic brain injury in
the mouse induces axotomy primarily within the axon initial segment. *Acta Neuropathol*
126, 59–74 (2013).

26. Ziogas, N. K. & Koliatsos, V. E. Primary Traumatic Axonopathy in Mice Subjected to
Impact Acceleration: A Reappraisal of Pathology and Mechanisms with High-Resolution
Anatomical Methods. J Neurosci **38**, 4031–4047 (2018).

27. Smith, D. H., Hicks, R. & Povlishock, J. T. Therapy Development for Diffuse Axonal
Injury. J. Neurotrauma 30, 307–323 (2013).

28. Arlotta, P. et al. Neuronal Subtype-Specific Genes that Control Corticospinal Motor
Neuron Development In Vivo. Neuron 45, 207–221 (2005).

29. Mo, A. et al. Epigenomic Signatures of Neuronal Diversity in the Mammalian Brain.
Neuron 86, 1369–84 (2015).

30. Thompson, M. R., Xu, D. & Williams, B. R. G. ATF3 transcription factor and its emerging
roles in immunity and cancer. *J. Mol. Med.* **87**, 1053 (2009).

31. Holland, S. D. & Ramer, M. S. Microglial activating transcription factor 3 upregulation: An
indirect target to attenuate inflammation in the nervous system. *Front. Mol. Neurosci.* **16**,
1150296 (2023).

32. Yao, Z. et al. A transcriptomic and epigenomic cell atlas of the mouse primary motor
cortex. *Nature* **598**, 103–110 (2021).

33. Kiryu-Seo, S. et al. Neuronal Injury-inducible Gene Is Synergistically Regulated by
ATF3, c-Jun, and STAT3 through the Interaction with Sp1 in Damaged Neurons*. J Biol
Chem **283**, 6988–6996 (2008).

34. Syc-Mazurek, S. B., Fernandes, K. A., Wilson, M. P., Shrager, P. & Libby, R. T.
Together JUN and DDIT3 (CHOP) control retinal ganglion cell death after axonal injury.
Mol Neurodegener 12, 71 (2017).

35. Sato, M., Chang, E., Igarashi, T. & Noble, L. J. Neuronal injury and loss after traumatic
brain injury: time course and regional variability. *Brain Res.* **917**, 45–54 (2001).

36. Holden, S. S. et al. Complement factor C1q mediates sleep spindle loss and epileptic
spikes after mild brain injury. https://www-science-
org.ezproxy.nihlibrary.nih.gov/doi/10.1126/science.abj2685 (2021).
37. Dmitrieva, N. I. & Burg, M. B. Analysis of DNA breaks, DNA damage response, and
apoptosis produced by high NaCl. *Am J Physiol-renal* **295**, F1678–F1688 (2008).
38. Rogakou, E. P., Nieves-Neira, W., Boon, C., Pommier, Y. & Bonner, W. M. Initiation of
DNA Fragmentation during Apoptosis Induces Phosphorylation of H2AX Histone at
Serine **139***. *J. Biol. Chem.* **275**, 9390–9395 (2000).
39. Momin, A., Cadiou, H., Mason, A. & McNaughton, P. A. Role of the hyperpolarization-
activated current Ih in somatosensory neurons. *J. Physiol.* **586**, 5911–5929 (2008).
40. Deng, P. & Xu, Z. C. Contribution of Ih to Neuronal Damage in the Hippocampus after
Traumatic Brain Injury in Rats. J. Neurotrauma 28, 1173–1183 (2011).
41. Kole, M. H. P. et al. Action potential generation requires a high sodium channel density
in the axon initial segment. *Nat. Neurosci.* **11**, 178–186 (2008).
42. Leterrier, C. The Axon Initial Segment: An Updated Viewpoint. *J. Neurosci.* **38**, 2135–
2145 (2018).
43. Teliska, L. H., Costa, I. D., Sert, O., Twiss, J. L. & Rasband, M. N. Axon initial segments
are required for efficient motor neuron axon regeneration and functional recovery of
synapses. J Neurosci 42, JN-RM-1261-22 (2022).
44. Marin, M. A. et al. Reassembly of Excitable Domains after CNS Axon Regeneration. J.
Neurosci. **36**, 9148–9160 (2016).
45. Kiryu-Seo, S. et al. Impaired disassembly of the axon initial segment restricts
mitochondrial entry into damaged axons. *Embo J* **41**, e110486 (2022).
46. Carron, S. F., Alwis, D. S. & Rajan, R. Traumatic Brain Injury and Neuronal Functionality
Changes in Sensory Cortex. *Frontiers Syst Neurosci* **10**, 47 (2016).
47. Eichel, K. & Shen, K. The function of the axon initial segment in neuronal polarity. *Dev.*
Biol. **489**, 47–54 (2022).
48. Tian, F. et al. Core transcription programs controlling injury-induced neurodegeneration
of retinal ganglion cells. Neuron (2022) doi:10.1016/j.neuron.2022.06.003.
49. Frias, E. S. et al. Aberrant cortical spine dynamics after concussive injury are reversed
by integrated stress response inhibition. *Biorxiv* 2022.05.31.494250 (2022)
doi:10.1101/2022.05.31.494250.
50. Krukowski, K. et al. Integrated Stress Response Inhibitor Reverses Sex-Dependent
Behavioral and *Cell*-Specific Deficits after Mild Repetitive Head Trauma. J Neurotraum
37, 1370–1380 (2020).
51. Marion, C. M., McDaniel, D. P. & Armstrong, R. C. Sarm1 deletion reduces axon
damage, demyelination, and white matter atrophy after experimental traumatic brain
injury. Exp Neurol **321**, 113040 (2019).
52. Alexandris, A. S. et al. Traumatic axonopathy in spinal tracts after impact acceleration
head injury: Ultrastructural observations and evidence of SARM1-dependent axonal
degeneration. Exp Neurol **359**, 114252 (2023).
53. Henninger, N. et al. Attenuated traumatic axonal injury and improved functional outcome
after traumatic brain injury in mice lacking Sarm1. Brain 139, 1094–1105 (2016).
54. Costa-Mattioli, M. et al. eIF2α Phosphorylation Bidirectionally Regulates the Switch from

Short- to Long-Term Synaptic Plasticity and Memory. Cell **129**, 195–206 (2007).

55. Waller, T. J. & Collins, C. A. Multifaceted roles of SARM1 in axon degeneration and
signaling. *Front. Cell. Neurosci.* **16**, 958900 (2022).

56. Watkins, T. A. et al. DLK initiates a transcriptional program that couples apoptotic and
regenerative responses to axonal injury. *Proc. National Acad. Sci.* **110**, 4039–4044
(2013).

57. Le Pichon, C. E. et al. Loss of dual leucine zipper kinase signaling is protective in animal
models of neurodegenerative disease. *Sci Transl Med* **9**, eaag0394 (2017).

58. Adib, E. A., Smithson, L. J. & Collins, C. A. An axonal stress response pathway:
degenerative and regenerative signaling by DLK. *Curr Opin Neurobiol* **53**, 110–119
(2018).

59. Farley, M. M. & Watkins, T. A. Intrinsic Neuronal Stress Response Pathways in Injury
and Disease. *Annu Rev Pathology Mech Dis* **13**, 93–116 (2018).

60. Larhammar, M. et al. Dual leucine zipper kinase-dependent PERK activation contributes
to neuronal degeneration following insult. Elife 6, e20725 (2017).

61. Summers, D. W., Frey, E., Walker, L. J., Milbrandt, J. & DiAntonio, A. DLK Activation
Synergizes with Mitochondrial Dysfunction to Downregulate Axon Survival Factors and
Promote SARM1-Dependent Axon Degeneration. *Mol Neurobiol* 1–13 (2019)
doi:10.1007/s12035-019-01796-2.

62. Figley, M. D. & DiAntonio, A. The SARM1 axon degeneration pathway: control of the
NAD+ metabolome regulates axon survival in health and disease. Curr Opin Neurobiol
63, 59–66 (2020).

63. Walker, L. J. et al. MAPK signaling promotes axonal degeneration by speeding the
turnover of the axonal maintenance factor NMNAT2. *eLife* **6**, e22540 (2017).

64. Wlaschin, J. J. et al. Dual leucine zipper kinase is required for mechanical allodynia and
microgliosis after nerve injury. Elife 7, e33910 (2018).

65. Hu, Z., Deng, N., Liu, K. & Zeng, W. DLK mediates the neuronal intrinsic immune
response and regulates glial reaction and neuropathic pain. Exp Neurol **322**, 113056
(2019).

66. Shin, J. E., Ha, H., Kim, Y. K., Cho, Y. & DiAntonio, A. DLK regulates a distinctive
transcriptional regeneration program after peripheral nerve injury. *Neurobiol Dis* **127**,
178–192 (2019).

67. Ogino, Y., Bernas, T., Greer, J. E. & Povlishock, J. T. Axonal injury following mild
traumatic brain injury is exacerbated by repetitive insult and is linked to the delayed
attenuation of NeuN expression without concomitant neuronal death in the mouse. *Brain
Pathol* **32**, e13034 (2022).

68. Pressl, C. et al. Selective vulnerability of layer 5a corticostriatal neurons in Huntington's
disease. Neuron (2024) doi:10.1016/j.neuron.2023.12.009.

69. Jara, J. H. et al. Corticospinal Motor Neurons Are Susceptible to Increased ER Stress
and Display Profound Degeneration in the Absence of UCHL1 Function. *Cereb. Cortex*
25, 4259–4272 (2015).

70. Stone, S. et al. Neuron-specific PERK inactivation exacerbates neurodegeneration
during experimental autoimmune encephalomyelitis. *JCI Insight* **4**, e124232 (2019).

71. Ozdinler, P. H. et al. Corticospinal Motor Neurons and Related Subcerebral Projection

Neurons Undergo Early and Specific Neurodegeneration in hSOD1G93A Transgenic
ALS Mice. J Neurosci 31, 4166–4177 (2011).

72. Maekawa, S. et al. Cortical selective vulnerability in motor neuron disease: a morphometric study. Brain 127, 1237–1251 (2004).

73. Kim, H. J. et al. Deep scRNA sequencing reveals a broadly applicable Regeneration Classifier and implicates antioxidant response in corticospinal axon regeneration. Neuron **111**, 3953-3969.e5 (2023).

74. Yadav, A. et al. A cellular taxonomy of the adult human spinal cord. Neuron 111, 328-344.e7 (2023).

75. Welsbie, D. S. et al. Functional genomic screening identifies dual leucine zipper kinase as a key mediator of retinal ganglion cell death. *Proc National Acad Sci* **110**, 4045–4050 (2013).

76. Pozniak, C. D. et al. Dual leucine zipper kinase is required for excitotoxicity-induced neuronal degeneration. *J Exp Medicine* **210**, 2553–2567 (2013).

77. Welsbie, D. S. et al. Targeted disruption of dual leucine zipper kinase and leucine zipper kinase promotes neuronal survival in a model of diffuse traumatic brain injury. Mol Neurodegener 14, 44 (2019).

78. Wlaschin, J. J. et al. Promoting regeneration while blocking cell death preserves motor neuron function in a model of ALS. Brain **146**, 2016–2028 (2022).

79. Saikia, J. M. et al. A critical role for DLK and LZK in axonal repair in the mammalian spinal cord. J Neurosci JN-RM-2495-21 (2022) doi:10.1523/jneurosci.2495-21.2022.

80. Miller, B. R. et al. A dual leucine kinase–dependent axon self-destruction program promotes Wallerian degeneration. *Nat Neurosci* **12**, 387–389 (2009).

81. Siu, M., Ghosh, A. S. & Lewcock, J. W. Dual Leucine Zipper Kinase Inhibitors for the Treatment of Neurodegeneration. *J Med Chem* **61**, 8078–8087 (2018).

82. Ferraris, D., Yang, Z. & Welsbie, D. Dual leucine zipper kinase as a therapeutic target for neurodegenerative conditions. *Future Med Chem* **5**, 1923–1934 (2013).

83. Alexandris, A. S. et al. Protective effects of NAMPT or MAPK inhibitors and NaR on Wallerian degeneration of mammalian axons. Neurobiol. Dis. **171**, 105808 (2022).

84. Katz, J. S. et al. A Phase 1 study of GDC-0134, a dual leucine zipper kinase inhibitor, in ALS. *Ann Clin Transl Neur* **9**, 50–66 (2022).

85. Harris, S. E. et al. Meox2Cre-mediated disruption of CSF-1 leads to osteopetrosis and osteocyte defects. *Bone* **50**, 42–53 (2012).

86. Alkaslasi, M. R. et al. Single nucleus RNA-sequencing defines unexpected diversity of cholinergic neuron types in the adult mouse spinal cord. Biorxiv 2020.07.16.193292 (2020) doi:10.1101/2020.07.16.193292.

87. Hao, Y. et al. Integrated analysis of multimodal single-cell data. Cell **184**, 3573-3587.e29 (2021).

88. Crow, M., Paul, A., Ballouz, S., Huang, Z. J. & Gillis, J. Characterizing the replicability of cell types defined by single cell RNA-sequencing data using MetaNeighbor. *Nat. Commun.* **9**, 884 (2018).

89. Yarur, H. E. et al. Dynorphin / kappa-opioid receptor regulation of excitation-inhibition balance toggles afferent control of prefrontal cortical circuits in a pathway-specific manner. *Mol. Psychiatry* 1–13 (2023) doi:10.1038/s41380-023-02226-5.

Figure 1

a

3 mm

b

7 dpi

-1.3
-1.1
-0.6
-0.1
+0.3
+0.6
+0.8
+1.1

GFAP

c

Righting time

Seconds

Sham   mTBI

d

Neurofilament Light

pg/mL

Baseline   1 dpi   9 dpi   14 dpi

e

Contralateral   Ipsilateral
                                                      7 dpi

Thy1-YFP-H

f Dendrites

g Cell bodies

h Axons

i

Dendrite degeneration

Degeneration Index

ns   *
        ***

Sham   Contra   Ipsi

j

Axon Beading

Bleb count/Cell count

p = 0.063
ns   ***

Sham   Contra   Ipsi

Axon Swelling

**
ns   **

Sham   Contra   Ipsi

k

Cell loss

Layer V neuron count Ipsilateral/Contralateral

*   **
        ns

Sham   7 dpi   14 dpi

Figure 1. Closed-skull mTBI induces layer V neuron degeneration and death. a. Schematic of injury model and location. A controlled cortical impact injury is delivered lateral to bregma. Coronal view (top) shows 3 mm tip positioned over left cortex. Horizontal view (bottom) shows 3 mm impact site relative to bregma. **b.** Example of extent of injury, representative of > 50 samples in which immunostaining for neuroinflammation was performed. GFAP staining is shown across sections from anterior to posterior. For each section, approximate mm from Bregma on the anterior-posterior axis is shown on the right. **c.** Quantification of righting times (time to wake from anesthesia) demonstrating loss of righting reflex in TBI animals consistent with mild TBI. N = 25 Sham, 218 mTBI. All wildtype mice in the study included, excluding those for which righting time was not accurately recorded. **d.** Longitudinal measurement of neurofilament light in serum of mTBI animals. Each shape represents the average of two replicates per mouse. * p = 0.0421 (1 dpi), p = 0.0352 (9 dpi), and p = 0.0381 (14 dpi) by Tukey's multiple comparisons test for each timepoint compared to baseline. **e.** Low magnification image of ipsilateral and contralateral cortex in Thy1-YFP-h mice. Layer V is outlined. High magnification images of **f.** dendrites, **g.** cell bodies, and **h.** axons in the contralateral and ipsilateral cortices. For ipsilateral cell bodies and axons, insets show DAPI expression in cell bodies, and lack of DAPI expression in axon swellings. **i.** Quantification of dendrite degeneration at 7 dpi in ipsilateral, contralateral, and sham (* p = 0.0302, *** p = 0.0007 by Kruskal-Wallis test). **j.** Quantifications of axon beading (fragments with area < 10 μm^2, *** p = 0.0009 by Kruskal-Wallis test) and axon swellings (fragments with area > 10 μm^2, ** p = 0.007 by Kruskal-Wallis test) at 7 dpi. **k.** Quantification of YFP+ neurons in ipsilateral compared to contralateral cortex in sham animals and at 7 and 14 dpi compared to sham (* p = 0.0378, ** p = 0.0073 by Kruskal-Wallis test). For **i-k,** points represent the average of 3-4 sections per mouse. ns: not significant, by unpaired t-test. Scale bars: b. 1 mm, e. 200 μm, f-h (shown in h). 50 μm.

Figure 2

Figure 2. mTBI activates an Atf3-associated transcriptional response in different subclasses of cortical neurons. a. Immunostaining of ATF3 (red) in a Thy1-YFP (white) mouse showing specific expression in the ipsilateral cortex, with higher expression in layer V. Inset highlights YFP+ ATF3+ neurons. ATF3+ neurons are outlined. **b.** Quantification of percent of YFP+ neurons expressing ATF3 at 7 dpi in the ipsilateral cortex. Points represent the average of 3-4 sections per animal. **c.** Immunostaining showing that CTIP2 (cyan) is not expressed in layer V ATF3+ (red) nuclei. ATF3+ nuclei are outlined. **d.** A schematic representation of the single nucleus RNA sequencing workflow, including unilateral closed-skull CCI, tamoxifen dosing, nuclear isolation and FACS, following by 10X Genomics sequencing. **e.** UMAP showing neurons collected by snRNAseq of ipsilateral cortex from pooled Atf3-CreER animals, annotated by mapping to a reference atlas. For subsequent analyses, cell types with fewer than 20 nuclei are excluded. **f.** Dotplot of marker genes for layer-specific excitatory neurons and subclasses of inhibitory neurons, showing downregulation of some markers in Atf3-CreER animals compared to the reference dataset. **g.** Dotplot of a panel of known stress response genes involved in axon growth and regeneration, cell stress and transcription factor regulation, apoptosis, and ER stress highlighting different responses between *Atf3*-expressing neuron subclasses. Genes were selected based on altered expression compared to reference dataset. Low magnification scale bars, 500 μm. High magnification scale bars, 50 μm.

Figure 3

a — Atf3-Cre::Snap25-eGFP

Avg Count vs dpi
- Layer 2/3
- Layer 5
- Layer 6

b — 7 dpi, 21 dpi, Atf3-Cre::Snap25-eGFP, Layer II/III, Layer V

c — Atf3-CreER::Ai14, Tamoxifen 4,5 dpi, Collect 7 dpi, Collect 21 dpi

d — Atf3-CreER::Ai14, Layer II/III, Layer V, Layer V, 7 dpi, 21 dpi

Figure 3. Atf3-expressing neurons in layer V die, while those in layer II/III survive following mTBI . a. Quantification of the average count per section of Atf3-GFP neurons by cortical layer at 3, 5, 7, 10, 14, 21, 42, and 70 dpi. N=6 per timepoint, 3-4 sections counted per animal. **b.** Examples of Atf3-GFP endogenous labeling (left) at 7 and 21 dpi. Layer V is outlined. High magnification examples of GFP-immunolabeled neurons (right) in layer II/III (top) and layer V (bottom) at 7 and 21 dpi. Panels a and b use the Atf3-Cre::Snap25-eGFP mouse, which labels all neurons in which Atf3 has ever been expressed. **c.** Schematic representation of tamoxifen dosing and tissue collection. The Atf3-CreER mouse was crossed to the Ai14 RFP reporter (not neuron-specific). **d.** Representative images of Ai14 signal in ipsilateral cortex at 7 dpi and 21 dpi. Layer V is outlined. Insets highlight neurons and other labeled cells in layer II/III and layer V. Insets for layer II/III suggest an earlier loss of projection complexity that is regained by 21 dpi. Inset for layer V at 21 dpi includes a neuron (left) and a glial cell (right). Panels **c** and **d** use the Atf3-CreER::Ai14 mouse, which labels any cell in which *Atf3* is activated in the presence of tamoxifen. For **a**, error bars represent standard error of the mean (SEM). Scale bars: b. Low magnification, 50 μm, high magnification, 10 μm; d. Low magnification, 500 μm, high magnification, 50 μm.

Figure 4

Figure 4. Layer V Atf3-GFP neurons are unable to fire and downregulate ion channels while layer II/III Atf3-expressing neurons are electrophysiologically functional following mTBI. a. Examples of electrophysiological traces from 7 dpi layer V ipsilateral GFP+ and GFP- neurons and contralateral GFP- neurons. Quantifications of **b.** IO curve, **c.** rheobase, **d.** max spike count (*** p = 0.0002, **** P <0.0001), **e.** resting membrane potential (**** p = <0.0001), and **f.** capacitance (* p = 0.0307) in 7 dpi layer V neurons. **g.** Dot plot of select ion channels in the reference dataset compared to the Atf3-CreER dataset collected in this study showing dysregulation of ion channels in layer II/III and layer V neurons. Genes in bold were validated by in situ hybridization. **h.** In situ hybridization validating downregulation of ion channels. Low magnification image of bilateral cortices from an Atf3-Cre::Snap25-GFP mouse at 7 dpi showing mRNA expression of Gfp (cyan) and Scn1a (yellow). Layer V is outlined. Inset shows mRNA of Gfp (cyan), Tubb3 (white), Scn1a (yellow), and Kcnq5 (red). Gfp+ neurons (outlined) lack expression of Scn1a and Kcnq5 and have little to no expression of Tubb3. Arrowheads highlight Tubb3+ Gfp- neurons with high expression of Scn1a and Kcnq5. A single Z-plane is shown in the insets. **i.** Examples of electrophysiological traces from 21 dpi layer II/III surviving ipsilateral GFP+ and GFP- neurons and contralateral GFP- neurons. Quantifications of **j.** IO curve, **k.** rheobase (** p = 0.0050), **l.** max spike count, **m.** resting membrane potential (* p = 0.0191, ** p = 0.0041), and **n.** capacitance (* p = 0.0217, ** p = 0.0034) in 21 dpi layer II/III neurons. For b and j, points represent the average of all neurons per group, error bars represent SEM. For **c-f** and **k-n**, each point represents one neuron recorded from N=2 or 3 animals. ns: not significant. Significance was determined by Tukey's multiple comparisons test. Low magnification scale bar, 50 μm. High magnification scale bar, 500 μm.

Figure 5

a p-c-Jun | Layer II/III | Layer V

b p-c-Jun

c Contralateral — Ipsilateral 14 dpi | Dlk WT | Dlk cKO | Layer V neuronal nuclei | Contra | Ipsi | WT | cKO

d Cell survival

e 7 dpi | Dendrites | Axons | WT Contralateral | WT Ipsilateral | cKO Ipsilateral

f Dendrite degeneration | WT | cKO

g Axon beading | Axon swelling

Figure 5. Layer V neurons can be rescued from mTBI-induced death and degeneration by deletion of DLK. a. P-c-Jun immunolabeling in the ipsilateral cortex of WT mice at 7 dpi (left). Layer V is outlined. Insets show high magnification images of immunolabeling in layer II/III and layer V. **b.** Quantification of p-c-Jun intensity in layer II/III and layer V at 7 dpi (** p = 0.0016 by paired t-test). Only p-c-Jun+ cells are included based on a threshold of 1.2-fold expression compared to background. Average per animal and value per cell are displayed. Each shape represents one animal. N=4, 2 sections per animal, 11-53 nuclei per animal. **c.** Overview of ipsilateral and contralateral cortices in DLK WT and DLK cKO mice showing layer V GFP+ nuclei. Insets shown on the right. **d.** Quantification of GFP+ neurons in ipsilateral compared to contralateral cortex in WT and cKO mice at 14 dpi based on Sun1-GFP expression shown in c (** p = 0.0025 by Mann-Whitney test). **e.** High magnification images of YFP+ dendrites (top) and axons (bottom) in WT contralateral and ipsilateral cortex, and cKO ipsilateral cortex. **f.** Quantification of dendrite degeneration at 7 dpi in WT and cKO mice (** p = 0p =0.0087 by Mann-Whitney test). **g.** Quantification of axon beading (** p = 0.0079) and swelling (** p = 0.0079 by Mann-Whitney test) at 7 dpi in WT and cKO mice. For **d, f, g,** each point represents the average of 3-4 sections per animal. Low magnification scale bars, 50 μm. High magnification scale bars, 500 μm.