The recovery of parabolic avalanches in spatially subsampled neuronal networks at criticality

Keshav Srinivasan, Tiago L. Ribeiro, Patrick Kells, and Dietmar Plenz*

Section on Critical Brain Dynamics, National Institute of Mental Health, Bethesda, MD 20892, USA

* Correspondence

Figures: 5 Figures, 1 Supplemental Figure

Running Title: Parabolic avalanches in subsampled networks

Correspondence: Dietmar Plenz, Ph.D., Section on Critical Brain Dynamics, National Institute of Mental Health, Porter Neuroscience Research Center, Rm 3A-1000, 35 Convent Drive, Bethesda, MD 20892. Office: (301) (301) 675-5311, plenzd@mail.nih.gov

Keywords: Neuronal avalanches, E/I balanced neural network, subsampling, thresholding, scaling exponent, criticality, ongoing activity, mouse, frontal cortex, 2-photon imaging

1   ABSTRACT

Scaling relationships are key in characterizing complex systems at criticality. In the brain, they are evident in

neuronal avalanches—scale-invariant cascades of neuronal activity quantified by power laws. Avalanches

manifest at the cellular level as cascades of neuronal groups that fire action potentials simultaneously. Such

spatiotemporal synchronization is vital to theories on brain function yet avalanche synchronization is often

underestimated when only a fraction of neurons is observed. Here, we investigate biases from fractional

sampling within a balanced network of excitatory and inhibitory neurons with all-to-all connectivity and

critical branching process dynamics. We focus on how mean avalanche size scales with avalanche duration.

For parabolic avalanches, this scaling is quadratic, quantified by the scaling exponent, χ = 2, reflecting rapid

spatial expansion of simultaneous neuronal firing over short durations. However, in networks sampled

fractionally, χ is significantly lower. We demonstrate that applying temporal coarse-graining and increasing a

minimum threshold for coincident firing restores χ = 2, even when as few as 0.1% of neurons are sampled.

This correction crucially depends on the network being critical and fails for near sub- and supercritical

branching dynamics. Using cellular 2-photon imaging, our approach robustly identifies χ = 2 over a wide

parameter regime in ongoing neuronal activity from frontal cortex of awake mice. In contrast, the common

'crackling noise' approach fails to determine χ under similar sampling conditions at criticality. Our findings

overcome scaling bias from fractional sampling and demonstrate rapid, spatiotemporal synchronization of

neuronal assemblies consistent with scale-invariant, parabolic avalanches at criticality.

INTRODUCTION

Complex systems, composed of many local components or agents that interact weakly, often exhibit event

cascades that cover a wide range of scales in both space and time. Such scale-invariant cascades, typically

identified by power-law distributions in their duration and size – defined as the total activity observed during its

lifetime, have been observed in many real systems, including solar flares[1], earthquakes[2,3], superconductors[4],

sandpiles[5], forest fires[6], and in the brain in the form of neuronal avalanches[7-9]. The scaling of such system

cascades, specifically how their mean size grows with their duration, has been particularly informative as to the

potential underlying dynamics of cascade generation. For example, if local system components activate

independently at a constant rate, the mean size, <S>, of observed cascades will scale linearly with duration, T,

exhibiting a scaling exponent of $\chi \cong 1$, i. e., temporally contiguous system events reflect a constant, yet random

probability of their continuation. Values of $\chi \cong 1$ are often found near a 1st order phase transition[10-12].

Conversely, if cascades exhibit $\chi = 2$, their mean growth is rapid and parabolic as can be found at the second-

order phase transition of a high-dimensional critical branching process, supporting local system interactions[13]. In

general, values of $\chi \gg 1$ are of particular interest as they suggest the ability of a system to bind spatially

distributed events rapidly and selectively into diverse internal system configurations which is generally

considered beneficial for a complex adaptive system such as the brain (e. g.,[14,15]).

Here, we focus on neuronal avalanches, which were first shown to spontaneously emerge in isolated

brain networks, i. e. acute cortex slices and long-term cortex cultures[7]. By measuring locally synchronized

neuronal activity using the local field potential (LFP) and tracking the spatiotemporal spread of LFP cascades

with microelectrode arrays, power-law distributions for both size, *S*, and duration, *T*, $P(S) \sim S^{-\alpha}$ and

$P(T) \sim T^{-\beta}$ were revealed, with exponents $\alpha \approx 3/2$ and $\beta \approx 2$ respectively, in addition to a critical branching

parameter of $\sigma \cong 1$, and significant correlation of neuronal activity within and between[16,17] neuronal cascades

(for review see[18]). These hallmarks of fast, neuronal synchronization in the form of avalanches were

subsequently confirmed in the normal brain for ongoing and evoked LFP cascades in the cortex of nonhuman

primates[19-22] and in humans using magnetoencephalography[23,24]. Their sensitivity to even slight pharmacological

alterations in the balance of excitation and inhibition[7,18,22,25] and changes during pathological conditions such as

epilepsy[26-29] or changes in brain states such as sleep[30,31] and sleep deprivation[32-35], firmly indicate that these

avalanches represent neuronal synchronization of cell assemblies in the normal brain.

To understand avalanche synchronization mechanistically, the contribution of single neuron firing to

avalanches needs to be properly identified. This task is extremely challenging as studies in nonhuman primates

and isolated slice preparations show that neurons participate sparsely and highly selectively even in spatially

extended, large LFP avalanches[22]. Accurate reconstruction of avalanche synchronization from single neuron

activity, therefore, requires the appropriate sampling of a significant portion of the brain network which poses

an enormous technical challenge in neuroscience. Conversely, spatial subsampling, that is, sampling of only a

fraction of neurons, is limited in identifying neuronal synchronization largely by failing to identify avalanche

continuation in non-sampled neurons[36]. This underestimates avalanche duration and size, biasing outcomes

towards expectations for decorrelated, random processes[9,37,38]. Accordingly, simulations of avalanche generating

critical branching processes and fractional sampling of neurons reduces the expected growth in mean avalanche

size with avalanche duration, χ, from the expected value of $\chi = 2$, to a value of χ closer to $1 - 1.3$[39-41]. Not

surprisingly, estimates of $\chi \cong 1.3$ were reported for fractionally sampled neuronal spikes of potential avalanche

activity *in vivo*[40,42-45] and *in vitro*[46].

If the exact start and end times of avalanches are known, for example in simulations under the

assumption of 'separation of time scales', spatially subsampled avalanches can be easily corrected for by

temporal integration thereby recovering the correct critical exponents in size and duration[41] (for a review, see

also[47]). Avalanche times, however, are largely unattainable in experimental data for which only a small fraction

of neurons is observed and temporal integration eventually will degenerate avalanche scaling by

indiscriminately concatenating successive avalanches[9]. Recently, Capek & Ribeiro et al.[9] demonstrated that,

even when the exact times of avalanches are not known, $\chi = 2$ can be recovered for subsampled excitation-

inhibition (E-I) balanced networks of integrate-and-fire neurons which are known to belong to the mean-field

directed percolation (MF-DP) universality class[48,49]. This rescue requires (1) reducing uncorrelated neuronal

action potential firing and (2) using temporal coarse-graining with an increase in minimal synchronization

requirements to identify synchronized population activity in the network. Under these conditions, the authors

could show $\chi = 2$ *in vivo* 2-photon imaging recordings of cellular activity with high signal-to-noise ratio for

sensory evoked and spontaneous spike avalanches in frontal and sensory cortices of awake mice in line with

their simulations[9]. This is in line with a recent EEG study[50] in humans that demonstrated $\chi = 2$.

In the current work, we extend our previous investigations of rescuing χ in subsampled, critical

networks. We systematically study the rescue of the scaling exponent χ to its critical value of 2 in a continuously

driven, subsampled all-to-all DP-model[48] for which we demonstrate the equivalence of fractional sampling and

thresholding. Importantly, we show that the rescue requires the network to be very close to criticality and fails

in the near sub- and supercritical regimes. We demonstrate the robust rescue of χ in the critical model over a

domain covering 4 orders of magnitude in fractional sampling within the temporal coarse graining vs.

thresholding parameter regime. Using this approach, we demonstrate the robust recovery of $\chi \cong 2$ in cellular 2-

photon imaging data of the frontal cortex of awake mice over a large range of varying thresholds and recording

conditions. We further show that, in contrast, the 'crackling noise' relationship and related measures which can

predict χ[51-53] fail under conditions of fractional sampling. Our findings demonstrate the pitfalls of assessing

avalanche synchronization when only parts of a critical network are observed and demonstrate the regime of

potential scaling rescue for synchronization in the form of parabolic avalanches under experimental conditions

of fractional sampling.

RESULTS

Neuronal avalanches were studied using a critically balanced excitatory-inhibitory (E-I) model with all-to-all

connectivity consisting of one million neurons driven externally using independent, stationary Poisson

processes[48] (**Fig. 1a**). The external driving triggered an average of 20 neurons to fire per time step, Δt, which

intermittently induced spreading of neuronal activity in the network. Simulations spanned across 10^8 time steps.

An avalanche was defined as a series of contiguous time periods, T, of coincident firing within Δt surpassing a

predetermined threshold, θ, over a specific number of time steps, k (**Fig. 1b, e**). Thus, our model-based

approach was characterized by two basic parameters: $k \cdot \Delta t$ and θ, which were explored in the context of the

subsampling fraction of observed neurons, f, ranging from 1 out of every 10,000 neurons observed (f = 0.01%) to

all neurons observed (f = 100%). We identified the statistical distribution of sequence size, S, and duration, T, as

a function of θ, f, and k and estimated the corresponding size exponent, α, and duration exponent, β, within the

constraints of the lower and upper cut-offs (**Fig. 1c**; see Methods;[54,55]). In our critical model, these distributions

take on the form of power laws at k = 1, f = 100% and θ = 100, the defining characteristics of avalanches[7]. The

scaling exponent, χ, governs the growth in average avalanche size, $\langle S \rangle$, for a given duration, T,

$$\langle S \rangle \sim T^{\chi},$$ (1)

and accordingly, we estimated the scaling factor χ from the slope of the average avalanche size, $\langle S \rangle$, for a given

avalanche duration, T, outside the minimal and maximal cut-off in logarithmic scales (**Fig. 1d**). This scaling factor

was previously referred to as χ_{sh} as it captures the scaling of short-duration avalanches that occur within the

power law regime of the size and duration distribution (see also[9]). The fitting procedure (see Methods) also

identifies a second slope, χ_{lg}, for long-duration avalanches that occupy the cut-off regime of the corresponding

size and duration distributions. As demonstrated previously and in line with expectations for uncorrelated

activity[9], χ_{lg} remains close to 1 independent of thresholding and coarse-graining and this trivial scaling factor will

not be considered further.

In critical systems, it has also been found that χ relates to the critical exponents α' and β' as

$$\chi_{cn} = \frac{(\beta' - 1)}{(\alpha' - 1)}$$ (2)

('cn' = crackling noise[51-53]). For comparison, we therefore calculated both χ_{cn} and the 'Deviation from Criticality

Coefficient' [56],

$$\text{DCC} = \chi - \chi_{cn} \tag{3}$$

from our slope estimates α and β. This latter construct has increasingly been used as a distance measure from a

critical point implying that the experimentally observed dynamics is close to criticality with low (< 0.2) DCC

Figure 1. Schematics of the balanced E/I-model and identification of suprathreshold sequences in population spiking activity with respect to minimal coincident spiking threshold and temporal coarse graining. (**a**) Schematic of the neuronal network consisting of N = 10^6 neurons (80% excitatory and 20% inhibitory) with a low external Poisson drive of rate $\lambda = 2*10^{-5}$ per time step per neuron. The E/I balance is controlled by g, which scales the inhibitory weight matrices (W_{II}, W_{EI}) as a function of the excitatory weight matrices (W_{EE}, W_{IE}). If not otherwise stated, we set g = 3.5 to obtain critical dynamics (see Methods). (**b**) High variability in intermittent population activity characterizes critical dynamics. Snapshot of the summed neuronal spiking activity as a function of time. (**c**) Size and duration of avalanches in the critical model follow power-laws with corresponding exponent estimates α and β, respectively. Note that the external Poisson drive and the finite size of the network introduce lower and higher cut-offs, respectively (*shaded areas*). (**d**) Scaling of mean avalanche size as a function of duration also follows a power-law. χ is the estimate of the scaling exponent for short avalanches (below cut-off point, ▲). Avalanches within the upper cut-off (above cut-off point, ▲) exhibit a trivial scaling exponent close to 1, denoted as χ_{lg}, that is largely independent of threshold and temporal coarse-graining and will not be considered further. *Purple*: Corresponding scaling relation and cut-off for temporal coarse-graining with k = 5 (see below). (**e**) Zoomed population activity from *B*. At the original temporal resolution Δt and given the coincident spiking threshold, θ (*left*), we can identify two sequences of suprathreshold activity, S_1 and S_2 with durations T_1 and T_2, respectively. Temporally coarse-graining the population activity (*right*; k = 5, binning the data into new bins of 5 time points) and increasing the threshold to $\theta' > \theta$, absorbs S_1 and S_2 into a new suprathreshold activity period T' with size S' with corresponding change in scaling (see also d).

(e.g. [44,57]), despite the fact that as one deviates from criticality the size and duration power laws break down,

making the slope estimates α and β, and therefore the χ_{cn} measure, unreliable.

High coincident firing thresholds underestimate χ in the fully sampled model

In systems characterized by a clear separation of timescales and an absence of noise, the necessity for a

coincident firing threshold diminishes. However, within real systems lacking such temporal separation and

affected by significant intrinsic and measurement noise, the statistical analysis becomes intricately tied to our

selection of threshold values. Setting the threshold too low poses the risk of incorporating substantial noise

from the system, while setting it too high leads to data sparsity, resulting in the exclusion of a significant portion

of the system's activity. To emulate the noisy environment and absence of timescales separation between

avalanche generation and propagation inherent in real systems, we introduce a minimal external Poisson drive

in our simulations. This Poisson drive establishes a noise floor within the spiking activity, serving as the origin of

avalanches within the system. This is clearly visible in the size distribution of avalanches, which exhibits a power

law only for avalanche sizes of $S_0 > 50 - 100$ spikes up to the finite-size of the network[48] (*cf*. **Fig. 1c** *left*). To

isolate avalanches above the noise floor (i. e., $S > S_0$), we first studied the dependence of α, β, χ on θ and k in the

fully sampled network (f = 100%; **Fig. 2**). In **Fig. 2a**, we show an overview in the form of color maps for all three

parameters when increasing θ. At the highest temporal resolution (k = 1), we observed a rapid decrease in χ

below its expected value of 2 as θ increases. This change largely follows the dependence of β while α remains

close to 1.5, the latter being in line with findings from LFP avalanches in non-human primates[19] (**Fig. 2b**). At high

thresholds, χ tends to approach 1.3, a value commonly reported in experiments[39-45], which is likely a

consequence of subsampling biases inherent in these experimental settings. We conclude that high thresholds in

weakly driven, noisy networks bias towards longer suprathreshold periods, significantly underestimating the

scaling exponent in critical systems.

A similar underestimation holds true for the χ_{cn} value. Inserting the corresponding values of exponents

extracted from our fully sampled system for small thresholds, we obtain $\chi_{cn} \cong 2$ and DCC $\cong 0$, in line with

expectations for the mean-field directed percolation (MF-DP) universality class[58].

Figure 2. Increasing the threshold in the fully sampled model underestimates the scaling exponent, χ, which can be rescued by temporal coarse graining.
(**a**) Consolidated view of the avalanche size (α), duration (β), and scaling (χ) exponents as a function of threshold θ and temporal coarse-graining factor k (f = 100%, fully sampled model). *Drive dominated*: Low threshold regime (*above the red broken line*) dominated by external Poisson drive. *White frames*: Parameter regions displayed in **b** and **c**.
(**b**) At the highest temporal resolution (k = 1), high thresholds underestimate the scaling exponent, χ, as well as χ_{cn}, with DCC remaining low. Size exponent, α (*left*), duration exponent, β (*middle*), scaling exponent, χ, and expected crackling noise relation, χ_{cn}, (*right*) as a function of θ.
(**c**) At high threshold (θ=1000), temporal coarse graining recovers χ, but the χ_{cn} exhibits a singularity leading to high DCC. Size exponent, α (*left*), duration exponent, β (*middle*), as a function of coarse graining factor, k, (θ = 1000). α passes through 1 (*red broken line; black triangle*), which causes a singularity in χ_{cn}. (*right*). Note that the scaling exponent, χ, stabilizes to a value of 2 (*right, black dashed line*), whereas χ_{cn} grows until it passes through a singularity at the temporal coarse graining value of k = 17 (*vertical broken line; triangle; see inset*).

When θ increases, χ_{cn} drops significantly below 2, which is similar to our observation for χ and accordingly the

DCC remains close to zero (see **Fig. 2a**). We attribute the small, yet consistently observed underestimation of χ

with respect to χ_{cn} (**Fig. 2b**) to the error-bounds of our fitting procedure. We conclude that avalanche scaling will

be underestimated even in fully sampled networks if there are technical constraints on minimal coincident spike

counts that require the use of high thresholds.

Temporal coarse-graining rescues χ at high thresholds in the fully sampled model

Next, we show that temporal coarse graining rescues χ in the presence of a high threshold in fully sampled

networks. This is shown in **Fig. 2c**, in which χ approaches 2 for $k > 8$ at $\theta = 1000$ (see also corresponding

projections in **Fig. 2a**). In contrast, the size and duration distributions exhibit a gradual reduction in steepness

when k increases, due to concatenating subsequent cascades, in line with experimental findings[7,19,41,59]. In the

model, this eventually causes α to cross the value of 1 (**Fig. 2c**), which forces χ_{cn} to undergo a singularity

(Equation 2; **Fig. 2c, inset**). Accordingly, we find a rapid breakdown in the agreement between χ and χ_{cn} with

increase in k, with diverging DCC (**Fig. 2c**). We conclude that while temporal coarse-graining can mitigate the

bias introduced by thresholding in the fully sampled model, the DCC remains as an unreliable estimate for

avalanche scaling even in fully sampled networks under high thresholding conditions.

Rescue of χ = 2 is only seen in critical dynamics that is fractionally sampled and fails for slightly
super- or sub-critical dynamics.

We recently demonstrated that the rescue of the critical scaling exponent, χ, was exclusive to the critical model

($g_c = 3.5$) and failed for subcritical conditions[9]. Here, we explored this dependency more systematically and show

that small deviations from criticality by changing the E/I-balance parameter, g (see Methods), significantly

impede rescue to the expected critical (MF-DP) value of 2. In **Fig. 3a,** we illustrate the maximum χ across various

temporal coarse graining factors k, plotted against g for subsampled models ($f = 0.1\%$) utilizing a 1-spike

threshold ($\theta = 1$). In subcritical models ($g > 3.5$), the restored scaling exponent swiftly deviates from its critical

value of 2 towards 1. Conversely, in slightly supercritical models ($3.45 < g < 3.5$), χ also rapidly degrades to below

2. We note that for $g < 3.45$, persistent activity (shaded gray regions in **Fig. 3a**) obstructs the identification of

avalanches at the specified threshold. This demonstration carried out for a critical branching model suggests

that a successful rescue of the critical value of χ implies the dynamics to be at an E/I-balance of the order of 0.05

or 1.4% close to criticality. While further analytical studies are warranted to establish the rescue of χ as a key

hallmark of critical dynamics, our findings suggest temporal coarse-graining combined with thresholding to

recover the expected value χ within a particular universality class might be a sensitive approach when

confronted with fractionally sampled critical dynamics.

Figure 3. Temporal coarse graining at critical value of g: Rescue of scaling exponent, equivalence between threshold and fractional sampling, and increased mean pairwise neuron correlation.
(**a**) The maximum scaling exponent across different temporal binning values as a function of the balance parameter, g, for a subsampled model ($f = 0.1\%$) employing a 1-spike threshold. Notably, the retrieval of $\chi = 2$ is exclusively observed in critical models ($g_c = 3.5$), but promptly diminishes in slightly subcritical or supercritical models. In supercritical models, we obtain ceaseless activity (*shaded gray region*), precluding the identification of avalanches at the specified threshold. Yellow circles mark g values used in **c**.
(**b**) Collapsed scaling exponent curves for different values of the sampling fraction, f, ranging from 100% to 0.1% for a threshold that is scaled by the sampling fraction as $\theta = 3{,}000{*}f$ (i.e., a threshold of 3,000 spikes for the fully sampled model translates to a threshold of 3 spikes for the 0.1% sampled model). *Inset*: The total collapse error as a function of collapse exponent, ξ. The error is taken over multiple different curves with θ ranging from 100 to 100,000. The error shows a clear minimum at $\xi = 1$, indicating the threshold can be scaled proportionally with the sampling fraction to obtain the best collapse.
(**c**) The rescue of $\chi = 2$ correlates with an increase of the mean delayed pair-wise correlation among neurons, which is independent of the fraction of neurons sampled. However, the temporal coarse graining regime of k for which $\chi = 2$ (*green area*) is not predicted by this pair-wise correlation. For a supercritical ($g = 3.475$) network, the mean pairwise correlation is lower, but still increases with temporal coarse graining. For a subcritical network ($g = 3.6$), the mean delayed pair-wise correlation is close to 0.

Scaling collapse demonstrates equivalence of thresholding and subsampling in the critically balanced

all-to-all E/I-model

Both, the increase in coincident spiking threshold θ as well as a decrease in sampling fraction, f, underestimate

the scaling exponent χ due to prematurely terminating avalanches by unobserved network activity. On the other

hand, temporal coarse graining allows for concatenating sequential activity periods potentially recovering the

proper duration for a subset of avalanches. We therefore examined the impact of temporal coarse graining on

thresholding and fractional sampling. To achieve this, we first collapsed the χ vs. k curves over a wide range of θ

($10^2 - 10^5$) by scaling θ with an exponent of f, ξ (see Methods). We found that the corresponding functions,

obtained for different values of f, exhibited a minimum total collapse error for $\xi = 1$, demonstrating a linear

relationship between θ and f in our all-to-all model. This implies that a fully sampled network (10^6 neurons) with

$\theta = 3,000$ spikes is qualitatively equivalent to a 0.1% sampled model (1,000 neurons) with $\theta = 3$ spikes (**Fig. 3b**).

This equivalence between θ and f allows us to rescale the rescue of χ for different f in the corresponding (k, θ)-

plane for ease of comparison (see **Fig. 4**).

The rescue of $\chi = 2$ indicates a recovery of mean pair-wise correlations

Subsampling as well as decorrelation of spiking activity have been shown to reduce the expected growth in

mean avalanche size with avalanche duration from $\chi = 2$, to a value of χ closer to $1 - 1.3$[39-41]. We therefore

studied how the mean delayed pair-wise correlation, $\bar{C}(k)$, changes with temporal coarse graining, k, when

recovering χ (see Methods). Temporal coarse graining increases $\bar{C}(k)$ in the critical balanced E/I-network and

saturates at high k, independent of the amount of fractional sampling (**Fig. 3c**). The increase in correlation did

not predict the temporal coarse graining value of k at which $\chi = 2$ and was not found in strongly subcritical

networks, for which the average pairwise correlation remains near zero and independent of k (**Fig. 3c**; g = 3.75).

In slightly supercritical models, the recovery of $\bar{C}(k)$ was significant but did not support $\chi = 2$ for the critical

model. Taken together, these findings demonstrate that $\chi = 2$ identifies the expansion of coincidental firing

among groups of neurons in the critical model and is not adequately captured by a particular value of $\bar{C}(k)$, a

measure limited to studying pairs of neurons.

Subsampling and spike-resolution introduce unobservable regions of scaling exponents

By taking advantage of the trade-off between thresholding and sub-sampling (see Subsection above), we now

can rescale our k vs. θ parameter space by $f \cdot \theta$ to account for different fractional sampling. In the consolidated

analysis (**Fig. 4**), we demonstrate reasonable agreement with the rescue of $\chi = 2$ from 0.1% - 100% fractional

sampling. When we increase the threshold (or equivalently, decrease f), we observe a deviation from $\chi = 2$, yet

the exponent can be reliably restored at higher values of k (**Fig. 4**, white broken lines). On the other hand,

Figure 4. Temporal coarse -graining identifies a robust rescue regime for χ = 2 covering 0.1% - 100% fractional sampling at decreasing thresholding level, which is absent for the *DCC*

(a) Consolidated view of χ as a function of θ and k for different values of f. For f = 100%, $\chi \cong 2$ for low θ and k (*star*) but as we make the data sparser by increasing the threshold (or reducing the sampling), we need a higher coarse graining factor, k, to compensate and rescue χ back to 2 (*square*). *White dotted region*: visual guide for χ close to 2 for increasing θ and k. The grey parts of the plots for f = 0.1% and 0.01% respectively are unobservable parameter regions since they would require fractional thresholds, below the 1 spike minimum resolution of the model. At f = 0.1% sampling, χ can be rescued to a value of 2, but not for more severe subsampling, f = 0.01%. For triangles see **b**. (b) The *DCC* remains close to 0 only for low k, and quickly breaks down at higher k values. For the fully sampled model, there exists a region with the correct scaling exponent $\chi \cong 2$ as well as low *DCC* (*star* in A and B) However, as we move to lower sampling, this region of correspondence becomes more and more difficult to maintain. For f = 1% and 0.1%, regions at low k, and moderate threshold (or equivalently lower sampling) remain at low *DCC*, but underestimate the true scaling exponent, χ (*up* and *down triangles*, respectively).

reducing f eventually introduces unobservable (k, θ)-regions which arise from the 1 spike resolution of the

model excluding fractional thresholds. We note that even when sampling only 0.1% of the neurons, one can still

recover $\chi = 2$. However, this approach fails to simultaneously restore all exponents while maintaining the

crackling-noise relationship (see **Fig. 2a** and Supplementary Fig. S1 online) because the point of concurrent

rescue lies within the unobservable regions of parameter space. Decreasing sampling further to f = 0.01% (100

neurons out of 1 million neurons), prevents rescue of $\chi = 2$.

Temporal coarse graining does not rescue the 'crackling noise' relationship of χ_{cn} (Equation 2). The DCC,

introduced recently as the difference between the data-based value of χ (Equation 3) and χ_{cn} should measure

close to 0 if both estimates agree. As can be seen in **Figure 4**, co-regions with DCC = 0 and $\chi = 2$ exist in the fully

sampled system (**Fig. 4**; star). In contrast, no such co-alignment is found for subsampled systems and temporal

coarse graining fails to establish a *DCC* = 0 regime for most values of k and θ (**Fig. 4**; square, triangles) and

accordingly *DCC* = 0 does not reflect a proper rescue of the true $\chi = 2$ under these conditions.

Robust recovery of χ = 2 for parabolic avalanches in frontal cortex of awake mice

In the previous section, we demonstrated that temporal coarse-graining and thresholding effectively identify a

robust subregion where the scaling exponent χ is rescued in the model. This rescue can be collapsed across

different levels of fractional sampling. We now present similar findings from cellular 2-photon imaging of

ongoing neuronal activity in the anterior cingulate cortex of awake mice (n = 17 experiments, each 30 minutes

long; n = 5 mice; see Methods for details). The avalanche number distribution, obtained for each recording,

follows a lognormal pattern as a function of the threshold (**Fig. 5a,d;** see also [9]) and guides our analysis. In

addition, this lognormal distribution varies with temporal coarse-graining and is thus a function of k (**Fig. 5a, d**).

In a first approach, we pooled all data sets for a given absolute spike threshold (**Fig. 5a**), which revealed a

consistent subregion of $\chi \cong 2$ for different spike density threshold levels (**Fig. 5b**). Specifically, we observe that

the recovery of $\chi \cong 2$ shows a θ vs. k dependency, consistent with our simulations, whereas the DCC remains

variable and noisy (**Fig. 5c**).

Figure 5. Parabolic avalanches in ongoing activity of frontal cortex exhibit threshold and temporal coarse-graining rescue of scaling exponent χ = 2 in line with critical model dynamics.
(**a**) Lognormal distribution of avalanche number as a function of threshold for 17 different recordings (*n* = 5 mice; 2-photon imaging of ongoing activity in frontal cortex) at the original temporal resolution (*k*=1, *black*) and after temporal coarse graining (*k*=10, *purple*). Note the experimental variability at a given *k*, as well as the systematic shift in the distribution with change in *k*. A sample threshold (*θ, grey dotted*) shows how a given fixed threshold relates to these distributions. (**b**) Scaling exponent, *χ*, and (**c**) crackling noise deviation, DCC, as a function of temporal coarse grain factor, *k* and the spike density thresholds. The subregion in which $χ \cong 2$ (*broken line*) is similar to the one identified in the critically balanced E-I model. At low thresholds, the scaling is rescued at smaller values of temporal coarse graining factor, k and at high thresholds, it is rescued at a larger value of *k*. Over this large range of parameters, the DCC is very noisy even a small change in the parameters can cause a large change in the DCC. (**d**) Z-scoring procedure for the lognormal distributions corrects for experimental variability as well as temporal coarse graining . A sample z-scored threshold ($θ_{z,k}$, *grey dotted*) shows how a given z-scored threshold relates to these distributions. (**e**) Scaling exponent, *χ*, and (**f**) crackling noise deviation, DCC, as a function of temporal coarse grain factor, *k,* and the z-scored threshold at the given *k*, $θ_{z,k}$. Figure D shows a robust rescue of $χ \cong 2$ (*broken line*) for a large range of z-scored threshold values (-1.5 to 1.5). The range of k values over which $χ \cong 2$ becomes smaller with an increase in the z-scored threshold. Like B, Figure E shows that the DCC remains unreliable and noisy even when the thresholds are z-scored and tuned for each *k*.

In a second approach, we z-scored the avalanche number distribution for each data set in order to account for

experimental variability as well as for variable shifts induced by temporal coarse graining (**Fig. 5d**). **Figure 5e**

shows that, regardless of the exact choice of z-scored threshold within a wide range (-2 to +1.5), *χ* is robustly

recovered to approximately 2. The range of *k* values that recover *χ* decreases as the threshold increases. This

robust recovery supports previous studies that adopted an avalanche-maximizing threshold (z-score = 0)[8] or

lower thresholds (z-score = -2)[9]. We note that at very low thresholds, noise becomes problematic, and at very

high thresholds, data scarcity leads to poor statistics. In contrast, the DCC measure remains highly variable and

unreliable even when the threshold is individually selected for each experiment and coarse-graining value (**Fig.**

5f).

DISCUSSION

The study of criticality in neuronal systems, particularly neuronal avalanches, has been extensively researched

and debated (for review see[18,53,60-63]). This study addresses the intricacies and challenges of estimating scaling

relationships in the brain, especially under conditions of fractional subsampling. A significant finding of our

research is the ability to retrieve scaling exponents with fractional subsampling by employing temporal coarse

graining and threshold adjustments. The sensitivity of this approach to deviations from criticality supports our

method to study potential universality classes of criticality in real-world systems such as the brain. We

demonstrate that temporal coarse-graining, combined with stricter criteria for coincident neuronal firing, allows

for accurate estimation of the critical scaling exponent ($\chi = 2$), even when observing as little as 0.1% of the

network's neurons. This suggests that it's possible to accurately determine critical scaling characteristics despite

heavy subsampling limitations. In contrast, previous studies on similar directed percolation (DP) models found a

reduction in the scaling exponent to 1.3 with spatial subsampling of 10% and below[39]. Additionally, subsampling

has been observed to falsely suggest population independence of correlations, impacting markers of criticality

such as specific heat[64]. It's important to note that our model, due to its all-to-all connectivity, does not allow for

exploration of spatial correlation dependencies.

Temporal coarse-graining when avalanche periods are unknown

Temporal coarse-graining has previously been suggested as a means to preserve the size exponents of

avalanches in subsampled systems, assuming an infinite separation of time scales[41]. This assumption would

theoretically allow researchers to distinguish sequentially occurring avalanches, a scenario which may not hold

in natural systems. In contrast, our analysis shows that temporal coarse-graining can effectively maintain the

scaling exponent in models lacking a clear separation of time scales, even with fractional sampling as low as

0.1%. This suggests its utility in analyzing highly subsampled experimental data within the directed percolation

(DP) class. Our approach introduces a threshold above the Poisson noise floor to approximate the separation of

successive avalanches, revealing a trade-off between subsampling intensity and the threshold level. In our all-to-

all model, increasing the threshold of the fully sampled network mimics the effects of subsampling combined

with a lower threshold, illustrating the interaction between these factors and their collective impact on criticality

evaluations. Leveraging this trade-off, we apply our methodology to cellular-resolution 2-photon imaging data

from the anterior cingulate cortex of awake mice, consistently recovering the scaling exponent to a value of 2, in

line with expectations for a critical branching process. Additionally, new avalanche detection methods that

utilize continuous time models have been developed, though their effectiveness compared to traditional

methods under experimental and subsampling conditions remains to be fully evaluated[65].

Our recovery of the scaling exponent $\chi \approx 2$ in ongoing neuronal spike synchronization of the awake mouse

frontal cortex provides compelling evidence for the critical role of $\chi \approx 2$ in capturing essential features of

neuronal avalanches (see also[9,20]). For instance, models with slow latent variables can generate heavy-tailed or

even power-law distributions of neuronal activity size and duration, but they typically show $\chi \approx 1$, suggesting a

lack of correlated activity[66-68]. While the critical branching process used here achieves $\chi \approx 2$, it does not account

for temporal correlations among successive avalanches (e. g.[16-18,69]), limiting its ability to fully model neuronal

avalanches in the brain.

It's important to distinguish temporal coarse-graining from renormalization procedures. While

renormalization in neuronal avalanches can preserve estimates of maximal mutual information during spatial

coarse-graining under complex LFP sampling conditions[70], our focus here is on addressing stochastic

subsampling bias through temporal coarse-graining, as detailed previously[9].

Limits in rescuing the scaling exponent under subsampling conditions

Our simulations also clearly show that subsampling in discrete models renders certain parts of the phase space

inaccessible (see **Fig. 4**), preventing the recovery of the scaling exponent under heavy subsampling ($f \ll 0.1\%$). A

similar limitation exist for temporal subsampling, especially when the required precision exceeds the system's

resolution[39]. Properly addressing these spatial and temporal constraints is crucial for the analysis of

experimental data with limitations in both resolution domains. Further limitations arise with extensive

subsampling of network topologies, particularly when estimating heavy-tailed or scale-free distributions[71-73].

While under-sampled critical branching processes have been explored in extracellular spike recordings with

microelectrode arrays in the anesthetized rat[36], these limitations shouldn't overshadow our findings' potential.

Careful consideration of experimental parameters can ensure feasible and informative criticality assessments,

providing valuable insights into the core mechanisms of neural synchronization.

The DCC and crackling noise relationship fails to recover the scaling exponent under subsampling

conditions

The Deviation from Criticality Coefficient (DCC) approach[40,44,56] has gained popularity for quantifying criticality

based on the "crackling noise relationship"[51-53]. However, concerns have been raised about its ability to identify

critical dynamics[37,68] given that random processes can exhibit a DCC close to 0 thus limiting general claims of this

approach to identify critical dynamics even in fully sampled systems. For instance, an unbiased random walk will

produce excursions above the mean that are power-law distributed in sizes, with exponent 1.5, durations, with

exponent 1.33 and scaling with exponent 1.5, which result in the random walk obeying the crackling noise

relationship (*DCC* = 0), despite not being a critical system. Our study demonstrates several limitations of the DCC

approach, particularly under subsampling conditions: (1) DCC estimates become unreliable with limited data, (2)

the DCC is highly sensitive to temporal coarsening, leading to noisy results, (3) the DCC metric has a singularity

under fractional sampling, making it a questionable marker for criticality. Furthermore, the DCC approach

assumes power-law slopes obtained from size and duration distributions directly translate to critical exponents.

This is flawed, as analysis parameters like temporal resolution and spatial sampling distance systematically

influence these slopes[7,19]. These factors must be considered when interpreting criticality from experimental data

using the DCC approach.

The sensitivity of temporal coarse-graining approach to thresholds in avalanche size definition

Our analysis emphasizes the scaling exponent's sensitivity to synchronization thresholds. As the threshold

increases, we observe a noticeable deviation of the scaling exponent and the crackling noise ratio (χ_{cn}) from the

expected value of 2 (see also[13]). This sensitivity highlights the importance of careful consideration when

choosing thresholding strategies in experimental data. Thresholding of time-series and ignoring the

subthreshold contribution during suprathreshold periods can lead to a change in the avalanche size exponent

itself as demonstrated in simulations of critical dynamics[13]. Capek, Ribeiro et al.[9] demonstrated experimentally

that including or excluding the subthreshold regime for size estimates provides narrow upper and lower bounds

for χ that center around 2. Ideally, threshold selection should strike a balance between preserving genuine

synchronization events and minimizing noise, especially when accurate scaling exponent estimates are crucial.

Synchronization, coincident firing and delayed neuronal correlations

While traditional studies of spatiotemporal synchronization in neural systems have focused on phase-locked

oscillations, our system operates in a unique scenario—one devoid of a dominant frequency. This lack of a

dominant frequency becomes particularly important when considering 'noise correlations,' which encapsulate

the intrinsic non-stimulus-induced correlation structure among neurons. These correlations exert a notable

influence on the decoding capabilities of neuronal populations, constituting a pivotal aspect of neural

information processing[74-76].

In this study, we investigate scale-free neuronal avalanches as a context for exploring neuronal

synchrony and criticality. Avalanches offer a powerful tool for understanding neuronal synchronization, from

local clusters to the entire brain. As we've shown, analyzing avalanches depends crucially on choosing thresholds

and temporal resolution. This dependence is similar to the analysis of spatial and temporal neuronal

correlations, which are heavily influenced by their respective coarse-graining[77,78]. Interestingly, as shown in

Figure 3, restoring the scaling exponent χ to its critical value of 2 doesn't directly correspond to the recovery in

pairwise correlations. This emphasizes that χ at 2 signifies a critical system that describes the coordination

among many neurons, not just an average estimate of correlation among pairs of neurons.

Our findings have implications for both neuroscience and complex systems research. Neuronal

avalanches are valuable markers of synchronization in the brain. Precise scaling exponent estimation, even with

limited data, is crucial for understanding mechanisms of neural synchronization in healthy and diseased brains.

This supports the idea that the brain operates within a critical regime, aligning with the concept of self-

organized criticality observed in many complex systems.

METHODS

Model Topology

In this work, we employed the model originally proposed by Girardi-Schappo et al.[48] and subsequently modified

in Capek & Ribeiro et al.[9]. This model represents a critically balanced system of stochastic integrate-and-fire (IF)

neurons, where the excitatory and inhibitory neurons maintain an equilibrium. Our neuronal network consisted

of $N = 10^6$ stochastic, non-leaky IF neurons, with a fully connected architecture. To reflect the prevalence of

excitatory neurons in the cortex, we set the ratio of excitatory to inhibitory neurons to 4:1. The connectivity

matrix, denoted as W, was initialized in a manner such that all outgoing synapses from excitatory neurons had

equal strength, with $W_{EE} = W_{IE} = J$. Similarly, the inhibitory neurons had equal strength for their outgoing

synapses, with $W_{II} = W_{EI} = -gJ$. Here, the balance parameter, g, plays a crucial role in tuning the model towards

an E/I-balanced state exhibiting critical dynamics. For our network, the critical value of g, g_c, was determined to

be 3.5. When g exceeds g_c, the network shifts towards an inhibition-dominated regime with subcritical

dynamics, whereas values of g below g_c cause the network to exhibit an excitation-dominated regime with

supercritical dynamics.

Model Dynamics

Each neuron in the model was described by two variables. The Boolean variable, X, indicated whether a neuron

was firing or in a quiescent state at time t. Specifically, X(t) = 1 represented firing, while X(t) = 0 denoted

quiescence at time t. The membrane potential, V, influenced the firing probability and evolved according to the

following equation:

374
$$V_i(t + 1) = \left(\mu V_i(t) + \sum_{j=1}^{N} W_{ij} X_j(t) \right)(1 - X_i(t))$$

375 In the equation, μ represented the leakage parameter, which we set to 0 (non-leaky) for our simulations. The

376 term $(1 - X_i(t))$ introduced an absolute refractory period of Δt, allowing the voltage to reset after a spike

377 occurred (when $X_i(t) = 1$). Additionally, the probability of a neuron firing linearly increased with V and was given

378 by:

379
$$P(X = 1 \mid V) = \begin{cases} V, \Gamma V < 1 \\ 1, \Gamma V \geq 1 \end{cases}$$

Here, Γ represented the neuronal gain, which we fixed at 1 for simplicity and supports probabilistic behavior.

Notably, this network exhibited significant non-conservativeness, with energy dissipation occurring through

inhibition and spike collision or spike coalescence[79,80].

Additionally, we note that this model includes an absorbing quiescent state, and therefore, the network activity

is initiated by very low-rate independent Poisson processes applied to each neuron. Specifically, the Poisson

process triggers approximately ~20 neurons (0.002%) on average per time step, which is negligible compared to

the system size of 10^6 neurons.

Avalanche Statistics

To comprehensively investigate the resulting population activity, we run 100 simulations of the network over an

extended duration of 10^6 time steps for a total of 10^8 time steps. For the case of full sampling, we compute the

sum of activity across the entire population of $N = 10^6$ neurons. However, to explore various sampling scenarios,

we consider a random subset of $N*f$ neurons, where f represents the sampling fraction. Following this, a

thresholding procedure is applied to the time-series data, setting all neuronal activity below the threshold value

θ to 0. We have the option to analyze the data within the thresholded time-series in their original form or

subject them to temporal coarse graining. When employing coarse graining, each generation represents the sum

of activity over k consecutive time bins, with k representing the coarse graining factor. This approach enables us

to investigate network dynamics across different temporal scales and analyze resulting patterns. Consequently,

we collect the sizes and durations of contiguous epochs from the time-series and examine their distributions.

Power-law fitting

In the context of an ideal, infinitely sized critical system, the distribution of size, duration, and average size as a

function of duration adheres to perfect power laws. However, it is important to consider statistical anomalies,

finite size effects, and the influence of the low Poisson drive in our simulations when analyzing these

distributions. Therefore, a cautious approach is required for their analysis within the model. In the average size

versus duration analyses, we observed that the distributions could be reasonably approximated by a double

power law. For this purpose, we employ the fitting function introduced by Capek & Ribeiro et al.[9]:

$$S(d) = \frac{C d^{\sigma_1}}{(1 + (d/\Phi)^\gamma)^{(\sigma_1 + \sigma_2)/\gamma}}$$

This function exhibits an initial slope σ_1 that transitions to a second slope σ_2 around the point $d = \Phi$. The

parameter γ governs the abruptness of this transition and has been set to 4 for all the analysis presented. The

remaining parameters were allowed to adjust to the data, and all fits were performed in log-space. In all plots

presented in this paper, σ_1 is employed as the representative slope, while σ_2, which is typically associated with

finite-size effects, is not discussed in this context.

We use the cutoff, Φ, that we obtain from the scaling curves to analyze the individual size and duration

distributions. The size exponent, α, is the slope of the size distribution in log-scale with a lower cutoff which is

safely set to be greater than the Poisson noise level (>100) and an upper cutoff equal to the average size

corresponding to the elbow point, Φ. For the duration exponent, β, we do a similar analysis, with a lower

duration cutoff of 3 time-bins and an upper cutoff equal to Φ. However, for the sake of simplicity and due to the

small variability of the fitting procedure for the experimental data, we choose to utilize only the first 4

generations to calculate estimates for all the exponents for Figure 5. Furthermore, we only consider exponents

for parameters which have at least one avalanche that exceeds 20 generations and cases that do not satisfy this

criterion are ignored.

Collapse exponent of threshold and fractional sampling

In our analysis of the effects of temporal coarse graining on thresholding and fractional sampling, we collapsed

the χ vs. k curves for k within the range $1 - 100$. In the fully (100%) sampled model, we examined various

thresholds (θ_{100}) ranging from 10^2 to 10^5. In the case of fractionally sampled models, the threshold was adjusted

proportionally to an exponent, ξ, of the sampling fraction, f, according to the equation $\theta = \theta_{100} * f^{\xi}$. For each ξ

value spanning from 0 to 2 in steps of 0.1, we computed the collapse error of the five curves representing

sampling rates from 0.01% to 100% of χ vs. k. The collapse exponent ξ is determined by the minimum collapse

error and was found to be close to 1.

Correlation analysis

In our examination of the mean delayed pairwise correlation within the system, we focus on a smaller temporal

segment of the simulated network (consisting of 10,000 time steps) to facilitate analysis of large neuron raster

datasets. For each neuron pair (designated as an ordered pair [i,j]) within the network, we compute the time-

lagged correlation between the activity of neuron *i* and that of neuron *j*, shifted by one time step, $k\Delta t$, $C_{ij}(k\Delta t)$.

The mean delayed correlation, $\bar{C}(k)$, of the system is then determined as the average correlation coefficient

computed for all such neuron pairs. Hence $\bar{C}(k) = \langle C_{ij}(k\Delta t)\rangle$. This correlation analysis spans temporal bins, *k*,

ranging from 1 to 100 and spatial sampling values from 0.01% up to 1%. Due to the quadratic increase in

duration and memory requirements with system size, higher sampling (10% sampling or more) values become

impractical for storing raster data and conducting pairwise analyses.

Ongoing activity in contralateral ACC/mPFC monitored with jRGECO1a

All experiments have been previously reported in Capek & Ribeiro et al.[9] and are available online (see Data

Availability). No additional experiments were conducted for the present study. In short, mice (C57BL/6; Jackson

Laboratories; age >6 weeks) were injected with a viral construct to express jRGECO1a in cortical neurons using

Syn promotor. Chronic 2PI started after ~2 weeks in the contralateral ACC/mPFC at an estimated depth of ~150–

300 μm using a microprism. Recordings were collected over the course of several days from n = 5 mice (3 males,

2 females; age 8–20 weeks) with jRGECO1a expression (n = 17 recordings; 30 min each). Recordings were

conducted over the course of several weeks and analyzed separately for each mouse. 2PI images were acquired

by a scanning microscope (Bergamo II series, B248, Thorlabs Inc.) coupled to a pulsed femtosecond Ti:Sapphire

2-photon laser (Chameleon Discovery NX, Coherent Inc.). The wavelength was tuned to 1120 nm to excite

jRGECO1a. The field of view was ~450 um x 450 μm. Imaging frames of 512 x 512 pixels were acquired at 45.527

Hz by bidirectional scanning of a 12 kHz Galvo-resonant scanner. The average power for imaging was <70 mW,

measured at the sample. Obtained tif-movies in uint16 format were rigid motion corrected via the python-based

software package suite2p[81]. Registered images were denoised using machine-learning-based,

DeepInterpolation[82] and then semi-automatically processed by suite2p for ROI selection and fluorescence signal

extraction. For each labeled neuron, raw soma and neuropil fluorescence signals over time were extracted for

each ROI. Spiking probabilities were obtained from neuropil corrected fluorescence traces (F_corrected = F_ROI

– 0.7*F_neuropil) via MLspike (https://github.com/MLspike) by utilizing its autocalibration feature to obtain

unitary spike event amplitude, decay time, and channel noise for individual ROIs.

ACKNOWLEDGEMENTS

This research was supported by the Division of the Intramural Research Program (DIRP) of the National Institute

of Mental Health (NIMH), USA, ZIAMH002797, ZIAMH002971, and the BRAIN initiative Grant U19 NS107464-01

to D.P. This research utilized the supercomputing resources of the National Institutes of Health (NIH, USA;

Biowulf, http://hpc.nih.gov) and the University of Maryland (UMD College Park, USA,

https://hpcc.umd.edu/hpcc/dt2.html).

AUTHOR CONTRIBUTIONS

K.S., T.L.R., D. P. conceived the study, K.S. did the simulations, P.K. provided the experimental data and

processing, K.S., T.L.R., D. P. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY

The preprocessed imaging data used in this study are available in the general repository Zenodo using the

following access https://doi.org/10.5281/zenodo.7703224 under 'jRGECO1a-ongoing'.

CODE AVAILABILITY

The code used to generate the model data is available at https://github.com/PlenzLab/ParabolicAvalanches/.

482 REFERENCES

483   1   Hamon, D., Nicodemi, M. & Jensen, H. Continuously driven OFC: A simple model of solar flare
484   statistics. *Astronomy & Astrophysics* **387**, 326-334 (2002).
485   2   Olami, Z., Feder, H. J. S. & Christensen, K. Self-organized criticality in a continuous,
486   nonconservative cellular automaton modeling earthquakes. *Phys.Rev.Lett.* **68**, 1244-1247 (1992).
487   3   Bak, P. & Tang, C. Earthquakes as a self-organized critical phenomenon. *Journal of Geophysical*
488   *Research: Solid Earth* **94**, 15635-15637 (1989).
489   4   Bassler, K. E. & Paczuski, M. in *Complexity from Microscopic to Macroscopic Scales:*
490   *Coherence and Large Deviations*   215-227 (Springer, 2002).
491   5   Held, G. A. *et al.* Experimental study of critical-mass fluctuations in an evolving sandpile. *Phys.*
492   *Rev. Lett.* **65**, 1120 (1990).
493   6   Drossel, B. & Schwabl, F. Self-organized critical forest-fire model. *Phys.Rev.Lett.* **69**, 1629-1632
494   (1992).
495   7   Beggs, J. M. & Plenz, D. Neuronal avalanches in neocortical circuits. *J. Neurosci.* **23**, 11167-
496   11177, doi:10.1523/JNEUROSCI.23-35-11167.2003 (2003).
497   8   Bellay, T., Klaus, A., Seshadri, S. & Plenz, D. Irregular spiking of pyramidal neurons organizes
498   as scale-invariant neuronal avalanches in the awake state. *eLife* **4**, e07224,
499   doi:10.7554/eLife.07224 (2015).
500   9   Capek, E. *et al.* Parabolic avalanche scaling in the synchronization of cortical cell assemblies.
501   *Nature communications* **14**, 2555, doi:10.1101/2022.11.02.514938 (2023).
502   10   Scarpetta, S., Apicella, I., Minati, L. & de Candia, A. Hysteresis, neural avalanches, and critical
503   behavior near a first-order transition of a spiking neural network. *Phys. Rev. E* **97**, 062305,
504   doi:10.1103/PhysRevE.97.062305 (2018).
505   11   Scarpetta, S. in *The Functional Role of Critical Dynamics in Neural Systems*   (eds Nergis
506   Tomen, J. Michael Herrmann, & Udo Ernst) 179-197 (Springer International Publishing, 2019).
507   12   Liang, J. & Zhou, C. Criticality enhances the multilevel reliability of stimulus responses in
508   cortical neural networks. *PLoS Comput. Biol.* **18**, e1009848, doi:10.1371/journal.pcbi.1009848
509   (2022).
510   13   Villegas, P., Di Santo, S., Burioni, R. & Muñoz, M. A. Time-series thresholding and the
511   definition of avalanche size. *Phys. Rev. E* **100**, doi:10.1103/PhysRevE.100.012133 (2019).
512   14   Morowitz, H. J. *The mind, the brain and complex adaptive systems*. (Routledge, 1994).
513   15   Holland, J. H. *Signals and boundaries: Building blocks for complex adaptive systems*. (Mit
514   Press, 2012).
515   16   Lombardi, F., Herrmann, H., Plenz, D. & de Arcangelis, L. On the temporal organization of
516   neuronal avalanches. *Front. Syst. Neurosci.* **8**, 204, doi:10.3389/fnsys.2014.00204 (2014).
517   17   Lombardi, F., Herrmann, H. J., Plenz, D. & de Arcangelis, L. Temporal correlations in neuronal
518   avalanche occurrence. *Scientific Reports* **6**, 24690, doi:10.1038/srep24690 (2016).
519   18   Plenz, D. *et al.* Self-Organized Criticality in the Brain. *FrPhy* **9**, doi:10.3389/fphy.2021.639389
520   (2021).
521   19   Petermann, T. *et al.* Spontaneous cortical activity in awake monkeys composed of neuronal
522   avalanches. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 15921-15926, doi:10.1073/pnas.0904089106
523   (2009).
524   20   Miller, S. R., Yu, S. & Plenz, D. The scale-invariant, temporal profile of neuronal avalanches in
525   relation to cortical γ–oscillations. *Scientific Reports* **9**, 16403, doi:10.1101/757278 (2019).
526   21   Yu, S. *et al.* Higher-order interactions characterized in cortical activity. *J. Neurosci.* **31**, 17514-
527   17526, doi:10.1523/JNEUROSCI.3127-11.2011 (2011).

22   Bellay, T., Shew, W. L., Yu, S., Falco-Walter, J. J. & Plenz, D. Selective participation of single
cortical neurons in neuronal avalanches. *Frontiers in Neural Circuits* **14**,
doi:10.3389/fncir.2020.620052 (2021).
23   Shriki, O. & Plenz, D. in *Criticality in Neural Systems* Vol. 7 *Reviews of Nonlinear Dynamics
and Complexity* (eds D. Plenz & E. Niebur) Ch. 8, 177 - 189 (Wiley-VCH, 2014).
24   Palva, J. M. *et al.* Neuronal long-range temporal correlations and avalanche dynamics are
correlated with behavioral scaling laws. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 3585-3590 (2013).
25   Gireesh, E. D. & Plenz, D. Neuronal avalanches organize as nested theta- and beta/gamma-
oscillations during development of cortical layer 2/3. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 7576-
7581 (2008).
26   Meisel, C. Antiepileptic drugs induce subcritical dynamics in human cortical networks. *Proc.
Natl. Acad. Sci. U. S. A.* **117**, 11118, doi:10.1073/pnas.1911461117 (2020).
27   Maturana, M. I. *et al.* Critical slowing down as a biomarker for seizure susceptibility. *Nature
communications* **11**, 2172, doi:10.1038/s41467-020-15908-3 (2020).
28   Meisel, C., Plenz, D., Schulze-Bonhage, A. & Reichmann, H. Quantifying antiepileptic drug
effects using intrinsic excitability measures. *Epilepsia* **57**, e210-e215, doi:10.1111/epi.13517
(2016).
29   Meisel, C. *et al.* Intrinsic excitability measures track antiepileptic drug action and uncover
increasing/decreasing excitability over the wake/sleep cycle. *Proc. Natl. Acad. Sci. U. S. A.* **112**,
14694-14699, doi:10.1073/pnas.1513716112 (2015).
30   Ribeiro, S. & Nicolelis, M. A. Reverberation, storage, and postsynaptic propagation of memories
during sleep. *Learn.Mem.* **11**, 686-696 (2004).
31   Ribeiro, S. *et al.* Long-lasting novelty-induced neuronal reverberation during slow-wave sleep in
multiple forebrain areas. *PLoS.Biol.* **2**, E24 (2004).
32   Meisel, C., Klaus, A., Vyazovskiy, V. V. & Plenz, D. The interplay between long- and short-
range temporal correlations shapes cortex dynamics across vigilance states. *J. Neurosci.* **37**,
10114, doi:10.1523/JNEUROSCI.0448-17.2017 (2017).
33   Meisel, C., Bailey, K., Achermann, P. & Plenz, D. Decline of long-range temporal correlations in
the human brain during sustained wakefulness. *Scientific Reports* **7**, 11825, doi:10.1038/s41598-
017-12140-w (2017).
34   Meisel, C., Olbrich, E., Shriki, O. & Achermann, P. Fading signatures of critical brain dynamics
during sustained wakefulness in humans. *J. Neurosci.* **33**, 17363-17372,
doi:10.1523/JNEUROSCI.1516-13.2013 (2013).
35   Meisel, C., Storch, A., Hallmeyer-Elgner, S., Bullmore, E. & Gross, T. Failure of adaptive self-
organized criticality during epileptic seizure attacks. *PLoS Comput. Biol.* **8**, e1002312 (2012).
36   Ribeiro, T. L., Ribeiro, S., Belchior, H., Caixeta, F. & Copelli, M. Undersampled critical
branching processes on small-world and random networks fail to reproduce the statistics of spike
avalanches. *PloS one* **9**, e94992, doi:10.1371/journal.pone.0094992 (2014).
37   Destexhe, A. & Touboul, J. D. Is there sufficient evidence for criticality in cortical systems?
eneuro, ENEURO.0551-0520.2021, doi:10.1523/ENEURO.0551-20.2021 (2021).
38   Touboul, J. & Destexhe, A. Power-law statistics and universal scaling in the absence of
criticality. *Phys. Rev. E* **95**, doi:10.1103/PhysRevE.95.012413 (2017).
39   Carvalho, T. T. A. *et al.* Subsampled directed-percolation models explain scaling relations
experimentally observed in the brain. *Frontiers in Neural Circuits* **14**,
doi:10.3389/fncir.2020.576727 (2021).
40   Fontenele, A. J. *et al.* Criticality between cortical states. *Phys. Rev. Lett.* **122**, 208101,
doi:10.1103/PhysRevLett.122.208101 (2019).

41   Levina, A. & Priesemann, V. Subsampling scaling. *Nature communications* **8**, 15140,
doi:10.1038/ncomms15140 (2017).
42   Xu, Y., Schneider, A., Wessel, R. & Hengen, K. B. Sleep restores an optimal computational
regime in cortical networks. *bioRxiv*, 2022.2012.2007.519478, doi:10.1101/2022.12.07.519478
(2022).
43   Mariani, B. *et al.* Neuronal avalanches across the rat somatosensory barrel cortex and the effect
of single whisker stimulation. *Front. Syst. Neurosci.* **15**, doi:10.3389/fnsys.2021.709677 (2021).
44   Ma, Z., Liu, H., Komiyama, T. & Wessel, R. Stability of motor cortex network states during
learning-associated neural reorganizations. *J. Neurophysiol.* **124**, 1327-1342,
doi:10.1152/jn.00061.2020 (2020).
45   Karimipanah, Y., Ma, Z., Miller, J.-e. K., Yuste, R. & Wessel, R. Neocortical activity is
stimulus- and scale-invariant. *PloS one* **12**, e0177396, doi:10.1371/journal.pone.0177396 (2017).
46   Friedman, N. *et al.* Universal critical dynamics in high resolution neuronal avalanche data. *Phys.*
Rev. Lett. **108**, 208102 (2012).
47   Levina, A., Priesemann, V. & Zierenberg, J. Tackling the subsampling problem to infer
collective properties from limited data. *Nature Reviews Physics* **4**, 770-784, doi:10.1038/s42254-
022-00532-5 (2022).
48   Girardi-Schappo, M., Brochini, L., Costa, A. A., Carvalho, T. T. A. & Kinouchi, O. Synaptic
balance due to homeostatically self-organized quasicritical dynamics. *Physical Review Research*
2, 012042, doi:10.1103/PhysRevResearch.2.012042 (2020).
49   Girardi-Schappo, M. *et al.* A unified theory of E/I synaptic balance, quasicritical neuronal
avalanches and asynchronous irregular spiking. *Journal of Physics: Complexity* **2**, 045001,
doi:10.1088/2632-072X/ac2792 (2021).
50   Scarpetta, S. *et al.* Criticality of neuronal avalanches in human sleep and their relationship with
sleep macro- and micro-architecture. *iScience* **26**, 107840, doi:10.1016/j.isci.2023.107840
(2023).
51   Mehta, A. P., Mills, A. C., Dahmen, K. A. & Sethna, J. P. Universal pulse shape scaling function
and exponents: Critical test for avalanche models applied to Barkhausen noise. *Phys. Rev. E* **65**,
doi:10.1103/PhysRevE.65.046139 (2002).
52   Kuntz, M. C. & Sethna, J. P. Noise in disordered systems: The power spectrum and dynamic
exponents in avalanche models. *PhRvB* **62**, 11699-11708 (2000).
53   Jensen, H. J. *Self-organized criticality*. (Cambridge University Press, 1998).
54   Chen, Y.-J., Papanikolaou, S., Sethna, J. P., Zapperi, S. & Durin, G. Avalanche spatial structure
and multivariable scaling functions: Sizes, heights, widths, and views through windows. *Phys.*
Rev. E **84**, 061103 (2011).
55   Magni, A., Durin, G., Zapperi, S. & Sethna, J. P. Visualization of avalanches in magnetic thin
films: temporal processing. *JSMTE* **2009**, P01020 (2009).
56   Ma, Z., Turrigiano, G. G., Wessel, R. & Hengen, K. B. Cortical circuit dynamics are
homeostatically tuned to criticality in vivo. *Neuron* **104**, 655-664,
doi:10.1016/j.neuron.2019.08.031 (2019).
57   Burrows, D. R. *et al.* Microscale neuronal activity collectively drives chaotic and inflexible
dynamics at the macroscale in seizures. *J. Neurosci.* **43**, 3259-3283 (2023).
58   Muñoz, M. A., Dickman, R., Vespignani, A. & Zapperi, S. Avalanche and spreading exponents
in systems with absorbing states. *Phys. Rev. E* **59**, 6175-6179 (1999).
59   Priesemann, V., Munk, M. H. & Wibral, M. Subsampling effects in neuronal avalanche
distributions recorded in vivo. *BMC.Neurosci.* **10**, 40 (2009).

60 Muñoz, M. A. Colloquium: Criticality and dynamical scaling in living systems. *Rev. Mod. Phys.*
90, 031001, doi:10.1103/RevModPhys.90.031001 (2018).
61 Cocchi, L., Gollo, L. L., Zalesky, A. & Breakspear, M. Criticality in the brain: A synthesis of
neurobiology, models and cognition. *Prog. Neurobiol.*, doi:10.1016/j.pneurobio.2017.07.002
(2017).
62 Hesse, J. & Gross, T. Self-organized criticality as a fundamental property of neural systems.
Front. Syst. Neurosci. **8**, doi:10.3389/fnsys.2014.00166 (2014).
63 Chialvo, D. R. Emergent complex neural dynamics. *Nat. Phys.* **6**, 744-750, doi:DOI:
10.1038/NPHYS1803 (2010).
64 Nonnenmacher, M., Behrens, C., Berens, P., Bethge, M. & Macke, J. H. Signatures of criticality
arise from random subsampling in simple population models. *PLoS Comput. Biol.* **13**, e1005718
(2017).
65 Pausch, J. From neuronal spikes to avalanches: Effects and circumvention of time binning.
Physical Review Research **4**, 023212, doi:10.1103/PhysRevResearch.4.023212 (2022).
66 Aitchison, L., Corradi, N. & Latham, P. E. Zipf's law arises naturally when there are underlying,
unobserved variables. *PLoS Comput. Biol.* **12**, e1005110, doi:10.1371/journal.pcbi.1005110
(2016).
67 Priesemann, V. & Shriki, O. Can a time varying external drive give rise to apparent criticality in
neural systems? *PLoS Comput. Biol.* **14**, e1006081 (2018).
68 Morrell, M. C., Nemenman, I. & Sederberg, A. Neural criticality from effective latent variables.
eLife **12**, RP89337, doi:10.7554/eLife.89337 (2024).
69 Plenz, D. Neuronal avalanches and coherence potentials. *The European Physical Journal Special
Topics* **205**, 259-301, doi:10.1140/epjst/e2012-01575-5 (2012).
70 Shew, W. L., Yang, H., Yu, S., Roy, R. & Plenz, D. Information capacity is maximized in
balanced cortical networks with neuronal avalanches. *J. Neurosci.* **5**, 55-63,
doi:10.1523/JNEUROSCI.4637-10.2011 (2011).
71 Stumpf, M. P., Wiuf, C. & May, R. M. Subnets of scale-free networks are not scale-free:
sampling properties of networks. *Proc.Natl.Acad.Sci.U.S.A* **102**, 4221-4224 (2005).
72 Serafino, M. *et al.* True scale-free networks hidden by finite size effects. *Proc. Natl. Acad. Sci.
U. S. A.* **118**, e2013825118, doi:10.1073/pnas.2013825118 (2021).
73 Bonzanni, M., Bockley, K. M. & Kaplan, D. L. On the effect of neuronal spatial subsampling in
small-world networks. *Eur. J. Neurosci.* **53**, 485-498, doi:10.1111/ejn.14937 (2021).
74 Ribeiro, T. L. *et al.* Trial-by-trial variability in cortical responses exhibits scaling of spatial
correlations predicted from critical dynamics. *Cell Reports* **43** (2024).
75 Averbeck, B. B., Latham, P. E. & Pouget, A. Neural correlations, population coding and
computation. *Nat. Rev. Neurosci.* **7**, 358-366 (2006).
76 Gawne, T. J., Kjaer, T. W., Hertz, J. A. & Richmond, B. J. Adjacent visual cortical complex cells
share about 20% of their stimulus-related information. *Cereb. Cortex* **6**, 482-489 (1996).
77 Smith, M. A. & Kohn, A. Spatial and Temporal Scales of Neuronal Correlation in Primary
Visual Cortex. *J. Neurosci.* **28**, 12591, doi:10.1523/JNEUROSCI.2929-08.2008 (2008).
78 Schulz, D. P. A., Sahani, M. & Carandini, M. Five key factors determining pairwise correlations
in visual cortex. *J. Neurophysiol.* **114**, 1022-1033, doi:10.1152/jn.00094.2015 (2015).
79 Zierenberg, J., Wilting, J., Priesemann, V. & Levina, A. Description of spreading dynamics by
microscopic network models and macroscopic branching processes can differ due to coalescence.
Phys. Rev. E **101**, 022301 (2020).

80   Zierenberg, J., Wilting, J., Priesemann, V. & Levina, A. Tailored ensembles of neural networks
optimize sensitivity to stimulus statistics. *Physical Review Research* **2**, 013115,
doi:10.1103/PhysRevResearch.2.013115 (2020).
81   Pachitariu, M. *et al.* Suite2p: beyond 10,000 neurons with standard two-photon microscopy.
bioRxiv, 061507, doi:10.1101/061507 (2017).
82   Lecoq, J. *et al.* Removing independent noise in systems neuroscience data using
DeepInterpolation. *Nat. Methods*, doi:10.1038/s41592-021-01285-2 (2021).

The recovery of parabolic avalanches in spatially subsampled neuronal networks at criticality

SUPPLEMENTARY INFORMATION

Keshav Srinivasan, Tiago L. Ribeiro, Patrick Kells, and Dietmar Plenz[†]

Section on Critical Brain Dynamics, National Institute of Mental Health, Bethesda, MD 20892, USA

[†] Correspondence

Suppl. Fig. S1. Consolidated view of the exponents α, β and χ as a function of threshold and temporal coarse-graining for different sampling fractions.

Consolidated view of α, β and χ as a function of θ and k for different values of f. For f = 100%, $\chi \cong 2$ for low θ and k (*star*) but as we make the data sparser by increasing the threshold (or reducing the sampling), we need a higher coarse graining factor, k, to compensate and rescue χ back to 2 (*square*). Bottom row is replotted for χ from Figure 4 for ease of comparison with the corresponding size slopes, *α*, and duration slopes, *β*. *White dotted region*: visual guide for χ close to 2 for increasing θ and k. The grey parts of the plots for f = 0.1% and 0.01% respectively are unobservable parameter regions since they would require fractional thresholds, below the 1 spike minimum resolution of the model (*cf.* Figure 4).