The recovery of parabolic avalanches in spatially subsampled neuronal networks at criticality

Keshav Srinivasan, Tiago L. Ribeiro, Patrick Kells, and Dietmar Plenz[†]

Section on Critical Brain Dynamics, National Institute of Mental Health, Bethesda, MD 20892, USA

[†] Correspondence

Figures: 5 Figures, 2 Supplemental Figures

Correspondence: Dietmar Plenz, Ph.D., Section on Critical Brain Dynamics, National Institute of Mental Health, Porter Neuroscience Research Center, Rm 3A-1000, 35 Convent Drive, Bethesda, MD 20892. Office: (301) (301) 675-5311, plenzd@mail.nih.gov

Keywords: Neuronal avalanches, E/I balanced neural network, subsampling, thresholding, scaling exponent, criticality, ongoing activity, mouse, frontal cortex, 2-photon imaging

ABSTRACT

Scaling relationships characterize complex systems at criticality. In the brain, these relationships are evident

in scale-invariant activity cascades, so-called neuronal avalanches, quantified by power laws in avalanche size

and duration. At the cellular level, neuronal avalanches are identified in spatially distributed groups of

neurons that participate in cascades of coincident action potential firing. Such spatiotemporal synchronization

is central to theories on brain function, yet scaling relationships in avalanche synchronization have been

challenging to study when only a fraction of neurons is observed, underestimating avalanche properties.

Here, we study these biases from fractional sampling in an all-to-all, balanced network of excitatory and

inhibitory neurons with critical branching process dynamics. We focus on the growth of mean avalanche size

with avalanche duration. For parabolic avalanches, this growth is quadratic, quantified by the scaling

exponent, χ = 2, which signifies rapid spatial expansion of coincident firing within a relatively short period of

time. In contrast, χ << 2 for fractionally sampled networks. We show that temporal coarse-graining combined

with a threshold for the minimally required coincident firing in the network recovers χ = 2, even when

sampling as few as 0.1% of the neurons. In contrast, a commonly proposed 'crackling noise' approach fails to

recover χ under those conditions. Our approach robustly identifies χ = 2 for ongoing neuronal activity in

frontal cortex of awake mice using cellular 2-photon imaging. Our findings demonstrate how to correct scaling

bias from fractional sampling and identifies rapid, scale-invariant synchronization of cell assemblies in the

brain.

AUTHOR SUMMARY

In the brain, groups of neurons often fire together which is considered essential for normal brain function. A

particular form of coincident firing has been identified for "neuronal avalanches" which are scale-invariant

cascades of neuronal activity that rapidly engage large numbers of neurons. However, studying these avalanches

when observing only a few neurons in the network is challenging. We used simulations and recordings from

ongoing brain activity to explore how to overcome this challenge. We found that adjusting our temporal

resolution and focusing on increasingly larger avalanches allows us to accurately detect rapid, parabolic

expansions of these patterns even when examining a minuscule fraction of neurons. This identifies a general

rule for how the brain operates best in engaging neurons to fire together, despite the limitations in our

observations.

INTRODUCTION

Complex systems, composed of many local components or agents that interact weakly, often exhibit event

cascades that cover a wide range of scales in both space and time. Such scale-invariant cascades, typically

identified by power-law distributions in their duration and size – defined as the total activity observed during its

lifetime, have been observed in many real systems, including solar flares [1], earthquakes [2, 3],

superconductors [4], sandpiles [5], forest fires [6], and in the brain in the form of neuronal avalanches [7-9]. The

scaling of such system cascades, specifically how their mean size grows with their duration, has been particularly

informative as to the potential underlying dynamics of cascade generation. For example, if local system

components activate independently at a constant rate, the mean size, <S>, of observed cascades will scale

linearly with duration, T, exhibiting a scaling exponent of $\chi \cong 1$, i. e., temporally contiguous system events reflect

a constant, yet random probability of their continuation. Values of $\chi \cong 1$ are often found near a 1st order phase

transition [10-12]. Conversely, if cascades exhibit $\chi = 2$, their mean growth is rapid and parabolic as can be found

at the second-order phase transition of a critical branching process, supporting local system interactions [13]. In

general, values of χ >>1 are of particular interest as they suggest the ability of a system to bind spatially

distributed events rapidly and selectively into diverse internal system configurations which is generally

considered beneficial for a complex adaptive system such as the brain (e. g., [14, 15]).

Here, we focus on neuronal avalanches, which were first shown to spontaneously emerge in isolated

brain networks, i. e. acute cortex slices and long-term cortex cultures [7]. By measuring locally synchronized

neuronal activity using the local field potential (LFP) and tracking the spatiotemporal spread of LFP cascades

with microelectrode arrays, power-law distributions for both size, *S*, and duration, *T*, $P(S) \sim S^{-\alpha}$ and

$P(T) \sim T^{-\beta}$ were revealed, with exponents α ≈ 3/2 and β ≈ 2 respectively, in addition to a critical branching

parameter of σ ≅ 1, and significant correlation of neuronal activity within and between [16, 17]neuronal

cascades (for review see [18]). These hallmarks of fast, neuronal synchronization in the form of avalanches were

subsequently confirmed in the normal brain for ongoing and evoked LFP cascades in the cortex of nonhuman

primates [19-22] and in humans using magnetoencephalography [23, 24]. Their sensitivity to even slight

pharmacological alterations in the balance of excitation and inhibition [7, 18, 22, 25] and changes during

pathological conditions such as epilepsy [26-29] or changes in brain states such as sleep [30, 31] and sleep

deprivation [32-35], firmly indicate that these avalanches represent neuronal synchronization of cell assemblies

in the critically tuned, normal brain.

To understand avalanche synchronization mechanistically, the contribution of single neuron firing to

avalanches needs to be properly identified. This task is extremely challenging as studies in nonhuman primates

and isolated slice preparations show that neurons participate sparsely and highly selectively in even spatially

extended LFP avalanches [22]. Accurate reconstruction of avalanche synchronization from single neuron activity,

therefore, requires the appropriate sampling of a significant portion of the brain network which poses an

enormous technical challenge in neuroscience. Conversely, spatial subsampling, that is, sampling of only a

fraction of neurons, is limited in identifying neuronal synchronization largely by failing to identify avalanche

continuation in non-sampled neurons. This underestimates avalanche duration and size, biasing outcomes

towards expectations for decorrelated, random processes [9, 36, 37]. Accordingly, simulations of avalanche

generating critical branching processes and fractional sampling of neurons reduces the expected growth in mean

avalanche size with avalanche duration, χ, from $\chi = 2$, to a value of χ closer to $1 - 1.3$ [38-40]. Not surprisingly,

estimates of $\chi \cong 1.3$ were reported for fractionally sampled neuronal spikes of potential avalanche activity *in*

vivo [39, 41-43] and *in vitro* [44].

If the exact start and end times of avalanches are known, for example in simulations under the

assumption of 'separation of time scales', spatially subsampled avalanches can be easily corrected for by

temporal integration thereby recovering the correct critical exponents in size and duration [40] (for a review,

see also [45]). Avalanche times, however, are largely unattainable in experimental data for which only a small

fraction of neurons is observed and temporal integration eventually will degenerate avalanche scaling by

indiscriminately concatenating successive avalanches [9]. Recently, Capek & Ribeiro et al. [9] demonstrated that,

even when the exact times of avalanches are not known, $\chi = 2$ can be recovered for subsampled excitation-

inhibition (E-I) balanced networks of integrate-and-fire neurons which are known to belong to the mean-field

directed percolation (MF-DP) universality class [46, 47]. This rescue requires (1) reducing uncorrelated neuronal

action potential firing and (2) using temporal coarse-graining with an increase in minimal synchronization

requirements to identify synchronized population activity in the network. Under these conditions, the authors

could show $\chi = 2$ *in vivo* 2-photon imaging recordings of cellular activity with high signal-to-noise ratio for

sensory evoked and spontaneous spike avalanches in frontal and sensory cortices of awake mice in line with

their simulations [9]. This is in line with a recent EEG study in humans that demonstrated $\chi = 2$ [48].

In the current work, we extend our previous investigations of rescuing χ in subsampled, critical

networks. We systematically study the rescue of the scaling exponent χ to its critical value of 2 in a continuously

driven, subsampled all-to-all DP-model [46] for which we demonstrate the equivalence of fractional sampling

and thresholding. This allows us to identify a robust domain within the 'temporal coarse graining' vs.

thresholding parameter regime over 4 orders of magnitude of fractional sampling. We identify this domain in

cellular 2-photon imaging data of the frontal cortex of awake mice. We further show that the 'crackling noise'

relationship and related measures which can predict χ [49-51] fail under conditions of subsampling. Our findings

demonstrate the pitfalls of assessing avalanche synchronization when only parts of a critical network are

observed and demonstrate the regime of potential scaling rescue for synchronization in the form of parabolic

avalanches under experimental conditions of fractional sampling.

RESULTS

Neuronal avalanches were studied using a critically balanced Excitatory-Inhibitory (E-I) model with all-to-all

connectivity consisting of one million neurons driven externally using independent, stationary Poisson processes

[46] (**Fig. 1A**). The external driving triggered an average of 20 neurons to fire per time step, Δt, which

intermittently induced spreading of neuronal activity in the network. Simulations spanned across 10^8 time steps.

An avalanche was defined as a series of contiguous time periods, T, of aggregated neuronal activity surpassing a

predetermined threshold, θ, over a specific number of time steps, k (**Fig. 1B – D**). Our analytical approach was

characterized by two basic parameters: $k \cdot \Delta t$ and θ, which were explored in the context of the subsampling

fraction of observed neurons, f, which ranged from 1 out of every 10,000 neurons observed ($f = 0.01\%$) to all

neurons observed ($f = 100\%$). We focused in our analysis on continuous sequences of coincident neuronal firing

above θ, i. e., suprathreshold activity (**Fig. 1,B D**). We identified the statistical distribution of sequence size, S,

and duration, T, as a function of θ, f, and k and estimated the corresponding size exponent, α, and duration

exponent, β, within the constraints of the lower and upper cut-offs (**Fig. 1C**; see Methods; [52, 53]). In our

critical model, these distributions take on the form of power laws at $k = 1$, f = 100% and $\theta = 100$, the defining

characteristics of avalanches [7]. The scaling exponent, χ, governs the growth in average avalanche size, $\langle S \rangle$, for

a given duration, T,

$$\langle S \rangle \sim T^{\chi},$$ (1)

and accordingly, we estimated the scaling factor χ from the slope of the average avalanche size, $\langle S \rangle$, for a given

avalanche duration, T, outside the minimal and maximal cut-off in logarithmic scales (**Fig. 1C**, see

Figure 1. Schematics of the balanced E/I-model and identification of suprathreshold sequences in population spiking activity with respect to minimal coincident spiking threshold and temporal coarse graining. (A) Schematic of the neuronal network consisting of N = 10^6 neurons (80% excitatory and 20% inhibitory) with a low external Poisson drive of rate $\lambda = 2*10^{-5}$ per time step per neuron. The E/I balance is controlled by g, which scales the inhibitory weight matrices (W_{II}, W_{EI}) as a function of the excitatory weight matrices (W_{EE}, W_{IE}). If not otherwise stated, we set g = 3.5 to obtain critical dynamics (see Methods). (B) High variability in intermittent population activity characterizes critical dynamics. Snapshot of the summed neuronal spiking activity as a function of time. (C) Duration, size and scaling of avalanches in the critical model follow power-laws with corresponding exponent estimates α, β and χ. Note that the external Poisson drive and the finite size of the network introduce lower and higher cut-offs, respectively (*beige areas*). (D) Zoomed population activity from *B*. At the original temporal resolution Δt and given the coincident spiking threshold, θ (*left*), we can identify two sequences of suprathreshold activity, S_1 and S_2 with durations T_1 and T_2, respectively. Temporally coarse-graining the population activity (*right*; k = 5, binning the data into new bins of 5 time points) and increasing the threshold to $\theta' > \theta$, absorbs S_1 and S_2 into a new suprathreshold activity period S' with duration T'.

117   Methods). In critical systems, it has also been found that χ relates to the critical exponents α' and β' as

118 $$\chi_{cn} = \frac{(\beta' - 1)}{(\alpha' - 1)}$$ (2)

119   ('cn = crackling noise' [49-51]). For comparison, we therefore calculated both χ_{cn} and

120   DCC = χ - χ_{cn} (3)

from our slope estimates α and β. This latter construct has been suggested as a distance measure from a critical

point implying that the network dynamics is close to criticality with vanishing DCC (e.g. [54]).

High coincident firing thresholds underestimate χ in the fully sampled model

In systems characterized by a clear separation of timescales and an absence of noise, the necessity for a

coincident firing threshold diminishes. However, within real systems lacking such temporal separation and

affected by significant intrinsic and measurement noise, the statistical analysis becomes intricately tied to our

selection of threshold values. Setting the threshold too low poses the risk of incorporating substantial noise

from the system, while setting it too high leads to data sparsity, resulting in the exclusion of a significant portion

of the system's activity. To emulate the noisy environment and absence of timescales separation between

avalanche generation and propagation inherent in real systems, we introduce a minimal external Poisson drive

in our simulations. This Poisson drive establishes a noise floor within the spiking activity, serving as the origin of

avalanches within the system. This is clearly visible in the size distribution of avalanches, which exhibits a power

law only for avalanche sizes of S_0 > 50 – 100 spikes up to the finite-size of the network (*cf*. Fig. 1C, *middle*) [46].

To isolate avalanches above the noise floor (i. e., $S > S_0$), we first studied the dependence of α, β, χ on θ and k in

the fully sampled network (f = 100%; **Fig. 2**). In Fig. 2A, we show an overview in the form of color maps for all

three parameters when increasing θ. At the highest temporal resolution (k = 1), we observed that α and β,

increase with θ > 100, with a corresponding rapid decrease of χ below its expected value of 2 (cf. **Fig. 2B**). We

conclude that high thresholds in weakly driven, noisy networks bias towards longer suprathreshold periods,

significantly underestimating the scaling exponent for the critical system.

A similar underestimation holds true for the χ_{cn} value. Inserting the corresponding values of exponents

extracted from our fully sampled system for small thresholds, we obtain $\chi_{cn} = 2$, in line with expectations for

the scaling exponent for the mean-field directed percolation (MF-DP) universality class [55]. However, χ_{cn} drops

Figure 2. Increasing the threshold in the fully sampled model underestimates the scaling exponent, χ, which can be rescued by temporal coarse graining.

(**A**) Consolidated view of the avalanche size (α), duration (β), and scaling (χ) exponents as a function of threshold (θ) and temporal coarse-graining factor, k (f = 100%, fully sampled model). *Drive dominated*: Low threshold regime (*above the red broken line*) dominated by external Poisson drive. *White frames*: Parameter regions displayed in *B* and *C*. (**B**) At the highest temporal resolution (k=1), high thresholds underestimate the scaling exponent, χ, as well as χ_{CN}, with DCC remaining low. Size exponent, α (*left*), duration exponent, β (*middle*), scaling exponent, χ, and expected crackling noise relation, χ_{CN}, (*right*) as a function of θ. (**C**) At high threshold (θ=1000), temporal coarse graining recovers χ, but the χ_{CN} exhibits a singularity leading to high DCC. Size exponent, α (*left*), duration exponent, β (*middle*), as a function of coarse graining factor, k, (θ = 1000). α passes through 1 (*red broken line*; *black triangle*), which causes a singularity in χ_{CN}. (*right*). Note that the scaling exponent, χ, stabilizes to a value of 2 (*right, black dashed line*), whereas χ_{CN} grows until it passes through a singularity at the temporal coarse graining value of k = 17 (*vertical broken line*; *triangle*; see also *inset*).

significantly below 2 when θ increases, largely following the dependence of β on θ (*c.f.* **Fig. 2A**). We also observe

a small systematic underestimation of χ_{cn} with respect to χ (**Fig. 2B**), which we attribute largely to the error-

bounds of our fitting procedure. We conclude that avalanche scaling will be underestimated even in fully

sampled networks if there are technical constraints on minimal coincident spike counts, which require the use of

high thresholds.

Temporal coarse-graining rescues χ at high thresholds in the fully sampled model

Next, we show that temporal coarse graining rescues χ in the presence of a high threshold in fully sampled

networks. This is shown in **Fig. 2C**, in which χ approaches 2 for $k > 8$ at $\theta = 1000$ (see also corresponding

projections in **Fig. 2A**). In contrast, the size and duration distributions exhibit a gradual reduction in steepness

when k increases, due to concatenating subsequent cascades, in line with experimental findings [7, 19, 40, 56].

In the model, this eventually causes α to cross the value of 1 (**Fig. 2C**), which forces χ_{cn} to undergo a singularity

Figure 3. Equivalence in threshold and fractional sampling and increase in mean pair-wise correlation among neurons in the critical model with temporal coarse graining.
(**A**) Collapsed scaling exponent curves for different values of the sampling fraction, f, ranging from 100% to 0.1% for a threshold that is scaled by the sampling fraction as $\theta = 3,000*f$ (i.e., a threshold of 3,000 spikes for the fully sampled model translates to a threshold of 3 spikes for the 0.1% sampled model). *Inset*: The total collapse error as a function of collapse exponent, ξ. The error is taken over multiple different curves with θ ranging from 100 to 100,000. The error shows a clear minimum at $\xi = 1$, indicating the threshold can be scaled proportionally with the sampling fraction to obtain the best collapse.
(**B**) The rescue of $\chi = 2$ correlates with an increase of the delayed mean pair-wise correlation among neurons, which is independent of the fraction of neurons sampled and not found in subcritical networks. The temporal coarse graining regime of k for which $\chi = 2$ (*green area*) is not predicted by the pair-wise correlation among neurons.

(Equation 2; **Fig. 2C**, **inset**). Accordingly, we find a rapid breakdown in the agreement between χ and χ_{cn} with

increase in k, with diverging *DCC* (**Fig. 2C**). We conclude that while temporal coarse-graining can mitigate the

bias introduced by thresholding in the fully sampled model, the *DCC* is an unreliable estimate for avalanche

scaling even in fully sampled networks under high thresholding conditions.

Scaling collapse demonstrates equivalence of thresholding and subsampling in the critically balanced

all-to-all E/I-model

Both, the increase in coincident spiking threshold θ as well as the decrease in sampling fraction f underestimate

the scaling exponent χ due to prematurely terminating avalanches by unobserved network activity. On the other

hand, temporal coarse graining allows for concatenating sequential activity periods potentially recovering the

proper duration for a subset of avalanches. We therefore examined the impact of temporal coarse graining on

thresholding and fractional sampling. To achieve this, we first collapsed the χ vs. k curves over a wide range of θ

($10^2 - 10^5$) by scaling θ with an exponent of f, ξ (see Methods). We found that the corresponding functions,

obtained for different values of f, exhibited a minimum total collapse error for $\xi = 1$, demonstrating a linear

relationship between θ and f in our all-to-all model. This implies that a fully sampled network (10^6 neurons) with

$\theta = 3,000$ spikes is qualitatively equivalent to a 0.1% sampled model (1,000 neurons) with $\theta = 3$ spikes (**Fig. 3A**).

This equivalence between θ and f allows us to rescale the recovery of χ for different f in the corresponding (k,

θ)-plane for ease of comparison (see below).

The rescue of $\chi = 2$ correlates with a recovery of mean pair-wise correlations

Subsampling as well as decorrelation of spiking activity have been shown to reduce the expected growth in

mean avalanche size with avalanche duration from $\chi = 2$, to a value of χ closer to $1 - 1.3$ [38-40]. We therefore

studied how the delayed mean pair-wise correlation, *<$C_{ij}(1,k)$>* changes with temporal coarse graining when

recovering χ (see Methods). Temporal coarse graining increases *<$C_{ij}(1,k)$>* in the critical balanced E/I-network

and saturates at high k, independent of the amount of fractional sampling (**Fig. 3B**). The increase in correlation

did not predict the temporal coarse graining value of k at which χ = 2 and was not found in subcritical networks,

for which the average pairwise correlation remains near zero and independent of k (**Fig. 3B**; g = 3.75). These

findings demonstrate that χ = 2 identifies the expansion of coincidental firing among groups of neurons, which is

not captured by a particular value of $<C_{ij}(1,k)>$, a measure limited to studying pairs of neurons.

Subsampling and spike-resolution introduce unobservable regions of scaling exponents

By taking advantage of the trade-off between thresholding and sub-sampling (see Subsection above), we now

can rescale our k vs. θ parameter space by $f \cdot \theta$ to account for different fractional sampling. In the consolidated

analysis (**Fig. 4**), we demonstrate reasonable agreement with the rescue of χ = 2 from 0.1% - 100% fractional

sampling. When we increase the threshold (or equivalently, decrease f), we observe a deviation from χ = 2, yet

the exponent can be reliably restored at higher values of k (**Fig. 4**, white broken lines). On the other hand,

reducing f eventually introduces unobservable (k, θ)-regions which arise from the 1 spike resolution of the

model excluding fractional thresholds. We note that even when sampling only 0.1% of the neurons, one can still

recover χ = 2. However, this approach fails to simultaneously restore all exponents while maintaining the

crackling-noise relationship (see **Fig. 2A** and **Suppl. Fig. S1**) because the point of concurrent rescue lies within

the unobservable regions of parameter space. Decreasing sampling further to f = 0.01% (100 neurons out of 1

million neurons), prevents rescue of χ = 2.

200   Temporal coarse-graining does not rescue the 'crackling noise' relationship of χ_{CN} (Equation 2). The DCC,

201   introduced recently as the difference between the data-based value of χ (Equation 3) and χ_{CN} should measure

202   close to 0 if both estimates agree. As can be seen in **Figure 4**, co-regions with DCC = 0 and χ = 2 exist in the fully-

203   sampled system (**Fig. 4**; asterisk). In contrast, no such co-alignment is found for subsampled systems and

204   temporal coarse graining fails to establish a DCC = 0 regime for most values of k and θ (**Fig. 4**; square, triangles)

205   and accordingly DCC = 0 does not reflect a proper rescue of the true χ = 2 under these conditions.

Figure 4. Temporal coarse -graining identifies a robust rescue regime for χ = 2 covering 0.1% - 100% fractional sampling at decreasing thresholding level, which is absent for the *DCC*

(**A**) Consolidated view of χ and DCC as a function of θ and k for different values of *f*. For *f* = 100%, $\chi \cong 2$ for low θ and k (*star*) but as we make the data sparser by increasing the threshold (or reducing the sampling), we need a higher coarse graining factor, *k*, to compensate and rescue χ back to 2 (*square*). *White dotted region*: visual guide for χ close to 2 for increasing θ and k. The grey parts of the plots for f = 0.1% and 0.01% respectively are unobservable parameter regions since they would require fractional thresholds, below the 1 spike minimum resolution of the model. At *f* = 0.1% sampling, χ can be rescued to a value of 2, but not for more severe subsampling, f = 0.01%. For triangles see *B*.

(**B**) The *DCC* remains close to 0 only for low *k*, and quickly breaks down at higher *k* values. For the fully sampled model, there exists a region with the correct scaling exponent $\chi \cong 2$ as well as low *DCC* (*star* in *A* and *B*) However, as we move to lower sampling, this region of correspondence becomes more and more difficult to maintain. For f = 1% and 0.1%, regions at low k, and moderate threshold (or equivalently lower sampling) remain a low *DCC*, but underestimate the true scaling exponent, χ (*up* and *down triangles* respectively).

Systematic rescue of the scaling exponent for parabolic avalanches in frontal cortex of awake mice

We demonstrate this performance of χ and failure of the DCC in ongoing cellular 2-photon imaging of awake

mice from the anterior cingulate cortex (for details see also [9]). In Figure 5, we demonstrate the robust rescue

of the scaling exponent χ = 2 in our data set for different spike density threshold levels (**Fig. 5**; for details see

Methods and [9]). We find that the recovery of χ = 2 exhibits a θ vs. k dependency as predicted by our

simulations, whereas the DCC is variable and noisy (**Fig. 5**). Similar results were obtained when using a z-scored

threshold (**Suppl. Fig. 2**).

Figure 5. Parabolic avalanches in ongoing activity of frontal cortex exhibit threshold and temporal coarse-graining rescue of scaling exponent χ = 2 in line with critical model dynamics.
Scaling exponent, χ, and crackling noise deviation, DCC, as a function of the spike density thresholds and temporal coarse grain factor, k (n = 5 mice; 2-photon imaging of ongoing activity in frontal cortex (anterior cingulate and medial prefrontal cortex). At low k, an underestimation of χ correspondents to noisy, low values of DCC. At higher k, and threshold θ, rescue of χ to 2 corresponds to high values of the DCC. Note the relationship between k and θ at which χ≅2 is similar to that identified in the critically balanced, E-I model (*broken line*).

DISCUSSION

The exploration of criticality in neuronal systems, particularly focusing on neuronal avalanches, has been subject

to extensive investigation and debate (for review see [18, 51, 57-60]). This study addresses the intricacies and

challenges associated with estimating scaling relationships in the brain, especially in the presence of fractional

subsampling. Our key finding of significance revolves around the retrieval of scaling exponents under fractional

subsampling conditions. Our findings illustrate that temporal coarse-graining facilitates the precise estimation of

the critical scaling exponent ($\chi \approx 2$), even when observing a minimal fraction (0.1%) of the network's neurons.

This discovery carries substantial implications, indicating the potential to accurately estimate critical scaling

characteristics despite severe subsampling constraints. Previous work on similar directed percolation (DP)

models demonstrated a degradation of the scaling exponent to 1.3 with spatial subsampling of 10% and below

[38]. Subsampling has also been shown to mimic apparent population independence of correlations, affecting

criticality markers such as specific heat [61]. We note that in our model, spatial correlations' dependencies

cannot be explored due to the all-to-all connectivity.

Temporal coarse-graining when avalanche periods are unknown

Temporal coarse-graining has been proposed previously as a potential solution to rescue the size exponents of

avalanches in subsampled systems [40] assuming an infinite separation of time scales. This assumption would

provide the experimentalist with the information to separate sequentially occurring avalanches, which is

unrealistic in the study of natural systems. In contrast, our present analysis demonstrates that temporal coarse-

graining can rescue the scaling exponent in models without a clear separation of time scales for fractional

sampling as low as 0.1%, suggesting its potential value in studying highly subsampled experimental data falling

within the DP class. Our study introduces threshold above the Poisson noise floor to approximate separation of

successive avalanches and uncovers a trade-off between subsampling and thresholding. Elevating the threshold

in a fully sampled network is akin to subsampling the network and adopting a lower threshold. This highlights

the interplay between these two factors and their cumulative influence on criticality assessments. By exploiting

this trade-off, we apply our methodology to cellular-resolution 2-photon imaging data obtained from the

anterior cingulate cortex of awake of mice. We underscore the consistent recovery of the scaling exponent to its

critical value of 2. Recently, avalanche detection methods have been introduced that replace temporal

discretization with a continuous time model, though it is currently not clear how these model compare with

experimental data and subsampling conditions [62].

We note that temporal coarse graining differs from a renormalization procedure, which for neuronal avalanches

has been shown to preserve estimates of maximal mutual information as one coarse-samples the spatial scale

under fairly complex macroscopic sampling conditions of the LFP [63]. The temporal coarse graining applied

here rather addresses a stochastic subsampling bias as detailed previously [9].

Limits in rescuing the scaling exponent under subsampling conditions

Our simulations also clearly show that subsampling in discrete models renders certain parts of the phase space

inaccessible (*c.f.* **Fig. 4**), preventing the recovery of the scaling exponent under highly subsampled conditions (f

$\ll 0.1\%$). A similar limitation has been demonstrated for subsampling in the temporal domain, particularly when

the temporal precision required exceeds the temporal resolution of the model [38]. These spatial and temporal

subsampling constraints are particularly relevant to experimental data, which often faces limitations in both

temporal and spatial resolution. Similar constraints have been found for heavy subsampling of network

topologies, particularly when estimating heavy-tail or scale-free distributions of graph properties [64-66].

Insights into under-sampled critical branching processes have been used to understand avalanche statistics

obtained from extracellular spike recordings using microelectrode arrays in the anesthetized rat [67]. Despite

these limits imposed by extreme subsampling, our finding holds promise for experimental neuroscientists

contending with the practical constraint of recording from only a subset of neurons within neural circuits. It

suggests that, with careful consideration of experimental parameters, criticality assessments can remain feasible

and informative, offering valuable insights into the fundamental mechanisms governing neural synchronization.

The DCC and crackling noise relationship fails to recover the scaling exponent under subsampling

conditions

The crackling noise relationship [49-51] has been increasingly proposed and used to quantify the distance of the

observed system from its critical regime (e.g. labeled as the Deviation from Criticality Coefficient (DCC) (e. g. [39,

42]) . However, as outlined by [36] with respect to the general claim by [39], random processes can exhibit a

DCC close to 0 thus limiting general claims of this approach to identify critical dynamics even in fully sampled

systems. In the present study, we clearly show that the DCC approach leads to biased estimates under

subsampling conditions. We also emphasize the high sensitivity of the DCC metric to temporal coarse-graining,

which appears noisy in both our subsampled model and experimental data. Additionally, the DCC metric exhibits

a singularity as the denominator of χ_{cn} passes through 1, complicating its use as an accurate marker of critical

systems. A major confounding factor when using the DCC approach is the assumption that slope estimates from

size and duration distributions equate with critical exponents. This is obviously not the case, as various analysis

parameters such as the temporal resolution Δt and spatial grid sampling distance Δd relate to the obtained

power law slopes in a predictive manner [7, 19], which has to be taken into account when estimating critical

exponents from experimental measures.

Our analysis further highlights the scaling exponent's sensitivity to synchronization thresholds. As the

threshold increases, we observe a noticeable deviation of the scaling exponent and the crackling noise ratio (χ_{cn})

from the expected value of 2 (see also [13]). This sensitivity underscores the importance of careful consideration

of thresholding strategies in experimental setups. Thresholding of time-series can lead to a change in the

avalanche size exponent itself as demonstrated in simulations of critical dynamics [13]. In general, a higher

threshold can slightly increase α, however, the effect on estimating χ appears to be small. In fact, Capek et al. [9]

show in their experimental analysis that including or excluding the subthreshold regime for size estimates

provides narrow upper and lower boundary estimates of χ that center around 2 (see their supplemental figure 6

in [9]). We suggest that the selection of thresholding parameters should ideally be guided by a judicious

equilibrium that ensures the preservation of genuine synchronization events while minimizing the introduction

of noise, especially when the accurate estimation of scaling exponents is the objective.

Synchronization, coincident firing and delayed neuronal correlations

Spatiotemporal synchronization in neural systems traditionally focused on phase-locked oscillation. However,

our system stands out due to the absence of a dominant frequency, presenting a unique scenario. This

peculiarity gains significance when considering 'noise correlations,' which encapsulate the intrinsic non-stimulus-

induced correlation structure among neurons. These correlations exert a notable influence on the decoding

capabilities of neuronal populations, constituting a pivotal aspect of neural information processing [68, 69].

In this study, we scrutinize scale-free neuronal avalanches as the context for exploring neuronal

synchrony and criticality. Neuronal avalanches furnish a comprehensive framework for comprehending neuronal

synchronization, spanning from local clusters of neurons to the global scale of the entire brain. As

demonstrated, the quantitative analysis of neuronal avalanches crucially hinges on the choice of thresholds and

temporal coarse-graining. This dependence is similarly observed in the analysis of spatial and temporal neuronal

correlations, which are substantially influenced by spatial and temporal coarse-graining, respectively [70, 71].

Nevertheless, as depicted in Figure 3, the restoration of the scaling exponent, χ, does not straightforwardly

relate to an increase in pairwise correlations. This underscores the restoration of χ to its critical value of 2 as an

indicator for critical systems. Our findings bear implications for neuroscience and complex systems. Neuronal

avalanches serve as valuable indicators of neural synchronization within the cortex. The precise estimation of

scaling exponents, even under subsampling conditions, is imperative for elucidating the mechanisms governing

neural synchronization in both physiological and pathological states. It bolsters the proposition that the brain

operates near critical points, aligning with the broader concept of self-organized criticality observed in various

complex systems.

Neuronal cell assemblies in the form of parabolic avalanches

In conclusion, our research aims to contribute significantly to the ongoing discourse surrounding

criticality in neural systems. We address the challenges and potential solutions associated with estimating

scaling relationships in neuronal avalanches, especially when faced with subsampling and thresholding

constraints. We present our results as a framework for future investigations into the nature of criticality within

neural networks, paving the way for deeper insights into the fundamental principles governing both normal

brain function and dysfunction.

METHODS

Model Topology

In this work, we employed the model originally proposed by Girardi-Schappo et al. [46] and subsequently

modified in Capek & Ribeiro et al. [9]. This model represents a critically balanced system of integrate-and-fire

(IF) neurons, where the excitatory and inhibitory neurons maintain an equilibrium. Our neuronal network

consisted of N = 10^6 stochastic, non-leaky IF neurons, with a fully connected architecture. To reflect the

prevalence of excitatory neurons in the cortex, we set the ratio of excitatory to inhibitory neurons to 4:1. The

connectivity matrix, denoted as W, was initialized in a manner such that all outgoing synapses from excitatory

neurons had equal strength, with $W_{EE} = W_{IE} = J$. Similarly, the inhibitory neurons had equal strength for their

outgoing synapses, with $W_{II} = W_{EI} = -gJ$. Here, the balance parameter, g, plays a crucial role in tuning the model

towards an E/I-balanced state exhibiting critical dynamics. For our network, the critical value of g, g_c, was

determined to be 3.5. When g exceeds g_c, the network shifts towards an inhibition-dominated regime with

subcritical dynamics, whereas values of g below g_c cause the network to exhibit an excitation-dominated regime

with supercritical dynamics.

Model Dynamics

Each neuron in the model was described by two variables. The Boolean variable, X, indicated whether a neuron

was firing or in a quiescent state at time t. Specifically, X(t) = 1 represented firing, while X(t) = 0 denoted

quiescence. The membrane potential, V, influenced the firing probability and evolved according to the following

equation:

341
$$V_i(t + 1) = \left(\mu V_i(t) + \sum_{j=1}^{N} W_{ij} X_j(t) \right)(1 - X_i(t))$$

342   In the equation, μ represented the leakage parameter, which we set to 0 (non-leaky) for our simulations. The

343   term $(1 - X_i(t))$ introduced an absolute refractory period of Δt, allowing the voltage to reset after a spike

344   occurred (when $X_i(t) = 1$). Additionally, the probability of a neuron firing linearly increased with V and was given

345   by:

346
$$P(X = 1 \mid V) = \begin{cases} V, \Gamma V < 1 \\ 1, \Gamma V \geq 1 \end{cases}$$

Here, Γ represented the neuronal gain, which we fixed at 1 to support probabilistic behavior. Notably, this

network exhibited significant non-conservativeness, with energy dissipation occurring through inhibition and

spike collision.

Additionally, we note that this model includes an absorbing quiescent state, and therefore, the network activity

is initiated by very low-rate independent Poisson processes applied to each neuron. Specifically, the Poisson

process triggers approximately ~20 neurons (0.002%) on average per time step, which is negligible compared to

the system size of 10^6 neurons.

Avalanche Statistics

To comprehensively investigate the resulting population activity, we run 100 simulations of the network over an

extended duration of 10^6 time steps for a total of 10^8 time steps. For the case of full sampling, we compute the

sum of activity across the entire population of $N = 10^6$ neurons. However, to explore various sampling scenarios,

we consider a random subset of N*f neurons, where f represents the sampling fraction. Following this, a

thresholding procedure is applied to the time-series data, setting all neuronal activity below the threshold value

θ to 0. We have the option to analyze the data within the thresholded time-series in their original form or

subject them to temporal coarse graining. When employing coarse graining, each generation represents the sum

of activity over k consecutive time bins, with k representing the coarse graining factor. This approach enables us

to investigate network dynamics across different temporal scales and analyze resulting patterns. Consequently,

we collect the sizes and durations of contiguous epochs from the time-series and examine their distributions.

Power-law fitting

In the context of an ideal, infinitely sized critical system, the distribution of size, duration, and average size as a

function of duration adheres to perfect power laws. However, it is important to consider statistical anomalies,

finite size effects, and the influence of the low Poisson drive in our simulations when analyzing these

distributions. Therefore, a cautious approach is required for their analysis within the model. In the average size

versus duration analyses, we observed that the distributions could be reasonably approximated by a double

power law. For this purpose, we employ the fitting function introduced by Capek & Ribeiro et al. [9]:

$$S(d) = \frac{C d^{\sigma_1}}{(1 + (d/\Phi)^\gamma)^{(\sigma_1 + \sigma_2)/\gamma}}$$

This function exhibits an initial slope σ_1 that transitions to a second slope σ_2 around the point d = Φ. The

parameter γ governs the abruptness of this transition and has been set to 4 for all the analysis presented. The

remaining parameters are allowed to adjust to the data, and all fits were performed in log-space. In all plots

presented in this paper, σ_1 is employed as the representative slope, while σ_2, which is typically associated with

finite-size effects, is not discussed in this context.

We use the cutoff, Φ that we obtain from the scaling curves to analyze the individual size and duration

distributions. The size exponent, β, is the slope of the size distribution in log-scale with a lower cutoff which is

safely set to be greater than the Poisson noise level (>100) and an upper cutoff equal to the average size

corresponding to the elbow point, Φ. For the duration exponent, β, we do a similar analysis, with a lower

duration cutoff of 3 time-bins and an upper cutoff equal to Φ.

Correlation analysis

In our examination of the mean pairwise correlation within the system, we focus on a smaller temporal segment

of the simulated network (consisting of 10,000 time steps) to facilitate analysis of large neuron raster datasets.

For each neuron pair (designated as an ordered pair [i,j]) within the network, we compute the time-lagged

correlation between the activity of neuron i and that of neuron j, shifted by one time step, k*Δt. The mean

correlation across the system is then determined as the average correlation coefficient computed for all such

neuron pairs. This correlation analysis spans temporal bins ranging from 0 to 100 and spatial sampling values

from 0.01% up to 1%. However, due to the quadratic increase in duration and memory requirements with

system size, higher sampling rates become impractical for storing raster data and conducting pairwise analyses.

Ongoing activity in contralateral ACC/mPFC monitored with jRGECO1a.

Mice (C57BL/6; Jackson Laboratories; age >6 weeks) were injected with a viral construct to express jRGECO1a in

cortical neurons using Syn promotor. Chronic 2PI started after ~2 weeks in the contralateral ACC/mPFC at an

estimated depth of ~150–300 µm using a microprism. Recordings were collected over the course of several days

from n = 5 mice (3 males, 2 females; age 8–20 weeks) with jRGECO1a expression (n = 17 recordings; 30min

each). Recordings were conducted over the course of several weeks and analyzed separately for each mouse.

2PI images were acquired by a scanning microscope (Bergamo II series, B248, Thorlabs Inc.) coupled to a pulsed

femtosecond Ti:Sapphire 2-photon laser (Chameleon Discovery NX, Coherent Inc.). The wavelength was tuned to

1120 nm to excite jRGECO1a. The field of view was ~450 um x 450 μm. Imaging frames of 512 x 512 pixels were

acquired at 45.527Hz by bidirectional scanning of a 12 kHz Galvo-resonant scanner. The average power for

imaging was <70mW, measured at the sample. Obtained tif-movies in uint16 format were rigid motion corrected

via the python-based software package suite2p [72]. Registered images were denoised using machine-learning-

based, DeepInterpolationLecoq [73]and then semi-automatically processed by suite2p for ROI selection and

fluorescence signal extraction. For each labeled neuron, raw soma and neuropil fluorescence signals over time

were extracted for each ROI. Spiking probabilities were obtained from neuropil corrected fluorescence traces

(F_corrected = F_ROI – 0.7*F_neuropil) via MLspike (https://github.com/MLspike) by utilizing its autocalibration

feature to obtain unitary spike event amplitude, decay time, and channel noise for individual ROIs.

ACKNOWLEDGEMENTS

This research was supported by the Division of the Intramural Research Program (DIRP) of the National Institute

of Mental Health (NIMH), USA, ZIAMH002797, ZIAMH002971, and the BRAIN initiative Grant U19 NS107464-01

to D.P. This research utilized the supercomputing resources of the National Institutes of Health (NIH, USA;

Biowulf, http://hpc.nih.gov) and the University of Maryland (UMD College Park, USA,

https://hpcc.umd.edu/hpcc/dt2.html).

AUTHOR CONTRIBUTIONS

K.S., T.L.R., D. P. conceived the study, K.S. did the simulations, P.K. provided the experimental data and

processing, K.S., T.L.R., D. P. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

SUPPLEMENTARY INFORMATION

Suppl. Fig. S1. Consolidated view of the exponents α, β and χ as a function of threshold and temporal coarse-graining for different sampling fractions.

(**A**) Consolidated view of α, β and χ as a function of θ and k for different values of f. For f = 100%, $\chi \cong 2$ for low θ and k (*star*) but as we make the data sparser by increasing the threshold (or reducing the sampling), we need a higher coarse graining factor, k, to compensate and rescue χ back to 2 (*square*). Bottom row is replotted for χ from Figure 4 for ease of comparison with the corresponding size slopes, α, and duration slopes, β. *White dotted region*: visual guide for χ close to 2 for increasing θ and k. The grey parts of the plots for f = 0.1% and 0.01% respectively are unobservable parameter regions since they would require fractional thresholds, below the 1 spike minimum resolution of the model (*cf.* Figure 4).

Suppl. Fig. S2. Parabolic avalanches in ongoing activity of frontal cortex exhibit threshold and temporal coarse-graining rescue of scaling exponent χ = 2 in line with critical model dynamics: z-scored threshold.
(**A**) Number of avalanches as a function of threshold for 17 different recordings of ongoing neuronal activity in anterior cingulate frontal cortex. Inset highlights the z-scored threshold which is calculated independently for each recording. Data pooled from 5 mice (3 males, 2 females) with jRGECO1a expression for a total of 17 recordings of 30 mins each (2-photon imaging).
(**B, C**) Scaling exponent and DCC as a function of the z-scored thresholds instead of absolute spike density threshold (*cf*. Figure 5) and temporal coarse grain factor, *k*.

REFERENCES

1.   Hamon D, Nicodemi M, Jensen H. Continuously driven OFC: A simple model of solar flare
statistics. Astronomy & Astrophysics. 2002;387(1):326-34.
2.   Olami Z, Feder HJS, Christensen K. Self-organized criticality in a continuous, nonconservative
cellular automaton modeling earthquakes. PhysRevLett. 1992;68:1244-7.
3.   Bak P, Tang C. Earthquakes as a self-organized critical phenomenon. Journal of Geophysical
Research: Solid Earth. 1989;94(B11):15635-7.
4.   Bassler KE, Paczuski M. Cellular Model of Superconducting Vortex Dynamics. Complexity
from Microscopic to Macroscopic Scales: Coherence and Large Deviations: Springer; 2002. p. 215-27.
5.   Held GA, Solina D, Solina H, Keane D, Haag W, Horn P, Grinstein G. Experimental study of
critical-mass fluctuations in an evolving sandpile. Phys Rev Lett. 1990;65(9):1120.
6.   Drossel B, Schwabl F. Self-organized critical forest-fire model. PhysRevLett. 1992;69(11):1629-
32.
7.   Beggs JM, Plenz D. Neuronal avalanches in neocortical circuits. J Neurosci. 2003;23(35):11167-
77. doi: 10.1523/JNEUROSCI.23-35-11167.2003.
8.   Bellay T, Klaus A, Seshadri S, Plenz D. Irregular spiking of pyramidal neurons organizes as
scale-invariant neuronal avalanches in the awake state. eLife. 2015;4: e07224. doi: 10.7554/eLife.07224.
9.   Capek E, Ribeiro TL, Kells P, Srinivasan K, Miller SR, Geist E, et al. Parabolic avalanche
scaling in the synchronization of cortical cell assemblies. Nature communications. 2023;14(1):2555. doi:
10.1101/2022.11.02.514938.
10.   Scarpetta S, Apicella I, Minati L, de Candia A. Hysteresis, neural avalanches, and critical
behavior near a first-order transition of a spiking neural network. Phys Rev E. 2018;97(6):062305. doi:
10.1103/PhysRevE.97.062305.
11.   Scarpetta S. Critical Behavior and Memory Function in a Model of Spiking Neurons with a
Reservoir of Spatio-Temporal Patterns. In: Tomen N, Herrmann JM, Ernst U, editors. The Functional
Role of Critical Dynamics in Neural Systems. Cham: Springer International Publishing; 2019. p. 179-97.
12.   Liang J, Zhou C. Criticality enhances the multilevel reliability of stimulus responses in cortical
neural networks. PLoS Comput Biol. 2022;18(1):e1009848. doi: 10.1371/journal.pcbi.1009848.
13.   Villegas P, Di Santo S, Burioni R, Muñoz MA. Time-series thresholding and the definition of
avalanche size. Phys Rev E. 2019;100(1). doi: 10.1103/PhysRevE.100.012133.
14.   Morowitz HJ. The mind, the brain and complex adaptive systems: Routledge; 1994.
15.   Holland JH. Signals and boundaries: Building blocks for complex adaptive systems: Mit Press;
2012.
16.   Lombardi F, Herrmann H, Plenz D, de Arcangelis L. On the temporal organization of neuronal
avalanches. Front Syst Neurosci. 2014;8:204. doi: 10.3389/fnsys.2014.00204.
17.   Lombardi F, Herrmann HJ, Plenz D, de Arcangelis L. Temporal correlations in neuronal
avalanche occurrence. Scientific Reports. 2016;6:24690. doi: 10.1038/srep24690.
18.   Plenz D, Ribeiro TL, Miller SR, Kells PA, Vakili A, Capek EL. Self-Organized Criticality in the
Brain. FrPhy. 2021;9(365). doi: 10.3389/fphy.2021.639389.
19.   Petermann T, Thiagarajan T, Lebedev MA, Nicolelis MA, Chialvo DR, Plenz D. Spontaneous
cortical activity in awake monkeys composed of neuronal avalanches. Proc Natl Acad Sci USA.
2009;106(37):15921-6. doi: 10.1073/pnas.0904089106.
20.   Miller SR, Yu S, Plenz D. The scale-invariant, temporal profile of neuronal avalanches in
relation to cortical γ–oscillations. Scientific Reports. 2019;9:16403. doi: 10.1101/757278.

21.   Yu S, Yang H, Nakahara H, Santos GS, Nikolic D, Plenz D. Higher-order interactions
characterized in cortical activity. J Neurosci. 2011;31(48):17514-26. doi: 10.1523/JNEUROSCI.3127-
11.2011.
22.   Bellay T, Shew WL, Yu S, Falco-Walter JJ, Plenz D. Selective participation of single cortical
neurons in neuronal avalanches. Frontiers in Neural Circuits. 2021;14(90). doi:
10.3389/fncir.2020.620052.
23.   Shriki O, Plenz D. Neuronal avalanches in the human brain. In: Plenz D, Niebur E, editors.
Criticality in Neural Systems. Reviews of Nonlinear Dynamics and Complexity. 7. 1 ed. Berlin: Wiley-
VCH; 2014. p. 177 - 89.
24.   Palva JM, Zhigalov A, Hirvonen J, Korhonen O, Linkenkaer-Hansen K, Palva S. Neuronal long-
range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws. Proc
Natl Acad Sci U S A. 2013;110(9):3585-90. Epub 2013/02/13. PubMed PMID: 23401536; PubMed
Central PMCID: PMC3587255.
25.   Gireesh ED, Plenz D. Neuronal avalanches organize as nested theta- and beta/gamma-
oscillations during development of cortical layer 2/3. Proc Natl Acad Sci USA. 2008;105:7576-81.
26.   Meisel C. Antiepileptic drugs induce subcritical dynamics in human cortical networks. Proc Natl
Acad Sci U S A. 2020;117(20):11118. doi: 10.1073/pnas.1911461117.
27.   Maturana MI, Meisel C, Dell K, Karoly PJ, D'Souza W, Grayden DB, et al. Critical slowing
down as a biomarker for seizure susceptibility. Nature communications. 2020;11(1):2172. doi:
10.1038/s41467-020-15908-3.
28.   Meisel C, Plenz D, Schulze-Bonhage A, Reichmann H. Quantifying antiepileptic drug effects
using intrinsic excitability measures. Epilepsia. 2016;57(11):e210-e5. doi: 10.1111/epi.13517.
29.   Meisel C, Schulze-Bonhage A, Freestone D, Cook MJ, Achermann P, Plenz D. Intrinsic
excitability measures track antiepileptic drug action and uncover increasing/decreasing excitability over
the wake/sleep cycle. Proc Natl Acad Sci U S A. 2015;112(47):14694-9. doi: 10.1073/pnas.1513716112.
30.   Ribeiro S, Nicolelis MA. Reverberation, storage, and postsynaptic propagation of memories
during sleep. LearnMem. 2004;11(6):686-96.
31.   Ribeiro S, Gervasoni D, Soares ES, Zhou Y, Lin SC, Pantoja J, et al. Long-lasting novelty-
induced neuronal reverberation during slow-wave sleep in multiple forebrain areas. PLoSBiol.
2004;2(1):E24.
32.   Meisel C, Klaus A, Vyazovskiy VV, Plenz D. The interplay between long- and short-range
temporal correlations shapes cortex dynamics across vigilance states. J Neurosci. 2017;37(42):10114.
doi: 10.1523/JNEUROSCI.0448-17.2017.
33.   Meisel C, Bailey K, Achermann P, Plenz D. Decline of long-range temporal correlations in the
human brain during sustained wakefulness. Scientific Reports. 2017;7(1):11825. doi: 10.1038/s41598-
017-12140-w.
34.   Meisel C, Olbrich E, Shriki O, Achermann P. Fading signatures of critical brain dynamics during
sustained wakefulness in humans. J Neurosci. 2013;33(44):17363-72. Epub 2013/11/01. doi:

10.1523/JNEUROSCI.1516-13.2013. PubMed PMID: 24174669; PubMed Central PMCID:
PMC3858643.
35.   Meisel C, Storch A, Hallmeyer-Elgner S, Bullmore E, Gross T. Failure of adaptive self-
organized criticality during epileptic seizure attacks. PLoS Comput Biol. 2012;8(1):e1002312.
36.   Destexhe A, Touboul JD. Is there sufficient evidence for criticality in cortical systems? eneuro.
2021:ENEURO.0551-20.2021. doi: 10.1523/ENEURO.0551-20.2021.
37.   Touboul J, Destexhe A. Power-law statistics and universal scaling in the absence of criticality.
Phys Rev E. 2017;95(012413). doi: 10.1103/PhysRevE.95.012413.
38.   Carvalho TTA, Fontenele AJ, Girardi-Schappo M, Feliciano T, Aguiar LAA, Silva TPL, et al.
Subsampled directed-percolation models explain scaling relations experimentally observed in the brain.
Frontiers in Neural Circuits. 2021;14(83). doi: 10.3389/fncir.2020.576727.
39.   Fontenele AJ, de Vasconcelos NAP, Feliciano T, Aguiar LAA, Soares-Cunha C, Coimbra B, et
al. Criticality between cortical states. Phys Rev Lett. 2019;122(20):208101. doi:
10.1103/PhysRevLett.122.208101.
40.   Levina A, Priesemann V. Subsampling scaling. Nature communications. 2017;8:15140. doi:
10.1038/ncomms15140.
41.   Xu Y, Schneider A, Wessel R, Hengen KB. Sleep restores an optimal computational regime in
cortical networks. bioRxiv. 2022:2022.12.07.519478. doi: 10.1101/2022.12.07.519478.
42.   Ma Z, Liu H, Komiyama T, Wessel R. Stability of motor cortex network states during learning-
associated neural reorganizations. J Neurophysiol. 2020;124(5):1327-42. doi: 10.1152/jn.00061.2020.
PubMed PMID: 32937084.
43.   Karimipanah Y, Ma Z, Miller J-eK, Yuste R, Wessel R. Neocortical activity is stimulus- and
scale-invariant. PloS one. 2017;12(5):e0177396. doi: 10.1371/journal.pone.0177396.
44.   Friedman N, Ito S, Brinkman BAW, Shimono M, DeVille REL, Dahmen KA, et al. Universal
critical dynamics in high resolution neuronal avalanche data. Phys Rev Lett. 2012;108(20):208102.
45.   Levina A, Priesemann V, Zierenberg J. Tackling the subsampling problem to infer collective
properties from limited data. Nature Reviews Physics. 2022;4(12):770-84. doi: 10.1038/s42254-022-
00532-5.
46.   Girardi-Schappo M, Brochini L, Costa AA, Carvalho TTA, Kinouchi O. Synaptic balance due to
homeostatically self-organized quasicritical dynamics. Physical Review Research. 2020;2(1):012042.
doi: 10.1103/PhysRevResearch.2.012042.
47.   Girardi-Schappo M, Galera EF, Carvalho TTA, Brochini L, Kamiji NL, Roque AC, Kinouchi O.
A unified theory of E/I synaptic balance, quasicritical neuronal avalanches and asynchronous irregular
spiking. Journal of Physics: Complexity. 2021;2(4):045001. doi: 10.1088/2632-072X/ac2792.
48.   Scarpetta S, Morisi N, Mutti C, Azzi N, Trippi I, Ciliento R, et al. Criticality of neuronal
avalanches in human sleep and their relationship with sleep macro- and micro-architecture. iScience.
2023;26(10):107840. doi: 10.1016/j.isci.2023.107840.
49.   Mehta AP, Mills AC, Dahmen KA, Sethna JP. Universal pulse shape scaling function and
exponents: Critical test for avalanche models applied to Barkhausen noise. Phys Rev E. 2002;65. doi:
10.1103/PhysRevE.65.046139. PubMed PMID: 12005958.
50.   Kuntz MC, Sethna JP. Noise in disordered systems: The power spectrum and dynamic exponents
in avalanche models. PhRvB. 2000;62(17):11699-708.
51.   Jensen HJ. Self-organized criticality: Cambridge University Press; 1998.
52.   Chen Y-J, Papanikolaou S, Sethna JP, Zapperi S, Durin G. Avalanche spatial structure and
multivariable scaling functions: Sizes, heights, widths, and views through windows. Phys Rev E.
2011;84(6):061103.

53.   Magni A, Durin G, Zapperi S, Sethna JP. Visualization of avalanches in magnetic thin films:
temporal processing. JSMTE. 2009;2009(01):P01020.
54.   Ma Z, Turrigiano GG, Wessel R, Hengen KB. Cortical circuit dynamics are homeostatically
tuned to criticality in vivo. Neuron. 2019;104:655-64. doi: 10.1016/j.neuron.2019.08.031.
55.   Muñoz MA, Dickman R, Vespignani A, Zapperi S. Avalanche and spreading exponents in
systems with absorbing states. Phys Rev E 1999;59(5 Pt B):6175-9.
56.   Priesemann V, Munk MH, Wibral M. Subsampling effects in neuronal avalanche distributions
recorded in vivo. BMCNeurosci. 2009;10(1):40.
57.   Muñoz MA. Colloquium: Criticality and dynamical scaling in living systems. Rev Mod Phys
2018;90(3):031001. doi: 10.1103/RevModPhys.90.031001.
58.   Cocchi L, Gollo LL, Zalesky A, Breakspear M. Criticality in the brain: A synthesis of
neurobiology, models and cognition. Prog Neurobiol. 2017. doi: 10.1016/j.pneurobio.2017.07.002.
59.   Hesse J, Gross T. Self-organized criticality as a fundamental property of neural systems. Front
Syst Neurosci. 2014;8. doi: 10.3389/fnsys.2014.00166.
60.   Chialvo DR. Emergent complex neural dynamics. Nat Phys. 2010;6:744-50. doi: DOI:
10.1038/NPHYS1803.
61.   Nonnenmacher M, Behrens C, Berens P, Bethge M, Macke JH. Signatures of criticality arise
from random subsampling in simple population models. PLoS Comput Biol. 2017;13(10):e1005718.
62.   Pausch J. From neuronal spikes to avalanches: Effects and circumvention of time binning.
Physical Review Research. 2022;4(2):023212. doi: 10.1103/PhysRevResearch.4.023212.
63.   Shew WL, Yang H, Yu S, Roy R, Plenz D. Information capacity is maximized in balanced
cortical networks with neuronal avalanches. J Neurosci. 2011;5:55-63. doi: 10.1523/JNEUROSCI.4637-
10.2011. PubMed Central PMCID: PMCPMC3082868.
64.   Stumpf MP, Wiuf C, May RM. Subnets of scale-free networks are not scale-free: sampling
properties of networks. ProcNatlAcadSciUSA. 2005;102(12):4221-4.
65.   Serafino M, Cimini G, Maritan A, Rinaldo A, Suweis S, Banavar JR, Caldarelli G. True scale-
free networks hidden by finite size effects. Proc Natl Acad Sci U S A. 2021;118(2):e2013825118. doi:
10.1073/pnas.2013825118.
66.   Bonzanni M, Bockley KM, Kaplan DL. On the effect of neuronal spatial subsampling in small-
world networks. Eur J Neurosci. 2021;53(2):485-98. doi: 10.1111/ejn.14937.
67.   Ribeiro TL, Ribeiro S, Belchior H, Caixeta F, Copelli M. Undersampled critical branching
processes on small-world and random networks fail to reproduce the statistics of spike avalanches. PloS
one. 2014;9(4):e94992. doi: 10.1371/journal.pone.0094992.
68.   Gawne TJ, Kjaer TW, Hertz JA, Richmond BJ. Adjacent visual cortical complex cells share
about 20% of their stimulus-related information. Cereb Cortex. 1996;6(3):482-9.
69.   Averbeck BB, Latham PE, Pouget A. Neural correlations, population coding and computation.
Nat Rev Neurosci. 2006;7(5):358-66.
70.   Smith MA, Kohn A. Spatial and Temporal Scales of Neuronal Correlation in Primary Visual
Cortex. J Neurosci. 2008;28(48):12591. doi: 10.1523/JNEUROSCI.2929-08.2008.
71.   Schulz DPA, Sahani M, Carandini M. Five key factors determining pairwise correlations in
visual cortex. J Neurophysiol. 2015;114(2):1022-33. doi: 10.1152/jn.00094.2015. PubMed PMID:
26019310.
72.   Pachitariu M, Stringer C, Dipoppa M, Schröder S, Rossi LF, Dalgleish H, et al. Suite2p: beyond
10,000 neurons with standard two-photon microscopy. bioRxiv. 2017:061507. doi: 10.1101/061507.

73.   Lecoq J, Oliver M, Siegle JH, Orlova N, Ledochowitsch P, Koch C. Removing independent
noise in systems neuroscience data using DeepInterpolation. Nat Methods. 2021. doi: 10.1038/s41592-
021-01285-2.