Prioritizing Maize Metabolic Gene Regulators through Multi-Omic Network Integration

Fabio Gomez-Cano[1,6*], Jonas Rodriguez[2], Peng Zhou[3], Yi-Hsuan Chu[1], Erika Magnusson[3], Lina Gomez-Cano[1], Arjun Krishnan[4], Nathan M Springer[3,5], Natalia de Leon[2], Erich Grotewold[1]

[1] Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI 48824-6473, USA.

[2] Department of Plant and Agroecosystem Sciences, University of Wisconsin Madison, Madison, WI 53706, USA

[3] Department of Plant and Microbial Biology, University of Minnesota, Saint Paul, MN 55108

[4] Department of Biomedical Informatics, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA

[5] Current address: Global Breeding, Bayer Crop Sciences, Chesterfield MO 63017, USA

[6] Current address: Department of Molecular, Cellular, and Development Biology, University of Michigan, Ann Arbor, MI, 48109, USA.

* To whom correspondence should be addressed: gomezcan@umich.edu

Author e-mail address:

Fabio Gomez-Cano, ORCID 0000-0002-2624-0112, gomezcan@umich.edu

Jonas Rodriguez, ORCID 0000-0001-8693-1022, jorod550@gmail.com

Peng Zhou, ORCID 0000-0001-5684-2256, zhoux379@umn.edu

Erika Magnusson, ORCID 0000-0001-7868-6215, magnu513@umn.edu

Lina Gomez-Cano, ORCID 0000-0002-9963-2942, gomezca5@msu.edu

Arjun Krishnan, ORCID 0000-0002-7980-4110, arjun.krishnan@cuanschutz.edu

Nathan M Springer, ORCID 0000-0002-7301-4759, springer@umn.edu

Natalia de Leon, ORCID 0000-0001-7867-9058, ndeleongatti@wisc.edu

Erich Grotewold, ORCID 0000-0002-4720-7290, grotewol@msu.edu

GRAPHICAL ABSTRACT

Integration

Evaluation
Perturbation assays

TF-funtion mapping
based on expression

ABSTRACT

Elucidating gene regulatory networks is a major area of study within plant systems biology. Phenotypic traits are intricately linked to specific gene expression profiles. These expression patterns arise primarily from regulatory connections between sets of transcription factors (TFs) and their target genes. Here, we integrated 46 co-expression networks, 283 protein-DNA interaction (PDI) assays, and 16 million SNPs used to identify expression quantitative trait loci (eQTL) to construct TF-target networks. In total, we analyzed ~4.6M interactions to generate four distinct types of TF-target networks: co-expression, PDI, *trans*-eQTL, and *cis*-eQTL combined with PDIs. To functionally annotate TFs based on their target genes, we implemented three different network integration strategies. We evaluated the effectiveness of each strategy through TF loss-of function mutant inspection and random network analyses. The multi-network integration allowed us to identify transcriptional regulators of several biological processes. Using the topological properties of the fully integrated network, we identified potential functionally redundant TF paralogs. Our findings retrieved functions previously documented for numerous TFs and revealed novel functions that are crucial for informing the design of future experiments. The approach here-described lays the foundation for the integration of multi-omic datasets in maize and other plant systems.

Keywords: *Transcription factor, PDI, eQTL, Co-expression, Gene regulation*

INTRODUCTION

Like other organisms, plants use interconnected molecular networks that collaboratively coordinate every cellular process, from cell division to metabolite synthesis and adaptation. Among these, regulatory networks play central roles by controlling the expression of every gene in the cell, function that is carried out mainly by transcription factor (TF) proteins (1). The transcriptional regulation of genes requires direct protein-DNA interactions (PDI) between TFs and specific *cis*-regulatory elements (CREs) located in proximal (promoters) or distal (enhancers/silencers) regulatory regions of the corresponding target gene (2). TFs can also regulate gene expression indirectly by engaging in protein-protein interactions (PPI) and being tethered to DNA through other PDIs. Together, collections of PDIs form highly interconnected gene

regulatory networks (GRNs). The architecture of GRNs is elucidated using combinations of gene- and TF-centered approaches (3–5). One commonly used gene-centered method is provided by yeast one-hybrid (Y1H) assays (3, 5). TF-centered strategies involve techniques like chromatin immunoprecipitation followed by sequencing (ChIP-seq) for *in vivo* PDI discovery and DNA affinity purification sequencing (DAP-Seq) for *in vitro* analyses (5, 6). However, ChIP-seq has limitations in scalability to multiple TFs, while DAP-seq captures interactions that ignore the native chromatin environment (5). The organization of GRNs has implications for phenotypic variation (7), plant responses to abiotic and biotic stress (8–10), development (11), speciation (12), as well as adaptation and diversification (11–13), among others. Hence, understanding the structure and dynamics of GRNs offers a valuable avenue and strategy for manipulating desired phenotypes.

Maize holds great agricultural significance due to its broad adaptation and wide range of applications (14). Part of what provides maize with its versatility is its extraordinary metabolic diversity (15–18), which is underpinned by its genetic diversity (19–21), and varies depending on endogenous variables (e.g., organs or development stages) (18) and environmental factors (16, 22). Similar to other plant and animal systems, quantitative traits in maize are genetically complex, illustrated by the large number of loci associated with a single trait and the minor contribution of any given association to the trait's variance (15–18, 23, 24). Hence, elucidating the molecular mechanisms underlying quantitative traits faces two primary obstacles when only simply considering trait-loci associations: I) the involvement of multiple genes governing a single phenotypic outcome, and II) the impact of additional genetic elements that determine and modulate the genetic contribution to phenotypic variation.

A distinctive genomic characteristic of maize is a recent whole genome duplication (WGD) event which occurred ~5-12 Mya, making maize an ancient allotetraploid (25). Maize has an abundance of tandem duplicated genes (26), and is highly-enriched in transposable elements (TEs; ~85% of the genome) (19). Furthermore, the WGD event resulted in the formation of two subgenomes (*maize1* and *maize2*), exhibiting unequal gene loss and expression patterns, primarily driven by the subgenome with a lower fractionation rate (27). The dominant subgenome, *maize1,* was shown to have a larger contribution to phenotypic

variations (28). Nevertheless, the precise molecular mechanisms underlying the asymmetric contributions of each subgenome remain largely unknown and are likely orchestrated at multiple molecular levels, including regulation, signaling, and interactome level, as evidenced by co-expression and multi-network comparisons (29, 30). Understanding these mechanisms holds significance for modeling, prioritizing, and unraveling the principal factors behind agriculturally relevant traits, while also advancing our fundamental understanding of maize evolution.

The generation of multi-omic datasets has gained recent traction as a means to understand the complexity of genetic and phenotypic variation observed in biological systems (31). Thus, the integration of diverse datasets is an evolving field, which may be grouped into conceptual integration (overlapping observations), statistical integration, network-based integration, and machine learning-based integration (32). In maize, like other model organisms, technological advancements have also favored the rapid generation of multi-omic datasets. These include TF binding profiles (33–35); identification of accessible chromatin regions (ACRs) (34, 36, 37); expression and co-expression atlases (38–41); and population-level transcriptomic, proteomic, and metabolic information (16–18, 24, 42–44). Consequently, there is a growing interest in establishing multi-omic integration (17, 45–51). However, most integration efforts involve the verification of each layer with one another (i.e., conceptual integration) (32). In a few exceptional cases, the layers are leveraged to enhance the integration (50, 52) or to learn from their combined information (30), emphasizing the need for a comprehensive assessment of integration strategies and their effectiveness in prioritizing gene-specific processes.

In this study, we analyzed genetic and gene expression variation across 304 maize inbred lines. We utilized data from over 300 publicly available ChIP- and DAP-seq experiments, along with 45 previously analyzed co-expression networks (41) with the goal to define a robust strategy to integrate the corresponding datasets to ultimately identify elusive regulators of maize metabolism. Combining these datasets, we built four molecular networks and employed three integration methods to evaluate the functional annotation of TFs based on their predicted target genes. Overall, we find significantly better predictions by performing integrations of each molecular network after extracting physical features –

embeddings - of the corresponding networks (aka, network-base integration). Our analysis also shows that the network-based strategy implemented here allows to identify a large number of known transcriptional regulator from a wide range of biological process, from transcriptional control of hormone signaling, developmental-related processes, and several metabolic pathways. Additionally, our analyses show how mapping the TF-biological process guided by condition-specific expression profiles allows the mapping of specification of TF-biological functions into defined conditions, which also highlights TFs with contracting biological regulation profiles on defined biological conditions as a function of the tissue or the genotype. Finally, we combined the embedding with expression and protein sequence variation; we found candidates of TF homologs resulting from functional divergence, either by subfunctionalization or neofunctionalization — an observation with a large impact on the design of future experimental validations.

MATERIAL AND METHODS

Identification of eQTL

A set of 304 diverse inbred lines with publicly available SNP and gene expression information were included in our eQTL analysis (24, 43, 53). Given that the previously published SNP marker data from whole genome sequencing and RNA-seq were mapped to the same B73 maize reference genome (AGPv4) (54), we performed a simple concatenation of the datasets using bcftools (v1.7) (55). In the case that an overlap was observed between the two datasets, the RNA-seq marker was preferentially kept. The expression datasets capture variation both at the genotypic and tissue level.

RNA-seq and co-expression analyses

All the RNA-seq and co-expression datasets utilized here were previously published (41), except for co-expression network 46, which was constructed using the 304 inbred lines analyzed for genetic markers (referred to as n304). Specifically, we gathered pre-mapped count per million (CPM) values for the respective inbred lines and employed the exact strategy outlined by Zhou et al. (2020) to construct the corresponding co-expression networks. All co-expression networks were based on RandomForestRegressor and using the top 100K association by TF.

eQTL identification and classification

eQTLs were identify using eight distinct tissue types encompassing different developmental stages from germination to plant maturity (GRoot, Gshoot, Kern, L3Base, L3Tip, LMAD, LMAN, and seedling) (24, 43, 53). SNPs were filtered by removing non-biallelic markers, and those with a minor allele frequency (<0.05). Each of the tissue-specific expression datasets were filtered independently by retaining genes with ≥ 6 reads in $\geq 20\%$ of samples and ≥ 0.1 TPM in $\geq 20\%$ of samples. After filtering, we tested between 15.5M - 16.7M SNPs against the expression of 15.3K - 26.4K genes across the eight tissue types. Briefly, to test SNP-gene associations, a series of eight candidate linear models were fitted beginning with a t-test then progressively controlling by using different levels of kinship and population structure in a mixed linear model. For each model tested, the association was deemed significant if the observed P-value surpassed the 10K permutation threshold computed for each gene. Non-significant eQTLs were discarded when the association was supported by fewer than two of the candidate linear models or when the association involved non-syntenic genes (56). These associations were classified as *cis*-eQTLt, *trans*/*cis*-eQTL, *cis*-eQTL, *trans*-eQTL, and unassigned eQTL according with the distance between each eQTL and its corresponding target gene as well as its co-location with annotated maize genes (genome B73-V4, Figure S3.2a).

Protein-DNA interactions data analysis

Raw reads from plant ChIP-seq experiments (57–63), ChIP-seq experiments from protoplast (pChIP-seq) (35), and DAP-seq (34, 64) were collected from publicly available datasets (Table S2). Quality control of reads and peak identification were performed as reported previously (65). Briefly, read quality control was performed using FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/, V0.11.5). Adapters and low quality reads were trimmed with Trimmomatic (66) using the following parameters: ILLUMINACLIP:Adapter.fastq:2:40:15 SLIDINGWINDOW:4:20 MINLEN:30. Cleaned reads were mapped to the B73 maize reference genome (AGPv4) (54) with Bowtie2 v2.3.4.1 (67) and only using nuclear chromosomes. Multi-mapping reads were filtered with Samtools v1.9 (68) (q 30). Peaks were called using GEM v3.4 (69). For DAP-seq datasets, sequences retrieved using the HALO protein alone were used as a negative control. Peaks from plant ChIP-seq experiments were called including replicates

and using the corresponding mutants or tag-protein as control. Finally, ChIP-seq from protoplasts was called including replicates. All of them used the following parameters: --d Read_Distribution_default.txt --k_min 6 --k_max 15 --k_seqs 2000 --outNP --sl. Only TFs with >500 predicted peaks were used for further analysis.

Peak quality control. We evaluated the overlap of TF peaks with accessible chromatin regions (ACRs) and scaled the number of CPM per peak and assay. Thus, TF binding peaks with a Z-score larger than -0.5 and mapping to ACRs were kept for further analysis. Specifically, to obtain CPMs per peak, we used the summit of each peak and extended them 50 bps to each side of the summit. Then, we converted extended summit files into SAF files and count mapped reads using Rsubread v1.32.2 (70), followed by CPM normalization. Publicly available ACRs were collected from (37).

Functional annotation

All functional annotations were performed after keeping only syntenic genes with sorghum (56). Metabolic pathways (PWYs) were collected from CornCYC (71), and GO terms were obtained from GAMER (72). Syntenic genes were defined based on Schnable et al. (2019). Enrichment analysis for PWYs and GO terms was conducted in R using the GeneOverlap (v1.30.0) and topGO (v2.46.0) packages, respectively. GO term semantic similarity was calculated using the GOSemSim (v2.20.0) package in R, employing the "Wang" option (73). For *common function* analyses, all GO comparisons were performed on the original set of enriched GO terms, and after the comparisons (GO semantic similarity), all GO terms were mapped to their closest parent GO term using the R package Rrvgo (v1.6) (74). Similarly, all GO terms significantly enriched from *common target* and *network-based* analyses were mapped to its closed parent to reduce their redundancy.

Network integration

Common interactions. We identified common interactions by counting how frequently TF-target associations were predicted across all layers (i.e., GAN, GRN, eGRN, and CEN networks). We considered interactions as common when they were present in at least two different layers for the same corresponding TF. Thus, with the defined list of target genes, we tested for enrichment with PWYs and GO terms within

the corresponding targets per each TF. All significant GO term associations were mapped to their closest parent terms using the R package Rrvgo to reduce redundancy (v1.6) (74).

Common function. Common functions were identified by testing the enrichment of target genes with PWYs and GO terms for each TF in each layer as follows. TFs that had at least one PWY/GO term enriched in at least two different layers were retained to assess common predictions across layers (Figure S3.5a). The similarity in PWY predictions among layers was performed by comparing all PWYs between layers for each TF using a Fisher exact test. We used the enrichment test because a single gene could be annotated in multiple PWYs. The similarity in GO term predictions were performed by measuring the semantic similarity between the corresponding terms. Significant GO terms are then mapped to their closest parent terms.

Network-based. All four layers were combined, and the interaction frequencies were scaled from 0.5 to 1 before the embedding's identification. Embeddings were identified with PecanPy (75) using the following parameters: --weighted --dimensions 50 --walk-length 80 --num-walks 10 --directed. Gene similarity was assessed by computing the mutual rank (MR) of the mutual information (MI) using the following formula: MR_{MI} = sqrt(MI_rank * tMI_rank), where MI_rank represents the rank of the MI matrix and tMI_rank represents the transpose matrix of MI_rank. The MI was calculated using the R package Parmigene (76). To select highly similar genes by TF based on its MR_{MI} we used a decay function as follows: $D = e^{-(MRMI - 1)/50}$. D values ≤ 0.05 were taken as highly similar (77). After identifying genes highly similar per TF, we proceed to test the enrichment of PWY and GO terms. Significant GO terms are then mapped to their closest parent terms.

Evaluation of functional predictions

The PWY and GO term predictions were contrasted with the enriched PWY and GO terms identified in loss of function mutants (aka, knockout) using the differential express genes (DEGs) and their corresponding $\log_2$ fold change (FC) values for each TF. We used data from previous studies for KN1 (57), RA1 (59), FEA4 (60), O2 (62), bZIP22 (61), and TB1 (63) reanalyzed in (41). Additional data for MYBR32_m1, WRKY82_m1, HSF13m1m2, HSF18m1, HSF20m1, HSF29m1, HSF29m2, WRKY2m2,

WRKY8m1, and WRKY8m2 were collected from (78). The enrichment of PWY and GO terms were performed with DEG selected based on adjusted P-value as reported by DESeq2 (Padj ≤ 0.05) (79) and following enrichment analyses described above (Methods, *Functional annotation*). The similarities between PWYs and GO terms predicted and those observed in the corresponding knockout were estimated using PWY overlapping and GO semantic similarity, as previously described (Methods, *Functional annotation*). Gene set enrichment analysis was conducted using the R package FGSEA (v1.20) (80) with the parameters: minSize = 5, maxSize = 1000, and eps = 0. The gene sets tested were defined based on the predicted PWYs and GO terms for each TF, considering the available knockout data. The fraction of recovered predictions was calculated by determining the number of significant (P-value ≤ 0.05) PWYs and GO terms out of the total tested.

The comparison of each method's prediction against the random networks was conducted by randomizing each of the four initial networks (GRN, eGRN, CEN, and GAN) 3,000 times, generating 3,000 random versions of each network type. Subsequently, we annotated and integrated each set of random networks following the procedures described in Methods sections *Network integration* and *Functional annotation*. All random networks were generated using the "rewire" function from the R package Igraph (v1.2.4.1), with the following parameters: avoided loops and with niter = NodesInNetwork * 10000.

Prioritization of transcriptional regulators-process associations

All prioritization analyses were conducted using *network-based* results. GO terms with less than 800 genes were retained, and after mapping excessively specific GO terms (≤ 50 genes) to their corresponding GO terms parent. Mapping to parent terms was performed following the procedures described in about (Methods *Functional annotation*). Then, we proceeded to calculate the enrichment score associated with each TF-GO association as follow:

$$E_{ij} = \mathrm{Log}_2[(c/t)/(p/u)] \tag{1}$$

Where E_{ij} is the enrichment score of the TFi with the GOj, c is the intersection of target genes of TFi and annotated genes on GOj, t is the total number of target genes of TFi, p is the total number of genes annotated on GOj, and u is the total number of genes in maize, which in this case refers to the total number of syntenic genes with sorghum (56). All Eij values were subsequently normalized by each TFi and GOj as follows:

$$Zi = (Eij - Ui)/\sigma i \qquad (2)$$

and

$$Zj = (Eij - Uj)/\sigma j \qquad (3)$$

Ui and Uj represent the average enrichment score value for all the GOj associated with TFi and all the TFi associated with GOj, respectively. Similarly, σi and σj represent the standard deviation of the enrichment score value for all the GOj associated with TFi and all the TFi associated with GOj, respectively. Finally, we calculated the reciprocal Z-score (rZ) as follow:

$$rZij = sqrt(max(0, Zi)^2 + max(0, Zj)^2) \qquad (4)$$

Determining sequence similarities among TF paralogs

Sequences for all transcripts associated with the corresponding pair of paralgos were collected from MaizeGDB (https://maizegdb.org/) using genome v4 (54). TF similarities were calculated by averaging the Hamming distance between all transcripts associated with the respective TFs. The Hamming distance was computed using the R package DECIPHER (v2.22) (81) and the "DistanceMatrix" function with the following parameters: includeTerminalGaps = TRUE, penalizeGapLetterMatches = TRUE, and correction = "none".

Mapping of TF-GO terms into specific conditions by gene set enrichment analysis (GSEA)

The GSEA test was performed as previously reported (65). Briefly, GSEA was performed using R package FGSEA v1.18.0 (80) and with Pearson Correlation Coefficients (PCCs) as scoring metric (80). PCC was calculated as the weighted PCC (wPCC) between the TFs and the corresponding genes annotated with the corresponding GO term. The expression dataset and conditions were defined previously (*RNA-seq,*

and co-expression data) and Zhou et al., (2022). Expression values (CPM) were log$_2$ transformed, and wPCCs were calculated using the R package wCorr (Version 1.9.1) (82) with an optimal threshold of 0.4. The genes for each GO term were collected from GAMER (72).

RESULTS

Construction of a maize regulatory network based on multiple layers of information.

To build a multi-layer TF-function association network, we collected previously published co-expression networks, single-nucleotide polymorphisms (SNPs), and reanalyzed publicly available expression, DAP-seq, and ChIP-seq datasets from maize. We included 46 co-expression networks, genetic variation data for 304 maize inbred lines (Table S1), and 283 PDI datasets corresponding to 142 TFs (Table S2 and Figure 1a). This resulted in the identification of ~3.4M TF-gene associations derived from co-expression networks (CENs), ~155.1K associations derived from the gene association networks based on *trans*-eQTL network (GAN), ~1.18M associations in the GRN, and another 112.46K associations based on the *cis*-eQTLs overlapped with GRN interactions (eGRN) (Figure 1b). Construction details for the corresponding networks are described below.

Co-expression network (CEN). To build the CEN layer, we started by collecting 45 previously-published CENs (41), and added an additional network constructed with a subset of expression datasets associated with 304 inbred lines from the Wisconsin Diversity (WiDiv) panel (Mazaheri et al., 2019). The 304 lines were selected based on availability of high-density whole genome sequencing derived SNPs (53), following consistent methods used previously to describe CENs (Zhou et al., 2020) (see *Methods*). Thus, in total, we utilized 46 different co-expression networks to define the TF-target CEN layer (Table S3). Each network was reduced to only maize genes in synteny with *Sorghum bicolor* (56) to avoid potential biases towards non-functional genes when conducting gene enrichment analyses. The syntenic gene filter was also applied to all other network types (*i.e.*, GRN, eGRN, and GAN). On average, we found 1,055 TFs per CEN (Figure S1a) which had - on average - 74 predicted target genes each (Figure S1b). Combining all 46 CENs, we identified ~3.4M TF-target associations involving 1,852 TFs and 23,788 targets (on average, ~1,350

targets per TF; targets include other TF genes). Some TFs had orders of magnitude more targets than the average TF (Figure S1b). For example, ABI3VP1-7 (ABI7) and C2C2-CO-like-8 (COL8) showed >400 targets each (Figure S1b), while COL13 and bHLH127 showed > 6,000 targets each (Figure S1c).

Gene association network based on *trans-e*QTLs (*GAN)*. This layer was built based on *trans*-eQTLs identified in eight distinct tissues encompassing several plant developmental stages. Overall, after quality control and data preprocessing (see *Methods*), we tested between 15.5M and 16.7M SNPs against the expression of 15.3K to 26.4K genes across the eight tissue types (Table S4). After discarding non-significant eQTLs (See *Methods*) and non-syntenic genes (56), we obtained a total of ~22.9M eQTL-gene associations comprising ~10M different SNPs and ~26.4k target genes. These associations were classified as *trans/cis*-eQTL, *cis*-eQTL, *trans*-eQTL, or unassigned eQTL depending on the distance between each eQTL and its corresponding target gene (Figure S2a). We included the additional category *cis*-eQTLt, which describes eQTLs co-localized with their target genes (Figure S2a, yellow). Using this classification, we identified 10.2M unassigned eQTL, 6.7M *trans*-eQTL, 1.20M *trans/cis*-eQTL, 1.18M *cis*-eQTLt, and 3.5M *cis*-eQTL (Figure S2a). Within them, *trans*-eQTLs were used to define the GAN. Specifically, the "source gene" was described as a gene that is co-localized with an eQTL, while the "target gene" is the gene whose expression was explained by the corresponding eQTL (Figure S2a, gene blue and gene yellow, respectively).

After removing redundant associations (*i.e.,* multiple eQTL supporting the same gene-to-gene connection), the resulting GAN harbored ~155K associations, including 23.9K source and 18.9K target genes. To further interrogate the genes captured by the predicted GAN, we classified source and target genes into five functional categories including transcription factors (TFs), co-regulatory factors (CoReg), transcriptional mediator complex proteins (mediator), kinases, enzymes, or others (Table S5) (83–85). The kinase and enzyme classes were the top two with the highest number of target genes, surpassing the TF class (Figure S2b). Similarly, the enzyme was the most frequent target gene class followed by Mediator and CoReg classes (Figure S2c). We counted the interaction frequency between the corresponding classes, and after *other*, enzyme was the functional class with more interactions (13.7K) (Figure S2d). Finally, we

noted that the GAN recovered gene-gene interactions that captured typical TF-target interactions and also PPIs. An example is provided by TF HSF20 which showed 354 targets, including 27 genes previously reported as heat-response related genes (86), as well as five known HSF20 physical interactors (Table S6) (87). Altogether, the GAN highlights an unexplored set of regulatory connections among genes at several hierarchical levels.

Gene-regulatory network (GRN), and *cis-eQTLs* overlapped with *GRN interactions (eGRN)*. To construct the GRN, we collected and reanalyzed 283 PDI experiments associated with 142 different TFs. All the reanalyzed datasets corresponded to TF-centered approaches, including 245 ChIP-seq and 38 DAP-seq experiments. We used a single data analysis pipeline to process all PDI results to control for technical variations in the analysis (See *Methods*). On average, we obtained ~52K TF binding peaks per TF which, in total, represented ~7.6M PDIs (Table S7). The large majority of the predicted peaks were contributed by 215 protoplast ChIP-seq experiments (35), which represented 75% of the analyzed data (on average, ~55K peaks per TF, Figure S3a). To identify high-confidence peaks, we applied two filtering criteria. First, we gathered accessible chromatin regions (ACRs) from the recently published single-nuclei ATAC-seq (snATAC-seq) atlas (37), retaining only TF peaks that overlapped with ACRs. Therefore, we compared all the DAP-seq and ChIP-seq datasets to a shared open chromatin regulatory maize space. Second, we removed peaks with low counts per million (CPM) (as defined by a Z-score ≤ -0.5) for each PDI assay. Overall, we eliminated ~3.8M peaks using the ACR criteria and additional ~1.1M peaks with the CPM cut-off (Figure S3b, c). As expected, most of the eliminated peaks were derived from DAP-seq assays (Figure S3b, c). Comparing low-coverage and peaks co-localized with ACR with their distance to the closest annotated transcription start site (TSS), we found that peaks with the highest Z-values mapped largely to ACRs near TSSs (~10 kb to each side of the TSS) (Figure S3d). These last patterns were observed in all data types (DAP-, ChIP-, and pChIP-seq), supporting the biological relevance of the high-confidence peaks retained. After filtering, we ended with a set of ~3.4M peaks that were used for downstream analyses.

To define target genes, we integrated the peak-TSS distance and their overlap with *cis*-eQTLs (declared when a peak summit and a *cis*-eQTL were ≤ 20 bp away). Integrating these metrics, we categorized the

peaks into three types when they were near TSSs, defined as ≤ 3 kb, and two types when they were further away (> 3 kb and ≤ 50 kb) (Figure S3e). After discarding peaks located far away and without *cis*-eQTL support, we built a GRN and *cis*-eQTL supporting GRN (eGRN) combining all peaks by TF irrespectively of the PDI source. In total, we captured ~1.12M (GRN) and ~1123.46K (eGRN) TF-target interactions, involving 138 TFs and ~23.9K and 13.9K target genes, respectively (Figure 1a, b).

Figure 1. Construction of maize gene regulatory network based on multiple data types.

a. Model indicating the different types of TF-gene associations used to define the network types analyzed

in this work. **b**. Summary of the metrics of the four types of network layers. **c**. Schematic representation of

the pipeline implemented to annotate and evaluate the corresponding functional predictions.

TF functional annotation

One significant contrast among the constructed networks lies in the varying count of TFs and their corresponding target or associated genes (herein interchangeably referred to as target genes), posing a challenge for inter-layer comparisons. For instance, while all four networks feature 111 TFs with at least one associated target gene (see Figure S4a, b), this count dwindles drastically to just 17 TFs when considering TFs with at least ten distinct target genes per network (see Figure S4a, c). This reduction is largely caused by the low number of predicted targets on the GAN layer (on average, ~6.5 targets by gene). Therefore, we implemented three different strategies (*common interactions, common integrations,* and *network-based*) to functionally annotate the TFs present in the corresponding networks. In all three approaches, the annotation was performed based on enrichment of metabolic pathways (PWYs) (71) and GO terms (72) (Figure 1c) (See *Methods*). Briefly, the most conservative *common interactions* approach assumes that only common TF-target interactions between layers (i.e., GAN, GRN, eGRN, and CEN) capture true targets of the corresponding TF, and by extension its function. *Common function* assumes that a TF function is most accurately captured by those functions commonly enriched across different network types. Thus, it prioritizes functions commonly enriched for the corresponding TF across layers. Finally, *network-based* combines all layers to then extract topological properties for each gene. It assumes that each interaction type bore equally valid information about the function of the corresponding TFs. Specifically, it combines all four layers (GAN, CEN, eGRN, and GRN) creating a dense combined network from which to extract physical parameters - embeddings - from each gene in the combined network (See *methods*). The transformation of the networks into a matrix of genes and embeddings allowed the grouping of genes based on similarity of their embeddings. Here, we used the mutual rank (MR) of the mutual information (MI) as the metric to identify highly similar genes in the embedding matrix, to subsequently test for enrichment with PWYs and GO terms between the corresponding genes. The topologically informed strategy allowed

us to functionally annotate TFs independently of the number of target genes predicted at the individual network layers.

Extended descriptions of the functional annotation *Common interactions* and *Common function* are presented as supplemental material (Supplemental data 1). Briefly, using *Common interactions*, we found 11,362 significant associations between TFs and biological processes (TF-process), including 2,812 associations between TFs and PWY (TF-PWY) and 8,550 between TFs and GOs (TF-GO) [False Discovery Rate (FDR) ≤ 0.1, Fisher's Exact Test] (Figure 2a). On average, we identified ~8 PWYs and ~80 GO terms by TF (Figure 2b, yellow boxplots). Combining the PWY and GO term results, we annotated 347 TFs, out of which 235 TFs showed enrichment only in the PWYs analysis (Table S8). The remaining 112 TFs showed enrichment with both PWYs and GO terms (Figure 2c). *Common function* identified 7,081 TF-process annotations (727 TF-PWY and 6,354 TF-GO) (Table S8) (Figure 2a), corresponding on average to 3.5 different PWYs and 57.7 different GO term associations per TF (Figure 2b). These associations encompass annotations for 204 TFs through PWY enrichment and 110 TFs through GO term enrichment (Figure 2c).

Network-based. We combined all four layers and scaled the interaction frequencies. Using the scaled interaction frequencies, we identified low-dimensional representations (embeddings) for each gene/node in the combined network that works as descriptors of the corresponding gene (Figure S6a) (See *Methods*). The combined network included 4.6M interactions associated with 36.4K genes. Unlike the previous two strategies, this method generated an equal number of embeddings for each gene in the network, thus uncovering genes with similar properties, including TFs present in the CEN and/or GAN layers without data on the GRN/eGRN layers. The similarity of the embeddings was used as a metric to assess the wiring similarities between genes within the combined network (See *Methods*) (Figure S6a). On average, we found 235 genes per TF with highly similar embedding profiles [Distance (D), ≤ 0.05, See *Methods*] (Figure S6b). As in previous approaches, we annotated the corresponding TFs by assaying the enrichment in PWYs and GO terms within the set of genes with highly similar embeddings (Figure S6a). In total, we found 23,796 TF-PWY and 7,722 TF-GO significant associations (FDR ≤ 0.1, Fisher's Exact) (Figure 2a) (Table S8),

which on average captured ~7 PWYs and ~8 GO terms per TF (Figure 2b). Combining both assays, we annotated 2,910 different TFs, out of which 1,030 showed enrichment with both PWYs and GO terms (Figure S6c). To note, these 1,030 TFs belong to 82 different families (including co-regulators) representing ~34% of the total proteins annotated in the corresponding families (Figure S6d). This highlights the potential of the method to annotate TFs with unobserved interactions, such as no previously published ChIP-seq results.

Comparing all three methods, the *network-based* approach enabled us to identify the largest number of TF-PWYs, and the lowest number of TF-GOs associations (Figure 2a). Additionally, this approach had the lowest average of PWYs and GO terms per TF (Figure 2b). Unexpectedly, *network-based* and *common target* methods predicted a similar number of PWYs per TFs, which contrasts with the significantly lower number of GOs between *network-based* and the other two methods (Figure 2b). Noteworthy, the number of TF annotated by the *network-based* method is >2.5 times larger than the other two methods (Figure 2.c, left panel). Thus, by using the *network-based* approach, we functionally annotated 2,917 TFs.

Evaluation of functional predictions

To validate the predictions derived from each of the three methods, we used 21 previously published knockout assays associated with 13 different TFs (Table S9) (41, 78) and assayed the accuracy of each method by two independent strategies. First, we questioned the overlapping between our predictions and observed PWYs/GO terms within DEGs for the corresponding knockouts by each method. Second, we tested the gene set enrichment analysis (GSEA) of the predicted PWYs/GO terms within the corresponding TF knockouts, using the $\log_2$FC of as ranking metric (88) (See *Methods*). Overall, predicted PWYs from the networks - without distinction of the methods - showed poor overlapping with PWYs observed in the knockout assays, as well as low recovery of PWY significantly enriched within DEGs as estimated by the GSEA (Figure S7). Conversely, comparisons between predicted and observed GO analyses showed similarities [measured by the GO semantic similarity (GSS)] different than the expected by chance (P-value ≤ 0.05) (Figure S8). Overall, the GO terms from knockouts and the GO terms predicted by *network-based*

and *common function* are significantly more similar than the *common targets* predictions (higher GSS values, P-value ≤ 0.05, Wilcoxon test) (Figure 2d). When a prediction was available, the *network-based* method recovered the GO terms with the highest GSS values among all the methods (Figure 2d, TB1 and FEA4 results). Additionally, we observed that seven knockouts lacked predictions from *network-based* methods, while eight others had predictions only with *network-based* methods (Figure 2d, e). This variability in predictions can be partly attributed to the low number of target genes (when the prediction is absent; Figure S9a, TFs with Z-score ≤ 0) and the absence of data in at least one of the four layers (i.e., GRN, eGRN, CEN, and GAN) (when the *network-based* method is the only one making the prediction, Figure S9b). Consistently with the GSS analysis, the GSEA results indicate that GO terms recovered with the *network-based* and the *common function* are more consistently identified across the different knockouts (Figure 2e). When taken together, our results indicate that *network-based* predictions are robust to presence-absence variation in the data across layers, but is susceptible to the number of targets by TF. By extension, our findings also indicate that *common targets* and *function* predictions are more sensitive to the absence of data in at least one of the layers.

Combining all GSEA results, we found that on average 25% of total GO predictions show significant GSEA scores (P-value ≤ 0.05, as calculated by FGSEA implementing in R), denoting a modest recovery rate of GO terms (Figure 2e). We used the tissue specific expression of TFs as a proxy to better explain the relationship between the fraction of GO recovered by GSEA and the tissue/condition/genotype variation among the corresponding TFs. Including all the TFs for which we obtained at least one PWY/GO term prediction and using the Tau expression-specific index as a metric (89), we found a bi-modal expression distribution with ~55% of the TFs trending into a sample-specific expression fashion (Figure S9c, Tau ≥ 0.65). Just four out of the 13 TFs tested in the knockout analyses are expressed in a tissue-specific fashion (P-value ≤ 0.05) (Figure S9d, labeled in green). The top four TFs included RAMOSA1 (RA1) (Tau 0.99) and TEOSINTE BRANCHED1 (TB1) (Tau 0.96), which are also the top two TFs with the largest fraction of GO term supported by the GSEA (Figure 2e). Hence, the results support the notion that part of the low fraction of overhauling between the GO terms predicted compared with observed annotation from the

knockout can be attributed to the differences in biological conditions developmental stages used on the knockout and compare with the RNA-seq datasets used on the prediction analyses.

We also evaluated the identification of GO terms from random networks to determine which method recovered the lowest fraction of false positives (Supplemental data 2 and Figure S10). Overall, we observed a high degree of overlap between the predictions made by *the common target* and the *common function* methods, suggesting that the two methods were equally noisy. Consequently, *network-based* method predictions always result in a lower number and have more differences in GO terms compared to the observed GO terms from our predictions (Figure S10). In conclusion, the *network-based* method is a superior approach within the given context of these datasets. Consequently, we exclusively relied on *network-based* predictions for subsequent analyses.

Figure 2. Annotation and evaluation of TF functional annotation by contrasting predictions with knockout assays and random networks.

Total pathways (PWYs) and GO terms predicted per integration method after combining all TFs predictions (**a**) and per TF (**b**). **c**. Upset plot comparing total TFs annotated by each method and annotation system. Colors indicate the groups of TFs functionally annotated by all three methods by enrichment with PWYs (fuchsia), GO terms (blue), and both PWYs and GO terms (green). Black groups indicate TFs annotated by at least one of the methods and annotation systems. **d**. Boxplot of the GO semantic similarity for the top 10 most similar GO terms observed in knockout assays for each of the predicted GO terms per TF and methods. **e**. Stacked bar plot indicating the fraction of the GO terms predicted and significantly enriched - by GSEA analysis - in the knockouts. **f**. Violin plot showing the fraction of random networks with at least one significant (FDR $\leq$ 0.1, Fisher exact test) GO term by TF. Asterisks indicate P-value significance (*: p $\leq$ 0.05, **: 633 p $\leq$ 0.01, ***: p $\leq$ 0.001, ****: p $\leq$ 0.0001, two-sided t-test). "TFm" denotes multiple mutant lines for the same TF.

Prioritization of regulators by biological process

The network-based method detected approximately 7.7K TF-GO associations, encompassing 1,036 TFs and 2,219 GO terms (Figure 2a, c & Figure S6b, c). For ease of TF comparison, we retained associations involving GO terms within the biological process (BP) category. Additionally, to minimize GO term redundancy, we mapped GO terms with less than 50 genes to their nearest GO term parent. After applying the filters, we continued with 4,337 TF-GO associations, comprising 902 TFs and 559 GO terms. The distribution of TF-GO associations obtained follows a scale-free distribution (Figure 3a, b). Typically, highly interconnected GO terms and TFs suggest a greater number of annotated and targeted genes, respectively. Nevertheless, we did not discover any evidence linking the gene count per GO term or the target gene count per TF to their respective degrees (Figure 3c, d). Therefore, these analyses highlight GO terms whose regulation may depend on multiple TFs, and TFs that may contribute to regulating several biological processes.

From a gene regulation perspective, the association of a TF with multiple GO terms is suggestive of the influence of additional TF interactions. To assess the contribution of individual TFs to their respective GO terms, we calculated a scaled enrichment score (Z-score of the enrichment) for each TF and GO term (See *Methods*). Utilizing the scores to assess the strength of individual TF-GO associations, we observed that only 3.4% (151/4,337) of the TF-GO associations exhibit high enrichment scores (Z-score ≥ 1) (Figure 3e, top right corner). This implies that most of the analyzed TFs/GO terms have multiple associations of comparable significance. We combined both Z-scores (per TF and GO term) into a reciprocal Z-score (rZ, See *Methods*) to define a single ranking metric for each TFs-GO term association. We evaluated the ranking after grouping GO terms into specific biological processes (Figure 3f, and Figure S13). We highlight here 46 GO terms for abscisic acid (ABA) metabolism-related targeted by 44 TFs, 62 GO terms for lipid metabolism-related targeted by 55 TFs, 47 GO terms for phenylpropanoid metabolism-related targeted by 47 TFs, and 50 GO terms for leaf development-related terms that were targets of 50 TFs (Figure 3f). Using the rZ score as filter (rZ ≥ 0.5, to facilitate data displayed in Figure 3f), we narrowed down the list to 25, 27, 15, and 19 TF candidates to control the corresponding processes (Figure 3f, dots with name label included). Examples included the top two TFs ABA for metabolism related, *NAC56* and *WRINKLED2* (*WRI2*), with *WRI2* also being on the top three TFs related to lipid-related metabolism (Figure 3f, second panel). Five TFs (*WOX9a*, *OFP39*, *Zm00001d024353*, *EREB149*, and *LBD24*) out of the initial 47 phenylpropanoid-related TFs were previously identified as maize regulators of phenolic-related genes by yeast-one hybrid assays (Y1H) (90). Altogether, this highlights the potential biological relevance of the associations predicted by this analysis.

To identify TFs that play a higher-level role in controlling a biological process by controlling other TF genes, we calculated the ratio of TFs targeted by other TFs within specific GO terms with respect to the total number of TFs targeted by the corresponding TF. Given that we searched for TF associated with a common function, this ratio represents the weighted proportion of feed-forward loops associated at the level of biological process compared to the overall TF targets of the corresponding TFs, defined here as upstream regulator score (URS). We calculated the URS for the twenty different processes, including eight hormone-

, seven metabolic-, and five developmental-related processes (Figure 3g). Cytokinin- and shoot-related GO terms were the top two processes with the highest score, with *COL13* and *RA1* as their top regulators, respectively (Figure 3g). To note, *COL13* was previously associated with carbon metabolism (35), and is also differentially expressed in loss-of-function mutants of the autonomous floral induction *indeterminate1* (*1d1*) (91, 92). Thus, our results suggest a role of maize *COL13* in connecting cytokinin, carbon metabolism, and flowering, associations previously reported in other plant systems (93, 94). Similarly, *RA1* was predicted as the top upstream regulator of shoot-related processes, and RA1 itself was linked with shoot system development (Figure 3g, process number 17), both of them functions previously associated with *RA1* (59).

To further understand the regulatory landscape of the four processes previously described (Figure 3f), we selected the top two predictions (based on URS score) for each process and traced their TF targets back to the original networks (*i.e.*, GRN, eGRN, CEN, and GAN) (Figure 3h). Specifically, we looked for regulators directly upstream of any of the top TFs as predicted by the rZ analysis (Figure 3f). Without exception, we found at least one upstream regulator directly targeting in the GRN network at least one of the top three TFs from the rZ analysis (Figure 3h). To highlight an example within the ABA-related process network, EREB17 targeted *NAC56* and *WRI2* (top TFs in rZ analysis), and bHLH43 (URS top gene) targeted *WRI2* and *EREB17* (Figure 3f, h, first panel). This configuration forms a feed-forward loop with bHLH43 on top. Within the lipid-related network, ARF14 targeted *WRI2* and *PRH65* (top two and three by rZ score) (Figure 3f, h, second panel), and *HB33* targeted LBD23 (top in rZ) in the phenylpropanoid-related network (Figure 3f, h, third panel). Finally, WRKY25 (top USR) targeted the *MYBR4* and *EREB126* (top two TFs in rZ), as well as BZR2 (top two in URS analysis) on the leaf-related network. Thus, our results predicted specific regulatory interactions for further experimental validation.

Figure 3. Prioritization of regulators by biological process using network-based prediction.

Out degree (**a**) and in degree (**b**) distributions of the TF-GO term predictions obtained from the network-based integration analysis. **c** and **d**, scatter plots indicating the frequency (as density) of the number of TFs by GO terms (in degree) and GOs per TF (out degree) as a function of the number gene annotated per GO

term (**c**) and target genes per TF (**d**), respectively. **e**, Scatter plot indicating the frequency (as density) of the TF-GO term scaled enrichment scores, which allows to rank GOs highly enriched with specific TF (GOz) and TFs highly enriched with specific GO term (TFz). The enrichment was calculated only with TF-GO term associations already predicted in previous analysis. The orange dotted line indicates TF-GO term associations with enrichment score one standard deviation over the observed average for the corresponding TF and GO term (Z-score of enrichment ≥ 1 for both GO term and TF). **f**. Scatter plot with reciprocal Z scores (rZ) of four different biological process mapped into the GO and TF scaled score coordinates as presented in **e**; GO terms were grouped as follow: ABA-related (GO:0009737, GO:0009738, GO:0009688, and GO:0009788), Lipid-related (GO:0031408, GO:0006099, GO:0006635, GO:0019915, GO:0006629, GO:0019375, GO:0044255, GO:0016042, GO:0051790, GO:0008610, GO:0009062, and GO:0045332), phenylpropanoid-related (GO:0009963, GO:0009698, GO:2000762, and GO:0009699), and leaf-related (GO:0009965, GO:0048366, GO:0010305, GO:0010150). TF name/gene id labels are included for TFs with rZ ≥ 0.5. **g**. Scatter plot with TFs ranked by their upstream regulator score (URS) by biological process. TF name labels are included for TFs with rank ≤ 2. Square brackets indicate an arbitrary biological process index which matches with the number in square brackets of the corresponding TF names. All URS scores were calculated based on the original GRN, eGRN, CEN and GAN networks. **h**. Heatmap with top two TFs (y axes) from the URS analysis (**g**) for the four biological processes presented in **f**. X axes indicate the corresponding TF targets. Colors indicating the network(s) source of the corresponding interactions.

Mapping TF-GO process into specific conditions

TFs generally work in a tissue/condition-specific manner. Thus, to gain a better understanding of the regulatory condition that most closely captures the functional space of the corresponding TFs, we performed a gene set enrichment analysis (GSEA) with the 4,337 TF-GO associations predicted to map these associations to specific developmental and environmental conditions. Specifically, we used the expression data utilized previously in the constructions of the 45 co-expression networks (41), as well as, additional six expression datasets of maize plants subjected to heat and cold stress (See *Methods*) (95). Unlike a typical

GSEA, we employed the Pearson correlation coefficient (PCC) as the gene ranking metric, as we described previously (65). It is worth noting that this analysis does not discover new TF-GO associations. Instead, this GSEA ranks each of the previously described TF-GO term associations in the context of the expression datasets tested (referred here as co-expression networks). Consequently, we first determined the percentage of the co-expression networks in which the corresponding TF showed a significant gene-set enrichment (FDR ≤ 0.1) with at least one of their associated GO terms. On average, 76% of the tested TF-GO terms exhibit significant co-expression (FDR ≤ 0.1) in at least one network (Figure S14a). This suggested that most of the TF-GO term associations tested can be mapped to a particular biological condition. Moreover, we calculated the average percentage of significantly co-expressed TF-GO term associations in each co-expression network per TF. This helped us understand the functional diversity of each TF. In general, each TF had 53% of GO terms that showed significant enrichment in each co-expression network (Figure S14b). TFs with the lowest and highest average percentages of significantly co-expressed GO terms also possessed the lowest number of associated GO terms (Figure S14c). Thus, this reveals TFs with a reduce number of associations which also holds a completely different functional space, i.e., TFs associated with few GOs and co-expressed with them in either a few conditions or in almost all of them.

To better understand the pattern of GO terms co-expressed by TF, we clustered TFs by the percentage of co-expressed GO terms. In total, we identified three clusters of TFs (Figure 4a). Cluster 1 (291 TFs) captured TFs with the highest average percentage (69%) of GO terms co-expressed by network, while Cluster 3 includes 235 TFs with the lowest percentage (45%). Cluster 2, on the other hand, encompasses 303 TFs where approximately half (50%) of their GO terms are significantly co-expressed in at least one of the tested networks (Figure S14d). To note, Cluster 3 has lower number of co-expression networks by TF (Figure S14e), and the largest number of GO terms by TF (Figure S14f). The latter suggests that a large fraction of the TF-GO terms association in Cluster 3 fail to provide relevant biological information, at least based on the condition tested here (Figure S14d). This contrasts with TFs in Clusters 1 and 2, which are associated with a smaller number of GO terms but have a potentially broader scope of influence.

To explore the biological significance of the TF-GO mapping strategy, we focused on *bHLH43*, *ARF14*, *HB33*, and *WRKY25*, which we identified as upstream regulators of different biological processes (Figure 3). Our analysis revealed that *bHLH43*, *ARF14*, *HB33* (Cluster 1, Figure 4a), and *WRKY25* (Cluster 2, Figure 4a) has significant TF-GO terms association in almost all the co-expression networks tested (Figure 4b), which validates our prediction of these TFs as upstream regulators of the corresponding process. Additionally, the GSEA results unravel potential condition-specific activators and processes as a function of the condition tested, as indicated by the normalized gene set enrichment score (NES) (Figure 4c and S15). We highlight here *ARF14*, predicted as an upstream regulator of lipid-related processes (specifically fatty acid beta-oxidation, GO:0006635), which also has an enrichment profile similar with tryptophan catabolic process (GO:0006569) and phenylpropanoid biosynthetic process (GO:0009699). All three metabolism pathways are known for their interdependency because fatty acids provide precursors for phenylpropanoids, and because phenylpropanoids and tryptophan share shikimate pathway precursors (96, 97). Specifically, we selected three networks in which the three pathways showed positive NES score (N28, genotype by root variations), negative NES score (N29, genotype by leaf variations), and combined effects (N11, genotype by seedling variations) (Figure 4d). In all three cases, the GSEA profile validated the dynamics suggested by the NES score in terms of the gene set rank accumulation. However, we noticed that the highest and lowest positive and negative NES peaks occurred around the 5,000 and 20,000 gene ranks, respectively (Figure 4d, marked with dashed pink lines), indicating that many of the pathway-related genes are not highly correlated with the corresponding TF. We also observed a double weak peak in the case of tryptophan catabolism and fatty acid beta-oxidation in seedlings (Figure 4d, right and left panel - network N28 and N11), suggesting that *ARF14* may have both positive and negative expression effects on the corresponding pathway, even within the same network. To test our hypothesis, we specifically examined network 11, which displayed a variable NES sign across all three pathways associated with *ARF14*. We grouped the genes corresponding to each pathway based on their expression profiles (Figure 4e). As anticipated, we found a subset of genes that exhibited expression changes not related to *ARF14* (labeled as C1 in Figure 4e). We also observed that there is a group of maize lines in which an increase in *ARF14*

expression aligns with increased expression of the phenylpropanoid pathway (positive NES peak in Figure 4d, N11 middle panel), which also exhibited an increase in *ARF14* expression (Figure 4d, Samples 50 to 150). This observation supports the idea of different regulatory associations even within the same pathway. Finally, since this network was constructed based on genotype variations, we can identify the specific lines where *ARF14* has lowest (left labeled dots in Figure 4e, top panel) and highest (right labeled dots in Figure 4e, top panel) expression levels. These would correspond to lines in which manipulating *ARF14* could be expected to have the greatest and lowest impact on the associated target pathways.

Figure 4. Mapping to TF-GO terms associations to specific conditions.

Heatmap showing the total (**a**) and a subset of four TFs (**b**), indicating the percentage of GO terms significantly enriched in the GSEA analysis for each TF across the respective co-expression networks tested. **c**. Heatmap displays the normalized sentiment score (NES) of ARF14 and its three associated GO terms across 39 networks where its GSEA was assessed. GSEA results are not shown in the remaining 12 networks due to the absence of ARF14 expression or the lack of statistically significant (FDR > 0.1) in all three GO terms. **d**. GSEA profile from three networks in **c**, displaying the actual accumulation of gene sets along the ranking established based on the co-expression of ARF14 with the whole maize genome. **e**. Locally weighted scatterplot smoothing (LOESS) line plot with the scale CPMs of the gene set in **d** across different samples within the respective networks. Genes were clustered after normalizing and scaling their expression values. The dashed line represents the scaled expression of *ARF14*, with samples on the x-axis organized according to *ARF14* expression levels.

Topological properties predict TF paralogs redundancy

Substantial efforts have been made to comprehend and anticipate the functional redundancy between maize paralogs in subgenomes (26, 27, 29, 30), yet this remains challenging. We anticipate that if a pair of paralogs exhibit functional redundancy, these differences may manifest in their topological properties, *i.e.*, functional redundant paralogs would display similar properties indicating a comparable network arrangement. To assess the similarity between paralogs, we generated a distance matrix from the embeddings using the mutual rank (MR) of mutual information as metric (Figure 5a). Next, we mapped TF paralogs (56) and analyzed their MR (Figure 5a, "1" in orange circle) and the similarity of their MR profiles (Figure 5a, "2" in pink circle) with all the genes in the embeddings matrix. We used these metrics as a proxy for understanding the similarity of their embeddings and the similarity of their resemblance with other TFs examined, respectively. We also differentiated between paralogs located on the same chromosome, serving as a reference for pre-speciation tandem duplicates. In total, we tested 932 TF pairs, and regardless of the metric used, TF paralogs situated on the same chromosome demonstrated greater similarity compared to TF paralogs on different chromosomes (Figure 5b, c). We combined both scaled metrics to identify highly

similar TF pairs (Figure 5d). As expected, both metrics were correlated (PCC 0.68), yet they effectively served the purpose of identifying TF paralog pairs that were highly similar. We tallied the number of interactions after the embedding integrations (Figure S6a, b) for the top ten TF pairs that were most and least similar (Figure 5e). The top ten most similar TF pairs have several TF-GO common associations (Figure 5e, TF pairs highlighted in light brown), contrary to those observed within the top least similar, which have none (Figure 5e, TF pairs highlighted in gray). Additionally, seven TF pairs mapped to the same chromosome out of the top ten most similar TF pairs (Figure 5e, TF pairs with asterisk). This last supports our prediction of similarity in topological properties as a potential predictor of paralogs functional redundancy.

To discriminate better TF pairs by the similarities in the embedding space, we organized all TF pairs into nine bins using scaled similarity metrics (Figure 5d, areas defined by the dashed black lines and represented top right corner of Figure 5f). These categories range from the least similar TF pairs in the first bin (I) (Figure 4d, left bottom corner) to the most similar pairs in the last bin (IX) (Figure 4f, right top corner). We then compared the number of paralogs that are tandem duplicates and those that are not, which confirmed the observation from the top ten TFs (Figure 5e), exemplified by bin IX containing 5-7 times more tandem duplicates than the other bins (Figure 5f). We also quantified shared TF-GO associations using the Jaccard index. By considering bin I as a reference, we detected significant differences ($p < 0.05$, two-sided t-test) across five distinct bins (Figure 5g), primarily categorized based on the scaled correlation (Z_{SCC}) between TF pairs (Figure 5d, x axes). Furthermore, bin IX exhibited the utmost values, validating the predictive capacity of embedding similarities for functional redundancy in TF paralogs.

We speculate that a low similarity in topological properties indicates functional divergence. On the other hand, a high similarity suggests conservation of function. Therefore, we anticipated observing similar patterns of variation either in the protein sequence or in its regulation, along the corresponding pairs of paralogs. Specifically, we expected to see greater differences in either the protein sequence or the expression of paralogs in bin I compared to the other bins. To investigate if this is the case, we focused only on TF pairs from bins I and IX, which represent the most different pairs. We calculated the Hamming

distance of the amino acid sequences, which measures the number of changes in amino acids between the corresponding paralog pairs (98, 99). Additionally, we analyzed TF paralogs co-expression as an indicator of variation or conservation at the *cis*-regulatory level (100–102). Overall, we did not observe any differences in the Hamming distances between TF paralogs highly similar or dissimilar at the topological level (Figure 5h). However, TF pairs in bin I showed slightly lower average co-expression (PCC = 0.4) compared to those observed for TFs in bin IX (PCC = 0.5). Yet, the mapping of the co-expression along the Hamming distance for the corresponding TF pairs, allowed us to differentiate TF paralogs that may be undergoing neofunctionalization/subfunctionalization due to variations in their protein sequences or their expression profiles (Figure 5i,j). A striking example of the former is observed in *MYBR1* and *MYBR81*, which have large numbers of changes in their sequences (Hamming distance close to 1), distinct embedding profiles (bin I), and yet display high co-expression (PCC > 0.9) (Figure 5i). Additionally, within the groups of TFs sharing similar embedding profiles (bin IX), we identified TFs exhibiting high sequence conservation (Hamming distance close to 0) and similar expression, implying a significant degree of redundancy (e.g., *MADS73* and *TU1*) (Figure 5j). In contrast, C3H53 and C3H36 had low co-expression but high conservation (Hamming distance close to 0), indicating differences in their regulation (Figure 3j). Taken together, the combination of embedding similarities, protein amino acid similarities, and co-expression enables the identification of TFs that are either potential variable or redundant, a key observation for understanding functional redundancy.

Figure 5. Network embedding as predictor of TF paralogs with functional variation.

a. Diagram illustrating the key stages of comparing TF paralogs through embedding similarities. **b** and **c**. Box plots displaying the MRMI of TF pairs (**b**) and the Spearman correlations (SCC) of the observed MRMI profiles (**c**) derived from the embedding. d. Combined scaled scores of the MRMI and SCC for TF pairs. Black dashed lines with Z-scores of -0.5 and 0.5 indicate values below and above the average observed standard deviation. **e**. Heatmap indicating the total number of associated genes for the top ten TFs, on the top right corner and the bottom left corner TF pair (**d**). G1 and G2 represent the number of unique genes associated with the first and second TFs in the corresponding pair. G1:G2 indicates the common associations between the corresponding pair. F. Bar plot indicating the total number of TF pairs by bin. Bins are indicated on the interval box, which is a map of the zones in the plot in (**d**). **g**. Box plot with Jaccard index (as an approximation of common associated genes) by TF pair by bin (as presented in **f**). **h**. Jitter plot displaying amino acid (AA) differences (Hamming distance) between TF pairs in bins I and IX. **i** and **j**, Jitter density of points representing AA Hamming distance and co-expression (measured as PCC) for TF pairs in bin I (**i**) and IX (**j**). Asterisks indicate P-value significance (*: $p \leq 0.05$, **: 633 $p \leq 0.01$, ***: $p \leq 0.001$, ****: $p \leq 0.0001$, two-sided t-test). "TFm" denotes multiple mutant lines for the same TF.

DISCUSSION

Cells utilize complex networks of molecules to integrate and synergistically regulate their activities. The rapid generation of multi-omic genomic data in plants has led to growing interest for stabilizing integration strategies (17, 45–51). In this study, we analyzed distinct data types (PDIs, expression, and SNPs) and constructed four different molecular networks (layers). Considering the inherent challenge posed by the variations in each respective network, we examined three distinct integration methods and employed two different strategies to functionally annotate TFs (Figure 1). Overall, our findings indicate that the integration of multiple layers based on *common targets* and *functions*, although more intuitive, are less accurate in recovering observed GO terms in knockouts (Figure 2d, e). Contrary, we find that the *network-based* approach is highly resiliant to noisy datasets (Figure 2d, e). we hypothesize that this resilience is rooted in the inherent nature of the embedding generation process, as it is highly improbable to observe

similar wiring patterns across layers, despite the expected presence of false positives within each respective layer. Of equal significance, the *network-based* approach facilitated predictions for a considerably larger number of TFs (Figure 2c), thereby influencing the design of future experiments aimed at uncovering and validating specific TF functions in maize.

Thus, focusing on the *network-based* integration to prioritize transcriptional regulators associated with specific biological processes, we gained valuable insights into potential regulatory mechanisms underlying maize metabolism and development processes. Also, we showed that only ~3% of our prediction had a single TF as the primary regulator of the corresponding GO term, consistent with the notion that the vast majority of TFs contribute to the regulation of multiple functions, and that the regulation of a biological process requires the involvement of multiple TFs (35, 90, 103, 104). This last is a distinctive feature of GRNs, the regulatory repertoire of TFs is leveraged through interactions with other TFs (105, 106). This phenomenon has been previously observed in maize, both at the level of TF-target genes (35, 90) and in the organization *cis*-regulatory elements across cell types (37), supporting our results. Additionally, we showed the presence of a feed-forward motif within our results, which is known as a mechanism for reinforcement of regulatory signals (107). Thus, our findings extend the concept to a broader framework, encompassing biological processes, and have significant implications for future biotechnological applications, such as the targeted modification of specific metabolic processes.

TFs generally work in a tissues/condition-specific manner (108). Here, we mapped our TF-GO function prediction into 51 different datasets (41, 95) using GSEA and TF-gene correlations as ranking metric (109). Our analysis revealed that most TFs, including those in clusters 1 and 2 (Figure 4a), regulate multiple GO terms across various conditions. However, a smaller percentage of TFs are involved in regulating their corresponding GO terms in a limited number of conditions (cluster 3, Figure 4a). This emphasizes the existence of a condition-specific complex regulatory landscape. Additionally, since multiple TFs can contribute to the same function (Figure 2f), the results suggest that the specific behavior in each condition is established by the combined action of multiple TFs instead of individual TFs, as suggested by previous studies (108). We use ARF14 as an example here to highlight the potential of our analyses, which is known

to play a role in auxin response factors (110). Our analyses suggest that ARF14 is a potential regulator of various metabolic and developmental processes (Figure 3h and Figure 4b-e). Specifically, it may play a major role in regulating three metabolic pathways, including tryptophan catabolism. Since tryptophan serves as an essential precursor for indole-3-acetic acid (IAA), the primary auxin in plants (111), this finding provides a plausible mechanistic functional connection between hormonal and metabolic control. Furthermore, our GSEA analysis suggests a co-expression coordination between each pathway and ARF14, but this coordination is specific to particular conditions, further supporting the concept of interaction between metabolic pathways (112–116).

The understanding of functional redundancy between maize subgenomes remains far from complete (26, 27, 29, 30). In this study, we used the similarity of embeddings generated after molecular network integration to identify functionally redundant TFs. Our findings show that TF paralogs with more similar embeddings have more common targets and are often found as duplicates in tandem (Figure 5e-g). While we did not find significant differences in the variation of amino acid sequences between TF paralogs with low or high similarity in embeddings space (Figure 5h), we noticed slightly higher co-expression similarities between TF paralogs with closer topological similarities. Thus, although we can't fully explain the reasons behind the similarity in network connections between TF paralogs, it is important to note that by integrating information about proteins sequence, expression, and embedding similarities, we can identify specific candidates for functional neofunctionalization/subfunctionalization and redundancy (Figure 5i, j). These observations need to be tested through experimental validation.

In summary, our findings demonstrated that integrating all multi-omic datasets through embedding identification followed by clustering allows us to identify genes with similar wiring on the fully integrated network. It is worth mentioning that the embedding similarities create an association network themselves, encompassing gene-gene associations for over 24K genes. Here, we used this strategy to annotate TFs. However, this also defined a conceptual framework for expanding this annotation strategy to the other types of genes, such as the annotation or definition of metabolic pathway based on the topological embedding similarities. Thus, our results pave the way for the multi-omic integration of similar datasets in other plant

systems, facilitating the selection of TFs for crop improvement, metabolic engineering, and basic gene regulation understanding of the corresponding processes.

DATA AVAILABILITY

Code availability.

Codes associated with analysis and figures of raw data processing, network integrations, and processing of results presented here are available at https://github.com/gomezcan/Multi-Omic_Network_Integration_Maize.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR online.

SUPPLEMENTAL DATA LEGENDS

Supplementary data 1: Expanded descriptions of the TF functional annotation based on Common interactions and Common function.

Supplementary data 2: Evaluation of functional prediction by comparing with random networks.

SUPPLEMENTARY FIGURES LEGENDS

Figure S1. Summary of total TFs and their interactions used in the co-expression network (CEN) layer.

a. Histogram showing the frequency of TFs with at least one target gene per CEN. The dotted gray line indicates average TFs along all 46 CEN. **b**. Boxplot indicating total target genes per TF across the different CENs. CEN are named following the Zhou et al., (2022) nomenclature. Orange labels highlight TFs with the largest number of target genes in several CENs. **c**. Histogram indicating the frequency of total target

genes per TF after combined results from all 46 CENs. Descriptions of CEN in (b) can be found in Table S3.

Figure S2. Defining a gene association network (GAN) based on *trans*-eQTL.

a. Model indicating total eQTLs identified and the classification schema used to define *trans*-eQTL, *trans*/*c*is-eQTL, *cis*-eQTLt, and *cis*-eQTL. Within them, trans-eQTLs were used to define the GAN. In the context of trans-eQTLs, a source gene (in blue) was defined as a gene whose promoter (2kb upstream from annotated TSS) or gene body overlapped with an eQTL. Genes whose expression is explained by the SNP variation were defined as gene targets (gene in yellow). **b** and **c**. Each source and target gene was classified into one of five functional categories to count the number of associations by category (unclassified genes defined as other). **Left panel**, Boxplot indicating the number of targets (**b**) and source (**c**) genes by each gene category. **Right panel**, stacked bar plots indicate the fraction of each gene category over the total genes in GAN. **d**. Bar plot indicating total interactions by gene category pair.

Figure S3. Establishing the maize gene regulatory network (GRN) layer based on protein-DNA interaction data.

a. Density plot with peak distribution by PDI data type. **b** and **c**. Stacked bar plot with fraction of peaks mapped to accessible chromatin region (ACR) (**b**) and with low peak coverage (**c**) CPM scaled and filtered; $Z \leq -0.5$. d. Locally weighted scatterplot smoothing (LOESS) line plot of Z-scores by peak in 10 kb bins around 200 kb of the closest transcription start site (TSS). **e**. Classification schema (top) and corresponding proportion of total combined peaks (bottom, first stacked bar plot) and peaks by method (bottom, second stacked bar plot) utilized for determining target genes. Specifically, peaks proximal ($\leq$ 3 kb) to the TSS were classified as follows: peaks lacking *cis*-eQTL support (light purple peaks), peaks with *cis*-eQTL support and consistent target prediction (light green peaks), and peaks with *cis*-eQTL support but discordant target prediction (yellow peaks, Figure S3e). These categories represented 54.9%, 1.9%, and 0.1% of the total analyzed peaks, respectively. Similarly, peaks located far away were classified as peaks with (3.3%) and without (39.5%) *cis*-eQTL support (light blue and gray peaks, respectively).

Figure S4. Overview of strategy to annotate TFs based on common targets.

a. Schema of pipeline used to annotate TFs based on common target genes amount layer (GAN, GRN, eGRN, and CEN). **b** and **c**. Venn diagram indicating the number of common TFs with at least one (**b**) and ten (**c**) target genes. **d**. Venn diagram indicating total common interactions (TF-target gene) among layers.

Figure S5. Overview of strategy to annotate TFs based on common functions.

a. Pipeline schematics used to annotate TFs based on common function amount layer (GAN, GRN, eGRN, and CEN). **b**. Bar plot indicating total TFs annotated by layer and by type of function. **c** and **d**. Venn diagram indicating the number of TFs with at least a PWY (**c**) and GO term (**d**) commonly enriched among the corresponding layers.

Figure S6. Network-based strategy to annotate TFs.

a. Pipeline schematics used on the integration of layers to identify TF with similar topological properties, defined here as network-based TF annotation. **b**. Histogram plot indicating the distribution of genes associated per TF. **c**. Stacked bar plot with total TFs annotated by enrichment with PWYs and GO terms. **d**. Bar plot indicating the percentage of TFs annotated for the 82 TF families (and co-regulator) with at least a TF annotated (**c**).

Figure S7. Predicted PWY overlapped poorly with PWY observed in knockouts assays.

a. Heatmap showing the count of overlapped PWYs between predicted and observed PWY in knockouts per method. The violet box signifies significant overlap (P-value 0.05, Fisher test). An empty box (white) denotes no predicted PWYs for the corresponding TF. Square braking indicates the number of PWYs significantly enriched in the corresponding knockout (P-value <= 0.05, Fisher test). **b**. Stacked bar plot indicating the fraction of predicted PWY significantly enriched on DEGs per knockout assay and method.

Figure S8. GO semantic similarities observed between predicted GO terms and enriched GO terms in knockout are not occurring by chance.

Density plot displaying the distribution of random GO term semantic similarity. The observed value on real GO term enrichment, along with its corresponding P-value concerning the random distribution, is highlighted by the horizontal purple line.

Figure S9. Target genes and expression distribution of TFs compared with knockout results.

a. Histogram and density plot display the scaled (Z-score) number of target genes for each layer. TFs utilized in the knockout analysis are indicated by dotted red lines. **b**. Heatmap shows the presence or absence of target genes in each of the four layers for every TF analyzed in the knockout assays. **c**. Histogram and density plot of Tau index distribution for TF 2,910 TFs annotated with at least PWY/GO term. TFs utilized in the knockout analysis are indicated by dotted purple lines. **d**. Histogram and density plot illustrate the null distribution of Tau after randomly sampling 13 TFs a thousand times. TFs used in the knockout analysis are represented by dotted purple lines. P-values were calculated using the null distribution as a reference.

Figure S10. Evaluation of TF functional annotation by contrasting predictions random networks.

a, **b**, **c,** and **d**. Boxplot showing the fraction of random network with significantly enrich GO terms (**a**), average number of GO terms (**b**), -log10FDR (**c**), and GSS (**d**) observed in 3000 random networks by each method. The GSS values were calculated by comparing each random network with the observed GO terms from the true TF-target interactions. Asterisks indicate P-value significance (*: $p \leq 0.05$, **: 633 $p \leq 0.01$, ***: $p \leq 0.001$, ****: $p \leq 0.0001$, two-sided t-test).

Figure S11. GO term significance and similarity distributions from random networks per TF.

a and **b**. Density ridges plot showing the average -$\log_{10}$FDR (**a**) and GSS (**b**) distributions in 3,00 random networks for each TF. The GSS values were calculated by comparing each random network with the observed GO terms from the true TF-target interactions.

Figure S12. Scale count of GO terms in random networks.

Density plot displaying the distribution of GO terms in random networks predicted by the *network-based* method for each corresponding TF. The observed number of significantly enriched GO terms for the corresponding TF is indicated by a dotted orange line. The p-value was calculated from the random distribution as the null distribution.

Figure S13. Enrichment score for TF and GO term in several biological processes.

a, **b**, and **c**. Scatter plot with reciprocal Z score (rZ) of hormones- (**a**), metabolism- (**b**), and development-related (**c**) process.

Figure S14. Differences in GSEA enrichment per TF cluster.

a and **b** Boxplot with the percentage of co-expression networks (**a**) and the number of GO terms significantly enriched (FDR <= 0.1) per TF (**b**). **c**. Scatter plot indicating the total number of GO terms annotated by TF versus the percentage of GO terms co-expressed per TF. The color indicates the density of TFs overlapped. **d**. Boxplot indicating the average percentage of significantly co-expressed GO term. **e**. Boxplot with number of networks with at least a significant GO term by TF. On average, cluster 1 have TF significant associations with up to 29 different networks, while cluster 2 and 3 showed signals in 18.4 and 6.2 different networks out of the 51 total networks analyzed, respectively. **f**. Boxplot representing the number of GOs terms annotated per TF grouped in function of their GSEA enrichment similarities (Figure 4a).

Figure S15. GSEA enrichment profiles uncover TF-GO term regulatory effects which are condition-specific.

a, b, and **c.** Heatmap indicates the normalized enrichment score (NES) between bHLH43, HB33, and WRKY25 and their corresponding GO term by co-expression network.

SUPPLEMENTARY TABLES LEGENDS

Table S1. List of inbred lines and their corresponding tissues used in the eQTL analysis.

Table S2. Summary of ChIP- and DAP-seq publicly available reanalyzed in this work.

Table S3. Description of datasets used as part of the co-expression networks.

Table S4. List of expression datasets used in the identification of eQTLs.

Table S5. Categories of genes used to define types of *trans*-eQTLs.

Table S6. List of *trans*-eQTLs associations identified for HSF20.

Table S7. Total PDI (peaks) identified and used in the construction of the GRN.

Table S8. List of annotated functions (GO term and PWY) per TF and the method used on the corresponding annotation.

Table S9. List of DEG per experiments used to evaluate TF annotation predicted in this work.

AUTHOR CONTRIBUTIONS

F.G.-C. performed most of the analyses, assisted by E.G, N.-M.S, N.-d.L, and A.K. The research design was developed by F.G.-C, J.R., and P.Z. C.Y., E.M, and L. G.-C. contributed with analytic tools. F.G.-C and E.G. wrote the manuscript.

FUNDING

This research was supported by grants IOS-1733633 from the National Science Foundation to N.-d.L, N.-M.S., and E.G and Foundational Knowledge of Plant Products grant no. 2022-67013-36388 from the United States Department of Agriculture National Institute of Food and Agriculture to E.G.. L.G.-C. was supported in part by a fellowship from Michigan State University under the Training Program in Plant Biotechnology for Health and Sustainability (National Institutes of Health T32-GM110523). F.G.-C was supported in part by a fellowship from Michigan State University under the - (National Science Foundation DGE-1828149). J.R. was supported by a University of Wisconsin-Madison SciMed GRS Fellowship and United States Department of Agriculture.

REFERENCES

1. Lambert, S.A., Jolma, A., Campitelli,L.F., Das,P.K., Yin,Y., Albu,M., Chen,X., Taipale,J., Hughes,T.R. and Weirauch,M.T. (2018) The Human Transcription Factors. *Cell*, **175**, 598–599.

2. Schmitz,R.J., Grotewold,E. and Stam,M. (2022) Cis-regulatory sequences in plants: Their importance, discovery, and future challenges. *Plant Cell*, **34**, 718–741.

3. Arda,H.E. and Walhout,A.J.M. (2010) Gene-centered regulatory networks. *Brief. Funct. Genomics*, **9**, 4–12.

4. Mejia-Guerra,M.K., Pomeranz,M., Morohashi,K. and Grotewold,E. (2012) From plant gene regulatory grids to network dynamics. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*, **1819**, 454–465.

5. Yang,F., Ouma,W.Z., Li,W., Doseff,A.I. and Grotewold,E. (2016) Establishing the Architecture of Plant Gene Regulatory Networks. *Methods Enzymol.*, **576**, 251–304.

6. O'Malley,R.C., Huang,S.S.C., Song,L., Lewsey,M.G., Bartlett,A., Nery,J.R., Galli,M., Gallavotti,A. and Ecker,J.R. (2016) Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape. *Cell*, **165**, 1280–1292.

7. Deplancke,B., Mukhopadhyay,A., Ao,W., Elewa,A.M., Grove,C.A., Martinez,N.J., Sequerra,R., Doucette-Stamm,L., Reece-Hoyes,J.S., Hope,I.A., *et al.* (2006) A gene-centered C. elegans protein-DNA interaction network. *Cell*, **125**, 1193–1205.

8. Nakashima,K., Yamaguchi-Shinozaki,K. and Shinozaki,K. (2014) The transcriptional regulatory network in the drought response and its crosstalk in abiotic stress responses including drought, cold, and heat. *Front. Plant Sci.*, **5**, 1–7.

9. Birkenbihl,R.P., Liu,S. and Somssich,I.E. (2017) Transcriptional events defining plant immune responses. *Curr. Opin. Plant Biol.*, **38**, 1–9.

10. Sun,Y., Oh,D.-H., Duan,L., Ramachandran,P., Ramirez,A., Bartlett,A., Tran,K.-N., Wang,G., Dassanayake,M. and Dinneny,J.R. (2022) Divergence in the ABA gene regulatory network underlies differential growth control. *Nat Plants*, **8**, 549–560.

11. Marand,A.P., Eveland,A.L., Kaufmann,K. and Springer,N.M. (2023) cis-Regulatory Elements in Plant Development, Adaptation, and Evolution. *Annu. Rev. Plant Biol.*, **74**, 111–137.

12. Mack,K.L. and Nachman,M.W. (2017) Gene Regulation and Speciation. *Trends Genet.*, **33**, 68–80.

13. Bowles,A.M.C., Bechtold,U. and Paps,J. (2020) The Origin of Land Plants Is Rooted in Two Bursts of Genomic Novelty. *Curr. Biol.*, **30**, 530–536.e2.

14. Erenstein,O., Jaleta,M., Sonder,K., Mottaleb,K. and Prasanna,B.M. (2022) Global maize production, consumption and trade: trends and R&D implications. *Food Security*, **14**, 1295–1319.

15. Riedelsheimer,C., Lisec,J., Czedik-Eysenberg,A., Sulpice,R., Flis,A., Grieder,C., Altmann,T., Stitt,M., Willmitzer,L. and Melchinger,A.E. (2012) Genome-wide association mapping of leaf metabolic profiles for dissecting complex traits in maize. *Proc. Natl. Acad. Sci. U. S. A.*, **109**, 8872–8877.

16. Wen,W., Li,D., Li,X., Gao,Y., Li,W., Li,H., Liu,J., Liu,H., Chen,W., Luo,J., *et al.* (2014) Metabolome-based genome-wide association study of maize kernel leads to novel biochemical insights. *Nat. Commun.*, **5**, 3438.

17. Wen,W., Liu,H., Zhou,Y., Jin,M., Yang,N., Li,D., Luo,J., Xiao,Y., Pan,Q. and Tohge,T. (2016) Combining quantitative genetics approaches with regulatory network analysis to dissect the complex metabolism of the maize kernel. *Plant Physiol.*, **170**, 136–146.

18. Zhou,S., Kremling,K.A., Bandillo,N., Richter,A., Zhang,Y.K., Ahern,K.R., Artyukhin,A.B., Hui,J.X., Younkin,G.C., Schroeder,F.C., *et al.* (2019) Metabolome-Scale Genome-Wide Association Studies Reveal Chemical Diversity and Genetic Control of Maize Specialized Metabolites. *Plant Cell*, **31**, 937–955.

19. Schnable,P.S., Ware,D., Fulton,R.S., Stein,J.C., Wei,F., Pasternak,S., Liang,C., Zhang,J., Fulton,L., Graves,T.A., *et al.* (2009) The B73 maize genome: complexity, diversity, and dynamics. *Science*, **326**, 1112–1115.

20. Hufford,M.B., Seetharam,A.S., Woodhouse,M.R., Chougule,K.M., Ou,S., Liu,J., Ricci,W.A., Guo,T., Olson,A., Qiu,Y., *et al.* (2021) De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes. *Science*, **373**, 655–662.

21. McMullen,M.D., Kresovich,S., Villeda,H.S., Bradbury,P., Li,H., Sun,Q., Flint-Garcia,S., Thornsberry,J., Acharya,C., Bottoms,C., *et al.* (2009) Genetic properties of the maize nested association mapping population. *Science*, **325**, 737–740.

22. Kusmec,A., Srinivasan,S., Nettleton,D. and Schnable,P.S. (2017) Distinct genetic architectures for phenotype means and plasticities in Zea mays. *Nat Plants*, **3**, 715–723.

23. Xiao,Y., Liu,H., Wu,L., Warburton,M. and Yan,J. (2017) Genome-wide Association Studies in Maize: Praise and Stargaze. *Mol. Plant*, **10**, 359–374.

24. Mazaheri,M., Heckwolf,M., Vaillancourt,B., Gage,J.L., Burdo,B., Heckwolf,S., Barry,K., Lipzen,A., Ribeiro,C.B., Kono,T.J.Y., *et al.* (2019) Genome-wide association analysis of stalk biomass and

anatomical traits in maize. *BMC Plant Biol.*, **19**, 45.

25. Wei,F., Coe,E., Nelson,W., Bharti,A.K., Engler,F., Butler,E., Kim,H., Goicoechea,J.L., Chen,M., Lee,S., *et al.* (2007) Physical and genetic structure of the maize genome reflects its complex evolutionary history. *PLoS Genet.*, **3**, e123.

26. Kono,T.J.Y., Brohammer,A.B., McGaugh,S.E. and Hirsch,C.N. (2018) Tandem Duplicate Genes in Maize Are Abundant and Date to Two Distinct Periods of Time. *G3* , **8**, 3049–3058.

27. Schnable,J.C., Springer,N.M. and Freeling,M. (2011) Differentiation of the maize subgenomes by genome dominance and both ancient and ongoing gene loss. *Proc. Natl. Acad. Sci. U. S. A.*, **108**, 4069–4074.

28. Renny-Byfield,S., Rodgers-Melnick,E. and Ross-Ibarra,J. (2017) Gene Fractionation and Function in the Ancient Subgenomes of Maize. *Mol. Biol. Evol.*, **34**, 1825–1832.

29. Li,L., Briskine,R., Schaefer,R., Schnable,P.S., Myers,C.L., Flagel,L.E., Springer,N.M. and Muehlbauer,G.J. (2016) Co-expression network analysis of duplicate genes in maize (Zea mays L.) reveals no subgenome bias. *BMC Genomics*, **17**, 875.

30. Han,L., Zhong,W., Qian,J., Jin,M., Tian,P., Zhu,W., Zhang,H., Sun,Y., Feng,J.-W., Liu,X., *et al.* (2023) A multi-omics integrative network map of maize. *Nat. Genet.*, **55**, 144–153.

31. Tolani,P., Gupta,S., Yadav,K., Aggarwal,S. and Yadav,A.K. (2021) Big data, integrative omics and network biology. *Adv. Protein Chem. Struct. Biol.*, **127**, 127–160.

32. Depuydt,T., De Rybel,B. and Vandepoele,K. (2023) Charting plant gene functions in the multi-omics and single-cell era. *Trends Plant Sci.*, **28**, 283–296.

33. Galli,M., Khakhar,A., Lu,Z., Chen,Z., Sen,S., Joshi,T., Nemhauser,J.L., Schmitz,R.J. and Gallavotti,A. (2018) The DNA binding landscape of the maize AUXIN RESPONSE FACTOR family. *Nat. Commun.*, **9**, 4526.

34. Ricci,W.A., Lu,Z., Ji,L., Marand,A.P., Ethridge,C.L., Murphy,N.G., Noshay,J.M., Galli,M., Mejía-Guerra,M.K., Colomé-Tatché,M., *et al.* (2019) Widespread long-range cis-regulatory elements in the maize genome. *Nature Plants*, **5**, 1237–1249.

35. Tu,X., Mejía-Guerra,M.K., Valdes Franco,J.A., Tzeng,D., Chu,P.-Y., Shen,W., Wei,Y., Dai,X., Li,P., Buckler,E.S., *et al.* (2020) Reconstructing the maize leaf regulatory network using ChIP-seq data of 104 transcription factors. *Nat. Commun.*, **11**, 5089.

36. Rodgers-Melnick,E., Vera,D.L., Bass,H.W. and Buckler,E.S. (2016) Open chromatin reveals the functional maize genome. *Proc. Natl. Acad. Sci. U. S. A.*, **113**, E3177–84.

37. Marand,A.P., Chen,Z., Gallavotti,A. and Schmitz,R.J. (2021) A cis-regulatory atlas in maize at single-cell resolution. *Cell*, **184**, 3041–3055.e21.

38. Sekhon,R.S., Lin,H., Childs,K.L., Hansey,C.N., Robin Buell,C., de Leon,N. and Kaeppler,S.M. (2011) Genome-wide atlas of transcription during maize development. *Plant J.*, **66**, 553–563.

39. Stelpflug,S.C., Sekhon,R.S., Vaillancourt,B., Hirsch,C.N., Buell,C.R., de Leon,N. and Kaeppler,S.M. (2016) An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to Explore Root Development. *Plant Genome*, **9**.

40. Hoopes,G.M., Hamilton,J.P., Wood,J.C., Esteban,E., Pasha,A., Vaillancourt,B., Provart,N.J. and Buell,C.R. (2019) An updated gene atlas for maize reveals organ-specific and stress-induced genes. *Plant J.*, **97**, 1154–1167.

41. Zhou,P., Li,Z., Magnusson,E., Gomez Cano,F., Crisp,P.A., Noshay,J.M., Grotewold,E., Hirsch,C.N., Briggs,S.P. and Springer,N.M. (2020) Meta Gene Regulatory Networks in Maize Highlight Functionally Relevant Regulatory Interactions. *Plant Cell*, **32**, 1377–1396.

42. Li,H., Peng,Z., Yang,X., Wang,W., Fu,J., Wang,J., Han,Y., Chai,Y., Guo,T., Yang,N., *et al.* (2013) Genome-wide association study dissects the genetic architecture of oil biosynthesis in maize kernels. *Nat. Genet.*, **45**, 43–50.

43. Kremling,K.A.G., Chen,S.-Y., Su,M.-H., Lepak,N.K., Romay,M.C., Swarts,K.L., Lu,F., Lorant,A., Bradbury,P.J. and Buckler,E.S. (2018) Dysregulation of expression correlates with rare-allele burden and fitness loss in maize. *Nature*, **555**, 520–523.

44. Shrestha,V., Yobi,A., Slaten,M.L., Chan,Y.O., Holden,S., Gyawali,A., Flint-Garcia,S., Lipka,A.E. and Angelovici,R. (2022) Multiomics approach reveals a role of translational machinery in shaping maize kernel amino acid composition. *Plant Physiol.*, **188**, 111–133.

45. Liu,H., Wang,F., Xiao,Y., Tian,Z., Wen,W., Zhang,X., Chen,X., Liu,N., Li,W., Liu,L., *et al.* (2016) MODEM: multi-omics data envelopment and mining in maize. *Database* , **2016**.

46. Walley,J.W., Sartor,R.C., Shen,Z., Schmitz,R.J., Wu,K.J., Urich,M.A., Nery,J.R., Smith,L.G., Schnable,J.C., Ecker,J.R., *et al.* (2016) Integration of omic networks in a developmental atlas of maize. *Science*, **353**, 814–818.

47. Jin,M., Zhang,X., Zhao,M., Deng,M., Du,Y., Zhou,Y., Wang,S., Tohge,T., Fernie,A.R., Willmitzer,L., *et al.* (2017) Integrated genomics-based mapping reveals the genetics underlying maize flavonoid biosynthesis. *BMC Plant Biol.*, **17**, 17.

48. de Abreu E Lima,F., Li,K., Wen,W., Yan,J., Nikoloski,Z., Willmitzer,L. and Brotman,Y. (2018) Unraveling lipid metabolism in maize with time-resolved multi-omics data. *Plant J.*, **93**, 1102–1115.

49. Lee,T., Lee,S., Yang,S. and Lee,I. (2019) MaizeNet: a co-functional network for network-assisted systems genetics in Zea mays. *Plant J.*, **99**, 571–582.

50. Schaefer,R.J., Michno,J.-M., Jeffers,J., Hoekenga,O., Dilkes,B., Baxter,I. and Myers,C.L. (2018) Integrating Coexpression Networks with GWAS to Prioritize Causal Genes in Maize. *Plant Cell*, **30**, 2922.

51. Wen,W., Jin,M., Li,K., Liu,H., Xiao,Y., Zhao,M., Alseekh,S., Li,W., de Abreu E Lima,F., Brotman,Y., *et al.* (2018) An integrated multi-layered analysis of the metabolic networks of different tissues uncovers key genetic components of primary metabolism in maize. *Plant J.*, **93**, 1116–1128.

52. Yang,Z., Xu,G., Zhang,Q., Obata,T. and Yang,J. (2022) Genome-wide mediation analysis: an empirical study to connect phenotype with genotype via intermediate transcriptomic data in maize. *Genetics*, **221**.

53. Bukowski,R., Guo,X., Lu,Y., Zou,C., He,B., Rong,Z., Wang,B., Xu,D., Yang,B., Xie,C., *et al.* (2018) Construction of the third-generation Zea mays haplotype map. *Gigascience*, **7**, 1–12.

54. Jiao,Y., Peluso,P., Shi,J., Liang,T., Stitzer,M.C., Wang,B., Campbell,M.S., Stein,J.C., Wei,X.,

Chin,C.-S., *et al.* (2017) Improved maize reference genome with single-molecule technologies. *Nature*, **546**, 524–527.

55. Danecek,P., Bonfield,J.K., Liddle,J., Marshall,J., Ohan,V., Pollard,M.O., Whitwham,A., Keane,T., McCarthy,S.A., Davies,R.M., *et al.* (2021) Twelve years of SAMtools and BCFtools. *Gigascience*, **10**.

56. Schnable,J. (2019) Pan-Grass Syntenic Gene Set (sorghum referenced) with both maize v3 and maize v4 gene models. *figShare*, 10.6084/m9.figshare.7926674.v1.

57. Bolduc,N., Yilmaz,A., Mejia-Guerra,M.K., Morohashi,K., O'Connor,D., Grotewold,E. and Hake,S. (2012) Unraveling the KNOTTED1 regulatory network in maize meristems. *Genes Dev.*, **26**, 1685–1690.

58. Morohashi,K., Casas,M.I., Ferreyra,L.F., Mejia-Guerra,M.K., Pourcel,L., Yilmaz,A., Feller,A., Carvalho,B., Emiliani,J., Rodriguez,E., *et al.* (2012) A genome-wide regulatory framework identifies maize pericarp color1 controlled genes. *Plant Cell*, **24**, 2745–2764.

59. Eveland,A.L., Goldshmidt,A., Pautler,M., Morohashi,K., Liseron-Monfils,C., Lewis,M.W., Kumari,S., Hiraga,S., Yang,F., Unger-Wallace,E., *et al.* (2014) Regulatory modules controlling maize inflorescence architecture. *Genome Res.*, **24**, 431–443.

60. Pautler,M., Eveland,A.L., LaRue,T., Yang,F., Weeks,R., Lunde,C., Je,B.I., Meeley,R., Komatsu,M., Vollbrecht,E., *et al.* (2015) FASCIATED EAR4 encodes a bZIP transcription factor that regulates shoot meristem size in maize. *Plant Cell*, **27**, 104–120.

61. Li,C., Yue,Y., Chen,H., Qi,W. and Song,R. (2018) The ZmbZIP22 Transcription Factor Regulates 27-kD γ-Zein Gene Transcription during Maize Endosperm Development. *Plant Cell*, **30**, 2402–2424.

62. Zhan,J., Li,G., Ryu,C.H., Ma,C., Zhang,S., Lloyd,A., Hunter,B.G., Larkins,B.A., Drews,G.N., Wang,X., *et al.* (2018) Opaque-2 Regulates a Complex Gene Network Associated with Cell Differentiation and Storage Functions of Maize Endosperm. *Plant Cell*, **30**, 2425–2446.

63. Dong,Z., Xiao,Y., Govindarajulu,R., Feil,R., Siddoway,M.L., Nielsen,T., Lunn,J.E., Hawkins,J., Whipple,C. and Chuck,G. (2019) The regulatory landscape of a core maize domestication module controlling bud dormancy and growth repression. *Nat. Commun.*, **10**, 3810.

64. Dong,Z., Xu,Z., Xu,L., Galli,M., Gallavotti,A., Dooner,H.K. and Chuck,G. (2020) Necrotic upper tips1 mimics heat and drought stress and encodes a protoxylem-specific transcription factor in maize. *Proc. Natl. Acad. Sci. U. S. A.*, **117**, 20908–20919.

65. Gomez-Cano,F., Chu,Y.-H., Cruz-Gomez,M., Abdullah,H.M., Lee,Y.S., Schnell,D.J. and Grotewold,E. (2022) Exploring Camelina sativa lipid metabolism regulation by combining gene co-expression and DNA affinity purification analyses. *Plant J.*, 10.1111/tpj.15682.

66. Bolger,A.M., Lohse,M. and Usadel,B. (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, **30**, 2114–2120.

67. Langmead,B. and Salzberg,S.L. (2012) Fast gapped-read alignment with Bowtie 2. *Nat. Methods*, **9**, 357–359.

68. Li,H., Handsaker,B., Wysoker,A., Fennell,T., Ruan,J., Homer,N., Marth,G., Abecasis,G., Durbin,R. and 1000 Genome Project Data Processing Subgroup (2009) The Sequence Alignment/Map format

and SAMtools. *Bioinformatics*, **25**, 2078–2079.

69. Guo,Y., Mahony,S. and Gifford,D.K. (2012) High resolution genome wide binding event finding and motif discovery reveals transcription factor spatial binding constraints. *PLoS Comput. Biol.*, **8**, e1002638.

70. Liao,Y., Smyth,G.K. and Shi,W. (2019) The R package Rsubread is easier, faster, cheaper and better for alignment and quantification of RNA sequencing reads. *Nucleic Acids Res.*, **47**, e47–e47.

71. Andorf,C.M., Cannon,E.K., Portwood,J.L.,2nd, Gardiner,J.M., Harper,L.C., Schaeffer,M.L., Braun,B.L., Campbell,D.A., Vinnakota,A.G., Sribalusu,V.V., *et al.* (2016) MaizeGDB update: new tools, data and interface for the maize model organism database. *Nucleic Acids Res.*, **44**, D1195–201.

72. Wimalanathan,K., Friedberg,I., Andorf,C.M. and Lawrence-Dill,C.J. (2018) Maize GO Annotation-Methods, Evaluation, and Review (maize-GAMER). *Plant Direct*, **2**, e00052.

73. Yu,G. (2020) Gene Ontology Semantic Similarity Analysis Using GOSemSim. *Methods Mol. Biol.*, **2117**, 207–215.

74. Sayols,S. (2023) rrvgo: a Bioconductor package for interpreting lists of Gene Ontology terms. *MicroPubl Biol*, **2023**.

75. Liu,R. and Krishnan,A. (2021) PecanPy: a fast, efficient, and parallelized Python implementation of node2vec. *Bioinformatics*, **37**, 3377–3379.

76. Sales,G. and Romualdi,C. (2011) parmigene—a parallel R package for mutual information estimation and gene network reconstruction. *Bioinformatics*, **27**, 1876–1877.

77. Wisecaver,J.H., Borowsky,A.T., Tzin,V., Jander,G., Kliebenstein,D.J. and Rokas,A. (2017) A Global Coexpression Network Approach for Connecting Genes to Specialized Metabolic Pathways in Plants. *Plant Cell*, **29**, 944–959.

78. Ellison,E.L., Zhou,P., Hermanson,P., Chu,Y.-H., Read,A., Hirsch,C.N., Grotewold,E. and Springer,N.M. (2023) Mutator transposon insertions within maize genes often provide a novel outward reading promoter. *Genetics*, **225**.

79. Love,M.I., Huber,W. and Anders,S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.*, **15**, 550.

80. Korotkevich,G., Sukhov,V., Budin,N., Shpak,B., Artyomov,M.N. and Sergushichev,A. (2021) Fast gene set enrichment analysis. *bioRxiv*, 10.1101/060012.

81. Wright,E. (2016) Using DECIPHER v2.0 to analyze big biological sequence data in R. *R J.*, **8**, 352.

82. Emad,A. and Bailey,P. (2017) wCorr: weighted correlations.–R package ver. 1.9. 1.

83. Yilmaz,A., Nishiyama,M.Y., Garcia-Fuentes,B., Souza,G.M., Janies,D., Gray,J. and Grotewold,E. (2009) GRASSIUS: A platform for comparative regulatory genomics across the grasses. *Plant Physiol.*, **149**, 171–180.

84. Zheng,Y., Jiao,C., Sun,H., Rosli,H.G., Pombo,M.A., Zhang,P., Banf,M., Dai,X., Martin,G.B., Giovannoni,J.J., *et al.* (2016) iTAK: A Program for Genome-wide Prediction and Classification of Plant Transcription Factors, Transcriptional Regulators, and Protein Kinases. *Mol. Plant*, **9**, 1667–

1670.

85. Mathur,S., Vyas,S., Kapoor,S. and Tyagi,A.K. (2011) The Mediator complex in plants: structure, phylogeny, and expression profiling of representative genes in a dicot (Arabidopsis) and a monocot (rice) during reproduction and abiotic stress. *Plant Physiol.*, **157**, 1609–1627.

86. Zhou,P., Enders,T.A., Myers,Z.A., Magnusson,E. and Crisp,P.A. (2021) Applying cis-regulatory codes to predict conserved and variable heat and cold stress response in maize. *bioRxiv*.

87. Zhu,G., Wu,A., Xu,X.-J., Xiao,P.-P., Lu,L., Liu,J., Cao,Y., Chen,L., Wu,J. and Zhao,X.-M. (2016) PPIM: A Protein-Protein Interaction Database for Maize. *Plant Physiol.*, **170**, 618–626.

88. Subramanian,A., Tamayo,P., Mootha,V.K., Mukherjee,S., Ebert,B.L., Gillette,M.A., Paulovich,A., Pomeroy,S.L., Golub,T.R., Lander,E.S., *et al.* (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci. U. S. A.*, **102**, 15545–15550.

89. Kryuchkova-Mostacci,N. and Robinson-Rechavi,M. (2017) A benchmark of gene expression tissue-specificity metrics. *Brief. Bioinform.*, **18**, 205–214.

90. Yang,F., Li,W., Jiang,N., Yu,H., Morohashi,K., Ouma,W.Z., Morales-mantilla,D.E., Gomez-cano,F.A., Mukundi,E., Prada-salcedo,L.D., *et al.* (2017) A Maize Gene Regulatory Network for Phenolic Metabolism. *Mol. Plant*, **10**, 498–515.

91. Minow,M.A.A., Ávila,L.M., Turner,K., Ponzoni,E., Mascheretti,I., Dussault,F.M., Lukens,L., Rossi,V. and Colasanti,J. (2018) Distinct gene networks modulate floral induction of autonomous maize and photoperiod-dependent teosinte. *J. Exp. Bot.*, **69**, 2937–2952.

92. Colasanti,J., Yuan,Z. and Sundaresan,V. (1998) The indeterminate gene encodes a zinc finger protein and regulates a leaf-generated signal required for the transition to flowering in maize. *Cell*, **93**, 593–603.

93. Bartrina,I., Otto,E., Strnad,M., Werner,T. and Schmülling,T. (2011) Cytokinin regulates the activity of reproductive meristems, flower organ size, ovule formation, and thus seed yield in Arabidopsis thaliana. *Plant Cell*, **23**, 69–80.

94. Wahl,V., Ponnu,J., Schlereth,A., Arrivault,S., Langenecker,T., Franke,A., Feil,R., Lunn,J.E., Stitt,M. and Schmid,M. (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis thaliana. *Science*, **339**, 704–707.

95. Zhou,P., Enders,T.A., Myers,Z.A., Magnusson,E., Crisp,P.A., Noshay,J.M., Gomez-Cano,F., Liang,Z., Grotewold,E., Greenham,K., *et al.* (2022) Prediction of conserved and variable heat and cold stress response in maize using cis-regulatory information. *Plant Cell*, **34**, 514–534.

96. Maeda,H. and Dudareva,N. (2012) The shikimate pathway and aromatic amino Acid biosynthesis in plants. *Annu. Rev. Plant Biol.*, **63**, 73–105.

97. Perez de Souza,L., Garbowicz,K., Brotman,Y., Tohge,T. and Fernie,A.R. (2020) The Acetate Pathway Supports Flavonoid and Lipid Biosynthesis in Arabidopsis. *Plant Physiol.*, **182**, 857–869.

98. Cocco,S., Feinauer,C., Figliuzzi,M., Monasson,R. and Weigt,M. (2018) Inverse statistical physics of protein sequences: a key issues review. *Rep. Prog. Phys.*, **81**, 032601.

99. Facco,E., Pagnani,A., Russo,E.T. and Laio,A. (2019) The intrinsic dimension of protein sequence evolution. *PLoS Comput. Biol.*, **15**, e1006767.

100. Zheng,X., Liu,T., Yang,Z. and Wang,J. (2011) Large cliques in Arabidopsis gene coexpression network and motif discovery. *J. Plant Physiol.*, **168**, 611–618.

101. Ruan,J., Perez,J., Hernandez,B., Lei,C., Sunter,G. and Sponsel,V.M. (2011) Systematic identification of functional modules and cis-regulatory elements in Arabidopsis thaliana. *BMC Bioinformatics*, **12 Suppl 12**, S2.

102. Kulkarni,S.R. and Vandepoele,K. (2019) Inference of plant gene regulatory networks using data-driven methods: A practical overview. *Biochim. Biophys. Acta Gene Regul. Mech.*

103. Ledezma-Tejeida,D., Altamirano-Pacheco,L., Fajardo,V. and Collado-Vides,J. (2019) Limits to a classic paradigm: most transcription factors in E. coli regulate genes involved in multiple biological processes. *Nucleic Acids Res.*, **47**, 6656–6667.

104. Tang,M., Li,B., Zhou,X., Bolt,T., Li,J.J. and Cruz,N. (2021) A genome-scale TF–DNA interaction network of transcriptional regulation of Arabidopsis primary and specialized metabolism. *Mol. Syst. Biol.*

105. Reményi,A., Schöler,H.R. and Wilmanns,M. (2004) Combinatorial control of gene expression. *Nat. Struct. Mol. Biol.*, **11**, 812.

106. Brkljacic,J. and Grotewold,E. (2017) Combinatorial control of plant gene expression. *Biochim. Biophys. Acta*, **1860**, 31–40.

107. Alon,U. (2007) Network motifs: theory and experimental approaches. *Nat. Rev. Genet.*, **8**, 450–461.

108. Sonawane,A.R., Platig,J., Fagny,M., Chen,C.-Y., Paulson,J.N., Lopes-Ramos,C.M., DeMeo,D.L., Quackenbush,J., Glass,K. and Kuijjer,M.L. (2017) Understanding Tissue-Specific Gene Regulation. *Cell Rep.*, **21**, 1077–1088.

109. Gomez-Cano,F., Chu,Y.H. and Cruz-Gomez,M. (2022) Exploring Camelina sativa lipid metabolism regulation by combining gene co-expression and DNA affinity purification analyses. *The Plant.*

110. Cancé,C., Martin-Arevalillo,R., Boubekeur,K. and Dumas,R. (2022) Auxin response factors are keys to the many auxin doors. *New Phytol.*, **235**, 402–419.

111. Zhao,Y. (2010) Auxin Biosynthesis and Its Role in Plant Development. *Annu. Rev. Plant Biol.*, **61**, 49–64.

112. Vanstraelen,M. and Benkova,E. (2012) Hormonal interactions in the regulation of plant development. *Annu. Rev. Cell Dev. Biol.*, **28**, 463–487.

113. Derksen,H., Rampitsch,C. and Daayf,F. (2013) Signaling cross-talk in plant disease resistance. *Plant Sci.*, **207**, 79–87.

114. Munné-Bosch,S. and Müller,M. (2013) Hormonal cross-talk in plant development and stress responses. *Front. Plant Sci.*, **4**, 529.

115. Pacifici,E., Polverari,L. and Sabatini,S. (2015) Plant hormone cross-talk: the pivot of root growth. *J. Exp. Bot.*, **66**, 1113–1121.

116. Sawicki,M., Aït Barka,E., Clément,C., Vaillant-Gaveau,N. and Jacquard,C. (2015) Cross-talk between environmental stresses and plant metabolism during reproductive organ abscission. *J. Exp. Bot.*, **66**, 1707–1719.