Both environmental conditions and intra- and interspecific interactions

influence the movements of a marine predator

Johann Mourier[1], Marc Soria[1], Matthew Silk[2,3,4], Angélique Demichelis[1], Laurent Dagorn[1],

Tarek Hattab[1]

[1] MARBEC, Univ Montpellier, CNRS, IFREMER, IRD, Sète, France

[2] Centre for Ecology and Conservation, University of Exeter Penryn Campus, Cornwall, UK

[3] NIMBioS, The University of Tennessee Knoxville, Knoxville, TN, USA

[4] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

Corresponding author: Johann Mourier

email: johann.mourier@umontpellier.fr

Address: MARBEC, Station Ifremer, Avenue Jean Monnet, BP 171, 34203 Sète Cedex -

France.

Telephone: +33 6 72 71 21 57

ABSTRACT

Animal movements are typically influenced by multiple environmental factors

simultaneously and individuals vary in their response to this environmental heterogeneity.

Therefore, understanding how environmental aspects, including biotic, abiotic and

anthropogenic factors, influence the movements of wild animals is an important focus of

wildlife research and conservation. We apply exponential random graph models (ERGMs) to

analyse movement networks of a bull shark population in a network of acoustic receivers and

identify the effects of environmental, social or other types of covariates on their movements.

We found that intra- and interspecific factors often had stronger effects on movements than

environmental variables. ERGMs proved to be a potentially useful tool for studying animal

movement network data especially in the context of spatial attribute heterogeneity.

Keywords: spatial-social interface, acoustic telemetry; *Carcharhinus leucas*; ERGM;

movement networks; Indian Ocean; Reunion Island; shark.

Understanding animal movement is not straightforward, as movement decisions result

from a combination of many factors including the internal state of the individual, its intra-

and interspecific neighbourhood, and spatio-temporal variation in abiotic environmental

conditions (Nathan et al., 2008). Additionally, the response of individuals to variation in their

surrounding environment can differ across spatial and temporal scales. Daily activities can be

influenced by environmental factors that vary over a diel cycle (e.g., wind or tidal currents),

while seasonal movement can be governed by changes acting at larger temporal scales (e.g.,

photoperiod, temperature). Quantifying how heterogeneity in an animal's surroundings drives

behaviour, and consequently movement patterns, provides insight into how natural and

anthropogenic changes may impact populations and ecosystems.

One potential way to understand the 'causes, mechanisms and spatiotemporal patterns

of movement and their role in various ecological and evolutionary processes' (Nathan et al.

2008) is to link observed movement patterns with spatial and temporal variability in the

underlying environmental conditions (Avgar et al., 2013). Movement patterns result from

interactions between organisms and their environments (Johnson et al., 1992; Morales et al.,

2010; Schick et al., 2008). For example, movement rates are expected to decrease where

resources are rich and decrease where food availability is low (Kuefler et al., 2012; Pyke et

al., 1977). Understanding movement patterns thus requires consideration of the temporally

dynamic nature of these environments (Avgar et al., 2013; Couriot et al., 2018; Mueller et al.,

2011; Riotte-Lambert & Matthiopoulos, 2020; Schick et al., 2008).

In addition to interacting with their physical environment (e.g. habitat features

offering essential resources, conditions or shelter), animals also interact with other

individuals of the same or different species. These spatial encounters with conspecific or

heterospecific individuals may be advantageous or unfavourable. For example, aggregations

form to avoid predation or to forage efficiently (Krause & Ruxton, 2002), although the

tendency for non-gregarious species to seek or avoid each other is less well known. For all

sexual species, individuals need to encounter conspecifics at least for mating, but avoiding

opposite-sex individuals (sexual segregation) may play a major role in preventing sexual

harassment in some species (Wearmouth & Sims, 2008). Additionally, avoidance strategies

can act at the intra or interspecific level in the context of competition for mates, space or

resources (Giuggioli & Kenkre, 2014), or in the context of predator–prey interactions

(Atwood et al., 2009). Because individuals may select habitats based on exogenous

environmental features (e.g. habitat quality and predation risk), and local population factors

(e.g., mating opportunities, competition or density dependence), we need to quantify the

mechanisms by which dynamic interactions between individuals occur (e.g. movement

towards or away from other individuals).

Therefore, social behaviour is intrinsically connected with spatial behaviour and a

burgeoning literature now addresses the link between spatial and social aspects of behaviour

defined as the "spatial-social interface" (Webber et al., 2023). Indeed, while spatial and social

processes have been predominantly considered independently, because movement behaviour

emerges from social and spatial processes, animal movement studies offer an opportunity to

consider and integrate them (Albery et al., 2021; He et al., 2019; Mourier et al., 2019;

Webber et al., 2023).

Finally, the human footprint has altered the spatial ecology of many species at

different spatial and temporal scales, for example by decreasing animal movements as a

result of behavioural changes, habitat fragmentation and barrier effects (Tucker et al., 2018),

or by modifying activity-timing (Gilbert et al., 2023). COVID-19 lockdowns provided an

empirical experiment where an abrupt reduction in human activity (so called Anthropause)

led to decreases in animal movement rates and avoidance patterns of human footprint (Tucker

et al., 2023). Human disturbance can also fundamentally alter the way that species interact,

such as by causing a spatiotemporal compression of species co-occurrences in disturbed

landscape which can lead to increases in competition, predation and infectious disease

transmission (Gilbert et al., 2022). As such, incorporating human footprint as a potential

factor affecting animal movements at the spatial-social interface is now more relevant than

ever.

In this study, we explore the connection between seascape attributes and animal

movement patterns for a population of bull shark (*Carcharhinus leucas*) that inhabit a coastal

zone in Reunion Island (Indian Ocean), which encompasses a variety of environmental

conditions. We evaluate the relative importance of spatiotemporal variables associated with

several main aspects of the seascape (abiotic conditions, social environment, predation risk or

interspecific competition and anthropogenic pressure) as potential drivers of shark movement

patterns. This approach allows us to identify fundamental relationships between local

environmental conditions and animal movement patterns.

To do so, we used a network-based modelling that account for the non-independence

of the data of movements between location. We applied exponential random graph models

(ERGMs; Lusher, Koskinen, & Robins, 2012; Robins, Pattison, Kalish, & Lusher, 2007) to

monthly individual movement networks of bull sharks between listening stations (i.e.

acoustic receivers) deployed along the west coast of Reunion Island (Mourier et al., 2021;

Soria et al., 2019), using varying covariate attributes at these locations. ERGMs are statistical

models of networks that treat the weight of network edges (i.e. number of movements

between two nodes) as the response variable and network node (i.e. acoustic receivers)

attributes as explanatory variables (Robins, Pattison, et al., 2007; Robins, Snijders, et al.,

2007; Snijders et al., 2006). These models account for the non-independence of the data

expressed as movements between sites acting as sources or sinks (Silk & Fisher, 2017).

Bull shark movements are known to be influenced by both biotic and abiotic factors

(Lee et al., 2019; Lubitz et al., 2023; Niella et al., 2022; Werry et al., 2018), and

anthropogenic factors (Hammerschlag et al., 2022; Werry et al., 2012), but less is known

about the importance of con- and heterospecifics in movement decisions. Even if most

studies on the movements of these marine predators have identified environmental factors as

most influential to their movement ecology, we expect that social and competition factors

may be just as important in explaining individual movements. We incorporated swell height,

turbidity and rainfall levels as abiotic predictive variables of movements, turtle densities as a

potential indicator of the presence of sharks as suggested by local people, cumulative number

of sea users as an anthropogenic variable, as well as various intra- and interspecific factors

such as residency times of the opposite sex, abundance of bull sharks and occupancy of tiger

sharks (a larger and likely competitor species). Our analysis was designed to test whether

individual bull sharks move according to the distribution of conspecifics and heterospecifics

in the seascape, and whether these effects were comparable with expected positive effects of

swell height, turbidity and rainfall on shark movements and negative effects of human

densities (with sharks avoiding human presence). While segregation, resource partitioning

and competition processes may all shape bull shark movements, we had no clear directional

predictions of their effect due to the limited previous research on this population on these

aspects.

METHODS

Study species and local context

The bull shark is a large Carcharhinidae with a wide cosmopolitan distribution along the

continental coasts of all tropical and sub-tropical waters of the world and is known to be

mobile and move across a large range of habitats and environmental conditions

(Brunnschweiler et al., 2010; Daly et al., 2014; Espinoza et al., 2016; Heupel et al., 2015; Lee

et al., 2019; Niella et al., 2022). Additionally, it is also an euryhaline elasmobranch that uses

a range of salinities throughout its life-cycle (Niella et al., 2022). Its behaviour varies across

spatial and temporal scales, as well as according to size and sex, with a high individual

variability in the tendency to move (Espinoza et al., 2016; Lee et al., 2019; Mourier, Soria, et

al., 2021). However, much remains unclear about adult bull shark movement decisions.

Reunion Island (21°07'S / 55°32'E) is a volcanic island located 700 km east of Madagascar

in the southwest Indian Ocean. The island is 2512 km^2 with 217 km of coastline and

characterized by steep underwater slopes (*ca.* 10-20%) to a depth of 2,000 m. Fringing reefs

stretch over 25 km along the west and south-west coast (Fig. 1) forming a natural coral

barrier that bounds the reef flats and back-reef depressions and lies no further than 500 m

from the beach.

Figure 1: Cumulative number of movements across all 44 monthly individual movements
used in the ERGM analyses concentrated on the western part of Reunion Island. Circles
represent acoustic receivers. Inset map represents the location of Reunion Island and the area
where the movements were recorded.

Acoustic telemetry and movement network

Sharks were captured along the west coast of Reunion Island between September 2012 and
March 2013 (Supplementary material 1 Table S1), using horizontal drifting long-lines 0.2 to
1 km in length and equipped with 20 to 200 baited 16/0 circle hooks (Blaison et al., 2015).
Set times were fixed at a maximum of 3 h to minimize shark and bycatch mortality. Once
captured, a shark was brought alongside the vessel and held still by rubber-encased ropes to
prevent skin lesions and burns and rolled onto its back to induce tonic immobility. The boat

moved forward slowly throughout the procedure to allow the shark to breathe. Sanitized

transmitters were implanted into the peritoneal cavity through a midventral incision using a

sterile scalpel. A 1cm incision was made, then enlarged with retractors. The muscle was then

gently torn in the direction of the fibers to allow faster muscle tissue repair. Once the tag was

in place, two stitches were done to close the incision and facilitate healing. A sterile, non-

absorbable synthetic monofilament suture (polyamide) was used. Absorbable sutures

disintegrate too quickly. Braided sutures were more likely to increase the risk of infection

allowing seawater to enter in the tissue. The shark was then freed by cutting the hook at its

base and removing it, untying the lasso, replacing it on its belly and oxygenating it by

moving it back and forth or slowly forward until the first signs of autonomous movement

appear. Sex and total length (TL) were recorded and transmitters (Vemco V16, transmission

interval 40–80 s, estimated battery life 845 days) were implanted into the peritoneal cavity

through a midventral incision.

An array of 46 Vemco VR2W acoustic receivers was deployed along the coast with receivers

installed an average of approximately 2 km apart at depths of 10–60 m, comprising 33 (71%)

offshore receivers and 13 (29%) inshore receivers placed less than 300 m from shore (Fig. 1).

For each shark visit at a receiver, we used detection records to calculate a continuous

residency time (CRT) corresponding to the duration within which a tagged shark was

continuously monitored at a specific receiver without one hour-scale (> 1h) absences

(Capello et al., 2015; Ohta & Kakuma, 2005). All detections of the same shark at one

receiver separated by less than a predefined period, called the maximum blanking period

(1 h), were grouped into one CRT and defined as a visit. Each time a tagged shark was

detected at a different receiver, a new visit started - ending the visit at the previous receiver -

even if the interval between detections was less than the maximum blanking period.

We then built monthly movement networks of each shark with each node representing a

receiver along the coast of Reunion Island and each weighted, directed edge represented the

number of movements of the individual (deduced from CRT) from node A to node B (Fig. 1)

within a given month.

Ethical Note

All the fieldwork and protocols of handling and tagging of sharks were approved by the

Ethics Committee (n° 114) for the CYROI (Cyclotron Réunion Océan Indien). Procedures

were adapted to minimize stress on animals and avoid mortality. The use of circular hooks

prevented the shark from swallowing them. All sharks swam away in good condition after

being released. The procedure usually lasted less than 30 min. Most sharks were detected and

therefore remained alive during the experiment (mean days of detections 88.3 ± 79.4; min =

1, max = 285). Two sharks were caught and removed by local fishermen and two others were

never detected after being released. All operations were carried out or supervised by

scientists with a certificate in animal experimentation and a certificate in experimental

surgery (Oniris, Ecole Nationale Vétérinaire de Nantes).

Explanatory variables

We gathered data of different nature (i.e. biological, abiotic, anthropogenic and spatial) that

were accessible in the study area. Although non exhaustive, explanatory variables recorded

included a number of environmental, biological and anthropogenic factors (Table 1; a

detailed description of how they were recorded can be found in Supplementary material 1) as

well as the geographical distances between receivers. As all data were not available at the

same spatial and temporal scale, we chose to standardize all available candidate explanatory

data at the scale of one month for each receiver. This choice of resolution of the data is a

compromise to keep a valuable temporal scale of the variability of site conditions and in the

same time having enough movements to construct individual networks that represent

biologically meaningful variations. For some environmental variables such as rainfall,

multiple nearby receivers shared values as data were available for a specific zone including

several receivers. All explanatory variables were standardized to have a mean of 0 and a

standard deviation of 1.

Table 1: Predictor parameters included in the ERGMs.

Variables	Category	Description	Range	Source
Occupancy Tiger sharks	Biological	Frequency of days at least a tiger shark was detected at the receiver for each month (%)	[0-0.409]	Current acoustic telemetry data
Same sex CRT	Biological	Total CRT of bull sharks of the same sex for each month (hour)	[0.033-118.425]	Current acoustic telemetry data
Opposite sex CRT	Biological	Total CRT of bull sharks of the opposite sex for each month (hour)	[0.033-118.425]	Current acoustic telemetry data
Bull Shark abundance	Biological	Number of bull sharks present at the listening station divided by the number of bull shark with an active tag for each month	[0-1.24]	Current acoustic telemetry data
Turtle density	Biological	Mean density of turtles in the area of the receiver from aerial survey (nb/km^2)	[0-15.873]	Aerial surveys
Turbidity	Environmental	Average monthly reflectance for each zone over the multiple images (%)	[4.101-8.886]	Satellite SPOT 4 & 5
Rain	Environmental	Mean rain fall for each watershed for each month (mm)	[0.008-25.460]	METEO-France
Swell	Environmental	Height of the swell near the listening station for each month (m)	[0.557-2.636]	AVISO portal, CANDHIS & METEOLAMER platform
Human activities	Anthropogenic	Cumulated number of human activities (boats, swimmers, nautical activities) for each month	[0-1276.5]	Aerial surveys
Geodist	Spatial	Distance between stations (km)	[0.9-120]	Current acoustic telemetry data

Exponential Random Graph Models

As care is required when conducting statistical analysis of network data because of issues

related to potential non-independence among neighbouring nodes and edges (Croft et al.,

2011), we used Exponential Random Graph Models (ERGMs). ERGMs are statistical models

of networks that treat the weight of network edges as the response variable and network node

and edge attributes as explanatory variables (Robins, Pattison, et al., 2007; Robins, Snijders,

et al., 2007; Snijders et al., 2006). Such models are analogical to Generalized Linear Models

(GLMs) except that they enable hypothesis testing about the processes driving network

structure and link formation. These models have been mostly applied in social science, but

their properties also make them useful for answering questions related to how and why

animals move between locations in the context of movement networks (Fletcher et al., 2011;

Jacoby & Freeman, 2016; López-Calderón et al., 2023). Effectively the movement network

becomes the response in a regression model, where the predictors are the propensity for

nodes (i.e. locations) of similar or dissimilar attributes to be linked by movements.

The general form for an ERGM can be written as:

$$P(Y = y; \theta, x) = \frac{\exp(\theta^T g(y,x))}{\kappa(\theta,x)} \qquad (1)$$

where:

- Y is the random variable for the state of the network (with realization y),

- $g(y,x)$ is a vector of model statistics for network y,

- θ is the vector of coefficients for those statistics, and

- $\kappa(\theta)$ is a normalizing term which ensures that equation (1) is a proper probability

distribution. It represents the quantity in the numerator summed over all possible

networks (typically constrained to be all networks with the same node set as y).

The numerator represents a formula that is linear in the log form:

$$\log\left(\exp\left(\theta' g(y)\right)\right) = \theta_1 g_1(y) + \theta_2 g_2(y) + \cdots + \theta_p g_p(y)$$

where p is the number of terms in the model. From this one can more easily observe the

analogy to a traditional statistical model. The functions $g(y)$ are counts of configurations in

the network y and the parameters θ weight the relative importance of the respective

configurations, effectively the size and direction of the effects of the covariates. Parameter

estimation in most specifications of ERGMs uses maximum pseudo-likelihood, an

approximation of maximum likelihood based on Monte Carlo estimation.

All ERGMs were fitted using R packages *ergm* and *ergm.count* (Hunter et al., 2008).

Model-fitting process

Node attributes were varying at the month scale so we built a model for each monthly-

individual network. To ensure that it was possible to fit ERGMs to monthly networks and

improve model convergence, we first removed from monthly movement networks all

receivers that were deployed less than 20 days during the month, as well as all individuals

that did not present an active tag for at least 20 days during a month. This empirical choice

was made as a compromise between data loss and data representativity. We also retained only

monthly individual networks containing at least 10 different directed movements between

receivers, as networks including low numbers of movements caused problems with model

convergence and parameter estimation. We also excluded two monthly networks that were

binary (i.e. only contained at most single movements along any particular edge). This left us

with 42 monthly networks of 13 individuals (9 females: 4 males; mean/median/range of

monthly networks per individual = 3.1/3/1-6).

For each monthly network for each shark, we then fitted two alternative versions of the full

model that each modelled different effects of the environmental variables being studied

(Table 1). The effect estimated for each factor represents the difference from the intercept as

with classical GLMs. All models included the term "non-zero" to control for zero-inflation

generated by the weighted movement networks being sparse and the term "sum",

corresponding to the sum of all link values, was included as the equivalent to the intercept in

a linear model (Dey & Quinn, 2014).

The two model sets differed as follows:

• Model set 1: we tested the association between the current state of the environment at

each receiver location and number of movements towards and away from it (i.e. effect

of an attribute on out- and in-degree). These models used the *nodecov* term in the

fitted ERGMs.

• Model set 2: we tested the association between the absolute difference in current state

of the environment at dyads (pairs) of receiver locations and number of movements

between them. These models used the *absdiff* term in the fitted ERGMs.

We used a *nodecov* model rather than separate *nodeicov* and *nodeocov* models (movements

towards and away from receivers respectively) as simulations indicated the results were

qualitatively identical (see Supplementary material 2).

We fitted the *nodecov* and *absdiff* models separately to facilitate model parsimony and

convergence and test the robustness of the movement patterns discovered while controlling

for the full suite of environmental variables. Indeed, incorporating all variables and models

sets combinations would prevent convergence of models and render interpretation of output

complex. To account for the effect of the spatial distribution of nodes in our models, we

included as a covariate the matrix of distances between nodes.

For each model set, we then wrote a model formula including all potential predictors as

follows:

graph~nonzero+sum+term(Variable 1)+term(Variable 2)+… +term(Variable n)+

edgecov(Geodist,'Geodist')

where *term* can be replaced by *nodecov* and *absdiff* in model sets 1 and 2 respectively. An

edgecov term was added to control for the distance between locations. Models were fitted

with a Poisson reference distribution for edge weights creating the familiar log-linear effect.

The final coefficient estimates indicate log-linear increase in the weight variable

(movements). MCMLE.Hummel.maxit was set to 1000 and MPLE.type was set to

"penalized" so that the maximum penalized likelihood estimate was calculated using a bias-

reduced method.

For each model set for each individual monthly network, we used Aikaike Information

Criterion (AIC) to compare the AIC of 49 candidate models nested within the full model and

keep only the model that most parsimoniously explained the movements of the individual

during the month considered. Any models with convergence issues were discarded.

We then re-fitted all selected ERGMs for each combination of both model sets (*nodecov* and

absdiff) and individual monthly networks. From these final models we extracted model

estimates and their standard errors, and also MCMC diagnostics (Gewecke's convergence

diagnostic and $\hat{R}$) to ensure convergence.

To summarize the results incorporating each selected models, we extracted parameter

estimates and associated standard errors from selected variables from each selected models,

and followed a meta-analysis procedure in which each sample (monthly-individual

movement network) was treated as a single 'Study'. Effect sizes were calculated using and

inverse-variance weighting meta-analysis for each sample and grouped by variables using a

sub-grouping analysis. This allows to test if differences in effect sizes exist only due to

sampling error, or because of true differences in the effect sizes. This procedure was made for

the overall samples as well as separating by sex and by season. The meta-analysis was

conducted using the package 'meta' in R (Schwarzer, 2007).

 Simulation methods

To ensure the correct interpretation of our model results and check for potential limitations of

the ERGM methods applied to our dataset we used a simulation approach. In short, we

simulated a series of individual movement trajectories that matched our empirically collected

data. We then fitted ERGMs in the same way we did for the empirical data.

We generated sets of 25 receiver locations (equivalent to the empirical data) positioned in 2D

space and generated three environmental properties for each receiver location termed

factorA, factorB and factorC. We then generated 21 simulation input parameter sets that

varied the effect of factorA on the probability of movements between locations while keeping

factorB and factorC fixed as having no effects on movement (Supplementary material 2

Table S2). We considered scenarios where movements were conducted according to a

gradient in factorA or occurred between similarly high or low factorA locations. For each

simulation input parameter set we simulated the movements of 10 sharks. The number of

movements for each shark was drawn from a Poisson distribution with a mean of 23

(equivalent to the empirical dataset). Detailed methods are in Supplementary materials 2.

RESULTS

Interpretations of model outputs from simulations

Our simulation study revealed ERGMs performed as expected when analysing movement

networks of a similar size and structure to our empirically-measured networks (see

Supplementary Materials 2). Model estimates for the *nodecov* and *absdiff* models typically

accurately represented the presence of movements up a gradient or between similar locations

respectively (Fig. 2). The statistical power to detect true effects was limited for individual

networks, and there were slightly elevated false positive rates for the *absdiff* models.

However, effect size estimates were largely unbiased (both with and without true simulated

effects), with the only exceptions being that *absdiff* models underestimated the tendency to

move between similar locations when there was also a gradient effect, and that *absdiff* models

overestimated the tendency to move between similar locations when sharks tended to start

moving from high value locations and had a strong tendency to move up a gradient (although

the latter would be expected if this resulted in sharks moving only between high value

locations). Collectively these results indicate that collating estimates from multiple models

(as we do in our main analysis) will provide the most informative results. Simulations also

indicated that *nodeicov* and *nodeocov* models provided closely correlated results leading to us

using a single *nodecov* model in our empirical analysis (see above).

Figure 2: Interpretation of the model output for each scenario based on simulations: (a, b, c)

movements up a gradient of Factor A (e.g. swell height), (d, e, f) movements between

locations of low Factor A and (g, h, i) movement up a gradient and between locations of high

Factor A. (a, d, g) Toy examples on the left and movement network for the first shark. (b, e,

h) model estimates for *nodecov* and (c, f, i) model estimates for *absdiff*. The model estimates

are for the three effects (highlighting the one where there is a true effect in red) and then for

each of the 10 sharks simulated.

Effects of abiotic conditions

Figure 3: Summary of the outputs from the ERGM models from the 42 selected individual

monthly movement networks. Effect sizes and their 95% confidence intervals pooled over

variables are reported for the parameter estimate of terms *nodecov* and *absdiff*. Variables

were grouped in several categories: (1) environmental factors, (2) turtle density, (3)

anthropogenic factors, (4) social or intraspecific interactions, and (5) competitive or

interspecific interactions. The variable "Geodistance" controls for the distance between

receivers. Circle size is proportional to the number of models in which the coefficient of the

term was significant and therefore selected.

Our models were designed to test whether sharks were moving between locations with

similar values (tested by the *absdiff* term) and/or towards locations with higher/lower values

(tested by the *nodecov* term) of an abiotic condition, specifically swell height, degree of

turbidity and level of rainfall. Model results indicated the environmental variables that were

associated with movement network structure (Fig. 3). While high swell at a location was not

significantly associated with more movements, movements between locations that differed

considerably in swell height were much less frequent than movements between locations with

similar swells as demonstrated by negative values of the *absdiff* model. Together, these

outputs indicate that sharks move between locations characterized by similar swell and not

across a gradient of swell heights. The *nodecov* model revealed a statistically negative effect

of turbidity on movements, indicating that more movements occurred through low turbidity

locations. Further, the *absdiff* model revealed that sharks were also more likely to move

between receiver locations that differed from each other in their turbidity. Collectively these

results suggest that sharks moved down turbidity gradients and then more frequently among

locations with lower turbidity. Sharks also tended to move between receiver locations with

similar and low rainfall levels.

Effect of turtle densities

Our models could also test whether sharks were moving toward high or low densities of

turtles (tested by the *nodecov* term) and/or were remaining in habitats with similar densities

(tested by the *absdiff* term). Model outputs indicated that movements were more likely

between receiver locations with more similar turtle densities (negative estimate from *absdiff*

model). However, parameter estimates were small indicating that the biological importance

of turtle distribution densities may be limited.

Effect of anthropogenic factors

Our models were also set up to test whether sharks were avoiding (tested by the *nodecov* term

with an expected negative coefficient) and/or remaining in areas of similar human activity

(tested by the *absdiff* term with an expected negative coefficient). Models indicated that there

were more movements among locations with high human activities (positive *nodecov* effect

and negative *absdiff* effect). However, parameter estimates from the models were close to

zero suggesting that the biological importance of anthropogenic factors such as human

activities on bull shark movements in the monitored area were relatively limited.

Influence of conspecifics

Our models were also used to test whether sharks moved towards or avoided conspecifics

(tested by the *nodecov* term) and/or remained within areas with similar populations of

conspecifics (tested by the *absdiff* term). Receiver locations in which bull sharks were present

or more abundant were more strongly connected in the movement network. Outputs from

absdiff model also indicated that movements tended to occur between locations with similar

bull shark abundance, suggesting individuals were predominantly moving among a subset of

preferred locations. While controlling for the abundance effect, there was (overall) some

evidence for a negative effect of the residency time of sharks of the opposite sex, suggesting

some spatial segregation between the sexes, but this effect varied seasonally (see below).

Effect of inter-specific competition

We also used our models to test whether the presence of larger tiger sharks was influencing

bull shark movements by moving to/away from areas used by tiger sharks (tested by the

nodecov term) and/or moving between sites with similar numbers of tiger sharks (tested by

the *absdiff* term). There was weak evidence (due to wide confidence intervals and the small

number of times this parameter was selected in the top model) for the presence of tiger sharks

affecting bull shark movement network structure. Positive effects in *absdiff* model indicated

individuals tended to move across gradients of tiger shark occupancy, perhaps indicating

active avoidance of tiger sharks, with some additional weak evidence of reduced movements

through locations with higher occupancy of tiger sharks.

Sex differences in movement network structure

Analysing movement network structure for the two sexes independently revealed similar

broad trends as the overall movement network, except in the case of social factors and the

tiger shark occupancy (Fig. 4). While females tended to move toward locations with high

residency times for other females, males showed a general pattern of avoiding locations with

high residency times of either sex (while controlling for overall bull shark abundance). The

weak overall evidence for reduced movements through locations with high tiger shark

occupancy was driven by divergent effects between females and males. Female movements

were directed towards locations with low tiger shark occupancy while male movements were

directed towards locations with high tiger shark occupancy, indicating sex differences in how

competitors influenced movement through receiver locations, although these effects have

high uncertainty around them.

Figure 4: Summary of the outputs from the ERGM models presented separately for females

and males. Effect sizes and their 95% confidence intervals pooled over variables are reported

for the coefficient of terms *nodecov* and *absdiff*. The variable "Geodistance" controls for the

distance between receivers. Circles represent the number of models in which the coefficient

of the term was significant and therefore selected.

Seasonal differences in movement network structure

The factors that explained movement network structure remained similar between winter and

summer, with only competition and rainfall changing qualitatively (Fig. 5). While rainfall

levels had effects on movements in both seasons, the pattern was less clear during winter,

which is expected as rainfall is a seasonally-driven factor. It appears there is a shift in

behaviour related to the presence of tiger sharks. While there was no apparent movement

pattern related to tiger shark occupancy during winter, there was a pattern of movements

towards locations with low occupancy of tiger shark in summer as demonstrated by positive

absdiff coefficients and negatives coefficient of *nodecov*.

Figure 5: Summary of the outputs from the ERGM models presented separately for summer

and winter. Effect sizes and their 95% confidence intervals pooled over variables are reported

for the coefficient of terms *nodecov* and *absdiff*. The variable "Geodistance" controls for the

distance between receivers. Circles represent the number of models in which the coefficient

of the term was significant and therefore selected.

DISCUSSION

We predicted that bull sharks would respond specifically to certain environmental factors,

being attracted to high levels of turbidity, rainfall and swell. We also expected that bull shark

movements would be influenced by the distribution of conspecifics and heterospecifics in the

seascape, although we could not predict the direction of these effects. We were thus

interested whether these con- and heterospecific effects were as important as abiotic variables

in explaining movement patterns. Using Exponential random graph models (ERGMs), we

found evidence that abiotic factors, turtle density, anthropogenic factors, intraspecific

interactions, and competitive or interspecific interactions all contribute to bull shark

movement network structure, albeit with varying importance. As we hypothesized, our

models revealed that social factors were at least as important as abiotic parameters.

Additionally, the direction of response to abiotic variables were not always as predicted,

indicating that the role of physical variables in shaping movement network structure may be

context dependent or vary across spatial and temporal scales.

Effects of abiotic factors on movements

Our results indicate that environmental factors impact the movements of bull sharks,

however, while we expected to find positive effects of swell height, turbidity and rainfall on

shark movements as suggested by previous studies, we found opposite patterns. Bull sharks

were found to be mostly moving through areas with lower rainfall. While bull sharks are

known to respond to rainfall and freshwater runoff and are able to transit between freshwater

and marine habitats within hours or remain in low salinity areas for days (Niella et al., 2022;

Werry et al., 2018), selection of lower rainfall levels could be due to the spatial and temporal

resolution at which they react to such rapid environmental changes. In addition, bull sharks

may prefer to predominantly remain (at our scale of observation) in the same conditions of

rain, turbidity and salinity as rain levels are often strong and fast in tropical areas. Werry et

al. (2018) suggested that bull sharks are attracted to estuarine and associated nearshore areas

following high rainfall events in part due to increased prey availability while Niella et al.

(2022) suggested a more complex pattern with differences in response between the sexes.

Matich et al., (2020) also highlighted that during exceptional events bull sharks can avoid

high rainfall and high turbidity, moving away from sites with fast and strong perturbations.

The rainfall data entered in our models represent distinct watershed values that can

potentially incorporate multiple receivers, and can potentially explain why our models found

disproportionately strong effect of movements between (nearby) locations with similar levels

of rainfall. In addition, Werry et al. (2018) found a one-week lag in the response of sharks to

rainfall which could reflect changes in food availability and foraging effectiveness with

changes in salinity, a pattern that would be difficult to reveal with our models. Previous

studies that found strong behavioural response of bull sharks to rainfall were conducted in

large estuarine system where rainfall levels and consequences (e.g. induced turbidity) could

contrast with the coral reef coast of Reunion Island. Finally, movements in response to

attractive effects of environmental factors (swell, turbidity and rainfall) that are ephemeral

and irregular may occur over shorter timescales and be difficult to detect with monthly data.

While we expected to find stronger seasonal differences because the variations in these

parameters are more important and persistent in summer than during winter, the lack of such

behavioural differences could be due to bull sharks being resident for less lengthy periods

during summer (Blaison et al., 2015). It is important to note that other potential factors that

we did not measure could also contribute significantly to movement decisions. For example,

salinity is known to be an important factor affecting movement and distribution of bull sharks

(Dwyer et al., 2020). Our results confirm that considering temporally dynamic environmental

variables in studies of animal movement is important because movement patterns reflect

dynamic interactions between animals and their physical environment, with environmental

heterogeneity driving movements of individuals and allowing them to remain in optimal

environmental conditions (Avgar et al., 2013; Mueller et al., 2011; Schick et al., 2008).

Effects of turtle density on movements

Our study did not find any effect of turtle density on bull shark movements. While we

used available data on turtle density to investigate its influence on movement patterns of bull

sharks in this study based on local people's perception, we acknowledge that turtles are not

the main prey of bull sharks and remain opportunistic items (Cliff & Dudley, 1991). Bull

sharks are known to feed mainly on teleost fish (Trystram et al., 2016) and future studies

could integrate dynamic species-specific fishery data or parallel underwater surveys to

investigate the role of abundance and composition of fish communities in driving predator

movements. Indeed, resource availability and heterogeneity are important drivers of animal

movement, especially for predators (i.e. prey abundance and distribution). Finding prey

requires a predator to make adaptive decisions about which movement patterns to adopt to

feed most profitably given a particular prey distribution (Riotte-Lambert & Matthiopoulos,

2020; Sims et al., 2006). Thus, interpreting predator movements within a prey landscape may

provide a clearer picture of why certain habitats are selected over others.

Effects of human densities on movements

While we detected a tendency for more movements through locations with high

human use, the effect sizes were small suggesting human activities did not strongly influence

movement patterns. This is not surprising as nautical human activities and density are spread

fairly evenly along the west coast of the island where the network of receivers was deployed.

In addition, it is now recognized that bull sharks can adapt to urbanized areas and do not

especially avoid these high human density areas and their activities (Hammerschlag et al.,

2022; Werry et al., 2012). The majority of large-bodied terrestrial carnivores tend avoid high

human densities and activities (Tucker et al., 2018, 2023). As opposed to most predators, it is

possible that human activities are not fundamentally avoided by bull sharks because ocean

landscapes maybe affected differently than terrestrial landscapes. This pattern could be

reinforced by the limited suitable habitat available around the island preventing bull sharks

from avoiding most human activities along the coast.

Effects of social environment on movements

We found that social factors sometimes had similar effects on movement networks to

environmental variables. While the bull shark has been found to form occasional

aggregations around fish farms (Loiseau et al., 2016) or at artificial provisioning sites

(Bouveroux et al., 2021), it is not recognized as displaying collective behaviours and strong

patterns of spatial segregation have been found in our study population (Mourier et al., 2021).

However, we found that more movements occurred through bull shark hotspots. When

focusing on the behaviour of each sex, we showed that males were less social and tended to

avoid other sharks regardless of their sex. Contrastingly, females' movements occurred

between locations where other females spent a lot of time but avoided areas heavily used by

males. This confirmed previous findings that females show stronger patterns of residency

providing the opportunity to co-occur with other females, while males favour roaming

behaviour (Mourier et al., 2021). This gregarious behaviour of females could also suggest the

presence of mating arenas where females gather to choose transient males allowing them to

avoid male harassment during the mating period. Further research investigating the spatio-

temporal interactions between bull sharks is required to better understand their aggregative

and avoidance behaviour. Our results thus confirm that an individual's movements are not

solely driven by environmental features. Individuals share space with other conspecifics,

linking spatial and social processes (Albery et al., 2021; Webber et al., 2023). For example,

the spatial distribution of resources inherently drives the proximity of individuals through

foraging aggregations, influencing many forms of social interaction (Macdonald, 1983). As

such, the distribution of individuals of a species in space generates population structure and

will influence movement decisions at the individual level. In fact, regardless of whether a

species is perceived to be solitary or social, individuals will have to decide whether to join or

avoid other conspecifics distributed unevenly in space. Such decisions will also depend on

social processes driven by individual phenotypes that alter social decisions (e.g. size, sex, or

genetic relatedness).

Effects of interspecific competition on movements

In our study we tested the influence of another large predator, the tiger shark, on the

movements of bull sharks. Our results indicated the presence of patterns of avoidance

between bull and tiger sharks. Bull shark movements were affected by the presence of tiger

sharks with female bull sharks directing their movements towards locations with lower tiger

shark occupancy while male sharks showed the opposite pattern. The tiger shark is an apex

predator that is larger than the bull shark and potentially dominant. It is therefore plausible

that bull sharks avoid interactions with tiger sharks. Female bull sharks, being larger than

males, could also compete with tiger sharks. Therefore, one plausible explanation to this

pattern is spatial and foraging niche separation between both species (Niella et al., 2021). We

previously highlighted that male and female bull sharks displayed patterns of spatial

segregation (Mourier et al., 2021; Niella et al., 2022), which could also explain the difference

in interaction opportunities with tiger sharks based on habitat utilization and movement

patterns. Indeed, bull sharks rely more on coastal food sources than tiger sharks which feed

on more coastal-pelagic and deep-water prey (Le Croizier et al., 2020; Trystram et al., 2016),

thus supporting our results and implying local spatial segregation between the two species.

These patterns of interference interactions are relatively common in large sharks, as

competitive species generally show spatial (Papastamatiou et al., 2018), temporal (Lear et al.,

2021) and trophic niche (Matich et al., 2017) partitioning to limit the negative effect of

competition and promote co-existence of predators. Similar avoidance strategies are found

between competiting carnivores in terrestrial ecosystems. Indeed, lions (*Panthera leo*)

usually remain in areas rich in prey, while subordinated carnivores like leopard (*P. pardus*)

and cheetah (*Acinonyx jubatus*) overlap with the home range of lions but use fine-scaled

avoidance behaviours, and small species like African wild dog (*Lycaon pictus*) employ

multiple tactics to avoid all other competitors (Vanak et al., 2013). Coexistence of multiple

carnivore species is typically explained by dietary niche separation as a consequence of

avoiding intraguild competition. Another potential explanation is the avoidance of tiger

sharks by females during summer corresponding to the parturition season (Pirog et al., 2019)

in order to avoid tiger shark predation on bull shark neonates.

Our results thus complement previous studies suggesting that interactions among

large predators involve a complex interplay of competition and predation, as large carnivores

can suppress populations of smaller carnivores through direct predation, resource

competition, or via other forms of interference competition. This may result in spatial and/or

temporal avoidance, reductions in the density of the subordinate species, or even competitive

exclusion from certain habitats (Berger & Gese, 2007; Linnell & Strand, 2000; Prugh et al.,

2009; Prugh & Sivy, 2020).

Conclusion

To investigate the drivers of movements of a marine predator, we used a network-

based approach of movement between fixed stations and employed Exponential Random

Graph Model in a way that could account for the dynamical nature of site attributes visited by

individual animals within a flexible framework that could be extended to test hypotheses

related to the structure of the network itself (extending beyond what is easily possible using

more conventional statistical approaches). While this modelling framework performed well in

our study, it could be refined further to test similar research questions at different spatial and

temporal scales. For example, one promising model is the separable temporal exponential-

family random graph model (tERGM), which treats the formation and dissolution of ties in

parallel at each time step as independent ERGMs (Carnegie et al., 2015; Krivitsky &

Handcock, 2014) and can provide a more realistic view of movement networks (e.g.,

including consecutive individual monthly or daily networks). This would make it possible to

better integrate the temporal interdependency of environmental changes in explaining animal

movement patterns.

Our results fit into a broader picture illustrating that animal movements arise from

complex interactions of individuals with their physical environment as well as with both

surrounding conspecific and heterospecific individuals. By including a diverse set of

variables that may influence bull shark movements within our analyses, we demonstrate that

to fully explain animal movement patterns requires the incorporation of multiple variables

associated with environmental heterogeneity, human footprint and the distribution of

individuals in space, both of the same species and other members of the community. A main

finding was that, although environmental conditions were important factors influencing

movement of bull sharks, interactions with other individuals in their surroundings was also

important to consider, opening new perspectives at the socio-spatial interface for these

marine predators.

Data availability

Data and R codes used in this study are available at

SEANOE: https://doi.org/10.17882/99080

References

Albery, G. F., Morris, A., Morris, S., Pemberton, J. M., Clutton-Brock, T. H., Nussey, D. H.,
        & Firth, J. A. (2021). Multiple spatial behaviours govern social network positions in a
        wild ungulate. *Ecology Letters*, *24*(4), 676–686. https://doi.org/10.1111/ele.13684

Atwood, T. C., Gese, E. M., & Kunkel, K. E. (2009). Spatial Partitioning of Predation Risk in
        a Multiple Predator-Multiple Prey System. *The Journal of Wildlife Management*,
        73(6), 876–884. https://doi.org/10.2193/2008-325

Avgar, T., Mosser, A., Brown, G. S., & Fryxell, J. M. (2013). Environmental and individual
        drivers of animal movement patterns across a wide geographical gradient. *Journal of
        Animal Ecology*, *82*(1), 96–106. https://doi.org/10.1111/j.1365-2656.2012.02035.x

Berger, K. M., & Gese, E. M. (2007). Does interference competition with wolves limit the
        distribution and abundance of coyotes? *Journal of Animal Ecology*, *76*(6), 1075–
        1085. https://doi.org/10.1111/j.1365-2656.2007.01287.x

Blaison, A., Jaquemet, S., Guyomard, D., Vangrevelynghe, G., Gazzo, T., Cliff, G., Cotel, P.,
        & Soria, M. (2015). Seasonal variability of bull and tiger shark presence on the west

677  coast of Reunion Island, western Indian Ocean. *African Journal of Marine Science*,

678  *37*(2), 199–208. https://doi.org/10.2989/1814232X.2015.1050453

679 Bouveroux, T., Loiseau, N., Barnett, A., Marosi, N. D., & Brunnschweiler, J. M. (2021).

680  Companions and Casual Acquaintances: The Nature of Associations Among Bull

681  Sharks at a Shark Feeding Site in Fiji. *Frontiers in Marine Science*, *8*.

682  https://doi.org/10.3389/fmars.2021.678074

683 Brunnschweiler, J. M., Queiroz, N., & Sims, D. W. (2010). Oceans apart? Short-term

684  movements and behaviour of adult bull sharks Carcharhinus leucas in Atlantic and

685  Pacific Oceans determined from pop-off satellite archival tagging. *Journal of Fish

686  Biology*, *77*(6), 1343–1358. https://doi.org/10.1111/j.1095-8649.2010.02757.x

687 Capello, M., Robert, M., Soria, M., Potin, G., Itano, D., Holland, K., Deneubourg, J.-L., &

688  Dagorn, L. (2015). A Methodological Framework to Estimate the Site Fidelity of

689  Tagged Animals Using Passive Acoustic Telemetry. *PLOS ONE*, *10*(8), e0134002.

690  https://doi.org/10.1371/journal.pone.0134002

691 Carnegie, N. B., Krivitsky, P. N., Hunter, D. R., & Goodreau, S. M. (2015). An

692  Approximation Method for Improving Dynamic Network Model Fitting. *Journal of

693  Computational and Graphical Statistics*, *24*(2), 502–519.

694  https://doi.org/10.1080/10618600.2014.903087

695 Cliff, G., & Dudley, S. F. J. (1991). Sharks caught in the protective gill nets off Natal, South

696  Africa. 4. The bull shark Carcharhinus leucas Valenciennes. *South African Journal of

697  Marine Science*, *10*(1), 253–270. https://doi.org/10.2989/02577619109504636

698 Couriot, O., Hewison, A. J. M., Saïd, S., Cagnacci, F., Chamaillé-Jammes, S., Linnell, J. D.

699  C., Mysterud, A., Peters, W., Urbano, F., Heurich, M., Kjellander, P., Nicoloso, S.,

700  Berger, A., Sustr, P., Kroeschel, M., Soennichsen, L., Sandfort, R., Gehr, B., &

701  Morellet, N. (2018). Truly sedentary? The multi-range tactic as a response to resource

heterogeneity and unpredictability in a large herbivore. *Oecologia*, *187*(1), 47–60.

https://doi.org/10.1007/s00442-018-4131-5

Croft, D. P., Madden, J. R., Franks, D. W., & James, R. (2011). Hypothesis testing in animal

social networks. *Trends in Ecology & Evolution*, *26*(10), 502–507.

https://doi.org/10.1016/j.tree.2011.05.012

Daly, R., Smale, M. J., Cowley, P. D., & Froneman, P. W. (2014). Residency Patterns and

Migration Dynamics of Adult Bull Sharks (Carcharhinus leucas) on the East Coast of

Southern Africa. *PLOS ONE*, *9*(10), e109357.

https://doi.org/10.1371/journal.pone.0109357

Dey, C. J., & Quinn, J. S. (2014). Individual attributes and self-organizational processes

affect dominance network structure in pukeko. *Behavioral Ecology*, *25*(6), 1402–

1408. https://doi.org/10.1093/beheco/aru138

Dwyer, R. G., Campbell, H. A., Cramp, R. L., Burke, C. L., Micheli-Campbell, M. A.,

Pillans, R. D., Lyon, B. J., & Franklin, C. E. (2020). Niche partitioning between river

shark species is driven by seasonal fluctuations in environmental salinity. *Functional*

Ecology, *34*(10), 2170–2185. https://doi.org/10.1111/1365-2435.13626

Espinoza, M., Heupel, M. R., Tobin, A. J., & Simpfendorfer, C. A. (2016). Evidence of

Partial Migration in a Large Coastal Predator: Opportunistic Foraging and

Reproduction as Key Drivers? *PLOS ONE*, *11*(2), e0147608.

https://doi.org/10.1371/journal.pone.0147608

Fletcher, R. J., Acevedo, M. A., Reichert, B. E., Pias, K. E., & Kitchens, W. M. (2011).

Social network models predict movement and connectivity in ecological landscapes.

Proceedings of the National Academy of Sciences, *108*(48), 19282–19287.

https://doi.org/10.1073/pnas.1107549108

Gilbert, N. A., McGinn, K. A., Nunes, L. A., Shipley, A. A., Bernath-Plaisted, J., Clare, J. D.

J., Murphy, P. W., Keyser, S. R., Thompson, K. L., Nelson, S. B. M., Cohen, J. M.,

Widick, I. V., Bartel, S. L., Orrock, J. L., & Zuckerberg, B. (2023). Daily activity

timing in the Anthropocene. *Trends in Ecology & Evolution*, *38*(4), 324–336.

https://doi.org/10.1016/j.tree.2022.10.008

Gilbert, N. A., Stenglein, J. L., Pauli, J. N., & Zuckerberg, B. (2022). Human disturbance

compresses the spatiotemporal niche. *Proceedings of the National Academy of*

Sciences, *119*(52), e2206339119. https://doi.org/10.1073/pnas.2206339119

Giuggioli, L., & Kenkre, V. M. (2014). Consequences of animal interactions on their

dynamics: Emergence of home ranges and territoriality. *Movement Ecology*, *2*(1), 20.

https://doi.org/10.1186/s40462-014-0020-7

Hammerschlag, N., Gutowsky, L. F. G., Rider, M. J., Roemer, R., & Gallagher, A. J. (2022).

Urban sharks: Residency patterns of marine top predators in relation to a coastal

metropolis. *Marine Ecology Progress Series*, *691*, 1–17.

https://doi.org/10.3354/meps14086

He, P., Maldonado-Chaparro, A. A., & Farine, D. R. (2019). The role of habitat configuration

in shaping social structure: A gap in studies of animal social complexity. *Behavioral*

Ecology and Sociobiology, *73*(1), 9. https://doi.org/10.1007/s00265-018-2602-7

Heupel, M. R., Simpfendorfer, C. A., Espinoza, M., Smoothey, A. F., Tobin, A., &

Peddemors, V. (2015). Conservation challenges of sharks with continental scale

migrations. *Frontiers in Marine Science*, *2*. https://doi.org/10.3389/fmars.2015.00012

Hunter, D. R., Handcock, M. S., Butts, C. T., Goodreau, S. M., & Morris, M. (2008). ergm: A

Package to Fit, Simulate and Diagnose Exponential-Family Models for Networks.

Journal of Statistical Software, *24*(1), Article 1. https://doi.org/10.18637/jss.v024.i03

Jacoby, D. M. P., & Freeman, R. (2016). Emerging Network-Based Tools in Movement

Ecology. *Trends in Ecology & Evolution*, *31*(4), 301–314.

https://doi.org/10.1016/j.tree.2016.01.011

Johnson, A. R., Wiens, J. A., Milne, B. T., & Crist, T. O. (1992). Animal movements and

population dynamics in heterogeneous landscapes. *Landscape Ecology*, *7*(1), 63–75.

https://doi.org/10.1007/BF02573958

Krause, J., & Ruxton, G. D. (2002). *Living in Groups*. OUP Oxford.

Krivitsky, P. N., & Handcock, M. S. (2014). A separable model for dynamic networks.

Journal of the Royal Statistical Society: Series B (Statistical Methodology), *76*(1), 29–

46. https://doi.org/10.1111/rssb.12014

Kuefler, D., Avgar, T., & Fryxell, J. M. (2012). Rotifer population spread in relation to food,

density and predation risk in an experimental system. *Journal of Animal Ecology*,

81(2), 323–329. https://doi.org/10.1111/j.1365-2656.2011.01917.x

Le Croizier, G., Lorrain, A., Sonke, J. E., Jaquemet, S., Schaal, G., Renedo, M., Besnard, L.,

Cherel, Y., & Point, D. (2020). Mercury isotopes as tracers of ecology and

metabolism in two sympatric shark species. *Environmental Pollution*, *265*, 114931.

https://doi.org/10.1016/j.envpol.2020.114931

Lear, K. O., Whitney, N. M., Morris, J. J., & Gleiss, A. C. (2021). Temporal niche

partitioning as a novel mechanism promoting co-existence of sympatric predators in

marine systems. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1954),

20210816. https://doi.org/10.1098/rspb.2021.0816

Lee, K. A., Smoothey, A. F., Harcourt, R. G., Roughan, M., Butcher, P. A., & Peddemors, V.

M. (2019). Environmental drivers of abundance and residency of a large migratory

shark, Carcharhinus leucas, inshore of a dynamic western boundary current. *Marine*

Ecology Progress Series, *622*, 121–137. https://doi.org/10.3354/meps13052

Linnell, J. D. C., & Strand, O. (2000). Interference interactions, co-existence and

conservation of mammalian carnivores. *Diversity and Distributions*, *6*(4), 169–176.

https://doi.org/10.1046/j.1472-4642.2000.00069.x

Loiseau, N., Kiszka, J. J., Bouveroux, T., Heithaus, M. R., Soria, M., & Chabanet, P. (2016).

Using an unbaited stationary video system to investigate the behaviour and

interactions of bull sharks Carcarhinus leucas under an aquaculture farm. *African*

Journal of Marine Science, *38*(1), 73–79.

https://doi.org/10.2989/1814232X.2016.1156578

López-Calderón, C., Martín-Vélez, V., Blas, J., Höfle, U., Sánchez, M. I., Flack, A., Fiedler,

W., Wikelski, M., & Green, A. J. (2023). White stork movements reveal the

ecological connectivity between landfills and different habitats. *Movement Ecology*,

11(1), 18. https://doi.org/10.1186/s40462-023-00380-7

Lubitz, N., Daly, R., Filmalter, J. D., Sheaves, M., Cowley, P. D., Naesje, T. F., & Barnett,

A. (2023). Context drives movement patterns in a mobile marine predator. *Movement*

Ecology, *11*(1), 28. https://doi.org/10.1186/s40462-023-00390-5

Lusher, D., Koskinen, J., & Robins, G. (Eds.). (2012). *Exponential Random Graph Models*

for Social Networks: Theory, Methods, and Applications. Cambridge University

Press. https://doi.org/10.1017/CBO9780511894701

Macdonald, D. W. (1983). The ecology of carnivore social behaviour. *Nature*, *301*(5899),

Article 5899. https://doi.org/10.1038/301379a0

Matich, P., Kiszka, J. J., Mourier, J., Planes, S., & Heithaus, M. R. (2017). Species co-

occurrence affects the trophic interactions of two juvenile reef shark species in

tropical lagoon nurseries in Moorea (French Polynesia). *Marine Environmental*

Research, *127*(Supplement C), 84–91.

https://doi.org/10.1016/j.marenvres.2017.03.010

Matich, P., Strickland, B. A., & Heithaus, M. R. (2020). Long-term monitoring provides

insight into estuarine top predator (Carcharhinus leucas) resilience following an

extreme weather event. *Marine Ecology Progress Series*, *639*, 169–183.

https://doi.org/10.3354/meps13269

Morales, J. M., Moorcroft, P. R., Matthiopoulos, J., Frair, J. L., Kie, J. G., Powell, R. A.,

Merrill, E. H., & Haydon, D. T. (2010). Building the bridge between animal

movement and population dynamics. *Philosophical Transactions of the Royal Society*

B: Biological Sciences, *365*(1550), 2289–2301.

https://doi.org/10.1098/rstb.2010.0082

Mourier, J., Lédée, E. J. I., & Jacoby, D. M. P. (2019). A multilayer perspective for inferring

spatial and social functioning in animal movement networks. *bioRxiv*, 749085.

https://doi.org/10.1101/749085

Mourier, J., Soria, M., Blaison, A., Simier, M., Certain, G., Demichelis, A., & Hattab, T.

(2021). Dynamic use of coastal areas by bull sharks and the conciliation of

conservation and management of negative human–wildlife interactions. *Aquatic*

Conservation: Marine and Freshwater Ecosystems, *31*(10), 2926–2937.

https://doi.org/10.1002/aqc.3674

Mueller, T., Olson, K. A., Dressler, G., Leimgruber, P., Fuller, T. K., Nicolson, C., Novaro,

A. J., Bolgeri, M. J., Wattles, D., DeStefano, S., Calabrese, J. M., & Fagan, W. F.

(2011). How landscape dynamics link individual- to population-level movement

patterns: A multispecies comparison of ungulate relocation data. *Global Ecology and*

Biogeography, *20*(5), 683–694. https://doi.org/10.1111/j.1466-8238.2010.00638.x

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E.

(2008). A movement ecology paradigm for unifying organismal movement research.

Proceedings of the National Academy of Sciences, *105*(49), 19052–19059.

https://doi.org/10.1073/pnas.0800375105

Niella, Y., Smoothey, A. F., Taylor, M. D., Peddemors, V. M., & Harcourt, R. (2022).

Environmental Drivers of Fine-Scale Predator and Prey Spatial Dynamics in Sydney

Harbour, Australia, and Adjacent Coastal Waters. *Estuaries and Coasts*, *45*(5), 1465–

1479. https://doi.org/10.1007/s12237-021-01020-2

Niella, Y., Wiefels, A., Almeida, U., Jaquemet, S., Lagabrielle, E., Harcourt, R., Peddemors,

V., & Guyomard, D. (2021). Dynamics of marine predators off an oceanic island and

implications for management of a preventative shark fishing program. *Marine*

Biology, *168*(4), 42. https://doi.org/10.1007/s00227-021-03852-9

Ohta, I., & Kakuma, S. (2005). Periodic behavior and residence time of yellowfin and bigeye

tuna associated with fish aggregating devices around Okinawa Islands, as identified

with automated listening stations. *Marine Biology*, *146*(3), 581–594.

https://doi.org/10.1007/s00227-004-1456-x

Papastamatiou, Y. P., Bodey, T. W., Friedlander, A. M., Lowe, C. G., Bradley, D., Weng, K.,

Priestley, V., & Caselle, J. E. (2018). Spatial separation without territoriality in shark

communities. *Oikos*, *127*(6), 767–779. https://doi.org/10.1111/oik.04289

Pirog, A., Magalon, H., Poirout, T., & Jaquemet, S. (2019). Reproductive biology, multiple

paternity and polyandry of the bull shark Carcharhinus leucas. *Journal of Fish*

Biology, *95*(5), 1195–1206. https://doi.org/10.1111/jfb.14118

Prugh, L. R., & Sivy, K. J. (2020). Enemies with benefits: Integrating positive and negative

interactions among terrestrial carnivores. *Ecology Letters*, *23*(5), 902–918.

https://doi.org/10.1111/ele.13489

Prugh, L. R., Stoner, C. J., Epps, C. W., Bean, W. T., Ripple, W. J., Laliberte, A. S., &

Brashares, J. S. (2009). The Rise of the Mesopredator. *BioScience*, *59*(9), 779–791.

https://doi.org/10.1525/bio.2009.59.9.9

Pyke, G. H., Pulliam, H. R., & Charnov, E. L. (1977). Optimal Foraging: A Selective Review

of Theory and Tests. *The Quarterly Review of Biology*, *52*(2), 137–154.

Riotte-Lambert, L., & Matthiopoulos, J. (2020). Environmental Predictability as a Cause and

Consequence of Animal Movement. *Trends in Ecology & Evolution*, *35*(2), 163–174.

https://doi.org/10.1016/j.tree.2019.09.009

Robins, G., Pattison, P., Kalish, Y., & Lusher, D. (2007). An introduction to exponential

random graph (p*) models for social networks. *Social Networks*, *29*(2), 173–191.

https://doi.org/10.1016/j.socnet.2006.08.002

Robins, G., Snijders, T., Wang, P., Handcock, M., & Pattison, P. (2007). Recent

developments in exponential random graph (p*) models for social networks. *Social*

Networks, *29*(2), 192–215. https://doi.org/10.1016/j.socnet.2006.08.003

Schick, R. S., Loarie, S. R., Colchero, F., Best, B. D., Boustany, A., Conde, D. A., Halpin, P.

N., Joppa, L. N., McClellan, C. M., & Clark, J. S. (2008). Understanding movement

data and movement processes: Current and emerging directions. *Ecology Letters*,

11(12), 1338–1350. https://doi.org/10.1111/j.1461-0248.2008.01249.x

Schwarzer, G. R. (2007). An R package for meta-analysis. *R News*, *7*, 40–45.

Silk, M. J., & Fisher, D. N. (2017). Understanding animal social structure: Exponential

random graph models in animal behaviour research. *Animal Behaviour*, *132*, 137–

146. https://doi.org/10.1016/j.anbehav.2017.08.005

Sims, D. W., Witt, M. J., Richardson, A. J., Southall, E. J., & Metcalfe, J. D. (2006).

Encounter success of free-ranging marine predator movements across a dynamic prey

landscape. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1591),

1195–1201. https://doi.org/10.1098/rspb.2005.3444

Snijders, T. A. B., Pattison, P. E., Robins, G. L., & Handcock, M. S. (2006). New

Specifications for Exponential Random Graph Models. *Sociological Methodology*,

36(1), 99–153. https://doi.org/10.1111/j.1467-9531.2006.00176.x

Soria, M., Heithaus, M. R., Blaison, A., Crochelet, E., Forget, F., & Chabanet, P. (2019).

Residency and spatial distribution of bull sharks Carcharhinus leucas in and around

Reunion Island marine protected area. *Marine Ecology Progress Series*, *630*, 101–

113. https://doi.org/10.3354/meps13139

Trystram, C., Rogers, K. M., Soria, M., & Jaquemet, S. (2016). Feeding patterns of two

sympatric shark predators in coastal ecosystems of an oceanic island. *Canadian

Journal of Fisheries and Aquatic Sciences*, *74*(2), 216–227.

https://doi.org/10.1139/cjfas-2016-0105

Tucker, M. A., Böhning-Gaese, K., Fagan, W. F., Fryxell, J. M., Moorter, B. V., Alberts, S.

C., Ali, A. H., Allen, A. M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar,

B., Belant, J. L., Bertassoni, A., Beyer, D., Bidner, L., Beest, F. M. van, Blake, S.,

Blaum, N., … Mueller, T. (2018). Moving in the Anthropocene: Global reductions in

terrestrial mammalian movements. *Science*, *359*(6374), 466–469.

https://doi.org/10.1126/science.aam9712

Tucker, M. A., Schipper, A. M., Adams, T. S. F., Attias, N., Avgar, T., Babic, N. L., Barker,

K. J., Bastille-Rousseau, G., Behr, D. M., Belant, J. L., Beyer, D. E., Blaum, N.,

Blount, J. D., Bockmühl, D., Pires Boulhosa, R. L., Brown, M. B., Buuveibaatar, B.,

Cagnacci, F., Calabrese, J. M., … Mueller, T. (2023). Behavioral responses of

terrestrial mammals to COVID-19 lockdowns. *Science*, *380*(6649), 1059–1064.

https://doi.org/10.1126/science.abo6499

Vanak, A. T., Fortin, D., Thaker, M., Ogden, M., Owen, C., Greatwood, S., & Slotow, R.

(2013). Moving to stay in place: Behavioral mechanisms for coexistence of African

large carnivores. *Ecology*, *94*(11), 2619–2631. https://doi.org/10.1890/13-0217.1

Wearmouth, V. J., & Sims, D. W. (2008). Chapter 2 Sexual Segregation in Marine Fish,

Reptiles, Birds and Mammals: Behaviour Patterns, Mechanisms and Conservation

Implications. In *Advances in Marine Biology* (Vol. 54, pp. 107–170). Academic Press.

https://doi.org/10.1016/S0065-2881(08)00002-3

Webber, Q. M. R., Albery, G. F., Farine, D. R., Pinter-Wollman, N., Sharma, N., Spiegel, O.,

Vander Wal, E., & Manlove, K. (2023). Behavioural ecology at the spatial–social

interface. *Biological Reviews*, *98*(3), 868–886. https://doi.org/10.1111/brv.12934

Werry, J. M., Lee, S. Y., Lemckert, C. J., & Otway, N. M. (2012). Natural or Artificial?

Habitat-Use by the Bull Shark, Carcharhinus leucas. *PLOS ONE*, *7*(11), e49796.

https://doi.org/10.1371/journal.pone.0049796

Werry, J. M., Sumpton, W., Otway, N. M., Lee, S. Y., Haig, J. A., & Mayer, D. G. (2018).

Rainfall and sea surface temperature: Key drivers for occurrence of bull shark,

Carcharhinus leucas, in beach areas. *Global Ecology and Conservation*, *15*, e00430.

https://doi.org/10.1016/j.gecco.2018.e00430