Both environmental conditions and intra- and interspecific interactions

influence the movements of a marine predator

Johann Mourier[1], Marc Soria[1], Matthew Silk[2,3,4], Angélique Demichelis[1], Laurent Dagorn[1],

Tarek Hattab[1]

[1] MARBEC, Univ Montpellier, CNRS, IFREMER, IRD, Sète, France

[2] Centre for Ecology and Conservation, University of Exeter Penryn Campus, Cornwall, UK

[3] NIMBioS, The University of Tennessee Knoxville, Knoxville, TN, USA

[4] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

Corresponding author: Johann Mourier

email: johann.mourier@umontpellier.fr

Address: MARBEC, Station Ifremer, Avenue Jean Monnet, BP 171, 34203 Sète Cedex -

France.

ABSTRACT

Animal movements are typically influenced by multiple environmental factors

simultaneously and individuals vary in their response to this environmental heterogeneity.

Therefore, understanding how environmental aspects, including biotic, abiotic and

anthropogenic factors, influence the movements of wild animals is an important focus of

wildlife research and conservation. We apply Exponential Random Graph Models (ERGMs)

to analyse movement networks of a bull shark population in a network of acoustic receivers

and identify the effects of environmental, social or other types of covariates on their

movements. We found that intra- and interspecific factors often had stronger effects on

movements than environmental variables. ERGMs proved to be a potentially useful tool for

studying animal movement network data especially in the context of spatial attribute

heterogeneity.

Keywords: spatial-social interface, acoustic telemetry; *Carcharhinus leucas*; ERGM;

movement networks; Indian Ocean; Reunion Island; shark.

Understanding animal movement is not straightforward, as movement decisions result

from a combination of many factors including the internal state of the individual, its intra-

and interspecific neighbourhood, and spatio-temporal variation in abiotic environmental

conditions (Nathan et al., 2008). Additionally, the response of individuals to variation in their

surrounding environment can differ across spatial and temporal scales, as well as along their

ontogeny. Daily activities can be influenced by environmental factors that vary over a diel

cycle (e.g., wind or tidal currents), while seasonal movement can be governed by changes

acting at larger temporal scales (e.g., photoperiod, temperature). Quantifying how

heterogeneity in an animal's surroundings drives behaviour, and consequently movement

patterns, provides insight into how natural and anthropogenic changes may impact

populations and ecosystems.

One potential way to understand the 'causes, mechanisms and spatiotemporal patterns

of movement and their role in various ecological and evolutionary processes' (Nathan et al.

2008) is to link observed movement patterns with spatial and temporal variability in the

underlying environmental conditions (Avgar et al., 2013). Movement patterns result from

interactions between organisms and their environments (Johnson et al., 1992; Morales et al.,

2010; Schick et al., 2008). For example, movement rates are expected to decrease where

resources are rich and increase where food availability is low (Kuefler et al., 2012; Pyke et

al., 1977). Understanding movement patterns thus requires consideration of the temporally

dynamic nature of these environments (Avgar et al., 2013; Couriot et al., 2018; Mueller et al.,

2011; Riotte-Lambert & Matthiopoulos, 2020; Schick et al., 2008).

In addition to interacting with their physical environment, animals also interact with

other individuals of the same or different species. These spatial encounters with conspecific

or heterospecific individuals may be advantageous or unfavourable. For example,

aggregations form to avoid predation or to forage efficiently (Krause & Ruxton, 2002),

although the tendency for non-gregarious species to seek or avoid each other is less well

known. For all sexual species, individuals need to encounter conspecifics at least for mating,

but avoiding opposite-sex individuals (sexual segregation) may play a major role in

preventing sexual harassment in some species (Wearmouth & Sims, 2008). Additionally,

avoidance strategies can act at the intra- or interspecific level in the context of competition

for mates, space or resources (Giuggioli & Kenkre, 2014), or in the context of predator–prey

interactions (Atwood et al., 2009). Because individuals may select habitats based on

exogenous environmental factors (e.g. habitat quality and predation risk), and local

population factors (e.g., mating opportunities, competition or density dependence), we need

to quantify the mechanisms by which dynamic social interactions between individuals occur

(e.g. movement towards or away from other individuals). In fact, interactions with

conspecifics are intrinsically connected with spatial behaviour and a burgeoning literature

now addresses the link between spatial and social aspects of behaviour defined as the

"spatial-social interface" (Webber et al., 2023). While spatial and social processes have been

predominantly considered independently, because movement behaviour emerges from social

and spatial processes, animal movement studies offer an opportunity to consider and integrate

them (Albery et al., 2021; He et al., 2019; Mourier et al., 2019; Webber et al., 2023).

Finally, the human footprint has altered the spatial ecology of many species at

different spatial and temporal scales, for example by decreasing animal movements as a

result of behavioural changes, habitat fragmentation and barrier effects (Doherty et al., 2021;

Tucker et al., 2018), or by modifying activity-timing (Gilbert et al., 2023). COVID-19

lockdowns provided an empirical experiment where an abrupt reduction in human activity (so

called Anthropause) led to decreases in displacements and a reduced avoidance of human

activity (Tucker et al., 2023). Human disturbance can also fundamentally alter the way that

species interact. For example, it can induce a spatiotemporal compression of species co-

occurrences in disturbed landscape which can lead to increases in competition, predation and

infectious disease transmission (Gilbert et al., 2022). As such, incorporating human footprint

as a potential factor affecting animal movements at the spatial-social interface is now more

relevant than ever.

In this study, we explore the connection between seascape attributes and animal

movement patterns for a population of bull shark (*Carcharhinus leucas*) that inhabit a coastal

zone in Reunion Island (Indian Ocean), which encompasses a variety of environmental

conditions. We evaluate the relative importance of spatiotemporal variables associated with

several main aspects of the seascape (abiotic conditions, social environment, predation risk or

interspecific competition and anthropogenic pressure) as potential drivers of shark movement

patterns. This approach allows us to identify fundamental relationships between local

environmental conditions and animal movement patterns.

The bull shark is a large Carcharhinidae species with a wide cosmopolitan distribution

along the continental coasts of all tropical and sub-tropical waters of the world, and is known

to be mobile and move across a large range of habitats and environmental conditions

(Brunnschweiler et al., 2010; Daly et al., 2014; Espinoza et al., 2016; Heupel et al., 2015; Lee

et al., 2019; Niella et al., 2022). Additionally, it is also an euryhaline elasmobranch that uses

a range of salinities throughout its life-cycle (Niella et al., 2022). Its behaviour varies across

spatial and temporal scales, as well as according to size and sex, with a high among-

individual variability in the tendency to move (Espinoza et al., 2016; Lee et al., 2019;

Mourier, Soria, et al., 2021). However, much remains unclear about adult bull shark

movement decisions. Bull shark movements are known to be influenced by both biotic and

abiotic factors (Lee et al., 2019; Lubitz et al., 2023; Niella et al., 2022; Werry et al., 2018),

and anthropogenic factors (Hammerschlag et al., 2022; Werry et al., 2012), but among the

biotic factors, less is known about the importance of con- and heterospecifics in movement

decisions. Even if most studies on the movements of these marine predators have identified

rainfall, sea surface temperature and food distribution as the most influential to their

movement ecology, we expect that social and competition factors may be just as important in

explaining individual movements. Our analysis was designed to test whether individual bull

sharks move according to the distribution of conspecifics and heterospecifics in the seascape,

and whether these effects were comparable with expected positive effects of swell height,

turbidity and rainfall on shark movements and negative effects of human densities (with

sharks avoiding human presence). While we predict that bull sharks will respond to certain

abiotic factors, being attracted to high levels of turbidity, rainfall and swell as revealed in

other studies, we also expect the distribution of conspecifics and heterospecifics in the

seascape may drive movement decisions and that male and female sharks may have different

strategies.

METHODS

Study location

Reunion Island (21°07'S / 55°32'E) is a volcanic island located 700 km east of Madagascar

in the southwest Indian Ocean. The island is 2512 km^2 with 217 km of coastline and

characterized by steep underwater slopes (*ca.* 10-20%) to a depth of 2,000 m. Fringing reefs

stretch over 25 km along the west and south-west coast (Figure 1) forming a natural coral

barrier that bounds the reef flats and back-reef depressions and lies no further than 500 m

from the beach.

Acoustic telemetry and movement network

Sharks were captured along the west coast of Reunion Island between September 2012 and

March 2013 (Supplementary material 1 Table S1), using horizontal drifting long-lines 0.2 to

1 km in length and equipped with 20 to 200 baited 16/0 circle hooks (Blaison et al., 2015).

Set times were fixed at a maximum of 3 h to minimize shark and bycatch mortality. Once

captured, a shark was brought alongside the vessel and held still by rubber-encased ropes to

prevent skin lesions and burns and rolled onto its back to induce tonic immobility. The boat

moved forward slowly throughout the procedure to allow the shark to breathe. Sanitized

transmitters (Innovasea V16, transmission interval 40–80 s, estimated battery life 845 days)

were implanted into the peritoneal cavity through a midventral incision using a sterile scalpel.

A 1cm incision was made, then enlarged with retractors. Once the tag was in place, two

stitches closed the incision and facilitated healing. A sterile, non-absorbable synthetic

monofilament suture (polyamide) was used. The shark was then freed by cutting the hook at

its base and removing it, untying the lasso, replacing it on its belly and oxygenating it by

moving it back and forth or slowly forward until the first signs of autonomous movement

appeared. Sex and total length (TL) were recorded during the procedure. A total of 23

individual sharks (i.e., 6 males and 17 females) were equipped with transmitters

(Supplementary materials 1 Table S1).

An array of 46 Innovasea VR2W acoustic receivers was deployed along the coast with

receivers installed an average of approximately 2 km apart at depths of 10–60 m, comprising

33 (71%) offshore receivers deployed between 300 m and 5 km from shore, and 13 (29%)

inshore receivers placed less than 300 m from shore (Figure 1).

For each shark visit at a receiver, we used detection records to calculate a continuous

residency time (CRT) corresponding to the duration within which a tagged shark was

continuously monitored at a specific receiver (Capello et al., 2015; Ohta & Kakuma, 2005).

All detections of the same shark at one receiver separated by less than a predefined period,

called the maximum blanking period (1 h), were grouped into one CRT and defined as a visit.

Each time a tagged shark was detected at a different receiver, a new visit started - ending the

visit at the previous receiver - even if the interval between detections was less than the

maximum blanking period. We then built monthly movement networks of each shark with

each node representing a receiver along the coast of Reunion Island and each weighted,

directed edge represented the number of movements of the individual (deduced from CRT)

from node A to node B (Figure 1) within a given month.

Explanatory variables

We gathered a variety of biological, abiotic, anthropogenic and spatial data that were

accessible in the study area. Although non-exhaustive, the explanatory variables recorded

included a number of environmental, biological and anthropogenic factors (Table 1; a

detailed description of how they were recorded can be found in Supplementary material 1) as

well as the geographical distances between receivers. We incorporated swell height, turbidity

and rainfall levels as abiotic explanatory variables of movements. Turtle densities were

included as a potential indicator of the presence of sharks, as suggested by local people.

Cumulative number of sea users (boats, swimmer and nautical activities) represents an

anthropogenic variable. We also included various intra- and interspecific factors such as

residency times of the opposite sex, abundance of bull sharks and occupancy of tiger sharks

(a larger and likely competitor species) as con- and heterospecific social interaction variables.

As all data were not available at the same spatial and temporal scale, we chose to standardize

all available candidate explanatory data at the scale of one month for each receiver, which

practically corresponded to calculating a mean for each explanatory variable at each receiver

for each month. This monthly resolution represents a compromise between biologically

meaningful variation in site conditions (i.e., node attributes) and retaining enough movements

to construct individual networks. For some environmental variables such as rainfall, multiple

nearby receivers shared values as data were available for a specific zone encompassing

several receivers. All explanatory variables were standardized to have a mean of 0 and a

standard deviation of 1.

Exponential Random Graph Models

Care is required when conducting statistical analysis of network data because of issues

related to potential non-independence among neighbouring nodes and edges (Croft et al.,

2011). Thus, we used Exponential Random Graph Models (ERGMs). ERGMs are statistical

models of networks that treat the weight of network edges as the response variable and

network node and edge attributes as explanatory variables (Robins, Pattison, et al., 2007;

Robins, Snijders, et al., 2007; Snijders et al., 2006). Such models are analogous to

Generalized Linear Models (GLMs) except that they enable hypothesis testing about the

processes driving network structure and link formation. These models have been mostly

applied in social science, but their properties also make them useful for answering questions

related to how and why animals move between locations in the context of movement

networks (Fletcher et al., 2011; Jacoby & Freeman, 2016; López-Calderón et al., 2023).

Effectively the movement network becomes the response in a regression model, where the

predictors are the propensity for nodes (i.e. locations) of similar or dissimilar attributes to be

linked by movements. In the current study, we applied ERGMs to our monthly individual

bull shark movement networks. These ERGMs are treat the weight of network edges (i.e.

number of movements between two nodes) as the response variable and network node (i.e.

acoustic receivers) attributes as explanatory variables.

The general form for an ERGM can be written as:

216
$$P(Y = y; \theta, x) = \frac{\exp(\theta^T g(y,x))}{\kappa(\theta,x)} \quad (1)$$

where:

• *Y* is the random variable for the state of the network (with realization *y*),

• *g(y,x)* is a vector of model statistics for network *y*,

• *θ* is the vector of coefficients for those statistics, and

• *κ(θ)* is a normalizing term which ensures that equation (1) is a proper probability

distribution. It represents the quantity in the numerator summed over all possible

networks (typically constrained to be all networks with the same node set as *y*).

The numerator represents a formula that is linear in the log form (equation 2):

225
$$\log\big(\exp\big(\theta' g(y)\big)\big) = \theta_1 g_1(y) + \theta_2 g_2(y) + \cdots + \theta_p g_p(y) \quad (2)$$

where *p* is the number of terms in the model. From this one we can more easily observe the

analogy to a traditional statistical model. The functions *g(y)* are counts of configurations in

the network *y* and the parameters *θ* weight the relative importance of the respective

configurations, effectively the size and direction of the effects of the covariates. Parameter

estimation in most specifications of ERGMs uses maximum pseudo-likelihood, an

approximation of maximum likelihood based on Monte Carlo estimation.

All ERGMs were fitted using R packages *ergm* and *ergm.count* (Hunter et al., 2008).

Model-fitting process

Node attributes were varying at the month scale so we built a model for each monthly-

individual network. To ensure that it was possible to fit ERGMs to monthly networks and

improve model convergence, we first removed from monthly movement networks all

receivers that were deployed less than 20 days during the month, as well as all individuals

that did not present an active tag for at least 20 days during a month. This empirical choice

was made as a compromise between data loss and data representativity. We also retained only

monthly individual networks containing at least 10 different directed movements between

receivers, as networks including low numbers of movements caused problems with model

convergence and parameter estimation. We also excluded two monthly networks that were

binary (i.e. only contained at most single movements along any particular edge). This left us

with 42 monthly networks of 13 individuals (9 females: 4 males; mean/median/range of

monthly networks per individual = 3.1/3/1-6).

For each monthly network and each shark, we then fitted two alternative versions of the full

model (using terms 'nodecov' and 'absdiff') accounting for different effects of the variables

being studied (Table 1). The effect estimated for each factor represents the difference from

the intercept as with classical GLMs. All models included the term "non-zero" to control for

zero-inflation generated by the weighted movement networks being sparse. The term "sum",

corresponding to the sum of all link values, was included as the equivalent to the intercept in

a linear model (Dey & Quinn, 2014).

The two model sets differed as follows:

• Model set 1: We tested the association between the current state of the environment at

each receiver location and number of movements towards and away from it (i.e. effect

of an attribute on out- and in-degree). These models used the *nodecov* term in the

fitted ERGMs.

• Model set 2: We tested the association between the absolute difference in current state

of the environment at dyads (pairs) of receiver locations and number of movements

between them. These models used the *absdiff* term in the fitted ERGMs.

We used a *nodecov* model rather than separate *nodeicov* and *nodeocov* models (movements

towards and away from receivers respectively) as simulations indicated the results were

qualitatively identical (see Supplementary material 2).

We fitted the *nodecov* and *absdiff* models separately to facilitate model parsimony and

convergence and test the robustness of the movement patterns discovered while controlling

for the full suite of environmental variables. Indeed, incorporating all variables and model

sets combinations would prevent convergence of model and render interpretation of output

complex. To account for the effect of the spatial distribution of nodes in our models, we

included as a covariate the matrix of distances between nodes.

For each model set, we then wrote a model formula (equation 3) including all potential

predictors as follows:

276
$$graph \sim nonzero + sum + term(Variable\ 1) + term(Variable\ 2) + \cdots +$$

277
$$term(Variable\ n) + edgecov(Geodist, 'Geodist') \qquad (3)$$

where *term* can be replaced by *nodecov* and *absdiff* in model sets 1 and 2, respectively. An

edgecov term was added to control for the distance between locations. Models were fitted

with a Poisson reference distribution for edge weights creating the familiar log-linear effect.

The final coefficient estimates indicate log-linear increase in the weight variable

(movements). MCMLE.Hummel.maxit was set to 1000 (Monte Carlo maximum likelihood

estimation using the Partial Stepping technique of Hummel et al., (2012)) and MPLE.type

was set to "penalized" so that the Maximum Penalized Likelihood Estimate(MPLE) was

calculated using a bias-reduced method.

For each model set for each individual monthly network, we used Aikaike Information

Criterion (AIC) to compare the AIC of 49 candidate models nested within the full model, and

keep only the model that most parsimoniously explained the movements of the individual

during the month considered. Any models with convergence issues were discarded.

We then re-fitted all selected ERGMs for each combination of both model sets (*nodecov* and

absdiff) and individual monthly networks. From these final models, we extracted model

estimates and their standard errors, and also MCMC diagnostics (Gewecke's convergence

diagnostic and $\hat{R}$) to ensure convergence.

To summarize the results incorporating each selected models, we extracted parameter

estimates and associated standard errors from selected variables from each selected models,

and followed a meta-analysis procedure in which each sample (monthly-individual

movement network) was treated as a single 'study'. Effect sizes were calculated using an

inverse-variance weighting meta-analysis for each sample and grouped by variables using a

sub-grouping analysis. This allows to test if differences in effect sizes exist only due to

sampling error, or because of true differences in the effect sizes. This procedure was made for

the overall samples, as well as separating by sex and by season. The meta-analysis was

conducted using the package '*meta*' in R (Schwarzer, 2007).

Simulation methods

To ensure the correct interpretation of our model results and check for potential limitations of

the ERGM methods applied to our dataset, we used a simulation approach. In short, we

simulated a series of individual movement trajectories that matched our empirically collected

data. We then fitted ERGMs in the same way we did for the empirical data.

We generated sets of 25 receiver locations (equivalent to the empirical data) positioned in 2D

space and generated three environmental properties for each receiver location termed

factorA, factorB and factorC. We then generated 21 simulation input parameter sets that

varied the effect of factorA on the probability of movements between locations while keeping

factorB and factorC fixed as having no effects on movement (Supplementary material 2

Table S2). We considered scenarios where movements were conducted according to a

gradient in factorA or occurred between similarly high or low factorA locations. For each

simulation input parameter set, we simulated the movements of 10 sharks. The number of

movements for each shark was drawn from a Poisson distribution with a mean of 23

(equivalent to the empirical dataset). Detailed methods are in Supplementary material 2.

RESULTS

Interpretations of model outputs from simulations

Our simulation study revealed ERGMs performed as expected when analysing movement

networks of a similar size and structure to our empirically-measured networks (see

Supplementary materials 2). Model estimates for the *nodecov* and *absdiff* models typically

accurately represented the presence of movements up a gradient or between similar locations,

respectively (Figure 2). The statistical power to detect true effects was limited for individual

networks, and there were slightly elevated false positive rates for the *absdiff* models.

However, effect size estimates were largely unbiased (both with and without true simulated

effects). The only exception was that *absdiff* models underestimated the tendency to move

between similar locations when there was also a gradient effect (Figure 2f). In addition,

absdiff models overestimated the tendency to move between similar locations when sharks

tended to start moving from high value locations and had a strong tendency to move up a

gradient (Figure 2i, although the latter would be expected if this resulted in sharks moving

only between high value locations). Collectively, these results indicate that collating

estimates from multiple models (as we do in our main analysis) will provide the most

informative results. Simulations also indicated that *nodeicov* and *nodeocov* models provided

closely correlated results leading to us using a single *nodecov* model in our empirical analysis

(see above).

Effects of abiotic conditions

Our models were designed to test whether sharks were moving between locations with

similar abiotic conditions (tested by the *absdiff* term with a negative coefficient indicating

movement between location with similar conditions, while a positive coefficient means that

an increase in the absolute differences of abiotic conditions increases the odds of movements)

and/or towards locations with higher/lower abiotic conditions (tested by the *nodecov* term, a

positive coefficient indicating higher values and a negative coefficient suggesting lower

values), specifically swell height, degree of turbidity and level of rainfall. Model results

indicated the environmental variables that were associated with movement network structure

(Figure 3). While high swell at a location was not significantly associated with more

movements (positive coefficients in the *nodecov* model), movements between locations that

differed considerably in swell height were much less frequent than movements between

locations with similar swells as demonstrated by negative values of the *absdiff* model.

Together, these outputs indicate that sharks move between locations characterized by similar

swell and not across a gradient of swell heights. The *nodecov* model revealed a negative

effect of turbidity on movements, indicating that more movements occurred through low

turbidity locations. Further, the *absdiff* model revealed that sharks were also more likely to

move between receiver locations that differed from each other in their turbidity. Collectively

these results suggest that low turbidity increased the frequency of movements between

receivers. Sharks also tended to move between receiver locations with similar and low

rainfall levels. The observed trends were more robustly represented for rain and turbidity than

swell as these variables were supported by more monthly networks and involved in a larger

number of individuals with statistically significant results (Supplementary material 1 Table

S2).

Effect of turtle densities

Our models could also test whether sharks were moving toward high or low densities of

turtles (tested by the *nodecov* term) and/or were remaining in habitats with similar densities

(tested by the *absdiff* term). Model outputs indicated that movements were more likely

between receiver locations with more similar turtle densities (negative estimate from *absdiff*

model). However, parameter estimates were small and the number of monthly networks and

individuals in which the coefficients were statistically significant were low (Supplementary

material 1 Table S2), indicating that the biological importance of turtle distribution densities

may be limited.

Effect of anthropogenic factors

Our models were also set up to test whether sharks were avoiding (tested by the *nodecov* term

with an expected negative coefficient) and/or remaining in areas of similar human activity

(tested by the *absdiff* term with an expected negative coefficient). Models indicated that there

were more movements among locations with high human activities (positive *nodecov* effect

and negative *absdiff* effect). However, parameter estimates from the models were close to

zero and the number of monthly networks and individuals in which the coefficients were

statistically significant were relatively high (Supplementary material 1 Table S2), suggesting

that the biological importance of anthropogenic factors such as human activities on bull shark

movements in the monitored area were relatively limited but consistent among sharks and

months.

Influence of conspecifics

Our models were also used to test whether sharks moved towards or avoided conspecifics

(tested by the *nodecov* term) and/or remained within areas with similar populations of

conspecifics (tested by the *absdiff* term). Receiver locations in which bull sharks were present

or more abundant were more strongly connected in the movement network. Outputs from

absdiff model also indicated that movements tended to occur between locations with similar

bull shark abundance, suggesting individuals were predominantly moving among a subset of

preferred locations. While controlling for the abundance effect, there was (overall) some

evidence for a negative effect of the residency time of sharks of the opposite sex, suggesting

some spatial segregation between the sexes, but this effect varied seasonally (see below). The

results were as robust, as the models describing responses to abiotic factors, characterized by

a relatively high number of monthly networks and individuals involved and consistency in

the direction of responses (positive or negative coefficients) across networks (Supplementary

material 1 Table S2).

Effect of inter-specific competition

We also used our models to test whether the presence of larger tiger sharks was influencing

bull shark movements by moving to/away from areas used by tiger sharks (tested by the

nodecov term) and/or moving between sites with similar numbers of tiger sharks (tested by

the *absdiff* term). There was weak evidence (due to wide confidence intervals and the small

number of times this parameter was selected in the top model, Supplementary material 1

Table S2) for the presence of tiger sharks affecting bull shark movement network structure.

Positive effects in *absdiff* model indicated individuals tended to move across gradients of

tiger shark occupancy, perhaps indicating active avoidance of tiger sharks, with some

additional weak evidence of reduced movements through locations with higher occupancy of

tiger sharks.

Sex differences in movement network structure

Analysing movement network structure for the two sexes independently revealed similar

broad trends as the overall movement network, except in the case of social factors and the

tiger shark occupancy (Figure 4). While females tended to move toward locations with high

residency times for other females, males showed a general pattern of avoiding locations with

high residency times of either sex (while controlling for overall bull shark abundance). The

weak overall evidence for reduced movements through locations with high tiger shark

occupancy was driven by divergent effects between females and males. Female movements

were directed towards locations with low tiger shark occupancy while male movements were

directed towards locations with high tiger shark occupancy, indicating sex differences in how

competitors influenced movement through receiver locations, although these effects have

high uncertainty around them.

Seasonal differences in movement network structure

The factors that explained movement network structure remained similar between winter and

summer, with only interspecific competition and rainfall changing qualitatively (Figure 5).

While rainfall levels had effects on movements in both seasons, the pattern was less clear

during winter, which is expected as rainfall is a seasonally-driven factor. It appears there is a

shift in behaviour related to the presence of tiger sharks. While there was no apparent

movement pattern related to tiger shark occupancy during winter, there was a pattern of

movements towards locations with low occupancy of tiger shark in summer as demonstrated

by positive *absdiff* coefficients and negatives coefficient of *nodecov*.

DISCUSSION

We predicted that bull sharks would be attracted to high levels of turbidity, rainfall and swell.

We also expected that bull shark movements would be influenced by the distribution of

conspecifics and heterospecifics in the seascape, although we did not predict the direction of

these effects. We were thus interested whether these con- and heterospecific effects were as

important as abiotic variables in explaining movement patterns. As we hypothesized, if bull

shark movements were driven by abiotic parameters, our models revealed that social factors

were also important factors even if the reproducibility and consistency across the sampled

individuals and months were slightly smaller (Supplementary material 1 Table S2). We also

acknowledge that other potential important factors such as temperature or prey distribution

not included in our model could contribute to explain shark movement. Additionally, the

direction of response to abiotic variables were not always as predicted, indicating that the role

of physical variables in shaping movement network structure may be context dependent or

vary across spatial and temporal scales.

Effects of abiotic factors on movements

Our results indicate that environmental factors impact the movements of bull sharks,

however, while we expected to find positive effects of swell height, turbidity and rainfall on

shark movements as suggested by previous studies, we found opposite patterns. Bull sharks

were found to be mostly moving through areas with lower rainfall. While bull sharks are

known to respond to rainfall and freshwater runoff and are able to transit between freshwater

and marine habitats within hours or remain in low salinity areas for days (Niella et al., 2022;

Werry et al., 2018), selection of lower rainfall levels could be due to the spatial and temporal

resolution at which they react to such rapid environmental changes. In addition, bull sharks

may prefer to predominantly remain (at our scale of observation) in the same conditions of

rain, turbidity and salinity as rain levels are often strong and fast in tropical areas. Werry et

al. (2018) suggested that bull sharks are attracted to estuarine and associated nearshore areas

following high rainfall events in part due to increased prey availability while Niella et al.

(2022) suggested a more complex pattern with differences in response between the sexes.

Matich et al., (2020) also highlighted that during exceptional events bull sharks can avoid

high rainfall and high turbidity, moving away from sites with fast and strong perturbations.

The rainfall data entered in our models represent distinct watershed values that can

potentially incorporate multiple receivers, and can potentially explain why our models found

disproportionately strong effects of movements between (nearby) locations with similar

levels of rainfall. In addition, Werry et al. (2018) found a one-week lag in the response of

sharks to rainfall which could reflect changes in food availability and foraging effectiveness

with changes in salinity, a pattern that would be difficult to reveal with our models. Previous

studies that found strong behavioural responses of bull sharks to rainfall were conducted in

large estuarine system where rainfall levels and consequences (e.g. induced turbidity) could

contrast with the coral reef coast of Reunion Island. Finally, movements in response to

attractive effects of environmental factors (swell, turbidity and rainfall) that are ephemeral

and irregular may occur over shorter timescales and be difficult to detect with monthly data.

While we expected to find stronger seasonal differences because the variations in these

parameters are more important and persistent in summer than during winter, the lack of such

behavioural differences could be due to bull sharks being resident for less lengthy periods

during summer (Blaison et al., 2015). It is important to note that other potential factors that

we did not measure could also contribute significantly to movement decisions. For example,

salinity is known to be an important factor affecting movement and distribution of bull sharks

(Dwyer et al., 2020). Our results confirm that considering temporally dynamic environmental

variables in studies of animal movement is important because movement patterns reflect

dynamic interactions between animals and their physical environment, with environmental

heterogeneity driving movements of individuals and allowing them to remain in optimal

environmental conditions (Avgar et al., 2013; Mueller et al., 2011; Schick et al., 2008).

Effects of turtle density on movements

Our study did not find any clear effect of turtle density on bull shark movements.

While we used available data on turtle density to investigate its influence on movement

patterns of bull sharks in this study based on local people's perception, we acknowledge that

turtles are not the main prey of bull sharks and remain opportunistic items (Cliff & Dudley,

1991). Bull sharks are known to feed mainly on teleost fish (Trystram et al., 2016) and future

studies could integrate dynamic species-specific fishery data or parallel underwater surveys

to investigate the role of abundance and composition of fish communities in driving predator

movements. Indeed, resource availability and heterogeneity are important drivers of animal

movement, especially for predators (i.e. prey abundance and distribution). Finding prey

requires a predator to make adaptive decisions about which movement patterns to adopt to

feed most profitably given a particular prey distribution (Riotte-Lambert & Matthiopoulos,

2020; Sims et al., 2006). Thus, interpreting predator movements within a prey landscape may

provide a clearer picture of why certain habitats are selected over others.

Effects of human densities on movements

While we detected a tendency for more movements through locations with high

human use, the effect sizes were small suggesting human activities did not strongly influence

movement patterns. This is not surprising as nautical human activities and density are spread

fairly evenly along the west coast of the island where the network of receivers was deployed.

In addition, it is now recognized that bull sharks can adapt to urbanized areas and do not

especially avoid these high human density areas and their activities (Hammerschlag et al.,

2022; Werry et al., 2012). The majority of large-bodied terrestrial carnivores tend to avoid

high human densities and activities (Tucker et al., 2018, 2023). COVID lockdowns revealed

the same trend in the marine realm with fish communities responding positively to the

absence of human activities (e.g., fish biomass increasing and predatory species increasing

usage of shallow habitat during tourism lockdown; Weng et al., 2023). However, it is less

clear with sharks as they have previously been shown to both respond positively to cessation

of human activities (Afonso, 2024) and adapt to high human densities (Hammerschlag et al.,

2022). The limited suitable habitat available around the island may also prevent bull sharks

from avoiding most human activities along the coast. In addition, human activities are

concentrated on the west coast of the island and overlap with other factors that could explain

bull shark habitat preferences, such as the presence of a marine protected area (Soria et al.,

2019). Further investigations are therefore necessary to better understand the influence of

human activities on bull shark behaviour.

Effects of social environment on movements

We found that social factors sometimes had similar effects on movement networks to

abiotic variables. While the bull sharks have been found to form occasional aggregations

around fish farms (Loiseau et al., 2016) or at artificial provisioning sites (Bouveroux et al.,

2021), they are not typically recognized as displaying collective behaviours. In addition,

strong patterns of sexual and ontogenetic spatial segregation have been found in our study

population (Mourier et al., 2021). However, we found that more movements occurred

through bull shark hotspots. When focusing on the behaviour of each sex, we showed that

males were less social and tended to avoid other sharks regardless of their sex. Contrastingly,

females' movements occurred between locations where other females spent a lot of time but

avoided areas heavily used by males. This confirmed previous findings that females show

stronger patterns of residency providing the opportunity to co-occur with other females, while

males favour roaming behaviour (Mourier et al., 2021). This gregarious behaviour of females

could also suggest the presence of mating arenas where females gather to choose transient

males allowing them to avoid male harassment during the mating period. Further research

investigating the spatio-temporal interactions between bull sharks is required to better

understand their aggregative and avoidance behaviour. Our results thus confirm that an

individual's movements are not solely driven by abiotic factors. Individuals share space with

other conspecifics, linking spatial and social processes (Albery et al., 2021; Webber et al.,

2023). For example, the spatial distribution of resources inherently drives the proximity of

individuals through foraging aggregations, influencing many forms of social interaction

(Macdonald, 1983). As such, the distribution of individuals of a species in space generates

population structure and will influence movement decisions at the individual level. In fact,

regardless of whether a species is perceived to be solitary or social, individuals will have to

decide whether to join or avoid other conspecifics distributed unevenly in space. Such

decisions will also depend on social processes driven by individual phenotypes that alter

social decisions (e.g. size, sex, or genetic relatedness).

Effects of interspecific competition on movements

In our study we tested the influence of another large predator, the tiger shark, on the

movements of bull sharks. Our results indicated the presence of patterns of avoidance

between bull and tiger sharks. Bull shark movements were affected by the presence of tiger

sharks, with female bull sharks directing their movements towards locations with lower tiger

shark occupancy while male sharks showed the opposite pattern. The tiger shark is an apex

predator that is larger than the bull shark and potentially dominant. It is therefore plausible

that bull sharks avoid interactions with tiger sharks. Female bull sharks, being larger than

males (Hoarau et al., 2021), could also compete with tiger sharks. Therefore, one plausible

explanation to this pattern is spatial and foraging niche separation between both species

(Niella et al., 2021). We previously highlighted that male and female bull sharks displayed

patterns of spatial segregation (Mourier et al., 2021; Niella et al., 2022), which could also

explain the difference in interaction opportunities with tiger sharks based on habitat

utilization and movement patterns. These patterns of interference interactions are relatively

common in large sharks, as competitive species generally show spatial (Papastamatiou et al.,

2018), temporal (Lear et al., 2021; Séguigne et al., 2023) and trophic niche (Matich et al.,

2017) partitioning to limit the negative effect of competition and promote co-existence of

predators. Similar avoidance strategies are found between competing carnivores in

terrestrial ecosystems. Indeed, lions (*Panthera leo*) usually remain in areas rich in prey, while

subordinated carnivores like leopard (*P. pardus*) and cheetah (*Acinonyx jubatus*) overlap with

the home range of lions but use fine-scaled avoidance behaviours, and small species like

African wild dog (*Lycaon pictus*) employ multiple tactics to avoid all other competitors

(Vanak et al., 2013). Coexistence of multiple carnivore species is typically explained by

dietary niche separation as a consequence of avoiding intraguild competition. Another

potential explanation is the avoidance of tiger sharks by females during summer

corresponding to the parturition season (Pirog et al., 2019) in order to avoid tiger shark

predation on bull shark neonates.

Although our results are to be treated with caution because we only studied a small

number of individuals, and only a subset of these individuals responded to tiger sharks (Table

2), the direction of response was consistent. Our results thus complement previous studies

suggesting that interactions among large predators involve a complex interplay of

competition and predation, as large carnivores can suppress populations of smaller carnivores

through direct predation, resource competition, or via other forms of interference

competition. This may result in spatial and/or temporal avoidance, reductions in the density

of the subordinate species, or even competitive exclusion from certain habitats (Berger &

Gese, 2007; Linnell & Strand, 2000; Prugh et al., 2009; Prugh & Sivy, 2020).

Conclusion

To investigate the drivers of movements of a marine predator, we used a network-

based approach of movement between fixed stations and employed Exponential Random

Graph Model in a way that could account for the dynamical nature of site attributes visited by

individual animals within a flexible framework that could be extended to test hypotheses

related to the structure of the network itself (extending beyond what is easily possible using

more conventional statistical approaches). While this modelling framework performed well in

our study, it could be refined further to test similar research questions at different spatial and

temporal scales. For example, one promising model is the separable temporal exponential-

family random graph model (tERGM), which treats the formation and dissolution of ties in

parallel at each time step as independent ERGMs (Carnegie et al., 2015; Krivitsky &

Handcock, 2014) and can provide a more realistic view of movement networks (e.g.,

including consecutive individual monthly or daily networks). This would make it possible to

better integrate the temporal interdependency of environmental changes in explaining animal

movement patterns.

Our results fit into a broader picture illustrating that animal movements arise from

complex interactions of individuals with their physical environment as well as with both

surrounding conspecific and heterospecific individuals with various level of consistency. By

including a diverse set of variables that may influence bull shark movements within our

analyses, we demonstrate that to fully explain animal movement patterns requires the

incorporation of multiple variables associated with environmental heterogeneity, human

footprint and the distribution of individuals in space, both of the same species and other

members of the community. A main finding was that, although environmental conditions

were important factors influencing movement of bull sharks, interactions with other

individuals in their surroundings was also important to consider, opening new perspectives at

the socio-spatial interface for these marine predators.

Author contributions

Johann Mourier: Conceptualization (Equal); Formal analysis (Equal); Investigation (Equal);

Methodology (Equal); Software (Equal); Validation (Equal); Visualization (Equal); Writing –

original draft (Lead); Writing – review & editing (Equal). **Marc Soria**: Conceptualization

(Equal); Data curation (Equal); Funding acquisition (Equal); Investigation (Equal);

Methodology (Equal); Project administration (Equal); Resources (Equal); Supervision

(Equal); Validation (Equal); Writing – review & editing (Equal). **Matthew Silk**: Formal

analysis (Equal); Investigation (Equal); Methodology (Equal); Software (Equal); Validation

(Equal); Visualization (Equal); Writing – review & editing (Equal). **Angelique Demichelis**:

Formal analysis (Equal); Investigation (Equal); Software (Equal); Writing – review & editing

(Equal). **Laurent Dagorn**: Conceptualization (Equal); Investigation (Equal); Validation

(Equal); Writing – review & editing (Equal). **Tarek Hattab**: Conceptualization (Equal);

Investigation (Equal); Methodology (Equal); Supervision (Equal); Validation (Equal);

Writing – review & editing (Equal).

Acknowledgments

We are grateful to the members of the institutions and associations involved in the CHARC

program (Connaissance de l'HAbitat des Requins Côtiers de la Réunion): IRD, CRPMEM,

CRESSM, University of Reunion Island, Globice, Kélonia, ARVAM, Squal'Idées, RNMR,

and IFREMER as well as the fishermen and volunteers who assisted with the shark tagging

and made our work possible. This study received financial support from the European Union

(convention FEDER ref. 2012-dossier Presage no. 33021), the Ministère de la Transition

écologique et Solidaire (BOP 113 n°2012/03), and the Regional Council of Reunion Island

(POLENV no. 20120257). MJS received funding from the European Union's Horizon 2020

research and innovation programme under the Marie Sklodowska-Curie grant agreement No

101023948. All the fieldwork and protocols of handling and tagging of sharks were approved

by the Ethics Committee (n° 114) for the CYROI (Cyclotron Réunion Océan Indien).

Procedures were adapted to minimize stress on animals and avoid mortality (more details in

Supplementary material 1). All operations were carried out or supervised by scientists with a

certificate in animal experimentation and a certificate in experimental surgery (Oniris, Ecole

Nationale Vétérinaire de Nantes).

Data availability

Data and R codes used in this study are available at

SEANOE: https://doi.org/10.17882/99080

Conflict of interest statement

All authors declare that they have no conflict of interest.

References

Afonso, A. S. (2024). Humans influence shark behavior: Evidence from the COVID-19

   lockdown. *Ocean & Coastal Management*, *248*, 106965.

   https://doi.org/10.1016/j.ocecoaman.2023.106965

Albery, G. F., Morris, A., Morris, S., Pemberton, J. M., Clutton-Brock, T. H., Nussey, D. H.,

   & Firth, J. A. (2021). Multiple spatial behaviours govern social network positions in a

   wild ungulate. *Ecology Letters*, *24*(4), 676–686. https://doi.org/10.1111/ele.13684

Atwood, T. C., Gese, E. M., & Kunkel, K. E. (2009). Spatial Partitioning of Predation Risk in

   a Multiple Predator-Multiple Prey System. *The Journal of Wildlife Management*, *73*(6),

   876–884. https://doi.org/10.2193/2008-325

Avgar, T., Mosser, A., Brown, G. S., & Fryxell, J. M. (2013). Environmental and individual

   drivers of animal movement patterns across a wide geographical gradient. *Journal of*

   Animal Ecology, *82*(1), 96–106. https://doi.org/10.1111/j.1365-2656.2012.02035.x

Berger, K. M., & Gese, E. M. (2007). Does interference competition with wolves limit the

   distribution and abundance of coyotes? *Journal of Animal Ecology*, *76*(6), 1075–1085.

   https://doi.org/10.1111/j.1365-2656.2007.01287.x

Blaison, A., Jaquemet, S., Guyomard, D., Vangrevelynghe, G., Gazzo, T., Cliff, G., Cotel, P.,

   & Soria, M. (2015). Seasonal variability of bull and tiger shark presence on the west

   coast of Reunion Island, western Indian Ocean. *African Journal of Marine Science*,

   37(2), 199–208. https://doi.org/10.2989/1814232X.2015.1050453

Bouveroux, T., Loiseau, N., Barnett, A., Marosi, N. D., & Brunnschweiler, J. M. (2021).

   Companions and Casual Acquaintances: The Nature of Associations Among Bull

   Sharks at a Shark Feeding Site in Fiji. *Frontiers in Marine Science*, *8*.

   https://doi.org/10.3389/fmars.2021.678074

Brunnschweiler, J. M., Queiroz, N., & Sims, D. W. (2010). Oceans apart? Short-term

688  movements and behaviour of adult bull sharks Carcharhinus leucas in Atlantic and

689  Pacific Oceans determined from pop-off satellite archival tagging. *Journal of Fish*

690  *Biology*, *77*(6), 1343–1358. https://doi.org/10.1111/j.1095-8649.2010.02757.x

691 Capello, M., Robert, M., Soria, M., Potin, G., Itano, D., Holland, K., Deneubourg, J.-L., &

692  Dagorn, L. (2015). A Methodological Framework to Estimate the Site Fidelity of

693  Tagged Animals Using Passive Acoustic Telemetry. *PLOS ONE*, *10*(8), e0134002.

694  https://doi.org/10.1371/journal.pone.0134002

695 Carnegie, N. B., Krivitsky, P. N., Hunter, D. R., & Goodreau, S. M. (2015). An

696  Approximation Method for Improving Dynamic Network Model Fitting. *Journal of*

697  *Computational and Graphical Statistics*, *24*(2), 502–519.

698  https://doi.org/10.1080/10618600.2014.903087

699 Cliff, G., & Dudley, S. F. J. (1991). Sharks caught in the protective gill nets off Natal, South

700  Africa. 4. The bull shark Carcharhinus leucas Valenciennes. *South African Journal of*

701  *Marine Science*, *10*(1), 253–270. https://doi.org/10.2989/02577619109504636

702 Couriot, O., Hewison, A. J. M., Saïd, S., Cagnacci, F., Chamaillé-Jammes, S., Linnell, J. D.

703  C., Mysterud, A., Peters, W., Urbano, F., Heurich, M., Kjellander, P., Nicoloso, S.,

704  Berger, A., Sustr, P., Kroeschel, M., Soennichsen, L., Sandfort, R., Gehr, B., &

705  Morellet, N. (2018). Truly sedentary? The multi-range tactic as a response to resource

706  heterogeneity and unpredictability in a large herbivore. *Oecologia*, *187*(1), 47–60.

707  https://doi.org/10.1007/s00442-018-4131-5

708 Croft, D. P., Madden, J. R., Franks, D. W., & James, R. (2011). Hypothesis testing in animal

709  social networks. *Trends in Ecology & Evolution*, *26*(10), 502–507.

710  https://doi.org/10.1016/j.tree.2011.05.012

711 Daly, R., Smale, M. J., Cowley, P. D., & Froneman, P. W. (2014). Residency Patterns and

712  Migration Dynamics of Adult Bull Sharks (Carcharhinus leucas) on the East Coast of

Southern Africa. *PLOS ONE, 9*(10), e109357.

https://doi.org/10.1371/journal.pone.0109357

Dey, C. J., & Quinn, J. S. (2014). Individual attributes and self-organizational processes

affect dominance network structure in pukeko. *Behavioral Ecology, 25*(6), 1402–1408.

https://doi.org/10.1093/beheco/aru138

Doherty, T. S., Hays, G. C., & Driscoll, D. A. (2021). Human disturbance causes widespread

disruption of animal movement. *Nature Ecology & Evolution*, *5*(4), 513–519.

https://doi.org/10.1038/s41559-020-01380-1

Dwyer, R. G., Campbell, H. A., Cramp, R. L., Burke, C. L., Micheli-Campbell, M. A.,

Pillans, R. D., Lyon, B. J., & Franklin, C. E. (2020). Niche partitioning between river

shark species is driven by seasonal fluctuations in environmental salinity. *Functional*

Ecology, *34*(10), 2170–2185. https://doi.org/10.1111/1365-2435.13626

Espinoza, M., Heupel, M. R., Tobin, A. J., & Simpfendorfer, C. A. (2016). Evidence of

Partial Migration in a Large Coastal Predator: Opportunistic Foraging and

Reproduction as Key Drivers? *PLOS ONE*, *11*(2), e0147608.

https://doi.org/10.1371/journal.pone.0147608

Fletcher, R. J., Acevedo, M. A., Reichert, B. E., Pias, K. E., & Kitchens, W. M. (2011).

Social network models predict movement and connectivity in ecological landscapes.

Proceedings of the National Academy of Sciences, *108*(48), 19282–19287.

https://doi.org/10.1073/pnas.1107549108

Gilbert, N. A., McGinn, K. A., Nunes, L. A., Shipley, A. A., Bernath-Plaisted, J., Clare, J. D.

J., Murphy, P. W., Keyser, S. R., Thompson, K. L., Nelson, S. B. M., Cohen, J. M.,

Widick, I. V., Bartel, S. L., Orrock, J. L., & Zuckerberg, B. (2023). Daily activity

timing in the Anthropocene. *Trends in Ecology & Evolution*, *38*(4), 324–336.

https://doi.org/10.1016/j.tree.2022.10.008

Gilbert, N. A., Stenglein, J. L., Pauli, J. N., & Zuckerberg, B. (2022). Human disturbance

compresses the spatiotemporal niche. *Proceedings of the National Academy of*

Sciences, *119*(52), e2206339119. https://doi.org/10.1073/pnas.2206339119

Giuggioli, L., & Kenkre, V. M. (2014). Consequences of animal interactions on their

dynamics: Emergence of home ranges and territoriality. *Movement Ecology*, *2*(1), 20.

https://doi.org/10.1186/s40462-014-0020-7

Hammerschlag, N., Gutowsky, L. F. G., Rider, M. J., Roemer, R., & Gallagher, A. J. (2022).

Urban sharks: Residency patterns of marine top predators in relation to a coastal

metropolis. *Marine Ecology Progress Series*, *691*, 1–17.

https://doi.org/10.3354/meps14086

He, P., Maldonado-Chaparro, A. A., & Farine, D. R. (2019). The role of habitat configuration

in shaping social structure: A gap in studies of animal social complexity. *Behavioral*

Ecology and Sociobiology, *73*(1), 9. https://doi.org/10.1007/s00265-018-2602-7

Heupel, M. R., Simpfendorfer, C. A., Espinoza, M., Smoothey, A. F., Tobin, A., &

Peddemors, V. (2015). Conservation challenges of sharks with continental scale

migrations. *Frontiers in Marine Science*, *2*. https://doi.org/10.3389/fmars.2015.00012

Hoarau, F., Darnaude, A., Poirout, T., Jannel, L.-A., Labonne, M., & Jaquemet, S. (2021).

Age and growth of the bull shark () around Reunion Island, South West Indian Ocean.

Journal of Fish Biology, *99*(3), 1087–1099. https://doi.org/10.1111/jfb.14813

Hummel, R. M., Hunter, D. R., & Handcock, M. S. (2012). Improving Simulation-Based

Algorithms for Fitting ERGMs. *Journal of Computational and Graphical Statistics*,

21(4), 920–939. https://doi.org/10.1080/10618600.2012.679224

Hunter, D. R., Handcock, M. S., Butts, C. T., Goodreau, S. M., & Morris, M. (2008). ergm: A

Package to Fit, Simulate and Diagnose Exponential-Family Models for Networks.

Journal of Statistical Software, *24*(1), Article 1. https://doi.org/10.18637/jss.v024.i03

Jacoby, D. M. P., & Freeman, R. (2016). Emerging Network-Based Tools in Movement

Ecology. *Trends in Ecology & Evolution*, *31*(4), 301–314.

https://doi.org/10.1016/j.tree.2016.01.011

Johnson, A. R., Wiens, J. A., Milne, B. T., & Crist, T. O. (1992). Animal movements and

population dynamics in heterogeneous landscapes. *Landscape Ecology*, *7*(1), 63–75.

https://doi.org/10.1007/BF02573958

Krause, J., & Ruxton, G. D. (2002). *Living in Groups*. OUP Oxford.

Krivitsky, P. N., & Handcock, M. S. (2014). A separable model for dynamic networks.

Journal of the Royal Statistical Society: Series B (Statistical Methodology), *76*(1), 29–

46. https://doi.org/10.1111/rssb.12014

Kuefler, D., Avgar, T., & Fryxell, J. M. (2012). Rotifer population spread in relation to food,

density and predation risk in an experimental system. *Journal of Animal Ecology*,

81(2), 323–329. https://doi.org/10.1111/j.1365-2656.2011.01917.x

Lear, K. O., Whitney, N. M., Morris, J. J., & Gleiss, A. C. (2021). Temporal niche

partitioning as a novel mechanism promoting co-existence of sympatric predators in

marine systems. *Proceedings of the Royal Society B: Biological Sciences*, *288*(1954),

20210816. https://doi.org/10.1098/rspb.2021.0816

Lee, K. A., Smoothey, A. F., Harcourt, R. G., Roughan, M., Butcher, P. A., & Peddemors, V.

M. (2019). Environmental drivers of abundance and residency of a large migratory

shark, Carcharhinus leucas, inshore of a dynamic western boundary current. *Marine*

Ecology Progress Series, *622*, 121–137. https://doi.org/10.3354/meps13052

Linnell, J. D. C., & Strand, O. (2000). Interference interactions, co-existence and

conservation of mammalian carnivores. *Diversity and Distributions*, *6*(4), 169–176.

https://doi.org/10.1046/j.1472-4642.2000.00069.x

Loiseau, N., Kiszka, J. J., Bouveroux, T., Heithaus, M. R., Soria, M., & Chabanet, P. (2016).

Using an unbaited stationary video system to investigate the behaviour and interactions of bull sharks Carcharhinus leucas under an aquaculture farm. *African Journal of Marine Science*, *38*(1), 73–79. https://doi.org/10.2989/1814232X.2016.1156578

López-Calderón, C., Martín-Vélez, V., Blas, J., Höfle, U., Sánchez, M. I., Flack, A., Fiedler, W., Wikelski, M., & Green, A. J. (2023). White stork movements reveal the ecological connectivity between landfills and different habitats. *Movement Ecology*, *11*(1), 18. https://doi.org/10.1186/s40462-023-00380-7

Lubitz, N., Daly, R., Filmalter, J. D., Sheaves, M., Cowley, P. D., Naesje, T. F., & Barnett, A. (2023). Context drives movement patterns in a mobile marine predator. *Movement Ecology*, *11*(1), 28. https://doi.org/10.1186/s40462-023-00390-5

Macdonald, D. W. (1983). The ecology of carnivore social behaviour. *Nature*, *301*(5899), Article 5899. https://doi.org/10.1038/301379a0

Matich, P., Kiszka, J. J., Mourier, J., Planes, S., & Heithaus, M. R. (2017). Species co-occurrence affects the trophic interactions of two juvenile reef shark species in tropical lagoon nurseries in Moorea (French Polynesia). *Marine Environmental Research*, *127*(Supplement C), 84–91. https://doi.org/10.1016/j.marenvres.2017.03.010

Matich, P., Strickland, B. A., & Heithaus, M. R. (2020). Long-term monitoring provides insight into estuarine top predator (Carcharhinus leucas) resilience following an extreme weather event. *Marine Ecology Progress Series*, *639*, 169–183. https://doi.org/10.3354/meps13269

Morales, J. M., Moorcroft, P. R., Matthiopoulos, J., Frair, J. L., Kie, J. G., Powell, R. A., Merrill, E. H., & Haydon, D. T. (2010). Building the bridge between animal movement and population dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2289–2301. https://doi.org/10.1098/rstb.2010.0082

Mourier, J., Lédée, E. J. I., & Jacoby, D. M. P. (2019). A multilayer perspective for inferring

spatial and social functioning in animal movement networks. *bioRxiv*, 749085.

https://doi.org/10.1101/749085

Mourier, J., Soria, M., Blaison, A., Simier, M., Certain, G., Demichelis, A., & Hattab, T.

(2021). Dynamic use of coastal areas by bull sharks and the conciliation of

conservation and management of negative human–wildlife interactions. *Aquatic*

Conservation: Marine and Freshwater Ecosystems, *31*(10), 2926–2937.

https://doi.org/10.1002/aqc.3674

Mueller, T., Olson, K. A., Dressler, G., Leimgruber, P., Fuller, T. K., Nicolson, C., Novaro,

A. J., Bolgeri, M. J., Wattles, D., DeStefano, S., Calabrese, J. M., & Fagan, W. F.

(2011). How landscape dynamics link individual- to population-level movement

patterns: A multispecies comparison of ungulate relocation data. *Global Ecology and*

Biogeography, *20*(5), 683–694. https://doi.org/10.1111/j.1466-8238.2010.00638.x

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E.

(2008). A movement ecology paradigm for unifying organismal movement research.

Proceedings of the National Academy of Sciences, *105*(49), 19052–19059.

https://doi.org/10.1073/pnas.0800375105

Niella, Y., Smoothey, A. F., Taylor, M. D., Peddemors, V. M., & Harcourt, R. (2022).

Environmental Drivers of Fine-Scale Predator and Prey Spatial Dynamics in Sydney

Harbour, Australia, and Adjacent Coastal Waters. *Estuaries and Coasts*, *45*(5), 1465–

1479. https://doi.org/10.1007/s12237-021-01020-2

Niella, Y., Wiefels, A., Almeida, U., Jaquemet, S., Lagabrielle, E., Harcourt, R., Peddemors,

V., & Guyomard, D. (2021). Dynamics of marine predators off an oceanic island and

implications for management of a preventative shark fishing program. *Marine Biology*,

168(4), 42. https://doi.org/10.1007/s00227-021-03852-9

Ohta, I., & Kakuma, S. (2005). Periodic behavior and residence time of yellowfin and bigeye

tuna associated with fish aggregating devices around Okinawa Islands, as identified

with automated listening stations. *Marine Biology*, *146*(3), 581–594.

https://doi.org/10.1007/s00227-004-1456-x

Papastamatiou, Y. P., Bodey, T. W., Friedlander, A. M., Lowe, C. G., Bradley, D., Weng, K.,

Priestley, V., & Caselle, J. E. (2018). Spatial separation without territoriality in shark

communities. *Oikos*, *127*(6), 767–779. https://doi.org/10.1111/oik.04289

Pirog, A., Magalon, H., Poirout, T., & Jaquemet, S. (2019). Reproductive biology, multiple

paternity and polyandry of the bull shark Carcharhinus leucas. *Journal of Fish Biology*,

95(5), 1195–1206. https://doi.org/10.1111/jfb.14118

Prugh, L. R., & Sivy, K. J. (2020). Enemies with benefits: Integrating positive and negative

interactions among terrestrial carnivores. *Ecology Letters*, *23*(5), 902–918.

https://doi.org/10.1111/ele.13489

Prugh, L. R., Stoner, C. J., Epps, C. W., Bean, W. T., Ripple, W. J., Laliberte, A. S., &

Brashares, J. S. (2009). The Rise of the Mesopredator. *BioScience*, *59*(9), 779–791.

https://doi.org/10.1525/bio.2009.59.9.9

Pyke, G. H., Pulliam, H. R., & Charnov, E. L. (1977). Optimal Foraging: A Selective Review

of Theory and Tests. *The Quarterly Review of Biology*, *52*(2), 137–154.

Riotte-Lambert, L., & Matthiopoulos, J. (2020). Environmental Predictability as a Cause and

Consequence of Animal Movement. *Trends in Ecology & Evolution*, *35*(2), 163–174.

https://doi.org/10.1016/j.tree.2019.09.009

Robins, G., Pattison, P., Kalish, Y., & Lusher, D. (2007). An introduction to exponential

random graph (p*) models for social networks. *Social Networks*, *29*(2), 173–191.

https://doi.org/10.1016/j.socnet.2006.08.002

Robins, G., Snijders, T., Wang, P., Handcock, M., & Pattison, P. (2007). Recent

developments in exponential random graph (p*) models for social networks. *Social*

Networks, *29*(2), 192–215. https://doi.org/10.1016/j.socnet.2006.08.003

Schick, R. S., Loarie, S. R., Colchero, F., Best, B. D., Boustany, A., Conde, D. A., Halpin, P.

N., Joppa, L. N., McClellan, C. M., & Clark, J. S. (2008). Understanding movement

data and movement processes: Current and emerging directions. *Ecology Letters*,

11(12), 1338–1350. https://doi.org/10.1111/j.1461-0248.2008.01249.x

Schwarzer, G. R. (2007). An R package for meta-analysis. *R News*, *7*, 40–45.

Séguigne, C., Mourier, J., Clua, É., Buray, N., & Planes, S. (2023). Citizen science provides

valuable data to evaluate elasmobranch diversity and trends throughout the French

Polynesia's shark sanctuary. *PLOS ONE*, *18*(3), e0282837.

https://doi.org/10.1371/journal.pone.0282837

Sims, D. W., Witt, M. J., Richardson, A. J., Southall, E. J., & Metcalfe, J. D. (2006).

Encounter success of free-ranging marine predator movements across a dynamic prey

landscape. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1591), 1195–

1201. https://doi.org/10.1098/rspb.2005.3444

Snijders, T. A. B., Pattison, P. E., Robins, G. L., & Handcock, M. S. (2006). New

Specifications for Exponential Random Graph Models. *Sociological Methodology*,

36(1), 99–153. https://doi.org/10.1111/j.1467-9531.2006.00176.x

Soria, M., Heithaus, M. R., Blaison, A., Crochelet, E., Forget, F., & Chabanet, P. (2019).

Residency and spatial distribution of bull sharks Carcharhinus leucas in and around

Reunion Island marine protected area. *Marine Ecology Progress Series*, *630*, 101–113.

https://doi.org/10.3354/meps13139

Trystram, C., Rogers, K. M., Soria, M., & Jaquemet, S. (2016). Feeding patterns of two

sympatric shark predators in coastal ecosystems of an oceanic island. *Canadian Journal*

of Fisheries and Aquatic Sciences, *74*(2), 216–227. https://doi.org/10.1139/cjfas-2016-

0105

Tucker, M. A., Böhning-Gaese, K., Fagan, W. F., Fryxell, J. M., Moorter, B. V., Alberts, S.

C., Ali, A. H., Allen, A. M., Attias, N., Avgar, T., Bartlam-Brooks, H., Bayarbaatar, B.,

Belant, J. L., Bertassoni, A., Beyer, D., Bidner, L., Beest, F. M. van, Blake, S., Blaum,

N., … Mueller, T. (2018). Moving in the Anthropocene: Global reductions in terrestrial

mammalian movements. *Science*, *359*(6374), 466–469.

https://doi.org/10.1126/science.aam9712

Tucker, M. A., Schipper, A. M., Adams, T. S. F., Attias, N., Avgar, T., Babic, N. L., Barker,

K. J., Bastille-Rousseau, G., Behr, D. M., Belant, J. L., Beyer, D. E., Blaum, N.,

Blount, J. D., Bockmühl, D., Pires Boulhosa, R. L., Brown, M. B., Buuveibaatar, B.,

Cagnacci, F., Calabrese, J. M., … Mueller, T. (2023). Behavioral responses of

terrestrial mammals to COVID-19 lockdowns. *Science*, *380*(6649), 1059–1064.

https://doi.org/10.1126/science.abo6499

Vanak, A. T., Fortin, D., Thaker, M., Ogden, M., Owen, C., Greatwood, S., & Slotow, R.

(2013). Moving to stay in place: Behavioral mechanisms for coexistence of African

large carnivores. *Ecology*, *94*(11), 2619–2631. https://doi.org/10.1890/13-0217.1

Wearmouth, V. J., & Sims, D. W. (2008). Chapter 2 Sexual Segregation in Marine Fish,

Reptiles, Birds and Mammals: Behaviour Patterns, Mechanisms and Conservation

Implications. In *Advances in Marine Biology* (Vol. 54, pp. 107–170). Academic Press.

https://doi.org/10.1016/S0065-2881(08)00002-3

Webber, Q. M. R., Albery, G. F., Farine, D. R., Pinter-Wollman, N., Sharma, N., Spiegel, O.,

Vander Wal, E., & Manlove, K. (2023). Behavioural ecology at the spatial–social

interface. *Biological Reviews*, *98*(3), 868–886. https://doi.org/10.1111/brv.12934

Weng, K. C., Friedlander, A. M., Gajdzik, L., Goodell, W., & Sparks, R. T. (2023).

Decreased tourism during the COVID-19 pandemic positively affects reef fish in a high

use marine protected area. *PLOS ONE*, *18*(4), e0283683.

https://doi.org/10.1371/journal.pone.0283683

Werry, J. M., Lee, S. Y., Lemckert, C. J., & Otway, N. M. (2012). Natural or Artificial?

Habitat-Use by the Bull Shark, Carcharhinus leucas. *PLOS ONE*, *7*(11), e49796.

https://doi.org/10.1371/journal.pone.0049796

Werry, J. M., Sumpton, W., Otway, N. M., Lee, S. Y., Haig, J. A., & Mayer, D. G. (2018).

Rainfall and sea surface temperature: Key drivers for occurrence of bull shark,

Carcharhinus leucas, in beach areas. *Global Ecology and Conservation*, *15*, e00430.

https://doi.org/10.1016/j.gecco.2018.e00430

Table 1: Predictor parameters included in the Exponential Random Graph Models for bull

sharks monitored along the west coast of Reunion Island.

Variables	Category	Description	Range	Source
Occupancy Tiger sharks	Biological	Frequency of days at least a tiger shark was detected at the receiver for each month (%)	[0-0.409]	Current acoustic telemetry data
Same sex CRT	Biological	Total CRT of bull sharks of the same sex for each month (hour)	[0.033-118.425]	Current acoustic telemetry data
Opposite sex CRT	Biological	Total CRT of bull sharks of the opposite sex for each month (hour)	[0.033-118.425]	Current acoustic telemetry data
Bull Shark abundance	Biological	Number of bull sharks present at the listening station divided by the number of bull shark with an active tag for each month	[0-1.24]	Current acoustic telemetry data
Turtle density	Biological	Mean density of turtles in the area of the receiver from aerial survey (nb/km^2)	[0-15.873]	Aerial surveys
Turbidity	Environmental	Average monthly reflectance for each zone over the multiple images (%)	[4.101-8.886]	Satellite SPOT 4 & 5
Rain	Environmental	Mean rain fall for each watershed for each month (mm)	[0.008-25.460]	METEO-France
Swell	Environmental	Height of the swell near the listening station for each month (m)	[0.557-2.636]	AVISO portal, CANDHIS & METEOLAMER platform
Human activities	Anthropogenic	Cumulated number of human activities (boats, swimmers, nautical activities) for each month	[0-1276.5]	Aerial surveys
Geodist	Spatial	Distance between stations (km)	[0.9-120]	Current acoustic telemetry data

Figure legend

Figure 1: Cumulative number of movements across all 44 monthly individual movements used in the ERGM analyses concentrated on the western part of Reunion Island. Circles represent acoustic receivers. Inset map represents the location of Reunion Island and the area where the movements were recorded.

Figure 2: Interpretation of the model output for each scenario based on simulations: (a, b, c) movements up a gradient of FactorA (e.g. swell height), (d, e, f) movements between locations of low FactorA and (g, h, i) movement up a gradient and between locations of high FactorA. (a, d, g) Toy examples on the left and movement network for the first shark. (b, e, h) model estimates for *nodecov* and (c, f, i) model estimates for *absdiff*. The model estimates are for the three effects (highlighting the one where there is a true effect in red) and then for each of the 10 sharks simulated.

Figure 3: Summary of the outputs from the Exponential Random Graph Models from the 42 selected individual monthly movement networks. Effect sizes and their 95% confidence intervals pooled over variables are reported for the parameter estimate of terms *nodecov* and *absdiff*. Variables were grouped in several categories: (1) environmental factors, (2) turtle density, (3) anthropogenic factors, (4) social or intraspecific interactions, and (5) competitive or interspecific interactions. The variable "Geodistance" controls for the distance between receivers. Circle size is proportional to the number of models in which the coefficient of the term was significant and therefore selected.

Figure 4: Summary of the outputs from the Exponential Random Graph Models presented

separately for females (n = 10) and males (n = 3). Effect sizes and their 95% confidence

intervals pooled over variables are reported for the coefficient of terms *nodecov* and *absdiff*.

The variable "Geodistance" controls for the distance between receivers. Circles represent the

number of networks in which the parameter estimate of the term was statistically significant

and therefore selected.

Figure 5: Summary of the outputs from the Exponential Random Graph Models presented

separately for summer and winter. Effect sizes and their 95% confidence intervals pooled

over variables are reported for the coefficient of terms *nodecov* and *absdiff*. The variable

"Geodistance" controls for the distance between receivers. Circles represent the number of

networks in which the coefficient of the term was significant and therefore selected.