1

Knockout mutations of *Nicotiana benthamiana* defenses reveal the relative importance of acylsugars, nicotine, and a serine protease inhibitor in a natural setting

Boaz Negin, Fumin Wang, Hillary D. Fischer, and Georg Jander

Boyce Thompson Institute, Ithaca, NY, 14853

*Georg Jander

Email: gj32@cornell.edu

Keywords: *Nicotiana benthamiana*, nicotine, acylsugars, insect, natural setting, specialized metabolites, snails, invertebrates

Summary

Plants produce an immense diversity of defensive specialized metabolites. However, despite extensive functional characterization, the relative importance of different defensive compounds is rarely examined in natural settings. Here, we compare the efficacy of three *Nicotiana benthamiana* defensive compounds, nicotine, acylsugars, and a serine protease inhibitor, by growing plants with combinations of knockout mutations in a natural setting, quantifying invertebrate interactions, and comparing relative plant performance. Among the three tested compounds, acylsugars had the greatest defensive capacity, affecting aphids, leafhoppers, spiders, and flies. Nicotine mutants displayed increased leafhopper feeding and aphid colonization. Plants lacking both nicotine and acylsugars were more susceptible to flea beetles and thrips. By contrast, knockout of the serine protease inhibitor did not affect insect herbivory in the field. Complementary experiments under controlled laboratory conditions with caterpillars grasshoppers, and aphids confirmed results obtained in a natural setting. We conclude that the three metabolite groups collectively provide broad-spectrum protection to *N. benthamiana*. However, there is a gradient in their effects on the interacting invertebrates present in the field. Furthermore, we demonstrate that, even if individual metabolites do not have a measurable defensive benefit on their own, they can have an additive effect when combined with other defensive compounds.

Main Text

Introduction

Plants synthesize an enormous diversity of specialized metabolites, with estimates ranging from 200,000 to 1,000,000 unique compounds in the plant kingdom (Wang *et al.*, 2019). Any individual plant produces roughly 1,750 – 3,500 specialized metabolites with diverse functions (Pichersky & Lewinsohn, 2011). Many families of specialized metabolites, such as benzoxazinoids (Meihls *et al.*, 2013), cardiac glycosides (Agrawal *et al.*, 2012), and glucosinolates (Hopkins *et al.*, 2009), have well-documented functions in plant defense, whereas others are yet to be investigated. All plants produce multiple classes of specialized metabolites with differing defensive functions. For instance, *Erysimum cheiranthoides* (wormseed wallflower) accumulates both cardenolides and glucosinolates (Züst *et al.*, 2020) and *Nicotiana attenuata* (coyote tobacco) produces nicotine (Steppuhn *et al.*, 2004), chlorogenic acid (Demkura *et al.*, 2010), protease inhibitors (Zavala *et al.*, 2004) and volatile terpenes (Yang *et al.*, 2023), which each have a distinct protective role. Furthermore, the effects of the different defensive metabolites are not only additive, but also can involve more complex interactions. Examples

include a protective function only in the presence of additional metabolites (Steppuhn & Baldwin,

2007), reciprocal reduction in toxicity (Heiling *et al.*, 2022), or co-option by specialist insect

herbivores for their own protection (Reichstein *et al.*, 1968; Hoogshagen *et al.*, 2023). Therefore,

altering the production of several defensive compounds simultaneously and examining their

ecological impacts in a natural setting are essential for gaining insight into their functions and

interplays in plant defense.

Biosynthesis of nicotine, a pyridine alkaloid built from the conjugation of a pyridine ring

and a nitrogen containing methylpyrrolidine ring (Kaminski *et al.*, 2020), is restricted to *Nicotiana*

species (Negin & Jander, 2023). The function of nicotine in protecting plants against insect

herbivores has been extensively demonstrated under both laboratory (Yang & Guthrie, 1969; Parr

& Thurston, 1972) and field conditions (Steppuhn *et al.*, 2004). Nicotine functions primarily by

targeting nicotinic acetylcholine receptors in animal nervous systems (Millar & Denholm, 2007),

and investigations of tritrophic interactions exemplify the complex relation between this

specialized metabolite, insects that consume it, and their predators (Kumar *et al.*, 2014; Xu *et al.*,

2017).

Acylsugars are a diverse family of compounds with the common feature of a sugar core

that has several aliphatic chains of different lengths connected to it (Fan *et al.*, 2019). These

metabolites, which are secreted from glandular trichomes of plants in the Solanaceae and several

other families (Fobes *et al.*, 1985; Kroumova & Wagner, 2003; Schilmiller *et al.*, 2012), have an

inhibitory effect on insect herbivores in controlled insect-plant interaction studies, both *in planta*

(Van Dam & Hare, 1998; Hare, 2005; Luu *et al.*, 2017; Feng *et al.*, 2022) and *in vitro* (Puterka *et*

al., 2003). Several studies have identified acylsugar biosynthesis genes in tomato and other

Solanaceae (Schilmiller *et al.*, 2012, 2016; Lou *et al.*, 2021), paving the way for targeted

manipulation of these defenses.

Protease inhibitors are proteins that inhibit insect digestive enzymes, rendering the

consumed plant material less nutritious (Green & Ryan, 1972). These plant proteins, which are

induced in response to caterpillar feeding, jasmonate treatment, or wounding (Van Dam *et al.*,

2001), reduce caterpillar growth (Zavala *et al.*, 2004; Hu *et al.*, 2018), slow aphid development

(Carrillo *et al.*, 2011), and reduce attractiveness of plants to mirids (Zavala *et al.*, 2004). Field-

grown *Solanum nigrum* (black nightshade), in which expression of three serine protease inhibitors

was silenced, was more susceptible to both thrips and caterpillars (Hartl *et al.*, 2010).

Nicotiana benthamiana is an Australian tobacco species that adapted to desert conditions

millions of years ago (Bally *et al.*, 2018). This species is widely used for plant molecular biology

research, largely due to its extreme viral susceptibility (Jassbi *et al.*, 2017), amenability for

transient gene expression assays (Bally *et al.*, 2015), and ease of heritable gene editing (Ellison

et al., 2020). By contrast, very few laboratory studies involve the investigation of insect

interactions in this species, likely due to the high level of endogenous herbivore resistance.

In the current study, we utilized existing genome-edited *N. benthamiana* plants, newly-

generated mutants, and mutant combinations to examine the function of three representative

defensive compounds, nicotine, acylsugars, and a serine protease inhibitor, in both a natural

setting and under controlled laboratory conditions. We found, that acylsugars had the strongest

effect, nicotine had a comparatively moderate effect, and the protease inhibitor had no discernible

effect on invertebrate interactions.

111
112
113
114 **Materials and Methods**
115

Plant material and growth conditions

Nicotiana benthamiana plants were grown in temperature-controlled growth chambers (Conviron,

Winnipeg, Canada) at 28°C or in a temperature-regulated greenhouse, at ~27°C for

approximately three weeks prior to transplanting in the field. For laboratory insect assays, *N.*

benthamiana plants were grown similarly in growth chambers at 28°C – 30°C. Plants used for

genome editing either expressed the *PCO-CAS9* gene (Li *et al.*, 2013), or the same *CAS9* while

also harboring a mutation in the *SerPIN-II* gene (Feng & Jander, 2023) described as the *SERP-*

II3-1 line (see Table S1 where this line, harboring a three BP deletion is shown to be named

serp2-2 in the current paper). These were grown in a growth room at 22°C prior to and following

viral infection. For assessment of the effects of growth temperature, on plant size, plants were

grown in a 22°C growth room, or in growth chambers at 28°C or 34°C.

Field-grown plants were planted during the summer of 2023 in a deer-excluded field in

Ithaca, NY (42.46 °N, 76.44 °W). Plants were planted in blocks of 14, each including two wildtype

plants and two independent lines of each mutant combination (Fig. S1). The plants' locations in

the blocks were randomized, but changes were made to ensure each mutation was equally

represented in the inner plants of the block and in the different compass directions. Distances

between individual plants were 40 cm and 110 cm between blocks (building on the field

experimental setup of Steppuhn et al (Steppuhn *et al.*, 2004)), and 24 blocks were planted in

three rows (Fig. S1B).

Insects and growth conditions

Chloridea virescens (tobacco budworm; formerly *Heliothis virescens*) and *Spodoptera exigua*

(beet armyworm) eggs were purchased from Benzon Research (Carlisle, PA, USA) and hatched

on beet armyworm diet (Southland Products, Lake Village, AR, USA) in a 28°C incubator. Two
strains of green peach aphid (*Myzus persicae*) were used in this study: a green strain that was
reported to be nicotine-sensitive and a red "USDA" strain that is nicotine adapted (Ramsey *et al.*,
2007, 2014). The green strain was reared on *Brassica rapa* var chinensis, whereas the red strain
was reared on *Nicotiana tabacum*. *Schistocerca americana* (American bird grasshopper) eggs
were kindly provided by Hojun Song at Texas A&M University. Eggs were hatched in a 30°C
growth chamber, and nymphs were raised at 30°C on lettuce, wheat grass, and wheat bran.

Mutant accessions and generation of the mutant population

The mutant population used in this study consisted of wildtype plants, mutants of three genes
encoding defensive compounds, and their combinations: *SerPIN-II* (Feng & Jander, 2023) (the
serp2-1 line corresponding to the *serpII3-2* line and the *serp2-2* line corresponding to *serpII3-1*),
ACYLSUGAR ACYL TRANSFERASE2 (*ASAT2*;(Feng *et al.*, 2022)), and the *N. benthamiana*
A622 gene, which was identified by a BLAST search of the *N. benthamiana* transcriptome with
known genes from other *Nicotiana* species (Deboer *et al.*, 2009) and was predicted to convert
nicotinic acid into 3,4-dihydronicotinic acid as an intermediate in nicotine biosynthesis. The *N.*
benthamiana gene with the highest similarity to the *A622* query sequence was
Nbe.v1.s00140g13970 (Supplemental Data S3). Three crRNA (RNA sequences transcribed from
the CRISPR locus) targets (TCTCTGATAAGAACGAAAGT, TGATTTCCACTGTCGGGGGA, and
AGAGTCTTATTCAATGTCCG) in the first, second and fourth exons, respectively, of *A622* were
identified using the CRISPR-P tool (Lei *et al.*, 2014). The three crRNAs were inserted into a
tobacco rattle virus (TRV) plasmid vector (Ellison *et al.*, 2020), either on their own or in
combination with an additional two crRNAs targeting *ASAT2*, which were identical to those used
by Feng et al., (Feng *et al.*, 2022). The two guide RNA combinations were expressed on two
different backgrounds, either *CAS9*-expressing plants (Ellison *et al.*, 2020) or *CAS9*-expressing
SerPIN-II mutants (Feng & Jander, 2023).

Genome editing of *N. benthamiana* plants was performed using TRV expressing guide RNAs
(Ellison *et al.*, 2020). In short, the PEE392 plasmid (Addgene #149282, www.addgene.org) was
used as a template to amplify a guide RNA fragment, a changing crRNA and a fragment from the
FLOWERING LOCUS T (*FT*) gene to drive meristem localization. PCR products were run on
electrophoreses gels, cut, cleaned and combined into the PEE083 plasmid (Addgene #149279)
using the Golden Gate assembly system (Engler *et al.*, 2008). PEE083 is a TRV plasmid
encoding TRV2 coat proteins, and between them the guide RNAs driven by a yellow leaf curl
virus promoter. This plasmid was transferred into the GV3101 *Agrobacterium* strain and co-
infiltrated with a TRV1 vector to drive viral infection of *N. benthamiana* (Sparkes *et al.*, 2006).

Plants were then grown to seed set, younger seed capsules were collected, and these were

grown for analysis by liquid chromatography-mass spectrometry (LC-MS).

Identification and relative quantification of nicotine and acylsugars

Nicotine and acylsugars were identified and relatively quantified using LC-MS. Acylsugars were

extracted using a 3:2:2 mix of acetonitrile (ACN), isopropanol, and water with 0.1% formic acid.

Leaves from the shoot apex were dipped in 500 µl of the solution and gently shaken for 2 min,

following which the leaves were removed, dried, and weighed. The solvent was then filtered and

analyzed using an Ultimate 3000 HPLC (Thermofisher Scientific) coupled to a Q Exactive

quadrupole - orbitrap mass spectrometer (Thermofisher Scientific) using a Supelco C18 column

(Sigma Aldrich; St. Louis, MO, USA). Buffers A and B were water and ACN respectively, to each

of which 0.1% formic acid was added. Buffer gradients and program settings were those used in

Feng et al. (Feng *et al.*, 2022). For nicotine extraction, 10 mm diameter leaf discs were cut from

leaves 3-4 of ~4-week-old plants and flash frozen in liquid N_2. Samples were ground using a

GenoGrinder 1600 MiniG homogenizer (SPEX SamplePrep, Metuchen NJ, USA) and metabolites

were extracted in a 500 µL solution composed of 89.9% MeOH, 10% water, and 0.1% FA. This

extract was filtered and run on the same LC-MS system and column as the acylsugars. The

column oven was set to 35°C, and initial flow was set to 0.1 ml/min at 99% water and 1% ACN.

From 5 - 8 min, the ACN concentration was increased to 100%. At that point, the pump's speed

was increased to 0.5 ml/min and the ACN concentration was kept at 100% until 9 min, then

reduced back to the initial 1% at 10 min, and held for an additional minute. Nicotine was identified

using a nicotine standard (Sigma Aldrich), exact mass, and fragmentation patterns.

Field observations

Interactions with insects, snails, slugs, and spiders were observed starting four days post planting

in the field, during the morning and early afternoon hours. Due to the large number of plants

initially observed, the observation cycle was divided to several days during which all the plants in

the field were examined. This examination included observation of plants without disturbing them,

from a distance of ~20 cm. Following this, interactions were examined more closely on both sides

of the leaves, the shoot apexes and the stems. For identification, interactors were photographed

using a Cannon G16 camera, or were sampled and further examined under a stereo microscope.

All 336 plants were examined between days 4-7 and 18-23. Following this, the first 8 blocks,

which were severely damaged in an uneven manner, were no longer examined for invertebrate

interactions. The remaining 16 blocks were examined between days 31-34, and blocks 10-13 and

18-22 were examined on day 45. Finally, blocks 12, 18 and 20 were examined on day 59. Given

the two-week period between observations and the extreme mobility of most interactors,

invertebrates that were observed during following visitations were scored individually. This

approach was validated by an analysis of observations of prominent invertebrates (leaf hoppers,

aphid colonies, flea beetles and spiders; Supplemental Data S2) in relation to the specific plants

on which they were found, which showed that in the vast majority of cases these insects were

observed on plants that did not have similar insects on them in the previous observation, did not

have similar insects in the following observation, and there was no case of insect types observed

continually on any plant. Furthermore, even when invertebrates were found on the same plant on

consecutive observations, they were at times from different species altogether (Fig. S2). In

addition to invertebrate interactions, plant height and survival were monitored. Plant height was

recorded at 24, 36-38, 52 and 66 days. Survival was monitored continually, *i.e.*, when a dead

plant was identified, it was recorded as such and added to the analysis.

Field grown plants were also sampled for nicotine and acylsugars. This process was

similar to sampling of these metabolites in the lab except that nicotine samples were whole young

leaves from the shoot apex and were weighed.

Insect assays under controlled conditions

For caterpillar growth assays, freshly hatched *S. exigua* neonates were placed in organza mesh

bags (Amazon item B073J4RS9C) on individual *N. benthamiana* leaves and returned to a 28˚C

growth chamber. After 11 days, survival was scored, and live caterpillars were weighed. This

timeframe was chosen following a previous trial with *Nicotiana glauca* in which *S. exigua* larvae

were confined on leaves for a similar duration (Negin *et al.*, 2024). Four-day-old *C. virescens*

larvae hatched on beet armyworm diet were weighed, and those weighing 5-12 mg were placed

in organza mesh bags on individual *N. benthamiana* leaves, with a single caterpillar in each bag.

Plants with caterpillars were then moved back to 28˚C growth chambers. After four days, the

caterpillars were weighed, and survival was assessed. Relative growth rate (RGR) was calculated

using the equation $\frac{\ln(w2)-\ln(w1)}{t}$ where *w1* is the weight on day 4, *w2* is the weight on day 8, and *t*

is the number of days between weighing (4 days in this case). This equation represents how

many mg the caterpillars gained every day for every mg they weighed.

Seven-week-old wildtype, *asat2*, a*622*, and *asat2+a622 N. benthamiana* plants, grown at

30˚C so that all plants are similarly-sized, were used for *Schistocerca americana* choice and no-

choice assays. For no-choice assays, four two-week-old grasshoppers were confined on

individual *N. benthamiana* plants using 10 x 16 inch micro-perforated polypropylene bags

(PrismPak; Berwick, PA, USA). Bagged plants with grasshoppers were placed in an incubator at

30˚C. After 16 days, grasshopper survival was assessed and the surviving grasshoppers were

weighed. For choice assays a cork borer was used to make 23 mm diameter leaf disks from
mature *N. benthamiana* leaves. Individual leaf disks of the two genotypes being compared were
placed on moistened filter paper in 10 cm diameter Petri dishes. Two two-week-old grasshoppers
were added to each Petri dish, and the Petri dishes were placed in a growth room at 23°C. After
14 hours, the leaf material was collected and scanned. The area of the remaining leaf material
was calculated using ImageJ (https://imagej.net/ij/). To account for leaf shrinkage during the
experiment, the amount eaten by the grasshoppers was calculated relative to leaves that had
received no grasshopper feeding.

Aphid reproduction assays were performed using both nicotine-sensitive and adapted *M.*
persicae strains. Five adult aphids were caged on the adaxial side of an intact leaf for
parthenogenetic reproduction. After 24 h, adults were removed and only three newly born
nymphs were left in each cage. After either eight or nine days (for nicotine-sensitive and adapted
strains, respectively), aphids were counted. Leaf cages were placed on individual plants in blocks
of all seven genotypes used in this study (wildtype, *a622*, *asat2*, *serp2*, *a622+asat2*, *a622+serp*
and *a622+asat2+serp2*). For choice assays, two leaf discs from different genotypes were placed
on top of Whatman filter paper in 100x15 mm Petri dishes, to which 1.1 ml water were added.
Five grown aphids were placed between the two discs and the Petri dishes were sealed with
Parafilm. These were left in a growth room for 24 h, after which the number of aphids on each of
the leaf discs was counted.

Statistical analysis.

Statistical analyses were performed using JMP 16 software (SAS Institute, Cary, NC, USA). The
frequency of field observed events was analyzed using a Chi-square test comparing the different
proportion of plants with a specific invertebrate observed on them between mutants and wildtype.
A Benjamini-Hochberg correction was performed to account for multiple comparisons. In this
analysis, events were considered as a type of invertebrate on a plant (*i.e.*, a plant with 3
leafhoppers was scored as one leafhopper event). Survival was similarly assessed using a Chi-
square test, whereas plant heights and field extracted nicotine and acylsugars were compared
using a Tukey's HSD test. Caterpillar and grasshopper weights on the different mutants were first
compared using a two-way ANOVA to examine the effects of the different metabolites and the
interactions between them. Next, caterpillar weights on individual mutants were compared to
those on wildtype leaves using Student's *t*-tests followed by a Benjamini-Hochberg correction.
This approach was chosen to balance a conservative approach, which would account for multiple
comparisons, but would not lose power through unnecessary comparisons (such as comparing
serp2 to *a622+asat2*), given the inherently large variance in insect assays. Aphid reproduction
was analyzed similarly. For choice assays, paired Student's *t*-tests were used.

275
276
277 **Results**
278
Generation of a622 and asat2 mutants

To generate *N. benthamiana* lines with reduced nicotine and acyl sugar content, the *A622* gene,

which is key to nicotine production (Deboer *et al.*, 2009), and *ASAT2*, which is essential for acyl

sugar production (Fan *et al.*, 2016), were targeted for CRISPR mutagenesis. Plants were infected

with tobacco rattle virus (TRV) carrying guide RNAs (gRNAs) targeting either the *A622* gene, or a

combination of the *A622* and *ASAT2* genes in both *CAS9*-expressing wildtype and *CAS9*-

expressing *serp2* mutants, which do not produce the Serp-II3 protease inhibitor (Feng & Jander,

2023). We assessed foliar nicotine and acylsugar content in subsequent generations using LC-

MS (Fig. S3-S4). Multiple independent lines from both backgrounds and infiltrated constructs

contained less than 5% of normal nicotine abundance (Fig. S3). When analyzing acylsugar

abundance, we found plants containing less than 5% of wildtype levels of acylsugars in the *CAS9*

background (Fig. S3D), but no plants with less than 5% of the acylsugar abundance in the *serp2*

background (Fig. S3F). However, there was one line close to the 5% threshold whose progeny

were grown for further examination.

When confirming the mutations using Sanger sequencing, we found that the *A622* gene

was often missing stretches of DNA between the three gRNA targets (1-3, 1-2, or 2-3), but single-

base pair indels were also formed (Fig. S5, Table S1). The *ASAT2* gene displayed a range of

mutations, including 136, 29, 3, and 2-base pair deletions (Fig. S6, Table S1). Based on these

results, additional plants were grown and screened (Fig. S3, S4) until the final panel for field

experiments was assembled. This panel did not include the *serp2+a622+asat2* triple mutant,

which was still being characterized and would be included later in controlled laboratory

experiments. Furthermore, when validating genome edits in the acylsugars + serpin double

mutants, both independent lines were not edited in the *ASAT2* gene. These plants were excluded

from further analyses.

Effects of nicotine, acylsugars and SERP2 on invertebrate interactions in a natural setting

Figure 1. Photographs of invertebrates comprising the main interactor classes observed in the field. (A) A leafhopper from the genus *Agallia*. (B) A leaf aphid (C) A leaf aphid colony (D) A flea beetle from the *Epitrix* genus. (E). A snail, probably from the genus *Succinea*. (F) A bumble bee (genus *Bombus*) pollinating a *N. benthamiana* flower. (G) A fly from the genus *Condylostylus*. (H) An orb weaver spider, probably from the species *Mangora acalypha*. (I) Stink bug (*Halyomorpha halys*) eggs during hatching. (J) An ant "farming" an aphid colony. (K) A lady bug (*Coccinella septempunctata*). (L) A parasitized aphid collected from *N. benthamiana* with a wasp larva that has almost entirely consumed the aphid's body.

Prior to transferring the mutant population to the field, we noticed that plants harboring mutations

in the *A622* gene were smaller than wildtype and the other *serp2* and *asat2* mutants. This

phenotype has been previously reported in field grown *Nicotiana tabacum*. However, we found

that this growth retardation was variable, with plants from the same genotype sometimes

exhibiting growth retardation and sometimes not. When searching for conditions that would

facilitate normal growth of the *a622* mutants, we found that growing the plants in higher

temperatures in growth chambers or a greenhouse reduced the growth retardation significantly

(Fig. S7). Whereas *a622* mutants grown at 22°C were significantly smaller than wildtype, those

grown at both 28°C and 34°C were similar in size to wildtype. Examining the response to the

changes in the growth temperature using a two-way ANOVA showed a significant interaction

between the line and growth temperature, meaning that wildtype and *a622* responded differently

to the increase in temperature. Wildtype plants were smaller as the temperature increased. By

contrast, *a622* plants were larger at 28°C than at 22°C, but slightly smaller at 34°C than when

grown at 28°C. Therefore, plants were grown in a temperature-controlled greenhouse at 27°C or

in growth chambers at 28°C. Wildtype *a622,* and all genotypes of the mutant population, which

showed similar growth under these conditions, were planted in the field.

The mutant population was planted in 24 blocks of 14 plants in a plowed 20 x 10 m field

section (Fig. S1). Invertebrate interactions were observed at five time points up to day 59 post-

planting (Fig. 1, S8; Supplemental Data S1). Throughout the wet summer months, snails and

slugs caused the greatest amount of plant damage. During the first observation cycle (4-7 d post-

planting) many leafhoppers and flies but no aphid colonies were observed. During the second

cycle (18-23 d post planting), six aphid colonies, fewer flies, and more than double the previous

number of flea beetles were observed. During the third observation cycle (31-34 d post-planting),

aphids and aphid colonies tripled, despite fewer plants being monitored due to high mortality in

the first row closest to the field margins. Far fewer leafhoppers and flea beetles were observed

during this cycle, but spiders tripled. A new predator, an assassin bug, appeared on eight plants,

totaling 10 individuals. At this point bumble bees pollinated plants, which had substantial numbers

of flowers. During the fourth observation cycle (45 d post planting), fewer plants were monitored.

However, aphids and aphid colony numbers were maintained whereas other herbivore numbers

were reduced. Assassin bug numbers increased to 29 individuals, whereas spiders maintained

their numbers. An additional observed interaction was four instances of aphids parasitized by

wasps. Finally, the fifth observation cycle (59 d post planting) included only three blocks. Despite

this small number, six parasitized aphids were observed, as well as five plants with whiteflies,

which were not seen in the first four observation cycles.

Figure 2. A breakdown of invertebrate interactions observed in a natural setting. (A) All the observed events separated to feeding and non-feeding interactions. Feeding interactions are further divided to (B) phloem feeders and (C) leaf chewers, with thrips and pollinators not falling strictly in either of those categories. (D) Non-feeding interactions. These include all interactions in which invertebrates interacted with the plant but did not directly feed on it. Within each invertebrate type, frequencies were compared between plant treatments and wildtype by a Chi-square test, as proportions of plants with events out of all the observed plants of the genotype and followed by a Benjamini-Hochberg correction (*p<0.05, **p<0.01). Interactions which were

observed at very low frequencies, such as a single caterpillar, were not included as individual

bars, but are accounted for in their group's proportion (such as leaf chewers). Wildtype (WT) n =

132, *a622* n = 107, *asat2* n = 131, *serp2* n = 132, *a622 + asat2* n = 107, *a622 + serp2* n = 101.

When analyzing all of the collected data, which included over 430 individual events

totaling over 560 individual invertebrates observed on plants (Fig. 2A), we discerned clear effects

of some of the mutations. Leafhoppers, by far the most commonly observed insects (121 events

totaling 174 individual insects), were found more frequently on *asat2*, *a622*, and *a622+asat2*

mutants (Fig. 2B), demonstrating that both nicotine and acylsugars protect plants from

leafhoppers (Fig. S9A). Aphids (52 events totaling 80 individuals) were found on significantly

more *asat2* and *a622+asat2* mutants (Fig. 2B), indicating that acylsugars but not nicotine are

important for protection against these phloem feeders (Fig. S9B). By contrast, aphid colonies

were found more often on *asat2*, *a622*, and *a622+asat2* plants, and none were found on wildtype

plants (Fig. 2B), indicating that despite not affecting individual aphids, nicotine does affect aphids'

ability to colonize *N. benthamiana* (Fig S9C). Flea beetles, the most abundantly observed leaf

chewers (45 events, 52 individuals; Fig. 2C), were significantly more abundant on *a622+asat2*

mutants, indicating that both acylsugars and nicotine provide protection, and plants only become

susceptible when losing both defenses (Fig. S9D). Similar results were obtained for thrips, which

were found more commonly on *a622+asat2* mutants. Slugs and snails showed a trend of being

found on more *asat2* mutants. Bumble bee pollinator visitations were statistically similar between

all plants due to the low numbers observed (Fig. 2C), However, the bees seemed to prefer plants

that had more flowers regardless of their chemotype. In non-feeding interactions (Fig. 2D), flies

and spiders preferred *asat2* and *a622+asat2* mutants.

Insect predators, including lady bugs and assassin bugs, were found significantly less

commonly on *a622+asat* and *a622+serp* plants. However, this is most likely due to the different

plant sizes, as plant height at the end of the experiment correlates significantly with average

assassin bug abundance on the different chemotypes (R^2=0.83, p=0.0042). Three other

observation classes, ants, egg-laying, and parasites, did not show differences between the

mutants due to their rare occurrence (6-10 events). However, it is interesting to note that the ratio

of parasitized aphids to aphids and aphid colony occurrence, is not correlated (R^2=0.17, p=0.35

and R^2=0.04, p=0.68 respectively). This lack of correlation could indicate that wasps parasitize

aphids with greater ease when aphids are on better-protected plants.

We next examined whether *asat2*, *a622*, and *a622+asat2* mutants were significantly

different from each other, indicating that effects seen in double mutants were additive, and

whether one of these specialized metabolites had a stronger effect than the other (Fig. S9). We

compared seven interactions in which some of these mutants were significantly different from

wildtype. There was no difference between the three mutants in leafhopper, flea beetle, and thrips

interactions, indicating that both nicotine and acylsugars protect plants similarly from these

insects. In the case of aphids, aphid colonies, and flies, both *asat2* and the double mutants were

significantly different from nicotine mutants, indicating that only acylsugars protect plants from

aphids and flies and that their protection from aphid colonization is stronger than nicotine, to the

extent that nicotine has no added effect on this interaction when acylsugars are affected. Spiders

were affected by acylsugars and the double mutants, indicating an effect of acylsugars but not

nicotine on these invertebrates.

401   When examining general fitness parameters, we monitored plant height and survival. In

general, all lines that harbored mutations in the *A622* gene were smaller (Fig. 3C). However, this

is an *a622*-specific phenotype, which also can be seen when growing *a622* mutants under

controlled conditions (Fig. S7). Since we found that cold temperatures contributed to the dwarfed

phenotype, it may very well be that the cold summer in the field in Ithaca, NY, contributed to *a622*

mutants growing more slowly, and it is difficult to attribute this phenotype to the effect of

invertebrate interactors. In addition to the effects of the *a622* gene, at the last measurement point

asat2 mutants were significantly smaller than wildtype, indicating a cumulative effect of their

increased invertebrate susceptibility. *Serp2* plants were similar to wildtype at all time points. In

addition to height, plant survival was assessed (Fig. 3D). Here we found that all plants harboring

the *a622* mutation survived at lower rates, whereas *asat2* mutations did not affect plant survival.

Seed production could not be assessed because, due to the transgenic status of the plants,

capsule development was terminated at early stages. However, assuming that the plants' harvest

index was not drastically altered, we can assume that the smaller biomass of *asat2* mutants

would correlate to a reduced seed yield.

Figure 3. Nicotine and acylsugar abundance of field-grown plants, plant height, and survival. (A)

Nicotine abundance determined by LC-MS with peak normalization by a bipyridine internal

standard and tissue weight. (B) Acylsugar abundance determined by LC-MS with the peak area

normalized to tissue weight. (C) Height of field grown plants throughout the growing season.

Dead plants (as scored by lack of stem greenness) were removed from the analysis. Different

letters indicate significance of $p < 0.05$ as determined by Tukey's HSD test. Results in A-C are

presented as the mean ± SE. (D) Survival frequency of field grown *N. benthamiana*. *$p < 0.05$ **p

< 0.01. as determined in a Chi-square test, comparing wildtype and each mutant. (A-B) N = 7-8,

(C) 24 d n = 40-46, 37 d n = 26-41, 52 d n = 11-30, 66 d n = 8-26. (D) N = 48.

Insect interactions under controlled conditions

Interactions observed in the field are the sum of many different factors combined into a single

phenotype. Therefore, we next examined the two main feeding guilds, phloem feeding and leaf

chewing, under controlled conditions. These experiments, in which plants were size-matched,

also aided us in understanding whether the smaller frequency of interactors found on *a622*

mutants compared to *asat2* mutants in the field was solely due to their smaller size. We initially

examined the interaction of the broad generalist caterpillar *Spodoptera exigua* (beet armyworm)

using a population that also included the *a622+asat2+serp2* triple mutant. However, only 10

caterpillars out of 350 neonates placed on 179 plants from the seven genotypes survived, and no
trend was seen of higher survival on plants that had fewer defensive compounds. Therefore, we
performed an additional trial with *Chloridea virescens* (tobacco budworm), with caterpillars 4 days
post-hatching, rather than first-instar larvae. Caterpillar weight (5 – 12 mg) was recorded and
insects were placed in mesh bags on individual leaves. Following four days of feeding, the
caterpillars reached weights of 4.5-108 mg. the average weight per treatment varied between
46.4 mg for wildtype and 59.8 mg for the triple mutants. Of the seven mutant combinations
examined, only the triple *a622+asat2+serp2* mutants had a significantly higher relative growth
rate (RGR) than wildtype (Fig. 4A). This result is intriguing since the *serp2* mutants showed no
effect in field and controlled conditions, but the addition of the *serp2* mutation beyond the
a622+asat2 mutations lead to caterpillars being heavier than on wildtype. When the effects of the
three defensive compounds on the caterpillars' RGR were examined in a two-way ANOVA, a
significant effect was found only for the *asat2* mutation ($p = 0.0079$), with no significant effect for
the interactions of the compounds, indicating that acylsugars affected *C. virescence* similarly
when only they were lacking and when combined with the other compounds. In this instance, only
13 out of 337 caterpillars placed on the plants were dead or had escaped at the end of the
experiment, with no trend seen for more death and escape on any of the examined genotypes.

Figure 4. *Chloridea virescens* growth and *Myzus persicae* reproduction under controlled conditions. (A) *C. virescens* weight following four days of feeding on *N. benthamiana* leaves. Larvae were placed in mesh bags on individual leaves four days post-hatching. N = 44-54. (B-C) The number of aphids in leaf cages, 8 (B) or 9 (C) days following the removal of all but three newly born nymphs. (B) Non-nicotine-tolerant *M. persicae* strain. (C) Nicotine-tolerant *M. persicae* strain. P values of student's *t*-tests are displayed above mutants that were compared to wildtype, with a Benjamini-Hochberg correction. Solid lines represent median values. Dashed lines represent quartiles. (B) N = 8. (C) N = 9.

Whereas phloem feeders belong to the Hemiptera order, leaf chewers are found across many orders, including Lepidoptera, Coleoptera and Orthoptera. For this reason, in addition to two lepidopteran species, we examined a leaf chewing orthopteran, *Schistocerca americana* (American bird grasshopper). Four grasshoppers were confined to single wildtype, *a622*, *asat2* and *a622+asat2* plants and, after 16 days of feeding, were weighed and their survival rates were scored. Only grasshoppers confined to the double *a622+asat2* mutants were heavier than those

confined to wildtype (Fig. 5A), and grasshoppers confined to all three mutants survived at higher

rates compared to wildtype (Fig. 5B). When the effects of nicotine and acylsugars on grasshopper

weight were examined in a two-way ANOVA, the effects of both nicotine (p=0.001) and

acylsugars (p=0.002) were significant, whereas the interaction between them was not. In addition,

choice assays between pairs of leaf discs of the four genotypes were performed (Fig. 5C-H).

Grasshoppers fed more on *a622* and *a622+asat2* leaves compared to wildtype (Fig. 5C,E) and

on *a622+asat2* leaves compared to *asat2* (Fig. 5H). The combination of these results indicates

that both acylsugars and nicotine protect *N. benthamiana* from *S. americana* herbivory, but that

nicotine has the greater effect of the two.

Figure 5. Growth, survival, and leaf choice of *Schistocerca americana* under controlled

conditions. (A) Average weight and (B) percent survival of groups of four *S. americana*

grasshoppers confined on whole wildtype and mutant *N. benthamiana* plants. N = 15; P values

are from Benjamini Hochberg corrected student's *t*-tests, comparing to the wildtype (WT) control.

(C-H) Percentage of total consumed leaf area in plates with leaf discs from two different

genotypes. The compared *N. benthamiana* lines are indicated beneath the panels. P values

indicated above panels were obtained using paired Student's *t*-tests. Solid and dashed lines

represent median and quartile values, respectively. N = 20.

We next examined the effects of our mutant population on aphid reproduction using both

nicotine-sensitive and nicotine-adapted *M. persicae* strains. Results were identical for the two

strains, with aphids reproducing faster on mutants containing an *asat2* knockout (Fig. 4B-C). All

other mutant combinations were similar to wildtype. When analyzing the effects of the compounds

using a two-way ANOVA, similar to the results in *C. virescence*, only acyl sugars had a significant

effect on aphid reproduction of both the non-adapted strain of *M, persicae* (p = 0.017) and the

nicotine adapted strain (p = 0.0007). Just as when the plants were grown in the field, acylsugars

dominated defense against aphids, and the other compounds were less important in this context.

The lower protective capacity of nicotine was not related to the plants' size in this case.

An additional question we wished to address, was whether aphids' preference of plants

harboring *asat2* mutations in the field was due to an increased "choice" of these plants, increased

growth, reproduction and survival of the aphids, or a combination of the factors. We therefore

performed dual choice assays using leaf discs. Twenty-one chemotype combinations were

examined using the two aphid strains. We found that the nicotine-adapted strain, which did not

show a different response compared to the nicotine sensitive strain in the reproduction assay, did

not significantly prefer any of the chemotypes (Fig. S10). However, when all leaves that have an

asat2 mutation were compared against those that do not, a significant preference for the *asat2*

mutants was revealed (Fig. S10W). Surprisingly, aphids from the "nicotine susceptible" strain did

not show a preference for nicotine-free leaves (Fig. 6E, I, H, M, N, O, Q). Furthermore, when

comparing all leaves containing normal nicotine abundance to those where nicotine was reduced,

the aphids' preference was similar (Fig. 6V). By contrast, all leaves harboring the *asat2* mutation

were preferred compared to wildtype (Fig. 6B, D, F), and also relative to many of the other

mutants (Fig. 6G, K, L, N, P, R, S, U). The combination of all *asat2* mutants compared to all leaf

discs not harboring an *asat2* mutation, was significant as well (Fig. 6W).

Figure 6. Non-nicotine-adapted *Myzus persicae* choice between 21 combinations of two leaf discs of wildtype and mutant plants. (A-U) comparison of single lines. The compared lines are indicated below the panels. (V-X) A comparison of all mutant discs harboring a mutation to all those that do not harbor it. P values indicated above panels were obtained from comparisons of the two-choice percentage using paired Student's *t*-tests. Solid and dashed lines represent median and quartile values, respectively. (A-U) N = 11-14. (V-X) N = 137–152.

Discussion

Chemical defenses are an extremely large group of plant protectants. A majority of these metabolites are assumed to be related to protection against pathogens and herbivores (Kortbeek *et al.*, 2019). However, the effects of very few of these metabolites on insect interactions have been examined in detail using mutant lines. Furthermore, since each plant produces many specialized metabolites and other defenses simultaneously, these compounds may have an altered effect in the context of the entire plant's chemotype.

For this reason, we chose three prominent *N. benthamiana* defensive compounds and examined their effects, both separately and combined. We found that, as might be expected, different defenses affected different interactors to varying extents. However, acylsugars stood out in almost all of the interaction types. Whether it was flea beetles, aphids, flies, or spiders, these sticky compounds protected plants from the initial visitation stage to the final colonization stage. Since the comparisons we performed were on plants that had a reduced abundance of the compounds, instances where only the double *a622+asat2* mutants were susceptible indicate that each compound on its own is enough to protect the plant. In this case, we might ask what advantage *N. benthamiana* gains from producing extremely high amounts of nicotine, which may reach up to 0.4% of the leaf's dry weight (Drapal *et al.*, 2021), which is similar to ~0.02%-0.04% of leaf fresh weight reported for *N. attenuata* (Keinänen *et al.*, 2001; Steppuhn *et al.*, 2004). One probable answer, is that nicotine protects *N. benthamiana* from herbivores which were not represented in our field site. For instance, Machado et al., (Machado *et al.*, 2016) found that jasmonate-deficient *Nicotiana attenuata* plants were more susceptible to vertebrate herbivores such as deer and rabbits. Our experimental setup, which excluded deer herbivory and attempted to exclude groundhog herbivory was not able to assess such an effect. Similarly, we found that nicotine has a greater effect than acylsugars on grasshopper herbivory under controlled conditions. However, no live grasshoppers were found in our field site. It is probable that, if our mutant population were grown in the native range of *S. americana*, greater effects of nicotine would be observed in the field.

It is possible that nicotine, like many other specialized metabolites (Zhou *et al.*, 2018; Erb
& Kliebenstein, 2020) has a developmental role, in addition to its insecticidal properties (Parr &
Thurston, 1972; Steppuhn *et al.*, 2004; Harvey *et al.*, 2007). This is in line with the smaller size of
our *N. benthamiana a622* mutants at lower temperatures, and a similar recent observation in *N.
tabacum* mutants (Burner *et al.*, 2022). By contrast targeting other parts of the nicotine
biosynthesis pathway, RNAi of the putrescine methyl transferase (*PMT*) gene required for
production of the methylpyrrolidine ring (Steppuhn *et al.*, 2004) and genome editing of the
berberine bridge like protein (*BBL*) gene (Vollheyde *et al.*, 2023), did not inhibit growth of the
nicotine-reduced plants. However, since the nicotine abundance in the *a622* mutants is reduced
to a greater extent than in the *pmt* and *bbl* mutants, in addition to expected reduction in three
other pyridine alkaloids, anabasine, anatabine, and nornicotine (Kaminski *et al.*, 2020), it cannot
be ruled out that pyridine alkaloids directly affect plant growth.

Nonetheless, the double *a622+asat2* mutants demonstrate that the small size of *a622*
mutants is not the main reason for the fewer insect interactions observed on these plants. In all
five interactions in which *asat2* mutants had an increased prevalence of invertebrates, the smaller
double mutants displayed a significant effect as well. Furthermore, there are two interactions
where the effects of double mutants were significant but those of single *asat2* mutants were not.
When observing the plants in the field, there were many cases in which smaller plants harboring
mutations in the *asat2* gene had far more interactors than nearby larger plants which did not.

Nicotine in *N. benthamiana* may also serve as a "second line of defense" against insect
herbivory. Almost every potent specialized metabolite that evolved in plants has driven the
evolution and adaptation of specialist herbivores, which are more tolerant or altogether resistant
to the metabolite. Furthermore, even when not looking at specialists, there are entire classes of
herbivores that could just be less susceptible to certain chemical defenses, as was seen with the
snails and grasshoppers in this study and was reported for slug resistance to alkaloids (Self *et al.*,
1964; Cornell & Hawkins, 2003; Aguiar & Wink, 2005; Ali & Agrawal, 2012; Altesor *et al.*, 2014;
Petschenka & Agrawal, 2015).

We may wonder, in that case, whether examining the effects of *N. benthamiana's*
defensive compounds so far away from its natural habitat, not only in terms of distance, but also
in terms of the growing conditions of this desert-adapted plant, are relevant. Indeed, examining
the ecology of *N. benthamiana* in such an ectopic setting may not shed light on many aspects of
N. benthamiana's ecology and native interactions with herbivores. However, it can elucidate the
function of the examined metabolites in a more precise way. In *N, benthamiana's* natural habitat,
herbivores have been exposed and adapted to its defenses for millions of years. In this habitat,
when an insect consumes a nicotine-rich leaf, we do not know whether this is because nicotine
does not affect the insect or whether it is a specialist herbivore attracted by the presence of

nicotine. In our conditions, we can be almost certain that the aphids that were found on the
nicotine-containing plants, were not there because of acquired tolerance through exposure, but
rather through more inherent properties of the compound's interaction with this insect. Similarly,
N. benthamiana is unlikely to suffer from slug consumption in its natural habitat. However, that
does not reduce the significance of our finding that nicotine has little effect on slugs and snails.

587   Plants lacking nicotine, acylsugars, and the serine protease inhibitor are nevertheless
lethal to *S. exigua* neonates, demonstrating that they must have additional potent protections.
When analyzing the specialized metabolites present in *Nicotiana* species, Drapal *et al.* showed
that a laboratory strain of *N. benthamiana* produced several pyridine alkaloids, chlorogenic acids,
sterols, terpenes, hydrocarbons, and other specialized metabolites (Drapal *et al.*, 2022). This
wide array of chemicals is consistent with the observation that combinations of compounds are
typically involved in protecting plants against any given herbivore (Chen, 2008). In contrast to *S.*
exigua, snails, which caused immense damage in the field, consumed wildtype plants as readily
as the mutant plants. We can therefore assume that, even if we didn't see an effect of nicotine or
SerPIN-II in the field, there are contexts in which these play a primary role in plant protection.
Finally, it is possible that the combination of compounds is greater than the sum of their individual
effects. This was elegantly demonstrated by Steppuhn et al. (Steppuhn & Baldwin, 2007), who
showed that herbivores could compensate for a protease inhibitor by increasing the amount of
plant material they consumed. However, nicotine prevented this compensation response. Such
an effect can be seen in the caterpillar growth assays. Nicotine alone, acylsugars alone, and even
the combination of the two does not affect tobacco budworm growth. However, adding the *serp2*
mutations, which alone was not found to affect insect interactions in any of our assays, results in
plants that are more susceptible to *C. virescens* (Fig. 4A). Similarly, *S. americana* nymphs grew
faster than on wildtype plants only in the absence of both acylsugars and nicotine (Fig. 5A).

606   In this study, we adopted an approach of hypothesis-based experimentation in a natural
setting (Negin & Aharoni, 2023). Using this approach, we recorded every interaction we were able
to score, including flies that were resting on leaves for mere seconds, parasitized aphids, and
bumblebees pollinating the *N. benthamiana* flowers. This type of experimentation is common
practice in many ecological circles but is not often used when more "molecular" trials that include
the generation of mutant populations are performed. An approach using mutant plants has the
advantage of allowing us to determine the defensive functions of the metabolite combinations,
while still leaving room for more complex answers beyond rejecting null hypothesis.

615
616

618 **Author Contributions:** GJ planned, guided, and oversaw the project, performed experiments,

619 analyzed data and revised the manuscript. BN planned the project, conducted the experiments,

620 analyzed the data, and wrote the manuscript. FW conducted experiments, aided with insect

621 identification and revised the manuscript. HDF assisted in insect identification, insect assay

622 setup, and manuscript revisions.

623
624 **Acknowledgments**
625
626 We thank Hojun Song for providing grasshopper eggs. This research was made possible by US

627 Department of Agriculture postdoctoral fellowship number 2022-67012-36739 to BN, and by US

628 Department of Agriculture award 2021-67013-33565 and a Triad Foundation award to GJ.

629
630
631 **References**

632
Agrawal AA, Petschenka G, Bingham RA, Weber MG, Rasmann S. 2012. Toxic cardenolides:
Chemical ecology and coevolution of specialized plant–herbivore interactions. *New Phytologist*
194: 28–45.

Aguiar R, Wink M. 2005. How do slugs cope with toxic alkaloids? *CHEMOECOLOGY* 15: 167–177.

Ali JG, Agrawal AA. 2012. Specialist versus generalist insect herbivores and plant defense. *Trends*
in Plant Science 17: 293–302.

Altesor P, García Á, Font E, Rodríguez-Haralambides A, Vilaró F, Oesterheld M, Soler R, González
A. 2014. Glycoalkaloids of wild and cultivated *Solanum*: Effects on specialist and generalist Insect
herbivores. *Journal of Chemical Ecology* 40: 599–608.

Bally J, Jung H, Mortimer C, Naim F, Philips JG, Hellens R, Bombarely A, Goodin MM, Waterhouse
PM. 2018. The rise and rise of *Nicotiana benthamiana*: A plant for all reasons. *Annual Review of*
Phytopathology 56: 405–426.

Bally J, Nakasugi K, Jia F, Jung H, Ho SYW, Wong M, Paul CM, Naim F, Wood CC, Crowhurst RN, *et*
al. 2015. The extremophile *Nicotiana benthamiana* has traded viral defence for early vigour.
Nature Plants 1: 1–6.

Burner N, Kernodle SP, Steede T, Lewis RS. 2022. Editing of *A622* genes results in ultra-low
nicotine whole tobacco plants at the expense of dramatically reduced growth and development.
Molecular Breeding 42: 20.

Carrillo L, Martinez M, Álvarez-Alfageme F, Castañera P, Smagghe G, Diaz I, Ortego F. 2011. A
barley cysteine-proteinase inhibitor reduces the performance of two aphid species in artificial
diets and transgenic Arabidopsis plants. *Transgenic Research* 20: 305–319.

Chen M-S. 2008. Inducible direct plant defense against insect herbivores: A review. *Insect*
Science 15: 101–114.

Cornell HV, Hawkins BA. 2003. Herbivore responses to plant secondary compounds: A test of
phytochemical coevolution theory. *The American Naturalist* 161: 507–522.

Deboer KD, Lye JC, Aitken CD, Su AK-K, Hamill JD. 2009. The *A622* gene in *Nicotiana glauca* (tree
tobacco): Evidence for a functional role in pyridine alkaloid synthesis. *Plant Molecular Biology*
69: 299–312.

Demkura PV, Abdala G, Baldwin IT, Ballare CL. 2010. Jasmonate-dependent and -independent
pathways mediate specific effects of solar ultraviolet B radiation on leaf phenolics and
antiherbivore defense. *Plant Physiology* 152: 1084–1095.

Drapal M, Enfissi EMA, Fraser PD. 2021. Metabolic changes in leaves of *N. tabacum* and *N.*
benthamiana during plant development. *Journal of Plant Physiology* 265: 153486.

Drapal M, Enfissi EMA, Fraser PD. 2022. The chemotype core collection of genus *Nicotiana*. *The*
Plant Journal 110: 1516–1528.

Ellison EE, Nagalakshmi U, Gamo ME, Huang P, Dinesh-Kumar S, Voytas DF. 2020. Multiplexed
heritable gene editing using RNA viruses and mobile single guide RNAs. *Nature Plants* 6: 620–
624.

Engler C, Kandzia R, Marillonnet S. 2008. A one pot, one step, precision cloning method with
high throughput capability. *PLOS ONE* 3: e3647.

Erb M, Kliebenstein DJ. 2020. Plant secondary metabolites as defenses, regulators, and primary
metabolites: The blurred functional trichotomy. *Plant Physiology* 184: 39–52.

Fan P, Leong BJ, Last RL. 2019. Tip of the trichome: Evolution of acylsugar metabolic diversity in
Solanaceae. *Current Opinion in Plant Biology* 49: 8–16.

Fan P, Miller AM, Schilmiller AL, Liu X, Ofner I, Jones AD, Zamir D, Last RL. 2016. In vitro
reconstruction and analysis of evolutionary variation of the tomato acylsucrose metabolic
network. *Proceedings of the National Academy of Sciences* 113: E239-E248

Feng H, Acosta-Gamboa L, Kruse LH, Tracy JD, Chung SH, Nava Fereira AR, Shakir S, Xu H, Sunter
G, Gore MA, *et al.* 2022. Acylsugars protect *Nicotiana benthamiana* against insect herbivory and
desiccation. *Plant Molecular Biology* 109: 505–522.

Feng H, Jander G. 2023. Serine proteinase inhibitors from *Nicotiana benthamiana*, a non-
preferred host plant, inhibit the growth of *Myzus persicae* (green peach aphid). *BioRxiv:*
2023.05.16.540980.

Fobes JF, Mudd JB, Marsden MP. 1985. Epicuticular lipid accumulation on the leaves of
Lycopersicon pennellii (Corr.) D'Arcy and *Lycopersicon esculentum* Mill. *Plant Physiology* 77:
567–570.

Green TR, Ryan CA. 1972. Wound-induced proteinase inhibitor in plant leaves: A possible
defense mechanism against insects. *Science* 175: 776–777.

Hare JD. 2005. Biological activity of acyl glucose esters from *Datura wrightii* glandular trichomes
against three native insect herbivores. *Journal of Chemical Ecology* 31: 1475–1491.

Hartl M, Giri AP, Kaur H, Baldwin IT. 2010. Serine protease inhibitors specifically defend *Solanum*
nigrum against generalist herbivores but do not influence plant growth and development. *The*
Plant Cell 22: 4158–4175.

Harvey JA, Van Dam NM, Witjes LMA, Soler R, Gols R. 2007. Effects of dietary nicotine on the
development of an insect herbivore, its parasitoid and secondary hyperparasitoid over four
trophic levels. *Ecological Entomology* 32: 15–23.

Heiling S, Li J, Halitschke R, Paetz C, Baldwin IT. 2022. The downside of metabolic diversity:
Postingestive rearrangements by specialized insects. *Proceedings of the National Academy of*
Sciences 119: e2122808119.

Hoogshagen M, Hastings AP, Chavez J, Duckett M, Pettit R, Pahnke AP, Agrawal AA, de Roode JC.
2023. Mixtures of milkweed cardenolides protect monarch butterflies against parasites. *Journal*
of Chemical Ecology.

Hopkins RJ, van Dam NM, van Loon JJA. 2009. Role of glucosinolates in insect-plant relationships
and multitrophic interactions. *Annual Review of Entomology* 54: 57–83.

Hu L, Ye M, Kuai P, Ye M, Erb M, Lou Y. 2018. OsLRR-RLK1, an early responsive leucine-rich
repeat receptor-like kinase, initiates rice defense responses against a chewing herbivore. *New*
Phytologist 219: 1097–1111.

Jassbi AR, Zare S, Asadollahi M, Schuman MC. 2017. Ecological roles and biological activities of
specialized metabolites from the genus *Nicotiana*. *Chemical Reviews* 117: 12227–12280.

Kaminski KP, Goepfert S, Ivanov NV, Peitsch MC. 2020. Production of valuable compounds in
tobacco. In: Ivanov NV, Sierro N, Peitsch MC, eds. Compendium of Plant Genomes. The Tobacco
Plant Genome. Cham: Springer International Publishing, 249–263.

Keinänen M, Oldham NJ, Baldwin IT. 2001. Rapid HPLC screening of jasmonate-induced
increases in tobacco alkaloids, phenolics, and diterpene glycosides in *Nicotiana attenuata*.
Journal of Agricultural and Food Chemistry 49: 3553–3558.

Kortbeek RWJ, van der Gragt M, Bleeker PM. 2019. Endogenous plant metabolites against
insects. *European Journal of Plant Pathology* 154: 67–90.

Kroumova AB, Wagner GJ. 2003. Different elongation pathways in the biosynthesis of acyl
groups of trichome exudate sugar esters from various solanaceous plants. *Planta* 216: 1013–
1021.

Kumar P, Pandit SS, Steppuhn A, Baldwin IT. 2014. Natural history-driven, plant-mediated RNAi-
based study reveals CYP6B46's role in a nicotine-mediated antipredator herbivore defense.
Proceedings of the National Academy of Sciences 111: 1245–1252.

Lei Y, Lu L, Liu H-Y, Li S, Xing F, Chen L-L. 2014. CRISPR-P: A web tool for synthetic single-guide
RNA design of CRISPR-system in plants. *Molecular plant* 7: 1494–1496.

Li J-F, Norville JE, Aach J, McCormack M, Zhang D, Bush J, Church GM, Sheen J. 2013. Multiplex
and homologous recombination–mediated genome editing in Arabidopsis and *Nicotiana*
benthamiana using guide RNA and Cas9. *Nature Biotechnology* 31: 688–691.

Lou Y-R, Anthony TM, Fiesel PD, Arking RE, Christensen EM, Jones AD, Last RL. 2021. It happened
again: Convergent evolution of acylglucose specialized metabolism in black nightshade and wild
tomato. *Science Advances* 7: eabj8726.

Luu VT, Weinhold A, Ullah C, Dressel S, Schoettner M, Gase K, Gaquerel E, Xu S, Baldwin IT.
2017. O-acyl sugars protect a wild tobacco from both native fungal pathogens and a specialist
herbivore. *Plant Physiology* 174: 370–386.

Machado RA, McClure M, Hervé MR, Baldwin IT, Erb M. 2016. Benefits of jasmonate-dependent
defenses against vertebrate herbivores in nature. *eLife* 5: e13720.

Meihls LN, Handrick V, Glauser G, Barbier H, Kaur H, Haribal MM, Lipka AE, Gershenzon J,
Buckler ES, Erb M, *et al.* 2013. Natural variation in maize aphid resistance is associated with 2,4-
dihydroxy-7-methoxy-1,4-benzoxazin-3-one glucoside methyltransferase activity. *The Plant Cell*
25: 2341–2355.

Millar NS, Denholm I. 2007. Nicotinic acetylcholine receptors: Targets for commercially
important insecticides. *Invertebrate Neuroscience* 7: 53–66.

Negin B, Aharoni A. 2023. Let's connect nature with hypothesis-based experimentation and
explore life in context. *The Plant Journal* 113: 23–25.

Negin B, Jander G. 2023. Convergent and divergent evolution of plant chemical defenses.
Current Opinion in Plant Biology 73: 102368.

Negin B, Shachar L, Meir S, Ramirez CC, Rami Horowitz A, Jander G, Aharoni A. 2024. Fatty
alcohols, a minor component of the tree tobacco surface wax, are associated with defence
against caterpillar herbivory. *Plant, Cell & Environment* 47: 664–681.

Parr JC, Thurston R. 1972. Toxicity of nicotine in synthetic diets to larvae of the tobacco
hornworm. *Annals of the Entomological Society of America* 65: 1185–1188.

Petschenka G, Agrawal AA. 2015. Milkweed butterfly resistance to plant toxins is linked to
sequestration, not coping with a toxic diet. *Proceedings of the Royal Society B: Biological*
Sciences 282: 20151865.

Pichersky E, Lewinsohn E. 2011. Convergent evolution in plant specialized metabolism. *Annual*
Review of Plant Biology 62: 549–566.

Puterka GJ, Farone W, Palmer T, Barrington A. 2003. Structure-function relationships affecting
the insecticidal and miticidal activity of sugar esters. *Journal of Economic Entomology* 96: 636–
644.

Ramsey JS, Elzinga DA, Sarkar P, Xin Y-R, Ghanim M, Jander G. 2014. Adaptation to nicotine
feeding in *Myzus persicae*. *Journal of Chemical Ecology* 40: 869–877.

Ramsey JS, Wilson AC, de Vos M, Sun Q, Tamborindeguy C, Winfield A, Malloch G, Smith DM,
Fenton B, Gray SM, *et al.* 2007. Genomic resources for *Myzus persicae*: EST sequencing, SNP
identification, and microarray design. *BMC Genomics* 8: 423.

Reichstein T, von Euw J, Parsons JA, Rothschild M. 1968. Heart poisons in the monarch butterfly.
Science 161: 861–866.

Schilmiller AL, Charbonneau AL, Last RL. 2012. Identification of a BAHD acetyltransferase that
produces protective acyl sugars in tomato trichomes. *Proceedings of the National Academy of*
Sciences 109: 16377–16382.

Schilmiller AL, Gilgallon K, Ghosh B, Jones AD, Last RL. 2016. Acylsugar acylhydrolases:
carboxylesterase-catalyzed hydrolysis of acylsugars in tomato trichomes. *Plant Physiology* 170:
1331–1344.

Self LS, Guthrie FE, Hodgson E. 1964. Adaptation of tobacco hornworms to the ingestion of
nicotine. *Journal of Insect Physiology* 10: 907–914.

Sparkes IA, Runions J, Kearns A, Hawes C. 2006. Rapid, transient expression of fluorescent fusion
proteins in tobacco plants and generation of stably transformed plants. *Nature Protocols* 1:
2019–2025.

Steppuhn A, Baldwin IT. 2007. Resistance management in a native plant: Nicotine prevents
herbivores from compensating for plant protease inhibitors. *Ecology Letters* 10: 499–511.

Steppuhn A, Gase K, Krock B, Halitschke R, Baldwin IT. 2004. Nicotine's defensive function in
nature. *PLOS Biology* 2: e217.

Van Dam NM, Hare JD. 1998. Biological activity of *Datura wrightii* glandular trichome exudate
against *Manduca sexta* larvae. *Journal of Chemical Ecology* 24: 1529–1549.

Van Dam NM, Horn M, Mareš M, Baldwin IT. 2001. Ontogeny constrains systemic protease
inhibitor response in *Nicotiana attenuata*. *Journal of Chemical Ecology* 27: 547–568.

Vollheyde K, Dudley QM, Yang T, Oz MT, Mancinotti D, Fedi MO, Heavens D, Linsmith G, Chhetry
M, Smedley MA, *et al.* 2023. An improved *Nicotiana benthamiana* bioproduction chassis
provides novel insights into nicotine biosynthesis. *New Phytologist* 240: 302–317.

Wang S, Alseekh S, Fernie AR, Luo J. 2019. The structure and function of major plant metabolite
modifications. *Molecular Plant* 12: 899–919.

Xu S, Brockmöller T, Navarro-Quezada A, Kuhl H, Gase K, Ling Z, Zhou W, Kreitzer C, Stanke M,
Tang H, *et al.* 2017. Wild tobacco genomes reveal the evolution of nicotine biosynthesis.
Proceedings of the National Academy of Sciences 114: 6133–6138.

Yang RSH, Guthrie FE. 1969. Physiological responses of insects to nicotine. *Annals of the*
Entomological Society of America 62: 141–146.

Yang C, Halitschke R, O'Connor SE. 2023. OXIDOSQUALENE CYCLASE 1 and 2 influence triterpene
biosynthesis and defense in *Nicotiana attenuata*. *Plant Physiology*.

Zavala JA, Patankar AG, Gase K, Hui D, Baldwin IT. 2004. Manipulation of endogenous trypsin
proteinase inhibitor production in *Nicotiana attenuata* demonstrates their function as
antiherbivore defenses. *Plant Physiology* 134: 1181–1190.

Zhou S, Richter A, Jander G. 2018. Beyond defense: Multiple functions of benzoxazinoids in
maize metabolism. *Plant and Cell Physiology* 59: 1528–1537.

Züst T, Strickler SR, Powell AF, Mabry ME, An H, Mirzaei M, York T, Holland CK, Kumar P, Erb M.
2020. Independent evolution of ancestral and novel defenses in a genus of toxic plants
(Erysimum, Brassicaceae). *Elife* 9.