The choice-wide behavioral association study: data-driven identification of interpretable behavioral components

David B. Kastner[1,4], Greer Williams[1], Cristofer Holobetz[1], Joseph P. Romano[2], Peter Dayan[3]

[1]*Department of Psychiatry and Behavioral Sciences, University of California, San Francisco, CA 94143, USA*

[2]*Department of Statistics, Stanford University, Stanford, CA 94305, USA*

[3]*Max Planck Institute for Biological Cybernetics, Tübingen 72076, Germany*

[4]Lead Contact

Correspondence: David B. Kastner: david.kastner2@ucsf.edu

Abstract

Behavior contains rich structure across many timescales, but there is a dearth of methods to identify relevant components, especially over the longer periods required for learning and decision-making. Inspired by the goals and techniques of genome-wide association studies, we present a data-driven method—the choice-wide behavioral association study: CBAS—that systematically identifies such behavioral features. CBAS uses powerful, resampling-based, methods of multiple comparisons correction[1–3] to identify sequences of actions or choices that either differ significantly between groups or significantly correlate with a covariate of interest. We apply CBAS to different tasks and species (flies[4], rats[5], and humans[6]) and find, in all instances, that it provides interpretable information about each behavioral task.

Understanding how behavior differs between different groups of humans or other animals is critical for generating and testing hypotheses about the functional role of genes, regions of the brain, and neural circuits, and is central to characterizing neurological and psychiatric dysfunction[7]. However, behavior is highly complex, evolving over multiple timescales and exhibiting substantial path dependencies due to individual experience[8–12]. It is increasingly possible to automate behavioral paradigms, and for computational methods to revolutionize behavioral analyses[13–22]. The latter come in two main flavors[23,24]: model-based, or top-down approaches, and data-driven, or bottom-up approaches. The former are substantially more prevalent than the latter; we offer a partial corrective.

In model-based analyses, behavioral data, such as choices in a decision-making task, are processed under the specific assumptions of a hypothesis or model. If the model or hypothesis is correct, this is highly efficient, since large volumes of data can be reduced to a handful of parameters that index semantically meaningful phenotypes, such as learning rates or differential sensitivity to rewards or punishments. These parameters can then be compared between the groups. However, even when substantial effort is put into building multiple alternative models, and comparing them in a statistically rigorous manner, it remains possible that the best fitting model nevertheless fails to characterize the behavior properly, rendering nugatory any interpretation of group comparisons. Additionally, with such approaches, confirmation bias[25] presents a significant challenge for accurate interpretation. Furthermore, in the worst of cases, model and hypothesis-based analyses provides a post-hoc framework for explaining any difference that can be found in a behavioral dataset[26]. Modern machine learning methods[27,28] can provide useful lower bounds for how well hypothesis-driven models should fit, but these approaches lack interpretability and data efficiency.

Data-driven analyses start from the other end, taking, and then characterizing, behavior without relying on parametric assumptions, and making very few assumptions about how the data is generated. Some of these approaches are unsupervised, for instance finding clusters in behavioral space and defining them as 'syllables' which can subsequently be compared between groups[18,29,30]; others are more supervised, directly looking for discriminative differences between populations or individuals[13–16]. By not trying to force data into a limited set of parameters, these methods should be more sensitive; however, comparisons between groups pose severe statistical challenges, because of the complexity and dimensionality of behavioral datasets.

Here, we present the choice-wide behavioral association study (CBAS), a data-driven
analysis method designed to identify relevant sequences of choices (or other discrete
behavioral features) made by subjects. CBAS has two components: 1) breaking down behavior
into a comprehensive language for comparison between two groups or correlation with a
covariate of interest; and 2) using rigorous, resampling-based, statistical corrections to account
for the resulting large number of comparisons and maintain statistical power despite
correlations in the data.

Choice as a common discretization for behavior

In developing our analysis, we were motivated by the data-driven approach of genome-
wide association studies (GWAS), whole exome sequencing (WES) and whole genome
sequencing (WGS)[31]. In some ways, the state of behavioral analysis in systems neuroscience
resembles the state of genetic analysis prior to GWAS/WES/WGS, where studies attempted to
associate candidate genes with phenotypes. Candidate gene studies were often underpowered
and failed to replicate[32–34], reminiscent of aspects of behavioral analyses[35].

GWAS/WES/WGS look for differences in base-pairs of the genome between groups of
subjects. The large numbers of base pairs being compared in these studies necessitates
statistical correction to enable reliable decisions about the significance of any differences
found. The discrete nature of base-pairs and the ability to compare that set across subjects
make GWAS/WES/WGS possible. To be able to develop a comparable method for behavioral
analyses it is critical to identify an appropriate discretization of behavioral tasks. In many cases,
choice provides just such a discretization (Fig 1). Indeed, there is evidence that, at a rather
fundamental level, behavior occurs through discrete choices[18,36]. We use three tasks, in three
diverse species—flies, humans, and rats—to show the breadth of applicability of CBAS.

Using choice as the basis of the comparison for the behavioral analysis requires an
additional consideration beyond what is done for the genome with base-pairs. For
GWAS/WES/WGS, an individual base pair can be a meaningful unit of information (although this
is only a partial story[37,38]) that can be compared between subjects. This need not be the case
for individual choices in behavior, whereby the choices that precede and follow a specific
choice can change the meaning of that choice. In this case, the relevant behavioral feature is a
whole sequence of choices. Therefore, CBAS does not just evaluate the occurrence of individual
choices, but, rather, the occurrence of all sequences up to a certain, user-defined, length. In
general, the longer the sequence length, the more data and computing time will be necessary.

Evaluating the occurrence of all sequences of choices up to a certain length, causes a
complexity, when it comes to correcting for the many comparisons, since, the sequences can be
highly correlated. Standard methods to correct statistically for multiple comparisons (e.g.
Bonferroni, Holm, Benjamini-Hochberg, Benjamini-Yekutieli) are either incorrect or
underpowered for correlated data. Therefore, we instead use an approach based on resampling
to correct for the multiple comparisons, which retains power in the face of correlations[1].

CBAS identifies interpretable differences for fly y-maze

The first task we considered involves drosophila walking on a y-maze (Fig 1a). The
movement of flies on the maze is tracked as they make the choice of going to either the left or

right arm after leaving the previous arm[4]. This left/right discretization of the task has enabled
many conclusions about the genetic nature of individual variability[4,39–43]. Here, we compare the
choices of two outbred strains of fly. Analyses in the original paper[4] identified some clear
indications about the difference; however, the data provide a useful testing ground for CBAS
since there are many subjects and the results are relatively low-dimensional.

When deciding to perform a data-driven genomics method, there are still decisions that
need to be made. For instance, deciding to focus on only exome sequences in WES, or SNPs in
GWAS. Similarly, to run CBAS, a few decisions need to be made about how to structure the data
(Fig 2a). These decisions are important, and are not normally pre-determined directly by the
data. Any conclusions drawn need to be interpreted in their light.

The first decision is the possible choices that will make up the sequences used in CBAS.
We refer to this as the basic language for the application of CBAS. For the fly task, the language
is left or right turns. Next, as described above, a decision needs to be made about the maximum
number of choices in a row that will make up all the sequences used in CBAS. For the fly task,
we chose a maximum sequence length of 10 choices in a row. That means that CBAS evaluates
all sequences from length 1 – 10 that exist in the dataset.

CBAS then works through evaluating the average sequence count for each sequence,
which we refer to as the rate of that sequence. The rate is calculated by counting the total
number of times that sequence occurs in the population divided by the number of individuals in
the population. To be precise about the occurrence of the sequence, a decision needs to be
made about the number of trials over which this is counted, i.e. a criterion. The same criterion
is applied to all subjects. For the fly task, we use the first 250 turns as that criterion. The last
decision that needs to be made is what to do with subjects that do not reach the criterion. For
the fly task, we exclude the subjects that do not reach criterion.

Given these decisions, we performed a CBAS comparing two outbred strains of flies,
Cambridge-A (CA) and w1118, from a publicly available dataset[4]. The w1118 strain is the
background strain for many transgenic flies[4]. For each sequence of left/right turns up to 10
turns long, the rate was calculated for the two strains (Fig 2b). Using the Romano-Wolf
resampling-based multiple comparisons correction[1,2] (see methods), we determined which
sequences the two strains utilize significantly differently (Fig 2c). As a significance threshold, we
use median control of the false discovery proportion at 5%, which is comparable to 5% false
discovery rate (FDR) for the resampling-based method[2] (see methods).

CBAS identifies many sequences that differ significantly between the CA and w1118
lines, some of which are shown in Fig. 2d. Upon inspection of the sequences, a clear,
interpretable, difference is apparent between those sequences that occur more in the CA line
and those that occur more in the w1118 line (Fig 2d&e). The CA line utilizes sequences with
extended numbers of the same turn in a row, whereas the w1118 line utilizes sequences with
more frequent changes in turn direction. Therefore, CBAS not only identifies that there is a
difference between the two fly lines, but it also provides information to support an
interpretation as to the nature of the difference.

Data-driven methods invariably need more data than hypothesis-driven methods. To
estimate the sample size needed for appropriately powered experiments to detect any
difference between these two fly lines, we took advantage of the large dataset to resample
groups of smaller sizes from the data and recalculate the CBAS for each set of resampled
groups. We generated an estimate of the power for different number of flies per group (Fig 2f)
(see methods), by comparing the number of significant sequences identified by CBAS when
comparing CA to w1118 to the number of significant sequences identified by CBAS when
comparing the lines to themselves (Fig S1b). With 40 flies per group, CBAS has an estimated
power >80% to distinguish the CA and w1118 lines.

Graceful decay of CBAS output with decreasing group size

The power calculation concerns the ability of CBAS to distinguish between the two
strains. A more refined question is what the nature and comprehensiveness of the collection of
significantly different sequences would be as smaller numbers of flies per group are analyzed.
This would determine our ability to derive interpretations from fewer subjects.

We first evaluated the number of significant sequences identified from smaller group
sizes as a fraction of the total number of significant sequences identified with the full dataset
(Fig 3a). As expected, with smaller group sizes, we find fewer overall sequences; however,
across the range of group sizes evaluated, the median fraction of sequences was larger than the
fraction of the population being used in the smaller group CBAS (Fig 3a). This indicates that, for
this dataset at smaller group sizes, the proportion of sequences identified by CBAS grows faster
than the proportion of subjects in the CBAS. This provides a rapid increase in the amount of
information provided by CBAS as sample size increases.

We next evaluated the fraction of the sequences identified by CBAS for the smaller sized
groups that are not identified in the CBAS on the full dataset. Since we control the false
discovery proportion and not the family-wise error rate, we do not expect this value to be zero;
however, it was consistently small across all the group sizes evaluated (Fig 3b). The medians
across the different repeats of the same sample sizes across the different sample sizes are all
less than 2%.

Then, we evaluated the similarity of the sequences identified by CBAS with
nonoverlapping sets of subjects with the same group size (Fig 3c). At smaller groups sizes, this
overlap could be quite low (median 24.4%), even though there was sufficient evidence to
discriminate the strains (Fig. 2f). This means that the specific sequences identified by CBAS can
be quite variable from one experiment to the next, especially at smaller group sizes. Given that,
we sought to understand if the structure of the sequences comported with the conclusion from
the full dataset—that the CA line uses more of the same turns in a row and the w1118 line
more frequent changes in direction of turn. Even though the overlap between CBAS on
different groups might not be that high (Fig. 3c), the sequences identified still follow the same
structure as the full dataset in regard to number of the same turns in a row, (Fig 3d). As the
group size increases the output of CBAS consistently becomes more similar to the results from
the full dataset (Fig 3e), licensing more generalized conclusions.

CBAS provides evidence for testing model-based analysis

The second task we considered, is the two-step task developed to test the interplay
between model-based and model-free (reinforcement) learning (Fig 1b)[44]. In this task, subjects
choose between pairs of images at two different stages. The image choice at the first stage
governs the pair of images from which the subject can chose at the second stage. Following
image choice at stage two, reward is delivered based on dynamic probabilities associated with
each of the images. Model-based analyses of variants of this task have led to conclusions about
the algorithms used in different brains regions[44,45] and differences that underlie psychiatric
symptoms[6]; however, the interpretations are not without controversy[46], and the complexity of
behavior makes it difficult to evaluate ways in which the model might be missing features in the
data.

To test the ability for CBAS to provide useful information in such a situation, we
evaluated the open-source dataset from Gillan et al[6]. In that work, a large population of human
subjects performed the two-step task, and also answered a series of psychiatric symptom
questionnaires. The authors performed factor analysis on the different questionnaires and
found that factor 2, which they associated with intrusive thoughts and compulsive behavior,
had significant association with the way subjects performed that task. They found that subjects'
factor 2 score was negatively associated with model-based decision making, leading to their
conclusion that the greater the loading on intrusive thoughts and compulsive behavior for a
person, the less likely they were to utilize model-based decision making on the two-step task.

There were two different groups to compare in the fly dataset. In this dataset the
relevant comparison is correlation with a covariate (i.e. factor 2 score from the psychiatric
symptom questionnaires)[6]. Therefore, we extended CBAS to identify the sequences that
significantly correlate with this particular factor score (see methods). As with the CBAS applied
to the fly data, there are decisions that need to be made to apply CBAS to this human dataset
(Fig 4a). For this CBAS, the language is comprised of 8 different units: choosing image 1 or 2;
making a choice within set A or set B, distinguishing A and B depending on whether a reward
was delivered (shown in the figure as bold and underlined); and making 'no choice' (either at
the first or second image), which occurs in the data, albeit rarely. Following Akam et al.[45] we
collapsed image 3 and 4 into set A and image 5 and 6 into set B because the specific images
within the set are not relevant for the critical decision at the first stage. We ran CBAS on all
sequences up to 4 choices long and evaluated the rate of the sequences over the 400 choices of
the dataset (200 trials of the two-step task). In the open-source dataset, there was no subject
who did not reach criterion.

CBAS evaluated the correlation between the usage of each sequence and the factor 2
score (Fig 4b&S2a). Multiple sequences were significantly correlated (Fig 4c&d). To understand
the output of CBAS, we review the expectation associated with the hypothesis-driven analysis
of this experiment. This task was designed to evaluate the interplay between model-based and
model-free decision making[44]. Model-based decision making develops an understanding of the
structure of the world (i.e. the model) and makes choices based on that understanding. Model-
free decision making makes choices based on reinforced past successful actions, without
developing an understanding of the structure of the world. A way to see the difference
between these two decision-making schemes is when a reward is delivered at set A ('**A**' in our
language) after having come from image 2. A regular model-free learner will reinforce the visits

to both 2 and A and will therefore be more likely to choose 2 on the next trial. The model-based
learner will have an understanding that choosing image 1 on the next opportunity makes it
more likely to return to set A (because of the common/rare transition structure in the
environment) and will therefore chose object 1[44].

Instead of identifying correlations with sequences that relate to either the model-free or
model-based learning, CBAS instead identifies a positive correlation with many sequences
involving being rewarded at A and then selecting image 2 or being rewarded at B and then
selecting image 1 (Fig 4d). Just as it was helpful with the fly data to find complete sets of
sequences that were significant (e.g. all of the sequences that occurred in the data with 10 left
or right turns in a row were identified as happening more in CA than w1118, etc.) we were also
able to find practically complete sets of sequences that were identified as being positively
correlated with the factor 2 score. Practically all of the sequences in the dataset that contained
1**A**2 or 2**B**1, were significantly correlated with this score; whereas none of the sequences with
1**B**1 was identified as significant (and there were no instances of 2**A**2 in the dataset)
(Fig4e&S2b).

Sequences 1**A**2 or 2**B**1 can be classified as anti-model-based decisions. The subjects get
rewarded after choosing from the common side (A from 1 or B from 2), but then selects the
image that will rarely bring them back to the previously rewarded side (2 from A or 1 from B).
CBAS therefore identifies that anti-model-based learning is a prominent feature that correlates
with intrusive thought and compulsive behavior loading. Further experimentation will be
needed to understand the interplay between this mode of decision making, the task, and these
symptoms scales.

As with the fly data, we could evaluate the sample size needed to reach a given
statistical power for identifying any significant correlation between the factor 2 score and this
task. We resampled smaller group sizes and compared the CBAS for the true relationship
between the sequence counts and factor 2 score to a randomly generated set of factor 2 score
values that was drawn from the same distribution as the data (Fig S2c). A sample size of 900
individual provides a power >80% to detect significant correlations between the factor 2 score
and the subject (Fig 4f), which compares favorably to the sample size of ~1,200 – 1,600 subjects
that Gillan et al. calculated to generate their dataset[6].

CBAS identifies a phenotype consistent with ASD in *Scn2a* haploinsufficient rats

The third task we considered, involves spatial alternation behavior in rats. For this task,
to get reward, the rats must alternate between pairs of arms of a track whilst visiting a different
arm of the track in between (Fig 1c). Spatial alternation is a common behavioral paradigm for
phenotyping and neurophysiology, and the discretization of the behavior into the arms chosen
by the animals forms the basis of many of the conclusions from these studies[47–51]. However, our
recent work calls into question the assumptions and hypothesis that motivate the standard
analysis for spatial alternation behavior[5,52]. Even though we developed reinforcement learning
agents to fit individual behavior, those agents showed clear differences from the way the
animals learned[5], limiting their use for phenotyping. Therefore, we applied CBAS.

We sought to discriminate wild-type (WT) rats from those haploinsufficient for *Scn2a*
(*Scn2a^+/-*), a high confidence, large effect, autism spectrum disorder (ASD) risk gene[53,54]. We
collected a dataset of over 200 rats performing six different spatial alternation contingencies
using our previously described automated behavioral system[5] (see methods). Each spatial
alternation contingency was defined by the three arms of the track where alternation needed
to occur to get reward. For example, if the contingency was at arms 2, 3, and 4, reward would
be provided for every arm visit in the sequence **3**-**4**-**3**-**2**-**3**-**4**.

The language for the spatial alternation CBAS was visiting each arm of the track and not
getting reward and visiting the goal arms and getting reward. That means that within each
contingency there were a total of 9 possibilities—6 unrewarded arms and 3 rewarded arms. For
example, if the contingency was at arms 2, 3, and 4, the language is composed of visiting arms
1, 2, 3, 4, 5, or 6 and not getting rewarded or visiting arms **2**, **3**, or **4**, and getting rewarded.
Each contingency was considered separately, as visiting a sequence of arms during one
contingency likely means something different than visiting those same arms in a different
contingency. We chose to evaluate all sequences up to six choices long. For the criterion, we
used the trial at which each animal reached 100 perfect performance sequences four choices
long for each contingency, and we included animals that did not reach the criterion (Fig 5a) (see
methods).

In running the CBAS on this dataset, we evaluated the rate difference of >86,000
sequences across the six different contingencies, with a total of 1,476 sequences being
identified as significantly different between the *Scn2a^+/-* and WT littermates (Fig 5b&c). To
interpret common features of those sequences, we sought categories of sequences for which a
substantial fraction was identified as being significantly different between the groups: the
strategy that was informative for the fly and human datasets (Fig 2e&4e). One common feature
of spatial alternation behavior is the 3-arm structure of each contingency (e.g. arms 2, 3, and 4
are the only arms with the potential to be rewarded during contingency A and E). Therefore, in
all contingencies, we evaluated all sequences that exclusively contained all sets of three arms.
For example, within a contingency we identified all sequences exclusively containing arms 2, 3,
and 4, which means that every sequence within that category only contains visits to arms 2, 3,
and 4 (but does not have to visit all of the arms). We did this with sets of three arms,
independent of whether the choice was rewarded or not, as well as the set of three arms in a
contingency that were all rewarded. There are 21 sets of three arms for each contingency, and,
consistent with the relevance of these sets for the behavior, across all contingencies they
contained only ~25% of all the sequences that the rats performed during the experiment, but
~75% of all of the significant sequences (Fig S3a).

Some of these sets of 3 arms are clearly interpretable. One such is the set of three arms
from the current contingency, which contains all sequences where the animal exclusively visits
the arms of the contingency but could make errors in the order of those arm visits (Fig 5d; left).
A second interpretable set consists of three arms from the current contingency that are
rewarded, which contains all sequences where the animal consistently performs the task
correctly. A third and fourth are the sets of three arms exclusively containing the previous (Fig
5d; right), or the one before previous, contingency arms, which contain sequences where the
animals repeat prior actions that are no longer optimally rewarded.

For all sets of three arms, in all the contingencies we tabulated the fraction of the total sequences in the set that were significant either for WT > $Scn2a^{+/-}$ or WT < $Scn2a^{+/-}$. We then asked if any of these fractions were larger than would be expected from randomly distributing the significant sequences across all possible sequence types within the entire category of sets of three arms (see methods) and found consistent patterns in how the WT and $Scn2a^{+/-}$ rats differed (Fig 5e). In 4 out of the 6 contingencies, $Scn2a^{+/-}$ rats showed increased usage of sequences containing the current contingency, and in 3 out of the 6 contingencies, $Scn2a^{+/-}$ rats also showed increased usage of sequences related to prior contingencies (either 1 or 2 contingencies back). By contrast, in 4 out of the 6 contingencies the WT rats showed increased usage of sequences containing the rewarded arms of the current contingency. This indicates that WT rats are ultimately better at performing spatial alternation behavior than $Scn2a^{+/-}$ rats, and that $Scn2a^{+/-}$ rats show difficultly transitioning from prior actions, repeating sequences from prior contingencies, possibly consistent with restrictive and repetitive actions, which forms one of the diagnostic criteria for ASD[55].

As with the fly and human data, we could evaluate the sample size needed to reach a given statistical power to detect any difference between these genotypes. We resampled smaller group sizes and compared the CBAS for the comparison between the WT and $Scn2a^{+/-}$ rats to the comparison between WT and itself and $Scn2a^{+/-}$ and itself (Fig S3b). A sample size of 30 rats per group provides a power >80% to detect a difference between the two genotypes (Fig 5f).

We have presented a general behavioral analysis method, CBAS, for identifying interpretable behavioral components that is grounded in sequences of choices made by subjects. There has been significant progress, in recent years, in tracking and analyzing, short timescale actions of subjects[13–22]; however, we lack methods for generally analyzing and interpreting sequences of these actions or long-run choices and decisions of subjects during behavior. CBAS provides just such a method and is applicable across a wide array of species and different behavioral paradigms (Fig 1). It can be used to test models and hypotheses (Fig 2&4), and, for instance, using generalizable simple principles about the relationship between sequences and task contingencies, it can generate hypotheses in complex behaviors where reliable computational understanding has yet to emerge (Fig 5). Through taking advantage of large-scale data collection and rigorous statistical methods, CBAS has the potential to transform our use of behavior in a comparable way to the ways that GWAS/WES/WGS changed the paradigm for genomic studies.

Methods

Animals: All experiments on rats were conducted in accordance with University of California San Francisco Institutional Animal Care and Use Committee and US National Institutes of Health guidelines. Rats were fed standard rat chow (LabDiet 5001). To motivate the rats to perform the task, reward was sweetened evaporated milk: 25 g of sugar per can (354 ml) of evaporated milk (Carnation). The rats were food restricted to ~85% of their basal body weight.

The $Scn2a$ mutant rats were generated due to funding from the Simons Foundation Autism Research Initiative. Long Evans $Scn2a$ mutant animals (LE-$Scn2a^{em1Mcwi}$;

RRID:RGD_25394530) were generated at the Medical College of Wisconsin and shipped to the
University of California San Francisco for this study. Briefly, a single guide RNA targeting the
sequence GTGAAATCCAACCAATTCCA sequence within exon 5 of *Scn2a* was mixed with Cas9 (*S.*
pyogenes) protein (QB3 MacroLab, UC Berkeley) and injected into the pronucleus of fertilized
Long Evans (Crl:LE, Charles River Laboratories) embryos. Among the resulting offspring, a
mutant founder was identified harboring a net 4-bp deletion allele consisting of a 10-bp
deletion (rn7: chr3:50,364,411-50,364,420) along with a 6-bp insertion of TTCACT, inducing a
frameshift in the coding sequence predicted to truncate the normal protein after 193 amino
acids. The founder was backcrossed to the parental Crl:LE strain to establish a breeding colony.

Spatial alternation behavior: The automated behavior system for spatial alternation behavior
was previously described[5]. There are different symbols on each arm of the track serving as
proximal cues, and there are distal cues distinguishing the different walls of the room.
Pneumatic pistons (Clippard) open and close the doors. Python scripts, run through Trodes
(Spike Gadgets), control the logic of the automated system. The reward wells contain an
infrared beam adjacent to the reward spigot. The automated system uses the breakage of that
infrared beam to progress through the logic of the behavior. In addition to the infrared beam
and the spigot to deliver the reward, each reward well has an associated white light LED.

Each cohort of rats is divided into groups of four (or three) animals. The same groups
were maintained throughout the duration of the experiment. Within a group, a given rat is
always placed in the same rest box, and the four rats of a group serially perform the behavior.
The rats have multiple sessions on the track each day. Prior to beginning the first spatial
alternation contingency, the rats experience multiple days and sessions where they get
rewarded at any arm that they visit (provided it is not an immediate repeat). During this period
of the behavior, the duration of a session is defined by a fixed number of rewards, or a fixed
amount of time on the track (15 minutes), whichever came first. During the alternation task the
duration of a session was defined either by a fixed number of center arm visits and at least one
subsequent visit to any other arm, or a fixed amount of time on the track (15 minutes),
whichever came first.

The algorithm underlying the spatial alternation task is such that three arms on the track
have the potential for reward within a given contingency, for example during a contingency at
arms 2-3-4, arms 2, 3, and 4 have the potential to be rewarded, and arms 1, 5, and 6 do not. Of
those three arms we refer to the middle of the three arms as the center arm (arm 3 in the
above example) and the other two arms as the outer arms (arms 2 and 4 in the above example).
Reward is delivered at the center arms if and only if: 1) the immediately preceding arm whose
reward well infrared beam was broken was not the center arm. Reward was delivered at the
outer two arms if and only if: 1) the immediately preceding arm whose reward well infrared
beam was broken was the center arm, and 2) prior to breaking the infrared beam at the center
arm, the most recently broken outer arm infrared beam was not the currently broken outer
arm infrared beam. The one exception to the outer arm rules was at the beginning of a session,
if no outer arm infrared beam was broken prior to the first infrared beam break at the center
arm, then only the first condition had to be met.

For the running of the behavior, the infrared beam break determined an arm visit;
however, the rats sometimes go down an arm, get very close to the reward wells, but do not
break the infrared beam. Therefore, for all the analyses described for the rats, an arm choice is
defined as when a rat gets close to a reward well. These times were extracted from a video
recording of the behavior. These, effective, missed pokes were more frequent at the beginning
of a contingency. This proximity-based definition of an arm visit added additional arm visits to
those defined by the infrared beam breaks, and none of them could ever be rewarded, nor alter
the logic of the underlying algorithm. However, because of the non-Markovian nature of the
reward contingency, they could affect the rewards provided for subsequent choices.

A total of 121 WT (66 males, 55 females) and 120 $Scn2a^{+/-}$ (66 males, 54 females) rats
were run on the spatial alternation task. WT and $Scn2a^{+/-}$ rats were littermates and were
housed together prior to their being food restricted before the behavior. During the behavior
and food restriction the rats were single housed. 1 WT rat died after finishing the first spatial
alternation contingency, 1 WT rat died after finishing the fourth spatial alternation contingency,
and 1 $Scn2a^{+/-}$ rat died after finishing the first spatial alternation contingency. The data from the
animals that died was included up until their expiration. For the first contingency: 5/121 WT
rats and 15/120 $Scn2a^{+/-}$ rats did not reach the CBAS criterion. For the second contingency:
4/120 WT rats and 19/119 $Scn2a^{+/-}$ rats did not reach the CBAS criterion; for the third
contingency: 4/120 WT rats and 12/119 $Scn2a^{+/-}$ rats did not reach the CBAS criterion; for the
fourth contingency: 2/120 WT rats and 12/119 $Scn2a^{+/-}$ rats did not reach the CBAS criterion;
for the fifth contingency: 4/119 WT rats and 14/119 $Scn2a^{+/-}$ rats did not reach the CBAS
criterion; and for the sixth contingency: 3/119 WT rats and 12/119 $Scn2a^{+/-}$ rats did not reach
the CBAS criterion.

Romano-Wolf resampling based multiple comparisons correction: We follow the terminology
and description laid out in Clarke et. al[56] to describe the Romano-Wolf multiple comparison
correction. First, we describe the way the method corrects for the family-wise error rate
(FWER) and then explain how the procedure is extended to provide median control of the false
discovery proportion. FWER control at a level of α means that across all comparisons there is a
α percent chance of having at least one false positive rejection of a null hypothesis. The
Romano-Wolf procedure provides FWER control through resampling the data[1]. It tests a total of
S hypotheses.

It is not generally known if the Romano-Wolf procedure controls for type III, or
directional, errors. Type III errors are errors in the sign, or direction, of the conclusion. For
example, if a statistical test provided information to reject the null hypothesis $\theta_1 = \theta_2$, and you
then concluded that $\theta_1 > \theta_2$, when in fact $\theta_1 < \theta_2$. Therefore, instead of running a single two-
tailed test for each sequence, we run two one-tailed tests for each sequence. Therefore, the
total number of hypotheses tested, S, is twice the total number of sequences being compared.
For CBAS those hypotheses take one of two forms: 1) the rate of each sequence (r_s) is the same
between two groups $\Delta r_s = 0$, or 2) that there is no correlation (ρ_s) between each sequence and
a covariate of interest, $\rho_s = 0$. For the one-tailed versions of each hypothesis, we ask if $r_{s_1} -$
$r_{s_2} > 0$ and $r_{s_2} - r_{s_1} > 0$ for the comparison CBAS, where r_{s_N} is the rate of sequence s for
group N, or if $\rho_s > 0$ and $\rho_s < 0$ for the correlational CBAS.

The first step in the procedure is to create a studentized test statistic for each
hypothesis. The studentization is different based on whether the CBAS is comparing two groups
or calculating a correlation. In the case where two groups are being compared:

$$t_s = \frac{\Delta r_s}{\sigma_s} \tag{7}$$

428   where σ_s is the standard error of Δr_s, which we calculate by combining the standard error of

429   the mean of the rate for each group using error propagation, i.e. $\sigma_s = \sqrt{\sigma_{s_1}{}^2 + \sigma_{s_2}{}^2}$, where σ_{s_N}

430   is the standard error of the occurrence rate of sequence s for group N.

431   In the case where the correlation is being calculated the studentized test statistic[3] is:

$$t_s = \frac{\sqrt{n}\hat{\rho}_n}{\hat{\tau}_n} \tag{8}$$

433   where:

$$\hat{\rho}_s = \frac{\sum X_i Y_i - n\bar{X}_s\bar{Y}}{\sqrt{\sum(X_i - \bar{X}_n)^2 \sum(Y_i - \bar{Y})^2}} \tag{9}$$

$$\hat{\tau}_s = \frac{\sqrt{\frac{1}{n}\sum_{i=1}^{n}(X_i - \bar{X}_s)^2(Y_i - \bar{Y})^2}}{\sqrt{\frac{1}{n}\sum_{i=1}^{n}(X_i - \bar{X}_s)^2}\sqrt{\frac{1}{n}\sum_{i=1}^{n}(Y_i - \bar{Y})^2}} \tag{10}$$

For eq. 8, 9, and 10, n is the number of subjects for which the correlation is being calculated, X_i
and Y_i and the values of the metrics being correlated (in our case, the sequence count and
factor 2 score for each individual respectively), $\bar{X}_s$ is the mean sequence count for the specific
sequence being considered, and $\bar{Y}$ is the mean of the covariate of interest (factor 2 score, which
is the same for any sequence).

When two groups are being compared, we resample from the entire population with
replacement (separately for each group) and build up a null distribution by bootstrapping M
times. The test statistic from the m^{th} bootstrap sample for $m = 1, \dots M$ is:

$$t_s^{*,m} = \frac{\Delta r_s^{*,m}}{\sigma_s^{*,m}} \tag{11}$$

where, $\Delta r_s^{*,m}$ is the difference in the rate of each sequence whilst resampling, with
replacement, from the entire population, ignoring the group labels. $\sigma_s^{*,m}$ is the accompanying
standard error of the resampled groups. The resampled group sizes are the same as the two
groups of interest.

In the case where the correlation is being calculated, the test statistic based on the m^{th}
bootstrap sample is:

$$t_s^{*,m} = \frac{\sqrt{n}\hat{\rho}_s^{*,m}}{\hat{\tau}_s^{*,m}} \qquad (12)$$

where, $\hat{\rho}_s^{*,m}$ is the correlations of each sequence whilst resampling, with replacement, from the
entire population, ignoring the group labels. $\hat{\tau}_s^{*,m}$ is the accompanying normalization factor of
the resampled groups. The resampled group size is the same as the original.

We used a value of $M = 10{,}000$. Importantly, for each individual resampling, m, the
same resampled set is used for all sequences.

The test statistics, and their accompanying estimators are ordered from largest to
smallest values. This creates a $M \times S$ matrix where each column contains all the estimators of
the test statistics. The first column contains the estimators from the largest test statistic, the
second column contains the estimators from the second largest, etc.

To define the distribution for which each test statistic is compared, which then
determines the adjusted p-value, the following algorithm is used. The first sequence considered
is the one with the maximum test statistic, t_s. Its comparison distribution is defined as the
maximum value within each row of the matrix of estimators of the test statistic:

$$\max(t^{*,m}) = \max\{t_1^{*,m}, \dots, t_S^{*,m}\} \qquad (13)$$

461 which provides a total of M values, $t^{*,m}$ (there is no longer an association with s, because these
462 values can come from a resampling of any of the sequences). Using those M values, the
463 adjusted p-value is calculated as follows:

$$p_s^{adj} = \frac{\#\{\max(t^{*,m}) \geq t_s\} + 1}{M + 1} \qquad (14)$$

After calculating p_s^{adj} for the first sequence, the column with the test statistic
estimators generated from the first sequence is removed from the matrix. This now leaves a
matrix that is $M \times (S - 1)$. The above procedure is then used to calculate p_s^{adj} for the
sequence with the second largest test statistic, and then its column of test statistic estimators is
removed, etc until all p_s^{adj} have been calculated. Following the algorithm described in Clarke et
al.[56], we enforce monotonicity of the p-values by resetting the p-value for each sequence:

$$p_s^{adj} = \max\{p_s^{adj}, p_{s-1}^{adj}\} \qquad (15)$$

This is done prior to calculating the p-value for the next sequence.

To control the false discovery proportion (i.e., control the number of false positives
divided by the total number of hypotheses rejected), the idea of k-FWER is introduced[2]. For

control of the FWER, k is equal to 1, and that leads to an α percent chance that there is at least
1 false positive among all hypotheses rejected (most commonly $\alpha = 0.05$). If k equals 2, then
there is an α percent chance that there are at least 2 false positives among all hypotheses
rejected. Therefore, to get control of the false discovery proportion we need to find the k that
provides the proportion of interest given the number of hypotheses rejected. So, if we want a
false discovery proportion, γ, of 0.05, we need $k \sim 0.05 \times$ number of hypotheses rejected.

Romano and Wolf also derived an algorithm to do just that. The algorithm is as follows.
Start with $k = 1$. Apply the k-FWER procedure, and note the total number of hypotheses
rejected, N. If $N < \frac{k}{\gamma} - 1$, stop and you have identified k. Otherwise, increase k by 1, and
repeat[2]. The way you determine k-FWER is in eq. 13, instead of taking the maximum value in
each row, you take the k[th] largest value. Finally, to get median control of the false discovery
proportion, $\alpha = 0.5$. This means that 50% of the time you will get a value greater than γ, and
50% of the time you will get a value less than γ, leading to control of the median[2]. This is a
similar decision to what is done when calculating the false discovery rate with Benjamini-
Hochberg or Benjamini-Yekutieli, except the false discovery rate controls the mean of the false
discovery proportion instead of the median.

Power estimation for CBAS. To estimate the statistical power of CBAS for a given sample size
we resampled the dataset without replacement and ran a CBAS to determine the number of
significant sequences. For each sample size we performed 20 repeats (Fig S1b, S2c, and S3b).
We also ran a CBAS comparing each group to itself (for the fly and rat datasets) (Fig S1b and
S3b), with 20 repeats for each group; or correlating the sequence counts with a randomly
generated set of factor 2 scores drawn from the same distribution as the actual factor 2 scores
(for the human dataset) (Fig S2c), with 40 repeats. Then for each sample size, the power is
estimated by identifying the largest 20 values of significant sequences, and determining the
fraction of those values that were generated by the comparison of the two groups or the
correlation with the data with the actual factor 2 scores.

Spatial alternation category fraction determination of significance. There are a total of 20
different sets of three arms, and 1 set of three arms with all choices being rewarded. For each
of the six contingencies the fraction of sequences that exclusively contains the set of three arms
that are significant is calculated separately for significant sequences greater in the WT and
greater in the *Scn2a*[+/-]. That means that there a total of $21 \times 2 \times 6 = 252$ fractions across all
contingencies. To determine whether the fraction is significantly larger than expected by
chance we proceeded as follows: Within a contingency, we determined the sequences that
belonged to any of the 21 categories. Then, separately for sequences greater in WT and
sequences greater in *Scn2a*[+/-], we permuted the association between those sequences and
whether or not they were significant. With that permuted association we recalculated the
fraction of sequences in each category that were significant. We repeated that process 25,000
times and determined significance by calculating the number of permuted fractions that were
greater than or equal to the actual fraction value for each category and divided that by 25,001
(as with eq. 15). That number was then corrected for multiple comparisons using the
Bonferroni method and multiplied by 252, the number of tests being performed. Any category
whose corrected p-value was < 0.05, was determined to be significantly larger than expected.

Data availability. Data will be made available upon reasonable request to the lead author.

Code availability. Code used to calculate CBAS will be posted to Github upon publication.

Acknowledgements. We thank Claire Gillan and Benjamin de Bivort for making their data
publicly available. We thank Thomas Akam and Adam Frank for helpful comments on the
manuscript. We thank the Simons Foundation Autism Research Initiative for supporting the
generation of the *Scn2a$^{+/-}$* rat line. This work was supported by grants from the Jane Coffin
Childs Memorial Fund for Medical Research (D.B.K.), the UCSF Physician Scientist Scholars
Program (D.B.K.), an NIH R25 (R25MH060482) (D.B.K), a Simons Foundation Autism Research
Initiative grant (899599) (D.B.K.), the Max Planck Society (P.D) and the Humboldt Foundation
(P.D.). Some of the data used in this study were collected in the laboratory of Loren Frank at the
University of California, San Francisco.

Author Contributions. D.B.K. and P.D. designed the study, D.B.K., J.P.R., and P.D. developed the
analysis method, D.B.K., G.W., and C.H. collected the data, D.B.K. analyzed the data, D.B.K and
P.D. wrote the manuscript, and D.B.K., J.P.R. and P.D. edited the manuscript.

Declaration of interests. The authors declare no competing interests.

1. Romano, J. P. & Wolf, M. Exact and Approximate Stepdown Methods for Multiple Hypothesis
Testing. *J Am Stat Assoc* **100**, 94–108 (2005).

2. Romano, J. P. & Wolf, M. Control of generalized error rates in multiple testing. *Ann Statistics*
35, 1378–1408 (2007).

3. DiCiccio, C. J. & Romano, J. P. Robust Permutation Tests For Correlation And Regression
Coefficients. *J. Am. Stat. Assoc.* **112**, 1211–1220 (2017).

4. Buchanan, S. M., Kain, J. S. & Bivort, B. L. de. Neuronal control of locomotor handedness in
Drosophila. *Proc National Acad Sci* **112**, 6700–6705 (2015).

5. Kastner, D. B. *et al.* Spatial preferences account for inter-animal variability during the
continual learning of a dynamic cognitive task. *Cell Reports* **39**, 110708–110708 (2022).

6. Gillan, C. M., Kosinski, M., Whelan, R., Phelps, E. A. & Daw, N. D. Characterizing a psychiatric
symptom dimension related to deficits in goal-directed control. *eLife* **5**, e11305 (2016).

7. Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A. & Poeppel, D.
Neuroscience Needs Behavior: Correcting a Reductionist Bias. *Neuron* **93**, 480–490 (2017).

8. Bialek, W. On the dimensionality of behavior. *P Natl Acad Sci Usa* **119**, e2021860119 (2022).

9. Meister, M. Learning, fast and slow. *Curr Opin Neurobiol* **75**, 102555 (2022).

10. Gomez-Marin, A. & Ghazanfar, A. A. The Life of Behavior. *Neuron* **104**, 25–36 (2019).

11. Trillmich, F., Günther, A., Müller, C., Reinhold, K. & Sachser, N. New perspectives in
behavioural development: adaptive shaping of behaviour over a lifetime? *Front Zool* **12**, S1
(2015).

12. Dayan, P., Roiser, J. P. & Viding, E. The first steps on long marches: the costs of active
observation. in *Rethinking Biopsychosocial Psychiatry.* (eds. Davies, W., Roache, R. & Savulescu,
J.) (Rethinking Biopsychosocial Psychiatry., 2018).

13. Hession, L. E., Sabnis, G. S., Churchill, G. A. & Kumar, V. A machine-vision-based frailty index
for mice. *Nat. Aging* **2**, 756–766 (2022).

14. Geuther, B. Q. *et al.* Action detection using a neural network elucidates the genetics of
mouse grooming behavior. *eLife* **10**, e63207 (2021).

15. Geuther, B. *et al.* High-throughput visual assessment of sleep stages in mice using machine
learning. *SLEEP* **45**, zsab260 (2021).

16. Sheppard, K. *et al.* Stride-level analysis of mouse open field behavior using deep-learning-
based pose estimation. *Cell Rep.* **38**, 110231 (2022).

17. Weinreb, C. *et al.* Keypoint-MoSeq: parsing behavior by linking point tracking to pose
dynamics. *bioRxiv* 2023.03.16.532307 (2023) doi:10.1101/2023.03.16.532307.

18. Wiltschko, A. B. *et al.* Mapping Sub-Second Structure in Mouse Behavior. *Neuron* **88**, 1121–
1135 (2015).

19. Pereira, T. D. *et al.* Fast animal pose estimation using deep neural networks. *Nat. Methods*
16, 117–125 (2019).

20. Pereira, T. D. *et al.* SLEAP: A deep learning system for multi-animal pose tracking. *Nat.*
Methods **19**, 486–495 (2022).

21. Marshall, J. D. *et al.* Continuous Whole-Body 3D Kinematic Recordings across the Rodent
Behavioral Repertoire. *Neuron* **109**, 420-437.e8 (2020).

22. Dunn, T. W. *et al.* Geometric deep learning enables 3D kinematic profiling across species
and environments. *Nat. Methods* **18**, 564–573 (2021).

23. Huys, Q. J. M., Maia, T. V. & Frank, M. J. Computational psychiatry as a bridge from
neuroscience to clinical applications. *Nat Neurosci* **19**, 404–413 (2016).

24. Kumar, P., Dayan, P. & Wolfers, T. From Complexity to Precision—Charting Decision-Making
Through Normative Modeling. *JAMA Psychiatry* **81**, (2024).

25. Nickerson, R. S. Confirmation bias: A ubiquitous phenomenon in many guises. *Rev Gen*
Psychol **2**, 175–220 (1998).

26. Kerr, N. L. HARKing: Hypothesizing After the Results are Known. *Pers Soc Psychol Rev* **2**,
196–217 (1998).

27. Dezfouli, A., Griffiths, K., Ramos, F., Dayan, P. & Balleine, B. W. Models that learn how
humans learn: The case of decision-making and its disorders. *PLoS Comput. Biol.* **15**, e1006903
(2019).

28. Eckstein, M. K., Summerfield, C., Daw, N. D. & Miller, K. J. Predictive and Interpretable:
Combining Artificial Neural Networks and Classic Cognitive Models to Understand Human
Learning and Decision Making. *bioRxiv* 2023.05.17.541226 (2023)
doi:10.1101/2023.05.17.541226.

29. Wiltschko, A. B. *et al.* Revealing the structure of pharmacobehavioral space through Motion
Sequencing. *Nat. Neurosci.* **23**, 1433–1443 (2020).

30. Gschwind, T. *et al.* Hidden behavioral fingerprints in epilepsy. *Neuron* **111**, 1440-1452.e5
(2023).

31. Uffelmann, E. *et al.* Genome-wide association studies. *Nat Rev Methods Primers* **1**, 59
(2021).

32. Farrell, M. S. *et al.* Evaluating historical candidate genes for schizophrenia. *Mol Psychiatr* **20**,
555–562 (2015).

33. Border, R. *et al.* No Support for Historical Candidate Gene or Candidate Gene-by-Interaction
Hypotheses for Major Depression Across Multiple Large Samples. *Am J Psychiat* **176**, 376–387
(2019).

34. Duncan, L. E., Ostacher, M. & Ballon, J. How genome-wide association studies (GWAS) made
traditional candidate gene studies obsolete. *Neuropsychopharmacol* **44**, 1518–1523 (2019).

35. Ioannidis, J. P. A. Why Most Published Research Findings Are False. *Plos Med* **2**, e124 (2005).

36. Markowitz, J. E. *et al.* Spontaneous behaviour is structured by reinforcement without
explicit reward. *Nature* **614**, 108–117 (2023).

37. Cui, T. *et al.* Gene–gene interaction detection with deep learning. *Commun. Biol.* **5**, 1238
(2022).

38. Fang, G. *et al.* Discovering genetic interactions bridging pathways in genome-wide
association studies. *Nat. Commun.* **10**, 4274 (2019).

39. Skutt-Kakaria, K., Reimers, P., Currier, T. A., Werkhoven, Z. & Bivort, B. L. de. A neural circuit
basis for context-modulation of individual locomotor behavior. *bioRxiv* 797126 (2019)
doi:10.1101/797126.

40. Ayroles, J. F. *et al.* Behavioral idiosyncrasy reveals genetic control of phenotypic variability.
Proc National Acad Sci **112**, 6706–6711 (2015).

41. Werkhoven, Z. *et al.* The structure of behavioral variation within a genotype. *eLife* **10**,
e64988 (2021).

42. Akhund-Zade, J., Ho, S., O'Leary, C. & Bivort, B. de. The effect of environmental enrichment
on behavioral variability depends on genotype, behavior, and type of enrichment. *J. Exp. Biol.*
222, jeb202234 (2019).

43. Bivort, B. de *et al.* Precise Quantification of Behavioral Individuality From 80 Million
Decisions Across 183,000 Flies. *Front. Behav. Neurosci.* **16**, 836626 (2022).

44. Daw, N. D., Gershman, S. J., Seymour, B., Dayan, P. & Dolan, R. J. Model-Based Influences on
Humans' Choices and Striatal Prediction Errors. *Neuron* **69**, 1204–1215 (2011).

45. Akam, T. *et al.* The Anterior Cingulate Cortex Predicts Future States to Mediate Model-
Based Action Selection. *Neuron* **109**, 149-163.e7 (2020).

46. Silva, C. F. da & Hare, T. A. Humans primarily use model-based inference in the two-stage
task. *Nat. Hum. Behav.* **4**, 1053–1066 (2020).

47. Kim, S. M. & Frank, L. M. Hippocampal Lesions Impair Rapid Learning of a Continuous Spatial
Alternation Task. *Plos One* **4**, e5494 (2009).

48. Singer, A. C., Carr, M. F., Karlsson, M. P. & Frank, L. M. Hippocampal SWR Activity Predicts
Correct Decisions during the Initial Learning of an Alternation Task. *Neuron* **77**, 1163–1173
(2013).

49. Kay, K. *et al.* Constant Sub-second Cycling between Representations of Possible Futures in
the Hippocampus. *Cell* **180**, 552-567.e25 (2020).

50. Fernández-Ruiz, A. *et al.* Long-duration hippocampal sharp wave ripples improve memory.
Science **364**, 1082–1086 (2019).

51. Sigurdsson, T., Stark, K. L., Karayiorgou, M., Gogos, J. A. & Gordon, J. A. Impaired
hippocampal–prefrontal synchrony in a genetic mouse model of schizophrenia. *Nature* **464**,
763–767 (2010).

52. Kastner, D. B., Gillespie, A. K., Dayan, P. & Frank, L. M. Memory alone does not account for
the speed of learning of a simple spatial alternation task in rats. *J Neurosci* **40**, JN-RM-0972-20
(2020).

53. Sanders, S. J. *et al.* De novo mutations revealed by whole-exome sequencing are strongly
associated with autism. *Nature* **485**, 237–241 (2012).

54. Satterstrom, F. K. *et al.* Large-Scale Exome Sequencing Study Implicates Both
Developmental and Functional Changes in the Neurobiology of Autism. *Cell* **180**, 568-584.e23
(2020).

55. Lord, C. *et al.* Autism spectrum disorder. *Nat. Rev. Dis. Prim.* **6**, 5 (2020).

56. Clarke, D., Romano, J. P. & Wolf, M. The Romano-Wolf Multiple Hypothesis Correction in
Stata. *Ssrn Electron J* (2020) doi:10.2139/ssrn.3513687.

Figure 1. Choice is a common discretization for behavior. In the three behavioral tasks under consideration in this work, the actions of the subjects are broken down into a series of choices. (**a**) Fly y-maze (top), and example set of choices of left (red) and right (blue) turns (bottom) of an individual fly. Data come from Buchanan et al.[4]. (**b**) Two-step task, performed by human subjects (top), and an example set of choices of an individual subject (bottom). The colors correspond to the different objects chosen (bottom). Data come from Gillan et al.[6]. (**c**) Spatial alternation behavior, performed by rats (top), and an example set of choices of an individual rat (bottom). The contingency is defined as the 3 arms that can be rewarded, and the 6-arm track enables different sets of three arms to be rewarded. In **b** and **c,** filled in circles indicate that reward was received.

Figure 2. CBAS applied to y-maze produces interpretable differences between fly strains.
CBAS was applied to flies tracked on a y-maze (Fig 1a) (**a**) This table demarcates the decisions
made to perform the CBAS on this dataset. There was a total of 1,225 Cambridge-A (CA) and
1,372 w1118 flies in the dataset, 466 CA and 565 w1118 were excluded from analysis due to
not reaching a total of at least 250 turns. (**b**) The occurrence of a single sequence of turns (10
right turns in a row) in the two strains during the first 250 turns. The rows show the occurrence
of that sequence of turns for all individual flies from each strain. The CA strain has a rate of 9.3,
meaning that, on average, each fly utilizes this sequence 9.3 times. The W1118 strain has a rate
of 2.1, meaning that, on average, each fly utilizes this sequence 2.1 times. (**c**) CBAS Manhattan
plot displays the p-value for each sequence. Each sequence has two p-values on this plot, one
for CA > w1118, and the other for CA < w1118. The sequences are ordered based on the
number of choices in the sequence, and they are displayed on a log scale to make all sequence
lengths visible. Within a given sequence length, the sequences are ordered based on frequency
of occurrence in the entire dataset. The horizontal dotted line indicates the significance
threshold of 5% control of the median false discovery proportion. A total of 10,000 resamplings
were used to calculate the p-values making the maximum value on the plot 4.00004
$(-\log(\frac{1}{10,001}))$. (**d**) Every sequence that either has a maximum of 1 turn in a direction in a row
(left) or a maximum of 7 turns in the same direction in a row (right). Sequences are colored
based on whether CBAS identifies them as occurring significantly more in the CA strain (purple),
w1118 strain (green) or as not significantly different (black). The sequences on the right are
aligned to the 7 right or left turns. (**e**) The maximum number of the same turns in a row was

calculated for every sequence in the dataset (2,046 total sequences), and for every number of
maximum turns in a row, the fraction of sequences that whose prevalence was significantly
greater in the CA strain (purple) or the w1118 strain (green) is plotted. The CA and w1118
strains showed separation with other related metrics even for the max turns that show both
significance for both direction of comparison (e.g. max same turn in a row of 4) (Fig S1a). (**f**)
Power estimate for different sized groups of flies in each strain (see methods). Horizontal
dotted line shows a value of 80%.

Figure 3. Common, but degraded output with CBAS on smaller sample sizes. For all panels in this figure, many repeats of smaller sample sizes were generated from the fly data used in Fig 2 by resampling subjects (without replacement within a group) from the full dataset. (**a**) Each CBAS run on the smaller sample size identified some number of significant sequences. Those sequences were compared to the sequences identified in the CBAS on the full dataset, and the graph shows the ratios of the number of sequences identified by the smaller sample size that were also in the full dataset to the total number of sequences identified in the full dataset. As a comparison, the ratio of the number of flies in the smaller CBAS to the total number of flies in the dataset is plotted in the maroon horizontal lines. (**b**) The number of significant sequences identified in the smaller sample size that were not also identified in the full dataset is plotted over the total number of significant sequences identified in the smaller sample size. (**c**) In creating the smaller samples sizes, 20 paired sets of animals were generated that had no overlapping individuals. The number of the same significant sequences that were identified with these nonoverlapping sets of flies is plotted over the average number of significant sequences in the pair. (**d**) For each repeat of each sample size, all of the sequences were

categorized based on the maximum number of turns in the same direction and the fraction of
significant sequences within those categories are plotted (as in Fig 2e). The bottom row of this
plot is from the full dataset and is identical to Fig 2e. (**e**) The Euclidean distance between each
row from panel **d** and the full dataset row is plotted. Colors correspond to the sample sizes as
shown in panel **d**. For panels **a** – **c** data points are overlayed by box and whisker plots. The
center line of the box displays the median of the data, the top and bottom lines of the box
show the 25th and 75th quartiles, respectively, and the end of the whiskers show the full range
of the data.

Figure 4. CBAS identifies unexpected sequences in human dataset. CBAS was applied to
humans performing the two-step task (Fig 1b) (**a**) This table demarcates the decisions made to
perform the CBAS on this dataset. There was a total of 1,413 human subjects in the dataset. (**b**)
The occurrence of a single sequence of choices ('1**A**2', meaning choosing object 1, then getting
rewarded for a choice in set A, then choosing object 2) across all subjects in the dataset.
Subjects are ordered based on their factor 2 ("intrusive thoughts and compulsive behaviors")
score. This sequence shows a correlation (ρ) of 0.12 with factor 2 score from the questionnaire
factor analysis (see S2a). (**c**) CBAS Manhattan plot displays the p-value for each sequence. Each
sequence has two p-values on this plot, one for positive correlation and one for negative
correlation. The sequences are ordered based on the number of choices in the sequence and
are displayed on a log scale to make all sequence lengths visible. Within a given sequence
length, the sequences are ordered based on frequency of occurrence in the entire dataset. The
horizontal dotted line indicates the significance threshold of 5% control of the median false
discovery proportion. A total of 10,000 resamplings were used to calculate the p-values making
the maximum value on the plot 4.00004 ($-\log\left(\frac{1}{10,001}\right)$). (**d**) Every sequence that was
significantly positively (left) or negatively (right) correlated with factor 2 score. The sequences
on the left are aligned to **B**1 or **A**2, and sequences on the right are aligned to **A**1. (**e**) For the
categories listed (2**B**1, 1**B**1, 1**A**2, 2**A**2), the number of significant sequences that contain the
category is plotted over total number of sequences that exist in the dataset that contain the
category. For 2**A**2 no subject in the dataset performed that sequence. (**f**) Power estimate for
different sized groups of subjects (see methods). Horizontal dotted line shows a value of 80%.

Figure 5. CBAS provide interpretable information to phenotype *Scn2a*^{+/-} rats. CBAS was
applied to rats performing a spatial alternation task (Fig 1c) (**a**) This table demarcates the
decisions made to perform the CBAS on this dataset. There was a total of 121 WT and 120
Scn2a^{+/-} rats for the analyses (see methods). (**b**) The occurrence of a single (unrewarded)
sequence of arm choices (642424; all from contingency D which rewards 246) in the two rat
genotypes during contingency E, which is rewarded at arms 234. The rows show the occurrence
of that sequence of choices for all individual rats from each genotype. WT rats used this
sequence at a rate of 0.12, meaning that on average about 1 out of every 10 rats used this
sequence. *Scn2a*^{+/-} rats used this sequence at a rate of 0.45, meaning that on average 1 out of
every 2 rats used this sequence. (**c**) CBAS Manhattan plot displays the p-value for each
sequence. Each sequence has two p-values on this plot, one for WT > *Scn2a*^{+/-}, and the other for
WT > *Scn2a*^{+/-}. The sequences are ordered based on the number of choices in the sequence, and
within each contingency are displayed on a log scale to make all sequence lengths visible.
Within a given sequence length, the sequences are ordered based on frequency of occurrence
in the entire dataset. The rewarded arms for each contingency are shown below the letter
symbol for each contingency. The horizontal dotted line indicates the significance threshold of
5% control of the median false discovery proportion. A total of 10,000 resamplings were used
to calculate the p-values making the maximum value on the plot 4.00004 ($-\log(\frac{1}{10,001})$). (**d**)

Every significant sequence that exclusively contains arms 1, 2, 3 (irrespective of reward) during
contingency B (left) or every significant sequence that exclusively contains arms 2, 4, 6
(irrespective of reward) during contingency E (right). Sequences that occur significantly more in
Scn2a$^{+/-}$ are shown in red, and those that occur significantly more in WT are shown in black. (**e**)
The sets of 3 arms which show a significantly greater fraction of significant sequences than
would be expected by chance (see methods). The categories above each group of bar plots
indicate the structure of the set of 3 arms. "Current" indicates the 3 arms from the current
contingency irrespective of reward, "current rewarded" indicates the 3 arms from the current
contingency all of which are rewarded, "1 back" indicates the 3 arms from the prior
contingency, "2 back" indicate the 3 arms from 2 contingencies prior, and "other" are
categories that show significant fractions that do not fit the other categories. Of note, the
sequences in the "current rewarded" category are a subset of the sequences in the "current"
category, as can be seen in panel **d**, left. (**f**) Power estimate for different sized groups of rats in
each genotype (see methods). Horizontal dotted line shows a value of 80%.

Supplementary Figure 1. (**a**) All sequences with a maximum length of 4 turns in the same direction in a row (Fig 2e) were evaluated for the fraction of L turns in the sequence (encompassing L = 4 – 8) and the rate of switching (number of changes in turn direction divided by the total number of turns in the sequence). These two metrics show a separation in the sequences that were significantly greater in the CA strain compared to those that were significantly greater in the w1118 strain: the CA strain had sequences with more extreme fraction of left turns and lower switch rate, consistent with more turns in the same direction. (**b**) Left: the number of significant sequences when randomly resampling the populations without replacement and calculating CBAS on the smaller sample sizes. Middle: the number of significant sequences when comparing smaller samples sizes of the CA strain to itself, with nonoverlapping individuals in each group. Right: the number of significant sequences when comparing smaller samples sizes of the w1118 strain to itself, with nonoverlapping individuals in each group. Data points are overlayed by box and whisker plots. The center line of the box displays the median of the data, the top and bottom lines of the box show the 25th and 75th quartiles, respectively, and the end of the whiskers show the full range of the data.

Supplementary Figure 2. (**a**) Correlation between the sequences count for each human subject
and their factor 2 score for sequence: 1**A**2, which means choosing object 1, then making a
choice in set A and getting rewarded, and then choosing object 2. Factor 2 reflects the intrusive
thoughts and obsessive behavior loading on the psychiatric symptom questionnaires. Red line
shows the linear fit to the data. (**b**) The correlation plotted as a function of the average
sequence rate for all sequences containing the unit 2**B**1 (left), 1**B**1 (middle), or 1**A**2 (right).
Filled in circles indicate those sequences that show a significant positive correlation. (**c**) Left:
the number of significant sequences when randomly resampling the populations without
replacement and calculating CBAS on the smaller sample sizes. Right: the number of significant
sequence when randomly resampling the populations without replacement and comparing it to
randomly generated factor 2 scores drawn from a distribution imputed from the original factor
2 scores and calculating CBAS on the smaller sample sizes. Data points are overlayed by box and
whisker plots. The center line of the box displays the median of the data, the top and bottom
lines of the box show the 25th and 75th quartiles, respectively, and the end of the whiskers show
the full range of the data.

Supplementary Figure 3. (**a**) For the sequences that exclusively contain all sets of 3 arms (Fig
5d&e) the fraction of the total number of sequences is plotted in black for each contingency,
and the fraction of significant sequences that are a part of the sequences that exclusively
contain all sets of 3 arms compared to the total number of significant sequences in each
contingency is plotted in green. (**b**) Left: the number of significant sequences when randomly
resampling the populations without replacement and calculating CBAS on the smaller sample
sizes. Middle: the number of significant sequence when comparing smaller samples sizes of the
WT genotype to itself, with nonoverlapping individuals in each group. Right: the number of
significant sequence when comparing smaller samples sizes of the $Scn2a^{+/-}$ genotype to itself,
with nonoverlapping individuals in each group. Data points are overlayed by box and whisker
plots. The center line of the box displays the median of the data, the top and bottom lines of
the box show the 25[th] and 75[th] quartiles, respectively, and the end of the whiskers show the full
range of the data.