The Bacterial Quorum-Sensing Signal 2-Aminoacetophenone Rewires Immune Cell

Bioenergetics through the PGC-1α/ERRα Axis to Mediate Tolerance to Infection

Arijit Chakraborty[1,2,3#], Arunava Bandyopadhaya[1,2,&#], Vijay K Singh[1,2], Filip Kovacic[1,3,4], Sujin

Cha[1], William M. Oldham[5], A. Aria Tzika[1,2], and Laurence G Rahme[1,2,3*]

[1]Department of Surgery, Massachusetts General Hospital, and Harvard Medical School,

Boston, Massachusetts, USA

[2]Shriners Hospitals for Children Boston, Boston, Massachusetts, USA

[3]Department of Microbiology, Harvard Medical School, Boston, Massachusetts, USA

[4]Institute of Molecular Enzyme Technology, Heinrich Heine University Düsseldorf,

Forschungszentrum Jülich GmbH, Jülich, Germany

[5]Department of Medicine, Brigham and Women's Hospital and Harvard Medical School,

Boston, Massachusetts, USA

[&] Present Address: Astellas Pharma Inc., USA

[#]These authors contributed equally

[*]**Corresponding author:** Laurence G. Rahme, Ph.D

e-mail: rahme@molbio.mgh.harvard.edu (LGR)

Address: 340 Their Research Building, 50 Blossom Street, Massachusetts General Hospital,
Boston, MA 02114

Phone: 617-724-5003

Author Contributions:

Author contributions: L.G.R. conceptualization, A.B., A.C. and L.G.R., designed research;

A.B., A.C. V.K.S., S. C., performed research; A.B., A.C. V.K.S., F.K., and L.G.R. analyzed

data; W.M.O and A.A.T., provided resources, and F.K. and L.G.R. wrote the paper.

Classification:

Biological, health, and Medical Sciences/Microbiology

Keywords: Pseudomonas aeruginosa, quorum sensing, infection, macrophages, bacterial

persistence, 2-aminoacetophenone (2-AA), MvfR, PqsR, immunometabolism, host

tolerization, histone deacetylation, epigenetic reprogramming, metabolic reprogramming,

oxidative phosphorylation**,** pyruvate, mitochondrial pyruvate carrier (MPC1), peroxisome

proliferator-activated receptor gamma co-activator 1-alpha (PGC-1α), estrogen-related

receptor (ERRα), acetyl-CoA, ATP, bioenergetics.

Abstract

How bacterial pathogens exploit host metabolism to promote immune tolerance and persist in infected hosts remains elusive. To achieve this, we show that *Pseudomonas aeruginosa (PA),* a recalcitrant pathogen, utilizes the quorum sensing (QS) signal 2-aminoacetophenone (2-AA). Here, we unveil how 2-AA-driven immune tolerization causes distinct metabolic perturbations in macrophages' mitochondrial respiration and bioenergetics. We present evidence indicating that these effects stem from decreased pyruvate transport into mitochondria. This reduction is attributed to decreased expression of the mitochondrial pyruvate carrier (MPC1), which is mediated by diminished expression and nuclear presence of its transcriptional regulator, estrogen-related nuclear receptor alpha (ERRα). Consequently, ERRα exhibits weakened binding to the MPC1 promoter. This outcome arises from the impaired interaction between ERRα and the peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1α). Ultimately, this cascade results in diminished pyruvate influx into mitochondria and, consequently reduced ATP production in tolerized macrophages. Exogenously added ATP in infected macrophages restores the transcript levels of *MPC1* and *ERR-α and* enhances cytokine production and intracellular bacterial clearance. Consistent with the *in vitro* findings, murine infection studies corroborate the 2-AA-mediated long-lasting decrease in ATP and acetyl-CoA and its association with *PA* persistence, further supporting this QS signaling molecule as the culprit of the host bioenergetic alterations and *PA* persistence. These findings unveil 2-AA as a modulator of cellular immunometabolism and reveal an unprecedented mechanism of host tolerance to infection involving the PGC-1α/ERRα axis in its influence on MPC1/OXPHOS-dependent energy production and *PA* clearance. These paradigmatic findings pave the way for developing treatments to bolster host resilience to pathogen-induced damage. Given that QS is a common characteristic of prokaryotes, it is likely that 2-AA-like molecules with similar functions may be present in other pathogens.

Introduction

Host tolerance is a fundamental mechanism of innate immunity. Studies have shown that innate immune cells following infection or exposure to microbial products may enter a state of immune tolerance characterized by reduced responsiveness towards microbial re-exposure a few hours later [1-3]. Epigenetic and signaling-based mechanisms have been proposed to be involved in host immune tolerance [1-3]. In recent years, several reports have highlighted the complex interplay between metabolic reprogramming and immunity [4, 5]. It was suggested that specific metabolic programs are activated in monocytes and macrophages upon exposure to infection and microbial products [6, 7]. However, in host tolerance, the molecular mechanisms underlying immunometabolic reprogramming mediated by bacterial pathogens remain poorly understood.

Metabolic pathways are essential for generating energy for various cellular functions, including those performed by immune cells [8, 9]. Macrophages use metabolic pathways to generate energy and metabolites to adapt to changing environments and stimuli, thereby enabling them to cope with the needs of a fluctuating immune response [7, 10]. Thus, properly functioning metabolic pathways are vital for immune cells to counteract pathogens. For instance, the crucial energy-carrying molecule adenosine triphosphate (ATP) is required to phagocytose pathogen-derived molecules efficiently.

Pseudomonas aeruginosa (PA), a recalcitrant ESKAPE (*Enterococcus faecium, Staphylococcus aureus, Klebsiella pneumoniae, Acinetobacter baumannii, PA,* and *Enterobacter sp.*) pathogen that causes acute and persistent infections, secretes virulence-associated low molecular weight signaling molecules, several of which are regulated by quorum sensing (QS) [11-14] and able to modulate host immune responses [15-18]. QS is a cell density-dependent signaling system bacteria use to synchronize their activities [11-14]. MvfR (a.k.a. PqsR), a critical QS transcription factor of *PA,* regulates the synthesis of many small molecules, including signaling molecules such as 2-aminoacetophenone (2-AA) [11, 19-21]. *In vivo* studies have demonstrated that 2-AA enables *PA* to persist in infected murine

tissues by promoting innate immune tolerance through histone deacetylase 1 (HDAC1)-mediated epigenetic reprogramming [17, 18]. 2-AA tolerization reprograms the host inflammatory signaling cascade by maintaining chromatin in a "silent" state through increased HDAC1 expression and activity and decreased histone acetyltransferase (HAT) activity [18]. These changes also impact the protein-protein interaction between HAT and cyclic AMP response element-binding protein/HDAC1 and p50/p65 nuclear factor (NF)-κB subunits [18, 22]. The 2-AA-mediated tolerization permits *PA* to persist in infected tissues [17, 18], underscoring the difference from lipopolysaccharide (LPS)-mediated tolerization, which instead leads to bacterial clearance and involves different HDACs [23].

Mitochondria, the "powerhouse" of the cells, are crucial for the regulation, differentiation, and survival of macrophages and other immune cells [8]. Our group's previous *in vivo* and *in vitro* studies indicated that 2-AA affects metabolic functions in skeletal muscle, which contains a high concentration of mitochondria [24-26]. Injection of 2-AA in murine skeletal muscle dampens the expression of genes associated with OXPHOS and the master regulator of mitochondrial biogenesis peroxisome proliferator-activated receptor-γ coactivator-1 beta (PGC-1β) [24]. Another *PA* QS molecule, 3-oxo-C12-HSL, attenuates the expression of PGC-1α [27]. PGC-1α and PGC-1β belong to the PPARγ family of inducible transcriptional coactivators and are known regulators of mitochondrial metabolism [28]. The coactivator PGC-1 proteins are involved in various cellular energy metabolic processes [29, 30], including but not limited to mitochondrial metabolism [31]. Recent studies have also emphasized the role of estrogen-related nuclear receptor α (ERRα) in coordinating metabolic capacity with energy demand in health and disease [32]. PGC-1α activates ERRα transcriptional activity through protein-protein interaction, which enhances the expression of ERRα [33, 34] and other mitochondrial genes involved in lipid metabolism and OXPHOS [35]. PGC-1α/ERRα axis has been extensively studied in cancer and established as a central regulatory node of energy metabolism that induces the global expression of genes involved in mitochondrial biogenesis and functions [36-38]. ERRα has been shown to occupy the

promoter regions of many genes participating in the TCA cycle and OXPHOS [39, 40],

including the mitochondrial pyruvate carrier (MPC1). In human renal carcinoma cells, PGC-

1α/ERRα interaction results in efficient activation of MPC1 expression and transport of

pyruvate into mitochondrion for efficient OXPHOS and energy production [41].

Given that our previous studies pointed to the 2-AA effect on mitochondrial functions

[24-26] and that mitochondria are crucial for the regulation, differentiation, and survival of

macrophages and other immune cells [8], we investigated how 2-AA may mediate cellular

metabolic reprogramming in tolerized immune cells by examining the link between the PGC-

1α/ERRα axis and 2-AA-mediated immune tolerance. Our findings uncovered the crucial

action of 2-AA on energy homeostasis and metabolism in tolerized immune cells through

perturbances of the ERRα/PGC-1α interaction, MPC1-mediated pyruvate transport and the

production of the key energy metabolism molecules, ATP and acetyl-CoA and their

association with the persistence of *PA* in mammalian tissues.

Results

Tolerization by 2-AA impacts the generation of crucial energy metabolites in

macrophages

Following infection or exposure to microbial products, innate immune cells may enter

a state of tolerance several hours later, as depicted by their diminished responsiveness or

unresponsiveness to microbial re-exposure [1-3, 42]. Given that 2-AA tolerized cells are non-

responsive to 2-AA re-exposure, we investigated the impact of 2-AA on energy homeostasis

and metabolism in non-tolerized and tolerized BMDM cells (Fig. 1). We quantified the levels

of ATP and acetyl-CoA, which is a fuel for ATP production, in a series of *in vitro* experiments

in which murine BMDM cells were exposed to 2-AA (1st 2-AA exposure) or re-exposed (2nd 2-

AA exposure) to respectively determine their activation, and tolerance as well as memory to

this QS molecule (Fig. 1A). Initially, following the 1st 2-AA exposure (black bars), a significant

increase in the concentrations of intracellular ATP (Fig. 1B) and acetyl-CoA (Fig. 1C) was

observed in BMDM cells exposed to 2-AA for 1 and 6 h compared to their corresponding

naïve control cells (grey bars). However, cells exposed to 2-AA for 48 h (black bar) appeared

to have entered a state of tolerance as ATP and acetyl-CoA levels were decreased compared

to 2-AA exposures for a shorter time (Figs. 1B and 1C). The tolerance of these cells as well

as their memory to 2-AA was determined by their responsiveness to a 2nd 2-AA exposure

(Fig. 1A). To achieve this, these cells were washed to remove 2-AA and allowed to rest for 24

or 106 h in the absence of 2-AA before receiving the 2nd 2-AA exposure for 1 or 6 h (Fig. 1A).

Figures 1B and 1C (red bars) show the unresponsiveness of BMDM cells to 2-AA re-

exposure as evidenced by the similar levels of ATP and acetyl-CoA compared to cells

exposed to 2-AA for 48 h (1st exposure black bars). The sustained unresponsiveness of the

cells to re-exposure indicates that the cells exhibited innate immunological memory, which

could not be reverted by 2-AA re-exposure. Similar responsiveness patterns of ATP and

acetyl-CoA as with BMDM cells were observed in murine macrophage RAW 264.7 cells

(Figs. S1A and S1B) or human monocyte THP-1 cells (Figs. S1C and S1D) following

exposure or re-exposure to 2-AA. These findings further support our observations with

BMDM cells and confirm that either cell type can be used to study responses to 2-AA. These

results reinforce the notion that 2-AA impacts crucial energy metabolites in macrophages.

Moreover, we sought to determine whether 2-AA-tolerized BMDM cells would respond

to heterologous stimulus. To test this, we re-exposed the cells to lipopolysaccharide (LPS),

an outer membrane bacterial component that strongly induces macrophages. LPS re-

exposure of the 2-AA tolerized BMDM cells did not significantly ($p = 0.9$) increase ATP levels

(Fig. S2), indicating that 2-AA cross-tolerized the cells to this heterologous bacterial

immunostimulant. On the other hand, as expected, we observed increased ATP levels

following LPS stimulation of the non-tolerized BMDM cells (Fig. S2). The broad spectrum of

2-AA cross-tolerization strongly suggests an interplay of epigenetic mechanism rather than

ligand-receptor mediated signaling-based mechanism in bringing out the observed 2-AA-

mediated tolerance.

Tolerization by 2-AA leads to a quiescent state by reducing cell bioenergetics and
compromising OXPHOS.

To further investigate the 2-AA's impact on the bioenergetics of macrophages, we

assessed by Seahorse assay the oxygen consumption rate (OCR) as an index of OXPHOS

in 2-AA exposed (for 1 and 24 h) and re-exposed (2^{nd} exposure for 1 h) BMDM cells (Fig.

1D). Values of basal (Fig. 1E) and maximal (Fig. 1F) mitochondrial respiration and spare

respiratory capacity (Fig. S3) interpreted from these measurements revealed that 1 h 2-AA

exposure of naïve cells significantly increased OCR compared to naïve control cells. At 24 h

post-2-AA exposure, cells had a significantly reduced basal OCR level than the control

BMDM cells, indicating decreased OXPHOS and an overall quiescent phenotype of

tolerized macrophages. Re-exposure of tolerized macrophages with 2-AA also did not

augment maximal and basal OCR (Figs. 1D-1F), indicating the unresponsiveness of

tolerized BMDM cells. Ultramicroscopic examination of RAW 264.7 cells exposed to 2-AA

for 48 h although shows the same number of mitochondria as in control cells, their

mitochondrial morphology is altered, appearing smaller and round and having reduced

cristae indicating dysfunctional mitochondria (Fig. S4).

These OCR data are consistent with previous direct measurements of ATP (Fig. 1B)

and acetyl-CoA (Fig. 1C) and support the notion that 2-AA tolerization induces a quiescent

phenotype in these cells characterized by defective OXPHOS. These findings confirm that

2-AA mediates energy homeostatic and metabolic alterations in the immune cells and that 2-

AA tolerization arrests these cells in a sustained 2-AA-unresponsive state.

The pyruvate transport into mitochondria is decreased in tolerized macrophages.

Since acetyl-CoA and ATP levels were reduced due to 2-AA tolerization, we

investigated whether the mitochondrial dysfunction observed is related to pyruvate

metabolism. Pyruvate links glycolysis with mitochondrial production of acetyl-CoA and ATP

via the tricarboxylic acid (TCA) cycle and OXPHOS [7]. RAW 264.7 cells initially exposed to

2-AA for 1 or 3 h (black bars) exhibited significantly higher concentrations of cytosolic and

mitochondrial pyruvate compared to the corresponding naïve control cells (Fig. 2A). Pyruvate

significantly decreased over time in cytosolic and mitochondrial fractions as cells were

entering into the tolerized state (Fig. 2A). Tolerized cells remain unresponsive to 2-AA re-

exposure (red bars), exhibiting no significant change in the concentration of pyruvate in the

cytosolic or mitochondrial fraction (Fig. 2A). A similar result was observed in THP-1 cells

(Figs. S1E and S1F).

To gain a better understanding of the 2-AA-mediated reductions in mitochondrial

pyruvate, ATP production, and acetyl-CoA levels, we sought to determine the involvement of

the MPC1 [43] that is crucial in transporting pyruvate from the cytosol into the mitochondria

[44]. Western blot studies of whole cell lysates of RAW 246.7 cells exposed to 2-AA showed

a significant decrease over time in the abundance of MPC1 protein, and tolerized cells

remain unresponsive to 2-AA re-exposure (Fig. 2B). Together, these findings suggest that 2-

AA tolerization dysregulates MPC1-mediated transport of the pyruvate into mitochondria.

216   2-AA tolerization suppresses ERRα binding to the *MPC1* promoter and the interaction

217   between PGC-1α and ERRα

Given that the PGC-1α/ERRα axis is the transcriptional regulatory node that activates

the expression of MPC1 [33], we determined the protein levels of ERRα. Western blot

studies of whole cell lysates showed a significant decrease over time in the abundance of

ERRα protein following 1[st] exposure of RAW 246.7 cells to 2-AA, while tolerized cells remain

unresponsive to 2-AA re-exposure (Fig. 3A). By assessing the cytosolic and nuclear

abundance of ERRα in tolerized cells we found that both fractions had lower levels of ERRα

protein than their corresponding control cells (Fig. 3B).

Furthermore, because an ERRα binding putative DNA motif (TNAAGGTCA) has been

identified upstream of the *MPC1* promoter in humans [43], and we found the same motif to

be present 1.5 kB upstream of the transcription start site of mouse *MPC1* promoter, we

analyzed if tolerization led to reduced ERRα binding to the *MPC1* promoter motif

TNAAGGTCA. Chromatin immunoprecipitation (ChIP) assay followed by qPCR analysis

revealed that in tolerized cells, ERRα binds approximately 4-fold less efficiently to the

putative binding site on the *MPC1* than in corresponding control cells (Fig. 3C).

Since the transcriptional activity of ERRα is regulated through a protein-protein

interaction with the transcriptional coactivator PGC-1α [33], we assessed whether the

formation of the PGC-1α/ERRα complex is affected in 2-AA tolerized cells. Using

paraformaldehyde fixed nuclear lysates of RAW 264.7 tolerized and naïve cells, we

performed a co-immunoprecipitation (co-IP) of PGC-1α and reversed co-IP with ERRα

followed by immunoblotting (Fig. 3D). As shown in Figure 3D less ERRα was detected when

PGC-1α was pulled down in presence of 2-AA than in naïve control cells. Moreover, reverse

co-IP with ERRα protein showed lower levels of PGC-1α in the presence of 2-AA than in

naïve control cells (Fig. 3D). Control IgG showed no detectable levels of ERRα following

nuclear fraction pull down. These results indicate that in the tolerized macrophages, PGC-

1α/ERRα interaction is impaired (Fig. 3D).

Moreover, since PGC-1α enhances the transcriptional activity of ERRα through their

interaction, and ERRα transcription is regulated *via* an autoregulatory loop [34], we examined

the effect of tolerization on the transcription of *PGC-1α, ERRα,* and its target gene *MPC1*.

Using RAW 264.7 cells infected with the wild type *PA* strain PA14 or isogenic Δ*mvfR* mutant*,*

which does not produce 2-AA [21] we observed lower RNA transcript levels of *MPC1* (Fig.

4A)*, ERRα* (Fig. 4B) and *PGC-1α* (4C) in PA14 compared to Δ*mvfR.* The addition of 2-AA to

Δ*mvfR* decreased the levels of these gene transcripts to levels similar to the PA14 infection

condition, supporting the role of 2-AA in effect. To test if the observed effects are due to

MPC1-dependent reduction of mitochondrial ATP generation, we supplemented the

macrophages with ATP. The findings indicate that the addition of ATP in PA14-infected cells

elevated the transcript levels of *ERRα*, *MPC1*, and *PGC-1α,* reaching levels similar to those

observed in Δ*mvfR* infected macrophages (Fig. 4A, 4B, and 4C). These results indicate that

reduced MPC1 function is due to 2-AA tolerization on the transcriptional activation of *MPC1*

through the PGC-1α/ERRα axis.

2-AA tolerization impairs macrophage-mediated intracellular bacterial clearance

through decrease in MPC1-mediated pyruvate import, ATP, and TNF-α levels.

2-AA-mediates persistence of *PA in vivo,* dampens proinflammatory responses, and

increases the intracellular burden of this pathogen in macrophages *via* epigenetic

modifications [17, 18, 26]. Here, we used RAW 264.7 cells infected with the PA14 or the 2-AA

deficient Δ*mvfR* mutant to assess the clearance of *PA* by macrophages. Macrophages

infected with PA14 showed increased bacterial burden than the cells infected with Δ*mvfR,*

and 2-AA addition to Δ*mvfR* led to increased bacterial burden (Fig. 4D). Furthermore, we

used UK5099 and ATP to interrogate whether MPC1 mediated mitochondrial pyruvate import

and bioenergetics are linked to the clearance of *PA* intracellular burden. The addition of the

UK5099 inhibitor strongly enhanced the bacterial intracellular burden in Δ*mvfR* infected

macrophages compared to the non-inhibited Δ*mvfR* infected cells, reaching a similar burden

to those infected with PA14 (Fig. 4D). Conversely, exogenously added ATP to macrophages

infected with PA14 strongly reduced the *PA* intracellular burden (Fig. 4D).

Given that 2-AA tolerization decreases the expression of pro-inflammatory cytokine

TNF-α by hypo-acetylating the core histone 3 lysine 18 acetylation (H3K18ac) mark at TNF-α

promoter [18], we investigated the link between bioenergetics and TNF-α expression in

infected macrophages. UK5099, or ATP was added exogenously to infected macrophages

with PA14 or ∆*mvfR* (Fig. 4E). As shown in Figure 4E, PA14 infected cells showed lower

TNF-α transcript levels compared to the ∆*mvfR* infected cells. Supplementation of 2-AA to

∆*mvfR* infected cells led to a decrease in the TNF-α transcript levels (Fig. 4E). The addition of

the MPC1 inhibitor, UK5099 in ∆*mvfR* infected cells counteracted the increase in TNF-α

transcript levels observed in ∆*mvfR* infected cells in the absence of UK5099 (Fig. 4E).

Conversely, exogeneous addition of ATP to PA14 infected cells enhanced the transcription of

TNF-α compared to the untreated PA14 infected cells, while no difference in TNF-α

expression levels were observed in ∆*mvfR* infected macrophages in presence or absence of

ATP (Fig. 4E).These findings strongly suggest that 2-AA tolerization severely alters

macrophages' ability to facilitate the clearance of *PA* intracellular burden, *via* the reduction in

MPC1-mediated pyruvate import into mitochondria, ATP levels and TNF-α transcription.

***In vivo* studies corroborate the 2-AA-mediated decrease *in vitro* of the central**

metabolic fuel acetyl-CoA and the energy-carrying molecule ATP and their association

with *PA* persistence.

To determine whether the decrease in the key metabolites mediated by 2-AA is also

observed *in vivo* during infection, we quantified the levels of ATP and acetyl-CoA in murine

spleen tissues at 1, 5, and 10 days (Fig. 5). This organ was selected for our

immunometabolic studies due to its role in regulating not only local but also systemic (whole-

body) immune responses, facilitated by various immune cells including macrophages [45].

Mice were infected with *PA* strain PA14, or isogenic mutant ∆*mvfR,* which does not produce

2-AA [21]. It is important to note that it is not possible to generate or utilize a bacterial mutant

that is defective in 2-AA only because 2-AA is formed by spontaneous decarboxylation rather

than by an enzyme-catalyzed reaction [46-48]. Therefore, animals that were either infected

with ∆*mvfR* and received 2-AA (∆*mvfR*+2-AA) at the time of infection or uninfected mice

injected with a single dose of 2-AA served as direct controls (Fig. 5). Naïve and sham mice

groups served as additional basal controls (Fig. 5).

Infection with PA14 or uninfected mice injected with 2-AA led to a significant decrease

in both ATP (Fig. 5A) and acetyl-CoA (Fig. 5B) concentrations in spleen tissues compared to

Δ*mvfR* infected mice that sustained higher ATP and acetyl-CoA levels similar to naïve and

sham control groups across the time points tested. However, in mice infected with Δ*mvfR*

and 2-AA injected *(Δ*mvfR*+2-AA)* at the time of infection, ATP and acetyl-CoA concentrations

in the spleen decreased to levels comparable to those of PA14-infected animals, strongly

indicating the biological function of 2-AA in decreasing these key metabolites. These *in vivo*

findings further support the adverse action of 2-AA on host energy homeostasis and

metabolism observed in *in vitro* studies.

The metabolic alterations observed are associated with the host tolerance to *PA*

persistence (Fig. 5C). Using *Pseudomonas* isolation agar plates, we evaluated the bacterial

load at the infection site over the course of 10 days, obtaining samples at 1, 5, and 10 days.

At one day post-infection, mice infected with PA14, Δ*mvfR,* or Δ*mvfR*+2-AA exhibited no

difference in bacterial burden (Fig. 5C), verifying the ability of all strains to establish infection.

At 5- and 10 days post-infection, mice infected with PA14 or Δ*mvfR*+2-AA sustained the

bacterial burden at a significantly higher bacterial burden over time compared to those

infected with Δ*mvfR* (Fig. 5C).

To strengthen the relevance of our *in vivo* data, we performed additional *in vivo*

experiments. In this set of *in vivo* studies, mice received the first exposure to 2-AA by

injecting 2-AA only and the 2^{nd} exposure through infection with PA14 or Δ*mvfR* four days

post-2-AA injection. As shown in the supplementary Figure S5, the levels of ATP and acetyl

CoA in the spleen of infected animals and the enumeration of the bacterial counts were

similar between PA14 or Δ*mvfR* receiving the 1^{st} 2-AA exposure and agree with the "one-

shot infection" findings presented in Figure 5 with the PA14 or Δ*mvfR* +2-AA infected mice or

those receiving 2-AA only. These results are consistent with our previous findings, showing

that 2-AA impedes the clearance of PA14 [17, 18] and provide compelling evidence that the

metabolic alterations identified may favor *PA* persistence in infected tissues.

Discussion

The *PA* signaling molecule 2-AA that is abundantly produced and secreted in human tissues

[21, 22] is the first QS molecule that epigenetically reprograms immune functions, promotes

immune tolerance, and sustains *PA's* presence in host tissues [17, 18, 22, 26]. The present

study provides insights into the mechanistic actions of 2-AA on cellular metabolism that

contribute to host immune tolerance to *PA* persistence. We uncover that this signaling

molecule causes distinct metabolic alterations in macrophages' mitochondrial respiration and

energy production promoted via the PGC-1α/ERRα axis and MPC1-dependent OXPHOS that

links the TCA cycle to the production of ATP and energy homeostasis (Fig. 6). Our results

show that 2-AA tolerization decreases ATP the crucial energy metabolite and the histone

acetylation metabolite acetyl-CoA and *in vivo*, implicating the importance of the key energy-

producing mitochondrial process in immune tolerance. Although macrophages respond to the

first exposure to 2-AA by enhancing ATP and acetyl-CoA production, long-term exposure to

2-AA leads to a tolerized state characterized by sustained reduced ATP and acetyl-CoA

concentrations, a quiescent bioenergetic state with reduced mitochondrial pyruvate uptake

and unresponsiveness to a 2^{nd} exposure.

Our results reveal that the decrease of ATP and acetyl-CoA in tolerized macrophages

results from the 2-AA-mediated perturbation of the physical interaction between ERRα and

PGC-1α. The PGC-1α/ERRα axis has not previously been implicated in tolerization but has

been extensively studied in cancer, underscoring the novelty of our findings. Studies in

cancer cells have shown that the physical interaction of human ERRα with PGC-1α strongly

enhances the binding of ERRα to DNA to induce the activation of the targeted genes [33],

including *MPC1,* responsible for the import of cytosolic pyruvate into mitochondria and PDH,

that catalyzes the conversion of pyruvate to acetyl-CoA in mitochondria. Inhibition of ERRα

has also been shown in cancer cell metabolism studies to interfere with pyruvate entry in

mitochondria by inhibiting the expression of MPC1 [49], and inhibition of ERRα led to

decreased expression of PGC-1α-controlled expression of mitochondrial genes in mice

brains [45]. Indeed, we find that the transcription of *ERRα, PGC-1α,* and *MPC1* genes and

pyruvate uptake into mitochondria is reduced in tolerized macrophages. Given that pyruvate

is a primary carbon source in the TCA cycle that fuels OXPHOS and the production of ATP

into mitochondria, it explains the reduced ATP and acetyl-CoA levels observed in tolerized

macrophages. Future studies will focus on unveiling the pathways by which macrophages

respond to the bioenergetic changes identified and decipher their effect on the pyruvate cycle

in host tolerance to infection [50-52].

We show that the unprecedented action of a QS bacterial small signaling molecule on

the interaction between ERRα and PGC-1α in the nucleus is hampered. Although it remains

to be elucidated whether 2-AA interferes directly or indirectly with the PGC-1α/ERRα axis,

our data bring up the possibility that the reduced production of acetyl-CoA may be

responsible since cholesterol is a known coactivator/ligand of ERRα [53], is generated from

acetyl-CoA [53]. This possibility may also explain the reduced expression of *ERRα* and lower

abundance since cholesterol is also required for its activation and autoregulation. However, it

is also possible that 2-AA may antagonize the interaction between ERRα and PGC-1α by

binding to ERRα. Small molecules as antagonists of ERRα have been reported previously

[54]. The weaker binding of ERRα to the *MPC1* promoter site, and consequently reduced

MPC1 expression, explains the reduced pyruvate presence in mitochondria and that of ATP

and mitochondrial quiescence. These findings open avenues for exploring novel therapeutic

strategies through the overexpression of ERRα to counteract the tolerization induced by 2-

AA and enhance clearance of *PA* infection. Interestingly, ERRα overexpression on breast

cancer metastases promotes an efficient antitumor immune response selectively in the bone
[55].

The observed cellular metabolic perturbances also align with our previous studies in
skeletal muscle, which pointed to mitochondrial dysfunction triggered by 2-AA [24, 25].
Interestingly, the *PA* QS LasR-regulated homoserine lactone molecule, 3-oxo-C12-HSL, was
reported to attenuate the expression of PGC-1α and decrease the mitochondrial respiratory
capacity in lung epithelial cells [27]. Moreover, another *PA* QS molecule, PQS, also regulated
by LasR, was recently shown to induce organelle stress, including mitochondria, by
disrupting the mitochondrial membrane potential in human macrophages [56]. As opposed to
2-AA, which dampens the pro-inflammatory response, PQS increases proinflammatory
cytokines [56], consistent with the fact that PQS is an acute infection type molecule unrelated
to chronic/persistent infections.

Previously, we reported that 2-AA tolerization induces histone deacetylation via
HDAC1, reducing H3K18ac at the TNF-α promoter [18]. The findings with acetyl-CoA
reduction, the primary substrate of histone acetylation, and the TNF-α transcription using
UK5099 and ATP in 2-AA treated macrophages are in support of the bioenergetics
disturbances observed in macrophages and their link to epigenetic modifications we have
shown to be promoted by 2-AA [18]. Macrophages exposed to 2-AA in the presence of
exogenous ATP showed improved intracellular bacterial clearance and enhanced TNF-α
levels, supporting the epigenetic interconnection of macrophages and cellular ATP
responsiveness levels against infection. The exogenous addition of UK5099 that reverted the
efficiency of the *mvfR* infected macrophages to clear the *PA* intracellular burden and the
counteraction to the 2-AA effect by ATP addition that increases the clearance of *PA*
intracellular burden support the importance of this energy-carrying molecule in *PA*
persistence.

The results with the MPC1 inhibitor, UK5099, suggest that the availability of pyruvate could

underlie the mechanism of 2-AA-regulated HDAC/HAT-dependent control of transcription of

pro-inflammatory mediators and bacterial clearance [57]. These results align with our

previous findings, showing that establishing bacterial intracellular burden in macrophages is

HDAC1 dependent [57]. It remains to be elucidated if HDAC/HAT-mediated histone

modification directly regulates the expression of OXPHOS genes in 2-AA tolerized

macrophages, as it was shown that histone deacetylation downregulates OXPHOS in

persistent *M. tuberculosis* infection [58, 59]. Given the complexity of 2-AA-mediated long-

term effects, future omics studies combined with immune profiling may aid in deciphering the

possible network of co-factors across different subpopulations of immune cells and the

immunometabolic reprogramming related to 2-AA.

We have shown that although 2-AA tolerization leads to a remarkable increase in the

survival rate of infected mice, it permits an HDAC1-dependent sustained presence of *PA* in

mice tissues and intracellularly in macrophages [17, 18, 22, 26]. Here, we use 2-AA-

producing and isogenic non-producing *PA* strains to confirm the 2-AA-mediated decrease in

the central metabolite acetyl-CoA and the energy-carrying molecule ATP during infection.

These murine infection studies provide strong evidence of the 2-AA biological relevance in

reducing these metabolites *in vivo*. Taking together our previous and current *in vivo* findings

provide compelling evidence that the metabolic alterations identified favor *PA* persistence in

infected tissues.

That 2-AA permits *PA* to persist in infected tissues despite rescuing the survival of

infected mice underscores its difference from that of LPS tolerization, which results in

bacterial clearance [23] *via* the mechanism that relies on a set of different HDAC enzymes.

LPS tolerization predominantly involves changes in H3K27 acetylation [60], while 2-AA

tolerization involves H3K18 modifications [22]. The 2-AA-mediated effects reported here are

also distinct from the epigenetic-metabolic reprogramming mediated by LPS [61], which

promotes endotoxin tolerance in immune cells by upregulating glycolysis and suppressing

OXPHOS [62]. Although 2-AA and LPS implicate different components and lead to different

outcomes, both involve epigenetic mechanisms and immune memory.

Metabolic reprogramming upon infection may be pathogen-specific, with each

pathogen impacting specific metabolic pathways that better fit its respective metabolic needs

[7]. This was shown for infections of various tissues with *Chlamydia pneumonia*, *Legionella*

pneumophila, *Mycobacterium tuberculosis*, and *Salmonella* [63-66] and, by our group, for *PA*

infections [17, 18, 24]. It would be interesting, however, to test whether 2-AA affects the

killing efficacy of macrophages against other pathogens, as emerging evidence shows that

synergistic or antagonistic interactions between clinically relevant microorganisms and host

have important implications for polymicrobial infectious diseases [67].

While in this study, we focused on the role of ERRα mainly in pyruvate metabolism;

future studies are needed to reveal other ERRα/PGC-1α axis-dependent metabolic and

immune pathways are modulated by 2-AA. In this view, the cholesterol synthesis pathway

that relies on acetyl-CoA precursor, as cholesterol is a known coactivator/ligand of ERRα [53]

and fatty acid catabolism, generating acetyl-CoA, and PGC-1α regulates fatty acid oxidation

[68].

This study unveils the unprecedented actions of a QS bacterial molecule in

orchestrating cellular metabolic reprogramming in addition to the epigenetic reprogramming

reported previously and has also been shown to promote a long-lasting presence of *PA* in the

host [17, 18, 26]. That 2-AA cross-tolerized macrophages to LPS corroborates our previous

findings on the implication of epigenetic mechanism [18] rather than ligand-receptor

mediated signaling-based mechanism and raises the possibility that this QS molecule may

confer non-specific cross-protection. This is an aspect we plan to investigate in the future.

The reported immunometabolic reprogramming contributes to a better understanding of the

molecular and cellular mechanisms, biomarkers, and functional significance that may be
involved in immune tolerance to persistent infection, providing for designing and developing
innovative therapeutics and interventions. These approaches can focus on promoting host
resilience against bacterial burden and safeguarding patients from recalcitrant persistent
infections.

Materials & Methods

Ethics statement

The animal protocol was approved by the Institutional Animal-Care and Use Committee
(IACUC) of Massachusetts General Hospital (protocol no. 2006N000093). No randomization
or exclusion of data points was applied.

Conditions used for the exposures of the cells to 2-AA

Mouse BMDM cells were isolated from the femur of CD1 6 weeks old male and female mice
and used to estimate the levels of various metabolites (Fig. 2A) following exposure and re-
exposure to 2-AA (Fig. 1A). The 1^{st} exposure of naïve BMDMs (10^6/mL in 6-well plates) was
achieved with 400 µM 2-AA for 1, 6, or 48 h. Cells exposed for 48 h were used for the 2^{nd} 2-
AA exposure. These cells were washed to remove residual 2-AA, left in the medium in the
absence of 2-AA for either 24 or 106 h, and re-exposed to 2-AA (200 µM) for 1 or 6 h,
respectively. For RAW 264.7 (10^6/mL in 6-well plates) and THP-1 cells (10^6/mL in 6-well
plates), the 2-AA concentration and conditions used were the same as those used for BMDM
cells. The hours of the 1^{st} round and 2^{nd} round of exposure to 2-AA are indicated in Figure
S1.

For the LPS stimulation studies, cells were stimulated with 400 µM 2-AA or 100 ng/mL LPS
for 6 h and 800 µM of 2-AA for 48 h. Cells receiving 1^{st} exposure for 48 h were washed and
re-exposed to 400 µM of 2-AA or 100 ng/mL LPS for 6 h.

ATP and acetyl-CoA quantifications

The levels of ATP acetyl-CoA, and were assessed in BMDM and RAW 264.7 macrophage cells following exposure to 2-AA at the times indicated and by utilizing the ATP Assay kit (cholorimetric/fluorometric) (#ab83355, Abcam) and the PicoProbe AcCoA assay kit (ab87546, Abcam), respectively, according to the manufacturer's instruction. Quantifications of ATP and acetyl-CoA were performed in triplicates.

ATP: For ATP determination, macrophages were lysed and subsequently centrifuged at 14,000 rpm for 10 min at 4 °C. The supernatants were transferred to a fresh Eppendorf tube. Standards and cell supernatants of 50 µL were added to a 96-well plate suitable for fluorescent analysis (black sides, clear bottom). A reaction mixture containing an ATP converter, probe, buffer, and developer mix was then added to all wells (50 µL) and incubated away from light at room temperature for 30 min. ATP quantification was conducted fluorometrically at 535/587 nm using a plate reader and the following settings: λex = 535 nm; λem = 587 nm. Fluorescence was measured using a microplate reader (Tecan Group Ltd, Männedorf, Switzerland)

Acetyl-CoA: Acetyl-CoA content was assessed by first deproteinizing total cell fractions of macrophages using the perchloric acid and then centrifuging at 14,000 rpm for 10 min at 4 °C. 50 µL of cells' supernatant sample CoASH were quenched to correct the background generated by free CoASH. Following the homogenization procedure described above, the samples were diluted with the reaction mix, and fluorescence was quantified using a plate reader and the settings as above with ATP.

Seahorse Assays

Seahorse analysis was performed according to previously published protocols [69]. Briefly, freshly prepared BMDM cells were reseeded in complete RPMI-1640 cells using Seahorse plates (Agilent cat. no. 103729100) at a density of 5×10^4 cells per well. Cells were exposed

to 400 µM 2-AA for 1 h or 24 hours (1st exposure). Cells receiving 1st exposure for 24 h were

washed and re-exposed to 200 µM of 2-AA for 1 h. Naïve cells were used as control. Prior to

initiating Seahorse measurements, cells were washed, and Seahorse XF DMEM

supplemented with 2 mM glutamine (Gibco cat. No. 25030-081) was added to each well.

Cells were allowed to stabilize in a 37°C incubator without CO_2 for 1 h. The Seahorse

cartridge was hydrated and calibrated as per the manufacturer's instructions. The

Mitochondrial stress test kit (Agilent cat. no. 10395-100) was used (oligomycin at 1 µM,

FCCP (1.5 µM), rotenone (0.5 µM), and antimycin A at (0.5 µM)) with slight modifications

according to the published protocol [69] that included injection of 25 mM glucose, and

sodium pyruvate (1 µM). All samples N= 4 were run in a Seahorse XFe96 Analyzer, and data

were analyzed using Wave and plotted using GraphPad Prism software.

Isolation of cytosolic and mitochondrial fraction and pyruvate quantification

Pyruvate is produced in the cytosol and is transported into the mitochondria. RAW 264.7

cells were plated at 1 ×□10^6 to incubate overnight at 37°C in a CO_2 incubator. Cells were

exposed to 2-AA for 1, 3, 24, or 48 h or re-exposed for 1 h as described above. Mitochondria

and cytosolic fractions of each group were isolated utilizing the Mitochondria Isolation Kit for

cultured cells (#ab110170, Abcam) following the manufacturer's protocol. Briefly, cells were

collected with a cell lifter and pelleted by centrifugation at 1,000 g, frozen, and then thawed

to weaken the cell membranes. The cells were resuspended in Reagent A and transferred

into a pre-cooled Dounce Homogenizer. The homogenates were centrifuged at 1,000 g for 10

min at 4°C and saved as supernatant #1 for the cytosolic fractions. The pellet was

resuspended in Reagent B, followed by repeat rupturing and centrifugation. The pellet was

collected and resuspended in 500 µL of Reagent C supplemented with Protease Inhibitor

cocktails (P8340, Sigma Aldrich). Following separation, a Bradford assay was conducted to

determine the protein concentration in each fraction. Pyruvate levels were determined in

cellular fractions and mitochondrial fractions by using the Pyruvate Assay Kit (#ab65342,

Abcam) according to the manufacturer's instructions. Briefly, after deproteinization using perchloric acid, the samples were neutralized in ice-cold 2 M KOH. 10 µL samples were incubated with reaction mix and kept on the plate at RT for 30 min in the dark. The absorbance was measured in a microplate reader (Tecan Group Ltd, Männedorf, Switzerland) at 570 nm, and the results were shown in three independent experiments.

Bacterial strains and growth conditions

The *PA* strain known as Rif[R] human clinical isolate UCBPP-PA14 (also known as PA14) was used [70]. The bacteria were grown at 37 °C in lysogeny broth (LB) under shaking and aeration or on LB agar plates containing appropriate antibiotics. PA14 and isogenic mutant $\Delta mvfR$ [19] cultures were grown in LB from a single colony to an optical density of 600 nm (OD_{600}) of 1.5, diluted 1:50,000,000 in fresh LB media, and grown overnight to an OD_{600} of 3.0.

Gentamicin protection assay

RAW 264.7 macrophage cells were plated on 6 well-cell culture-treated plates overnight in pyruvate-free DMEM. After 3 h of incubation with 10 µM UK5099, 20 µM ATP or 400 µM 2-AA, cells were infected with PA14 and isogenic mutant $\Delta mvfR$ at 5 MOI for 30 min at 37 °C in 5% CO_2. Unbound bacteria were removed by washing once with cold DMEM medium. Afterward, the cells were incubated with 100 µg/ml of gentamicin for 30 min to eliminate residual extracellular bacteria. The cells were washed with DMEM, transferred to free medium without gentamicin and kept for 3 h at 37°C in 5% CO_2. The infected cells were scraped after 3 h, centrifuged at 500 × g, and lysed in distilled water. The lysed cells were immediately diluted in PBS and plated on LB agar plates to assess bacterial presence. Bacterial colony-forming units (CFUs) were counted after incubating the plates overnight at 37°C.Untreated cells infected with PA14 were set as 100 % and the reduction of the bacterial load was expressed as %CFU.

Pharmacological inhibitors and ATP supplementations

For the OXPHOS inhibition assay, RAW 264.7 macrophage cells were treated with UK5099

(10 µM, Sigma-Aldrich, dissolved in DMSO 3 h prior to 1^{st} or 2^{nd} rounds of 2-AA exposure.

RAW 264.7 macrophages supplemented with ATP (20 µM, Sigma-Aldrich, dissolved in PBS)

received ATP at the time of 1^{st} and 2^{nd} 2-AA exposure.

Co-Immunoprecipitation assay

For all immunoprecipitation assays, protein A/G agarose Magnetic beads (Pierce) were used

after washing in 1X IP buffer. Nuclear lysates from RAW 264.7 tolerized cells exposed to 2-

AA for 24 h or non-exposed cells were diluted in 1X IP buffer, and 500 µg of protein was

taken for each experiment. The lysates were precleared by using unbound 50 µl protein A/G

magnetic beads for 2 h at room temperature on a rotator. Precleared lysates were either

incubated with ERRα, PGC-1α or rabbit IgG antibody overnight at 4 °C. 100 µl protein A/G

magnetic beads were used to pull down the antibody-protein complex and washed twice with

IP buffer to remove unbound proteins. Magnetic beads were then eluted in 2X Laemmli SDS-

PAGE loading buffer. Immunoprecipitated ERRα and PGC-1α was detected by Western blot

analyses, using conformational specific Anti Rabbit antibody (TruBlot)**.**

Immunoblotting analysis

Cells were plated at $6×10^5$ cells per well in six-well plates. Cells were washed with PBS and

subsequently lysed using RIPA lysis buffer containing 1 mM phenylmethylsulfonyl fluoride

(PMSF). 20µg of proteins were separated by electrophoresis on any KD (Kilo Dalton) (Bio-

Rad, Cat no. 4569033) SDS-polyacrylamide gel. Proteins were transferred to a 0.2 µm

polyvinylidene fluoride membrane (PVDF, Millipore, Billerica, USA) using a Bio-Rad semi-dry

instrument. After blocking with 5% BSA in TBS containing 0.1% Tween-20 for 1 h at room

temperature, the membranes were incubated with a primary antibody ERR-a (Abcam,

#ab76228), MPC1 (D2L91, Cell signaling), and anti-β-actin (cat no: sc-47778) (Santa Cruz

Biotechnology) overnight at 4°C. Following washing, the membranes were incubated with an

anti-rabbit secondary antibody, and the bands were detected by SuperSignal West Pico

Chemiluminescent Substrate (Thermo Scientific) reaction, according to the manufacturer's

instructions. The blots were visualized in the ChemiDOC Imaging system (Bio-Rad

Laboratories, Inc., Hercules, CA, USA). The bands were analyzed densitometrically using

QuantityOne software (Bio-Rad).

Chromatin immunoprecipitation (ChIP) and ChIP-qPCR

For ChIP studies, RAW 264.7 macrophage cells exposed to 2-AA for 24 h were cross-linked

in 1% (v/v) methanol-free formaldehyde for 10 min and then placed in 0.125 M glycine for 5

min at RT. Using the truChIP High Cell Chromatin Shearing kit (Covaris, United States), cells

were prepared for sonication according to the Covaris protocol. Approximately 1×10^7 cells

were plated in a 12 mm × 12 mm tube and subjected to shearing with the Covaris S220

sonicator for 8 min (140 peak power, 5 duty factor, 200 cycles/burst). The Magna ChIP A/G

kit (Millipore, United States) was used for the subsequent immunoprecipitations according to

the manufacturer's protocol. Briefly, chromatin from approximately 10^6 cells was incubated

overnight at 4°C with 2 µg of anti-ERRα or anti-PGC-1α (Abcam, USA) ChIP-grade antibody

and 20 µL of A/G magnetic beads. The beads were washed serially (5 min each) with low-

salt wash buffer, high-salt wash buffer, LiCl wash buffer, and TE buffer from the kit at 4°C.

Chromatin was eluted with elution buffer containing Proteinase K at 62°C for 4 h, then

incubated at 95°C for 10 min. DNA was isolated by column purification (QIAquick PCR

purification kit).

Real-time ChIP-qPCR was performed with the Brilliant II SYBR green super mix (Agilent,

USA). Forward (AGTGGTGACCTTGAACTTCCC) and reverse

(CTGAAGACGACCTTCCCCTT) primers were chosen to amplify a genomic locus of *MPC1*

promoter, which had a putative ERR binding site (TNAAGGTCA) at 1582 bp upstream of the

start site. Normalized values were calculated using the percent-input method relative to the

IgG. The assay was performed three times.

RNA extraction and RT-qPCR

Total RNA from all the groups mentioned above was isolated from approximately 2×10^6

cells with the RNeasy minikit (Qiagen, USA), and cDNA was prepared with the iScript

Reverse transcription kit (Bio-Rad, USA), as per the manufacturer's instruction. Real-time

PCR was conducted using the PowerUP SYBR Green Master mix (Applied Biosystem, USA)

and primer sets for mouse ERR-α (forward: ACTACGGTGTGGCATCCTGTGA; reverse:

GGTGATCTCACACTCATTGGAGG), *PGC-1α* (forward:

GAATCAAGCCACTACAGACACCG; reverse: CATCCCTCTTGAGCCTTTCGTG), MPC-1

(forward: CTCCAGAGATTATCAGTGGGCG; reverse: GAGCTACTTCGTTTGTTACATGGC),

TNF-α (forward: GGTGCCTATGTCTCAGCCTCTT; reverse:

GCCATAGAACTGATGAGAGGGAG) and mouse 18S rRNA (forward:

GTTCCGACCATAAACGATGCC; reverse: TGGTGGTGCCCTTCCGTCAAT). The transcript

levels of all the genes were normalized to 18S rRNA with the ΔΔCT method. The relative

expression was calculated by normalizing transcript levels to those of PA14-infected cells.

The assay was conducted in triplicate; means and standard deviations were calculated for

each group.

Animal infection and metabolites assessment experiments

The full-thickness thermal burn injury and infection (BI) model [17, 18, 71] was used to

assess the effect of 2-AA on metabolic alterations in the spleens of six-week-old CD1 male

mice and bacterial burden (Charles River Labs, USA). A full-thickness thermal burn injury

involving 5-8% of the total body surface area was produced on the shaved mouse abdomen

dermis, and an inoculum of ~1×10^4 PA14 and Δ*mvfR* [19] cells in 100 µL of MgSO$_4$ (10 mM)

was injected intradermally into the burn eschar. For the groups that received 2-AA, mice

were injected intraperitoneally (IP) with 100 µL of 2-AA (6.75 mg/kg). The entire procedure

was done under the influence of anesthesia. One of the groups of mice infected with the

PA14 isogenic mutant $\Delta mvfR$ [19] also received 100 μL of 2-AA in PBS (6.75 mg/kg) at the

time of infection ($\Delta mvfR$ +2-AA) and served as an additional control. CFU counts from rectus

abdominus muscle (underlying the burn-infected tissue) were assessed in groups of 4 mice

each at 1, 5, and 10 d post-BI by plating diluted muscle homogenate on Pseudomonas

isolation agar (Sigma-Aldrich) plates containing rifampicin (50 mg/L). Spleen samples from

mice were collected for metabolite analyses from all mice groups at 1, 5, and 10 d post-BI to

assess ATP (#ab83355 Abcam) and acetyl-CoA (#ab87546, Abcam) as described above.

Tissues were homogenized, 100 mg of homogenate was centrifuged at 4°C for 15 min at

10,000 g, and supernatants were collected. The supernatant was mixed with 400 μL of 1 M

perchloric acid. The deproteinized supernatant was neutralized by 3 M $KHCO_3$. The ATP and

acetyl-CoA assays were performed as described in the *in vitro* section.

Electron microscopy studies

RAW 264.7 macrophages exposed to 2-AA and corresponding controls were fixed with 2%

glutaraldehyde in 0.1 M cacodylate buffer and postfixed in 1% OsO4 in 0.1 M cacodylate

buffer for 1 h on ice. The cells were stained all at once with 2% uranyl acetate for 1 h on ice,

after which they were dehydrated in a graded series of ethanol (50–100%) while remaining

on ice. Ultrathin (70 nm) sections were cut using a Leica EMUC7 ultramicrotome and

collected onto formvar-coated grids (EMS, Hatfield, PA). Sections were contrast-stained

using 2.0% aqueous uranyl acetate. Grids were examined at 80 kV in a JEOL 1011

transmission electron microscope (Peabody, MA) equipped with an AMT digital camera and

proprietary image capture software (Advanced Microscopy Techniques, Danvers, MA).

Acknowledgments and Funding sources

We thank Dr. Vamsi Mootha and Sneha Rath for access and help with using the Seahorse

instrument, respectively. We also thank Dr. Diane Capen and Dr. Dennis Brown for the

guidance and processing of the ultramicroscopy images. This work was supported by the

NIH award R01AI134857, The John Lawrence Massachusetts General Hospital Research

Scholar Award and Shriner's grant 83009 to L.G.R., and the Shriner's grant 85132 to A.A.T.

The funders had no role in the study design, data collection, analysis, decision to publish, or

manuscript preparation.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

 Figure Legends

Figure 1. 2-AA tolerization decreases crucial metabolites of cellular energy and affects

mitochondrial respiration in mouse BMDM. (A) Schematic representation showing

experimental design: naïve cells were exposed to 2-AA for 1, 6, or 48 h (black). Cells

exposed to 2-AA (400 µM) for 48 h were washed, rested for 24 or 106 h, and re-exposed

(200 µM) for 1 or 6 h (red), respectively. The same color code, black cells after 1^{st} exposure

and red cells after 2nd exposure, was kept throughout the manuscript with corresponding

controls in grey and pink, respectively. The levels of (B) ATP and (C) acetyl-CoA in BMDM

cells after 1^{st} and 2^{nd} 2-AA exposure. (D) Real-time oxygen consumption rate (OCR) traces

were recorded using a Seahorse XF analyzer and normalized to protein content. Cells were

exposed to 2-AA for 1 or 24 h (black), washed and rested for 24 h and re-exposed for 1 h

(red). Mitochondrial respiratory parameters, (E) basal respiration, and (F) maximal

respiration. Data are presented as mean ± SD, n ≥ 4, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and

ns indicates no significant difference. One-way ANOVA followed by Tukey's post hoc test was

applied.

Figure 2. 2-AA perturbs the mitochondrial MPC1-mediated import and metabolism of

pyruvate. (A) Cytosolic and mitochondrial pyruvate levels following 2-AA exposure (black) or

re-exposure (red) and corresponding controls in grey and pink, respectively, for indicated

time points. (B) Representative Western blot and results of densitometric analysis of MPC1

protein levels following 2-AA exposure or re-exposure for indicated time points. β-actin was

used as a control. Corresponding controls are shown in grey or pink, respectively. Mean ±

SD is shown, $n = 3$, $***p < 0.001$, $****p < 0.0001$, and ns indicates no significant difference.

One-way ANOVA followed by Tukey's post hoc test was applied.

Figure 3: 2-AA-mediated macrophage tolerization deranges PGC-1α/ERRα-dependent

metabolic programing. (A) Representative Western blot and results of densitometric

analysis of ERRα protein levels following 2-AA exposure or re-exposure for indicated time

points. β-actin was used as a control. (B) Western blots and its corresponding densitometric

analysis of ERRα in cytoplasmic or nuclear lysates of tolerized macrophages exposed to 2-

AA for 24 h or not exposed to 2-AA. (C) ChIP-qPCR assay of ERRα binding at the MPC1

promoter in RAW 264.7 tolerized macrophages exposed to 2-AA (200 µM) for 24 h (black)

compared to untreated control macrophages (grey). IgG served as a negative control (D)

Representative Western blot of co-immunoprecipitation (co-IP) studies of ERRα and PGC-1α

in nuclear extracts of 2-AA tolerized (24 h) and control RAW 264.7 cells. Pull down with IgG

served as a negative control. 2-AA tolerized macrophages are shown in black, and untreated

control macrophages in grey. Mean ± SD is shown, n ≥ 3, *$p < 0.05$, **$p < 0.01$, ****$p <$

0.0001, and ns indicates no significant difference. One-way ANOVA followed by Tukey's post

hoc test was applied.

Figure 4: Increased intracellular burden in macrophages is associated with decreased

expression of *MPC1, ERR-α,* and *TNF-α genes*. Real-time PCR analysis of *MPC1* (A),

ERR-α (B), and *PGC-1α* (C) expression in RAW 246.7 macrophages infected with PA14 or

ΔmvfR in the presence or absence of exogenous addition of 2-AA or ATP for 6 h as indicated.

Transcript levels were normalized to 18srRNA. PA14-infected cells served as controls. (D)

The intracellular burden of PA14 or *ΔmvfR* of infected macrophages in the presence or

absence of exogenous addition of 2-AA, UK5099, or ATP. Untreated cells infected with PA14

were set as 100%. (E) Real-time PCR analysis of TNF-α expression in RAW 246.7

macrophages infected with PA14 or *ΔmvfR* in the presence or absence of exogenous

addition of 2-AA, ATP, or UK5099. Transcript levels were normalized to 18srRNA. PA14-

infected cells served as controls. The compound concentration used for UK5099 was 10 µM

and ATP 20 µM. Mean ± SD is shown, *n* = 3, * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$

0.0001***, and ns indicates no significant difference. One-way ANOVA followed by Tukey's

post hoc test was applied.

Figure 5. 2-AA promotes a long-lasting decrease in ATP, acetyl-CoA levels, and

bacterial persistence in *PA*-infected mice. (A) ATP and (B) acetyl-CoA concentrations in

the spleens of mice infected with *PA* wild type (PA14), the isogenic mutant Δ*mvfR*, Δ*mvfR*

injected with 2-AA at the time of infection (Δ*mvfR* + 2-AA), or non-infected but injected with 2-

AA (6.75 mg/kg), (C) Bacterial burden in muscles expressed as CFU count was analyzed

using the Kruskal–Wallis non-parametric test with Dunn's post-test; ***$p < 0.001$, and ns

indicates no significant difference. Control mice groups: naïve were not given 2-AA; mice

receiving 2-AA were given a single intraperitoneal injection of 2-AA; sham represents a

burn/PBS group since the burn and infection model was used. Results of three independent

replicates with 4 mice per group for 1, 5, and 10 d are shown. Means ± SD are shown, *$p <$

0.05, **$p < 0.01$, ***$p < 0.001$, and ns indicate no significant difference. One-way ANOVA

followed by Tukey's post hoc test was applied.

Figure 6: Proposed mechanism by which 2-AA impairs bioenergetics through the

inhibition of MPC1-mediated pyruvate transport into mitochondria and its impact on

the PGC-1α/ERRα axis. 2-AA tolerized macrophages exhibit diminished pyruvate levels in

mitochondria due to the decreased expression of *MPC1*, a consequence of the 2-AA impact

on the interaction of ERRα with the transcriptional coactivator PGC-1α for the effective

transcription of ERRα since ERRα controls its own transcription and that of MPC1. In the

presence of 2-AA, the weakened interaction between ERRα and PGC-1α results in reduced

expression of MPC1 and ERRα. The reduction in mitochondrial pyruvate levels leads to

decreased acetyl-CoA and ATP levels, which modulate HDAC1- and HAT-catalyzed

remodeling of H3K18 acetylation. The diminished levels of this epigenetic mark have

previously been associated with an increased intracellular presence of bacteria in

macrophages, as demonstrated by our group [18]. Pathways, proteins, and metabolites that

are negatively affected are indicated in red, while positively affected are denoted in black.

The figure was created using BioRender.com.

Figure S1. 2-AA tolerization decreases metabolites in murine RAW 264.7 (A-B) and human

THP-1 (C-D) cells. (A and C) ATP, (B and D) acetyl-CoA, and (E) cytosolic and (F)

mitochondrial pyruvate levels were quantified in 2-AA (400 μM) exposed and re-exposed

cells for the hours indicated using the same condition as shown in Fig. 1A. Mean ± SD is

shown (n = 3); *** p < 0.001, and ns indicates no significant difference. One-way ANOVA

followed by Tukey's post hoc test was applied. The same color code, black cells after 1st

exposure and red cells after 2nd exposure, was kept throughout the main manuscript, with

corresponding controls shown in grey and pink, respectively.

Figure S2: ATP levels in murine BMDM (A) cells with and without 2-AA (400 μM) or LPS

stimulation (100 μg/mL). Each dot represents 1 experimental replicate of 4 independent

experiments (n = 4). Data are presented as mean ± SD, ** p < 0.01, *** p < 0.001, and ns

indicates no significant difference. One-way ANOVA followed by Tukey's post hoc test was

applied.

Figure S3: Effect of 2-AA on mitochondrial spare respiratory capacity in BMDM non-tolerized

(black) and tolerized (red) macrophages. Controls are shown in grey and pink, respectively.

Mitochondrial spare respiratory capacity data extrapolated from the OCR profiles shown in

Figure 1D. Unstimulated BMDM macrophages were used as a control (c). Means ± SDs are

shown, n = 6, * p < 0.05, *** p < 0.001, and ns indicates no significant difference. One-way

ANOVA followed by Tukey's post hoc test was applied.

Figure S4: Transmission electron microscopy (TEM) images showing structural alterations of

mitochondria in 2-AA exposed macrophages for 48 h. High-magnification (30,000X) TEM

images showing mitochondrial abundance and structural changes in 2-AA exposed

macrophages as compared to the non-exposed macrophages.

Figure S5: Exposure and re-exposure to 2-AA promotes a long-lasting decrease in ATP and

acetyl-CoA levels and sustains bacterial presence in mice receiving 1st exposure to 2-AA by

injecting 2-AA and 2nd exposure through infection with PA14 or Δ*mvfR* four days post-2-AA

injection. (A) ATP and (B) acetyl-CoA concentrations in the spleens of mice. (C) Bacterial

burden in muscles expressed as CFU counts was analyzed using the Kruskal–Wallis non-

parametric test with Dunn's post-test; ***$p < 0.001$, and ns indicates no significant difference.

Control mice groups: naïve were not given 2-AA; mice receiving 2-AA were given a single

intraperitoneal injection of 2-AA four days prior to infection; sham represents a burn/PBS

group since the burn and infection model was used. N = 4 mice per group, and 10 days post-

infection is shown. Means ± SD are shown, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ns

indicate no significant difference. One-way ANOVA followed by Tukey's post hoc test was

applied.

References

1.　Foster, S.L., D.C. Hargreaves, and R. Medzhitov, *Gene-specific control of inflammation by tlr-*
induced chromatin modifications. Nature, 2007. **447**(7147): p. 972.
2.　Bagchi, A., et al., *Myd88-dependent and myd88-independent pathways in synergy, priming,*
and tolerance between tlr agonists. Journal of immunology, 2007. **178**(2): p. 1164.
3.　Masyutina, A.M., et al., *Inhibition of specific signaling pathways rather than epigenetic*
silencing of effector genes is the leading mechanism of innate tolerance. Frontiers in
Immunology, 2023. **14**.
4.　O'Neill, L.A., R.J. Kishton, and J. Rathmell, *A guide to immunometabolism for immunologists.*
Nat Rev Immunol, 2016. **16**(9): p. 553.
5.　Chi, H., *Immunometabolism at the intersection of metabolic signaling, cell fate, and systems*
immunology. Cell Mol Immunol, 2022. **19**(3): p. 299.
6.　Tannahill, G.M., et al., *Succinate is an inflammatory signal that induces il-1beta through hif-*
1alpha. Nature, 2013. **496**(7444): p. 238.
7.　Galli, G. and M. Saleh, *Immunometabolism of macrophages in bacterial infections.* Front Cell
Infect Microbiol, 2020. **10**: p. 607650.
8.　Wang, Y., et al., *Mitochondrial metabolism regulates macrophage biology.* J Biol Chem, 2021.
297(1): p. 100904.
9.　Karan, K.R., et al., *Leukocyte cytokine responses in adult patients with mitochondrial DNA*
defects. Journal of Molecular Medicine, 2022. **100**(6): p. 963.
10.　Bird, L., *Getting enough energy for immunity.* Nat Rev Immunol, 2019. **19**(5): p. 269.
11.　Deziel, E., et al., *The contribution of mvfr to pseudomonas aeruginosa pathogenesis and*
quorum sensing circuitry regulation: Multiple quorum sensing-regulated genes are modulated
without affecting lasri, rhlri or the production of n-acyl-l-homoserine lactones. Mol Microbiol,
2005. **55**(4): p. 998.
12.　Xiao, G., et al., *Mvfr, a key pseudomonas aeruginosa pathogenicity lttr-class regulatory*
protein, has dual ligands. Mol Microbiol, 2006. **62**(6): p. 1689.
13.　Eickhoff, M.J. and B.L. Bassler, *Snapshot: Bacterial quorum sensing.* Cell, 2018. **174**(5): p.
1328.
14.　Fuqua, C. and E.P. Greenberg, *Listening in on bacteria: Acyl-homoserine lactone signalling.*
Nature Reviews Molecular Cell Biology, 2002. **3**(9): p. 685.
15.　Kariminik, A., M. Baseri-Salehi, and B. Kheirkhah, *Pseudomonas aeruginosa quorum sensing*
modulates immune responses: An updated review article. Immunol Lett, 2017. **190**: p. 1.
16.　Liu, Y.C., K.G. Chan, and C.Y. Chang, *Modulation of host biology by pseudomonas aeruginosa*
quorum sensing signal molecules: Messengers or traitors. Front Microbiol, 2015. **6**: p. 1226.
17.　Bandyopadhaya, A., et al., *The quorum sensing volatile molecule 2-amino acetophenon*
modulates host immune responses in a manner that promotes life with unwanted guests.
PLoS pathogens, 2012. **8**(11): p. e1003024.
18.　Bandyopadhaya, A., et al., *A quorum-sensing signal promotes host tolerance training through*
hdac1-mediated epigenetic reprogramming. Nat Microbiol, 2016. **1**: p. 16174.
19.　Cao, H., et al., *A quorum sensing-associated virulence gene of pseudomonas aeruginosa*
encodes a lysr-like transcription regulator with a unique self-regulatory mechanism. Proc Natl
Acad Sci U S A, 2001. **98**(25): p. 14613.
20.　Deziel, E., et al., *Analysis of pseudomonas aeruginosa 4-hydroxy-2-alkylquinolines (haqs)*
reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication. Proc Natl Acad Sci
U S A, 2004. **101**(5): p. 1339.
21.　Kesarwani, M., et al., *A quorum sensing regulated small volatile molecule reduces acute*
virulence and promotes chronic infection phenotypes. PLoS pathogens, 2011. **7**(8): p.
e1002192.

22.   Bandyopadhaya, A., A. Tsurumi, and L.G. Rahme, *Nf-kappabp50 and hdac1 interaction is*
implicated in the host tolerance to infection mediated by the bacterial quorum sensing signal
2-aminoacetophenone. Front Microbiol, 2017. **8**: p. 1211.
23.   Wheeler, D.S., et al., *Induction of endotoxin tolerance enhances bacterial clearance and*
survival in murine polymicrobial sepsis. Shock, 2008. **30**(3): p. 267.
24.   Tzika, A.A., et al., *A small volatile bacterial molecule triggers mitochondrial dysfunction in*
murine skeletal muscle. PloS one, 2013. **8**(9): p. e74528.
25.   Bandyopadhaya, A., et al., *Bacterial-excreted small volatile molecule 2-aminoacetophenone*
induces oxidative stress and apoptosis in murine skeletal muscle. International Journal of
Molecular Medicine, 2016. **37**(4): p. 867.
26.   Chakraborty, A., et al., *Quorum-sensing signaling molecule 2-aminoacetophenone mediates*
the persistence of pseudomonas aeruginosa in macrophages by interference with autophagy
through epigenetic regulation of lipid biosynthesis. mBio, 2023: p. e0015923.
27.   Maurice, N.M., et al., *Pseudomonas aeruginosa induced host epithelial cell mitochondrial*
dysfunction. Scientific Reports, 2019. **9**(1).
28.   Lelliott, C.J. and A. Vidal-Puig, *Pgc-1beta: A co-activator that sets the tone for both basal and*
stress-stimulated mitochondrial activity. Adv Exp Med Biol, 2009. **646**: p. 133.
29.   Supruniuk, E., A. Mikłosz, and A. Chabowski, *The implication of pgc-1α on fatty acid transport*
across plasma and mitochondrial membranes in the insulin sensitive tissues. Front Physiol,
2017. **8**: p. 923.
30.   Coppi, L., et al., *Pgc1s and beyond: Disentangling the complex regulation of mitochondrial*
and cellular metabolism. Int J Mol Sci, 2021. **22**(13): p. 6913.
31.   Lin, J., C. Handschin, and B.M. Spiegelman, *Metabolic control through the pgc-1 family of*
transcription coactivators. Cell Metab, 2005. **1**(6): p. 361.
32.   Huss, J.M., W.G. Garbacz, and W. Xie, *Constitutive activities of estrogen-related receptors:*
Transcriptional regulation of metabolism by the err pathways in health and disease. Biochim
Biophys Acta, 2015. **1852**(9): p. 1912.
33.   Schreiber, S.N., et al., *The transcriptional coactivator pgc-1 regulates the expression and*
activity of the orphan nuclear receptor estrogen-related receptor alpha (erralpha). J Biol
Chem, 2003. **278**(11): p. 9013.
34.   Laganiere, J., et al., *A polymorphic autoregulatory hormone response element in the human*
estrogen-related receptor alpha (erralpha) promoter dictates peroxisome proliferator-
activated receptor gamma coactivator-1alpha control of erralpha expression. J Biol Chem,
2004. **279**(18): p. 18504.
35.   Villena, J.A., *New insights into pgc-1 coactivators: Redefining their role in the regulation of*
mitochondrial function and beyond. FEBS J, 2015. **282**(4): p. 647.
36.   Ranhotra, H.S., *The estrogen-related receptor alpha: The oldest, yet an energetic orphan with*
robust biological functions. J Recept Signal Transduct Res, 2010. **30**(4): p. 193.
37.   Deblois, G. and V. Giguere, *Functional and physiological genomics of estrogen-related*
receptors (errs) in health and disease. Biochim Biophys Acta, 2011. **1812**(8): p. 1032.
38.   Ranhotra, H.S., *The estrogen-related receptors: Orphans orchestrating myriad functions.* J
Recept Signal Transduct Res, 2012. **32**(2): p. 47.
39.   Charest-Marcotte, A., et al., *The homeobox protein prox1 is a negative modulator of*
erralpha/pgc-1alpha bioenergetic functions. Genes Dev, 2010. **24**(6): p. 537.
40.   Eichner, L.J. and V. Giguere, *Estrogen related receptors (errs): A new dawn in transcriptional*
control of mitochondrial gene networks. Mitochondrion, 2011. **11**(4): p. 544.
41.   Koh, E., et al., *Mpc1 is essential for pgc-1alpha-induced mitochondrial respiration and*
biogenesis. Biochem J, 2018. **475**(10): p. 1687.
42.   Divangahi, M., et al., *Trained immunity, tolerance, priming and differentiation: Distinct*
immunological processes. Nat Immunol, 2021. **22**(1): p. 2.

43.   Koh, E., et al., *Mpc1 is essential for pgc-1α-induced mitochondrial respiration and biogenesis.*
Biochem J, 2018. **475**(10): p. 1687.

44.   Xue, C., et al., *Mitochondrial pyruvate carrier 1: A novel prognostic biomarker that predicts*
favourable patient survival in cancer. Cancer Cell Int, 2021. **21**(1): p. 288.

45.   Bronte, V. and M.J. Pittet, *The spleen in local and systemic regulation of immunity.* Immunity,
2013. **39**(5): p. 806.

46.   Starkey, M., et al., *Identification of anti-virulence compounds that disrupt quorum-sensing*
regulated acute and persistent pathogenicity. PLoS pathogens, 2014. **10**(8): p. e1004321.

47.   Dulcey, C.E., et al., *The end of an old hypothesis: The pseudomonas signaling molecules 4-*
hydroxy-2-alkylquinolines derive from fatty acids, not 3-ketofatty acids. Chem Biol, 2013.
20(12): p. 1481.

48.   Fetzner, S. and S.L. Drees, *Old molecules, new biochemistry.* Chem Biol, 2013. **20**(12): p. 1438.

49.   Park, S., et al., *Inhibition of erralpha prevents mitochondrial pyruvate uptake exposing nadph-*
generating pathways as targetable vulnerabilities in breast cancer. Cell Rep, 2019. **27**(12): p.
3587.

50.   Caslin, H.L., et al., *Lactate is a metabolic mediator that shapes immune cell fate and function.*
Front Physiol, 2021. **12**: p. 688485.

51.   Lee, T.Y., *Lactate: A multifunctional signaling molecule.* Yeungnam Univ J Med, 2021. **38**(3): p.
183.

52.   Li, X., et al., *Lactate metabolism in human health and disease.* Signal Transduct Target Ther,
2022. **7**(1): p. 305.

53.   Wei, W., et al., *Ligand activation of erralpha by cholesterol mediates statin and*
bisphosphonate effects. Cell Metab, 2016. **23**(3): p. 479.

54.   Chisamore, M.J., et al., *Identification of small molecule estrogen‐related receptor α‐*
specific antagonists and homology modeling to predict the molecular determinants as the
basis for selectivity over errβ and erry. Drug development research, 2008. **69**(4): p. 203.

55.   Bouchet, M., et al., *Errα expression in bone metastases leads to an exacerbated antitumor*
immune response. Cancer Research, 2020. **80**(13): p. 2914.

56.   Kushwaha, A., V. Kumar, and V. Agarwal, *Pseudomonas quinolone signal induces organelle*
stress and dysregulates inflammation in human macrophages. Biochim Biophys Acta Gen
Subj, 2023. **1867**(2): p. 130269.

57.   Pietrocola, F., et al., *Acetyl coenzyme a: A central metabolite and second messenger.* Cell
Metab, 2015. **21**(6): p. 805.

58.   Chandran, A., et al., *Mycobacterium tuberculosis infection induces hdac1-mediated*
suppression of il-12b gene expression in macrophages. Front Cell Infect Microbiol, 2015. **5**: p.
90.

59.   Shi, L. and B.P. Tu, *Acetyl-coa and the regulation of metabolism: Mechanisms and*
consequences. Curr Opin Cell Biol, 2015. **33**: p. 125.

60.   Lauterbach, M.A., et al., *Toll-like receptor signaling rewires macrophage metabolism and*
promotes histone acetylation via atp-citrate lyase. Immunity, 2019. **51**(6): p. 997.

61.   Saeed, S., et al., *Epigenetic programming of monocyte-to-macrophage differentiation and*
trained innate immunity. Science, 2014. **345**(6204): p. 1251086.

62.   Liu, T.F., et al., *Nad+-dependent sirtuin 1 and 6 proteins coordinate a switch from glucose to*
fatty acid oxidation during the acute inflammatory response. J Biol Chem, 2012. **287**(31): p.
25758.

63.   Ishida, K., et al., *Chlamydia pneumoniae effector chlamydial outer protein n sequesters*
fructose bisphosphate aldolase a, providing a benefit to bacterial growth. BMC Microbiol,
2014. **14**: p. 330.

64.   Kunze, M., et al., *Metabolic adaption of legionella pneumophila during intracellular growth in*
acanthamoeba castellanii. Int J Med Microbiol, 2021. **311**(4): p. 151504.

65.   Shi, L., et al., *Infection with mycobacterium tuberculosis induces the warburg effect in mouse*
lungs. Sci Rep, 2015. **5**: p. 18176.

66.   Perez-Morales, D. and V.H. Bustamante, *Cross-kingdom metabolic manipulation promotes*
salmonella replication inside macrophages. Nat Commun, 2021. **12**(1): p. 1862.

67.   Dhamgaye, S., Y. Qu, and A.Y. Peleg, *Polymicrobial infections involving clinically relevant gram-*
negative bacteria and fungi. Cell Microbiol, 2016. **18**(12): p. 1716.

68.   Vega, R.B., J.M. Huss, and D.P. Kelly, *The coactivator pgc-1 cooperates with peroxisome*
proliferator-activated receptor alpha in transcriptional control of nuclear genes encoding
mitochondrial fatty acid oxidation enzymes. Molecular and cellular biology, 2000. **20**(5): p.
1868.

69.   Van den Bossche, J., J. Baardman, and M.P. de Winther, *Metabolic characterization of*
polarized m1 and m2 bone marrow-derived macrophages using real-time extracellular flux
analysis. J Vis Exp, 2015(105): p. e53424.

70.   Rahme, L.G., et al., *Common virulence factors for bacterial pathogenicity in plants and*
animals. Science, 1995. **268**(5219): p. 1899.

71.   Maura, D., A. Bandyopadhaya, and L.G. Rahme, *Animal models for pseudomonas aeruginosa*
quorum sensing studies. Methods Mol Biol, 2018. **1673**: p. 227.