The Bacterial Quorum-Sensing Signal 2-Aminoacetophenone Rewires Immune Cell Bioenergetics through the PGC-1α/ERRα Axis to Mediate Tolerance to Infection

Arijit Chakraborty[1,2,3#], Arunava Bandyopadhaya[1,2,&#], Vijay K Singh[1,2], Filip Kovacic[1,3,4], Sujin Cha[1], William M. Oldham[5], A. Aria Tzika[1,2], and Laurence G Rahme[1,2,3*]

[1]Department of Surgery, Massachusetts General Hospital, and Harvard Medical School, Boston, Massachusetts, USA

[2]Shriners Hospitals for Children Boston, Boston, Massachusetts, USA

[3]Department of Microbiology, Harvard Medical School, Boston, Massachusetts, USA

[4]Institute of Molecular Enzyme Technology, Heinrich Heine University Düsseldorf, Forschungszentrum Jülich GmbH, Jülich, Germany

[5]Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA

[&] Present Address: Astellas Pharma Inc., USA

[#]These authors contributed equally

[*]**Corresponding author:** Laurence G. Rahme, Ph.D

e-mail: rahme@molbio.mgh.harvard.edu (LGR)

Address: 340 Their Research Building, 50 Blossom Street, Massachusetts General Hospital, Boston, MA 02114

Phone: 617-724-5003

Author Contributions:

Author contributions: L.G.R. conceptualization, A.B., A.C. and L.G.R., designed research; A.B., A.C. V.K.S., S.H.W., S. C., performed research; A.B., A.C. V.K.S., F.K., and L.G.R. analyzed data; W.M.O and A.A.T., provided resources, and F.K. and L.G.R. wrote the paper.

Competing Interest Statement:

L.G.R. has a financial interest in Spero Therapeutics, a company developing therapies to treat bacterial infections. L.G.R.'s financial interests are reviewed and managed by Massachusetts General Hospital and Partners Health Care in accordance with their conflict-of-interest policies. No funding was received from Spero Therapeutics, and it had no role in study design, data collection, analysis, interpretation, or the decision to submit the work for publication. The remaining authors declare no competing interests.

Classification:

Biological, health, and Medical Sciences/Microbiology

Keywords: Pseudomonas aeruginosa, quorum sensing, infection, macrophages, bacterial persistence, 2-aminoacetophenone (2-AA), MvfR, PqsR, immunometabolism, host tolerization, histone deacetylation, epigenetic reprogramming, metabolic reprogramming, oxidative phosphorylation, pyruvate, mitochondrial pyruvate carrier (MPC1), peroxisome proliferator-activated receptor gamma co-activator 1-alpha (PGC-1α), estrogen-related receptor (ERRα), acetyl-CoA, ATP, bioenergetics.

Abstract

How bacterial pathogens exploit host metabolism to promote immune tolerance and persist in infected hosts remains elusive. To achieve this, we show that *Pseudomonas aeruginosa (PA),* a recalcitrant pathogen, utilizes the quorum sensing (QS) signal 2-aminoacetophenone (2-AA). Here, we unveil how 2-AA-driven immune tolerization causes distinct metabolic perturbations in macrophages' mitochondrial respiration and bioenergetics. We found that the 2-AA tolerization impairs oxidative phosphorylation (OXPHOS), leading to decreased generation of the crucial energy metabolite ATP and histone acetylation acetyl-CoA. We provide evidence that these effects result from reduced pyruvate flux into mitochondria due to the decreased expression of the mitochondrial pyruvate carrier (MPC1) mediated *via* the reduced expression and nuclear presence of its transcriptional regulator estrogen-related nuclear receptor (ERRα), leading to the weaker binding of ERRα to *MPC1* promoter. This is the outcome of the hampered interaction of ERRα with the peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1α) that ultimately leads to reduced pyruvate flux into mitochondria and ATP production in tolerized macrophages. Exogenously added ATP in 2-AA exposed macrophages restores the transcript levels of *MPC1* and *ERR-α and* enhances cytokine production and intracellular bacterial clearance. Consistent with the *in vitro* findings, murine infection studies corroborate the 2-AA-mediated long-lasting decrease in ATP and acetyl-CoA and its association with PA persistence, further supporting this QS signaling molecule as the culprit of the host bioenergetic impairment and *PA* persistence. These findings unveil 2-AA as a negative modulator of cellular immunometabolism implicating the PGC-1α/ERRα axis in its influence on MPC1/OXPHOS-dependent energy production and *PA* clearance. These findings shed light on the underlying mechanisms of host tolerance and on potential therapeutic strategies to combat persistent *PA* infections.

Introduction

Host tolerance is a fundamental mechanism of innate immunity. Studies have shown that innate immune cells following infection or exposure to microbial products may enter a state of immune tolerance characterized by reduced responsiveness towards microbial re-exposure a few hours later [1-3]. Epigenetic and signaling-based mechanisms have been proposed to be involved in host immune tolerance [1-3]. In recent years, several reports have highlighted the complex interplay between metabolic reprogramming and immunity [4, 5]. It was suggested that specific metabolic programs are activated in monocytes and macrophages upon exposure to infection and microbial products [6, 7]. However, in host tolerance, the molecular mechanisms underlying immunometabolic reprogramming mediated by bacterial pathogens remain poorly understood.

Metabolic pathways are essential for generating energy for various cellular functions, including those performed by immune cells [8, 9]. Macrophages use metabolic pathways to generate energy and metabolites to adapt to changing environments and stimuli, thereby enabling them to cope with the needs of a fluctuating immune response [7, 10]. Thus, properly functioning metabolic pathways are vital for immune cells to counteract pathogens. For instance, the crucial energy-carrying molecule adenosine triphosphate (ATP) is required to phagocytose pathogen-derived molecules efficiently.

Pseudomonas aeruginosa (PA), a recalcitrant ESKAPE (*Enterococcus faecium, Staphylococcus aureus, Klebsiella pneumoniae, Acinetobacter baumannii, PA,* and *Enterobacter sp.*) pathogen, secretes virulence-associated low molecular weight signaling molecules, several of which are regulated by quorum sensing (QS) [11-14] and able to modulate host immune responses [15-18]. QS is a cell density-dependent signaling system bacteria use to synchronize their activities [11-14]. MvfR (a.k.a. PqsR), a critical QS transcription factor of *PA,* regulates the synthesis of many small molecules, including signaling molecules such as 2-aminoacetophenone (2-AA) [11, 19-21]. *In vivo* studies have demonstrated that 2-AA enables *PA* to persist in infected murine tissues by promoting innate

immune tolerance through histone deacetylase 1 (HDAC1)-mediated epigenetic

reprogramming [17, 18]. 2-AA tolerization reprograms the host inflammatory signaling cascade

by maintaining chromatin in a "silent" state through increased HDAC1 expression and activity

and decreased histone acetyltransferase (HAT) activity [18]. These changes also impact the

protein-protein interaction between HAT and cyclic AMP response element-binding

protein/HDAC1 and p50/p65 nuclear factor (NF)-κB subunits [18, 22]. The 2-AA-mediated

tolerization permits *PA* to persist in infected tissues [17, 18] underscoring the difference from

lipopolysaccharide (LPS)-mediated tolerization, which instead leads to bacterial clearance and

involves different HDACs [23].

Mitochondria, the "powerhouse" of the cells, are crucial for the regulation, differentiation, and

survival of macrophages and other immune cells [8]. Our group's previous *in vivo* and *in vitro*

studies indicated that 2-AA affects metabolic functions in skeletal muscle, which contains a

high concentration of mitochondria [24-26]. Injection of 2-AA in murine skeletal muscle

dampens the expression of genes associated with OXPHOS and the master regulator of

mitochondrial biogenesis peroxisome proliferator-activated receptor-γ coactivator-1 beta

(PGC-1β) [24]. Another *PA* QS molecule, 3-oxo-C12-HSL, attenuates the expression of PGC-

1α [27]. PGC-1α and PGC-1β belong to the PPARγ family of inducible transcriptional

coactivators and are known regulators of mitochondrial metabolism [28]. The coactivator PGC-

1 proteins are involved in various cellular energy metabolic processes [29, 30], including but

not limited to mitochondrial metabolism [31]. Recent studies have also emphasized the role of

estrogen-related nuclear receptor α (ERRα) in coordinating metabolic capacity with energy

demand in health and disease [32]. PGC-1α activates ERRα transcriptional activity through

protein-protein interaction, which enhances the expression of ERRα [33, 34] and other

mitochondrial genes involved in lipid metabolism and OXPHOS [35]. PGC-1α/ERRα axis has

been extensively studied in cancer and established as a central regulatory node of energy

metabolism that induces the global expression of genes involved in mitochondrial biogenesis

and functions [36-38]. ERRα has been shown to occupy the promoter regions of many genes

participating in the TCA cycle and OXPHOS [39, 40], including the mitochondrial pyruvate

carrier (MPC1). In human renal carcinoma cells, PGC-1α/ERRα interaction results in efficient

activation of MPC1 expression and transport of pyruvate into mitochondrion for efficient

OXPHOS and energy production [41].

Given that our previous studies pointed to the 2-AA effect on mitochondrial functions [24-26]

and that mitochondria are crucial for the regulation, differentiation, and survival of

macrophages and other immune cells [8] we investigated how 2-AA may mediate cellular

metabolic reprogramming in tolerized immune cells by examining the link between the PGC-

1α/ERRα axis and 2-AA-mediated immune tolerance. Our findings uncovered the crucial action

of 2-AA on energy homeostasis and metabolism in tolerized immune cells through

perturbances of the ERRα/PGC-1α interaction, MPC1-mediated pyruvate transport and the

production of the key energy metabolism molecules, ATP and acetyl-CoA and their association

with the persistence of *PA* in mammalian tissues.

Results

Tolerization by 2-AA impacts the generation of crucial energy metabolites in

macrophages

Following infection or exposure to microbial products, innate immune cells may enter

a state of tolerance several hours later, as depicted by their diminished responsiveness or

unresponsiveness to microbial re-exposure [1-3, 42]. Given that 2-AA tolerized cells are non-

responsive to 2-AA re-exposure, we investigated the impact of 2-AA on energy homeostasis

and metabolism in non-tolerized and tolerized BMDM cells (Fig. 1). We quantified the levels of

ATP and acetyl-CoA, which is a fuel for ATP production, in a series of *in vitro* experiments in

which murine BMDM cells were exposed to 2-AA (1st 2-AA exposure) or re-exposed (2nd 2-AA

exposure) to respectively determine their activation, and tolerance as well as memory to this

QS molecule (Fig. 1A). Initially, following the 1st 2-AA exposure (black bars), a significant increase in the concentrations of intracellular ATP (Fig. 1B) and acetyl-CoA (Fig. 1C) was observed in BMDM cells exposed to 2-AA for 1 and 6 h compared to their corresponding naïve control cells (grey bars). However, cells exposed to 2-AA for 48 h (black bar) appeared to have entered a state of tolerance as ATP and acetyl-CoA levels were decreased compared to 2-AA exposures for a shorter time (Figs. 1B and 1C). The tolerance of these cells as well as their memory to 2-AA was determined by their responsiveness to a 2nd 2-AA exposure (Fig. 1A). To achieve this, these cells were washed to remove 2-AA and allowed to rest for 24 or 106 h in the absence of 2-AA before receiving the 2nd 2-AA exposure for 1 or 6 h (Fig. 1A). Figures 1B and 1C (red bars) show the unresponsiveness of BMDM cells to 2-AA re-exposure as evidenced by the similar levels of ATP and acetyl-CoA compared to cells exposed to 2-AA for 48 h (1st exposure black bars). The sustained unresponsiveness of the cells to re-exposure indicates that the cells exhibited innate immunological memory, which could not be reverted by 2-AA re-exposure. Similar responsiveness patterns of ATP and acetyl-CoA as with BMDM cells were observed in murine macrophage RAW 264.7 cells (Figs. S1A and S1B) or human monocyte THP-1 cells (Figs. S1C and S1D) following exposure or re-exposure to 2-AA. These findings further support our observations with BMDM cells and confirm that either cell type can be used to study responses to 2-AA. These results reinforce the notion that 2-AA impacts crucial energy metabolites in macrophages.

Moreover, we sought to determine whether 2-AA-tolerized BMDM cells would respond to heterologous stimulus. To test this, we re-exposed the cells to lipopolysaccharide (LPS), an outer membrane bacterial component that strongly induces macrophages. LPS re-exposure of the 2-AA tolerized BMDM cells did not significantly ($p = 0.9$) increase ATP levels (Fig. S2), indicating that 2-AA cross-tolerized the cells to this heterologous bacterial immunostimulant. On the other hand, as expected, we observed increased ATP levels following LPS stimulation of the non-tolerized BMDM cells (Fig. S2). The broad spectrum of 2-AA cross-tolerization

strongly suggests an interplay of epigenetic mechanism rather than ligand-receptor mediated

signaling-based mechanism in bringing out the observed 2-AA-mediated tolerance.

Tolerization by 2-AA leads to a quiescent state by reducing cell bioenergetics and

compromising OXPHOS.

To further investigate the 2-AA's impact on the bioenergetics of macrophages, we

assessed by Seahorse assay the oxygen consumption rate (OCR) as an index of OXPHOS

in 2-AA exposed (for 1 and 24 h) and re-exposed (2^{nd} exposure for 1 h) BMDM cells (Fig. 1D).

Values of basal (Fig. 1E) and maximal (Fig. 1F) mitochondrial respiration and spare

respiratory capacity (Fig. S3) interpreted from these measurements revealed that 1 h 2-AA

exposure of naïve cells significantly increased OCR compared to naïve control cells. At 24 h

post-2-AA exposure, cells had a significantly reduced basal OCR level than the control BMDM

cells, indicating decreased OXPHOS and an overall quiescent phenotype of tolerized

macrophages. Re-exposure of tolerized macrophages with 2-AA also did not augment

maximal and basal OCR (Figs. 1D-1F), indicating the unresponsiveness of tolerized BMDM

cells.

These OCR data are consistent with previous direct measurements of ATP (Fig. 1B)

and acetyl-CoA (Fig. 1C) and support the notion that 2-AA tolerization induces a quiescent

phenotype in these cells characterized by defective OXPHOS. These findings confirm that 2-

AA mediates energy homeostatic and metabolic alterations in the immune cells and that 2-AA

tolerization arrests these cells in a sustained 2-AA-unresponsive state.

The pyruvate transport into mitochondria is decreased in tolerized macrophages.

Since acetyl-CoA and ATP levels were reduced due to 2-AA tolerization, we

investigated whether the mitochondrial dysfunction observed is related to pyruvate

metabolism. Pyruvate links glycolysis with mitochondrial production of acetyl-CoA and ATP *via*

the tricarboxylic acid (TCA) cycle and OXPHOS [7]. RAW 264.7 cells initially exposed to 2-AA

for 1 or 3 h (black bars) exhibited significantly higher concentrations of cytosolic and

mitochondrial pyruvate compared to the corresponding naïve control cells (Fig. 2A). Pyruvate

significantly decreased over time in cytosolic and mitochondrial fractions as cells were entering

into the tolerized state (Fig. 2A). Tolerized cells remain unresponsive to 2-AA re-exposure (red

bars), exhibiting no significant change in the concentration of pyruvate in the cytosolic or

mitochondrial fraction (Fig. 2A). A similar result was observed in THP-1 cells (Figs. S1E and

S1F).

To gain a better understanding of the 2-AA-mediated reductions in mitochondrial

pyruvate, ATP production, and acetyl-CoA levels, we sought to determine the involvement of

the MPC1 [41] that is crucial in transporting pyruvate from the cytosol into the mitochondria

[43]. Western blot studies of whole cell lysates of RAW 246.7 cells exposed to 2-AA showed a

significant decrease over time in the abundance of MPC1 protein, and tolerized cells remain

unresponsive to 2-AA re-exposure (Fig. 2B). Together these findings suggest that 2-AA

tolerization dysregulates MPC1 mediated transport of the pyruvate into mitochondria.

2-AA tolerization suppresses ERRα binding to the *MPC1* promoter and the interaction

between PGC-1α and ERRα

Given that the PGC-1α/ERRα axis is the transcriptional regulatory node that activates

the expression of MPC1 [33] we determined the protein levels of ERRα. Western blot studies

of whole cell lysates showed a significant decrease over time in the abundance of ERRα

protein following 1st exposure of RAW 246.7 cells to 2-AA, while tolerized cells remain

unresponsive to 2-AA re-exposure (Fig. 3A). By assessing the cytosolic and nuclear

abundance of ERRα in tolerized cells we found that both fractions had lower levels of ERRα

protein than their corresponding control cells (Fig. 3B).

Furthermore, because an ERRα binding putative DNA motif (TNAAGGTCA) has been

identified upstream of the *MPC1* promoter in humans [41], and we found the same motif to be

present 1.5 kB upstream of the transcription start site of mouse *MPC1* promoter, we analyzed

if tolerization led to reduced ERRα binding to the *MPC1* promoter motif TNAAGGTCA.

Chromatin immunoprecipitation (ChIP) assay followed by qPCR analysis revealed that in

tolerized cells, ERRα binds approximately 4-fold less efficiently to the putative binding site on

the *MPC1* than in corresponding control cells (Fig. 3C).

Since the transcriptional activity of ERRα is regulated through a protein-protein interaction with

the transcriptional coactivator PGC-1α [33] we assessed whether the formation of the PGC-

1α/ERRα complex is affected in 2-AA tolerized cells. Using paraformaldehyde fixed nuclear

lysates of RAW 264.7 tolerized and naïve cells, we performed a co-immunoprecipitation (co-

IP) of PGC-1α and reverse co-IP with ERRα followed by immunoblotting (Fig. 3D). As shown

in Figure 3D less ERRα was detected when PGC-1α was pulled down in presence of 2-AA

than in naïve control cells. Moreover, reverse co-IP with ERRα protein showed lower levels of

PGC-1α in the presence of 2-AA than in naïve control cells (Fig. 3D). Control IgG showed no

detectable levels of ERRα following nuclear fraction pull down. These results indicate that in

the tolerized macrophages, PGC-1α/ERRα interaction is impaired (Fig. 3D).

Since PGC-1α enhances the transcriptional activity of ERRα through their interaction,

and ERRα transcription is regulated *via* an autoregulatory loop [34], we examined the effect of

tolerization on the transcription of *ERRα* and its target gene *MPC1*. Using RAW 264.7 cells

infected with the wild type *PA* strain PA14 or isogenic Δ*mvfR* mutant*,* which does not produce

2-AA [21] we observed lower RNA transcript levels of *MPC1* (Fig. 4A) and *ERRα* (Fig. 4B) in

PA14 compared to Δ*mvfR*. The addition of 2-AA to Δ*mvfR* decreased the levels of both gene

transcripts to the levels similar to PA14 infection condition, supporting the role of 2-AA in effect.

To test if the observed effects are due to MPC1 dependent reduction of mitochondrial ATP

generation, we supplemented the macrophages with ATP. The findings indicate that the

addition of ATP in PA14 infected cells elevated the transcript levels of both *ERRα* and *MPC1*,

but not in the Δ*mvfR* infected macrophages (Fig. 4A and 4B). These results indicate that

reduced MPC1 function is due to negative effect of 2-AA tolerization on the transcriptional

activation of *MPC1* through PGC-1α/ERRα axis.

2-AA tolerization impairs macrophage-mediated intracellular bacterial clearance

through decrease in MPC1-mediated pyruvate import, ATP, and TNF-α levels.

2-AA-mediates persistence of *PA in vivo,* dampens proinflammatory responses and

increases the intracellular burden of this pathogen in macrophages *via* epigenetic

modifications [17, 18, 26]. Here we used RAW 264.7 cells that were infected with the PA14 or

the 2-AA deficient Δ*mvfR* mutant to assess the clearance of *PA* by macrophages.

Macrophages infected with PA14 showed increased bacterial burden than the cells infected

with Δ*mvfR,* and 2-AA supplementation to Δ*mvfR* lead to increase bacterial burden (Fig. 4C).

Furthermore, we used UK5099 and ATP to interrogate whether MPC1 mediated mitochondrial

pyruvate import and bioenergetics are linked to the clearance of *PA* intracellular burden. The

addition of the UK5099 inhibitor strongly enhanced the bacterial intracellular burden in Δ*mvfR*

infected macrophages compared to the non-inhibited cells (Fig. 4C). UK5099 had no effect on

PA14 infected macrophages. Conversely, exogenously added ATP to macrophages infected

with PA14 strongly reduced the *PA* intracellular and had no change Δ*mvfR* infected cells (Fig.

4C).

Given that 2-AA tolerization decreases the expression of pro-inflammatory cytokine

TNF-α by hypo-acetylating the core histone 3 lysine 18 acetylation (H3K18ac) mark at TNF-α

promoter [18], we investigated the link between bioenergetics and TNF-α expression in

infected macrophages. UK5099, or ATP was added exogenously to infected macrophages with

PA14 or Δ*mvfR* (Fig. 4D). As shown in Figure 4D, PA14 infected cells showed lower TNF-α

transcript levels compared to the Δ*mvfR* infected cells. Supplementation of 2-AA to Δ*mvfR*

infected cells led to decrease in the TNF-α transcript levels. Moreover, the addition of the MPC1

inhibitor, UK5099 in Δ*mvfR* infected counteracted the increase in TNF-α transcript levels, while

had no effect in PA14 infected macrophages (Fig. 4D). Conversely, exogeneous addition of

ATP to PA14 infected cells enhanced the transcription of TNF-α compared to the untreated

PA14 infected cells, while no changes in TNF-α expression levels were observed in Δ*mvfR*

infected macrophages (Fig. 4D).

These findings strongly suggest that 2-AA tolerization severely impairs macrophages'

ability to facilitate the clearance of *PA* intracellular burden, *via* mechanism which involves

reduction in MPC1-mediated pyruvate import into mitochondria, ATP levels and TNF-α

transcription.

***In vivo* studies corroborate the 2-AA-mediated decrease *in vitro* of the central metabolic**

fuel acetyl-CoA and the energy-carrying molecule ATP and their association with *PA*

persistence.

To determine whether the decrease in the key metabolites mediated by 2-AA is also

observed *in vivo* during infection, we quantified the levels of ATP and acetyl-CoA in murine

spleen tissues at 1, 5, and 10 days (Fig. 5). This organ was selected for our immunometabolic

studies due to its role in regulating not only local but also systemic (whole-body) immune

responses, facilitated by various immune cells including macrophages [44]. Mice were infected

with *PA* strain PA14, or isogenic mutant Δ*mvfR,* which does not produce 2-AA [21]. It is

important to note that it is not possible to generate or utilize a bacterial mutant that is defective

in 2-AA only because 2-AA is formed by spontaneous decarboxylation rather than by an

enzyme-catalyzed reaction [45-47]. Therefore, animals that were either infected with Δ*mvfR*

and received 2-AA (Δ*mvfR*+2-AA) at the time of infection or uninfected mice injected with a

single dose of 2-AA served as direct controls (Fig. 5). Naïve and sham mice groups served as

additional basal controls (Fig. 5).

Infection with PA14 or uninfected mice injected with 2-AA led to a significant decrease

in both ATP (Fig. 5A) and acetyl-CoA (Fig. 5B) concentrations in spleen tissues compared to

Δ*mvfR* infected mice that sustained higher ATP and acetyl-CoA levels similar to naïve and

sham control groups across the time points tested. However, in mice infected with Δ*mvfR* and

2-AA injected *(ΔmvfR+2-AA)* at the time of infection, ATP and acetyl-CoA concentrations in the

spleen decreased to levels comparable to those of PA14-infected animals, strongly indicating

the biological function of 2-AA in decreasing these key metabolites. These *in vivo* findings

further support the adverse action of 2-AA on host energy homeostasis and metabolism

observed *in vitro* studies.

The metabolic alterations observed are associated with the persistence of *PA* (Fig. 5C). Using

Pseudomonas isolation agar plates, we evaluated the bacterial load at the infection site over

the course of 10 days, obtaining samples at 1, 5, and 10 days. At one day post-infection, mice

infected with PA14, Δ*mvfR,* or Δ*mvfR*+2-AA exhibited no difference in bacterial burden (Fig.

5C), verifying the ability of all strains to establish infection. At 5- and 10-days post-infection,

mice infected with PA14 or Δ*mvfR*+2-AA sustained the bacterial burden at a significantly higher

bacterial burden over time compared to those infected with Δ*mvfR* (Fig. 5C). These results are

consistent with our previous findings [17, 18] and provide compelling evidence that the

metabolic alterations identified may favor *PA* persistence in infected tissues.

Discussion

The *PA* signaling molecule 2-AA that is abundantly produced and secreted in human tissues

[21, 22] is the first QS molecule that epigenetically reprograms immune functions, promotes

immune tolerance, and sustains *PA's* presence in host tissues [17, 18, 22, 26]. The present

study provides insights into the mechanistic actions of 2-AA on cellular metabolism that

contribute to host immune tolerance to infection. We uncover that this signaling molecule

causes distinct metabolic alterations in macrophages' mitochondrial respiration and energy

production promoted via the PGC-1α/ERRα axis and MPC1-dependent OXPHOS that links

the TCA cycle to the production of ATP and energy homeostasis (Fig. 6). Our results show that

2-AA tolerization decreases ATP the crucial energy metabolite and the histone acetylation

metabolite acetyl-CoA and *in vivo*, implicating the importance of the key energy-producing

mitochondrial process in immune tolerance. Although macrophages respond to the first

exposure to 2-AA by enhancing ATP and acetyl-CoA production, long-term exposure to 2-AA

leads to a tolerized state characterized by sustained reduced ATP and acetyl-CoA

concentrations, a quiescent bioenergetic state with reduced pyruvate flux and

unresponsiveness to a 2nd exposure.

Our results reveal that the decrease of ATP and acetyl-CoA in tolerized macrophages

results from the 2-AA-mediated perturbation of the physical interaction between ERRα and

PGC-1α. The PGC-1α/ERRα axis has not previously been implicated in tolerization but has

been extensively studied in cancer, underscoring the novelty of our findings. Studies in cancer

cells have shown that the physical interaction of human ERRα with PGC-1α strongly enhances

the binding of ERRα to DNA to induce the activation of the targeted genes [33] including *MPC1,*

responsible for the import of cytosolic pyruvate into mitochondria and PDH, that catalyzes the

conversion of pyruvate to acetyl-CoA. Inhibition of ERRα has been shown in cancer cell

metabolism studies to interfere with pyruvate entry in mitochondria by inhibiting the expression

of MPC1 [48]. Indeed, we find that the transcription of *ERRα* and *MPC1* genes and pyruvate

entry into mitochondria is reduced in tolerized macrophages. Given that pyruvate is a primary

carbon source in the TCA cycle that fuels OXPHOS and the production of ATP into

mitochondria, it explains the reduced ATP and acetyl-CoA levels observed in tolerized

macrophages. Future studies will focus on unveiling the pathways by which macrophages

respond to the bioenergetic changes identified and decipher their effect on pyruvate cycle in

host tolerance to infection [49-51].

We show that the unprecedented action of a QS bacterial small signaling molecule on

the interaction between ERRα and PGC-1α in the nucleus is hampered. Although it remains to

be elucidated whether 2-AA interferes directly or indirectly on the PGC-1α/ERRα axis, our data

bring up the possibility that the reduced production of acetyl-CoA may be responsible, since

cholesterol is a known coactivator/ligand of ERRα [52] is generated from acetyl-CoA [52]. This

possibility may also explain the reduced expression of *ERRα* and its lower abundance since

cholesterol is also required for its activation and autoregulation. However, it is also possible

that 2-AA may antagonize the interaction between ERRα and PGC-1α by binding to ERRα.

Small molecules as antagonist of ERRα have been reported previously [53]. The weaker

binding of ERRα to the *MPC1* promoter site, and consequently reduced MPC1 expression

explains the reduced pyruvate presence in mitochondria and that of ATP and mitochondrial

quiescence.

The observed cellular metabolic perturbances also align with our previous studies in

skeletal muscle, which pointed to mitochondrial dysfunction triggered by 2-AA [24, 25].

Interestingly, the *PA* QS LasR-regulated homoserine lactone molecule, 3-oxo-C12-HSL, was

reported to attenuate the expression of PGC-1α and decreasing the mitochondrial respiratory

capacity [27] and promoting apoptosis in lung epithelial cells [54]. Moreover, another *PA* QS

molecule PQS also regulated by LasR was recently shown to induce organelle stress, including

mitochondria, by disrupting the mitochondrial membrane potential in human macrophages [55].

As opposed to 2-AA, which dampens the pro-inflammatory response, PQS increases

proinflammatory cytokines [55], consistent with the fact that PQS is an acute infection type

molecule unrelated to chronic/persistent infections.

Previously, we reported that 2-AA tolerization induces HDAC1 expression and activity,

reducing H3K18ac at the TNF-α promoter [18]. The findings with TNF-α transcription using

UK5099 and ATP in 2-AA   treated macrophages are in support of the bioenergetics

disturbances observed in macrophages and their link to epigenetic modifications we have

shown to be promoted by 2-AA [18]. Macrophages exposed to 2-AA in the presence of

exogenous ATP showed improved intracellular bacterial clearance and enhanced TNF-α

levels, supporting the epigenetic interconnection of macrophages and cellular ATP

responsiveness levels against infection. The exogenous addition of UK5099 that reverted the

efficiency of the *mvfR* infected macrophages to clear the *PA* intracellular burden and the

counteraction to the 2-AA effect by ATP addition that increases the clearance of *PA* intracellular

burden support the importance of this energy-carrying molecule in *PA* persistence.

The results with the MPC1 inhibitor, UK5099, suggest that the availability of pyruvate

could underlie the mechanism of 2-AA-regulated HDAC/HAT-dependent control of transcription

of pro-inflammatory mediators and bacterial clearance [56]. These results align with our

previous findings, showing that establishing bacterial intracellular burden in macrophages is

HDAC1 dependent [56]. It remains to be elucidated if HDAC/HAT-mediated histone

modification directly regulates expression of OXPHOS genes in 2-AA tolerized macrophages

as it was shown that histone deacetylation downregulates OXPHOS in persistent *M.*

tuberculosis infection [57, 58]. Given the complexity of 2-AA-mediated long-term effects, future

omics studies combined with immune profiling may aid in deciphering the possible network of

co-factors across different subpopulations of immune cells and the immunometabolic

reprogramming related to 2-AA.

We have shown that although 2-AA tolerization leads to a remarkable increase in the

survival rate of infected mice, it permits an HDAC1-dependent sustained presence of *PA* in

mice tissues and intracellularly in macrophages [17, 18, 22, 26]. Here, we use 2-AA-producing

and isogenic non-producing *PA* strains to confirm the 2-AA-mediated decrease in the central

metabolite acetyl-CoA and the energy-carrying molecule ATP during infection. These murine

infection studies provide strong evidence of the 2-AA biological relevance in reducing these

metabolites *in vivo*. Taking together our previous and current findings, these *in vivo* studies

provide compelling evidence that the metabolic alterations identified may favor *PA* persistence

in infected tissues.

That 2-AA permits *PA* to persist in infected tissues despite rescuing the survival of

infected mice underscores its difference from that of LPS tolerization, which results in bacterial

clearance [23] *via* the mechanism that relies on a set of different HDAC enzymes. LPS

tolerization predominantly involves changes in H3K27 acetylation [59], while 2-AA tolerization

involves H3K18 modifications [22]. The 2-AA-mediated effects reported here are also distinct

from Sirtuin 1 and 6-dependent epigenetic-metabolic reprogramming mediated by LPS [60],

which promotes endotoxin tolerance in immune cells by upregulating glycolysis and

suppressing OXPHOS [61]. Although short-term exposure of BMDMs to LPS enhances ATP

production, we could not observe this effect in 2-AA tolerized cells.

Metabolic reprogramming upon infection may be pathogen-specific, with each

pathogen impacting specific metabolic pathways that better fit its respective metabolic needs

[7]. This was shown for infections of various tissues with *Chlamydia pneumonia*, *Legionella*

pneumophila, *Mycobacterium tuberculosis*, and *Salmonella* [62-65] and, by our group, for *PA*

infections [17, 18, 24]. It would be interesting, however, to test whether 2-AA affects the killing

efficacy of macrophages against other pathogens, as emerging evidence shows that

synergistic or antagonistic interactions between clinically relevant microorganisms and host

have important implications for polymicrobial infectious diseases [66].

While in this study, we focused on the role of ERRα mainly in pyruvate metabolism,

future studies are needed to reveal other ERRα/PGC-1α axis-dependent metabolic and

immune pathways are modulated by 2-AA. In this view, the cholesterol synthesis pathway that

relies on acetyl-CoA precursor, as cholesterol is a known coactivator/ligand of ERRα [52] and

fatty acid catabolism, generating acetyl-CoA, and PGC-1α regulates fatty acid oxidation [67].

In summary, our study highlights the role of 2-AA in orchestrating cellular metabolic

reprogramming in addition to the epigenetic reprogramming reported previously. This

immunometabolic reprogramming that ultimately leads to host tolerance favoring the prolonged

presence of *PA* in the infected host resembles trained immunity, a nonspecific form of immune

memory in which an initial exposure to a pathogen "trains" innate immune cells to respond to

a re-exposure differently than their unexposed counterparts with the help of epigenetic

changes [68, 69]. The reported immunometabolic reprogramming contributes to a better

understanding of the molecular and cellular mechanisms, biomarkers, and functional

significance that may be involved in immune tolerance to infection, providing for designing and developing innovative therapeutics and interventions. These approaches can focus on promoting host resilience against bacterial burden and safeguarding patients from recalcitrant persistent infections.

Materials & Methods

Ethics statement

The animal protocol was approved by the Institutional Animal-Care and Use Committee (IACUC) of Massachusetts General Hospital (protocol no. 2006N000093). No randomization or exclusion of data points was applied.

Conditions used for the exposures of the cells to 2-AA

Mouse BMDM cells were isolated from femur of CD1 6 weeks old male and female mice and used to estimate the levels of various metabolites (Fig. 2A) following exposure and re-exposure to 2-AA (Fig. 1A). The 1st exposure of naïve BMDMs (10^6/mL in 6-well plates) was achieved with 400 µM 2-AA for 1, 6, or 48 h. Cells exposed for 48 h were used for the 2nd 2-AA exposure. These cells were washed to remove residual 2-AA, left in the medium in the absence of 2-AA for either 24 or 106 h, and re-exposed to 2-AA (200 µM) for 1 or 6 h, respectively. For RAW 264.7 (10^6/mL in 6-well plates) and THP-1 cells (10^6/mL in 6-well plates) the 2-AA concentration and conditions used were as with BMDM cells. The hours of the 1st round and 2nd round of exposure to 2-AA are indicated in Figure S1.

For the LPS stimulation studies, cells were stimulated with 400 µM 2-AA or 100 ng/mL LPS for 6 h, and 800 µM of 2-AA for 48 h. Cells receiving 1st exposure for 48 h were washed and re-exposed to 400 µM of 2-AA or 100 ng/mL LPS for 6 h.

ATP and acetyl-CoA quantifications

The levels of ATP, acetyl-CoA, and were assessed in BMDM and RAW264.7 macrophage cells

following exposure to 2-AA at the times indicated and by utilizing the ATP Assay kit

(cholorimetric/fluorometric) (#ab83355, Abcam) and the PicoProbe AcCoA assay kit (ab87546,

Abcam), respectively, according to the manufacturer's instruction. Quantifications of ATP and

acetyl-CoA were performed in triplicates.

ATP: For ATP determination, macrophages were lysed and subsequently centrifuged at 14,000

rpm for 10 min at 4 °C. The supernatants were transferred to a fresh Eppendorf tube.

Standards and cell supernatants of 50 µL were added to a 96-well plate suitable for fluorescent

analysis (black sides, clear bottom). A reaction mixture containing an ATP converter, probe,

buffer, and developer mix was then added to all wells (50 µL) and incubated away from light at

room temperature for 30 min. ATP quantification was conducted fluorometrically at 535/587 nm

using a plate reader and the following settings: $\lambda ex = 535\,nm$; $\lambda em = 587\,nm$. Fluorescence

was measured using a microplate reader (Tecan Group Ltd, Männedorf, Switzerland)

Acetyl-CoA: Acetyl-CoA content was assessed by first deproteinizing total cell fractions of

macrophages using the perchloric acid and then centrifuging at 14,000 rpm for 10 min at 4 °C.

50 µL of cells' supernatant sample CoASH were quenched to correct the background

generated by free CoASH. Following the homogenization procedure described above, the

samples were diluted with the reaction mix, and fluorescence was quantified using a plate

reader and the settings as above with ATP.

Seahorse Assays

Seahorse analysis was performed according to previously published protocols [70]. Briefly,

freshly prepared BMDM cells were reseeded in complete RPMI-1640 cells using Seahorse

plates (Agilent cat. no. 103729100) at a density of 5×10^4 cells per well. Cells were exposed

to 400 µM 2-AA for 1 h or 24 hours (1st exposure). Cells receiving 1st exposure for 24 h were

washed and re-exposed to 200 µM of 2-AA for 1 h. Naïve cells were used as control. Prior to

initiating Seahorse measurements cells were washed and Seahorse XF DMEM supplemented

with 2 mM glutamine (Gibco cat. No. 25030-081) was added to each well. Cells were allowed

to stabilize in a 37°C incubator without CO_2 for 1 h. The Seahorse cartridge was hydrated and

calibrated as per the manufacturer's instructions. The Mitochondrial stress test kit (Agilent cat.

no. 10395-100) was used (oligomycin at 1 µM, FCCP (1.5 µM), rotenone (0.5 µM), and

antimycin A at (0.5 µM)) with slight modifications according to the published protocol [70] that

included injection of 25 mM glucose, and sodium pyruvate (1 µM). All samples N= 4 were run

in a Seahorse XFe96 Analyzer, and data were analyzed using Wave and plotted using

GraphPad Prism software.

Isolation of cytosolic and mitochondrial fraction and pyruvate quantification

Pyruvate is produced in the cytosol and is transported into the mitochondria. RAW 264.7 cells

were plated at 1×10^6 to incubate overnight at 37°C in a CO2 incubator. Cells were exposed

to 2-AA for 1, 3, 24 or 48 h or re-exposed for 1 h as described above. Mitochondria and

cytosolic fractions of each group were isolated utilizing the Mitochondria Isolation Kit for

cultured cells (#ab110170, Abcam) following the manufacturer's protocol. Briefly, cells were

collected with a cell lifter and pelleted by centrifugation at 1,000 g, frozen, and then thawed to

weaken the cell membranes. The cells were resuspended in Reagent A and transferred into a

pre-cooled Dounce Homogenizer. The homogenates were centrifuged at 1,000 g for 10 min at

4°C and saved as supernatant #1 for the cytosolic fractions. The pellet was resuspended in

Reagent B, followed by repeat rupturing and centrifugation. The pellet was collected and

resuspended in 500 µL of Reagent C supplemented with Protease Inhibitor cocktails (P8340,

Sigma Aldrich). Following separation, a Bradford assay was conducted to determine the

protein concentration in each fraction. Pyruvate levels were determined in cellular fractions

and mitochondrial fractions by using the Pyruvate Assay Kit (#ab65342, Abcam) according to

the manufacturer's instructions. Briefly, after deproteinization using perchloric acid, the

samples were neutralized in ice-cold 2 M KOH. 10 µL samples were incubated with reaction

mix and kept on the plate at RT for 30 min in the dark. The absorbance was measured in a

microplate reader (Tecan Group Ltd, Männedorf, Switzerland) at 570 nm and results are shown

from three independent experiments.

Bacterial strains and growth conditions

509   The *P. aeruginosa* strain known as Rif[R] human clinical isolate UCBPP-PA14 (also known as

510   PA14) was used [71]. The bacteria were grown at 37 °C in lysogeny broth (LB) under shaking

511   and aeration or on LB agar plates containing appropriate antibiotics. PA14 and isogenic mutant

512    $\Delta mvfR$ [19] cultures were grown in LB from a single colony to an optical density of 600 nm

513   (OD_{600}) of 1.5, diluted 1:50,000,000 in fresh LB media, and grown overnight to an OD_{600} of 3.0.

Gentamicin protection assay

RAW264.7 macrophage cells were plated on 6 well cell culture-treated plates overnight in

pyruvate free DMEM. After 3 h of incubation with 10 μM UK5099, 20 μM ATP or 400 μM 2-AA,

cells were infected with PA14 and isogenic mutant $\Delta mvfR$ at 5 MOI for 30 min at 37 °C in 5%

CO2. Unbound bacteria were removed by washing once with cold DMEM medium. Afterwards,

the cells were incubated with 100 μg/ml of gentamicin for 30 min to eliminate residual

extracellular bacteria. The cells were washed with DMEM and transferred to free medium

without gentamicin and kept for 3 h at 37°C in 5% CO2. The infected cells were scraped after

3 h, centrifuged at 500 × g, and lysed in distilled water. The lysed cells were immediately diluted

in PBS and plated on LB agar plates to assess bacterial presence. Bacterial colony-forming

units (CFUs) were counted after incubating the plates overnight at 37°C.Untreated cells

infected with PA14 were set as 100 % and the reduction of Bacterial load was expressed as

%CFU.

Pharmacological inhibitors and ATP supplementations

For the OXPHOS inhibition assay, RAW264.7 macrophage cells were treated with UK5099 (10

µM, Sigma-Aldrich, dissolved in DMSO 3 h prior to 1st or 2nd rounds of 2-AA exposure.

RAW264.7 macrophages supplemented with ATP (20 µM, Sigma-Aldrich, dissolved in PBS)

received ATP at the time of 1st and 2nd 2-AA exposure.

Co-Immunoprecipitation assay

For all immunoprecipitation assays, protein A/G agarose Magnetic beads (Pierce) were used

after washing in 1X IP buffer. Nuclear lysates from RAW 264.7 tolerized cells exposed to 2-AA

for 24 h or non-exposed cells were diluted in 1X IP buffer and 500 µg of protein was taken for

each experiment. The lysates were precleared by using unbound 50 µl protein A/G magnetic

beads for 2 h at room temperature on a rotator. Precleared lysates were either incubated with

ERRα, PGC-1α or rabbit IgG antibody overnight at 4 °C. 100 µl protein A/G magnetic beads

were used to pull down the antibody-protein complex and washed twice with IP buffer to

remove unbound proteins. Magnetic beads were then eluted in 2X Laemmli SDS-PAGE

loading buffer. Immunoprecipitated ERRα and PGC-1α was detected by Western blot

analyses, using conformational specific Anti Rabbit antibody (TruBlot)**.**

Immunoblotting analysis

Cells were plated at 6×10^5 cells per well in six-well plates. Cells were washed with PBS and

subsequently lysed using RIPA lysis buffer containing 1 mM phenylmethylsulfonyl fluoride

(PMSF). 20µg of proteins were separated by electrophoresis on any KD (Kilo Dalton) (Bio-

Rad, Cat no. 4569033) SDS-polyacrylamide gel. Proteins were transferred to a 0.2 µm

polyvinylidene fluoride membrane (PVDF, Millipore, Billerica, USA) using a Bio-Rad semi-dry

instrument. After blocking with 5% BSA in TBS containing 0.1% Tween-20 for 1 h at room

temperature, the membranes were incubated with a primary antibody ERR-a (Abcam,

#ab76228), MPC1 (D2L91, Cell signaling), and anti-β-actin (cat no: sc-47778) (Santa Cruz

Biotechnology) overnight at 4°C. Following washing, the membranes were incubated with an

anti-rabbit secondary antibody, and the bands were detected by SuperSignal West Pico Chemiluminescent Substrate (Thermo Scientific) reaction, according to the manufacturer's instructions. The blots were visualized in the ChemiDOC Imaging system (Bio-Rad Laboratories, Inc., Hercules, CA, USA). The bands were analyzed densitometrically using QuantityOne software (Bio-Rad).

Chromatin immunoprecipitation (ChIP) and ChIP-qPCR

For ChIP studies, RAW264.7 macrophage cells exposed to 2-AA for 24 h were cross-linked in 1% (v/v) methanol-free formaldehyde for 10 min and then placed in 0.125 M glycine for 5 min at RT. Using the truChIP High Cell Chromatin Shearing kit (Covaris, United States), cells were prepared for sonication according to the Covaris protocol. Approximately 1×10^7 cells were plated in a 12 mm × 12 mm tube and subjected to shearing with the Covaris S220 sonicator for 8 min (140 peak power, 5 duty factor, 200 cycles/burst). The Magna ChIP A/G kit (Millipore, United States) was used for the subsequent immunoprecipitations according to the manufacturer's protocol. Briefly, chromatin from approximately 10^6 cells was incubated overnight at 4°C with 2 µg of anti-ERRα or anti-PGC-1α (Abcam, USA) ChIP-grade antibody and 20 µL of A/G magnetic beads. The beads were washed serially (5 min each) with low-salt wash buffer, high-salt wash buffer, LiCl wash buffer, and TE buffer from the kit at 4°C. Chromatin was eluted with elution buffer containing Proteinase K at 62°C for 4 h, then incubated at 95°C for 10 min. DNA was isolated by column purification (QIAquick PCR purification kit).

Real-time ChIP-qPCR was performed with the Brilliant II SYBR green super mix (Agilent, USA). Forward (AGTGGTGACCTTGAACTTCCC) and reverse (CTGAAGACGACCTTCCCCTT) primers were chosen to amplify a genomic locus of *MPC1* promoter which had a putative ERR binding site (gtgaccttg) at 1582 bp upstream of the start site. Normalized values were calculated by the percent-input method relative to the IgG. The assay was performed three times.

RNA extraction and RT-qPCR

Total RNA from all the groups mentioned above was isolated from approximately 2×10^6 cells with the RNeasy minikit (Qiagen, USA), and cDNA was prepared with the iScript Reverse transcription kit (Bio-Rad, USA), as per the manufacturer's instruction. Real-time PCR was conducted using the PowerUP SYBR green Master mix (Applied Biosystem, USA) and primer sets for mouse ERR-α (forward: ACTACGGTGTGGCATCCTGTGA; reverse: GGTGATCTCACACTCATTGGAGG), MPC-1 (forward: CTCCAGAGATTATCAGTGGGCG; reverse: GAGCTACTTCGTTTGTTACATGGC), TNF-α (forward: GGTGCCTATGTCTCAGCCTCTT; reverse: GCCATAGAACTGATGAGAGGGAG) and mouse 18S rRNA (forward: GTTCCGACCATAAACGATGCC; reverse: TGGTGGTGCCCTTCCGTCAAT). The transcript levels of all the genes were normalized to 18S rRNA with the ΔΔCT method. The relative expression was calculated by normalizing transcript levels to those of PA14 Infected cells. The assay was conducted in triplicate; means and standard deviations were calculated for each group.

Animal infection and metabolites assessment experiments

The full-thickness thermal burn injury and infection (BI) model [17, 18, 72] was used to assess the effect of 2-AA on metabolic alterations in the spleens of six-week-old CD1 male mice and bacterial burden (Charles River Labs, USA). A full-thickness thermal burn injury involving 5-8% of the total body surface area was produced on the shaved mouse abdomen dermis, and an inoculum of ~1×10^4 PA14 and Δ*mvfR* [19] cells in 100 μL of $MgSO_4$ (10 mM) was injected intradermally into the burn eschar. For the groups that received 2-AA, mice were injected intraperitoneally (IP) with 100 μL of 2-AA (6.75 mg/kg). The entire procedure was done under the influence of anesthesia. One of the groups of mice infected with the PA14 isogenic mutant Δ*mvfR* [19] also received 100 μL of 2-AA in PBS (6.75 mg/kg) at the time of infection (Δ*mvfR* +2-AA) and served as an additional control. CFU counts from rectus abdominus muscle (underlying the burn-infected tissue) were assessed in groups of 4 mice each at 1, 5, and 10

d post-BI by plating diluted muscle homogenate on Pseudomonas isolation agar (Sigma-

Aldrich) plates containing rifampicin (50 mg/L). Spleen samples from mice were collected for

metabolite analyses from all mice groups at 1, 5, and 10 d post-BI to assess ATP (#ab83355

Abcam) and acetyl-CoA (#ab87546, Abcam) as described above. Tissues were homogenized,

and 100 mg of homogenate was centrifuged at 4°C for 15 min at 10,000 g, and supernatants

were collected. The supernatant was mixed with 400 µL of 1 M perchloric acid. The

deproteinized supernatant was neutralized by 3 M $KHCO_3$. The ATP and acetyl-CoA assays

were performed as described in the *in vitro* section.

Acknowledgments and Funding sources

614   We thank Dr. Vamsi Mootha and Sneha Rath for access and help with the Seahorse

615   instrument, respectively. This work was supported by the NIH award R01AI134857, The John

616   Lawrence Massachusetts General Hospital Research Scholar Award and Shriner's grant

617   83009 to L.G.R., and the Shriner's grant 85132 to A.A.T. The funders had no role in the study

618   design, data collection, analysis, decision to publish, or manuscript preparation.

Figures

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3

Figure Legends

Figure 1. 2-AA tolerization decreases crucial metabolites of cellular energy and affects mitochondrial respiration in mouse BMDM. (A) Schematic representation showing experimental design: naïve cells were exposed to 2-AA for 1, 6, or 48 h (black). Cells exposed to 2-AA (400 µM) for 48 h were washed, rested for 24 or 106 h, and re-exposed (200 µM) for 1 or 6 h (red), respectively. The same color code: black: cells after 1^{st} exposure; red: cells after 2^{nd} exposure, was kept throughout the manuscript with corresponding controls in grey and pink, respectively. The levels of (B) ATP and (C) acetyl-CoA in BMDM cells after 1^{st} and 2^{nd} 2-AA exposure. (D) Real-time oxygen consumption rate (OCR) traces were recorded using a Seahorse XF analyzer and normalized to protein content. Cells were exposed to 2-AA for 1 or 24 h (black), washed and rested for 24 h and re-exposed for 1 h (red). Mitochondrial respiratory parameters, (E) basal respiration, and (F) maximal respiration. Data are presented as mean ± SD. n ≥ 4, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ns indicates no significant difference. One-way ANOVA followed by Tukey's post hoc test was applied.

Figure 2. 2-AA perturbs the mitochondrial MPC1-mediated import and metabolism of pyruvate. (A) Cytosolic and mitochondrial pyruvate levels following 2-AA exposure (black) or re-exposure (red) and corresponding controls in grey and pink respectively, for indicated time points. (B) Representative Western blot and results of densitometric analysis of MPC1 protein levels following 2-AA exposure or re-exposure for indicated time points. β-actin was used as a control. Corresponding controls are shown in grey or pink, respectively. Mean ± SD is shown; $n = 3$; ***$p < 0.001$, ****$p < 0.0001$, and ns indicates no significant difference. One-way ANOVA followed by Tukey's post hoc test was applied.

Figure 3: 2-AA-mediated macrophage tolerization deranges PGC-1α/ERRα-dependent metabolic programing. (A) Representative Western blot and results of densitometric analysis of ERRα protein levels following 2-AA exposure or re-exposure for indicated time points. β-

actin (Fig. 2B) was used as a control. (B) Western blots and its corresponding densitometric

analysis of ERRα in cytoplasmic or nuclear lysates of tolerized macrophages exposed to 2-AA

for 24 h or not exposed to 2-AA. (C) ChIP-qPCR assay of ERRα binding at the MPC1 promoter

in RAW 264.7 tolerized macrophages exposed to 2-AA (200 μM) for 24 h (black) compared to

untreated control macrophages (grey). IgG served as a negative control (D) Representative

Western blot of co-immunoprecipitation (co-IP) studies of ERRα and PGC-1α in nuclear

extracts of 2-AA tolerized (24 h) and control RAW 264.7 cells. Pull down with IgG served as a

negative control. 2-AA tolerized macrophages are shown in black, and untreated control

macrophages in grey. Mean ± SD in shown, n ≥ 3, *$p < 0.05$, **$p < 0.01$, ****$p < 0.0001$, and

ns indicates no significant difference. One-way ANOVA followed by Tukey's post hoc test was

applied.

Figure 4: Increased intracellular burden in macrophages is associated with decreased

expression of *MPC1, ERR-α,* and *TNF-α genes*. Real-time PCR analysis of *MPC1* (A) and

ERR-α (B) expression in RAW 246.7 macrophages infected with PA14 or *ΔmvfR* in the

presence or absence of exogenous addition of 2-AA or ATP for 6 h as indicated. Transcript

levels were normalized to 18srRNA. PA14 infected cells served as controls. (C) Intracellular

burden of PA14 or *ΔmvfR* of infected macrophages in the presence or absence of exogenous

addition of 2-AA, UK5099, or ATP. Untreated cells infected with PA14 were set as 100%. D)

Real-time PCR analysis of TNF-α expression in RAW 246.7 macrophages infected with PA14

or *ΔmvfR* in the presence or absence of exogenous addition of 2-AA, ATP, or UK5099.

Transcript levels were normalized to 18srRNA. PA14 infected cells served as controls. The

compound concentration used for UK5099 was 10 μM and ATP 20 μM. Mean ± SD is shown

($n = 3$); * $p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$***, and ns indicates no significant

difference. One-way ANOVA followed by Tukey's post hoc test was applied.

Figure 5. 2-AA promotes a long-lasting decrease in ATP and acetyl-CoA levels and bacterial persistence in *PA*-infected mice. (A) ATP and (B) acetyl-CoA concentrations in the spleens of mice infected with *PA* wild type (PA14), the isogenic mutant Δ*mvfR*, Δ*mvfR* injected with 2-AA at the time of infection (Δ*mvfR* + 2-AA), or non-infected but injected with 2-AA (6.75 mg/kg). (C) Bacterial burden in muscles expressed as CFU count was analyzed using the Kruskal–Wallis non-parametric test with Dunn's post-test; ***$p < 0.001$, and ns indicates no significant difference. Control mice groups: naïve were not given 2-AA; mice receiving 2-AA were given a single intraperitoneal injection of 2-AA; sham represents a burn/PBS group since the burn and infection model was used. Results of three independent replicates with 4 mice per group for 1, 5, and 10 d are shown. Means ± SD are shown: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ns indicate no significant difference. One-way ANOVA followed by Tukey's post hoc test was applied.

Figure 6: Proposed mechanism by which 2-AA impairs bioenergetics through the inhibition of MPC1-mediated pyruvate transport into mitochondria and its impact on the PGC-1α/ERRα axis. 2-AA tolerized macrophages exhibit diminished pyruvate levels in mitochondria due to the decreased expression of *MPC1*, a consequence of the 2-AA impact on the interaction of ERRα with the transcriptional coactivator PGC-1α for the effective transcription of ERRα, since ERRα controls its own transcription and that of MPC1. In the presence of 2-AA, the weakened interaction between ERRα and PGC-1α results in reduced expression of MPC1 and ERRα. The reduction in mitochondrial pyruvate levels leads to decreased acetyl-CoA and ATP levels, which modulate HDAC1- and HAT-catalyzed remodeling of H3K18 acetylation. The diminished levels of this epigenetic mark have previously been associated with an increased intracellular presence of bacteria in macrophages, as demonstrated by our group [18]. Pathways, proteins, and metabolites that

are negatively affected are indicated in red, while positively affected are denoted in black. The

figure was created using BioRender.com.

Figure S1. 2-AA tolerization decreases metabolites in murine RAW 264.7 (A-B) and human

THP-1 (C-D) cells. (A and C) ATP, (B and D) acetyl-CoA and (E) cytosolic and (F) mitochondrial

pyruvate levels were quantified in 2-AA (400 μM) exposed and re-exposed cells for the hours

indicated using the same condition as shown in Fig. 1A. Mean ± SD is shown (n = 3); *** p <

0.001, and ns indicates no significant difference. One-way ANOVA followed by Tukey's post

hoc test was applied. The same color code: black, cells after 1st exposure; red, cells after 2nd

exposure was kept as throughout the main manuscript with corresponding controls shown in

grey and pink, respectively.

Figure S2: ATP levels in murine BMDM (A) cells with and without 2-AA (400 μM) or LPS

stimulation (100 μg/mL). Each dot represents 1 experimental replicate of 4 independent

experiments (n = 4). Data are presented as mean ± SD. **p < 0.01, ***p < 0.001, and ns

indicates no significant difference. One-way ANOVA followed by Tukey's post hoc test was

applied.

Figure S3: Effect of 2-AA on mitochondrial spare respiratory capacity in BMDM non-tolerized

(black) and tolerized (red) macrophages. Controls are shown in grey and pink, respectively.

Mitochondrial spare respiratory capacity data extrapolated from the OCR profiles shown in

Figure 1D. Unstimulated BMDM macrophages were used as a control (c). Means ± SDs are

shown (n = 6); * p < 0.05, *** p < 0.001, and ns indicates no significant difference. One-way

ANOVA followed by Tukey's post hoc test was applied.

References

1. Foster, S.L., D.C. Hargreaves, and R. Medzhitov, *Gene-specific control of inflammation by tlr-induced chromatin modifications.* Nature, 2007. **447**(7147): p. 972.
2. Bagchi, A., et al., *Myd88-dependent and myd88-independent pathways in synergy, priming, and tolerance between tlr agonists.* Journal of immunology, 2007. **178**(2): p. 1164.
3. Masyutina, A.M., et al., *Inhibition of specific signaling pathways rather than epigenetic silencing of effector genes is the leading mechanism of innate tolerance.* Frontiers in Immunology, 2023. **14**.
4. O'Neill, L.A., R.J. Kishton, and J. Rathmell, *A guide to immunometabolism for immunologists.* Nat Rev Immunol, 2016. **16**(9): p. 553.
5. Chi, H., *Immunometabolism at the intersection of metabolic signaling, cell fate, and systems immunology.* Cellular & Molecular Immunology, 2022. **19**(3): p. 299.
6. Tannahill, G.M., et al., *Succinate is an inflammatory signal that induces il-1beta through hif-1alpha.* Nature, 2013. **496**(7444): p. 238.
7. Galli, G. and M. Saleh, *Immunometabolism of macrophages in bacterial infections.* Front Cell Infect Microbiol, 2020. **10**: p. 607650.
8. Wang, Y., et al., *Mitochondrial metabolism regulates macrophage biology.* J Biol Chem, 2021. **297**(1): p. 100904.
9. Karan, K.R., et al., *Leukocyte cytokine responses in adult patients with mitochondrial DNA defects.* Journal of Molecular Medicine, 2022. **100**(6): p. 963.
10. Bird, L., *Getting enough energy for immunity.* Nature Reviews Immunology, 2019. **19**(5): p. 269.
11. Deziel, E., et al., *The contribution of mvfr to pseudomonas aeruginosa pathogenesis and quorum sensing circuitry regulation: Multiple quorum sensing-regulated genes are modulated without affecting lasri, rhlri or the production of n-acyl-l-homoserine lactones.* Mol Microbiol, 2005. **55**(4): p. 998.
12. Xiao, G., et al., *Mvfr, a key pseudomonas aeruginosa pathogenicity lttr-class regulatory protein, has dual ligands.* Mol Microbiol, 2006. **62**(6): p. 1689.
13. Eickhoff, M.J. and B.L. Bassler, *Snapshot: Bacterial quorum sensing.* Cell, 2018. **174**(5): p. 1328.
14. Fuqua, C. and E.P. Greenberg, *Listening in on bacteria: Acyl-homoserine lactone signalling.* Nature Reviews Molecular Cell Biology, 2002. **3**(9): p. 685.
15. Kariminik, A., M. Baseri-Salehi, and B. Kheirkhah, *Pseudomonas aeruginosa quorum sensing modulates immune responses: An updated review article.* Immunol Lett, 2017. **190**: p. 1.
16. Liu, Y.C., K.G. Chan, and C.Y. Chang, *Modulation of host biology by pseudomonas aeruginosa quorum sensing signal molecules: Messengers or traitors.* Front Microbiol, 2015. **6**: p. 1226.
17. Bandyopadhaya, A., et al., *The quorum sensing volatile molecule 2-amino acetophenon modulates host immune responses in a manner that promotes life with unwanted guests.* PLoS pathogens, 2012. **8**(11): p. e1003024.
18. Bandyopadhaya, A., et al., *A quorum-sensing signal promotes host tolerance training through hdac1-mediated epigenetic reprogramming.* Nat Microbiol, 2016. **1**: p. 16174.
19. Cao, H., et al., *A quorum sensing-associated virulence gene of pseudomonas aeruginosa encodes a lysr-like transcription regulator with a unique self-regulatory mechanism.* Proceedings of the National Academy of Sciences of the United States of America, 2001. **98**(25): p. 14613.
20. Deziel, E., et al., *Analysis of pseudomonas aeruginosa 4-hydroxy-2-alkylquinolines (haqs) reveals a role for 4-hydroxy-2-heptylquinoline in cell-to-cell communication.* Proc Natl Acad Sci U S A, 2004. **101**(5): p. 1339.

21.   Kesarwani, M., et al., *A quorum sensing regulated small volatile molecule reduces acute*
virulence and promotes chronic infection phenotypes. PLoS Pathog, 2011. **7**(8): p. e1002192.

22.   Bandyopadhaya, A., A. Tsurumi, and L.G. Rahme, *Nf-kappabp50 and hdac1 interaction is*
implicated in the host tolerance to infection mediated by the bacterial quorum sensing signal
2-aminoacetophenone. Front Microbiol, 2017. **8**: p. 1211.

23.   Wheeler, D.S., et al., *Induction of endotoxin tolerance enhances bacterial clearance and*
survival in murine polymicrobial sepsis. Shock, 2008. **30**(3): p. 267.

24.   Tzika, A.A., et al., *A small volatile bacterial molecule triggers mitochondrial dysfunction in*
murine skeletal muscle. Plos One, 2013. **8**(9): p. e74528.

25.   Bandyopadhaya, A., et al., *Bacterial-excreted small volatile molecule 2-aminoacetophenone*
induces oxidative stress and apoptosis in murine skeletal muscle. International Journal of
Molecular Medicine, 2016. **37**(4): p. 867.

26.   Chakraborty, A., et al., *Quorum-sensing signaling molecule 2-aminoacetophenone mediates*
the persistence of pseudomonas aeruginosa in macrophages by interference with autophagy
through epigenetic regulation of lipid biosynthesis. mBIo, 2023: p. e0015923.

27.   Maurice, N.M., et al., *Pseudomonas aeruginosa induced host epithelial cell mitochondrial*
dysfunction. Scientific Reports, 2019. **9**(1).

28.   Lelliott, C.J. and A. Vidal-Puig, *Pgc-1beta: A co-activator that sets the tone for both basal and*
stress-stimulated mitochondrial activity. Adv Exp Med Biol, 2009. **646**: p. 133.

29.   Supruniuk, E., A. Mikłosz, and A. Chabowski, *The implication of pgc-1α on fatty acid transport*
across plasma and mitochondrial membranes in the insulin sensitive tissues. Front Physiol,
2017. **8**: p. 923.

30.   Coppi, L., et al., *Pgc1s and beyond: Disentangling the complex regulation of mitochondrial*
and cellular metabolism. International Journal of Molecular Sciences, 2021. **22**: p. 6913.

31.   Lin, J., C. Handschin, and B.M. Spiegelman, *Metabolic control through the pgc-1 family of*
transcription coactivators. Cell Metab, 2005. **1**(6): p. 361.

32.   Huss, J.M., W.G. Garbacz, and W. Xie, *Constitutive activities of estrogen-related receptors:*
Transcriptional regulation of metabolism by the err pathways in health and disease. Biochim
Biophys Acta, 2015. **1852**(9): p. 1912.

33.   Schreiber, S.N., et al., *The transcriptional coactivator pgc-1 regulates the expression and*
activity of the orphan nuclear receptor estrogen-related receptor alpha (erralpha). J Biol
Chem, 2003. **278**(11): p. 9013.

34.   Laganière, J., et al., *A polymorphic autoregulatory hormone response element in the human*
estrogen-related receptor α (errα) promoter dictates peroxisome proliferator-activated
receptor γ coactivator-1α control of errα expression*. Journal of Biological Chemistry, 2004.
279(18): p. 18504.

35.   Villena, J.A., *New insights into pgc-1 coactivators: Redefining their role in the regulation of*
mitochondrial function and beyond. The FEBS journal, 2015. **282**(4): p. 647.

36.   Ranhotra, H.S., *The estrogen-related receptor alpha: The oldest, yet an energetic orphan with*
robust biological functions. J Recept Signal Transduct Res, 2010. **30**(4): p. 193.

37.   Deblois, G. and V. Giguère, *Functional and physiological genomics of estrogen-related*
receptors (errs) in health and disease. Biochim Biophys Acta, 2011. **1812**(8): p. 1032.

38.   Ranhotra, H.S., *The estrogen-related receptors: Orphans orchestrating myriad functions.* J
Recept Signal Transduct Res, 2012. **32**(2): p. 47.

39.   Charest-Marcotte, A., et al., *The homeobox protein prox1 is a negative modulator of*
err{alpha}/pgc-1{alpha} bioenergetic functions. Genes Dev, 2010. **24**(6): p. 537.

40.   Eichner, L.J. and V. Giguère, *Estrogen related receptors (errs): A new dawn in transcriptional*
control of mitochondrial gene networks. Mitochondrion, 2011. **11**(4): p. 544.

41.   Koh, E., et al., *Mpc1 is essential for pgc-1α-induced mitochondrial respiration and biogenesis.*
Biochem J, 2018. **475**(10): p. 1687.

42. Divangahi, M., et al., *Trained immunity, tolerance, priming and differentiation: Distinct*
immunological processes. Nat Immunol, 2021. **22**(1): p. 2.
43. Xue, C., et al., *Mitochondrial pyruvate carrier 1: A novel prognostic biomarker that predicts*
favourable patient survival in cancer. Cancer Cell Int, 2021. **21**(1): p. 288.
44. Bronte, V. and M.J. Pittet, *The spleen in local and systemic regulation of immunity.* Immunity,
2013. **39**(5): p. 806.
45. Starkey, M., et al., *Identification of anti-virulence compounds that disrupt quorum-sensing*
regulated acute and persistent pathogenicity. PLoS pathogens, 2014. **10**(8): p. e1004321.
46. Dulcey, C.E., et al., *The end of an old hypothesis: The pseudomonas signaling molecules 4-*
hydroxy-2-alkylquinolines derive from fatty acids, not 3-ketofatty acids. Chem Biol, 2013.
20(12): p. 1481.
47. Fetzner, S. and S.L. Drees, *Old molecules, new biochemistry.* Chem Biol, 2013. **20**(12): p. 1438.
48. Park, S., et al., *Inhibition of erralpha prevents mitochondrial pyruvate uptake exposing nadph-*
generating pathways as targetable vulnerabilities in breast cancer. Cell Rep, 2019. **27**(12): p.
3587.
49. Caslin, H.L., et al., *Lactate is a metabolic mediator that shapes immune cell fate and function.*
Front Physiol, 2021. **12**: p. 688485.
50. Lee, T.Y., *Lactate: A multifunctional signaling molecule.* Yeungnam Univ J Med, 2021. **38**(3): p.
183.
51. Li, X., et al., *Lactate metabolism in human health and disease.* Signal Transduction and
Targeted Therapy, 2022. **7**(1): p. 305.
52. Wei, W., et al., *Ligand activation of errα by cholesterol mediates statin and bisphosphonate*
effects. Cell metabolism, 2016. **23**(3): p. 479.
53. Chisamore, M.J., et al., *Identification of small molecule estrogen-related receptor α–specific*
antagonists and homology modeling to predict the molecular determinants as the basis for
selectivity over errβ and erry. Drug development research, 2008. **69**(4): p. 203.
54. Mayer, M.L., et al., *The pseudomonas aeruginosa autoinducer 3o-c12 homoserine lactone*
provokes hyperinflammatory responses from cystic fibrosis airway epithelial cells. PloS one,
2011. **6**(1): p. e16246.
55. Kushwaha, A., V. Kumar, and V. Agarwal, *Pseudomonas quinolone signal induces organelle*
stress and dysregulates inflammation in human macrophages. Biochim Biophys Acta Gen
Subj, 2023. **1867**(2): p. 130269.
56. Pietrocola, F., et al., *Acetyl coenzyme a: A central metabolite and second messenger.* Cell
Metab, 2015. **21**(6): p. 805.
57. Chandran, A., et al., *Mycobacterium tuberculosis infection induces hdac1-mediated*
suppression of il-12b gene expression in macrophages. Front Cell Infect Microbiol, 2015. **5**: p.
90.
58. Shi, L. and B.P. Tu, *Acetyl-coa and the regulation of metabolism: Mechanisms and*
consequences. Curr Opin Cell Biol, 2015. **33**: p. 125.
59. Lauterbach, M.A., et al., *Toll-like receptor signaling rewires macrophage metabolism and*
promotes histone acetylation via atp-citrate lyase. Immunity, 2019. **51**(6): p. 997.
60. Saeed, S., et al., *Epigenetic programming of monocyte-to-macrophage differentiation and*
trained innate immunity. Science, 2014. **345**(6204): p. 1251086.
61. Liu, T.F., et al., *Nad+-dependent sirtuin 1 and 6 proteins coordinate a switch from glucose to*
fatty acid oxidation during the acute inflammatory response. J Biol Chem, 2012. **287**(31): p.
25758.
62. Ishida, K., et al., *Chlamydia pneumoniae effector chlamydial outer protein n sequesters*
fructose bisphosphate aldolase a, providing a benefit to bacterial growth. BMC Microbiol,
2014. **14**: p. 330.

63.   Kunze, M., et al., *Metabolic adaption of legionella pneumophila during intracellular growth in*
898   *acanthamoeba castellanii.* Int J Med Microbiol, 2021. **311**(4): p. 151504.
899   64.   Shi, L., et al., *Infection with mycobacterium tuberculosis induces the warburg effect in mouse*
900   *lungs.* Sci Rep, 2015. **5**: p. 18176.
901   65.   Perez-Morales, D. and V.H. Bustamante, *Cross-kingdom metabolic manipulation promotes*
902   *salmonella replication inside macrophages.* Nat Commun, 2021. **12**(1): p. 1862.
903   66.   Dhamgaye, S., Y. Qu, and A.Y. Peleg, *Polymicrobial infections involving clinically relevant gram-*
904   *negative bacteria and fungi.* Cellular microbiology, 2016. **18**(12): p. 1716.
905   67.   Vega, R.B., J.M. Huss, and D.P. Kelly, *The coactivator pgc-1 cooperates with peroxisome*
906   *proliferator-activated receptor alpha in transcriptional control of nuclear genes encoding*
907   *mitochondrial fatty acid oxidation enzymes.* Molecular and cellular biology, 2000. **20**(5): p.
908   1868.
909   68.   Bekkering, S., et al., *Trained immunity: Reprogramming innate immunity in health and*
910   *disease.* Annual Review of Immunology, 2021. **39**(1): p. 667.
911   69.   Ochando, J., et al., *Trained immunity — basic concepts and contributions to*
912   *immunopathology.* Nature Reviews Nephrology, 2023. **19**(1): p. 23.
913   70.   Au - Van den Bossche, J., J. Au - Baardman, and M.P.J. Au - de Winther, *Metabolic*
914   *characterization of polarized m1 and m2 bone marrow-derived macrophages using real-time*
915   *extracellular flux analysis.* JoVE, 2015(105): p. e53424.
916   71.   Rahme, L.G., et al., *Common virulence factors for bacterial pathogenicity in plants and*
917   *animals.* Science, 1995. **268**(5219): p. 1899.
918   72.   Maura, D., A. Bandyopadhaya, and L.G. Rahme, *Animal models for pseudomonas aeruginosa*
919   *quorum sensing studies.* Methods Mol Biol, 2018. **1673**: p. 227.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure S1

Figure S2

Figure S3