1

Experimental evolution in maize with replicated divergent selection identifies plant-height associated SNPs

Mila Tost[1,2], Cathy Westhues[1,2], Ginnie Morrison[3], Dietrich Kaufmann[1] and Timothy Beissinger[1,2,*]

[1]Department of Crop Science, Division of Plant Breeding Methodology, University of Goettingen, Carl-Sprengel-Weg 1, Goettingen, 37075, Germany

[2]Center for Integrated Breeding Research, University of Goettingen, Carl-Sprengel-Weg 1, Goettingen, 37075, Germany

[3]University of Missouri, Division of Biological Sciences, 105 Tucker Hall, Columbia, MO, 65211, USA

[*]**Corresponding author:** Timothy Beissinger (timbeissinger@gmail.com)

Running title: Replicated experimental evolution maize

Key words: Experimental evolution, Population genetics, maize, selection signature mapping, plant height

Abstract

Experimental evolution studies are common in agricultural research, where they are often deemed "long term selection". These are often used to perform selection mapping, which involves identifying markers which were putatively under selection based on finding signals of selection left in the genome. A challenge of previous selection mapping studies, especially in agricultural research, has been the specification of robust significance thresholds. This is in large part because long term selection studies are rarely replicated. Usually, significance thresholds in long term selection experiments are based on outliers from an empirical distribution or based on drift simulations. Both of these approaches are prone to missing true positives or including false positives. Under laboratory conditions with model species, replicated selection has been shown to be a powerful tool, especially for the specification of significance thresholds. In this study, we conducted divergent, replicated selection for short and tall plant-height in a random mating maize population under real field conditions. Selection of the 5% tallest or shortest plants was conducted for three generations. Significance thresholds were specified using the false discovery rate for selection (FDRfS) based on a single marker-based statistic leveraging replicated selection ($F_{ST}Sum$). Overall, we found 44 significant SNPs. Thirty-four were located within a region of 1.02 Mb comprising the coding region of the plant-height genes *Dwarf1* and *iAA8*. We applied a window-based analysis and haplotype block analysis to further dissect the pattern of selection in significant regions of the genome. We observed patterns of strong selection in the subpopulations selected for short plant height.

Introduction

The purpose of experimental evolution studies is to understand how populations evolve in response to a selective pressure (Janick, 2010a, 2010b). In agricultural research, they are commonly employed and often deemed "long term selection" studies (Dudley and Lambert, 2010; Moose, Dudley and Rocheford, 2004, Beissinger *et al.,* 2014). For instance, the Illinois long-term selection experiment for maize protein content began in 1896 at the University of Illinois and has been conducted ever since (Dudley and Lambert, 2010; Moose et al. 2004). Experimental evolution studies are also common in model organisms like mice (Firman and Simmons, 2009; Firman *et al.*, 2015), *Drosophila melanogaster* (Teotónio *et al.*, 2009; Burke *et al.*, 2010; Turner *et al.*, 2011; Phillips *et al.*, 2016)*, E. coli* (Herring *et al.*, 2006; Barrick *et al.*, 2009)*, Mimulus guttatus* (monkey flowers) (Kelly *et al.*, 2006; Angert, Bradshaw Jr and Schemske, 2008), and yeast (Payen *et al.*, 2016; Linder *et al.*, 2021). Early experimental evolution studies focused on the evolution of phenotypes alone, but now sequencing technologies allow researchers to study the effect of selection at the DNA level (Burke *et al.,* 2010; Parts *et al.*, 2011; Orozco-terwengel *et al.,* 2012; Remolina *et al.,* 2012).

Many experimental evolution studies in model organisms work with replicates (Teotónio and Rose, 2000; Teotónio *et al.*, 2009; Phillips *et al.*, 2016). These studies are often referred to as Evolve & Resequence (E&R) experiments (Turner *et al.*, 2011). E&R experiments commonly use pooled-based sequencing of the different subpopulations (Turner *et al.*, 2011, Orozco-terWengel *et al.,* 2012; Long *et al.*, 2015; Barghi *et al.*, 2017). In agricultural species, experimental evolution studies that leverage replicated selection populations are exceedingly rare (Dudley and Lambert, 2010, Beissinger *et al.*, 2014; Kumar *et al.*, 2021). On the other hand,

replication plays a very important role in E&R studies because it can help to identify putatively

selected sites and differentiate those from sites caused by drift (Baldwin-Brown, Long and

Thornton, 2014; Kofler and Schlötterer, 2014; Turner and Miller, 2012; Linder *et al.*, 2021). This

assumes that selection is repeatable, while randomly acting forces like drift are not (Teotónio

and Rose, 2000; Turner and Miller, 2012; Burke, Liti and Long, 2014).

Genetic drift is a complicating factor in experimental evolution studies because it can be difficult

to differentiate from selection (Krimbas and Tsakas, 1971; Turner *et al.*, 2011; Beissinger *et al.*,

2014). This is a recurring challenge in unreplicated selection mapping studies, because without

replicating the evolutionary process, no amount of sampling can estimate the variability

associated with genetic drift (Krimbas and Tsakas, 1971; Turner *et al.*, 2011; Hirsch *et al.*, 2014;

Barghi, Hermisson and Schlötterer 2020, Kumar *et al.*, 2021). The variability introduced by

genetic drift depends on population size and the number of generations of selection (Johannson

et al., 2010, Turner and Miller, 2012; Kessner and Novembre, 2015). Replicated selection

populations can help differentiate neutral loci subject to drift alone, from putatively selected

loci (Turner and Miller, 2012; Kofler and Schlötterer, 2014; Barghi *et al.,* 2019; Linder *et al.,*

2021). Replicates can be analyzed to calculate a null expectation and to derive an approximation

of the number of false positive observations in the experiment.

In previous experimental evolution studies on crops, significance thresholds for identifying

signals of selection were frequently derived based on outlier quantiles of the empirical

distribution (Beissinger *et al.*, 2014; Hirsch *et al.*, 2014; Lorenz *et al.*, 2015; Wang *et al.*, 2018;

Ayalew *et al.*, 2020). This assumes that selection is a locus-specific force; putatively selected loci

should be found at the tail of the empirical distribution (Akey *et al.*, 2002; Akey, 2009). Drift is

assumed to act uniformly genome-wide, so drifting loci dictate the bulk of the empirical

distribution (Akey *et al.*, 2002; Akey, 2009). However, it has been recognized for a very long time

that outlier-based thresholds are arbitrary (Akey *et al.*, 2002; Akey, 2009).

Another common approach for specifying significance thresholds for selection is to conduct drift

simulations (Kumar *et al.*, 2021; Turner *et al.*, 2011). This approach is an improvement over

arbitrary outlier thresholds, but it is impossible to know with certainty that simulations of

genetic drift fully mimic the complex demographic scenarios experienced by a real-world

population, including an experimental one.

For this study, we performed replicated divergent selection for plant height on a random mating

maize population for three generations. Plant height is an ideal model trait because it is highly

heritable and highly polygenic (Pfeiffer *et al*., 2014, Mazaheri *et al.,* 2019). We replicated

selection on four subpopulations, two of which were selected for tall plant height and two

which were selected for short plant height. Each subpopulation was maintained with a

population size of approximately 5,000 individuals, making this the largest replicated

experimental evolution study conducted on a crop species to date. Every generation, we

selected the 5% tallest or shortest plants (~250 plants). The truncation threshold of 5% tallest or

shortest plants was calculated based on 96 randomly sampled plants in from each

subpopulation (in total 384 plants). The 384 randomly sampled plants from the fourth

generation were also used for genotyping-by-sequencing (GBS) to conduct selection mapping.

We calculated F_{ST} for each identified marker in the genome (Weir and Cockerham, 1984). Then,

we calculated a version of F_{ST} leveraging replicated selection. Finally, we generated significance

thresholds based on Turner and Miller's (2012) false discovery rate for selection (FDRfS). Finally,

we compared these replication-based significance thresholds to those generated from the

empirical distribution of F_{ST} and from drift simulations.

Regardless of the method used to specify significance thresholds, regions corresponding to

selection can be quite large and contain neutral linked loci (Barghi and Schlötterer, 2019;

Franssen, Barton and Schlötterer, 2016; Tobler *et al.*, 2014). Single marker-based statistics like

F_{ST} are prone to identify neutral loci that have hitchhiked along with those that are under

selection (Franssen, Barton and Schlötterer, 2016; Tobler *et al.*, 2014). Several experimental

evolution studies have additionally investigated reconstructed haplotype blocks to disentangle

the complex pattern of linkage (Franssen, Barton and Schlötterer, 2016; Michalak *et al.*, 2018).

These investigations showed that often a large number of putatively significant sites

corresponds to only a small number of significant haplotype blocks (Franssen, Barton and

Schlötterer, 2016; Michalak *et al.*, 2018). Therefore, we applied a cubic smoothing spline

approach to identify window boundaries for selection regions (Beissinger *et al.*, 2015). This

procedure may also reduce the inflation of observations due to linked loci (Barghi and

Schlötterer, 2019). Additionally, we phased our data and investigated the haplotype blocks

under selection.

Material and methods

Plant materials and experimental design

The Shoepeg population is a maize Southern dent landrace population (Doebley *et al.*, 1988),

which was collected as open-pollinating variety in 1960 in Missouri, USA. We ordered Shoepeg

material from the National Plant Germplasm System as accession PI 269743 (U.S. National Plant

Germplasm System, 2021). Seeds were greenhouse-propagated to generate our starting

population for selection. The formation of the Shoepeg experimental evolution population and

results preceding the first generation of selection are described in Gyawali *et al.* (2019). To

summarize, in 2016 the Shoepeg population was divided into four subpopulations, each

consisting of approximately 5000 plants which were allowed to open-pollinate. In each

subpopulation, plant height was measured (Figure 1). In total, 384 (96 x 4) individual plants out

of approximately 20,000 (5000 x 4) plants were measured across all four subpopulations (Figure

1). Based on these plant height measurements, a truncation threshold corresponding to

approximately the 5% shortest or tallest plants was calculated for each subpopulation (Figure 1)

(Gyawali *et al.*, 2019). Two subpopulations were selected for short plants, and two were

selected for tall plants. The ~5% shortest or tallest plants from each subpopulation were

harvested as females to generate the next generation (Figure 1) (Gyawali *et al.*, 2019).

This procedure was repeated in 2017 and 2018 (Figure 1). Ultimately, we generated four

experimentally evolved subpopulations, two comprised of short-selected plants and two

comprised of tall-selected plants (Figure 1). The populations were grown again in 2020 and plant

height was measured for 96 randomly-chosen representative plants in each subpopulation (in

total 384 plants). Additionally, tissue was collected from the same 384 plants for genotyping

(Figure 1).

Figure 1: Simplified scheme for the development of subpopulations selected in divergent directions with their replicates for the three generations of selection. The subpopulations short 1 and 2 were selected for short plant height and the subpopulations tall 1 and 2 were selected for tall plant height.

Every year after the first, the subpopulations were grown in isolated fields at least 200 m apart

and away from any other maize to avoid contamination via cross-pollination (Nieh *et al.*, 2014;

Gyawali *et al.*, 2019). In 2016, 2017 and 2018, the subpopulations were grown in Columbia,

Missouri, USA (Gyawali *et al.*, 2019). In 2020 the subpopulations were grown in Göttingen,

Germany. In Missouri, each field had an approximate size of 0.1 ha (Gyawali *et al.*, 2019). The

row-spacing was 91 cm and the space between the plants was 15 cm (Gyawali *et al.*, 2019). In

Göttingen, the field size for three of the fields was 0.25 ha. One field was only 0.125 ha due to

fewer seeds caused by a lower-than-expected harvest from one of the subpopulations in 2018.

The distance between the seeds in Göttingen was approximately 33.3 cm and the row-spacing

was approximately 1 m. The fields in Goettingen experienced noteworthy but non-devastating

crow damage.

Plant height measurements were taken after flowering, from the ground up to the ligule of the

flag leaf. Phenotypic data was analyzed using R version 4.0.3 (R Core Team, 2021). For

phenotypic data analysis, data from 2016, 2017, 2018 and 2020 was used. The realized

heritability from the breeder's equation (h^2_{bs}) was calculated as $h^2_{bs} = \frac{R}{S}$ (Lush, 1937), where S is

the selection differential and R is the response to selection. The selection differential was

calculated as the difference between the mean of the population and the mean of the selected

portion (Lush, 1937). The response to selection was calculated as difference between the mean

of the previous generation and the mean of the next generation (Lush, 1937). This was done

between all subsequent generations.

Flowering time was measured because in maize it tends to be correlated with plant height

(Peiffer *et al.*, 2014; Mazaheri *et al.,* 2019). This could lead to indirect selection for flowering

time, thereby effecting the ability of plants to mate randomly. The possibility of this impacting

the effective population size is addressed below.

Genotyping, quality control and filtering

In 2020, the 96 randomly chosen plants from each subpopulation in generation four (384

individual plants in total) were genotyped using a genotyping-by-sequencing (GBS) approach.

Leaf tissue from those 384 plants was freeze-dried and shipped to the University of Wisconsin

Biotechnology center, Madison, WI, USA for DNA extraction, library preparation, and

sequencing via one lane on a NovaSeq6000 machine. Each subpopulation was sequenced on its

own sequencing run. Total sequencing yield from each run ranged from 125,000,000 to

300,000,000 fragments. Fragments had read lengths between 73 and 151 base pairs. GBS was

conducted using paired-end sequencing with the restriction enzyme *Ape*KI (Elshire *et al.*, 2011).

For SNP variant calling the GB-eaSy pipeline was used (Wickland *et al.*, 2017, Tange, 2011).

Reads were aligned to the 5th version of the B73 reference genome (Maize GDB, 2021) with the

Burrows Wheeler Aligner (Li, 2013). In total, 15,390,958 SNPs were called by the GB-eaSy

pipeline (Wickland *et al.*, 2017). In the GB-eaSy pipeline, the SNPs were filtered for a minimum

mapping and base quality of 30 with the BCFtools software (Danecek *et al.*, 2021). After filtering,

8,302,778 SNPs were kept. The mapping quality ranged from 30 to 60, and was on average

57.44. The raw read depth ranged from 161 to 4,999, and was on average 1,705 reads. The

average distance between adjacent markers was 257.67 bp.

Subsequent SNP filtering was conducted in R version 4.0.3 (R Core Team, 2021). Six individual

samples with a marker coverage below 0.1 were removed from the analysis. The 10% quartile of

marker coverage 0.6042, and 90% quartile was 0.7014 (supplementary figure S1). Markers with

an average sample read depth below 1 or above 10 were removed. This filtering threshold was

determined based on the distribution of read depth across all samples and markers

(supplementary figure S2). Furthermore, we required that every SNP marker was observed at

least 40 times in each subpopulation. Monomorphic and multi-allelic (>2) markers were

removed from the data set. The final data set contained 4,029,092 SNP markers, which were

used for the downstream analysis.

Allele frequency estimation

The allele frequency at each SNP was calculated in R according to its maximum-likelihood

estimate, which is computed by dividing the number of observations of a particular allele by the

number of observations in total (Beissinger *et al.*, 2014). The script is available at

(https://github.com/milaleonie/ExpEvo_with_replicated_selection).

Estimating effective population size

Effective population size was calculated using the known demographic parameters of the

populations and based off the formula, $N_e = \frac{4 * N_f * N_m}{N_f + N_m}$, where N_m is the number of male and N_f

the number of female individuals (Crow and Kimura, 1970). In every generation the ~5%

shortest or tallest plants were harvested, amounting to ~250 plants. Under the assumption of

complete random mating, all 5,000 plants could contribute as male parents for the next

generation. In field trials, the assumption of random mating of all 5,000 plants is violated

because of the effects of assortative mating (Allard, 1999). The number of male parents

contributing to the next generation is reduced due to varying flowering dates and limited spatial

pollen dispersion (Allard, 1999). Therefore, we evaluated the flowering dates of the 96 randomly

chosen plants from each subpopulation to approximate the number of maximal simultaneously

flowering tassels and silks in the entire population. We looked for the quantile in which the

majority of silks is mature and receptive, because it was assumed that silks remain receptive up

to five days after silk emergence (Nieh *et al.*, 2014). We calculated the number of flowering

tassels during this time interval and projected it onto the entire subpopulation. This procedure

is explained in more detail in supplementary figure S3.

Calculation of linkage disequilibrium (LD) decay

The extent of linkage disequilibrium (LD) was estimated based on 100,000 SNP markers with

TASSEL v5 using the squared correlation between markers, r^2, over a window size of 2,000 bp

(Bradbury *et al.*, 2007). We filtered the individuals out with a marker coverage below 0.6. For

this analysis, we required that every SNP marker was observed at least 80 times in each

subpopulation. This filtering resulted in 1,243,604 SNP markers. Additionally, we thinned the

data set by randomly sampling 10,000 markers per chromosome. LD decay was modeled with a

nonlinear regression model as expected value r^2 under drift-recombination equilibrium as

$E(r^2) = \frac{1}{(1+C)}$ and $C = 4Nc$ with N as the number of individuals at each site and C as the

recombination coefficient between sites (Remington *et al.*, 2001). We also created a pairwise LD

heatmap with the R package LDheatmap (Shin *et al.*, 2006).

Scan for selection signatures

Our scan for selection was based on a statistic derived from F_{ST}. F_{ST} was calculated according to

$$F_{ST} = \frac{s^2}{\mu(p) * \mu(1-p) + \frac{s^2}{r}}$$, where s^2 is the sample variance of the allele frequency between

populations, $\mu(p)$ is the mean allele frequency between the subpopulations and r is the number

of populations (Weir and Cockerham, 1984). The F_{ST} statistic was calculated between the

divergently selected subpopulations (Short 1 vs Tall 1; Short 2 vs Tall 2). Additionally, the F_{ST}

statistic was calculated between the subpopulations selected in the same direction (Short 1 vs

Short 2; Tall 1 vs Tall 2).

Calculation of $F_{ST}Sum$

265   The sum of F_{ST} was calculated between the two non-redundant comparisons of divergently

266   selected subpopulations (Short 1 vs Tall 1; Short 2 vs Tall 2) and subpopulation selected in the

267   same direction (Short 1 vs Short 2; Tall 1 vs Tall 2). Assessing divergence among replicates can

268   help in the distinction between putatively selected loci and false positives (Remolina *et al.,*

269   2012; Kofler and Schlötterer, 2014)

Window-based analysis

We implemented a window-based analysis to assess linked selection. A cubic smoothing spline

was applied to the single-marker based $F_{ST}Sum$ values*.* From the fitted spline, inflection points

were calculated and used as window boundaries for newly defined regions. This analysis was

performed with the R package GenWin (Beissinger *et al.,* 2015). Briefly, the GenWin package

returns window boundaries that we used to define regions with linked markers that may be

analyzed together (Beissinger *et al.*, 2015). However, those regions are of variable sizes, so we

used the W statistic (*Wstat*), as defined by Beissinger *et al.* (2015), to correct for this. Wstat is

defined according to $Wstat = \frac{(X-\mu)}{\sqrt{\frac{s^2}{n}}}$, where X is the mean $F_{ST}Sum$ of a window, μ is the mean

$F_{ST}Sum$ of the entire data set, s^2 is the sample variance of $F_{ST}Sum$ values across the entire data

set and *n* is the number of markers. *Wstat* can therefore be used to compare regions, despite

each consisting of a different number of markers (Beissinger *et al.*, 2015).

Calculation of significance thresholds

Significance thresholds for selection were calculated in three ways: 1) based on the empirical

distribution; 2) based on drift simulations; and 3) based on the FDRfS (Turner and Miller, 2012).

1) The significance thresholds based on the empirical distribution were calculated by taking

288   the 99.9[th] and the 99.99[th] percentile of the entire distribution of the $F_{ST}Sum$ values (Akey,

289   2009; Beissinger *et al.*, 2014; Hirsch *et al.*, 2014; Ma *et al.*, 2015; Cesconeto *et al.*, 2017;

290   Wang *et al.*, 2018; Ayalew *et al.*, 2020).

291   2) To calculate the significance thresholds based on drift simulations, we first simulated

292   drift 1,000,000 times based on the demographic parameters of the replicated

293   populations. Next, we calculated $F_{ST}Sum$ for the simulated allele frequencies. The

294   99.9999[th] percentile of all values generated by the drift simulations was used as the

295   significance thresholds (Kumar *et al.*, 2021). The drift simulations are described in more

296   detail below.

3) High $F_{ST}Sum$ values calculated between divergently selected subpopulations are

observed when a site shows the same pattern of differentiation in both comparisons.

Sites with high $F_{ST}Sum$ calculated between subpopulations selected in the same direction

are false positive observations for selection for plant height. By dividing the number of

sites diverged between the subpopulations selected in the same direction by the number

of sites diverged between the divergently selected subpopulation for a certain $F_{ST}Sum$,

we obtained the false discovery rate for selection (FDRfS) for $F_{ST}Sum$. The FDRfS was

calculated for all possible $F_{ST}Sum$ values (Turner and Miller, 2012). We identified the

$F_{ST}Sum$ which gave an FDRfS <5%. The corresponding $F_{ST}Sum$ value was used as our

significance threshold. The FDRfS was also used with *Wstat* to generate window-based

values. The significance threshold was chosen as the *Wstat* value that excluded all

comparisons between subpopulations selected in the same direction.

Simulations of drift

Drift was simulated with an R function

(https://github.com/milaleonie/ExpEvo_with_replicated_selection), which was implemented in

R version 4.0.3 (R Core Team, 2021). The function was implemented using a wrapper-script

based on that previously published by Kumar *et al.* (2021), which enabled the implementation of

the drift simulator over a large set of markers (Kumar *et al.*, 2021). We conducted 1,000,000

simulations. For each simulation, initial allele frequencies were set based on the allele frequency

spectrum observed in generation 0 (Gyawali *et al.*, 2019). Drift was simulated with a population

size of 5,000 individuals with 250 female and 5,000 male individuals for three generations. We

assumed that every ear contributed ~500 kernels. Every kernel could have been pollinated by

one of the male parents, which resulted in much higher harvest than 5,000 kernels. Therefore,

5,000 kernels were randomly drawn from the entire harvest to represent the seeds planted for

the next generation.

Haplotype estimation and haplotype block calculation

The data were phased using fastPHASE version 1.4.8 (Sheet and Stephens, 2006) for further

dissection. We phased the genomic data of the region from 9.437 to 10.457 Mb on chromosome

3 with 10 iterations of the expectation-maximization (EM) algorithm (Sheet and Stephens,

2006). The model was supplied with labels indicating the different subpopulations. Haplotype

blocks were calculated with the R package HaploBlocker (Pook *et al.*, 2019). Additionally, we

conducted the same haplotype block analysis in 39 randomly selected regions across the entire

genome, with the same physical length and marker density like the region from 9.437 to 10.457

Mb on chromosome 3. We also investigated the region from 9.437 to 10.457 Mb on

chromosome 3 in generation 0 in the haplotype block analysis.

Results

Phenotypic evaluation

We observed notable and significant plant height differences between the subpopulations

selected in opposite directions (Figure 2). After three generations of selection, the mean plant

height difference between subpopulations selected in the opposite direction was 66.55 cm (P ~

0***). The mean plant height decreased in the subpopulations selected for short plant height by

22.92 cm (P ~ 0***) and increased in the subpopulations selected for tall plant height by 30.61

cm (P ~ 0***). Throughout the experiment, the response to selection was always negative in the

subpopulations selected for short plants and always positive in the subpopulations selected for

tall plants (Figure 2).

346
347   **Figure 2:** Mean plant height in each generation of selection. Error bars indicate a confidence interval for our
348   estimation of means.
349

Realized heritability was computed by using the breeder's equation (h^2_{bs}) (Lush, 1937), and was

estimated to be 0.3. Since each population was evaluated in a different environment (i.e. the

year and location where it was selected), this application of the breeder's equation should be

considered only approximate. Plant height and flowering time were correlated in the

subpopulations short 1, short 2, tall 1, and tall 2 with correlation coefficients of 0.2183 (P ~

0.0376**), -0.0384 (P ~ 0.7165), 0.2709 (P ~ 0.0107**) and 0.4572 (P ~ 0***), respectively.

Effective population size and inbreeding

The effective population size under the assumption of complete random mating was 952

individuals. Due to differences in flowering time between plants, the assumption of random

mating is violated. Based on observed flowering times, we estimated that ~ 59.38% and ~

82.29% in the short and ~ 56.25% and ~ 38.54% in the tall subpopulations had the potential to

pollinate one another. When we re-estimated effective population size based on this number of

male plants, the effective population size was estimated ~ 922 and ~ 943 individuals in the short

subpopulations and ~ 918 and ~ 885 individuals in the tall subpopulations.

Test for selection

To visualize the distribution of F_{ST} values observed between subpopulations selected in the

same and opposite directions, we plotted the F_{ST} values calculated between the different

comparisons against each other (Figure 3). This type of plot, but based on allele frequency

differences, was previously named a "Sauron plot" by Turner and Miller (2012). We created this

plot with F_{ST}, which resulted in a "Quarter-Sauron" plot, because F_{ST} ranges from 0 to 1, whereas

allele frequency differences range from -1 to 1. A "Sauron" plot based on the allele frequency

differences, as was originally implemented by Turner and Miller (2012) is provided as

supplemental file S4. This plot depicts how the FDRfS was computed and provides a visualization

of the scope of drift and selection.

In the plot, blue points correspond to the sum of F_{ST} values calculated between the

subpopulations selected in the same direction for each marker, while red points correspond to

the sum of F_{ST} values calculated between the subpopulations selected in opposite directions for

each marker (Figure 3). Therefore, the FDRfS can be calculated for every area of the plot, which

corresponds to a certain $F_{ST}Sum$ value (Figure 3). The region of the plot where the FDRfS is <5%

is indicated by the semi-transparent red region (Figure 3). In this region, 44 SNPs with $F_{ST}Sum$

>0.662 were identified for subpopulations selected in opposite directions, while only two SNPs

displayed a $F_{ST}Sum$ >0.662 between subpopulations selected in the same direction (Figure 3).

This leads to an exact FDRfS of 4.5%. The FDRfS for every possible $F_{ST}Sum$ and the number of

markers observed at this value are available in supplementary figure S5. $F_{ST}Sum$ was only

calculated for comparisons without missing data, to make it less sensitive to outliers.

Figure 3: "Quarter-Sauron" plot of genetic differentiation for $F_{ST}Sum$ observed between the subpopulations
selected in the same direction (blue) and in opposite directions (red). Each dot represents one SNP. The semi-
transparent red region corresponds to a false discovery rate for selection (FDRfS) <5%.

Comparison of significance thresholds and candidate gene identification

We also evaluated significance thresholds based on outlier thresholds and drift simulations.

Thresholds based on the 99.9^{th} and the 99.99^{th} percentile of the empirical distribution were

exceeded, respectively, by 1,422 and 143 $F_{ST}Sum$ values observed at single SNP markers (Figure

4A). In total, 93 and 10 regions exceeded the thresholds based on the 99.9^{th} and the 99.99^{th}

percentile of the empirical distribution of *Wstat* values (Figure 4B). Significance thresholds

based on drift simulations were exceeded by 9,998 $F_{ST}Sum$ values observed at single SNP

markers (Figure 4A). Significance thresholds based on the FDRfS were exceeded by the $F_{ST}Sum$

observed at 44 single SNP markers (Figure 4A) and by *Wstat* observed across 3 different regions

(Figure 4B). $F_{ST}Sum$ and *Wstat* values calculated between the subpopulations selected in the

same direction are depicted in Figure 4C and Figure 4D.

Figure 4: Differentiation along the entire genome expressed by F_{ST} at each SNP marker between the subpopulations
selected in opposite directions (A) and the same direction (B) and expressed by *Wstat* values across regions (C, D)
with the significance thresholds based on the 99.9[th] percentile of the empirical distribution (light purple), based on
the 99.99[th] percentile of the empirical distribution (dark purple), drift simulations (pink), and by the false discovery
rate for selection (FDRfS) (yellow).

Candidate plant height SNPs

All three regions identified based on *Wstat* overlapped with coding regions in the B73 v5

annotated gene models (Maize GDB, 2021). For two of the regions, gene models were

associated with annotated gene IDs (Table 1). For the gene *impb24 - importin beta24*, no

information was found based on a literature review (Table 1). The gene *pip2e* belongs to the

plasma membrane intrinsic protein subfamily PIP2, which has been described as associated with

salt stress (Fox *et al.*, 2021). The gene model *Zm00001eb122360*, which coding region was

located within the identified region on chromosome 3, did not have an associated candidate

gene (Table 1).

Twenty-seven out 44 SNPs identified by $F_{ST}Sum$ were located within or 85 kb from B73 v5 (Maize

GDB, 2021) annotated gene models. Many markers correspond to the same gene models and

appear to be located in the same region (Table 1). On chromosome 3 we found 34 significant

markers, which were located in a region from 9.437 to 10.457 Mb (Table 1). All 34 markers in

that region show very similar allele frequency changes and resulting differentiation statistic

(Table 1). The plant height candidate genes *Dwarf1/GA 3-oxidase 1* and the *iAA8 - Aux/iAA-*

transcription factor 8 are located in this region.

The plant height candidate genes *Dwarf1* and the *iAA8* have already been shown to be directly

involved in plant height differences in many studies (Teng *et al.*, 2013; Peiffer *et al.*, 2014; Liu,

Fernie and Yan, 2020; Ramos Báez *et al.*, 2020). Therefore, these candidate genes were used as

ground truth to test the method. Other identified gene models did not show a direct connection

to plant height based on the literature review.

Four markers were located directly within the coding region of the *iAA8* gene (Table 1). Two

markers were found near the *Dwarf1* gene and were located ~ 85 kb upstream from this gene

(Table 1). The window-based region with the highest *Wstat* value was also located on

chromosome 3 within the previously described region ranging from 9.437 to 10.457 Mb (Table

1). This 0.12 Mb region is ~ 138 kb away from *iAA8,* and is ~ 518 kb away from *Dwarf 1* (Table 1).

The region is also 31 kb away from a significant marker identified in the Gyawali *et al.,* (2019)

study (Table 1). Other single SNP markers were also within or close to the coding region of other

annotated gene models, but for these no previous evidence for their impact on plant height was

found based on a literature review (Table 1).

Table 1: Significant SNP markers and regions with their associated gene models and candidate genes.

Chr	Position (bp)	F_{ST}*Sum* value	Region (bp)	*Wstat* value	Nr of SNPs	Maize gene annotation and candidate gene nearby	Distance to gene (bp)
2	19993292	0.7187				nothing	
2	19993690	0.8316				nothing	
2	19993788	0.7849				nothing	
3	8055976	0.7648				**Zm00001eb121570:** IDP7380	within the coding region
3	8055983	0.7909					
3	8055985	0.7388					
3	9435824	1.0224				**Zm00001eb122210:** No candidate gene associated	within the coding region
3	9435914	0.9061					
3	9474493	0.8283				**Zm00001eb122220:** No candidate gene associated	within the coding region
3	9529777	0.8925				nothing	
3	9529813	0.8413				nothing	
3	9529816	0.7980				nothing	
3	9529823	0.8103				nothing	
3	9529845	0.7946				nothing	
3	9529863	0.7903				nothing	
3	9544069	0.7926				**Zm00001eb122230:** No candidate gene associated	within the coding region
3	9544090	0.7879					
3	9544092	0.7849					
3	9545220	0.7307					
3	9548992	0.6977				**Zm00001eb122250:** No candidate gene associated	within the coding region
3	9781288	0.7179				**Zm00001eb122280:** pco120183	within the coding region
3	9781314	0.7361					within the coding region
3	9784253	0.7092				**Zm00001eb122290:** No candidate gene associated	within the coding region
3	9784255	0.6966					
3	9784270	0.7150					
3	9792757	0.6828				nothing	
3	9830677	0.7018				**Zm00001eb122300:** No candidate gene associated	within the coding region
3	9909752	0.6876				**Zm00001eb122350:** No candidate gene associated	within the coding region
3			9911000-9923000	52.24	62	**Zm00001eb122360:** No candidate gene associated	overlaps with the coding region
3	9921856	0.6766					
3	9921871	0.6833					

3	9921927	0.6749					
3	9921995	0.8183					
3	9922080	0.7083					
3	9922083	0.7052					
3	10053938	0.6698				**Zm00001eb122410:** iAA8 - Aux/iAA-transcription factor 8	within the coding region
3	10053961	0.6796					
3	10053964	0.6857					
3	10062671	0.6967					
3	10355793	0.8104				**Zm00001eb122500:** Dwarf1 - dwarf plant1	
3	10355846	0.8589					
3	15011305	0.6630				nothing	
5			30103000-30115000	42.61	22	**Zm00001eb221820:** impb24 - importin beta24	overlaps with the coding region
7	79405796	0.6624				nothing	
7			42687000-42725000	36.55	24	**Zm00001eb306380:** pip2e - plasma membrane intrinsic protein2	overlaps with the coding region
8	93026326	0.7371				nothing	
8	93026386	0.6803				nothing	

Haplotype block analysis

We investigated the region from 9.437 to 10.457 Mb on chromosome 3 in more detail by

dissecting the haplotypes in the population (Figure 5). The haplotype blocks located within and

around the coding region of the gene *Dwarf1* indicate allele fixation in almost all subpopulations

(Figure 5B). Only in subpopulation tall 2 did we observe three different haplotype blocks around

and within the coding region of *Dwarf1* (Figure 5B). But, two of the haplotype blocks contained

only one haplotype each (Figure 5B). The majority of observed haplotypes belonged to a single

block (Figure 5B). Within and approximately 10 kb from the coding region of the gene *iAA8*, we

observed three different haplotype blocks in the two short subpopulations and four in the two

tall subpopulations (Figure 5B). The majority of haplotypes identified in the short

subpopulations were located in one block (Figure 5B). The other two haplotype blocks contained

only one haplotype each (Figure 5B). In the short plant subpopulations, we observed larger

haplotype blocks (Figure 5B) than in the tall subpopulations. The block size was on average

646.4 bp in subpopulation short 1 and 809 bp in subpopulation short 2. In subpopulation tall 1

and tall2, we observed average block sizes of 160.9 bp and 160.97 bp, respectively (Figure 5B).

To supplement our haplotype investigation, we looked at LD across putatively selected regions

in the different subpopulations. A pairwise LD heatmap indicates strong LD across the entire

region from 9.437 to 10.457 Mb in the subpopulations selected for short plant height (Figure

5C). We observed a low LD across the entire region from 9.437 to 10.457 Mb in the

subpopulations selected for tall plant height (Figure 5C).

Figure 5: The region on chromosome 3 (from 9.437 Mb to 10.457 Mb) with the window boundary-based regions and their corresponding *Wstat* (A), the haplotype blocks (B) and pairwise LD heatmap for the different subpopulations(C).

471
472
473
474

Discussion

In contrast to most agriculture long-term selection experiments, our study included replicated selection populations. This allowed us to directly compare the commonly used approaches to significance threshold specification leveraging replications. Our results demonstrate that significance thresholds specified using evolutionary replicates can widely vary from thresholds based on empirical outliers or simulations.

In our study we used $F_{ST}Sum$ values to specify significance thresholds, which can only be calculated when replicates are available. If we had used conventional F_{ST} without leveraging replications, ~2,733 and ~274 would have passed the outlier thresholds of the 99.9[th] and 99.99[th] percentile. This is 2-fold the number of markers compared to thresholds calculated on $F_{ST}Sum$. In some studies, empirical outlier thresholds as low as the 99[th] percentile have been used (Wang *et al.*, 2018), which may comprise many false positives. Overall in this experiment, the number of markers identified using the FDRfS approach corresponds to the 99.99891[st] percentile of the empirical distribution. This percentile of the empirical distribution is much higher than the used percentiles of the empirical distribution used in most other studies (Wang *et al.*, 2018; Ayalew *et al.*, 2020).

FDRfS based on $F_{ST}Sum$ is mathematically equivalent to calculating an FDRfS value based on mean F_{ST} values. However, $F_{ST}Sum$ can be plotted in a much more interpretable way, so it is the statistic we chose to work with.

Despite our best efforts to include all relevant factors in our drift simulations, including

biological factors like mating scheme, setting initial allele frequencies based on the observed

allele frequency distribution, and including technical factors such as the sequencing procedure,

we were not able to identify a significance threshold which excluded false positive observations

(supplementary figure S6). Significance thresholds based on the FDRfS applied to $F_{ST}Sum$, on the

other hand, identified fewer markers than outlier-based or simulation-based significance

thresholds. Because these thresholds are based on replications, markers exceeding them are

expected to correspond to true signals of selection.

Previous studies applied significance thresholds based on the FDRfS on statistics like Chi-Square

(Linder *et al.*, 2021) and allele frequency differences (Turner and Miller, 2012). We adapted the

calculation of the FDRfS to $F_{ST}Sum$, a version of F_{ST} which leverages replications. For comparison,

we also explored calculating FDRfS from allele frequency differences (DiffStat) as was done by

Turner and Miller (2012). The results are available in supplementary figures S5 and S7. We

identified more significant markers using the $F_{ST}Sum$ approach at an FDRfS <5%. This was also

observed by Kofler and Schlötterer (2014).

Most of the 44 markers identified based on $F_{ST}Sum$ correspond to same region under selection.

On chromosome 3, we found 34 significant markers, all of which were located in a region

ranging from 9.437 to 10.457 Mb. All 34 markers show very similar allele frequency changes.

Because linkage leads to selection operating on regions instead of individual markers, we used

the R package GenWin (Beissinger *et al.*, 2015) to calculate *Wstat* values based on a cubic

smoothing spline applied to $F_{ST}Sum.$ This allowed us to define window boundaries of regions

which were putatively under selection. When we compared *Wstat* to $F_{ST}Sum$ values, we

observed a few significant regions, rather than many significant markers within regions.

Furthermore, we observed smaller differences between outlier-based and replication-based

significance thresholds when we calculated the thresholds for *Wstat*. The region identified by

Wstat on chromosome 3 from 9.911 to 9.923 Mb was very close to a marker from previous

study by Gyawali *et al.* (2019).

The plant height candidate gene *iAA8* is located within the putatively selected region from 9.437

to 10.457 Mb on chromosome 3 (Ramos Báez *et al.*, 2020). The plant height gene *Dwarf1* is also

located in this region (Teng *et al.*, 2013; Peiffer *et al.*, 2014; Liu, Fernie and Yan, 2020).

Dwarf1 is involved in the expression of gibberellin 3-oxidase (Teng *et al.*, 2013). High expression

of the gene leads to high levels of bioactive gibberellin and results in taller plants (Teng *et al.*,

2013). When the *Dwarf1* gene is not expressed, plants exhibit the *Dwarf1* phenotype, which is

characterized by dwarfism (Teng *et al.*, 2013). Multiple gene expression studies have previously

reported that this gene has a large effect on plant height (Teng *et al.*, 2013; Liu, Fernie and Yan,

2020). In one experiment, this gene explained 32.3% of the phenotypic variance for plant height

(Teng *et al.*, 2013). The *iAA8* gene regulates auxin biosynthesis, transport, and signaling

(Matthes *et al.*, 2019; Ramos Báez *et al.*, 2020). Auxin is a plant hormone and plays a key role in

plant growth and development (Ramos Báez *et al.*, 2020). The *iAA8* gene was also observed to

regulate maize inflorescence architecture (Galli *et al.*, 2015; Ramos Báez *et al.*, 2020). Both

genes appear to be causal for plant height differences based on the literature (Teng *et al.*, 2013;

Peiffer *et al.*, 2014; Liu, Fernie and Yan, 2020; Ramos Báez *et al.*, 2020). Therefore, these

candidate genes were used as ground truth to test the method.

In the haplotype block analysis, all alleles located in the coding region of *Dwarf1* show limited

genetic variation in all subpopulations except subpopulation tall 2. In subpopulation tall 2, the

remaining variation appears to be limited. Within and around 10 kb of the coding region for the

gene *iAA8*, we observe more genetic variation in the subpopulations selected for tall plants than

in the subpopulations selected for short plants. Overall, we observe larger blocks and

consequently fewer blocks in our candidate gene region in the subpopulations selected for short

plants. The large differences in block sizes and number of blocks may indicate stronger selection

in the subpopulations selected for short plants.

This could theoretically be a result of more inbreeding in the short subpopulations, which we

tested by evaluating haplotype blocks in 39 random regions of the genome. The random regions

were of equal size and marker density to the candidate gene region. We observe on average 31

blocks in short 1, 30 blocks in short 2, 31 blocks in tall 1 and 34 blocks in tall 2. The blocks have

an average block size of 402.8 bp in short 1, 427.36 bp in short 2, 409.01 bp in tall 1 and 404.76

bp in tall 2. This indicates that the tall subpopulations show similar levels of inbreeding to the

short populations. Results from the calculated haplotype blocks from the 39 random regions are

available in supplementary figure S8 and S9. Exemplary haplotype block figures from two

random regions are available in supplementary figures S10 and S11.

Overall, a slow decline in LD was observed with increasing physical distance across all

chromosomes (Supplemental S12). R^2 of 0.1 was observed at a distance of 2 Mb (Supplemental

S12). But when we consider the subpopulations separately in the pairwise LD heatmap, we

observe very different results (Figure 5C). In the pairwise LD heatmap calculated for each

subpopulation, strong LD was observed in the subpopulations short 1 and short 2 across the

entire region ranging from 9.437 to 10.457 Mb (Figure 5C). In the subpopulations tall 1 and tall

2, we observe low LD in the entire region (Figure 5C).

The observation of strong LD in the short plant populations supports the argument that strong

selection may act on the entire candidate gene region in theses populations (Franssen, Barton

and Schlötterer, 2016; Barghi and Schlötterer, 2019). The strength of selection and the

experimental design influence hitchhiking (Baldwin-Brown, Long and Thornton, 2014; Kofler and

Schlötterer, 2014). We applied a strong selective regime in our experiment in comparison to

previous studies (Kofler and Schlötterer, 2014; Long *et al*., 2015). Strong selection leads to more

hitchhiking and a lower mapping resolution (Kofler and Schlötterer, 2014), which we observe in

the short subpopulations. However, strong LD may also persist due to more inbreeding in the

short subpopulations. But we only observe strong LD in the short subpopulations in our

candidate gene region. When we calculated the pairwise LD heatmap for the different

subpopulations in two random regions (supplementary figures S10 and S11), there was no

similarly strong difference in LD between the subpopulations as we observed for our candidate

gene region (Figure 5)

It is not only the strength of selection that can influence the number of variants selected in

experimental evolution studies. The number of selected variants also depends highly on

experimental design factors such as the population size, the duration of the experiment, the

applied selective regime (Kofler and Schlötterer, 2014; Kessner and Novembre, 2015), and the

species (Tobler *et al.*, 2014; Barghi *et al.*, 2017). Divergent selection can increase the power of

the experiment in comparison to a single direction experiment (Kessner and Novembre, 2015).

We observe differences regarding the resolution between the subpopulations selected in

divergent directions. Even though the same strength of selection is applied in the divergently

selected subpopulations, differences between the subpopulations can be observed (Moose,

Dudley and Rocheford, 2004). But, these differences indicate that there is an increase in power

due to divergent selection (Kessner and Novembre, 2015).

Previous simulation studies investigating experimental evolution have demonstrated that

replication is one of the most powerful tools to increase power (Kofler and Schlötterer, 2014). In

fact, Kofler and Schlötterer (2014) observed that replication is more important for detecting

variants than population size (Kofler and Schlötterer, 2014; Kessner and Novembre, 2015).

Regions under selection might become smaller due to more recombination over time, and

subsequently we may find selection on additional regions as generations progress. This

phenomenon is visible in our "Quarter-Sauron" and "Sauron" plots (supplemental file S4)

(supplemental file S4). It was observed that the blue points do not appear completely random.

We observed that 682 markers diverged at F_{ST} of 0.3 between the subpopulations selected for

short plant height. There was no divergence shown in the subpopulations selected for tall plant

height at the same positions (supplemental file S4). At these markers, the frequency of the

reference allele was much higher in the subpopulation short 1. This could have been caused by

selection for specific environmental conditions (Dudley, 2007; Beissinger *et al.,* 2014). The

clustering of markers could have been caused by genetic hitchhiking of linked markers (Tobler *et*

al., 2014; Michalak *et al.*, 2018; Barghi and Schlötterer, 2019).

Conclusion

In conclusion, our research demonstrates that replicated selection is a powerful tool in

agricultural experimental evolution studies. We observed selection across replicates, even

though we only selected for three generations. Moreover, we saw that a significance threshold

based on the FDRfS is more reliable than existing methods for significance threshold

identification. The application of a cubic smoothing spline to single marker-based statistics like

$F_{ST}Sum$ to return window boundaries of regions which are putatively under selection helped

collapse signal from multiple hitchhiking markers. Haplotype block analysis additionally helped

to investigate the regions putatively under selection. We observe very different patterns of

selection in divergently selected subpopulations, which might be due to selection. After more

generations of selection, and a reduction of linkage between putatively selected loci and neutral

loci, we may draw more certain conclusions about selection for the plant height candidate

genes. Furthermore, we might be able to identify significant markers close to our plant height

candidate genes and novel gene loci without any associated gene models.

Availability of data and materials

All the data and statistics about the present study has been included in the current manuscript

in the form of figure and tables. Raw data are publicly available at figshare;

https://figshare.com/projects/Shoepeg_ExpEvo2020/139981. Our analysis code is available

publicly on github: https://github.com/milaleonie/ExpEvo_with_replicated_selection.

Acknowledgements

Abiskar Gyawali contributed substantially to this study but died before submission. Journal

policy prohibits him from being listed as a co-author because he could not approve the final

manuscript. We are grateful for Abi's contributions and miss him tremendously. Kate Guill was

instrumental in conducting early-generation seed increase and field management in Missouri.

Funding

This work was supported by startup funds provided by the University of Goettingen. The first

two generations of selection were funded by the USDA Agricultural Research Service. We

utilized the computational resources of the University of Goettingen's GWDG.

Conflict of interest

The author declares that there is no conflict of interest.

References

Aggrey S, Ankra-Badu GA, and Marks H. Effect of long-term divergent selection on growth characteristics
in Japanese quail. Poultry Science. 2003; 82(4): 538–542.

Akey JM, G Zhang, K Zhang, L Jin, and MD Shriver. Interrogating a High-Density SNP Map for Signatures of
Natural Selection. Genome Research. 2002;12(12): 1805–1814.

Akey JM. Constructing genomic maps of positive selection in humans: Where do we go from here?
Genome Research. 2009;19(5): 711–722.

Allard RW. Principles of plant breeding. 2nd ed. New York: J. Wiley. 1999.

Angert AL, Bradshaw HD Jr., and Schemske DW. Using Experimental Evolution to Investigate Geographic
Range Limits in Monkeyflowers. Evolution. 2008;62(10): 2660–2675.

Ayalew H, ME Sorrells, BF Carver, PS Baenziger, XF Ma. Selection signatures across seven decades of hard
winter wheat breeding in the Great Plains of the United States. The Plant Genome. 2020;13(3).

Baldwin-Brown JG, A Long and KR Thornton. The power to detect quantitative trait loci using
resequenced, experimentally evolved populations of diploid, sexual organisms. Molecular Biology and
Evolution. 2014; 31(4): 1040-55.

Barghi N, R Tobler, V Nolte, C Schlötterer. Drosophila simulans: A Species with Improved Resolution in
Evolve and Resequence Studies. G3 Genes|Genomes|Genetics. 2017; 7(7): 2337–2343.

Barghi N, R Tobler, V Nolte, AM Jakšić, F Mallard, KA Otte, M Dolezal,T Taus, R Kofler,C Schlötterer.
Genetic redundancy fuels polygenic adaptation in Drosophila. PLOS Biology. 2019; 17(2): 1–31.

Barghi N, C Schlötterer. Shifting the paradigm in Evolve and Resequence studies: From analysis of single
nucleotide polymorphisms to selected haplotype blocks. Molecular Ecology. 2019; 28: 521–524.

Barghi N, J Hermisson, C Schlötterer. Polygenic adaptation: a unifying framework to understand positive
selection. Nature reviews. 2020; 21: 769–781.

Barrick JE, DS Yu, SH Yoon, H Jeong, TK Oh, D Schneider, RE Lenski, JF Kim. Genome evolution and
adaptation in a long-term experiment with Escherichia coli. Nature. 2009; 461(7268): 1243–1247.

Beissinger TM, GJM Rosa, SM Kaeppler, D Gianola, N de Leon. Defining window-boundaries for genomic
analyses using smoothing spline techniques. Genetics Selection Evolution. 2015; 47(30):1-9.
676
Beissinger TM, CN Hirsch, B Vaillancourt, S Deshpande, K Barry, CR Buell, SM Kaeppler, D Gianola, N de
Leon. A Genome-Wide Scan for Evidence of Selection in a Maize Population Under Long-Term Artificial
Selection for Ear Number. Genetics. 2014; 196(3): 829–840.

Burke MK, JP Dunham, P Shahrestani, KR Thornton, MR Rose, AD Long. Genome-wide analysis of a long-
term evolution experiment with Drosophila. Nature. 2010; 467(7315): 587–590.

Burke MK. How does adaptation sweep through the genome? Insights from long-termselection
experiments. Proceedings of the Royal Society B: Biological Sciences. 2012; 279(1749): 5029–5038.

Burke MK, Liti G, Long AD. Standing Genetic Variation Drives Repeatable Experimental Evolution in
Outcrossing Populations of Saccharomyces cerevisiae. Molecular Biology and Evolution. 2014;31(12):
3228–3239.

Bradbury PJ, ZW Zhang, DE Kroon, TM Casstevens, Y Ramdoss, ES Buckler. TASSEL: software for
association mapping of complex traits in diverse samples. Bioinformatics. 2007; 23(19):2633-2635.

Cesconeto RJ, S Joost, CM McManus, SR Paiva, JA Cobuci, J Braccini. Landscape genomic approach to
detect selection signatures in locally adapted Brazilian swine genetic groups. Ecology and Evolution.
2017; 7(22): 9544–9556.

Crow JF, M Kimura. An introduction to population genetics theory. 1970. Harper & Row Publishers.

Danecek P, JK Bonfield, J Liddle, J Marshall, V Ohan, MO Pollard, A Whitwham, T Keane, SA McCarthy, RM
Davies, H Li. Twelve years of SAMtools and BCFtools. GigaScience. 2021; 10(2): giab008.

Deardorff ER, KA Fitzpatrick, GVS Jerzak, PY Shi, LD Kramer, GD Ebel. West Nile Virus Experimental
Evolution in vivo and the Trade-off Hypothesis. PLoS Pathogens. 2011; 7(11): e1002335.

Doebley J, JD Wendel, JSC Smith, CW Stuber, MM Goodman. The origin of cornbelt maize: the isozyme
evidence. Economic Botany. 1988; 42(1): 120 – 131.

Dudley JW, RJ Lambert. 100 Generations of Selection for Oil and Protein in Corn. 2003. Wiley & Sons.

Elshire RJ, JC Glaubitz, Q Sun, JA Poland, K Kawamoto, ES Buckler, SE Mitchell. A Robust, Simple
Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. PLoS ONE. 2011; 6(5), p. e19379.

Firman RC, F Garcia-Gonzalez, E Thyer, S Wheeler, Z Yamin, M Yuan, LW Simmons. Evolutionary change
in testes tissue composition among experimental populations of house mice. Evolution. 2015; 69(3):
848–855.

Firman RC, LW Simmons. Experimental evolution of sperm quality via postcopulatory sexual selection in
house mice. Evolution. 2009; 64-5: 1245–1256.

Fox AR, F Scochera, T Laloux, K Filik, H Degand, P Morsomme, K Alleva, F Chaumont. Plasma membrane
aquaporins interact with the endoplasmic reticulum resident VAP27 proteins at ER–PM contact sites
andendocytic structures. New Phytologist. 2020; 228:973-988.

Franssen SU, NH Barton, C Schlötterer. Reconstruction of Haplotype-Blocks Selected during Experimental
Evolution. Molecular Biology and Evolution. 2016; 34(1): 174–184.
712
Galli M, Q Liua, B L. Moss, S Malcomber, W Lia, C Gainese, S Federicia, J Roshkovana, R Meeley, JL
Nemhauserb, A Gallavottia. Auxin signaling modules regulate maize inflorescence architecture.
Proceedings of the National Academy of Sciences. 2015; 112(43): 13372–13377.

Gyawali A, V Shrestha, KE Guill, S Flint-Garcia, TM Beissinger. Single-plant GWAS coupled with bulk
segregant analysis allows rapid identification and corroboration of plant-height candidate SNPs. BMC
Plant Biology. 2019; 19(1):412.

Herring CD, A Raghunathan, C Honisch, T Patel, M Kenyon Applebee, AR Joyce, TJ Albert, FR Blattner, D
van den Boom, CR Cantor, BØ Palsson. Comparative genome sequencing of Escherichia coli allows
observation of bacterial evolution on a laboratory timescale. Nature Genetics. 2006; 38(12): 1406–1412.

Hirsch CN, SA Flint-Garcia, TM Beissinger, SR Eichten, S Deshpande, K Barry, MD McMullen, JB Holland,
ES Buckler, N Springer, CR Buell, N de Leon, SM Kaeppler. Insights into the Effects of Long-Term Artificial
Selection on Seed Size in Maize. Genetics. 2014;198(1): 409–421.

Janick, J. Plant Breeding Reviews: Long-Term Selection: Maize, Volume 24, Part 1. 2010a. John Wiley &
Sons, Inc.

Janick, J. (2010b) Plant Breeding Reviews, Volume 24, Part 2 Long-term Selection: Crops, Animals, and
Bacteria. 2010b. John Wiley & Sons, Inc.

Johansson AM, ME Pettersson, PB Siegel, Ö Carlborg. Genome-Wide Effects of Long-Term Divergent
Selection. PLoS Genetics. 2010; 6(11), e1001188.

Kelly SA, PP Czech, JT Wight, KM Blank, T Garland Jr. Experimental evolution and phenotypic plasticity of
hindlimb bones in high-activity house mice. Journal of Morphology. 2006; 267(3): 360–374.

Kessner D, J Novembre. Power Analysis of Artificial Selection Experiments Using Efficient Whole Genome
Simulation of Quantitative Traits. Genetics. Genetics, 2015; 199, 991–1005.

Kofler R, C Schlötterer. A Guide for the Design of Evolve and Resequencing Studies. Molecular Biology
and Evolution. 2014;31(2): 474–483.

Krimbas CB and S Tsakas. The Genetics of Dacus oleae. V. Changes of Esterase Polymorphism in a Natural
Population Following Insecticide Control-Selection or Drift? Evolution. 1971; 25(3): 454–460.

Kumar R, A Gyawali, GD Morrison, CA Saski, DJ Robertson, DD Cook, N Tharayil, RJ Schaefer, TM
Beissinger, RS Sekhon. Genetic Architecture of Maize Rind Strength Revealed by the Analysis of
Divergently Selected Populations. Plant and Cell Physiology. 2021; 62(7): 1199–1214.

Lenski RE. Experimental evolution and the dynamics of adaptation and genome evolution in microbial
populations. The ISME Journal. 2017; 11(10): 2181–2194.

Levy SF, JR Blundell, S Venkataram, DA Petrov, DS Fisher, G Sherlock. Quantitative evolutionary dynamics
using high-resolution lineage tracking. Nature. 2015; 519(7542): 181–186.

Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. 2013.

Linder RA, B Zabanavar, A Majumder, H Chiao-Shyan Hoang, VG Delgado, R Tran, VT La, SW Leemans, AD
Long. Adaptation in outbred sexual yeast is repeatable, polygenic, and favors rare haplotypes. Molecular
Biology and Evolution, 2022; 39(12): 1–25.

Liu J, AR Fernie, J Yan. The Past, Present, and Future of Maize Improvement: Domestication, Genomics,
and Functional Genomic Routes toward Crop Enhancement. Plant Communications. 2020;1(1),100010.

Long A, G Liti, A Luptak, O Tenaillon. Elucidating the molecular architecture of adaptation via evolve and
resequence experiments. Nature Reviews Genetics. 2015; 16(10): 567–582.

Lorenz AJ, TM Beissinger, R Rodrigues Silva, N de Leon. Selection for Silage Yield and Composition Did
Not Affect Genomic Diversity Within the Wisconsin Quality Synthetic Maize Population.G3
Genes|Genomes|Genetics. 2015; 5(4): 541–549.

Lush JL. Animal Breeding Plans. Ames, Ia., Collegiate Press. 1937.

Ma Y, X Ding, S Qanbari, S Weigend, Q Zhang, H Simianer. Properties of different selection signature
statistics and a new strategy for combining them. Heredity. 2015; 115(5): 426–436.

Maize GDB. Maize genetics and genomics database. 2021. www.maizegdb.org/.

Marks HL. Long-Term Selection for Body Weight in Japanese Quail Under Different Environments. Poultry
Science. 1996; 75(10): 1198–1203.

Matthes MS, NB Best, JM Robil, S Malcomber, A Gallavotti, P McSteen. Auxin EvoDevo: Conservation and
Diversification of Genes Regulating Auxin Biosynthesis, Transport, and Signaling. Molecular Plant. 2019;
12(3): 298–320.

Mazaheri M, M Heckwolf, B Vaillancourt, JL Gage, B Burdo, S Heckwolf, K Barry, A Lipzen, C Bastos
Ribeiro, TJY Kono, HF. Kaeppler, EP. Spalding, CN Hirsch, CR Buell, N de Leon, SM. Kaeppler. Genome-
wide association analysis of stalk biomass and anatomical traits in maize. BMC Plant Biology. 2019;
19(45): 1–19.

Moose SP, JW Dudley, TR Rocheford. Maize selection passes the century mark: a unique resource for
21st century genomics. Trends in plant science. 2014; 9: 358–364.

Michalak P, L Kang, MF Schou, HR Garner, V Loeschcke. Genomic signatures of experimental adaptive
radiation in Drosophila. Molecular Ecology. 2019; 28: 600–614.

Nieh SC, WS Lin, YH Hsu, GJ Shieh, BJ Kuo. The effect of flowering time and distance between pollen
source and recipient on maize. GM Crops & Food. 2014; 5(4): 287–295.

Orozco-terwengel P, M Kapun, V Nolte, R Kofler, T Flatt, C Schlötterer. Adaptation of Drosophila to a
novel laboratory environment reveals temporally heterogeneous trajectories of selected alleles.
Molecular Ecology. 2012; 21: 4931-4941.
779
780 Parts L, FA Cubillos, J Warringer, K Jain, F Salinas, SJ Bumpstead, M Molin, A Zia, JT Simpson, MA Quail, A
781 Moses, EJ Louis, R Durbin, G Liti. Revealing the genetic structure of a trait by sequencing a population
782 under selection. Genome Research. 2011; 21(7): 1131–1138.
783
Payen C, AB Sunshine, GT Ong, JL Pogachar, W Zhao, MJ Dunham. High-Throughput Identification of
Adaptive Mutations in Experimentally Evolved Yeast Populations. PLOS Genetics. 2016; 12(10), p.
e1006339.

Peiffer JA, MC Romay, MA Gore, SA Flint-Garcia, Z Zhang, MJ Millard, CAC Gardner, MD McMullen, JB
Holland, PJ Bradbury, ES Buckler. The genetic architecture of maize height. Genetics. 2014; 196(4): 1337–
1356.

Phillips MA, AD Long, ZS Greenspan, LF Greer, MK Burke, B Villeponteau, KC Matsagas, CL Rizza, LD
Mueller, MR Rose. Genome-wide analysis of long-term evolutionary domestication in Drosophila
melanogaster. Scientific Reports. 2016; 6: 39281.

Pook, T, M Schlather, G de los Campos, M Mayer, CC Schoen, H Simianer. HaploBlocker: Creation of
Subgroup-Specific Haplotype Blocks and Libraries. Genetics. 2019; 212(4): 1045–1061.

R Core Team. R: A Language and Environment for Statistical. 2021. https://CRAN.R-
project.org/package=doParallel.

Ramos Báez R, RY Buckley, H Yu, Z Chen, A Gallavotti, JL Nemhauser, BL Moss. A Synthetic Approach
Allows Rapid Characterization of the Maize Nuclear Auxin Response Circuit. Plant Physiology. 2020;
182(4): 1713–1722.

Remington DL, JM Thornsberry, Y Matsuoka, LM Wilson, SR Whitt, J Doebley, S Kresovich, MM Goodman,
ES Buckler. Structure of linkage disequilibrium and phenotypic associations in the maize genome. PNAS.
2001; 98(20): 11479–11484.

Remolina SC, PL Chang, J Leips, SV. Nuzhdin, KA Hughes. GENOMIC BASIS OF AGING AND LIFE-HISTORY
EVOLUTION IN DROSOPHILA MELANOGASTER. Evolution, 2012; 66: 3390-3403.

Romay MC, MJ Millard, JC Glaubitz, JA Peiffer, KL Swarts, TM Casstevens, RJ Elshire, CB Acharya, SE
Mitchell, SA Flint-Garcia, MD McMullen, JB Holland, ES Buckler, CA Gardner. Comprehensive genotyping
of the USA national maize inbred seed bank. Genome Biology. 2013; 14(R55).

Ross AJ, AR Hallauer, M Lee. Genetic analysis of traits correlated with maize ear length. Maydica. 2006;
51: 301–313.

Tange, O. GNU Parallel - The Command-Line Power Tool. The USENIX Magazine. 2011; 42-47.

Teng F, L Zhai, R Liu, W Bai, L Wang, D Huo, Y Tao, Y Zheng, Z Zhang. ZmGA3ox2, a candidate gene for a
major QTL, qPH3.1, for plant height in maize. The Plant Journal. 2013; 73(3): 405–416.

Teotónio H, IM Chelo, M Bradić, MR Rose, AD Long. Experimental evolution reveals natural selection on
standing genetic variation. Nature Genetics. 2009; 41(2): 251–257.

Teotónio H, MR Rose. Variation in the reversibility of evolution. Nature. 2000; 408(6811): 463–466.

Tobler R, SU Franssen, R Kofler, P Orozco-terWengel, V Nolte, Joachim Hermisson, C Schlötterer. Massive
Habitat-Specific Genomic Response in D. melanogaster Populations during Experimental Evolution in Hot
and Cold Environments. Molecular Biology and Evolution. 2014; 31(2): 364–375.

Turner TL, AD Stewart, AT Fields, WR Rice, AM Tarone. Population-Based Resequencing of Experimentally
Evolved Populations Reveals the Genetic Basis of Body Size Variation in Drosophila melanogaster. PLoS
Genetics. 2011; 7(3): e1001336.

Turner TL, Miller PM. Investigating Natural Variation in Drosophila Courtship Song by the Evolve and
Resequence Approach. Genetics. 2012; 191(2): 633–642.

Scheet P, M Stephens. A fast and flexible statistical model for large-scale population genotype data:
applications to inferring missing genotypes and haplotypic phase. American Journal of Human Genetics.
2006; 78(4): 629-644.

Silverman BW. Some Aspects of the Spline Smoothing Approach to Non-Parametric Regression Curve
Fitting. Journal of the Royal Statistical Society. 1985; 47(1): 1-52

Shin J, S Blay, B McNeney, J Graham. "LDheatmap: An R Function for Graphical Display of Pairwise
Linkage Disequilibria Between Single Nucleotide Polymorphisms." Journal of Statistical Software. 2006;
16(3).

U.S. National Plant Germplasm System. Details for: PI 269743, Zea mays L. subsp. mays, Shoepeg. 2021.
https://www.ars-grin.gov/.

Wallace JG, X Zhang, Y Beyene, K Semagn, M Olsen, BM Prasanna, ES Buckler. Genome-wide Association
for Plant Height and Flowering Time across 15 Tropical Maize Populations under Managed Drought
Stress and Well-Watered Conditions in Sub-Saharan Africa. Crop Science. 2016; 56(5): 2365–2378.

Wang K, P Wu, Q Yang, D Chen, J Zhou, A Jiang, J Ma, Q Tang, W Xiao, Y Jiang, L Zhu, X Li and G Tang.
Detection of Selection Signatures in Chinese Landrace and Yorkshire Pigs Based on Genotyping-by-
Sequencing Data. Frontiers in Genetics. 2018; 9:119.

Weir BS and CC Cockerham. Estimating F-Statistics for the Analysis of Population Structure. Evolution.
1984; 38(6): 1358– 1370.

Wichman HA, MR Badgett, LA Scott, CM Boulianne, JJ Bull. Different Trajectories of Parallel Evolution
During Viral Adaptation. Science. 1999; 285(5426): 422–424.

Wickland DP, G Battu, KA Hudson, BW Diers, ME Hudson. A comparison of genotyping-by-sequencing
analysis methods on low-coverage crop datasets shows advantages of a new workflow, GB-eaSy. BMC
Bioinformatics. 2017; 18:586.
847

848