Overexpressing NRT2.7 induces nitrate export from the vacuole and increases

growth of Arabidopsis

Patrick Armengaud[1*], Alexis De Angeli[2**], Patrick Berquin[1], Virginie Bréhaut[1], Mickaël
Durand[1***], Françoise Daniel-Vedele[1], Anne Krapp[1#] and Sophie Filleur[2,3#]

[1] Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin (IJPB),
78000, Versailles, France; [2] Université Paris-Saclay, CEA, CNRS, Institute for
Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France ; [3] Université Paris
Cité, UFR Sciences du Vivant, 35 rue Hélène Brion, F-75205 Paris Cedex 13, France.

*Present address: INRIA Rocquencourt, Domaine de Voluceau, Rocquencourt -
BP105, 78153 Le Chesnay Cedex, France
**Present address: Institute for plant sciences of Montpellier (IPSiM), Université de
Montpellier, CNRS, INRAE, Institut Agro, Montpellier, 34060, France.
***Present address: Biomolécules et Biotechnologies Végétales, EA2106, Université
de Tours, Tours, France
15

Email addresses of the authors :
Patrick Armengaud: patrick.armengaud@inria.fr
Alexis De Angeli: alexis.deangeli@cnrs.fr
Patrick Berquin: patrick.berquin@inrae.fr
Virginie Bréhaut : virginie.brehaut@inrae.fr
Mickaël Durand : midurand@univ-tours.fr
Françoise Daniel-Vedele : fravedele@yahoo.fr
Anne Krapp ([#] co-corresponding authors) : anne.krapp@inrae.fr
Sophie Filleur ([#] co-corresponding authors) : sophie.filleur@i2bc.paris-saclay.fr
25
26
27

28 **Date of submission: 26/02/2024**
29
30 **Number of tables: 0**
31
32 **Number of figures: 5**
33
34 **Word count: 4592**
35
36 **Number of supplemental figures: 4**
37
38 **Number of supplementary table: 1**
39
40
41
42
43 **Running title:** Physiological impact of vacuolar nitrate transport
44
45

Highlight :

The overexpression of the nitrate transporter NRT2.7 stimulates growth when nitrogen supply is limiting by increasing the export of nitrate from the vacuole, the main cell compartment for nitrate storage.

Abstract

Nitrogen nutrition is essential for crop yield but applying fertilizers has detrimental effects on the environment. Improved nutrient use efficiency is therefore a highly desired trait for developing a more sustainable agriculture. Compartmenting nitrate into vacuoles is one of the option to develop N-efficient crop adapted to less fertilizers. Only few proteins involved in nitrate transport on the tonoplast have been identified. CLCa is the major transporter involved in nitrate storage in Arabidopsis. Several other nitrate transporters amongst NRT2.7 have been localized in this membrane. The transport mechanism of NRT2.7 has not yet been defined as this protein is present mainly in seed cells that are not easily amenable for electrophysiology analysis. We then investigated its function by ectopically overexpressing it in a *clca* knock-out mutant. Although the growth on nitrogen sufficient medium was complemented, nitrate homeostasis was not restored by NRT2.7 activity like for CLCa overexpression. Moreover, NRT2.7 ectopic overexpression in wild-type background increased growth under limited nitrogen supply, suggesting that NRT2.7 stimulates nitrate efflux from vacuoles. This result was confirmed by electrophysiology performed on isolated vacuoles. Possible means of the growth stimulation by NRT2.7 versus CLCa are discussed based on nitrate fluxes through plasma membrane and nitrate homeostasis.

Keywords: Arabidopsis, Electrophysiology, Nitrate homeostasis, Nitrate transporter, NRT, vacuolar membrane

Abbreviations:

CaMV: Cauliflower Mosaic Virus

CLC: ChLoride Channel

KO: Knock-out

N: Nitrogen

NPF: Nitrate transporter 1/Peptide transporter Family

NRT2: High affinity Nitrate transporter

NUE: Nitrogen Use Efficiency

WS: Wassilewskija Arabidopsis accession

Introduction

Nitrogen (N) is an essential macronutrient acquired by plants from the soil or the air to synthesize amino acids, nucleotides and proteins. Most cultivated plant species rely on soil soluble ammonium and nitrate, the latter being the most abundant oxidized form of nitrogen found in temperate agricultural soils in amounts ranging from 0.1 to 10 mM (Miller *et al.*, 2007). The effect of nitrogen supply on plant growth has long been evidenced and this nutritional benefit was the basis of the fertilizer-driven yield increases (Epstein and Bloom, 2005). Crops are therefore abundantly provided with nitrogen fertilizers to insure best yield. This practice is at high economic and ecological costs since it has been shown that plants use only 25 to 50 % of nitrogen applied in the field (Cassman *et al.*, 2002). Because of both nitrate mobility and soil structure, nitrate leaks from the field, contaminates water effluents and stimulates growth of algae and microorganisms leading to ecological problems such as river eutrophication (Howarth and Marino, 2006). To avoid these problems and move toward a more sustainable agriculture, one means is to reduce fertilizer inputs without affecting crop yields by improving plant nitrogen use efficiency (NUE). This objective is challenging since NUE is a complex trait that includes uptake, utilization and remobilization efficiencies of the whole plant or of specific organs such as grains (Liu *et al.*, 2022). Therefore, research on NUE improvement has been tackled through approaches engineering known molecular players (transporters, enzymes) or heuristically by QTL, GWAS and transcriptome candidate genes approaches. However, to date, crop varieties with optimized NUE are still rare (see for reviews: (Liu *et al.*, 2022; Melino *et al.*, 2022; Govindasamy *et al.*, 2023).

The large number of nitrate transport mechanisms and their multiple level of regulation form a complex network coordinating nitrate fate and homeostasis. The latter is illustrated by remarkably stable cytosolic nitrate levels ranging between 3 and 5 mM (Cookson *et al.*, 2005). Once absorbed into root cells, nitrate has three possible paths (Dechorgnat *et al.*, 2011). A first one is a direct export to vascular tissue partially through xylem associated nitrate transporters (Wang *et al.*, 2018 for review). A second one is the assimilation path where cytosolic nitrate is reduced first by the nitrate reductase to nitrite and then by the plastidial nitrite reductase to form ammonium. Ammonium is then incorporated into organic acids to produce amino acids. The third path of cytosolic nitrate is vacuolar loading. The vacuolar compartment seems to play

important roles in controlling of the plant nitrate status as vacuolar nitrate accumulation has two major functions (Han *et al.*, 2016; Hodin *et al.*, 2023). First, vacuolar nitrate represents a temporary reserve of nitrogen allowing a near constant cytoplasmic provision independently of external fluctuations (van der Leij *et al.*, 1998; Matt *et al.*, 2001). Second, vacuolar nitrate acts as an osmoticum contributing to the osmotic potential, including during stomata movements (McIntyre, 1997; Wege *et al.*, 2014; Hodin *et al.*, 2023).

The first nitrate/proton exchanger mediating nitrate influx into the vacuole, CLCa, was characterized in Arabidopsis (De Angeli *et al.*, 2006). This protein is part of a family of seven members in this species. CLCa and CLCb were shown to preferentially exchange nitrate over chloride against proton (De Angeli *et al.*, 2006; Bergsdorf *et al.*, 2009; Wege *et al.*, 2010; Barbier-Brygoo *et al.*, 2011). Both of these exchangers are located on the vacuolar membrane. CLCa activity accounts for up to 50% of vacuolar nitrate content whereas it was proposed that CLCb mediates nitrate efflux from the vacuoles when plants are starved (Geelen *et al.*, 2000; Shi *et al.*, 2023). Impaired nitrate flux into vacuoles through CLCa was shown to reduce nitrate uptake, nitrate content and expression level of several nitrate transporter genes but to increase nitrate assimilation and consequently NUE (Monachello *et al.*, 2009; Hodin *et al.*, 2023).

Next to CLC transporters, some members of NRT2 and NPF transporters family are localized in the vacuolar membrane (Hu *et al.*, 2016; He *et al.*, 2017; Wang *et al.*, 2018*b*; Lu *et al.*, 2022). Among them, NRT2.7 belongs to the seven members of the NRT2 gene family in Arabidopsis that encodes major high affinity nitrate transporters. All NRT2 proteins except NRT2.7 are localized at the plasma membrane. NRT2.7 forms indeed a distinct monophyletic group within the Arabidopsis NRT2 family. The expression of the *NRT2.7* gene occurs mainly in seeds, but its expression has been also described for leaves and roots (Orsel *et al.*, 2002; Okamoto *et al.*, 2003; Chopin *et al.*, 2007). NRT2.7 is able to mediate nitrate fluxes in Xenopus oocytes. The main phenotype of *nrt2.7* knock-out mutants is a greatly decreased nitrate content in seeds which highlights the major roles of NRT2.7 in nitrate accumulation at reproductive stage (Chopin *et al.*, 2007).

Regarding nitrate as a signal, a nutrient or an electro-osmotic charge, further work in characterizing transporters and channels involved in vacuolar fluxes will shed light on the putative sensing and signaling properties of the largest compartment of plant cells.

The mechanism of nitrate transport by NRT2.7 was not investigated *in planta* as this protein is essentially present in seeds. So, in this report, we first investigated the impact of a NRT2.7 overexpression in Arabidopsis on nitrate homeostasis, activities of plasma membrane nitrate uptake systems, growth and vacuolar nitrate currents. Functional characterization of nitrate transporters *in planta* often requires the genetic removal of the quantitatively most important components to uncover more subtle or fine-tuning mechanisms. For example, the nitrate transport activity of NRT2 family members are masked by the NRT2.1 activity and has to be assessed in the *nrt2*.1 deficient background (Chopin *et al.*, 2007; Kiba *et al.*, 2012; Lezhneva *et al.*, 2014). Here we applied a similar strategy and used the *clca* mutant background in addition to a wild-type background, to test the physiological impact of NTR2.7 activities *in planta* and to test by electrophysiology the effect the overexpression of NRT2.7 in *clca* on vacuolar nitrate currents. We thus elucidate further the function of NRT2.7 by asking the following questions: (i) whether or not overexpression of NRT2.7 nitrate transporter would give similar effects as overexpression of the CLCa nitrate exchanger in the *clca* knock-out genetic background, (ii) how the NRT2.7 overexpression in wild-type background would affect nitrate homeostasis and growth under nitrate sufficient and limiting conditions and, (iii) what are the consequences of *NRT2.7* overexpression on nitrate fluxes through the vacuolar membrane.

Material and methods

Plant material and growth conditions

All *Arabidopsis thaliana* genotypes used in this study were obtained or generated in the Wassilewskija (WS) background. The *clca-2* T-DNA knock-out plant line originates from the Versailles collection (FST 171A06, line EAE89] whereas *clca-1* was isolated in Geelen *et al.* (2000). The *CLCa-OX* (*clca-2*) line (*clca-2* knock-out mutant overexpressing *CLCa* under the 35S promoter of the Cauliflower Mosaic Virus, CaMV) originates from Wege *et al.* (2010). The *NRT2.7-OX* (*clca*) lines (*clca* knock-out mutant overexpressing *NRT2.7* under the 35S promoter of the CaMV) was obtained by crossing NRT2.7 overexpressors (in the WS background, Chopin *et al.*, 2007) with the *clca* mutants (*clca-1* and *clca-2*).

Seeds were sterilized, stored for 3 days at 4°C to synchronize germination and sown

in 120 x 120 mm square Petri dishes containing either 35 or 70 mL nutrient medium to

monitor root architecture (4-5 plants/plate) or to produce material for analysis (12-15

plants/plate). Media differed in their nitrate content (0.2|2|20 mM) and consisted

respectively of 0.2|0.7|7 mM KNO_3; 0|0.65|6.5 mM $Ca(NO_3)_2$; 0.5|0.5|1.5 mM $MgSO_4$;

0.625|0.625|0 mM KH_2PO_4; 0|0|1.875 mM NaH_2PO_4; 0.75|1.25|1.25 mM $NaCl$;

0.65|0.25|0 mM $CaCl_2$ and 1.05|0.55|0 mM KCl. For all media, micronutrients (in μM)

consisted of 42.5 FeNaEDTA, 0.16 $CuSO_4$, 45 H_3BO_3, 0.015 $(NH_4)_6Mo_7O_{24}$, 0.01

$CoCl_2$, 0.38 $ZnSO_4$, 1.8 $MnSO_4$. All media were supplemented with 1% sucrose, pH

balanced at 5.8 before autoclaving and solidified using 1% agar (Sigma A1296). Petri

dishes were sealed using 3M adhesive tape and placed below tubes delivering 65 μE

irradiance for 16h/day in a tissue culture room with a thermoperiod day/night of

22/18°C. At 12 days after germination (14 days after sowing), plants were harvested

(or sampled for ^{15}N uptake) between 10 and 12 am (after 3 to 5h light).

Growth measurements

Root architecture was measured at 3, 6 and 9 days after germination by scanning Petri

dishes using a flat-bed scanner (Epson Perfection 4490 Photo) at a 200 dots per inch

resolution. Total root length was acquired by analyzing root morphology with the EZ-

Rhizo software (Armengaud *et al.*, 2009). Plant biomass was measured directly on a

microbalance (Mettler Toledo XS3DU) for fresh weight or after sampling in tin capsules,

dried 48h at 80°C and weighted for dry matter measurements.

RNA extraction and RT-qPCR analysis

RNA was extracted from both roots and shoots using the Trizol reagent (Invitrogen)

according to the provider recommendations. One microgram was treated with DNase

and reverse transcribed using the Quantitect kit (Qiagen) following the manufacturer

instructions. For qPCR analysis, specific primers were used for NRT2.7 (5'-

CCTTCATCCTCGTCCGTTTC-3' (forward) and 5'- AATTCGGCTATGGTGGAGTA-3'

(reverse)) and EF1α (At5g60390) 5'- CTGGAGGTTTTGAGGCTGGTAT-3' (forward)

and 5'- CCAAGGGTGAAAGCAAGAAGA-3' (reverse). The reactions were performed

in a Realplex mastercycler (Eppendorf) using commercial SYBR green mastermix

(Roche). Samples were subjected to 2 min pre-incubation at 95°C followed by 40

amplification cycles of 15 s at 95°C, 15 s at the melting temperature of the primers set

and 20 s at 68°C. NRT2.7 transcript levels were quantified using linear regression of quantified amplicon range and expressed relatively to EF1α.

Nitrate, C/N analysis

Frozen plant material was ground in liquid nitrogen using a Retsch mill grinder (MM301, Retsch GmbH, Germany). An aliquot of 20 mg of frozen powder was used to prepare an ethanol-water extract as described in Cross *et al.* (2006). All metabolites measurements were performed from this extract. Nitrate content was assayed after its reduction by vanadium(III) combined with detection by the acidic Griess reaction as described in Miranda *et al.* (2001). Total carbon and nitrogen content were determined using an elemental analyzer (EA, Flash 2000, Thermo Scientific) on aliquots of 0.5 to 2 mg dry weight in tin capsules. EA settings consisted of 20 sec of oxygen flux at 80 mL/min and helium at 140 mL/min for 5 min.

^{15}N uptake measurements

Influx of $^{15}NO_3^-$ was assayed as previously described (Orsel *et al.*, 2004). Two to three 12 day-old plants were first transferred to 0.1 mM $CaSO_4$ for 1 min. To measure high affinity ^{15}N nitrate uptake, plants were then transferred to a liquid 0.2 mM nitrate containing medium (the same as described earlier but without sucrose) enriched at 99% with $^{15}NO_3^-$ (Sigma). To measure low affinity transport capacity, plants were instead transferred to a 6 mM nitrate medium consisting of 3 mM $K^{15}NO_3$; 1.5 mM $Ca(^{15}NO_3)_2$; 0.5 mM $MgSO_4$; 0.625 mM KH_2PO_4; 1.5 mM NaCl and the same micronutrients as described before. ^{15}N nitrate uptake was performed for 5 min and plants were finally incubated for 1 min in 0.1 mM $CaSO_4$. Roots were then placed into tin capsules, dried at least 48h at 80°C and sent for ^{15}N elemental analysis using an ANCA–MS system (PDZ Europa, Crewe, UK) at INRAE Montpellier. Nitrate uptake was calculated from the ^{15}N content of the roots (0.5 to 1.5 mg DW). Each uptake experiment included at least 5 measurements per condition and genotype.

Patch-clamp analyses

Patch-clamp experiments were made in the whole-vacuole configuration with an access resistance between 1.5-3 MΩ. Patch-clamp recordings were made with an EPC8 amplifier (HEKA, Lambrecht-Pfalz Germany). Data were acquired with an AD/DA converter Instrutech LIH 8+8 under the control of the software PatchMaster (HEKA, Lambrecht-Pfalz Germany) and analysed with FitMaster software (HEKA,

Lambrecht-Pfalz Germany). Applied membrane potentials were always corrected by
the liquid junction potential (Neher, 1992). Current to voltage characteristics were
obtained measuring the amplitude of the stationary current. Measurements were done
after waiting that the vacuolar lumen solution was equilibrated with the pipette solution,
typically 10 minutes after the establishment of the whole-vacuole configuration.
Cytosolic side solutions were changed using a gravity driven perfusion system. Bis-
Tris-propane (BTP) was chosen as a large impermeable cation in order to minimize
cation currents. The solution used for patch-clamp experiments were (in mM): Nitrate
vacuolar solution: 100 BTP, 200 HNO_3, 0.1 $CaCl_2$, 5 $MgCl_2$, 5 MES, pH = 5.5. Chloride
vacuole solution : 475 MES, 15 HCl, 60.5 BTP, 0.1 $CaCl_2$, pH = 5.5. Chloride cytosolic
solutions: 7.5 BTP, 15 HCl, 0.1 $CaCl_2$, 2 $MgCl_2$, 15 MES, pH = 7. Nitrate cytosolic
solution: 0.1 $Ca(NO_3)_2$, 2 $Mg(NO_3)_2$, 15 MES, pH = 7 with BTP. Osmotic pressures of
the cytosolic and vacuolar solutions were adjusted with sorbitol to 640 and 650 mOsm,
respectively.

Statistical analysis
Statistical analyses were performed using the rcompanion and coin packages in the R
environment except for the electrophysiology results. For those, a Mann-Whitney test
was applied using GraphPad Prism 8 software. All the results of the statistical analysis
can be found in the supplementary table.
266

Results

Both CLCa and NRT2.7 overexpression restore growth of the clca-2 mutant
To set a suitable genetic background for NRT2.7 functional studies, we characterized
the phenotypes of *clca* mutants on a range of nitrate supplied as a sole nitrogen source
in vertical Petri dishes. We used *clca-1* and *clca-2* knock-outs (Geelen *et al.*, 2000;
De Angeli *et al.*, 2006) which both showed a significant decrease in shoot fresh weight
(-13 to -45%) and root growth rate (-8 to -40%) when compared to the WS wild-type
(Supplementary Fig. S1A-B and S2A-B). This decrease was more important under
non-limiting nitrate supply (2 and 20 mM). As expected *clca mutants* showed reduced
nitrate content compared to WS under all N supplies in both shoot and root
(Supplemental Fig. S1C-D and S2C-D).

To study the activity of NRT2.7 we overexpressed ectopically NRT2.7 in the *clca-2*
background and as control in the WT. These lines have been obtained by crossing the
clca mutants with the NRT2.7 overexpressing lines *NRT2.7-Ox1 (WS)* and *NRT2.7-*
Ox2 (WS) (Chopin *et al.*, 2007). Shoot NRT2.7 expression was increased 9 and 10
times, respectively compared to WS (Supplementary Fig. S3). Both lines NRT2.7-Ox1
(*clca-2*) and NRT2.7-Ox2 (*clca-2*) showed similar phenotypes and representative data
for one line are presented. For comparison, we included in our studies a *clca-2* mutant
overexpressing the *CLCa* cDNA under the control of the strong 35S promoter of the
CaMV (referred thereafter as CLCa-Ox (*clca-2*)) (Wege *et al.*, 2010). In this line, *CLCa*
transcript levels were increased 30-fold compared to WS (Wege *et al.*, 2014).
Based on the more pronounced differences in growth between *clca-2* and WS under
non-limiting external N supply, we cultivated the plants on 2 mM nitrate medium for all
further analyses using both *clca-2* lines overexpressing either *CLCa* or *NRT2.7*
compared to the controls WS, *NRT2.7* over-expressing lines (NRT2.7-Ox1 (WS)) and
clca-2. Both *CLCa* and *NRT2.7* overexpression restored the *clca-2* growth phenotypes
for shoot biomass and total root length (Fig. 1A and 1C). This growth restoration was
also observed in both lines overexpressing *NRT2.7* in the *clca-1* knock-out background
and occurred regardless external nitrate availability (Supplementary Fig. S2A-B). Wild-
type plants overexpressing NRT2.7 showed no significant difference in shoot fresh
weight and an important increase of 160% in total root length compared to the wild
(Fig. 1B-C). The growth stimulation observed for NRT2.7 overexpressing lines both in
the *clca-2* and the WS background, seems thus to be independent of the *clca-2*
mutation. The physiological changes triggered by NRT2.7 overexpression in the *clca*
mutants might thus differ from the complementation by CLCa while both led to restored
growth.

NRT2.7 and CLCa overexpression have opposite effects on both nitrate content and
nitrate uptake capacities
To decipher the underlying mechanism that impact growth of these overexpressing
lines, we determined nitrate content in roots and shoots. As described earlier, nitrate
content of *clca-2* was 53 and 60 % lower than the WT level in shoots and roots,
respectively. Expression of *CLCa* in the *clca-2* mutant restored nitrate levels in shoots
and roots, whereas *NRT2.7* overexpression in the *clca-2* background led to an about
two times decrease of the nitrate content when compared with the *clca-2* mutant (Fig.

2A and 2C). Likewise, *NRT2.7* overexpression does not complement *clca-1* phenotype for shoot and root nitrate content whatever is the external nitrate concentration for plant growth (Supplementary Fig. S2C-D). This means that the increased growth observed in the *NRT2.7-Ox (clca-2)* and *NRT2.7-OX (clca-1)* lines occurred without restoring nitrate content. Similar to the effect of *NRT2.7* overexpression observed in the *clca-2* mutant background, overexpression of *NRT2.7* in wild-type background was accompanied by a slight decrease in nitrate content in roots but no significant modification was observed in shoot (Fig. 2B and 2D).

In order to explain the impact on nitrate levels, we then investigated how nitrate uptake was affected in these lines. Nitrate uptake was measured using ^{15}N labelled nitrate at concentration of 0.2 and 6 mM to monitor both high and low affinity transport system activities (HATS and LATS, respectively, this latter activity being obtained as the difference between uptake measurements at 6 mM and 0.2 mM). Figure 3 shows that the HATS activity was decreased in *clca-2* whereas the LATS activity was not significantly different from that of wild-type plants. The *CLCa-Ox* (*clca-2*) line exhibited the same uptake capabilities for both transport activities as wild-type plants, illustrating a full complementation of the *clca-2* mutant phenotype. The *NRT2.7-Ox* (*clca-2*) line showed similar HATS activity as the *clca-2* knock-out mutant and an increase LATS activity (Fig. 3B and 3C). The *NRT2.7-Ox* (WS) lines displayed comparable tendencies with a 2-fold increased activity of the LATS compared to WT (Fig. 3C). Thus, whereas *CLCa* overexpression in the *clca* mutants increased mainly HATS activity compared to wild-type, *NRT2.7* overexpression had the opposite effect, supporting the hypothesis that these two nitrate transporters have different functions *in planta* when overexpressed ectopically.

NRT2.7 overexpression leads to increased biomass under limited and non- limited conditions

Plants stimulate their root growth when their N nutritional status is low (Jia and von Wirén, 2020). Given the important increase in root growth of the *NRT2.7-Ox* (WS) lines in wild-type background under non-limiting N supply (Fig. 1), we further studied the growth-promoting effect of *NRT2.7* overexpression under limiting N supply to better understand the changes of nitrate fluxes through the vacuolar membrane on N status. Plants were grown on vertical agar medium containing 0.2 mM nitrate for 12 days. Shoot biomass and total root length of *NRT2.7-Ox* (WS) lines were increased

compared to WT by 50 to 65% and 18 to 32% respectively (Fig. 4A and 4B). Conversely, root nitrate content of the *NRT2.7-OX* (WS) lines was dramatically reduced by 75 to 83% compared to WT, whereas shoot nitrate content was not modified (Fig. 4C and 4D). The carbon to nitrogen (C/N) ratio is a good indicator for plant growth, development and metabolism (Coruzzi and Zhou, 2001). The C/N ratio of *NRT2.7-Ox* (WS) compared to WT was increased in shoots and roots mainly due to a decrease in total N percentage (Fig. 4E). These data suggest that the ectopic expression of *NRT2.7* drives nitrate use for biomass production by increasing nitrate assimilation. This difference may probably due to a lower vacuolar storage capacity and/or an increased nitrate flux out of the vacuole as it was previously showed for other vacuolar nitrate transporters (He *et al.*, 2017; Lu *et al.*, 2022; Hodin *et al.*, 2023; Shi *et al.*, 2023) and no difference in nitrate uptake though the HATS was previously observed (Fig. 3).

NRT2.7 overexpression in clca-2 mutant leads to increased nitrate fluxes out of the vacuole

To determine if nitrate fluxes across the vacuolar membrane of the *NRT2.7-Ox* (*clca-2*) line were different from those of the knock-out mutant *clca-2*, we extracted vacuoles from mesophyll protoplasts of the *clca-2* and the two *NRT2.7-Ox* lines (*clca-2*) and performed patch clamp experiments in whole-vacuole configuration (Fig. 5). In order to maximize the fluxes of nitrate directed from the vacuole to the cytosol and from the cytosol to the vacuole, we performed two independent series of experimentations designed as follow. In both experiment series, we used the non-permeant cation BisTriPropane in order to limit the vacuolar currents mediated by cations. With the aim to highlight the efflux of nitrate from the vacuole (positive ionic currents), in the first series of experiments (Fig. 5B), we used bi-ionic conditions with 200 mM nitrate in the vacuole and 19.2 mM chloride in the cytosolic-side solution. The vacuoles extracted from *clca-2* knock-outs presented a mean current density of +1.3 ± 0.4 pA/pF at +43 mV and a mean reversal potential (E_{rev}=-8 ± 3 mV) that were comparable to those previously observed in similar ionic conditions (De Angeli *et al.*, 2006; Wege *et al.*, 2010). Interestingly, in the same ionic conditions vacuoles from *NRT2.7-Ox (clca-2)* plants presented significantly higher current densities compared to *clca-2* currents only at positive vacuolar membrane potentials (from +3 mV to +83 mV, Fig. 5B). These positive currents are compatible with an increased nitrate efflux capacity of the

vacuoles from *NRT2.7-Ox (clca-2)* plants. In the second set of experiments, we used

an ionic configuration suited to highlight nitrate currents directed from the cytosol to

the vacuolar lumen. We performed cytosolic-side solution exchange experiments

starting from a chloride based buffer (19.2 mM Cl^- pH 7) to a nitrate based buffer (4

mM NO_3^- pH 7) keeping the vacuolar-side buffer constant and with only chloride as a

permeant anion (15 mM Cl^- pH 5.5) (Supplementary Fig. S4). Upon the exchange of

the cytosolic-side solution a similar increase of the negative current was observed in

vacuoles obtained from the tested genotypes. Indeed, the mean current ratios at -60

mV were I_{NO3}/I_{Cl}=1.84±0.4 in *NRT2.7-Ox (clca-2)* vacuoles and I_{NO3}/I_{Cl}=1.77 ± 0.44 in

clca-2 vacuole*s*. Conversely, the current densities in presence of nitrate did not show

a significant difference between the genotypes, being I_{NO3}=-2.4±1 pA/pF at -60 mV in

NRT2.7-Ox (*clca-2*) and I_{NO3}=-2.7±0.9 pA/pF at -60 mV in *clca-2.* This second series

of experiments indicates that the vacuoles obtained from *NRT2.7-Ox* (*clca-2*) do not

display an increased capacity of driving NO_3^- fluxes directed to the vacuolar lumen

compared to the vacuoles from *clca-2* under our experimental conditions. Altogether,

the results obtained by patch-clamp experiments suggest that, compared to vacuoles

from *clca-2*, *NRT2.7-Ox* (*clca-2*) plant vacuoles present a higher capacity to drive

nitrate fluxes directed into the cytosol but are not modified concerning nitrate influx into

the vacuole.

Discussion

Recent studies underline the importance of nitrate fluxes through the vacuolar

membrane to enhance NUE by increasing efflux of nitrate from vacuole and/or reducing

nitrate storage in this latter compartment (Han *et al.*, 2016; Wang *et al.*, 2018*b*; Hodin

et al., 2023; Shi *et al.*, 2023). Therefore, it became essential to characterize the

different vacuolar nitrate transporter activities to get an overview of the specificity of

each. In this report, we analyzed the function of the vacuolar NRT2.7 transporter by

introducing its cDNA under the control of the strong *35S* promoter in *clca* KO mutant

or in wild-type. The combined results of on one hand the electrophysiological analyses

and on the other hand the phenotypes of the *CLCa* and *NRT2.7* overexpressing lines

strikingly demonstrate that both NRT2.7 and CLCa impacts the nitrogen use but

through distinct means or signals. The NRT2.7 overexpression in the *clca-2* mutant

bypassed the CLCa-dependent reduced growth via a mechanism that uncoupled

growth and nitrate accumulation probably related to modified nitrate efflux from the

vacuole. This mechanism when acting in a wild type background overexpressing

NRT2.7 led to an increase of biomass production mainly through a higher root biomass.

NRT2.7 stimulates nitrate export rather than nitrate import into the vacuole when

overexpressed ectopically.

We confirmed in this study that *CLCa* mutant physiology was deeply affected by its

limited capacity of loading nitrate into vacuoles (Fig. 1 and 2; Supplementary Fig. S1

and S2). Besides CLCa, the NRT2.7 nitrate transporter is located at the tonoplast.

NRT2.7 translocates nitrate in Xenopus oocytes and its loss-of-function reduced nitrate

accumulation in seeds where the gene is chiefly expressed (Chopin *et al.*, 2007).

Although *NRT2.7* mRNA were detected in leaves and root, no phenotype could be

noticed in the KO mutant for these organs (Orsel *et al.*, 2002; Okamoto *et al.*, 2003;

Chopin *et al.*, 2007). To better characterize this transporter, we then overexpressed it

in wild-type and *clca* KO mutant backgrounds. The overexpression of *NRT2.7* fully

restores the reduced *clca-2* plant growth and even exceeds wild-type biomass but,

surprisingly, further diminished the low *clca-2* nitrate content (Fig. 1 and 2). It led to a

35% and 23% decreases of shoot and root nitrate pools, respectively, when compared

to *clca-2*. Therefore, these results showed that NRT2.7 could indeed rescue the *clca-*

2 growth reduction but through an alternative mechanism rather than a true functional

complementation. This hypothesis was confirmed as the NRT2.7 overexpressors

(*NRT2.7-Ox* (WS)) showed a 50 to 60% increase in total root length whereas a further

depletion of root nitrate content compared to wild-type plants was observed.

The decrease of root nitrate content caused by NRT2.7 ectopic overexpression is

difficult to conceive with a vacuolar nitrate importer. Chopin *et al.* (2007) reported an

influx of ^{15}N labeled nitrate into NRT2.7 expressing oocytes. This result indicates that,

in this biological system, NRT2.7 was targeted to the plasma membrane with

presumably its cytoplasmic terminus oriented to the oocyte cytoplasm. In this same

study, GFP-NRT2.7 translational fusion driven by the 35S promoter in Arabidopsis

cells clearly showed a tonoplastic localization. As NRT2.7 cytoplasmic terminus is

probably maintained in the plant cell cytoplasm (Jacquot *et al.*, 2020), NRT2.7 would

function as a nitrate efflux system from the vacuole to the cytoplasm. This orientation

would be more conceivable with the fact that NRT2s are assumed to be H^+/NO_3^-

symporters (Forde and Clarkson, 1999) and that roots of NRT2.7 overexpressors become depleted of nitrate at low external nitrate availability (Fig. 4). Supporting this hypothesis, the determination of nitrate fluxes across isolated vacuoles by patch clamp techniques, showed that vacuoles from *NRT2.7-Ox* (*clca-2*) plants present a higher capacity to drive nitrate fluxes directed into the cytosol than vacuoles from *clca-2* plants while the capacity to drive nitrate influx into the vacuole did not vary between these genotypes (Fig. 5 and Supplementary Fig. S4). However, based on these experiments, we cannot exclude that NRT2.7 may regulate positively the activity of nitrate exporters localized in the tonoplast and is not itself the active transporter driving the fluxes.

Chopin *et al.* (2007) observed a decrease in seed nitrate content in *nrt2.7* mutants, which was interpreted as a reduced capacity to import nitrate into the vacuole of the embryo. NRT2.7 is highly expressed in seeds and the NRT2.7 promoter drives expression in the embryos where the protein is located at the tonoplast. A decreased nitrate content is thus difficult to explain by the loss-of-function of a vacuolar nitrate exporter protein. However, the transport direction of NRT2.7 in seeds and in shoot/root vacuoles might be different because of distinct types of vacuoles existing in these organs (Shimada *et al.*, 2018). In addition, tonoplast membrane potential might vary and the transport direction may reflect the organ specific ionic and osmotic conditions as it has been discussed previously for NPF6.3 (NRT1.1/CHL1, (Léran *et al.*, 2014) or CLCa (Wege *et al.*, 2014). Besides, nitrate transport properties are regulated via posttranslational protein modifications (e.g. (Wege *et al.*, 2014; Jacquot *et al.*, 2017), a mechanism that should be explored in the future for NRT2.7. However no phosphorylation site in NRT2.7 is found in the PhosPhAt databases (Heazlewood *et al.*, 2008). Finally, another hypothesis would be that *nrt2.7* seed nitrate content is decreased because nitrate transport from the roots or shoots towards the seeds is diminished in the *nrt2.7* mutants. Indeed, in Arabidopsis seeds nitrate content seems to rely directly on the nitrate taken up by roots (Schulze *et al.*, 1994). Inflorescence grafting experiments are required to distinguish between seed prone or root/shoot prone determination of the decreased nitrate level in *nrt2.7* seeds. Altogether, these results support that NRT2.7 is essential for nitrate fluxes through the vacuolar membrane. It stimulates nitrate export from the vacuole in seedlings but the contrasting phenotype of modified NRT2.7 expression in reproductive and vegetative organs is a challenging hypothesis and needs further investigation.

Nitrate fluxes through CLCa and NRT2.7 alter root nitrate uptake

As previously reported, we confirmed that the mutation in *CLCa* reduced nitrate uptake activity in the high affinity range (Monachello *et al.*, 2009, Fig. 3) whereas the level of LATS activity was not modified. These results illustrate the existence of a coordinated signaling crosstalk between vacuolar and plasma membrane nitrate fluxes. An obvious signal for this coordinated transmembrane transport would be the nitrate anion itself. It was shown that loss of function of CLCa leads to a faster increase of the cytosolic nitrate concentration in response to an increase of the extracellular NO_3^- concentration in guard cells (Demes *et al.*, 2020). In this scenario, a transitory elevated cytosolic nitrate level would decrease nitrate uptake, increase the nitrate efflux from the cells and/or the assimilation pathways through the nitrate reductase (NR). This was confirmed in a recent study demonstrating that the inhibition of nitrate storage in the vacuole leads to an increase of the NR activity and a better NUE (Hodin *et al.*, 2023). In addition to nitrate, nutrient uptake activity is controlled by a wide range of biophysical (membrane potential, pH) or physiological signals (phosphorylation, metabolites) (see Amtmann and Blatt, 2009 for review). CLCa through its activity as a NO_3^-/H^+ exchanger is known for its function in adjusting the cytosolic pH in guard cells (Demes *et al.*, 2020). More recently, Jain and Schmidt (2024) showed that NRT2.1 phosphorylation level varies according to the environmental pH certainly to adjust cytosolic pH. Then the loss of function of *CLCa* could lead to the inactivation of NRT2.1 to compensate the change in cytosolic pH.

The increase of nitrate export from the vacuole due to the NRT2.7 overexpression leads probably to change in nitrate cytosolic content and/or cytosolic pH, as for CLCa. Even if this hypothesis needs to be tested using the nitrate/pH biosensor, ClopHensor (Demes *et al.*, 2020), these modifications could explain the decrease in HATS activity observed in *NRT2.7-Ox (clca-2)* like in *clca-2* plants (Fig. 3). However, the *NRT2.7* overexpression led to an increase of the LATS activity that was not the case when *CLCa* is invalidated. Lejay *et al.* (1999) showed that HATS activity is under the control of both NR activity and N status of the plant whereas that of LATS is controlled solely by NR activity. The discrepancy observed between CLCa and NRT2.7 on the nitrate uptake could suggest that nitrate influx and efflux in and out the vacuole do not have exactly the same effect on N assimilation and N status. The coordination between the direction of vacuolar nitrate fluxes and nitrate uptake at the plasma membrane will need to be further studied in the future.

NRT2.7 overexpression increases growth at low external and internal nitrate

The overexpression of *NRT2.7* led to a higher plant growth when it stimulates nitrate export from the vacuole. This phenotype could be explained not only by the increase of LATS activity but also by a higher nitrate long-distance transport from root to shoot. A study characterizing two *Brassica napus* genotypes differing by their capacity in nitrate storage in root previously showed that the decrease of root nitrate content induces a higher nitrate translocation from root to shoot leading to a better NUE (Han *et al.*, 2016). We can assume that a similar phenomenon occurred in the *NRT2.7* overexpressors. Further studies like, for example, analyzing the expression of *NRT1.5* and *NRT1.8*, two nitrate transporters involved in the long-distance nitrate transport in Arabidopsis (Lin *et al.*, 2008; Li *et al.*, 2010) will be needed to confirm our hypothesis.

Interestingly, shoot and root biomass increased in *NRT2.7-Ox* (WS) grown under limiting N nutrition even that a severe decrease of nitrate in root (-75 to -83% compared to wild-type) and a slight decrease of total nitrogen content (-8 to −19% compared to wild-type) occurred in these growth conditions (Fig. 4). These results confirm an apparent disconnection between growth and nitrate content in *NRT2.7* overexpressors plants. C and N metabolism are interconnected and regulate each other (Zheng, 2009). Interestingly, the C/N ratio was higher in the overexpressors compared to wild-type indicating that, even if plants are grown under low external N availability, the photosynthesis outputs higher and could facilitate the increase in biomass. The low internal nitrate levels could be then the consequence of N dilution occurring under accelerated growth. At the opposite, the increase of nitrate export from the vacuole could stimulate photosynthesis as shown in a *Brassica napus* genotype presenting reduced root nitrate storage (Han *et al.*, 2016) or, more recently when Arabidopsis plants overexpressing the CLCb nitrate vacuolar exporter are grown on low nitrate (Shi *et al.*, 2023).

In conclusion, we show that the two vacuolar nitrate transporters CLCa and NRT2.7 share limited functionality and are not exchangeable. Our results suggest that NRT2.7 works as or induces the activity of a vacuolar nitrate exporter when overexpressed ectopically. By exploiting genotypes with modified levels of vacuolar transporters, we show that changes in the nitrate influx or efflux from the vacuolar compartment lead to differences in root nitrate uptake suggesting the existence of a fine regulation between vacuolar and plasma membranes through biophysical and physiological networks of

vacuolar nitrate sensing and homeostasis. NRT2.7, whose function *in vivo* seemed to be restricted to metabolism in seeds, provides new physiological features when ectopically overexpressed. The link between NRT2.7 nitrate fluxes and carbon content requires further studies but may be the basis for improving plants with higher C/N ratios accompanied by increased growth when nitrogen is limiting.

Supplementary data

The following supplementary data are available at JXB online.

Fig. S1. Growth and nitrate content of *clca-2* knock-out mutant grown on different nitrate concentrations.

Fig. S2. Growth and nitrate content of *NRT2.7* overexpressors in the c*lca-1* knock-out mutant.

Fig. S3. Level of NRT2.7 expression in NRT2.7-Ox (WS) lines.

Fig. S4. Nitrate influx in vacuoles from *clca-2 35S:NRT2.7* expressing plants and from *clca-2* knock-out.

Acknowledgements

The authors are very grateful to Pascal Tillard for servicing the [15]N measurements, Anne Marmagne for helping with the elementary C/N analyzer and the IJPB platform 'The Plant Observatory', in particular Joël Talbotec and Philippe Maréchal for taking care of greenhouse cultures.

Author contributions

FDV, AK and SF : conceptualization design ; PA, FDV, ADA, AK and SF: experimental design ; PA, ADA, PB and VB performed the experimental work ; PA, MD, FDV, ADA, AK and SF analyzed the data; PA, FDV, AK and SF wrote the manuscript. All authors reviewed the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Funding statement

This work benefits from the support of the LabEx Saclay Plant Sciences-SPS (ANR-10-LABX-0040-SPS) and of the Agence Nationale de la Recherche (ANR-Nitrapool ANR08-Blan-008).

Data availability

573   All the raw data supporting the findings of this work are available upon request from

574   the corresponding authors, Anne Krapp and Sophie Filleur. Statistical analyses are

575   available in the supplementary table.

References

Amtmann A, Blatt MR. 2009. Regulation of macronutrient transport. The New Phytologist **181**, 35–52.

Armengaud P, Zambaux K, Hills A, Sulpice R, Pattison RJ, Blatt MR, Amtmann A. 2009. EZ-Rhizo: integrated software for the fast and accurate measurement of root system architecture. The Plant Journal: For Cell and Molecular Biology **57**, 945–956.

Barbier-Brygoo H, De Angeli A, Filleur S, Frachisse J-M, Gambale F, Thomine S, Wege S. 2011. Anion channels/transporters in plants: from molecular bases to regulatory networks. Annual Review of Plant Biology **62**, 25–51.

Bergsdorf E-Y, Zdebik AA, Jentsch TJ. 2009. Residues important for nitrate/proton coupling in plant and mammalian CLC transporters. The Journal of Biological Chemistry **284**, 11184–11193.

Cassman KG, Dobermann A, Walters DT. 2002. Agroecosystems, nitrogen-use efficiency, and nitrogen management. Ambio **31**, 132–140.

Chopin F, Orsel M, Dorbe M-F, Chardon F, Truong H-N, Miller AJ, Krapp A, Daniel-Vedele F. 2007. The Arabidopsis ATNRT2.7 nitrate transporter controls nitrate content in seeds. The Plant Cell **19**, 1590–1602.

Cookson SJ, Williams LE, Miller AJ. 2005. Light-dark changes in cytosolic nitrate pools depend on nitrate reductase activity in Arabidopsis leaf cells. Plant Physiology **138**, 1097–1105.

Coruzzi GM, Zhou L. 2001. Carbon and nitrogen sensing and signaling in plants: emerging 'matrix effects'. Current Opinion in Plant Biology **4**, 247–253.

Cross JM, von Korff M, Altmann T, Bartzetko L, Sulpice R, Gibon Y, Palacios N, Stitt M. 2006. Variation of enzyme activities and metabolite levels in 24 Arabidopsis accessions growing in carbon-limited conditions. Plant Physiology **142**, 1574–1588.

De Angeli A, Monachello D, Ephritikhine G, Frachisse JM, Thomine S, Gambale F, Barbier-Brygoo H. 2006. The nitrate/proton antiporter AtCLCa mediates nitrate accumulation in plant vacuoles. Nature **442**, 939–942.

Dechorgnat J, Nguyen CT, Armengaud P, Jossier M, Diatloff E, Filleur S, Daniel-Vedele F. 2011. From the soil to the seeds: the long journey of nitrate in plants. Journal of Experimental Botany **62**, 1349–1359.

Demes E, Besse L, Cubero-Font P, Satiat-Jeunemaitre B, Thomine S, De Angeli A. 2020. Dynamic measurement of cytosolic pH and [NO3 -] uncovers the role of the vacuolar transporter AtCLCa in cytosolic pH homeostasis. Proceedings of the National Academy of Sciences of the United States of America **117**, 15343–15353.

Forde BG, Clarkson DT. 1999. Nitrate and Ammonium Nutrition of Plants: Physiological and Molecular Perspectives. In: Callow JA, ed. Advances in Botanical Research. Academic Press, 1–90.

Epstein E, Bloom AJ. 2005. Mineral nutrition in plants. Ed 2nd. Sinauer Associates, Sunderland.

Geelen D, Lurin C, Bouchez D, Frachisse JM, Lelièvre F, Courtial B, Barbier-Brygoo H, Maurel C. 2000. Disruption of putative anion channel gene AtCLC-a in Arabidopsis suggests a role in the regulation of nitrate content. The Plant Journal: For Cell and Molecular Biology **21**, 259–267.

Govindasamy P, Muthusamy SK, Bagavathiannan M, *et al.* 2023. Nitrogen use efficiency-a key to enhance crop productivity under a changing climate. Frontiers in Plant Science **14**, 1121073.

Han Y-L, Song H-X, Liao Q, *et al.* 2016. Nitrogen Use Efficiency Is Mediated by Vacuolar Nitrate Sequestration Capacity in Roots of Brassica napus. Plant Physiology **170**, 1684–1698.

He Y-N, Peng J-S, Cai Y, Liu D-F, Guan Y, Yi H-Y, Gong J-M. 2017. Tonoplast-localized nitrate uptake transporters involved in vacuolar nitrate efflux and reallocation in Arabidopsis. Scientific Reports **7**, 6417.

Heazlewood JL, Durek P, Hummel J, Selbig J, Weckwerth W, Walther D, Schulze WX. 2008. PhosPhAt: a database of phosphorylation sites in Arabidopsis thaliana and a plant-specific phosphorylation site predictor. Nucleic Acids Research **36**, D1015-1021.

Hodin J, Lind C, Marmagne A, *et al.* 2023. Proton exchange by the vacuolar nitrate transporter CLCa is required for plant growth and nitrogen use efficiency. The Plant Cell **35**, 318–335.

Howarth RW, Marino R. 2006. Nitrogen as the limiting nutrient for eutrophication in coastal marine ecosystems: Evolving views over three decades. Limnology and Oceanography **51**, 364–376.

Hu R, Qiu D, Chen Y, Miller AJ, Fan X, Pan X, Zhang M. 2016. Knock-Down of a Tonoplast Localized Low-Affinity Nitrate Transporter OsNPF7.2 Affects Rice Growth under High Nitrate Supply. Frontiers in Plant Science **7**, 1529.

Jacquot A, Chaput V, Mauries A, *et al.* 2020. NRT2.1 C-terminus phosphorylation prevents root high affinity nitrate uptake activity in Arabidopsis thaliana. The New Phytologist **228**, 1038–1054.

Jacquot A, Li Z, Gojon A, Schulze W, Lejay L. 2017. Post-translational regulation of nitrogen transporters in plants and microorganisms. Journal of Experimental Botany **68**, 2567–2580.

Jain D, Schmidt W. 2024. Protein Phosphorylation Orchestrates Acclimations of Arabidopsis Plants to Environmental pH. Molecular & cellular proteomics: MCP **23**, 100685.

Jia Z, von Wirén N. 2020. Signaling pathways underlying nitrogen-dependent changes in root system architecture: from model to crop species. Journal of Experimental Botany **71**, 4393–4404.

Kiba T, Feria-Bourrellier A-B, Lafouge F, *et al.* 2012. The Arabidopsis nitrate transporter NRT2.4 plays a double role in roots and shoots of nitrogen-starved plants. The Plant Cell **24**, 245–258.

van der Leij M, Smith SJ, Miller AJ. 1998. Remobilisation of vacuolar stored nitrate in barley root cells. Planta **205**, 64–72.

Lejay L, Tillard P, Lepetit M, Olive F d, Filleur S, Daniel-Vedele F, Gojon A. 1999. Molecular and functional regulation of two NO3- uptake systems by N- and C-status of Arabidopsis plants. The Plant Journal: For Cell and Molecular Biology **18**, 509–519.

Léran S, Varala K, Boyer J-C, *et al.* 2014. A unified nomenclature of NITRATE TRANSPORTER 1/PEPTIDE TRANSPORTER family members in plants. Trends in Plant Science **19**, 5–9.

Lezhneva L, Kiba T, Feria-Bourrellier A-B, Lafouge F, Boutet-Mercey S, Zoufan P, Sakakibara H, Daniel-Vedele F, Krapp A. 2014. The Arabidopsis nitrate transporter NRT2.5 plays a role in nitrate acquisition and remobilization in nitrogen-starved plants. The Plant Journal: For Cell and Molecular Biology **80**, 230–241.

Li J-Y, Fu Y-L, Pike SM, *et al.* 2010. The Arabidopsis nitrate transporter NRT1.8 functions in nitrate removal from the xylem sap and mediates cadmium tolerance. The Plant Cell **22**, 1633–1646.

Lin S-H, Kuo H-F, Canivenc G, *et al.* 2008. Mutation of the Arabidopsis NRT1.5 nitrate transporter causes defective root-to-shoot nitrate transport. The Plant Cell **20**, 2514–2528.

Liu Q, Wu K, Song W, Zhong N, Wu Y, Fu X. 2022. Improving Crop Nitrogen Use Efficiency Toward Sustainable Green Revolution. Annual Review of Plant Biology **73**, 523–551.

Lu Y-T, Liu D-F, Wen T-T, Fang Z-J, Chen S-Y, Li H, Gong J-M. 2022. Vacuolar nitrate efflux requires multiple functional redundant nitrate transporter in Arabidopsis thaliana. Frontiers in Plant Science **13**, 926809.

Matt P, Geiger M, Walch-Liu P, Engels C, Krapp A, Stitt M. 2001. The immediate cause of the diurnal changes of nitrogen metabolism in leaves of nitrate-replete tobacco: a major imbalance between the rate of nitrate reduction and the rates of nitrate uptake and ammonium metabolism during the first part of the light period. Plant, Cell & Environment **24**, 177–190.

McIntyre GI. 1997. The role of nitrate in the osmotic and nutritional control of plant development. Australian Journal of Plant Physiology **24**, 103–118.

Melino VJ, Tester MA, Okamoto M. 2022. Strategies for engineering improved nitrogen use efficiency in crop plants via redistribution and recycling of organic nitrogen. Current Opinion in Biotechnology **73**, 263–269.

Miller AJ, Fan X, Orsel M, Smith SJ, Wells DM. 2007. Nitrate transport and signalling. Journal of Experimental Botany **58**, 2297–2306.

Miranda KM, Espey MG, Wink DA. 2001. A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. Nitric Oxide: Biology and Chemistry **5**, 62–71.

Monachello D, Allot M, Oliva S, Krapp A, Daniel-Vedele F, Barbier-Brygoo H, Ephritikhine G. 2009. Two anion transporters AtClCa and AtClCe fulfil interconnecting but not redundant roles in nitrate assimilation pathways. The New Phytologist **183**, 88–94.

Neher E. 1992. Correction for liquid junction potentials in patch clamp experiments. Ion Channels. Methods in Enzymology. Academic Press, 123–131.

Okamoto M, Vidmar JJ, Glass ADM. 2003. Regulation of NRT1 and NRT2 gene families of Arabidopsis thaliana: responses to nitrate provision. Plant & Cell Physiology **44**, 304–317.

Orsel M, Eulenburg K, Krapp A, Daniel-Vedele F. 2004. Disruption of the nitrate transporter genes AtNRT2.1 and AtNRT2.2 restricts growth at low external nitrate concentration. Planta **219**, 714–721.

Orsel M, Filleur S, Fraisier V, Daniel-Vedele F. 2002. Nitrate transport in plants: which gene and which control? Journal of Experimental Botany **53**, 825–833.

Schulze W, Schulze E-D, Stadler J, Heilmeier H, Stitt M, Mooney H a. 1994. Growth and reproduction of Arabidopsis thaliana in relation to storage of starch and nitrate in the wild-type and in starch-deficient and nitrate-uptake-deficient mutants. Plant, Cell & Environment **17**, 795–809.

Shi Y, Liu D, He Y, Tang J, Chen H, Gong P, Luo J-S, Zhang Z. 2023. CHLORIDE CHANNEL-b mediates vacuolar nitrate efflux to improve low nitrogen adaptation in Arabidopsis. Plant Physiology, kiad438.

Shimada T, Takagi J, Ichino T, Shirakawa M, Hara-Nishimura I. 2018. Plant Vacuoles. Annual Review of Plant Biology **69**, 123–145.

Wang Y-Y, Cheng Y-H, Chen K-E, Tsay Y-F. 2018*a*. Nitrate Transport, Signaling, and Use Efficiency. Annual Review of Plant Biology **69**, 85–122.

Wang W, Hu B, Yuan D, *et al.* 2018*b*. Expression of the Nitrate Transporter Gene OsNRT1.1A/OsNPF6.3 Confers High Yield and Early Maturation in Rice. The Plant Cell **30**, 638–651.

Wege S, De Angeli A, Droillard M-J, *et al.* 2014. Phosphorylation of the vacuolar anion exchanger AtCLCa is required for the stomatal response to abscisic acid. Science Signaling **7**, ra65.

Wege S, Jossier M, Filleur S, Thomine S, Barbier-Brygoo H, Gambale F, De Angeli A. 2010. The proline 160 in the selectivity filter of the Arabidopsis NO(3)(-)/H(+) exchanger AtCLCa is essential for nitrate accumulation in planta. The Plant Journal: For Cell and Molecular Biology **63**, 861–869.

Zheng Z-L. 2009. Carbon and nitrogen nutrient balance signaling in plants. Plant Signaling & Behavior **4**, 584–591.

Figures Legends

Fig. 1. Growth effect of *NRT2.7* overexpression in *clca-2* and wild-type backgrounds. Plant were grown on 2 mM nitrate for 12 days. Shoot fresh weight (A, B) and total root length (C, D) were measured on wild-type (WS), *clca-2*, complemented line (*CLCa-Ox* (*clca-2*)) and supplemented line (*NRT2.7-Ox* (*clca-2*)) (A, C) or WS and the two *NRT2.7* over-expressing lines, *NRT2.7-Ox1* (WS) and *NRT2.7-Ox2* (WS) (B, D). The number of samples were 4-5 for shoot fresh weight (one sample = a pool of 3-4 plants) when n=6-17 for total root length on individual plants. Presented data are means ± SEM. Different letters at the top of the bars represent significant difference (p value<0.05).

Fig. 2. Nitrate content of *NRT2.7* overexpression in *clca-2* and wild-type backgrounds. Wild-type (WS), *clca-2*, complemented line (*CLCa-Ox* (*clca-2*)) and *NRT2.7* over-expressing line in *clca-2* (*NRT2.7-Ox* (*clca-2*)) (A, C) or WS and the two *NRT2.7* over-expressing lines, *NRT2.7-Ox1* (WS) and *NRT2.7-Ox2* (WS) (B, D), were grown on 2 mM nitrate. After 12 days on the medium, shoot (A, B) and root (C, D) nitrate contents were measured. Two to three biological replicates were performed on each genotype, one sample corresponding to a pool of at least 50 plants. Presented data are means ± SEM. Different letters at the top of the bars represent significant difference (p value<0.05).

Fig. 3. Effect of *NRT2.7* overexpression in *clca-2* and wild-type backgrounds on nitrate uptake activities. Wild-type (WS), *clca-2*, complemented line (CLCa-Ox (*clca-2*))*, NRT2.7* over-expressing line in *clca-2* (NRT2.7-Ox (*clca-2*)) and the two *NRT2.7* over-expressing lines, NRT2.7-Ox1 (WS) and NRT2.7-Ox2 (WS), were grown on vertical Petri dishes on a medium 2 mM nitrate for 12 days. Uptake experiments consisted of four to five measurements of $^{15}NO_3^-$ for 5 min in a liquid medium containing 6 mM (A=HATS+LATS) or 0.2 mM (B=HATS). The activity of LATS (C) was obtained by subtracting the activity measured at 0.2 mM to the one at 6 mM (see material and methods section for details). Presented data are averages of three independent experiments (n=3-5 by independent experiment). Bars represents means ± SEM. Different letters at the top of the bars represent significant difference (p value<0.05).

Fig. 4. Effect of limiting N supply on the phenotype of *NRT2.7* over-expressers. Wild-type (WS) and over-expressing lines, NRT2.7-Ox1 (WS) and NRT2.7-Ox2 (WS), were

grown in petri dishes for 12 days on 0.2 mM nitrate. Shoot fresh weight (A), total root length (B) shoot and root nitrate contents (C, D) were measured as described in Figure 1. C/N ratios, percentages of N and C contents were measured in shoot (E) and root (F) using an elementary analyser. Data are averages of three to four measurements (bars being the standard errors). Different letters shows significant differences between genotypes ($p < 0.05$).

Fig. 5. Nitrate efflux from mesophyll vacuoles is increased in NRT2.7-Ox (*clca-2*) expressing plants compared to clca-2 knock-outs. A. Representative whole-vacuole currents measured in vacuoles extracted from *clca-2* knock-out mutant (left) and *NRT2.7-Ox* (*clca-2*) transgenic line (right) in presence of nitrate only in the vacuolar lumen (200 mM NO_3^- pH 5.5 in the vacuolar buffer, 19.2 mM Cl^- in the cytosolic buffer, see Materials and Methods). Currents were evoked in response to 2 s voltage pulses ranging from -77 mV to +83 mV in +20 mV steps, followed by a tail pulse at +33 mV, with a holding potential at -17 mV. B. Box plot of the mean current density (pA/pF) measured at different positive (left) and negative (right) membrane potentials in vacuoles from *NRT2.7-Ox* (*clca-2*) (vertical diamonds line *NRT2.7-Ox1* (*clca-2*) n=11; horizontal diamonds line *NRT2.7-Ox2* (*clca-2*) n=5) and *clca-2* (n=10) knock-out plants (ionic conditions as in A). Boxes defines the first and the third quartile and the median, the whiskers represent the ninth and the ninety-first percentiles. The two asterisks indicate a statistical significance at 2% level with a Mann-Whitney test.

Fig. 1. Growth effect of *NRT2.7* overexpression in *clca-2* and wild-type backgrounds. Plant were grown on 2 mM nitrate for 12 days. Shoot fresh weight (A, B) and total root length (C, D) were measured on wild-type (WS), *clca-2*, complemented line (*CLCa-Ox* (*clca-2*)) and supplemented line (*NRT2.7-Ox* (*clca-2*)) (A, C) or WS and the two *NRT2.7* over-expressing lines, *NRT2.7-Ox1* (WS) and *NRT2.7-Ox2* (WS) (B, D). The number of samples were 4-5 for shoot fresh weight (one sample = a pool of 3-4 plants) when n=6-17 for total root length on individual plants. Presented data are means ± SEM. Different letters at the top of the bars represent significant difference (p value<0.05).

Fig. 2. Nitrate content of *NRT2.7* overexpression in *clca-2* and wild-type backgrounds. Wild-type (WS), *clca-2*, complemented line (*CLCa-Ox* (*clca-2*)) and *NRT2.7* over-expressing line in *clca-2* (*NRT2.7-Ox* (*clca-2*)) (A, C) or WS and the two *NRT2.7* over-expressing lines, *NRT2.7-Ox1* (WS) and *NRT2.7-Ox2* (WS) (B, D), were grown on 2 mM nitrate. After 12 days on the medium, shoot (A, B) and root (C, D) nitrate contents were measured. Two to three biological replicates were performed on each genotype, one sample corresponding to a pool of at least 50 plants. Presented data are means ± SEM. Different letters at the top of the bars represent significant difference (p value<0.05).

Fig. 3. Effect of *NRT2.7* overexpression in *clca-2* and wild-type backgrounds on nitrate uptake activities. Wild-type (WS), *clca-2*, complemented line (CLCa-Ox (*clca-2*)), *NRT2.7* over-expressing line in *clca-2* (NRT2.7-Ox (*clca-2*)) and the two *NRT2.7* over-expressing lines, NRT2.7-Ox1 (WS) and NRT2.7-Ox2 (WS), were grown on vertical Petri dishes on a medium 2 mM nitrate for 12 days. Uptake experiments consisted of four to five measurements of $^{15}NO_3^-$ for 5 min in a liquid medium containing 6 mM (A=HATS+LATS) or 0.2 mM (B=HATS). The activity of LATS (C) was obtained by subtracting the activity measured at 0.2 mM to the one at 6 mM (see material and methods section for details). Presented data are averages of three independent experiments (n=3-5 by independent experiment). Bars represents means ± SEM. Different letters at the top of the bars represent significant difference (p value<0.05).

Fig. 4. Effect of limiting N supply on the phenotype of *NRT2.7* over-expressers. Wild-type (WS) and over-expressing lines, NRT2.7-Ox1 (WS) and NRT2.7-Ox2 (WS), were grown in petri dishes for 12 days on 0.2 mM nitrate. Shoot fresh weight (A), total root length (B) shoot and root nitrate contents (C, D) were measured as described in Figure 1. C/N ratios, percentages of N and C contents were measured in shoot (E) and root (F) using an elementary analyser. Data are averages of three to four measurements (bars being the standard errors). Different letters shows significant differences between genotypes (p<0.05).

Fig. 5. Nitrate efflux from mesophyll vacuoles is increased in NRT2.7-Ox (*clca-2*) expressing plants compared to clca-2 knock-outs. A. Representative whole-vacuole currents measured in vacuoles extracted from *clca-2* knock-out mutant (left) and *NRT2.7-Ox* (*clca-2*) transgenic line (right) in presence of nitrate only in the vacuolar lumen (200 mM NO_3^- pH 5.5 in the vacuolar buffer, 19.2 mM Cl^- in the cytosolic buffer, see Materials and Methods). Currents were evoked in response to 2 s voltage pulses ranging from -77 mV to +83 mV in +20 mV steps, followed by a tail pulse at +33 mV, with a holding potential at -17 mV. B. Box plot of the mean current density (pA/pF) measured at different positive (left) and negative (right) membrane potentials in vacuoles from *NRT2.7-Ox* (*clca-2*) (vertical diamonds line *NRT2.7-Ox1* (*clca-2*) n=11; horizontal diamonds line *NRT2.7-Ox2* (*clca-2*) n=5) and *clca-2* (n=10) knock-out plants (ionic conditions as in A). Boxes defines the first and the third quartile and the median, the whiskers represent the ninth and the ninety-first percentiles. The two asterisks indicate a statistical significance at 2% level with a Mann-Whitney test.