Long-term Culturing of *Pseudomonas aeruginosa* in Static, Minimal Nutrient Medium

Results in Increased Pyocyanin Production, Reduced Biofilm Production, and Loss of

Motility

Rhiannon E. Cecil[1], Elana Ornelas[1], Anh Phan[1], Nahui Olin Medina-Chavez[2], Michael

Travisano[2], Deborah R. Yoder-Himes[1]

[1] Department of Biology, University of Louisville, Louisville, KY, U.S.A. 40292

[2] Department of Ecology, Evolution and Behavior, University of Minnesota, St. Paul, MN,

U.S.A 55108

<u>Abstract:</u>

Pseudomonas aeruginosa is a multidrug-resistant opportunistic human pathogen that can

survive in many natural and anthropogenic environments. It is a leading cause of morbidity in

individuals with cystic fibrosis and is one of the most prevalent pathogens associated with

nosocomial infections in the United States. It has been shown that this organism can survive and

persist in low nutrient environments, such as sink drains. How adaptation to these types of

environments influences the phenotypic traits of this organism has not been well studied. Here

we implemented an experimental evolution system in which six strains of *P. aeruginosa* were

subjected to low nutrient conditions over the course of 12-weeks and assessed phenotypic and

genotypic changes that occurred as a result of adaptation to such environments. We observed that

adaptation to low nutrient environments resulted in decreased generation time, reduced cell size,

reduced biofilm formation, increased pyocyanin production, and decreased motility for some of

the strains. Further, some of the evolved isolates were significantly more virulent/competitive

against a phagocytic predator. This study is significant as it allows us to predict how this

organism will evolve in hospital and domestic environments and can help us improve treatment

options for patients.

Importance:

Human commensal and pathogenic organisms undergo dynamic cycles across human and non-

human environments. Despite the crucial implications for human health, the understanding of

bacterial adaptations to these diverse environments and their subsequent impact on human-

bacterial interactions remains underexplored. This study shows how *P. aeruginosa*, an

opportunistic human pathogen, adapts phenotypically in response to a shift from high nutrients

(like those found in the human body) to low nutrients (like those found in many other

environments, like sink drains). This work also shows that, in some cases, resistance to

predatory forces can evolve in the absence of a predator. This work is important as it contributes

to the growing body of knowledge concerning how external, non-host-related abiotic conditions

influence host-pathogen interactions.

Introduction:

Adaptations to environments with fluctuating nutrients, otherwise known as feast and

famine cycles, are vitally important for microbial survival in various environmental niches

including natural environments such as soil, aquatic, and the intestinal and respiratory tract of

host organisms; as well as anthropogenic environments such as drains and hospital equipment

(1). Adaptations employed by some microorganisms in response to such environments, such as

the stringent response (1-3), cellular dormancy via reduced cell growth and the formation of

spores (4, 5), and metabolic shifting (fermentation vs respiration) (6, 7), are already well-studied.

Our understanding of how these adaptations evolve in opportunistic human pathogens and the

implications such adaptations have on virulence potential is relatively understudied. Uncovering

the mechanisms behind these adaptations holds significant potential for ecological, evolutionary,

environmental, industrial, and pharmaceutical applications. For example, these insights can be

leveraged to better understand microbial diversity, microbial food webs, speciation, the evolution

of bacterial communities, improve bioremediation techniques for cleaning polluted

environments, develop more efficient industrial processes for biofuel production, and even

design new antibiotics by targeting the pathways crucial for bacterial survival under nutrient

limitations.

P. aeruginosa is a member of the ESKAPE group of bacterial pathogens (8). It is an

opportunistic pathogen and is known for its intrinsic multidrug resistance, genomic plasticity,

ability to survive and form biofilms on abiotic surfaces such as hospital equipment, and ability to

evade the immune systems of humans, plants, and non-human animals (9-13). *P. aeruginosa*

expresses with a wide arsenal of phenotypes that serve as a means of offensive and defensive

protection against competitors and predators in the environment (e.g., protozoa, nematodes)

including pyocyanin production, biofilm formation, and T3SS expression (14-16). Many of these

phenotypes coincidentally act as virulence factors should the bacterium encounter a human host.

P. aeruginosa is a well-studied organism and how it adapts to host-associated niches, such as the

nutrient-rich cystic fibrosis lung, has been researched in detail (17); however, how this organism

evolves when forced to adapt to a low-nutrient, non-host associated environment has not been

thoroughly investigated. Furthermore, comparisons between adaptations to such an environment

between *P. aeruginosa* isolates from clinical origin and those from an environmental origin

remains under-explored with many studies utilizing clinical isolates and excluding environmental

isolates (18). Understanding how *P. aeruginosa* isolates, from clinical and environmental origin,

adapt to survive in a low-nutrient medium could increase our understanding of how *P.*

aeruginosa adapts in other low nutrient environments including sink drains, intubation tubes and

other types of hospital equipment, human tissues, and other such reservoirs where this organism

is able to thrive.

Experimental evolution provides a powerful means of understanding fundamental

evolutionary processes and can also be used to increase our understanding of how virulence

factors in opportunistic pathogens arise or change over time (19, 20). In this study, we used an

experimental evolution system to identify how six *P. aeruginosa* strains (four clinical strains and

two environmental strains) adapted to a low-nutrient, static environment over the course of

multiple generations. Whether adaptations to this type of environment would result in changes to

phenotypes commonly associated with growth and/or pathogenicity was then explored. We also

sought to determine if long-term culture in a low nutrient environment under static conditions

would result in increased or decreased fitness against a phagocytic predator, *Acanthamoeba*

castellanii, which would suggest that adaptation to such environments could result in increased

virulence in other organisms.

<u>Materials and Methods</u>

<u>Bacterial strains and culture maintenance</u>

P. aeruginosa strains used in this study were chosen based on their colony phenotype,

pyocin profile, and source of isolation. Strain details are listed in Table 1. Cells were routinely

maintained on LB (Lennox) agar plates at room temperature (22°C) or grown in LB (Lennox)

broth. Prior to initiation of an experiment, a single colony would be transferred from a LB agar

plate to either glass test tubes containing 5 mL of LB broth, 96-well deep plates containing 1%

HL5 medium (per L: 0.5 g KH_4PO_4; 0.5 g Na_4HPO_4; 7 g yeast extract; 14 g proteose peptone,

13.5 g glucose; purchased premixed from Formedium, Catalog #HLG0102), or 96-well

polystyrene plates containing 180 µL of LB broth and incubated overnight at 37°C with or

without aeration as indicated. When indicated, mid-log cultures were generated from overnight

liquid culture by transferring 100 µL of overnight culture to 5 mL of sterile LB broth and

incubated at 37°C with aeration for 90 minutes.

<u>Experimental Evolution Methods</u>

Each strain was cultured in 8 mL of 1% HL5 in 6-well tissue culture plates. The plates

were incubated at room temperature under static conditions. Every 2 to 3 days the medium was

removed with vacuum suction and replaced with 8 mL of fresh sterile 1% HL5 medium. The

evolving populations were also passaged every 7 days by removing the supernatant, adding 4 mL

of sterile 1% HL5, scraped with a cell scraper to dislodge the adherent cells. The suspended cells

were transferred to a sterile 15 mL conical tube and centrifuged at 6,976 x *g* for 15 minutes at

room temperature. The pellets were resuspended in 1 mL of 1% HL5 and 500 µl was added to 8

mL of sterile 1% HL5 in fresh 6-well tissue culture plates. A strain collection was generated at

bi-weekly time points by transferring 500 µL of culture to 500 µL of sterile LB+40% glycerol

(glycerol 20% final concentration) with subsequent storage at -80°C. At each dilution step,

lineages were tested for contamination via colony PCR for pyocin biosynthesis genes. Pyocins

are proteins exclusively produced by *P. aeruginosa.* The pyocin gene profiles were used to

ensure that no cross contamination occurred between the strains throughout the experimental

evolution via colony PCR using published primer sequences for the specific pyocin genes (21).

Three biological replicate lineages for each strain were evolved for 12 weeks total time.

On the rare occasion that a deviation of expected pyocin profiles was observed for any

given gene, the pyocin profiles were examined in the following week determine if the trend was

conserved. In all cases, the pyocin profiles returned to the expected pattern. Two strains that

were originally included in the dataset (not shown) were removed due to repeated loss of pyocin

gene positive results, which could result from contamination or simply selection against these

genes. However, we note that the final, analyzed isolates used in this study (the 12-week isolates)

all matched the respective ancestral pyocin profile.

Generation time

Overnight cultures of ancestral and evolved isolates were diluted (1:100 B83098, 17-047,

17-055 or 1:200 for B84725, PA3, and B80427) into 15 mL in 125-ml baffled flasks. Flasks

were incubated with shaking at 37°C. Every hour, the O.D.$_{600}$ was measured from 1 mL of

culture and plotted in GraphPad Prism v 5.04. At time points within exponential growth phase,

samples were taken for survival analysis via serial dilution and drip plating on LB square petri

dishes. Average bacterial concentrations were calculated for each of three biological replicates

for each ancestral and evolved isolate.

<u>Cell Size</u>

Overnight cultures of ancestral and evolved isolates were generated in 5 mL of LB broth

in glass test tubes with aeration. 20 µL of overnight culture was transferred onto a glass

microscope slide along with a drop (~20 µL) of 1% congo red staining solution. The mixture was

smeared across the slide using a coverslip then allowed to dry for 5 minutes in a biosafety

cabinet. The slides were imaged under a 1000X magnification light microscope with an ocular

micrometer for scale. ImageJ was used to quantify the length of cells. At least 100 cells from at

least two independent experiments (n > 200 for each isolate/strain) each were measured.

<u>Biofilm biomass quantification</u>

Biofilms were quantitated via a slightly modified version of a previously published

protocol (22). Briefly, overnight cultures of ancestral and evolved lineages in replicates of four

of each isolate were generated in 180 µL of LB in 96-well dilution plates without aeration.

Biofilms were set-up by first using a 96-well transfer apparatus to transfer inocula of overnight

cultures to fresh 96-well PVC plates containing 100 µL of 1% HL5. The inoculated plates were

sealed with porous adhesive film (VWR, 60941-086), loosely wrapped with aluminum foil, and

then placed in a humidity chamber with sterile water at the bottom. The lid of the humidity

chamber was loosely sealed to allow for air flow while also maintaining a moist environment.

The plates were incubated at 37°C for 48 hours or 7 days to allow for biofilm formation.

Biofilms were washed two times with water then 125 µL of 0.1% crystal violet was added to

each well and the plates were allowed to incubate at room temperature for 10 minutes.

Afterwards, the biofilms were washed four more times with water. Then 200 µL of 30% acetic

acid solution was added to each well and allowed to incubate at room temperature for 15

minutes. A 96-well plate reader was used to quantify crystal violet in solution at an optical

density of 500 nm.

<u>Pyocyanin and pyoverdine production</u>

Pigment levels from *P. aeruginosa* strains were determined using protocols modified

from a previously published study (23). Overnight cultures were generated in 5 mL of LB broth

with aeration in glass test tubes. 100 µL of overnight culture was transferred to 1 mL of LB broth

in 24-well plates. The plates were sealed with porous adhesive film (VWR, 60941-086) and

incubated at 37°C for 48 hours under static conditions. Post-incubation, 1 mL of each sample

was transferred from the 24-well plates to 1.5 mL microfuge tubes. The tubes were then

centrifuged at 21,130 x *g* for 30 minutes to pellet the bacteria, and 100 µL of supernatant from

each culture was transferred to either 96-well clear plates (VWR, #10861-562) for pyocyanin

quantitation or 96-well black plates (Greiner, #655077) for pyoverdine. A 96-well plate reader

was used to quantify the amount of pyocyanin (absorbance at 686 nm) or pyoverdine

(fluorescence 395ex/460em).

<u>Motility assays</u>

The motility protocol used in this study was modified from the motility assay described

in (23). Overnight cultures of ancestral and evolved isolates (each n = 3) were grown in 5 mL of

LB broth in glass test tubes with aeration. Briefly, a 2 – 10 µL pipet tip was dipped in overnight

culture then was used to stab approximately half-way deep into the motility agar (10 mL of 0.3%

LB agar in 6-well plates). The inoculated plates were incubated at room temperature for 24

hours. The diameter of spread was measured using a metric ruler to quantify swimming distance.

Isolates from each replicate lineage were tested at 12 weeks. For those that decreased, isolates

were then tested at 2, 3, 5, and 7 weeks but only until the time when motility appears to be

significantly reduced.

Acanthamoeba castellanii culture and maintenance

A. castellanii strain 30010 was obtained from the ATCC and cells were routinely

maintained at room temperature in T-75 tissue culture flasks containing 15 mL of 100% HL5

growth medium. The cells were passaged routinely when they reached confluence,

approximately every 3 days, using a cell scraper to dislodge the cells from the bottom of the

flask, then removing all but 1 mL of culture from the flask and replacing with 14 mL of sterile

100% HL5. Cell stock cultures were maintained up to 4 passages.

Co-culture experiment with *Acanthamoeba castellanii*

After reaching confluence, approximately 6 x 10^3 *A. castellanii* cells, based on

hemocytometer counts, were transferred to wells of 6-well tissue culture plates along with 4 mL

of 100% HL5 medium. The cells were incubated at room temperature for 1 hour to allow the

cells time to anneal to the surface of the wells in the 6-well tissue culture plates. The 4 mL of

100% HL5 was then removed, and the wells were washed 2 times with sterile 1X PBS then 8 mL

of 1% HL5 was added to the wells. Overnight cultures of ancestral *P. aeruginosa* and one of the

evolved lineages of each *P. aeruginosa* strain were generated in triplicate in 5 mL of LB broth.

Mid-log cultures were generated as previously described above until an O.D.$_{600}$ between 0.8 and

1.0 was reached for all strains. *P. aeruginosa* cells were added to the 6-well tissue culture plates

in a ratio of 1 *P. aeruginosa* cell to 10 *A. castellanii* cells in 3 biological replicates per *P.*

aeruginosa isolate.

The plates were incubated at room temperature for 16 days. Every three days the medium

was replaced by using vacuum suction and replaced with 8 mL of sterile 1% HL5. *A. castellanii*

cells were enumerated on days 0, 3, 6, 8, 11, 13, and 16 via direct cell count at 400X

magnification. *P. aeruginosa* cells were enumerated on day 16 via serial dilution and plating on

LB agar plates.

<u>Genome sequencing</u>

One mL of each ancestral and one of the evolved lineages for B80398, B80427, 17-047,

and 17-055 was grown in LB medium and genomic DNA was isolated using a commercial kit

(Promega Wizard Genomic DNA Purification Kit) according to the manufacturer's instructions.

DNA was shipped on dry ice to Novogene for genome sequencing. A total amount of 0.2 μg

DNA per sample was used as input material for the DNA library preparations. Briefly, genomic

DNA sample was fragmented by sonication to a size of 350 bp. Then DNA fragments were

endpolished, A-tailed, and ligated with the full-length adapter for Illumina sequencing, followed

by further size selection and PCR amplification. After PCR products were purified by AMPure

XP system (Beverly, USA). Subsequently, library quality was assessed on the Agilent 5400

system (Agilent, USA) and quantified by QPCR (1.5 nM). The qualified libraries were pooled

and sequenced on Illumina NovaSeq platforms with PE150 strategy at ~100X coverage,

according to effective library concentration and data amount required.

<u>Variant Calling Analysis</u>

To compare ancestral and evolved lineages, and measure punctual changes related to

adaptation, we performed variant calling analysis. After sequencing, raw read quality control was

performed using FastQC v.0.20.0 (24) to identify adapters and then remove them using Cutadapt

v4.2 (25). We then proceed to trim sequences with low quality bases using Trimmomatic (26)

with the following parameters LEADING:10 TRAILING:3 SLIDINGWINDOW:4:30

MINLEN:35. Paired-end high quality reads were mapped to the reference genome, *P.*

aeruginosa UCBPP-PA14, using Burrows-Wheeler Aligner (BWA) v.0.7.17 (27). Post-

alignment processing, including sorting, indexing, and duplicate removal, was conducted with

Samtools (28) to ensure clean and accurate input data for variant analysis. Base recalibration and

indel realignment were performed using GATK's Best Practices workflow to correct systematic

errors introduced during sequencing (29-31). Variants were called using GATK's

HaplotypeCaller, configured to detect both SNPs and indels across the entire genome. Stringent

filtering criteria were applied to identify high-confidence variants, including minimum mapping

quality scores, read depth thresholds, and variant quality scores. SnpEff (32) was used to identify

the genes where the variants were located or the closest genes to a variant and to predict the

effects and impacts of the variants on the genes and protein products.

Data Availability

Raw sequencing data can be found in NCBI at the following BioProject ID: PRJNA1194290

Statistics:

Data was analyzed with one-way ANOVA with Tukey's or Dunnett's post-test or with t-tests as described below in GraphPad Prism 5.0.4 as indicated. Spearman correlations were calculated in GraphPad Prism 9.0.

Table 1. Strains used in study and their pyocin profile.

Strain	Origin	Colony Phenotype	S1-type	S2-type	S1S2 immunity	R-type	F-type	Reference
PA B80398	*CF sputum	Normal Colony	1 [a]	0	0	1	0	(33)
PA B80427	CF sputum	Small Colony	0	1	1	1	1	(33)
PA B84725	CF sputum	Mucoid	0	0	0	1	1	(33)
PA3	CF sputum	Normal Colony	0	1	0	1	0	(33)
SRP 17-047	Bathroom sink drain	Normal Colony	1	0	1	1	0	(34-36)
SRP 17-055	Kitchen sink drain	Normal Colony	1	0	1	1	1	(34-36)

[a]1 indicates presence of the pyocin gene in the genome, 0 indicates absence from the genome.

*CF indicates isolates taken from sputum samples from individuals with cystic fibrosis

Results:

To test how a variety of *P. aeruginosa* phenotypes change in response to forced growth in a low-nutrient environment, we conducted a 12-week-long experimental evolution on six *P.*

aeruginosa strains. Four of these strains were of clinical origin (PA B80398, PA B80427, PA

B84725, and PA3) and two strains were of environmental origin (SRP 17-047 and SRP 17-055)

as they were isolated from household sink drains. These strains were chosen based on their

colony phenotypes, pigment production, and pyocin profiles and these strains represent a wide

range of *P. aeruginosa* morphotypes. Four of the strains (PA B80398, PA3, SRP 17-047, and

SRP 17-055) present with "normal" colony morphology. The normal colony phenotype of *P.*

aeruginosa is frequently isolated from acute infections or environmental samples and is

associated with more aggressive virulence factors such as pyocyin production and T3SS as well

as increased motility and a predisposition to planktonic lifestyle (37-39). The other two strains,

PA B80427 and PA B84725, are small colony and mucoid respectively, both of which are often

isolated from chronic infections and are not known to be isolated from environmental samples.

The small colony and mucoid phenotypes commonly present with defensive virulence factors to

avoid detection by the host's immune system such as reduction in motility and increased biofilm

formation (38, 40). The mucoid phenotype also overproduces the carbohydrate polymer alginate

making them difficult to phagocytose (41).

An experimental evolution was designed to mimic the conditions that *P. aeruginosa*

would encounter in a non-host environment, such as the soil or an abiotic surface such as a sink

drain. To mimic these environments, *P. aeruginosa* was cultured over time under static

conditions (to encourage niche specialization) in a low-nutrient medium (1% HL5 with glucose),

which contains dilute yeast extract and peptone along with sodium and potassium buffers. This

medium was chosen as *P. aeruginosa* readily utilizes glucose and amino acids as carbon sources,

amino acids stimulate biofilm formation and swarming activities, and for subsequent amoebae

studies (33, 42-44). Every two days, the medium was removed via vacuum suction [simulating

the flow of liquid down a sink drain (i.e. frequent mechanical removal)]. This removes most of

the planktonic cells, thus selecting for cells that can attach to the surface of the wells. Every 7

days the medium was removed, and the adherent cells were scraped from the wells and diluted

back into sterile 6-well tissue culture plates containing sterile 1% HL5. This process was

continued for 12 weeks generating three biological replicate lineages of the six *P. aeruginosa*

strains. Interestingly, the normal and mucoid colony strains remained the same in terms of

colony morphology over the course of the experiment. However, all three replicate lineages of

the small colony variant, B80427, underwent a morphological shift to a normal colony

phenotype during the evolution experiment. All the starting ancestral isolates and isolates from

the end of the evolution experiment (i.e. 12-week evolved isolates) were tested for a variety of

phenotypes associated with growth, long-term survival, or virulence.

Overall cellular changes during evolution in a low nutrient medium

During evolution experiments, an increase in fitness of the evolved lineage compared to

its ancestor usually occurs as the organisms adapt to their conditions. One global indicator of

fitness in bacteria is generation time. A decreased generation time in 1% HL5 at room

temperature would indicate that the evolved isolates have an increased fitness in this

environment compared to their ancestors and thus this measurement serves as a control for the

experimental evolution. We assessed growth of all ancestral and representative isolates from

evolved lines in 1% HL5 over time and calculated the generation times for each isolate during

exponential phase growth. Both generation times and growth curves in 1% HL5 were used to

assess overall growth patterns at two temperatures, 22°C which is the temperature used for the

experimental evolution and at 37°C, which is the optimal growth temperature for most *P.*

aeruginosa strains. At 22°C, generation times were on the orders of hours, not minutes, due to

the low nutrient conditions as well as the temperature. Most evolved isolates from three strains,

B80427, PA3, and 17-047, showed a decreased generation time compared to their associated

ancestral strain. This suggests that they adapted to the conditions of the experimental evolution.

Two individual isolates from the other strains, in B80398 and B84725, showed an increase in

generation times while the other two evolved isolates did not significantly change compared to

their ancestral parent. One strain, 17-055 did not show a change in generation times for any of

the evolved isolates.

We then assessed whether evolving in 1% HL5 for 12 weeks at room temperature also

conveyed an advantage at a higher temperature due to simply the medium alone. We calculated

the generation time for all strains grown in 1% HL5 at 37°C, generally considered the optimal

temperature for *P. aeruginosa* (though we note that we did not identify the true optimal

temperatures for all strains in this study). Three strains, B84725, PA3, and 17-055, have the

majority of evolved isolates showing a reduced generation time while almost all of the others are

not statistically different than their respective ancestors. Comparing the two temperature studies,

we also observed that *P. aeruginosa* B3098 evolved line 2 showed an increase in generation time

(i.e. slower growth) at both 22° and 37°, suggesting that this isolate did not adapt to the low

nutrient conditions of the experimental evolution for some reason. Interestingly, the B84725

showed a different pattern between the two temperatures with an increased generation time at

22°C and a slightly decreased generation time at 37°C. This particular strain is the only mucoid

isolate in our strain panel and it has previously been shown that the expression of alginate

biosynthesis genes is higher at 20°C than 37°C (45).

By examining the growth curves of each lineage, we observed that there were many

significant differences between evolved and ancestral lineages at 22°C but not at 37°C based on

repeated measures one-way ANOVAs (Fig. S1). Taken all the growth metrics together, this

suggests that many of the genetic changes that occurred during the evolution served to optimize

growth at room temperature and some of these changes may have additional benefits, even at

higher temperatures.

Cell size is also linked to metabolic fitness as an increased surface volume ratio results in

amplified nutrient diffusion into the cell resulting in increased metabolic efficiency, especially in

low nutrient conditions (46, 47). We hypothesized that an overall decrease in cell size would be

observed in the evolved isolates as the low-nutrient concentration of the growth medium would

encourage a smaller surface area to volume ratio. Many of the evolved isolates appeared to have

reduced cell length compared to the ancestor at 22°C (Fig. 2). A few evolved isolates that

increased in cell length at this temperature, including B80427 evolved lineage 1 (E1) and

B84725 evolved lineage 3 (E3). Curiously, the evolved lineages of PA3 all increased in size

compared to the ancestor; however, this ancestral strain was relatively small to begin with, which

might imply that PA3 was already adapted to lower nutrient conditions.   At 37°C, most of the

differences in cell length were not observed. However, all evolved lineages of B84725 showed a

decrease in cell length while a few evolved isolates, PA3 E3, 17-047 E2 and E3, 17-055 E2, had

a modest increase in cell length. Our cell size data overall suggests that a small cell size was not

strongly selected for but that individual lineages had slight increases or decreases in cell length.

This data is confirmed when aggregating the data from the evolved lineages as well (Fig. S2).

Another *P. aeruginosa* phenotype assessed was biofilm formation. Based on the design of

our experimental evolution, we hypothesized that the evolved lineages would be superior biofilm

formers compared to their ancestors because our experimental evolution design strongly selected

for cells that were adhered to the tissue culture surface. We assessed biofilm formation with

crystal violet assays which provide a broad quantitative means to assess the volume of biofilm

biomass produced.

Surprisingly, only two strains, PA B80398 and PA3, exhibited significantly different

biofilm densities compared to their ancestor strains (Fig. 3). In both instances, the evolved

lineages exhibited significantly reduced biofilm biomass compared to their ancestors, though the

overall degree of loss is quite small (5-25%). This result is consistent when considering each

evolved lineage separately as all lineages showed reduced biofilm biomass (Fig. S3) which

suggests that this change was not driven by data from a single lineage but likely represents a true

shift in the phenotype for these strains. All the remaining evolved lineages tested were not

significantly different from their ancestor with respect to biofilm formation (Fig. 3 and S3).

Individual molecular phenotypes

Pyocyanin is a redox-active toxin secreted by *P. aeruginosa.* It is toxic to eukaryotic cells

and provides *P. aeruginosa* protection against predators, but it also serves as a secondary

metabolite for *P. aeruginosa* and is associated with promoting survival in low-oxygen and low-

nutrient environments (48-50). We hypothesized that the evolved lineages would produce

significantly more pyocyanin than their ancestors due to the low-nutrient conditions present in

our experimental evolution system. Our hypothesis was supported for three of the strains: PA

B80427, SRP 17-047, and SRP 17-055, of which there was a small but significant increase in

pyocyanin production in PA B80427 and SRP 17-047, and a large increase in SRP 17-055 (Fig.

4). Two strains, PA B80398 and PA B84725, did not exhibit significant differences in pyocyanin

production compared to their ancestors; however, two of the individual evolved lineages of PA

B80398 did produce more pyocyanin than their ancestor (Fig. S4). Interestingly, all three

evolved lineages of PA3 produced significantly less pyocyanin compared to their ancestor

suggesting that pyocyanin production was not selected for in this strain. The reasons for this

remain unclear. From this data, it suggests that for most of our *P. aeruginosa* strains pyocyanin

production was selected for in our experimental evolution system and increased pyocyanin

increased the fitness of the organisms in these low-nutrient environments.

Pyoverdine is a siderophore secreted by *P. aeruginosa* which functions to sequester iron

from other cells as well as the environment (51). The 1% HL5 used for the experimental

evolution is undoubtedly low in most macronutrients. However, it likely does contain some iron

as it contains yeast extract and protease peptone but whether the concentration was sufficient for

P. aeruginosa growth is unknown. Therefore, we could not make a hypothesis on whether

pyoverdine production would change over time in the evolved isolates. Pyoverdine is highly

implicated in *P. aeruginosa* pathogenicity (52) so pyoverdine production was assessed using

spectrophotometry in the ancestral and evolved lineages.

Pyoverdine remained unchanged in five of the six strains tested (Fig. 5). Only one of the

evolved lineages of the environmental isolate, SRP 17-047 produced significantly less

pyoverdine than its ancestor, while the other two lineages of 17-047 were not significantly

different (Fig. S5). This suggests that pyoverdine was likely neither selected for nor against

under the conditions in our experimental evolution system.

The last molecular phenotype tested was swimming motility. We hypothesized that there

would be a reduction in motility in our evolved lineages as our experimental evolution design

selected against planktonic cells. The swimming abilities of the ancestral isolates were variable

from completely non-motile to highly motile (Fig. 5). There was a significant reduction in

swimming motility observed for almost all the evolved lineages compared to their ancestors for

B80398, PA3, 17-047, and 17-055. The other two strains, B80427 and B84725, showed very

small changes in motility compared to their ancestors; however, the ancestors for these two

strains were either amotile (B84725), or only weakly motile (B80427), under these conditions.

Surprisingly, two of the evolved lineages, E1 and E3 of B80427, displayed small but

significantly greater motility than the ancestor B80427 strain. There was no correlation between

biofilm biomass and motility for all strains aggregated (Spearman r, p=0.5720) or for any strain

individually.

We tested the earlier isolates of the four strains that showed the greatest reduction in

motility over the course of the experiment, B80398, PA3, 17-047, and 17-055, to determine

when motility was lost by selecting isolates from earlier time points in the experimental

evolution. Isolates from the second week of the experimental evolution were tested first, then

isolates from additional weeks were tested as needed. We found that motility was lost at week 2

for 17-047, week 3 for 17-055, week 5 for PA3, and week 7 for B80398 (Fig. S5). Together

these data suggest that for strains that were highly motile at the start of the experiment, motility

was selected against and loss of 49-90% of motility between 2 and 7 weeks of culture in low-

nutrient medium. The degree and timing of the loss varied for each motile strain.

<u>Host-associated phenotypes over the course of evolution in a low nutrient environment</u>

We also sought to determine if long-term adaptation to a low-nutrient environment would

result in an increased or decreased ability to survive and/or kill a phagocytic predator as this

could be a strong indicator of how *P. aeruginosa* virulence can increase in low-nutrient

environments. To test this, we co-cultured the ancestral isolates and one evolved lineage of each

strain with the phagocytic amoeba species *A. castellanii* over time to examine the survival of

both organisms. *A. castellanii,* an avid predator of *P. aeruginosa,* is a model organism for

studying phagocytosis and is often used as a proxy to study mammalian macrophages engulfment

(53-56). *A. castellanii* exists in two forms, trophozoites (the actively growing and metabolizing

form) and cysts (dormant).

We examined trophozoite and *P. aeruginosa* survival for individual ancestral and evolved

strains. After 16 days of co-culture, *A. castellanii* trophozoite survival was significantly reduced

when co-cultured with the evolved lineage of the clinical isolates B80398 and PA3 when

compared with the ancestral isolate of these strains (Fig. 6A). The remaining strains showed

neither a statistical increase nor decrease in the number of trophozoites found, perhaps due to the

large variability in the individual lineages for the most part. Trophozoite survival tended to be

reduced over time when in co-culture with the evolved lineage compared to the ancestral isolates

for most strains but this rarely reached statistical significance for most lineages/strains (Fig. S6).

The co-culture dynamics of the clinical strains PA3 and B80398 over time were particularly

interesting. Trophozoite levels in presence of the ancestral strains increased dramatically over

the 16 days of observation but trophozoite levels remained very low when co-cultured with the

evolved isolate of this strain (Fig. S6).

In comparison, all the *P. aeruginosa* isolates tested were able to survive in co-culture

with *A. castellanii* for at least 16 days (Fig. 6B). There were no significant differences between

the numbers of surviving evolved cells compared to ancestors at 16 days except for the clinical

isolate, PA3, for which the number of surviving cells at 16 days was significantly greater for the

evolved lineage than the ancestor. Therefore, ancestral and evolved lineages of *P. aeruginosa*

were able to persist in the presence of amoebae. *P. aeruginosa* cells were only enumerated on

day 16 as we were able to confirm the presence of live *P. aeruginosa* cells in the wells using

bright-field microscopy at each of the other time points.

<u>Genomic changes in the evolved lineages compared to ancestors</u>

To better understand the genetic mechanisms underlying the observed changes in

phenotype and to better identify other potential changes, the genomes of the ancestor and one of

the evolved lineages from four of the strains, B83098, B80427, 17-047, and 17-055, were

sequenced and compared. The genomes sizes ranged from 6.60 Mbp to 6.99 Mbp (average 6.81

Mbp) and the number of predicted genes was between 6149 and 6456 (average 6,318) (Table 2).

The number of SNPs in coding regions between ancestor and evolved isolates varied between

212 (17-055) and 312 (B80427).

Table 2. Genome analysis summary

Strain	Ancestor (A) or Evolved (line)	Mapped reads	Total reads	Mapping rate (%)	Average depth(X)	Coverage at least 1X (%)	Size (Mbp)	# predicted genes	# SNPs	# unique genes
B83098	A	9,766,129	11,003,760	88.75	187.1	96.11	6.89	6,410	254	66
B83098	Line 2	9,119,114	10,276,000	88.74	172.79	96.11	6.89	6,410		
B80427	A	8,921,569	9,610,204	92.83	168.34	98.18	6.60	6,149	312	47
B80427	Line 2	10,037,516	10,847,040	92.54	188.48	98.19	6.60	6,157		
17-047	A	8,676,544	9,777,748	88.74	162.52	96.65	6.83	6,253	263	62
17-047	Line 3	16,760,4	18,566,9	90.27	339.24	97.69	6.99	6,434		

		25	44							
17-055	A	8,866,855	10,048,604	88.24	174	97.67	6.76	6,281	W212	53
17-055	Line 2	9,461,172	10,879,064	86.97	182.51	96.65	6.92	6,456		

We identified coding genes that contained SNPs between ancestral and evolved lineages

(Fig. 7). There were 6 genes or homologous gene pairs that were mutated in all four evolved

lineages. They encoded: phenazine biosynthesis proteins PhzA2/PhzA1; Type VI secretion

system tip proteins VgrG/VgrG1; a short chain dehydrogenase; outer membrane protein OprM;

glutamate 5-kinase (involved in the urea cycle); and 3-carboxy-cis,cis-muconate cycloisomerase

(part of the protocatechuate pathway that is a central catabolic route for aromatic compounds). A

number of other genes were mutated in 3 of the four strains. Notably, genes that encode putative

virulence factors like VirS [a transcriptional regulator of virulence proteins] and TpsA/B proteins

[involved in two partner secretion of virulence proteins (57)], genes that encode histidine

biosynthesis proteins. There were no genes containing SNPs that were only shared between

B80398 and the two environmental strains, 17-047 and 17-055.

Additionally, SNPs that were shared between clinical strains or environmental strains

were also identified. The clinical isolates B80398 and B80427 had only two unique coding

regions with SNPs – a SDR oxidoreductase gene and a STY4528-family pathogenicity island

replication protein. The environmental strains showed SNPs in four different genes, encoding

acetylaldehyde 2, a NAD(P)H-dependent oxidoreductase, the 23S rRNA, and a gene that was

annotated as YycB, a uncharacterized transporter from *Bacillus subtilis*.

In the molecular phenotype analyses, motility showed the greatest level of changes

between ancestral and evolved lineages. We assessed the genomic DNA results and found that

only one strain, B80398, had SNPs in a known flagella-associated protein. FliR is a core part of

the FliP/FliQ/FliR complex that forms a proton motive force-dependent channel that exports

flagellar proteins across the cytoplasmic membrane (58) and has also been implicated in the

secretion of virulence factors (59) and in adhesion and biofilm formation (60) in other

organisms. DmlR, a transcription factor implicated in carbohydrate metabolism and flagellar

motility regulation contained SNPs in three of the four strains, B80398, B80427, and 17-047 (61)

though this association is only hypothetical. Evolved lineages of these strains may also have

SNPs in intergenic, regulatory regions or in other types of regulatory proteins that could explain

the changes in motility observed.

Discussion:

The adaptations that occur in response to long periods without sufficient food sources

could impact how bacteria grow (both in terms of how fast and how to divide their limited

resources), sense and respond to their environment, their lifestyle, or how they defend

themselves against predation. Culturing *P. aeruginosa* for 3 months under low-nutrient

conditions resulted in alterations to multiple phenotypes associated with fitness and/or virulence.

Some of these observed changes in phenotype were shared amongst most of the evolved

lineages, regardless of ancestral strain. These changes in phenotypes include faster generation

time, reduced cell size, increased pyocyanin production, and decreased motility. Since these

phenotypes were observed in most lineages, it could be inferred that these phenotypic alterations

lead to increased fitness in low nutrients and thus are likely to occur in these types of

environments. Two other phenotypes tested were biofilm formation and pyoverdine production,

and we observed variable changes between strains and within replicate evolved lineages and

within each strain. This indicates that these phenotypes were likely not as strongly selected for in

our experimental evolution system (62).

The environments where these particular *P. aeruginosa* strains were isolated likely had

higher levels of nutrients (i.e. the clinical isolates) or fluctuating levels of nutrients (household

drains). Therefore, the ancestral strains used in our experiments were likely not well adapted to

constant low nutrient conditions. Evolved lineages of three of the six strains tested exhibited

significantly faster growth compared to their ancestors (at either 22°C, 37°C, or both), indicating

that the evolved lineages became better adapted to the environment in which they were evolved

(Fig. 1). These results are consistent with results observed in other experimental evolution

systems such as those observed in the Long-Term Evolution Experiment (LTEE) of *Escherichia*

coli where the doubling times of the evolved lineages reduced from 55 minutes to 23 minutes in

the minimal nutrient medium used in the experiment after ~50,000 generations (62). In contrast,

a few of the evolved isolates (e.g. B80398 evolved line 2) exhibited significantly increased

generation times compared to their ancestor (Fig. 1). Surprisingly, we did not observe any direct

correlation between cell size and generation time suggesting that cell size does not play a factor

in doubling time for *P. aeruginosa* as it does for *E. coli* (46), at least not over the short-term

evolution experiment described here.

Our experimental evolution system theoretically was expected to have strongly selected

for increased biofilm formation. Surprisingly, evolved lineages in two of the six strains produced

significantly less biofilm biomass than their ancestors while the evolved lineages of the

remaining four strains were not significantly different from their ancestors (Fig. 3). One potential

explanation for this could be the low-nutrient culture medium used in the evolution experiment.

Biofilms require the production and secretion of polysaccharides and extracellular proteins to

form the biofilm matrix (63-65). These matrix components are undoubtedly metabolically costly

to produce as the responsibility for exopolysaccharide matrix generation has been shown to be a

shared responsibility within a biofilm in other species of bacteria such as *Bacillus subtilis* (66).

Thus, it is possible that the cells either acquired mutations that blocked the biosynthesis or export

of some of the necessary biofilm components – perhaps choosing to use the limited carbon for

growth rather than matrix production. In addition, the evolved cells could have used other

methods of attachment to the plate surface, such as pili or fimbriae, to avoid the expense of

secreting polysaccharides though no SNPs were found in the coding regions of known pilin or

fimbriae genes. Alternatives could include proteins like FliR, which has been shown to be

involved in biofilm and adhesion (60) or be due to SNPs in transcriptional factors. Thus, there

are several alternatives that could explain how *P. aeruginosa* could have attached to the surface

but not initiated polysaccharide secretion. The lack of increase in biofilm formation in low-

nutrient environments is similar to those observed in *P. aeruginosa* strains co-cultured long-term

with the amoebal predator, *A. castellanii,* in a minimal nutrient medium (67). In this latter study,

biofilm biomass was also found to decrease in *P. aeruginosa* strains cultured in monoculture as

well as in co-culture with *A. castellanii* in minimal medium for 42 days.

We expected that flagellar motility would be selected against in our experimental

evolution because planktonic cells were removed during every passage. A reduction in motility

was observed in the evolved lineages of strains where the ancestors exhibited high motility (Fig.

5). This result is similar to results observed in *P. aeruginosa* PAO1 cultured in artificial CF

medium for 120 days under aerated conditions both in monoculture and in the presence of phage

competitors (68). In our study, the evolved lineages of two strains, PA B80427 and PA B84725,

did not exhibit a reduction in motility, perhaps because the ancestors were poorly motile/amotile

and possibly do not have sufficient loci for evolution to act upon. We did not observe a

significant correlation, either negatively or positively, between biofilm formation and motility as

others have previously (69, 70). The reasons for this remain unclear though there have been a

few studies that also did not find correlations between these phenotypes in *P. aeruginosa*,

particularly when *P. aeruginosa* forms aggregates (71, 72), or suggest that the flagellar

dependence for biofilm formation may depend on the growth medium (73).

We also had hypothesized that pyocyanin production would be increased in the evolved

populations compared to their ancestors because outside of its toxic effects on host and other

prokaryotic cells, pyocyanin serves as a secondary metabolite, stabilizing the redox state of *P.*

aeruginosa and can act as a final electron acceptor in the electron transport chain (38). Increased

expression/production of pyocyanin should therefore increase metabolic fitness in low nutrient

environments. Pyocyanin has also been shown to play a role in pyruvate excretion and

metabolism under anoxic conditions which are common in the lung environment and in biofilms

(48-50). *P. aeruginosa* has been shown to be able to ferment excreted pyruvate to survive in

anoxic/low-nutrient conditions for at least 18 days (49). Our pyocyanin hypothesis was

supported for half of the evolved lineages when three of the six strains showed a significant

increase in pyocyanin production compared to their ancestors. Because our cultures were static,

it could be that increases in pyocyanin were observed for those strains that needed additional

electron acceptors for their particular electron transport chain while other strains did not.

In contrast, production of pyoverdine, a siderophore, was not significantly altered in

almost all evolved lineages compared to their ancestors. This result indicates that pyoverdine was

not selected for or against in our experimental evolution design. This result is in opposition to

results observed in previous studies which found that pyoverdine production decreased over the

course of 42 days in a minimal nutrient medium (67). It is possible that the 1% HL5 used in our

study contained more freely available iron and thus was not selected for or against in our

experimental evolution system compared to the supplemented M9 medium used in the previous

study.

In addition to identifying phenotypic changes that occurred in our evolved isolates, we

also sought to determine if adaptations to a low-nutrient environment could have implications on

virulence against host organisms. *A. castellanii* a bacterivorous amoebae species was identified

as a model organism to study phagocytosis because of its commonalities with phagocytic human

immune cells, such as macrophages (53-56). In our study, trophozoite survival was sometimes

decreased when co-cultured with evolved isolates compared to ancestors, particularly strains

B83098 and PA3, but never significantly increased when co-cultured with the ancestral isolates

(Figs. 6 & S6). Additionally, all the evolved lineages of *P. aeruginosa* were just as capable of

surviving predation from the amoebae predator as their ancestors were (Fig. 6B). One potential

reason that might underlie why *A. castellanii* trophozoites tended to survive better with the

ancestors than with the evolved lineages over time could be due to the reduced motility observed

in our evolved lineages. *Campylobacter jejuni, Pseudomonas fluorescens,* and *Proteus mirabilis*

flagellin are recognized by *A. castellanii* to initiate phagocytosis (74, 75). Thus, if there are

fewer/less flagella expressed in the evolved isolates, there would be less predation by *A.*

castellanii resulting in starvation of the amoebae predator when in co-culture with the evolved

lineages.

The genomes of the evolved and ancestral isolates showed many differences in the

presence of SNPs. Whether or not these SNPs actually relate to changes in the amino acid

structure of a protein or whether they influence the expression of these genes remains to be seen.

However, understanding the precise mutations that occurred can shed light on the genes where

natural selection may be occurring. We also note that individual isolates of evolved lines were

chosen for genome sequencing, and this may have limited the resolution of the SNPs that

occurred across the population. Previous studies have shown that long term starvation, nutrient

exhaustion, or feast-and-famine lifestyles can lead to increased genomic heterogeneity in

bacterial populations rather than communities dominated by one phenotype produced from

beneficial SNP(s) (76-78). Further, we observed evidence of mutational parallelism among the

strains after 12 weeks as was observed in (78), despite the feast and famine cycles somewhat

mirrored in our study were much shorter than the 100 day cycles used in that study. For example,

strains PA B80398 and PA 80427 showed mutations in the gene encoding the STY45280family

pathogenicity island replication protein and

In summary, a variety of *P. aeruginosa* phenotypes were observed to change over the

course of long periods with limited resources, but there was a great deal of variability in how *P.*

aeruginosa chose to invest its resources based on individual lineage data. This may not be

surprising, given the genomic plasticity within *P. aeruginosa* as a species. These results are

significant as they contribute to fundamental scientific knowledge into how *P. aeruginosa* might

evolve in other low nutrient environments (such as sink drains and faucets, hospital surfaces, or

freshwaters). This work could also yield insights into expected changes in *P. aeruginosa* over

time in low-nutrient environments and this could have implications for virulence properties or

treatment efficacy for human infections. It could help us to understand the mechanisms of

adaptation in such non-host environments (as in static cultures) and optimize sanitation

protocols, especially in healthcare settings, and/or designing surfaces or materials that inhibit

bacterial attachment and persistence. Lastly, this work could set the stage for future evolution

experiments in which *P. aeruginosa* could be evolved with predators, such as macrophage or

amoeba, to examine the effect of nutrient concentration on host-pathogen interactions.

Figure legends:

Figure 1. Relative generation time of ancestral and evolved *P. aeruginosa* isolates in 1% HL5.

Generation times were calculated for each strain growing at 22°C (panel A) or 37°C (panel B).

Grey bars represent the ancestral strains and colored bars represent the average of three replicates

of each evolved lineage. Data analyzed with one-way ANOVA with Dunnett's post-test

comparing all evolved lineages to the ancestor. Error bars represent standard deviation. *

$p<0.05$, ** $p<0.01$, *** $p<0.001$.

Figure 2. Cell size of ancestral and evolved lineages grown in 1%HL5 medium. Panels show

cell sizes for strains growing at (A) 22°C or (B) 37°C. Grey bars represent average values for the

ancestor strain while colored bars represent the average of three evolved isolates. An (A) in the

x-axis indicates the ancestral isolate, and (E) represents evolved isolates. Data analyzed with

one-way ANOVA with Dunnett's post-test comparing all evolved lineages to the ancestor. Error

bars represent standard deviation. * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

Figure 3. Biofilm formation of ancestral and evolved isolates in 1% HL5. Panels show crystal

violet retention after (A) 2 days or (B) 7 days of incubation. Note the different y-axis scales.

Grey bars represent ancestors and colored bars represent evolved isolates. Each bar represents

the average of three biological replicates for each isolate. An (A) in the x-axis indicates the

ancestral isolate, and (E) represents evolved isolates. Data analyzed with t-tests between the

ancestor and evolved lineages. Error bars represent standard deviation.   * $p<0.05$, ** $p<0.01$,

*** $p<0.001$.

Figure 4. Pigment production in ancestral and evolved *P. aeruginosa* isolates. Panels show (A)

pyocyanin or (B) pyoverdine production from 12-week-old evolved lineages compared to

ancestors. Grey bars represent ancestors and colored bars represent the evolved lineages. Each

bar represents the average of three biological replicates for each isolate. An (A) in the x-axis

indicates the ancestral isolate, and (E) represents evolved isolates. Data analyzed with t-tests

between the ancestor and evolved lineages. Error bars represent standard deviation. * $p<0.05$, **

$p<0.01$.

Figure 5. Swimming motility of ancestral and 12-week evolved isolates in 0.3% LB agar. Grey

bars represent ancestors and colored bars represent evolved isolates. Each bar represents the

average of three biological replicates for each isolate. The dashed line indicates the diameter of

the stab mark. An (A) in the x-axis indicates the ancestral isolate, and (E) represents evolved

isolates. Data analyzed with t-tests between the ancestor and evolved lineages. Error bars

represent standard deviation. * $p<0.05$, ** $p<0.01$, *** $p<0.001$.

Figure 6. *A. castellanii* and *P. aeruginosa* survival after 16 days of co-culture. (Panel A) Bars

represents the average number of A. castellanii trophozoites counts from microscopy images.

(Panel B) *P. aeruginosa* viable cell counts from ancestral or a single evolved lineage in each co-

culture condition (line 3 for strain SRP 17-047 and line 2 for strains B80398, B80427, B84725,

PA3, and SRP 17-055) are shown. Grey bars represent ancestors (A) and colored bars represent

the evolved lineage (E). Error bars represent standard deviation (n=3). Data analyzed with t-tests

between ancestor and evolved isolates. *** $p<0.0005$, **** $p<0.0001$.

Figure 7. Venn diagram of single nucleotide polymorphisms between ancestors and their

respective evolved lineages after the 12-week experimental evolution. Genes with at least two

SNPs in at least one of the evolved lineages are represented.

Supplemental figure legends:

Figure S1. Growth curves of the evolved lineages and their ancestors over time in 1% HL5 at

room temperature (panels A-F) or 37°C (panels G-L). Lines represent the mean of 3 biological

replicates. Error bars are not shown for image clarity. Data analyzed with repeat measures one-

way ANOVA with Dunnett's post-test comparing evolved lineages to their ancestor. * $p<0.05$,

** $p<0.01$, *** $p<0.001$.

Figure S2. Cell size of individual evolved lineages compared to their ancestor are shown with

each cell measure represented by a dot. Cells grown at room temperature (panels A-F) or 37°

(panels G-L) to exponential phase were measured. Panels M and N show aggregated data for

ancestral (A) and evolved (E) isolates. Horizontal bars for each strain represent the median cell

size of 100-600+ cells and error bars represent the interquartile range. Data were analyzed with

one-way ANOVA with Dunnett's post-test comparing each evolved line to its ancestor. *

$p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Figure S3. Biofilm biomass of individual evolved lineages compared to their ancestor after 3-month experimental evolution. Data for evolved lineages was aggregated to assess global patterns of evolution. Error bars represent standard deviation from four independent colonies from each lineage. Data analyzed with one-way ANOVA with Dunnett's posttest comparing each evolved line to its ancestor. *** $p < 0.001$, **** $p < 0.0001$.

Figure S4. Aggregate data for evolved *P. aeruginosa* lineages for pyocyanin (panel A) or pyoverdine (panel B) production of individual isolates compared to their ancestor after the 12-week experimental evolution. Error bars represent standard deviation from three independent colonies from each lineage. Data analyzed with one-way ANOVA with Dunnett's post-test comparing each evolved line to its ancestor. * $p < 0.05$, ** $p < 0.01$.

Figure S5. Swimming motility of individual isolates compared to their ancestor for isolates frozen at various points during the experimental evolution. Week 2 isolates were first tested, then week 3, 5, and/or 7 as needed until the loss of motility was observed. Error bars represent standard deviation from three independent colonies from each lineage. 35 mm was the largest measurable size in this analysis. N.D. indicates non-detectable growth past the stab line. Data analyzed with two-way ANOVA with Bonferroni post-test comparing each evolved line to its ancestor across time points. Error bars indicate standard deviation. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Figure S6. Competition assay between the ancestor and evolved lineages (line 3 for strain SRP

17-047 and line 2 for strains B80398, B80427, B84725, PA3, and SRP 17-055) of *P. aeruginosa*

against *A. castellanii* at all timepoints tested. Grey bars with or colored bars represent the

number of trophozoites observed when co-cultured with each ancestor (A in the x-axis) or

evolved lineage (E in the x-axis) over the course of 16 days (D) Panels represent each strain with

(A) B80398, (B) B80427, (C) B84725, (D) PA3, (E) 17-047, and (F) 17-055 measured at 0-, 3-,

6-, 8-, 11-, 13-, and 16 days of co-culture. Error bars represent standard deviation from three

independent colonies from each lineage. Data analyzed with t-tests between ancestor and

evolved lineage at each timepoint. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

References

1. Zhu M, Dai X. 2023. Stringent response ensures the timely adaptation of bacterial growth to nutrient downshift. Nat Commun 14:467.

2. Jain V, Kumar M, Chatterji D. 2006. ppGpp: stringent response and survival. J Microbiol 44:1-10.

3. Wang X, Wang J, Liu SY, Guo JS, Fang F, Chen YP, Yan P. 2023. Mechanisms of survival mediated by the stringent response in *Pseudomonas aeruginosa* under environmental stress in drinking water systems: Nitrogen deficiency and bacterial competition. J Hazard Mater 448:130941.

4. Gray DA, Dugar G, Gamba P, Strahl H, Jonker MJ, Hamoen LW. 2019. Extreme slow growth as alternative strategy to survive deep starvation in bacteria. Nat Commun 10:890.

5. Errington J. 1993. *Bacillus subtilis* sporulation: regulation of gene expression and control of morphogenesis. Microbiol Rev 57:1-33.

6. Huberts DH, Niebel B, Heinemann M. 2012. A flux-sensing mechanism could regulate the switch between respiration and fermentation. FEMS Yeast Res 12:118-28.

7. van Hoek MJ, Merks RM. 2012. Redox balance is key to explaining full vs. partial switching to low-yield metabolism. BMC Syst Biol 6:22.

8. Santajit S, Indrawattana N. 2016. Mechanisms of Antimicrobial Resistance in ESKAPE Pathogens. Biomed Res Int 2016:2475067.

9. Hancock RE, Speert DP. 2000. Antibiotic resistance in *Pseudomonas aeruginosa*: mechanisms and impact on treatment. Drug Resist Updat 3:247-255.

10. Fernandez-Billon M, Llambias-Cabot AE, Jordana-Lluch E, Oliver A, Macia MD. 2023. Mechanisms of antibiotic resistance in *Pseudomonas aeruginosa* biofilms. Biofilm 5:100129.

11. Botelho J, Tuffers L, Fuss J, Buchholz F, Utpatel C, Klockgether J, Niemann S, Tummler B, Schulenburg H. 2023. Phylogroup-specific variation shapes the clustering of antimicrobial resistance genes and defence systems across regions of genome plasticity in *Pseudomonas aeruginosa*. EBioMedicine 90:104532.

12. Mendes OR. 2023. Chapter 48 - The challenge of pulmonary *Pseudomonas aeruginosa* infection: How to bridge research and clinical pathology, p 591-608. *In* Bagchi D, Das A, Downs BW (ed), Viral, Parasitic, Bacterial, and Fungal Infections doi:https://doi.org/10.1016/B978-0-323-85730-7.00019-9. Academic Press.

13. Walker TS, Bais HP, Deziel E, Schweizer HP, Rahme LG, Fall R, Vivanco JM. 2004. *Pseudomonas aeruginosa*-plant root interactions. Pathogenicity, biofilm formation, and root exudation. Plant Physiol 134:320-31.

14. Thi MTT, Wibowo D, Rehm BHA. 2020. *Pseudomonas aeruginosa* Biofilms. Int J Mol Sci 21.

15. Britigan BE, Roeder TL, Rasmussen GT, Shasby DM, McCormick ML, Cox CD. 1992. Interaction of the *Pseudomonas aeruginosa* secretory products pyocyanin and pyochelin

generates hydroxyl radical and causes synergistic damage to endothelial cells.
Implications for *Pseudomonas*-associated tissue injury. J Clin Invest 90:2187-96.

16. Hauser AR. 2009. The type III secretion system of *Pseudomonas aeruginosa*: infection
by injection. Nat Rev Microbiol 7:654-65.

17. Hogardt M, Heesemann J. 2010. Adaptation of *Pseudomonas aeruginosa* during
persistence in the cystic fibrosis lung. Int J Med Microbiol 300:557-62.

18. Frimmersdorf E, Horatzek S, Pelnikevich A, Wiehlmann L, Schomburg D. 2010. How
Pseudomonas aeruginosa adapts to various environments: a metabolomic approach.
Environ Microbiol 12:1734-47.

19. Ebert D. 1998. Experimental evolution of parasites. Science 282:1432-5.

20. Wong A, Rodrigue N, Kassen R. 2012. Genomics of adaptation during experimental
evolution of the opportunistic pathogen *Pseudomonas aeruginosa*. PLoS Genet
8:e1002928.

21. Bakkal S, Robinson SM, Ordonez CL, Waltz DA, Riley MA. 2010. Role of bacteriocins
in mediating interactions of bacterial isolates taken from cystic fibrosis patients.
Microbiology (Reading) 156:2058-2067.

22. Merritt JH, Kadouri DE, O'Toole GA. 2005. Growing and analyzing static biofilms. Curr
Protoc Microbiol Chapter 1:Unit 1B 1.

23. Gajdacs M, Barath Z, Karpati K, Szabo D, Usai D, Zanetti S, Donadu MG. 2021. No
Correlation between Biofilm Formation, Virulence Factors, and Antibiotic Resistance in
Pseudomonas aeruginosa: Results from a Laboratory-Based In Vitro Study. Antibiotics
(Basel) 10.

24. S A. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Accessed

25. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011 17:3.

26. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics 30:2114-20.

27. Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754-60.

28. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane
T, McCarthy SA, Davies RM, Li H. 2021. Twelve years of SAMtools and BCFtools.
Gigascience 10.

29. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K,
Altshuler D, Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: a
MapReduce framework for analyzing next-generation DNA sequencing data. Genome
Res 20:1297-303.

30. DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA,
del Angel G, Rivas MA, Hanna M, McKenna A, Fennell TJ, Kernytsky AM, Sivachenko
AY, Cibulskis K, Gabriel SB, Altshuler D, Daly MJ. 2011. A framework for variation

discovery and genotyping using next-generation DNA sequencing data. Nat Genet
43:491-8.

31.   Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A,
Jordan T, Shakir K, Roazen D, Thibault J, Banks E, Garimella KV, Altshuler D, Gabriel
S, DePristo MA. 2013. From FastQ data to high confidence variant calls: the Genome
Analysis Toolkit best practices pipeline. Curr Protoc Bioinformatics 43:11 10 1-11 10 33.

32.   Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden
DM. 2012. A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaste*r strain w1118;
iso-2; iso-3. Fly (Austin) 6:80-92.

33.   Cecil RE, Yoder-Himes DR. 2024. Examining the influence of environmental factors on
Acanthamoeba castellanii and *Pseudomonas aeruginosa* in co-culture. PLoS One
19:e0305973.

34.   Mojesky AA, Remold SK. 2020. Spatial structure maintains diversity of pyocin inhibition
in household *Pseudomonas aeruginosa*. Proc Biol Sci 287:20201706.

35.   Purdy-Gibson ME, France M, Hundley TC, Eid N, Remold SK. 2015. *Pseudomonas
aeruginosa* in CF and non-CF homes is found predominantly in drains. J Cyst Fibros
14:341-6.

36.   Remold SK, Brown CK, Farris JE, Hundley TC, Perpich JA, Purdy ME. 2011.
Differential habitat use and niche partitioning by *Pseudomonas* species in human homes.
Microb Ecol 62:505-17.

37.   Hall CL, Lee VT. 2018. Cyclic-di-GMP regulation of virulence in bacterial pathogens.
Wiley Interdiscip Rev RNA 9.

38.   Coggan KA, Wolfgang MC. 2012. Global regulatory pathways and cross-talk control
Pseudomonas aeruginosa environmental lifestyle and virulence phenotype. Curr Issues
Mol Biol 14:47-70.

39.   Hou L, Debru A, Chen Q, Bao Q, Li K. 2019. AmrZ Regulates Swarming Motility
Through Cyclic di-GMP-Dependent Motility Inhibition and Controlling Pel
Polysaccharide Production in *Pseudomonas aeruginosa* PA14. Front Microbiol 10:1847.

40.   Romling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a
universal bacterial second messenger. Microbiol Mol Biol Rev 77:1-52.

41.   Ramsey DM, Wozniak DJ. 2005. Understanding the control of *Pseudomonas aeruginosa*
alginate synthesis and the prospects for management of chronic infections in cystic
fibrosis. Mol Microbiol 56:309-22.

42.   Bernier SP, Ha DG, Khan W, Merritt JH, O'Toole GA. 2011. Modulation of
Pseudomonas aeruginosa surface-associated group behaviors by individual amino acids
through c-di-GMP signaling. Res Microbiol 162:680-8.

43.   Palmer KL, Mashburn LM, Singh PK, Whiteley M. 2005. Cystic fibrosis sputum
supports growth and cues key aspects of *Pseudomonas aeruginosa* physiology. J
Bacteriol 187:5267-77.

44. Palmer KL, Aye LM, Whiteley M. 2007. Nutritional cues control *Pseudomonas*
aeruginosa multicellular behavior in cystic fibrosis sputum. J Bacteriol 189:8079-87.

45. MJ A, LO M, I SC. 1991. Temperature profiles and alginate synthesis in mucoid and
non☐mucoid variants of *Pseudomonas aeruginosa*. Letters in Applied Microbiology
12:244-248.

46. Westfall CS, Levin PA. 2017. Bacterial Cell Size: Multifactorial and Multifaceted. Annu
Rev Microbiol 71:499-517.

47. Wang JD, Levin PA. 2009. Metabolism, cell growth and the bacterial cell cycle. Nat Rev
Microbiol 7:822-7.

48. Price-Whelan A, Dietrich LE, Newman DK. 2007. Pyocyanin alters redox homeostasis
and carbon flux through central metabolic pathways in *Pseudomonas aeruginosa* PA14. J
Bacteriol 189:6372-81.

49. Eschbach M, Schreiber K, Trunk K, Buer J, Jahn D, Schobert M. 2004. Long-term
anaerobic survival of the opportunistic pathogen *Pseudomonas aeruginosa* via pyruvate
fermentation. J Bacteriol 186:4596-604.

50. Costa KC, Glasser NR, Conway SJ, Newman DK. 2017. Pyocyanin degradation by a
tautomerizing demethylase inhibits *Pseudomonas aeruginosa* biofilms. Science 355:170-
173.

51. Cox CD, Adams P. 1985. Siderophore activity of pyoverdin for *Pseudomonas*
aeruginosa. Infect Immun 48:130-8.

52. Jeong GJ, Khan F, Khan S, Tabassum N, Mehta S, Kim YM. 2023. *Pseudomonas*
aeruginosa virulence attenuation by inhibiting siderophore functions. Appl Microbiol
Biotechnol 107:1019-1038.

53. Kuburich NA, Adhikari N, Hadwiger JA. 2016. *Acanthamoeba* and *Dictyostelium* Use
Different Foraging Strategies. Protist 167:511-525.

54. Kumar A, Molli PR, Pakala SB, Bui Nguyen TM, Rayala SK, Kumar R. 2009. PAK
thread from amoeba to mammals. J Cell Biochem 107:579-85.

55. Al-Quadan T, Price CT, Abu Kwaik Y. 2012. Exploitation of evolutionarily conserved
amoeba and mammalian processes by *Legionella*. Trends Microbiol 20:299-306.

56. Siddiqui R, Khan NA. 2012. Biology and pathogenesis of Acanthamoeba. Parasit Vectors
5:6.

57. Guerin J, Bigot S, Schneider R, Buchanan SK, Jacob-Dubuisson F. 2017. Two-Partner
Secretion: Combining Efficiency and Simplicity in the Secretion of Large Proteins for
Bacteria-Host and Bacteria-Bacteria Interactions. Front Cell Infect Microbiol 7:148.

58. Kinoshita M, Namba K, Minamino T. 2021. A positive charge region of *Salmonella* FliI
is required for ATPase formation and efficient flagellar protein export. Commun Biol
4:464.

59. Zhuang WY, Shapiro L. 1995. *Caulobacter* FliQ and FliR membrane proteins, required
for flagellar biogenesis and cell division, belong to a family of virulence factor export
proteins. J Bacteriol 177:343-56.

60. Qi X, Xu X, Li H, Pan Y, Katharine Kraco E, Zheng J, Lin M, Jiang X. 2022. fliA, flrB,
and fliR regulate adhesion by controlling the expression of critical virulence genes in
Vibrio harveyi. Gene 839:146726.

61. Flores-Bautista E, Cronick CL, Fersaca AR, Martinez-Nunez MA, Perez-Rueda E. 2018.
Functional Prediction of Hypothetical Transcription Factors of Escherichia coli K-12
Based on Expression Data. Comput Struct Biotechnol J 16:157-166.

62. Lenski RE. 2017. Experimental evolution and the dynamics of adaptation and genome
evolution in microbial populations. ISME J 11:2181-2194.

63. Ma L, Conover M, Lu H, Parsek MR, Bayles K, Wozniak DJ. 2009. Assembly and
development of the Pseudomonas aeruginosa biofilm matrix. PLoS Pathog 5:e1000354.

64. Ryder C, Byrd M, Wozniak DJ. 2007. Role of polysaccharides in Pseudomonas
aeruginosa biofilm development. Curr Opin Microbiol 10:644-8.

65. Toyofuku M, Roschitzki B, Riedel K, Eberl L. 2012. Identification of proteins associated
with the Pseudomonas aeruginosa biofilm extracellular matrix. J Proteome Res 11:4906-
15.

66. York A. 2018. Sharing the burden to build a matrix. Nat Rev Microbiol 16:454-455.

67. Leong W, Poh WH, Williams J, Lutz C, Hoque MM, Poh YH, Yee BYK, Chua C,
Givskov M, Sanderson-Smith M, Rice SA, McDougald D. 2022. Adaptation to an
Amoeba Host Leads to Pseudomonas aeruginosa Isolates with Attenuated Virulence.
Appl Environ Microbiol 88:e0232221.

68. Davies EV, James CE, Brockhurst MA, Winstanley C. 2017. Evolutionary diversification
of Pseudomonas aeruginosa in an artificial sputum model. BMC Microbiol 17:3.

69. Valentin JDP, Straub H, Pietsch F, Lemare M, Ahrens CH, Schreiber F, Webb JS, van
der Mei HC, Ren Q. 2022. Role of the flagellar hook in the structural development and
antibiotic tolerance of Pseudomonas aeruginosa biofilms. The ISME Journal 16:1176-
1186.

70. Du B, Gu Y, Chen G, Wang G, Liu L. 2020. Flagellar motility mediates early-stage
biofilm formation in oligotrophic aquatic environment. Ecotoxicology and Environmental
Safety 194:110340.

71. Valentin JDP, Altenried S, Varadarajan AR, Ahrens CH, Schreiber F, Webb JS, van der
Mei HC, Ren Q. 2023. Identification of Potential Antimicrobial Targets of Pseudomonas
aeruginosa Biofilms through a Novel Screening Approach. Microbiol Spectr
11:e0309922.

72. Demirdjian S, Sanchez H, Hopkins D, Berwin B. 2019. Motility-Independent Formation
of Antibiotic-Tolerant Pseudomonas aeruginosa Aggregates. Appl Environ Microbiol 85.

73. Klausen M, Heydorn A, Ragas P, Lambertsen L, Aaes-Jørgensen A, Molin S, Tolker-
Nielsen T. 2003. Biofilm formation by Pseudomonas aeruginosa wild type, flagella and
type IV pili mutants. Molecular Microbiology 48:1511-1524.

74. Nasher F, Wren BW. 2023. Flagellin O-linked glycans are required for the interactions
between Campylobacter jejuni and Acanthamoebae castellanii. Microbiology (Reading)
169.

75. Preston TM, King CA. 1984. Binding Sites for Bacterial Flagella at the Surface of the
Soil Amoeba Acanthamoeba. Microbiology 130:1449-1458.

76. Finkel SE, Kolter R. 1999. Evolution of microbial diversity during prolonged starvation.
Proc Natl Acad Sci U S A 96:4023-7.

77. Zion S, Katz S, Hershberg R. 2024. Escherichia coli adaptation under prolonged resource
exhaustion is characterized by extreme parallelism and frequent historical contingency.
PLoS Genet 20:e1011333.

78. Behringer MG, Ho WC, Miller SF, Worthan SB, Cen Z, Stikeleather R, Lynch M. 2024.
Trade-offs, trade-ups, and high mutational parallelism underlie microbial adaptation
during extreme cycles of feast and famine. Curr Biol 34:1403-1413 e5.