Genome assembly variation and its implications for gene discovery in nematode species

Grace M. Mariene[1,2], James D. Wasmuth[1,2,*]

1. Faculty of Veterinary Medicine, University of Calgary, Calgary, Alberta, Canada

2. Host-Parasite Interactions Research Training Network, University of Calgary, Calgary, Alberta, Canada

* Corresponding author: jwasmuth@ucalgary.ca

Abstract

Genome assemblers are a critical component of genome science, but the choice of assembly software and

protocols can be daunting. Here, we investigate genome assembly variation and its implications for gene

discovery across three nematode species—*Caenorhabditis bovis*, *Haemonchus contortus*, and *Heligmosomoides*

bakeri—highlighting the critical interplay between assembly choice and downstream genomic analysis.

Selecting popular genome assemblers, we generated multiple assemblies for each species, analyzing their

structure, completeness, and effect on gene family analysis. Our findings demonstrate that assembly variations

can significantly affect gene family composition, with notable differences in critical gene families like *cyp*, *gst*,

ugt, and *nhr*. Despite broadly similar performance using various assembly metrics, comparisons of assemblies

with a single species revealed underlying structural rearrangements and inconsistencies in gene content. This

emphasizes the imperative for continuous refinement of genomic resources. Our findings advocate for a

cautious and informed approach to genome assembly and annotation to ensure reliable and insightful genomic

interpretations.

Introduction

Genome assembly software serves a critical, often underappreciated role in the life sciences. Its purpose is to reconstruct, *de novo*, the complete sequence of an organism's DNA—its genome—from shorter DNA fragments obtained through one or more sequencing technologies. A typical assembler performs various tasks, such as error correction, read alignment and overlap detection, and consensus building. The decrease cost of sequencing has empowered the launch of large BioGenome consortia which aim to sequence, assemble and annotation the genomes of as many animal and plant species as possible (Mieszkowska et al., 2022; Ebenezer et al., 2022). Individual research groups are also able to sequence the genomes of their favourite species. This places the evolution and refinement of genome assemblers at the core of current and future breakthroughs in health, agricultural, and environmental sciences.

Understanding accuracy of the output of the genome assembler—long strings of As, Gs, Cs, Ts and the occasional N—is helpful, as errors in the produced assembly will carry into and magnify through the annotation and analysis, often without notice. There are many sources of error. One example is high and/or inconsistent levels of heterozygosity in the organism's genome leading to sequence reads being assembled as separate loci rather than collapsed in the haploid assembly. A second example is long regions of repetitive DNA and transposable elements scattered through the genome being incorrectly collapsed or rearranged, leading to gaps. While some genome assemblers specialise in one particular problem, for example heterozygosity (Kajitani et al., 2014), most assemblers are generalists and seek to identify and resolve as many potential errors as possible. There are different approaches to solving these challenges and the result is that different assemblers—when given the same sequence reads as input—will produce different assemblies.

Evaluations of genome assembler performance show that even within a phylum, there is no single optimum: the Assemblathon2 competition used 102 metrics to compare assemblies from three chordate species (Bradnam et al., 2013); within the molluscs genome size dictated the choice of preferred assembler (Sun et al., 2021); and others have demonstrated the challenges of constructing a haploid assembly from tetraploid protozoa and plants (Pollo et al., 2020; Jung et al., 2020). There studies, and many others, demonstrate the need to continually evaluate assemblers for one's favourite species, which for us are the nematodes.

The phylum Nematoda is home to species that parasites of humans, livestock, and crops, free-living contributors to environmental health, and pivotal model organisms that help in the understanding of human biology. Accordingly, nematodes have often led the way in animal genomes: the first animal species to have its genome sequenced was *Caenorhabditis elegans* (*C. elegans* Sequencing Consortium, 1998); at least 177 genome assemblies—generated from a broad range of assemblers—are available for at least 135 species

(WormBase Parasite release 18) (Howe et al., 2017). This availability of genomic information has led to

comparative genomic studies on various scales, much of it focused on the evolution of gene families (Wasmuth

et al., 2008; Coghlan et al., 2019; Baker et al., 2021). For this present study, in addition to understanding how

different genome assemblers perform with different nematode species, we wanted to look deeper at the

downstream consequences of the technical variation. It has been shown that the number of apparent lineage-

specific genes is inflated if datasets are generated by different gene finding algorithms (Weisman, Murray &

Eddy, 2022), motivated us to examine the effect of genome assembler heterogeneity on gene family

comparisons.

We examined assemblies from three nematode species—*Caenorhabditis bovis*, *Haemonchus contortus*, and

Heligmosomoides bakeri—which reflect different genomic features—size and heterozygosity—and sequencing

platforms—Oxford Nanopore Technologies (ONT) and Pacific Biosciences (PacBio).

C. bovis is associated with bovine parasitic otitis disease and is the only pathogenic member of its genus (Kreis,

1964). Its genome assembly is approximately 63Mb, making it the smallest *Caenorhabditis* and one of the

smallest nematode genomes so far sequenced (Stevens et al., 2020). *H. contortus* parasitizes small ruminants

(sheep and goats) and the emergence of drug resistant strains motivated the study of its genome to

understand mechanisms of resistance and develop new drugs and vaccines (Laing et al., 2013). The *H.*

contortus genome assembly has undergone multiple rounds of improvement—a long-read assembly was

scaffolded with optical mapping and manually curated—to give seven scaffolds, with a total length of ~283Mb

(Doyle et al., 2020). Finally, *H. bakeri* is an intestinal parasite of mice that serves as a model for studying host-

parasite interactions and testing anthelmintic efficacy (Hu et al., 2013; Colomb et al., 2024). Note that earlier

descriptions of the *H. bakeri* genome, were described as *Heligmosomoides polygyrus*, a long-standing miss-

classification of the study species which has been recently corrected through genomic comparisons (Chow et

al., 2019; Stevens et al., 2023).

We generated genome assemblies for each species using several popular genome assemblers and annotated

these assemblies with a single, widely used pipeline. We found that most programs returned assemblies with

encouraging and similar assembly metrics—size, number of scaffolds, gene completeness—but intra-species

comparisons revealed structural rearrangements in the scaffolds and diverse complements of conserved

orthologues. We explored the effect of assembly differences on annotation more deeply by carrying out a

phylogenomic comparison of large gene families—the cytochrome P450s (CYP), glutathione S-transferases

(GST), UDP glucuronosyltransferases (UGT), and nuclear hormone receptors (NHR)—and found that despite

having the same sequence reads as input, the assemblies' annotations could vary by as much as 51%.

Methods

Nematode species and sequence data

DNA sequence reads for *C. bovis* and *H. contortus* were retrieved from the Sequence Read Archive, and their genomes downloaded from WormBase Parasite (release 17) (Leinonen, Sugawara & Shumway, 2011; Howe et al., 2017). The sequence reads and assembly for *H. bakeri* are new for this study. *C. bovis* DNA (BioProject: PRJEB34497) was sequenced using the Oxford Nanopore MinION platform (Stevens et al., 2020). *H. contortus* DNA (BioProject PRJEB506) was sequenced using the PacBio RS and Sequel platforms (Doyle et al., 2020). We generated *H. bakeri* DNA as part of a *de novo* sequencing project. Briefly: we extracted adult *H. bakeri* worms from experimentally infected mice; separated females and males; pooled 20 worms of each sex. PacBio Hi-Fi sequencing was done at the University of Delaware DNA Sequencing & Genotyping Center. For the assemblies in this study, we combined sequence reads from female and male worms. The sequence reads are available from the NCBI SRA: SRX23938287 and SRX23938287.

Genome assembly and quality control.

Generation of preliminary assemblies

For *C. bovis* and *H. contortus* we used six long-read assemblers: Redbean v2.5 (Ruan & Li, 2020), Canu v2.1.1 (Koren et al., 2017), Flye v2.8.2 (Lin et al., 2016; Kolmogorov et al., 2019), SMARTdenovo v1.0.0 (Liu et al., 2021), Falcon-kit v1.8.1 and Falcon-unzip v1.3.7 (Chin et al., 2016). For *H. bakeri*, we used three HiFi-compatible assemblers: Flye v2.8.2 (Lin et al., 2016; Kolmogorov et al., 2019), Hifiasm v0.16.1-r375 (Cheng et al., 2021) and HiCanu v2.1.1 (Nurk et al., 2020). All the commands and parameters can be found in File S1.

Decontamination and polishing of preliminary assemblies

For the *C. bovis* and *H. contortus*, we used Blobtools v1.1.1 (Laetsch & Blaxter, 2017). Long reads were aligned to assembly contigs using minimap2 v2.17-r941 (Li, 2018) and the taxonomic classification determined by searching contigs against the Uniprot Proteome database using DIAMOND v2.0.6.144 (Buchfink, Xie & Huson, 2015; Pundir, Martin & O'Donovan, 2017). For polishing—identification and correction of small errors—the *C. bovis* and *H. contortus* assemblies underwent multiple rounds of polishing using tools appropriate for the sequencing platforms used. Briefly:

1. Four rounds with Racon with long reads for both *C. bovis* and *H. contortus* assemblies (Vaser et al., 2017),

2. One round with Medaka v1.3.2 (Oxford Nanopore Technologies., 2018) with long reads for *C. bovis* assemblies (https://github.com/nanoporetech/medaka),

3. One round with Arrow v2.3.3 with long reads for *H. contortus* assemblies (installed via the GenomicConsensus package v2.3.3, http://github.com/PacificBiosciences/pbbioconda),

4. Two rounds of Racon and Pilon v1.14 Assemblies with Illumina reads for both *C. bovis* and *H. contortus*

assemblies (Walker et al., 2014; Vaser et al., 2017).

All the commands and parameters can be found in File S1.

Removal of residual haplotype duplications

This was done for all *H. contortus* and *H. bakeri* assemblies generated as part of this study. We used the

Purge_dups v1.2.5 tool (Guan et al., 2020). As recommended in the Purge_dups documentation, we initially

applied the default cut-off threshold, followed by two manually set cut-offs derived from the default cutoffs

(Table S10). All the commands and parameters can be found in File S1.

Assembly evaluation

For all three species, all the generated assemblies were evaluated for contiguity and completeness using

various metrics including: number of fragments, assembly size, N50, length of the longest contig, and BUSCO

v4.1.4 (Simão et al., 2015). All the commands and parameters can be found in File S1. Briefly, we used BUSCO

v4.1.4 to evaluate completeness of single copy orthologues (Manni et al., 2021) and Inspector v1.0.2 to identify

both large- and small-scale errors (Chen et al., 2021).

Genome synteny

We used Nucmer (from MUMmer v3.23) to create assembly-to-assembly alignments (Marçais et al., 2018). We

used the following assemblies as the reference for Nucmer: Redbean v2.5 of *C. bovis*; Doyle for *H. contortus*;

Hifiasm for *H. contortus*. For *C. bovis*, we used the dnadiff script (part of Nucmer) to report one-to-one

alignment coordinates and viewed the alignments with Cirocs v0.69-8 (Krzywinski et al., 2009). For all species,

we used NucDiff v2.0.3 to measure large- and small scale differences between each assembly and the

reference (Khelik et al., 2017). All the commands and parameters can be found in File S1.

Visualising BUSCO content analysis

We generated Sankey diagrams using the SankeyMATIC builder (available at http://sankeymatic.com/build/) to

visualize the transitions of BUSCO genes, i.e., those genes that changed categories between assemblies. Set-

based BUSCO genes data was displayed using UpSet plots (Lex et al., 2014). For *H. contortus*, we created a

heatmap of missing BUSCO genes with the ggplot2 package in R v3.4.2 (Wickham, 2016; R Core Team, 2022).

Most figures were edited with Adobe Illustrator (available at https://www.adobe.com). Whenever possible, a

colour-blind accessible palette was chosen using (https://davidmathlogic.com/colorblind/).

Functional annotation of BUSCO genes

We used BlastKOALA webserver (Kanehisa, Sato & Morishima, 2016).

Genome annotation

Soft masking assemblies

We used RepeatModeler v2.0.3 and RepeatMasker v4.1.2-p1 (Tarailo-Graovac & Chen, 2009; Flynn et al., 2020).

For RepeatModeler, we used NCBI/RMBLAST v2.11.0+ to create repeat library from the Dfam database of

transposable elements and DNA repeats (Storer et al., 2021). Next, we used the RepeatMasker util

(queryRepeatDatabase.pl script) to create a custom library specific to nematode species. Both repeat libraries

were combined as input for the RepeatMasker program.

Gene prediction

Protein coding genes (gene models) in the generated assemblies were predicted using Braker 3 (Gabriel et al.,

2023) installed using the Singularity container (available at https://hub.docker.com/r/teambraker/braker3).

For *H. contortus* and *H. bakeri*, we used short-read RNA-seq data from the Sequence Read Archive as external

evidence for the gene prediction (BioProjects: PRJEB506 (Laing et al., 2013) and PRJNA750155 (Pollo et al.,

2023) respectively). RNA-seq data was aligned using the STAR v2.7.10a aligner (Dobin et al., 2013) and the

generated alignments, in bam-format, used to produce a training set for AUGUSTUS v3.5.0, using GeneMark-ET

v4.71 (Hoff et al., 2016). We filtered redundant training gene structures with DIAMOND v0.9.24 (Buchfink, Xie

& Huson, 2015). For *C. bovis*—for which no RNA-Seq is available—we used the longest isoforms from *C. elegans*

gene models (BioProject: PRJNA13758) retrieved from WormBase Parasite (release 17) (Howe et al., 2017), and

the default protein sequences retrieved from UniProtKB/Swiss-Prot database (Boutet et al., 2007) as the

protein evidence for gene prediction. Braker 3 used the ProtHint pipeline to produce hints using GeneMark-EP

v4.71, DIAMOND v0.9.24 and Spaln v2.3.3d tools (Lomsadze et al., 2005; Iwata & Gotoh, 2012; Gotoh, Morita &

Nelson, 2014; Buchfink, Xie & Huson, 2015; Brůna, Lomsadze & Borodovsky, 2020). All the commands and

parameters can be found in File S1.

Identification of gene family members

Members of the Cytochrome P450 (CYP), Glutathione S-transferase (GST), UDP-glucuronosyltransferase (UGT)

and nuclear hormone receptors (NHR) gene families were identified using BLASTP v2.9.0+ against the full *C.*

elegans protein set (Camacho et al., 2009). For each species, we clustered the proteins for each family using

OrthoFinder2 ('-S blast', all other parameters were default) and included the *C. elegans* proteins to serve as

assumed outgroups (Emms & Kelly, 2019).

For the TGM protein family, the protein sequences were taken from the publication (Smyth et al., 2018). The

latest ARI and BARI protein sequences from provided by Dr. Henry McSorley (University of Dundee) (Osbourn

et al., 2017). These protein sequences were mapped to the genome assemblies using the splice-aware aligners

exonerate and miniprot (Slater & Birney, 2005; Li, 2023). For both, we a scoring threshold which only accepted

an alignment which scored within a specified percentage of the top score for a given query. For miniprot we

179   used '--outs 90', and for exonerate we used '--percent 90' and '--percent 70'. We found that the results for

180   'exonerate --percent 70' were broadly similar to 'miniprot --outs 90' (Table S69).

Results

We generated fifteen new assemblies—six for *C. bovis*, six for *H. contortus*, and three for *H. bakeri*—which were used, along with the published genomes for each species, for intraspecies comparisons. All the assemblies are available in File S2. For readability, we will refer to the published genomes by their papers' first author— Stevens for *C. bovis*, Doyle for *H. contortus*, and Chow for *H. bakeri*—and by the name of the genome assembly program for the others (Chow et al., 2019; Stevens et al., 2020; Doyle et al., 2020). Each new assembly was subject to decontamination and polishing protocols with an additional step of purging duplicates in the highly heterozygous *H. contortus* and *H. bakeri* assemblies. Decontamination—the removal of contigs from bacteria and other heterogenous sources—removed approximately 20% of the nucleotides of any given intermediate assembly and, in *C. bovis*, up to 90% of the scaffolds (Tables S7-S9). Polishing—using the sequence reads to identify and fix any small errors—led to notable improvements in the ONT *C. bovis* assemblies; for example, the Redbean2.5 assembly was improved from 79% to 96% of the BUSCO genes were found as complete (Table S7). However, polishing had a minor impact on the PacBio-based assemblies, *H. contortus* (CSII) and *H. bakeri* (HiFi), presumably because the genome assemblers used had their own in-built polishing steps. Purging duplicates— phasing out allelic repeats in highly heterozygous genomes based on read depth—led to notable reductions in the genome sizes and duplication of BUSCO genes; for example, the BUSCO genes' duplication in the *H. contortus*' Canu assembly was reduced by 48% (Table S8).

Below, we compare the assemblies of each species using general assembly statistics, synteny, and BUSCO gene completeness. We then use four gene families to demonstrate how differences in assembly impact downstream analysis.

C. bovis assemblies

The Stevens *C. bovis* assembly was generated using ONT reads with the Redbean assembler v2.3 and polished with Racon and Medaka. We generated assemblies using an updated Redbean (v2.5, hereafter Redbean2.5), Flye, Canu, SMARTdenovo, Falcon, and Falcon-Unzip (Chin et al., 2016; Lin et al., 2016; Koren et al., 2017; Kolmogorov et al., 2019; Ruan & Li, 2020; Liu et al., 2021). Following decontamination and polishing (Table S7), the overall standard assembly metrics were relatively similar (Table 1) . Compared to the Stevens assembly, the Redbean2.5 assembly was near identical in size but on fewer scaffolds (21 vs 35) and the N50 was 400 kb larger. The Canu assembly size was 10% larger and on almost four times as many scaffolds. The other four assemblies were comparable to the Stevens assembly in terms of assembly size and the number of scaffolds, but the N50s were between 25% and 50% smaller.

211 Through BUSCO analysis, we found a similar percentage of genes were complete across the seven assemblies

212 (Table 1 and Tables S11-S17). A notable difference was the duplication rate in the Canu assembly, which was

213 five times that of the Stevens assembly: 18 genes vs 97 genes.

Assembly Stats	Stevens	Redbean v2.5	Flye	Canu	SMARTdenovo	Falcon	Falcon-unzip
# Scaffolds	35	21	34	134	32	48	41
Assembly size (Mb)	62.73	62.72	62.93	69.16	63.39	64.93	64.40
N50 (Mb)	7.60	8.03	5.86	4.01	3.56	4.41	4.41
Longest contig (Mb)	10.86	10.08	10.12	12.16	6.29	9.90	9.79
BUSCO % n=3131	C:96.2 S:95.6 D:0.6	C:96.0 S:95.3 D:0.7	C:96.4 S:95.5 D:0.7	C:96.3 S:93.2 D:3.1	C:96.1 S:93.2 D:0.9	C:96.1 S:95.0 D:1.1	C:96.2 S:95.1 D:1.1
Inspector QV	25.1	25.1	25.0	25.3	25.1	25.0	25.2
Structural errors	12	8	9	10	16	21	9

Table 1: **Assembly statistics for *Caenorhabditis bovis.*** Complete 'C' gene the sum of single copy 'S' and

duplicate 'D' (all % from 3131 genes).

Of the 3131 BUSCO orthologues, 3040 genes (97%) were complete (single copy or duplicated) in at least one

assembly and 94.9% (2972) were complete in all seven assemblies (Figure S1). Of these complete genes, 15

were duplicated in all assemblies. Ninety-one genes were missing or fragmented in all seven assemblies, which

suggests they are undergoing pseudogenization or have been lost as a consequence of a genome size reduction

in *C. bovis* (Stevens et al., 2020). Ninety-one (3%) were absent (fragmented or missing) in all seven assemblies.

Six genes were complete in only one assembly. In total, the BUSCO classification differed for 178 genes, *e.g.*

from Duplicated to Fragmented (Figure 1A). Expectedly, there was strong agreement between the Stevens

(Redbean2.3) and the new Redbean2.5 assemblies; nine genes complete and single in the Stevens assembly

were fragmented or missing in the Redbean2.5 assembly with six genes in the reverse situation (Figure 1A).

Disagreements on BUSCO classification was more striking between the other assemblies. For example,

SMARTdenovo and Falcon had near identical BUSCO scores, but 36 single copy genes in the SMARTdenovo

assembly were classified as duplicated (16), fragmented (13) or missing (7) in the Falcon assembly. In a direct

comparison of Redbean2.5 and Canu assemblies, 18 genes considered fragmented or missing in Redbean2.5

but were found complete and single copy in Canu (Figure 1B).

Figure 1: Transitions of BUSCO genes across multiple *C. bovis* assemblies. The Sankey diagrams show how
BUSCO genes switch categories: red – complete and single copy genes; blue – complete and duplicated; yellow
– fragmented; green – missing. (A) The 178 genes that changed BUSCO categories in at least one assembly. The
order of the assemblies is arbitrary and follows Table 1. The data for this panel is in Table S1 and Table S2. (B)
Comparison of the 110 genes which differed between Redbean2.5 and Canu assemblies. The data for this panel
is in Table S3.

We used the original ONT reads to evaluate the reads alignment to each of the assemblies, identifying large

structural and small errors (Chen et al., 2021). All the assemblies had near identical QV scores, the Redbean2.5

assembly had the fewest large structural errors, and the Flye assembly had the fewest small-scale errors (Table

1 and Table S18).

Next, we investigated the structural variation—insertions, deletions, and translocations—between assemblies.

We needed a reference assembly, and considered Redbean2.5 assembly to have the best accuracy metrics,

albeit slightly. As expected, the Stevens and Flye assemblies—both use the DBG algorithm—had the fewest

number of differences (Table 2 and Table S19). These were followed by the OLC-based SMARTdenovo

assembly, the related HGAP Falcon and Falcon-Unzip assemblies, with the OLC-based Canu assembly having the

most differences to the Redbean2.5 assembly. For all assemblies, at least 97% of the variants were insertions

or deletions of typically short regions (1-to-10bp). However, we did find longer duplications and tandem

duplications which could impact accurate curation of genes. Using Braker3 to annotate the assemblies with

gene models, we found five genes from the Stevens assembly—four genes in tandem duplications and one

258   gene on a duplication—and 19 genes from the Redbean2.5—17 genes in tandem duplications and two on

259   duplications—in these regions (Table S20).

	Stevens	Flye	Canu	SMARTdenovo	Falcon	Falcon-unzip
Total number	8,215	8,527	62,971	12,895	19,319	16,540
Insertions	2,421	3,350	25,809	4,913	8,002	6,718
Deletions	5,553	4,916	36,399	7,684	10,929	9,363
Translocation	194	207	478	218	290	336
Relocations	29	18	142	38	39	57
Reshufflings	8	11	32	15	21	18
Inversions	10	25	109	27	38	48
Unaligned seq	0	0	2	0	0	0

Table 2: NucDiff results for *C. bovis* assemblies. Each assembly was aligned to the reference assembly,

Redbean2.5.

Larger structural variants—translocations, relocations, reshufflings, and inversions—were present even

between Redbean assemblies; we found that the RedBean2.5 Scaffold 3 was split across the Stevens Scaffolds

6, 7, and 11 (Figure 2). However, for five of the assemblies, all scaffolds could be aligned to the reference. The

exception was the Canu assembly; two scaffolds did not align to the Redbean2.5 assembly. Through a BLASTX

search against NCBI NR database (Camacho et al., 2009), we found that these were most likely of bacterial

origin and were not removed by the decontamination protocols.

269
270 **Figure 2: Synteny across six genome assemblies for the *C. bovis*.** In each Circos plot, the reference assembly
271 (Redbean2.5) is represented by the coloured scaffolds and the labeled assembly is represented by the gray
272 scaffolds. The lines linking syntenic regions are coloured according to the reference scaffold. The scaffolds are
273 ordered by length (longest to shortest) and for clarity we only label the first ten scaffolds.
274

H. contortus assemblies

The published *H. contortus* assembly is the result of an ongoing multinational consortium and involved deep

sequencing using multiple platforms and labour-intensive manual identification and correction of assembly

errors (Doyle et al., 2020). We generated six assemblies using only the PacBio reads that were used in Doyle

assembly. These six assemblies, once decontaminated and polished, varied considerably in their length,

fragmentation, and completeness (Table 3). There was a 23% difference between the longest (Flye 331 Mb)

and shortest (Falcon-unzip 262 Mb) assemblies. The SMARTdenovo assembly was the most contiguous, with

the fewest scaffolds and largest N50. The Redbean2.5 and Falcon assemblies were the closest to the Doyle

assembly in terms of assembly length.

	Doyle	Redbean v2.5	Flye	Canu	SMARTdenovo	Falcon	Falcon-unzip
# Scaffolds	7	1844	3249	2364	1473	2402	1975
Assembly size (Mb)	283.44	296.37	331.35	341.72	314.72	267.14	262.37
N50 (Mb)	47.40	0.31	0.24	0.34	0.45	0.23	0.25
Longest contig (Mb)	51.80	3.45	4.85	2.76	4.47	3.22	3.24
BUSCO % n=3131	C:94.8 S:93.7 D:1.1	C:93.9 S:86.7 D:7.2	C:94.1 S:85.0 D:9.1	C:95.2 S:86.6 D:8.6	C:94.7 S:86.5 D:8.2	C:88.2 S:84.2 D:4.0	C:87.7 S:84.2 D:3.5
Inspector Reference-free (mapping reads-to-assembly):							
Inspector QV	22.0	42.9	39.6	44.6	39.1	31.8	42.8
Structural errors	170	53	146	67	180	826	62
Inspector results Reference-guided (mapping assemblies-to-Doyle):							
Coverage of Doyle (%)	98	84	80	84	84	79	79
Reference bases with depth 0; 1; 2+ (%)	2; 98; 0	16; 82; 2	20; 76; 4	16; 78; 6	16; 81; 3	21; 77; 2	21; 77; 2
Number of collapses	0	1197	1633	1512	1234	6370	1019
Number of expansions	0	1004	1136	1127	1113	845	888
Number of inversions	0	1	0	0	0	2	0

Table 3: Assembly statistics for *H. contortus*. Complete 'C' gene the sum of single copy 'S' and duplicate 'D' (all
% from 3131 genes).

For the Inspector reports, we used both the reads and the Doyle assembly for reference-free and reference-

guided metrics, respectively (Table 3 and Table S21). We were initially surprised to see that the Doyle assembly

had the lowest QV score. Here, it is important to note that this score is reference-free, and a function of read

alignments to the assembly. The Doyle assembly has undergone extensive curation, in which data, including

optical mapping, was used to identify mistakes in the preliminary long-read driven assembly. So, a higher level

of disagreement with the raw sequence reads might be expected. For example, the Doyle assembly had the

highest number of haplotype switches likely a consequence of the manual improvement that sought to

maximize scaffold continuity (Table S21). For the six assemblies that we generated, the Inspector metrics were

similar. The coverage of the assembly-to-Doyle alignments were relatively low (79%-to-84%). The effect of the

alignment algorithm is demonstrated with the 98% coverage Doyle-to-Doyle assembly. Therefore, the regions

not covered predominantly are due to differences between each of the six assemblies and the Doyle reference.

The duplication rate—where the assembly-to-Doyle coverage was two or more—was lowest for the Redbean,

Falcon and Falcon-unzip assemblies and highest in the Canu assembly.

In considering the BUSCO results, the six new assemblies varied more than those of the other two species, with

between 87.7% to 95.2% of genes found complete. The duplication rate was notably higher in the six

assemblies compared to the Doyle reference. Of the 3131 pan-nematode orthologous genes, 3020 genes

(96.5%) were complete in at least one assembly, with 2525 genes (80.6%) complete in all seven assemblies and

94 genes (3.0%) missing or fragmented in all seven (Figures S2-S4). It is striking that fewer genes were missing

in the Canu and SMARTdenovo assemblies compared to the Doyle assembly. We looked more broadly and

found that of the 141 genes missing in the Doyle assembly, 56 (39.7%) were complete at least one of the new

assemblies, 27 in all six assemblies (Figure 3). It is likely that these represent errors in the current reference

assembly.

314
315   **Figure 3: BUSCO genes missing from the *H. contortus* Doyle assembly but complete in at least one other**
316   **assembly**. Red – complete and single copy genes; blue – complete and duplicated; yellow – fragmented; white
317   – missing. The data for this figure is in Table S6.
318

We used BlastKOALA to annotate these 141 genes and found at least three noteworthy examples (Table S22)

(Kanehisa, Sato & Morishima, 2016). One example is *let-756* which is involved in the fibroblast growth factor

receptor signaling pathway essential in nematode larval development. The knock-down of *let-756* in *C. elegans*

by RNA interference (RNAi) induces larval arrest in worms (Roubin et al., 1999). A second example is *nck-1*,

which enables kinase activity required for neuronal guidance. The knock-down of *nck-1* in *C. elegans* negatively

disrupts in axon guidance, neuronal cell positioning, and abnormalities in the excretory canal cell, gonad, and

male mating (Mohamed & Chin-Sang, 2011). A third example is *gcn-2*, a kinase involved in the nutrient-sensing

and, while knock out is not fatal in *C. elegans*, it does severally impact *C. elegans* survival during nutrient stress,

specifically amino acid limitation (Rousakis et al., 2013).

Of the 35 BUSCO genes duplicated in the Doyle reference assembly, 29 were also duplicated in most of the new

assemblies and three were single copy in all six new assemblies (Figure S2). There were large exchanges

between single copy and duplicated categories, suggesting that assembly algorithms successfully collapsed

divergent haplotypes of different genes at different rates (Figure S5 and Table S4). The presence or absence of

BUSCOs between the new assemblies was striking. For example, 44 genes considered missing or fragmented in

the Doyle reference assembly ('red') were classified as complete and single copies in the Redbean2.5 assembly

(Figure S5). Another example is, 62 genes considered missing or fragmented in the Redbean2.5 assembly single

copy in Flye (Figure S5), with 66 genes reclassified in the other direction (Figure S6 and Table S5). Both

assemblers use the DBG algorithm. The full list of BUSCO genes for each assembly can be found in Tables S23-

S29).

H. bakeri assemblies

At the start of our study, there were two published *H. bakeri* assemblies; one generated from short-reads only

and one that used a low-coverage of PacBio reads to scaffold together a short-read assembly (Coghlan et al.,

2019; Chow et al., 2019). We selected the Chow assembly as it was more contiguous and had a higher BUSCO

score. As we finished our study, a chromosome-scale assembly for *H. bakeri* was published (Stevens et al.,

2023). While more contiguous than the Chow assembly, we note that the Chow assembly had a higher BUSCO

score (Table 4).

As part of our own *H. bakeri* genome project, we generated HiFi PacBio reads and assembled them with three

different software: Flye, HiCanu, and Hifiasm. The assembly lengths differed less than 6%, but there was

considerable variation of N50 and longest contig (Table 4).

Assembly	Chow	Flye	HiCanu	Hifiasm	Stevens *et al.* (Chromosome-scale)
# Scaffolds	23647	2380	1115	284	321
Assembly size (Mb)	696.95	658.60	685.42	678.45	649.16
N50 (Mb)	0.18	0.94	1.9	4.94	110.8
Longest contig (Mb)	1.25	5.94	11.70	20.45	114.7
BUSCO % n=3131	C:94.3 S:93.1 D:1.2	C:94.6 S:92.5 D:2.1	C:95.0 S:92.2 D:2.8	C:94.9 S:92.0 D:2.9	C:93.8 S:92.6 D:1.2
Inspector QV	22.1	36.0	35.5	32.5	-
Structural errors	34	116	74	268	-

Table 4: Assembly statistics for *H. bakeri*. Complete 'C' gene the sum of single copy 'S' and duplicate 'D' (all % from 3131 genes). The chromosome-level Stevens *et al.* assembly was only published as we completed our study and was not included in our comparisons.

The three newly generated assemblies exhibited comparable performance in the reference-free Inspector

analysis. However, the Hifiasm assembly, despite its higher contiguity, exhibited more structural errors (Tables

4 and S30).

Similar to *C. bovis* and *H. contortus*, the *H. bakeri* assemblies, despite extremely similar BUSCO scores,

contained different cohorts of BUSCO genes (Figure 4 and Table S31). A total of 120 genes were complete in

three assemblies and absent (fragmented or missing) in one (Figure 5A. We note that while 58 of these in-

three-out-one genes were absent in the Chow assembly, 62 genes were fragmented or missing in one of the

HiFi-based assemblies. We explored the functions of these 62 genes to determine their potential impact on

hypotheses regarding *H. bakeri* biology (Tables S32-S38).

We found several genes whose absence, if real, could significantly affect a *H. bakeri* worm. One example is

mogs-1, absent only in the Flye assembly, which encodes a mannosyl-oligosaccharide glucosidase which is

involved in N-glycan biosynthesis and plays a pivotal role in digestion of bacteria (Geng et al., 2022). A second

example is *gpb-1,* also absent in the Flye assembly, which is a G protein beta subunit and whose RNAi knock-

down is embryonic lethal (Kamath et al., 2000). A third example, absent only in the HiCanu assembly, is *fmo-4*,

which encodes for flavin-containing monooxygenase crucial for osmoregulation and RNAi knock-down is also

embryoic lethal (Rual et al., 2004; Hirani et al., 2016).

While, marginally more BUSCO genes were found in the HiFi assemblies compared to the Chow assembly, it

came at the cost of a higher duplication. We found that 165 BUSCO genes were duplicated in at least one

assembly; the majority of these were assembly specific with 19 found duplicated in all four assemblies and a

378 further 10 found in the HiFi-based assemblies (Figure 5B). This tempts speculation that the ~29 genes are

379 indeed duplicated.

381
Figure 4: Transitions of BUSCO genes across multiple *H. bakeri* assemblies. The Sankey diagram shows how
351 genes changed categories in at least one *H. bakeri* assembly: red – complete & single copy genes; blue –
complete & duplicated; yellow – fragmented; green – missing. The order of the assemblies is arbitrary and
follows Table 4. The data for this figure is in Table S31.

Figure 5: UpSet plots of (A) Complete BUSCOs present in *H. bakeri* assemblies, and (B) Duplicated BUSCOs in
the assemblies. The figures were generated with https://asntech.shinyapps.io/intervene/.
390

Impact of assembly variation on gene family analysis

We generated protein-coding gene models for all assemblies using Braker3 (Table 5 and File S2) (Gabriel et al., 2023). For consistency, we used Braker3-generated gene models for the published genomes (see Discussion).

Species	Assembly	Number of protein-coding genes	BUSCO % n=3131
C. elegans	WS285	19,981	C:99.7 [S:99.3, D:0.4], F:0.1, M:0.2
C. bovis	Stevens – published	13,128	C:93.2 [S:92.4, D:0.8], F:1.0, M:5.8
	Stevens – Braker3	14,945	C:97.3 [S:96.5, D:0.8], F:0.7, M:2.0
	Canu	15,010	C:97.3 [S:94.3, D:3.0], F:0.5, M:2.2
	Falcon	14,143	C:96.9 [S:95.8, D:1.1], F:0.5, M:2.6
	Flye	14,070	C:97.1 [S:96.2, D:0.9], F:0.6, M:2.3
	Redbean2.5	14,001	C:97.2 [S:96.4, D:0.8], F:0.5, M:2.3
	SMARTdenovo	13,863	C:97.2 [S:96.2, D:1.0], F:0.5, M:2.3
	Falcon-unzip	13,988	C:97.1 [S:95.8, D:1.3], F:0.5, M:2.4
H. contortus	Doyle – published	19,621	C:96.2 [S:94.6, D:1.6], F:0.5, M:3.3
	Doyle – Braker3	14,824	C:92.7 [S:91.1, D:1.6], F:0.8, M:6.5
	Canu	18,808	C:93.0 [S:83.1, D:9.9], F:1.1, M:5.9
	Falcon	16,259	C:86.0 [S:81.7, D:4.3], F:2.4, M:11.6
	Flye	19,225	C:91.6 [S:81.6, D:10.0], F:1.4, M:7.0
	Redbean2.5	17,220	C:92.7 [S:84.7, D:8.0], F:1.1, M:6.2
	SMARTdenovo	17,589	C:93.7 [S:85.1, D:8.6], F:1.2, M:5.1
	Falcon-unzip	15,890	C:85.8 [S:81.9, D:3.9], F:1.7, M:12.5
H. bakeri	Chow – published	23,471	C:90.5 [S:88.8, D:1.7], F:2.9, M:6.6
	Chow – Braker3	23,218	C:92.1 [S:90.5, D:1.6], F:2.5, M:5.4
	Flye	20,801	C:95.1 [S:92.7, D:2.4], F:1.3, M:3.6
	HiCanu	20,259	C:94.9 [S:92.0, D:2.9], F:1.2, M:3.9
	Hifiasm	20,079	C:95.1 [S:92.0, D:3.1], F:1.0, M:3.9

Table 5: Protein-coding genes predicted for the different assemblies. For the published genomes, we report the both the published protein-coding gene models and our reannotation. For all other assemblies, the protein-coding gene models were generated with Braker3. BUSCO scores (C=complete, S=single copy, D=duplicated, F=fragmented, M=missing) are percentages where *n*=3131.

For all protein sets, we identified members of four gene families—cytochrome P450s (CYP), glutathione S-transferase (GST), UDP glucuronosyltransferases (UGT), and nuclear hormone receptors (NHR)—which have been implicated in anthelmintic studies and are among the largest in *C. elegans* (Tables S39-S43). For each gene family, within each species, we clustered the proteins with those from C. elegans (Emms & Kelly, 2019). Here, our assumption is that if the differences in genome assemblies did not affect the genome annotation, we would expect that every cluster would have an identical number of proteins from each assembly and that these would form perfect one-to-one orthologous groups to the exclusion of *C. elegans*. Any deviation from this

would demonstrate that differences in the assemblies led to different gene model predictions. We considered cluster membership across all assemblies ('all') and just for those assemblies generated in this study ('new') (Table 6).

	CYP	GST	UGT	NHR
C. bovis:				
Number found	min: 44 max: 58 diff: 27%	min: 38 max: 41 diff: 8%	min: 39 max: 51 diff: 27 %	min: 144 max: 160 diff: 11%
Number of clusters	42	35	16	144
Identical size	all: 26 new: 26	all: 32 new: 32	all: 9 new: 10	all: 120 new: 122
H. contortus:				
Number found	min: 35 max: 59 diff: 51%	min: 37 max: 49 diff: 28%	min: 31 max: 37 diff: 18%	min:103 max: 136 diff: 28 %
Number of clusters	35	29	14	107
Identical size	all: 14 new: 14	all: 12 new: 13	all: 8 new: 9	all: 56 new: 57
H. bakeri:				
Number found	min: 33 max: 40 diff: 19%	min: 48 max: 50 diff: 4%	min: 32 max: 43 diff: 29%	min: 121 max: 141 diff: 15%
Number of clusters	29	31	18	128
Identical size	all: 17 new: 20	all: 19 new: 24	all: 9 new: 11	all: 86 new: 100

Table 6: Showing percentage differences between the minimum and maximum number of the predicted members of the CYP, GST, UGT and NHR gene families across assemblies. We considered cluster membership across all assemblies ('all') and also restricted to those assemblies generated in this study ('new').

Overall, we found considerable variation within gene family membership (Table 6 and Tables S45-S68). We looked at the correlation between the size of the gene families and the size of the assembly or the BUSCO duplication rate. The results were inconsistent. We saw a strong and positive correlated (ρ>0.7) for: GSTs, UGTs, and NHRs in _C. bovis_ against both assembly size and duplication rate; and NHRs in _H. contortus_ against both assembly size and duplication rate. However, we saw a strong and negative correlation (ρ<-0.7) for CYPs, GTS, NHRs in _H. bakeri_ against the duplication rate (Table S44). The other comparisons could not be considered noteworthy ($|\rho|$<0.7). Across the 628 clusters that we generated, approximately 60% contained the same number of members for each assembly. Put another way, ~40% of clusters contained a variable number of proteins and often this variation was an additional member in just one assembly (Tables 6 and S45-S68).

We provide two examples of variable cluster memberships from the CYPs. The first is cluster bovis-cyp-0000003 (Table S45 and Table S46), in which: _C. elegans_ and the Redbean2.5 assembly have one member each; the Stevens, Canu and Flye assemblies have two members each; the Falcon and Falcon-unzip assemblies have 3

members each. The *C. elegans* protein is CYP-33C9 whose function has been associated with fatty-acid

desaturation hence reducing susceptibility of xenobiotics by increasing desiccation tolerance (Erkut et al.,

2013). The second example is cluster contortus-cyp-0000026, in which: *C. elegans*, in which: *C. elegans* and the

Flye, Falcon, and Falcon-unzip assemblies have one member each; the Doyle, Canu, Redbean2.5, and

SMARTdenovo assemblies have no members (Table S47 and Table S48). The *C. elegans* protein is CYP-43A1 and

is highly expressed in the intestine and serotonergic neuron ADF (Packer et al., 2019). Interestingly, an

orthologue to C. elegans CYP-43A1 was found in an earlier version of the *H. contortus* assembly and shown to

be the most highly expressed CYP in the *H. contortus* intestine (Laing et al., 2015).

In *H. bakeri*, three protein families—TGF-b mimics (TGM), ARI, and BARI—have been shown to modulate the

host's immune system (Osbourn et al., 2017; Johnston et al., 2017; Smyth et al., 2018). The available protein

sequences for these families aligned poorly to the *H. bakeri* Braker3 gene models (data not shown). We

determine the likely locations of these proteins on the genome assemblies using two splice-aware aligners,

exonerate and miniprot (Slater & Birney, 2005; Li, 2023). Similar to the other protein families, we found

variation in gene counts across the three HiFi long-read assemblies: nine-to-16 TGMs and 13-to-23 ARIs/BARIs

(Table 7 and Table S69). We checked whether the protein-to-genome alignments overlapped with Braker3

gene models and found that while most alignments (TGM 81-100%; ARI/BARI 68-100%) overlapped with a gene

model, far fewer alignments (TGM 40-91%; ARI/BARI 27-67%) overlapped gene models across by greater than

50% of their length (Table 7). Focusing on scaffold ptg000126l from the Hifiasm assembly, we saw a full set of

examples, from complete congruence between Braker3, exonerate, and miniprot, to evidence from just one

source (Figure 6).

Figure 6: Location of TGM protein-coding genes on *H. bakeri* scaffold ptg000126l. TGM genes were found to

cluster in two locations on this scaffold. The purple track is the Braker3 gene model predictions, the orange

track is the Exonerate protein-to-genome alignments, the green track is the miniprot protein-to-genome

alignments. For both exonerate and miniprot alignments, an * signifies ≥50% reciprocal overlap with a Braker3

gene model. The black bar is the scaffold, with nucleotide coordinates given above the bar.

Assembly	Chow		Flye		HiCanu		Hifiasm	
Algorithm	Miniprot	Exonerate	Miniprot	Exonerate	Miniprot	Exonerate	Miniprot	Exonerate
TGM-1	0	0	1	1	2	1	4	2
TGM-2	1	1	2	1	1	1	1	1
TGM-3	0	0	0	0	1	1	1	1
TGM-4	1	1	1	1	1	1	1	1
TGM-5	1	1	1	1	1	1	3	1
TGM-6	1	1	1	1	1	1	1	1
TGM-7	1	1	1	1	1	2	1	2
TGM-8	1	1	2	1	1	0	2	0
TGM-9	2	2	1	1	1	1	1	0
TGM-10	1	1	1	1	1	1	1	1
Total TGMs	**9**	**9**	**11**	**9**	**11**	**10**	**16**	**10**
Overlap gene model (any)	**9**	**9**	**11**	**9**	**11**	**10**	**13**	**9**
Overlap gene model (≥50%)	**7**	**6**	**10**	**6**	**10**	**8**	**9**	**4**
ARI1	1	1	3	2	7	3	8	4
ARI2	1	1	1	2	1	2	1	2
ARI3	4	3	3	3	3	3	3	3
BARI	2	1	2	2	3	3	3	3
BARI_Hom1	2	4	3	5	7	4	7	4
BARI_Hom2	1	1	1	1	1	1	1	1
Total ARI/BARI	**11**	**11**	**13**	**15**	**22**	**16**	**23**	**17**
Overlap gene model (any)	**10**	**10**	**13**	**14**	**15**	**14**	**17**	**14**
Overlap gene model (≥50%)	**3**	**3**	**6**	**10**	**10**	**8**	**8**	**8**

453   **Table 7: Immunomodulators annotated in *H. bakeri* assemblies.**

Discussion

Improvements in sequencing read length and accuracy is fuelling the demand for more contiguous genomes and led to the development of many novel computational strategies. The technologies are rapidly evolving, and we have not attempted to be comprehensive; rather, we selected genome assemblers based on their use at the start of our study. Benchmarking in other organisms has shown that there remains value in comparative analyses, especially given the remarkable genetic diversity encountered across the genomes sequenced to date (Pollo et al., 2020; Jung et al., 2020; Sun et al., 2021; Koo et al., 2023). Our objectives here have been to determine the relative efficacy of different assembly programs, particularly their effect on gene identification within nematodes species. These species cover a wide range of genomic characteristics, which should make our findings of interest to those sequencing and assembling their own favourite, non-nematode, organisms.

Long-read assemblers predominantly use two core algorithmic frameworks: Overlap-Layout-Consensus (OLC) and De-Bruijn Graph (DBG) (reviewed in (Li et al., 2012)). Each framework is decorated with a myriad of developer adaptations to address specific genome features or challenges, such as repetitive sequences, varying levels of ploidy, and differential sequencing error profiles. The process of implementing and optimising these adaptations is not trivial; they significantly affect the assemblers' effectiveness, especially in regions exhibiting intra-population structural complexities or high heterogeneity.

Arguably, the least complex genome in this study was *C. bovis* and DBG-based assemblers—Redbean2.5 and Flye—performed best, yielding less fragmented assemblies and higher BUSCO scores. They also did better with the more granular measures of assembly performance: read mapping rates, alignment depths, and structural errors (large and small). Determining the best assemblies for *H. contortus* was less obvious. Here, OLC-based assemblers—SMARTdenovo and Canu—achieved larger N50s and marginally better BUSCO scores. The Redbean2.5 (DBG) assembly had the lowest BUSCO duplication rate and fewest structural errors. For *H. bakeri*, the Hifiasm assembler, produced the most contiguous assembly but according to the Inspector report displayed the highest number of structural errors. This suggests potential challenges in handling highly heterozygous regions by the assembly algorithm, despite its default haplotype-resolving capabilities.

From our reading of the literature, the BUSCO score may be the most widely used metric to determine assembly accuracy, and we too relied heavily on it in the presented analysis. Often, we found that the scores were largely similar between assemblers, suggesting that the gene complements of each genome would be similar. However, diving deeper into the classification of the 3131 nematode single copy orthologous demonstrated a surprising situation, where a gene may be found as a single copy in half the assemblies and missing in another half. We saw differences between Redbean versions 2.3 and 2.5 which should motivate

researchers to carefully consider updated annotations; that is to ask, what gets lost when you improve your

annotation? We found at least 27 genes which were potentially incorrectly absent in the published *H. contortus*

assembly. This is one of the most carefully curated parasite genome assemblies and demonstrated the need to

fund ongoing curation for important species (Doyle et al., 2020; Doyle, 2022).

The reliance on BUSCO scores is that small variations in score can be considered trivial. For example, 1% change

in nematodes is 31 genes. Does this matter? It is important to remember that BUSCO genes are the most

conserved genes in an organism. They typically perform critical functions and their absence from an assembly, if

true *in vivo*, would point to a significant change in an organism's biology. However, If the absence is due to a

technical problem correctly assembling the reads, one might assume that assembly of regions containing less

well conserved single copy genes would be more error-prone, particularly for multi-copy gene families. This is

what we find with the *cyp*, *gst*, *ugt*, and *nhr* gene family. Comparative analyses often involve species assembled

and annotated using different methods across different laboratories (Coghlan et al., 2019). This annotation

heterogeneity inflates the number of lineages-specific genes (Weisman, Murray & Eddy, 2022). An alternative is

to reannotate the assemblies with a uniform approach, as we did here. A limitation of this approach, and our

study, is that we then ignore expert curation. For example, our reannotation of the Doyle *H. contortus* assembly

resulted in approximately 4,800 genes then the published annotation, and a lower BUSCO score. In contrast,

our reannotation of the published *C. bovis* and *H. bakeri* assemblies resulted in more genes, and importantly,

improved BUSCO scores. Together, this demonstrates the requirement to validate the absence of a gene, which

may actually be present in the assembly but score just the wrong side of a gene finder's score cut-off (Gilabert

et al., 2016; Stroehlein, Young & Gasser, 2018; Baker et al., 2021).

The TGM proteins were discovered through a proteomic survey of *H. bakeri* secreted products and later

mapped to the Chow assembly, where only two of the ten proteins could be confidently assigned to gene

models and another two proteins mapped directly to the genome (Johnston et al., 2017; Smyth et al., 2018).

Similarly, the alarmin release inhibitors (ARI and BARI) evaded direct mapping to the published Chow assembly

gene models. We find that even with improved assemblies, these proteins could not be reliably reconstructed

with arguably the most popular gene finding software. However, targeted searches did provide strong evidence

for genes for all proteins, some with multiple gene copies. However, as the variation across the assemblies

shows, a more careful curation is required.

Determining the usefulness of an assembly will remain a personal choice; one that should be guided by a range

of metrics. It is noteworthy that in a comparison of assemblies generated for several invertebrates, the

assemblies with the highest BUSCO score were sometimes the most fragmented (Sutton et al., 2021).

Conclusion

We encourage those charged with assembling genomes—be it *Heligmosomoides*, *Haemaphysalis*, or *Henneguya*—to be aware of the important differences between seemingly very similar assemblies. For users of genome assemblies and annotation, we emphasize the importance of exploring alternative assemblies generated by multiple assemblers to validate biological hypotheses. Finally, we note that much of the above discussion can be summed up neatly by the first recommendation from Assemblethon2: "Don't trust the results of a single assembly (Bradnam et al., 2013)."

Acknowledgements

We acknowledge the high-performance computing resources made available by the Faculty of Veterinary Medicine and Research Computing at the University of Calgary. We thank Drs. Stephen Doyle, John Gilleard, Graham Plastow, John Soghigian, and Frank van der Meer for discussions around this topic.

Competing Interests

The authors declare there are no competing interests.

Author contribution

GMM: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Validation; Visualization; Writing – original draft; Writing – review & editing

JDW: Conceptualization; Formal analysis; Funding acquisition; Investigation; Methodology; Supervision; Validation; Visualization; Writing – original draft; Writing – review & editing

Funding

This work was supported by grants to JDW from Results Driven Agricultural Research (RDAR, Alberta) and the Natural Sciences and Engineering Research Council of Canada (NSERC).

Data availability

539   Sequence reads for the *Heligmosomoides bakeri* genome are available on the sequence read archive (SRA):

540   SRX23938287 and SRX23938287. File S2 contains the genome assemblies and Braker3 gene models generated

541   in this study and is available on Datadryad (doi.org/10.5061/dryad.p2ngf1vzh)

References

References

Baker EA, Gilbert SPR, Shimeld SM, Woollard A. 2021. Extensive non-redundancy in a recently duplicated

developmental gene family. *BMC Ecology and Evolution* 21:33. DOI: 10.1186/s12862-020-01735-z.

Boutet E, Lieberherr D, Tognolli M, Schneider M, Bairoch A. 2007. UniProtKB/Swiss-Prot. *Methods in Molecular*

Biology (Clifton, N.J.) 406:89–112. DOI: 10.1007/978-1-59745-535-0_4.

Bradnam KR, Fass JN, Alexandrov A, Baranay P, Bechner M, Birol I, Boisvert S, Chapman JA, Chapuis G, Chikhi R,

Chitsaz H, Chou W-C, Corbeil J, Del Fabbro C, Docking TR, Durbin R, Earl D, Emrich S, Fedotov P, Fonseca

NA, Ganapathy G, Gibbs RA, Gnerre S, Godzaridis É, Goldstein S, Haimel M, Hall G, Haussler D, Hiatt JB,

Ho IY, Howard J, Hunt M, Jackman SD, Jaffe DB, Jarvis ED, Jiang H, Kazakov S, Kersey PJ, Kitzman JO,

Knight JR, Koren S, Lam T-W, Lavenier D, Laviolette F, Li Y, Li Z, Liu B, Liu Y, Luo R, MacCallum I,

MacManes MD, Maillet N, Melnikov S, Naquin D, Ning Z, Otto TD, Paten B, Paulo OS, Phillippy AM,

Pina-Martins F, Place M, Przybylski D, Qin X, Qu C, Ribeiro FJ, Richards S, Rokhsar DS, Ruby JG, Scalabrin

S, Schatz MC, Schwartz DC, Sergushichev A, Sharpe T, Shaw TI, Shendure J, Shi Y, Simpson JT, Song H,

Tsarev F, Vezzi F, Vicedomini R, Vieira BM, Wang J, Worley KC, Yin S, Yiu S-M, Yuan J, Zhang G, Zhang H,

Zhou S, Korf IF. 2013. Assemblathon 2: evaluating de novo methods of genome assembly in three

vertebrate species. *GigaScience* 2:10. DOI: 10.1186/2047-217X-2-10.

Brůna T, Lomsadze A, Borodovsky M. 2020. GeneMark-EP+: eukaryotic gene prediction with self-training in the

space of genes and proteins. *NAR genomics and bioinformatics* 2:lqaa026. DOI:

10.1093/nargab/lqaa026.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND. *Nature Methods*

12:59–60. DOI: 10.1038/nmeth.3176.

C. elegans Sequencing Consortium. 1998. Genome sequence of the nematode *C. elegans*: a platform for

investigating biology. *Science (New York, N.Y.)* 282:2012–2018. DOI: 10.1126/science.282.5396.2012.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture

and applications. *BMC Bioinformatics* 10:421. DOI: 10.1186/1471-2105-10-421.

Chen Y, Zhang Y, Wang AY, Gao M, Chong Z. 2021. Accurate long-read de novo assembly evaluation with

Inspector. *Genome Biology* 22:312. DOI: 10.1186/s13059-021-02527-4.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo assembly using phased

assembly graphs with hifiasm. *Nature Methods* 18:170–175. DOI: 10.1038/s41592-020-01056-5.

Chin C-S, Peluso P, Sedlazeck FJ, Nattestad M, Concepcion GT, Clum A, Dunn C, O'Malley R, Figueroa-Balderas

R, Morales-Cruz A, Cramer GR, Delledonne M, Luo C, Ecker JR, Cantu D, Rank DR, Schatz MC. 2016.

Phased diploid genome assembly with single-molecule real-time sequencing. *Nature Methods*

13:1050–1054. DOI: 10.1038/nmeth.4035.

Chow FW-N, Koutsovoulos G, Ovando-Vázquez C, Neophytou K, Bermúdez-Barrientos JR, Laetsch DR,

Robertson E, Kumar S, Claycomb JM, Blaxter M, Abreu-Goodger C, Buck AH. 2019. Secretion of an

Argonaute protein by a parasitic nematode and the evolution of its siRNA guides. *Nucleic Acids*

Research 47:3594–3606. DOI: 10.1093/nar/gkz142.

Coghlan A, Tyagi R, Cotton JA, Holroyd N, Rosa BA, Tsai IJ, Laetsch DR, Beech RN, Day TA, Hallsworth-Pepin K,

Ke H-M, Kuo T-H, Lee TJ, Martin J, Maizels RM, Mutowo P, Ozersky P, Parkinson J, Reid AJ, Rawlings ND,

Ribeiro DM, Swapna LS, Stanley E, Taylor DW, Wheeler NJ, Zamanian M, Zhang X, Allan F, Allen JE,

Asano K, Babayan SA, Bah G, Beasley H, Bennett HM, Bisset SA, Castillo E, Cook J, Cooper PJ, Cruz-

Bustos T, Cuéllar C, Devaney E, Doyle SR, Eberhard ML, Emery A, Eom KS, Gilleard JS, Gordon D, Harcus

Y, Harsha B, Hawdon JM, Hill DE, Hodgkinson J, Horák P, Howe KL, Huckvale T, Kalbe M, Kaur G, Kikuchi

T, Koutsovoulos G, Kumar S, Leach AR, Lomax J, Makepeace B, Matthews JB, Muro A, O'Boyle NM,

Olson PD, Osuna A, Partono F, Pfarr K, Rinaldi G, Foronda P, Rollinson D, Samblas MG, Sato H, Schnyder

M, Scholz T, Shafie M, Tanya VN, Toledo R, Tracey A, Urban JF, Wang L-C, Zarlenga D, Blaxter ML,

Mitreva M, Berriman M, International Helminth Genomes Consortium. 2019. Comparative genomics of

the major parasitic worms. *Nature Genetics* 51:163–174. DOI: 10.1038/s41588-018-0262-1.

Colomb F, Ogunkanbi A, Jamwal A, Dong B, Maizels RM, Finney CAM, Wasmuth JD, Higgins MK, McSorley HJ.

2024. IL-33-binding HpARI family homologues with divergent effects in suppressing or enhancing type 2

immune responses. *Infection and Immunity* 0:e00395-23. DOI: 10.1128/iai.00395-23.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR:

ultrafast universal RNA-seq aligner. *Bioinformatics (Oxford, England)* 29:15–21. DOI:

10.1093/bioinformatics/bts635.

Doyle SR. 2022. Improving helminth genome resources in the post-genomic era. *Trends in Parasitology* 38:831–

840. DOI: 10.1016/j.pt.2022.06.002.

Doyle SR, Tracey A, Laing R, Holroyd N, Bartley D, Bazant W, Beasley H, Beech R, Britton C, Brooks K, Chaudhry

U, Maitland K, Martinelli A, Noonan JD, Paulini M, Quail MA, Redman E, Rodgers FH, Sallé G, Shabbir

MZ, Sankaranarayanan G, Wit J, Howe KL, Sargison N, Devaney E, Berriman M, Gilleard JS, Cotton JA.

2020. Genomic and transcriptomic variation defines the chromosome-scale assembly of Haemonchus

contortus, a model gastrointestinal worm. *Communications Biology* 3:656. DOI: 10.1038/s42003-020-

01377-3.

Ebenezer TE, Muigai AWT, Nouala S, Badaoui B, Blaxter M, Buddie AG, Jarvis ED, Korlach J, Kuja JO, Lewin HA,

Majewska R, Mapholi N, Maslamoney S, Mbo'o-Tchouawou M, Osuji JO, Seehausen O, Shorinola O,

Tiambo CK, Mulder N, Ziyomo C, Djikeng A. 2022. Africa: sequence 100,000 species to safeguard

biodiversity. *Nature* 603:388–392. DOI: 10.1038/d41586-022-00712-4.

Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome*

Biology 20:238. DOI: 10.1186/s13059-019-1832-y.

Erkut C, Vasilj A, Boland S, Habermann B, Shevchenko A, Kurzchalia TV. 2013. Molecular Strategies of the

Caenorhabditis elegans Dauer Larva to Survive Extreme Desiccation. *PLOS ONE* 8:e82473. DOI:

10.1371/journal.pone.0082473.

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. 2020. RepeatModeler2 for automated

genomic discovery of transposable element families. *Proceedings of the National Academy of Sciences*

117:9451–9457. DOI: 10.1073/pnas.1921046117.

Gabriel L, Brůna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, Stanke M. 2023. BRAKER3: Fully automated

genome annotation using RNA-Seq and protein evidence with GeneMark-ETP, AUGUSTUS and TSEBRA.

DOI: 10.1101/2023.06.10.544449.

Geng S, Li Q, Zhou X, Zheng J, Liu H, Zeng J, Yang R, Fu H, Hao F, Feng Q, Qi B. 2022. Gut commensal E. coli outer

membrane proteins activate the host food digestive system through neural-immune communication.

Cell Host & Microbe 30:1401-1416.e8. DOI: 10.1016/j.chom.2022.08.004.

Gilabert A, Curran DM, Harvey SC, Wasmuth JD. 2016. Expanding the view on the evolution of the nematode

dauer signalling pathways: refinement through gene gain and pathway co-option. *BMC Genomics*

17:476. DOI: 10.1186/s12864-016-2770-7.

Gotoh O, Morita M, Nelson DR. 2014. Assessment and refinement of eukaryotic gene structure prediction with

gene-structure-aware multiple protein sequence alignment. *BMC bioinformatics* 15:189. DOI:

10.1186/1471-2105-15-189.

Guan D, McCarthy SA, Wood J, Howe K, Wang Y, Durbin R. 2020. Identifying and removing haplotypic

duplication in primary genome assemblies. *Bioinformatics (Oxford, England)* 36:2896–2898. DOI:

10.1093/bioinformatics/btaa025.

Hirani N, Westenberg M, Seed PT, Petalcorin MIR, Dolphin CT. 2016. C. elegans flavin-containing

monooxygenase-4 is essential for osmoregulation in hypotonic stress. *Biology Open* 5:537–549. DOI:

10.1242/bio.017400.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1: Unsupervised RNA-Seq-Based Genome

Annotation with GeneMark-ET and AUGUSTUS. *Bioinformatics (Oxford, England)* 32:767–769. DOI:

10.1093/bioinformatics/btv661.

Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. 2017. WormBase ParaSite – a comprehensive resource for

helminth genomics. *Molecular and Biochemical Parasitology* 215:2–10. DOI:

10.1016/j.molbiopara.2016.11.005.

Hu Y, Ellis BL, Yiu YY, Miller MM, Urban JF, Shi LZ, Aroian RV. 2013. An Extensive Comparison of the Effect of

Anthelmintic Classes on Diverse Nematodes. *PLOS ONE* 8:e70702. DOI: 10.1371/journal.pone.0070702.

Iwata H, Gotoh O. 2012. Benchmarking spliced alignment programs including Spaln2, an extended version of

Spaln that incorporates additional species-specific features. *Nucleic Acids Research* 40:e161. DOI:

10.1093/nar/gks708.

Johnston CJC, Smyth DJ, Kodali RB, White MPJ, Harcus Y, Filbey KJ, Hewitson JP, Hinck CS, Ivens A, Kemter AM,

Kildemoes AO, Le Bihan T, Soares DC, Anderton SM, Brenn T, Wigmore SJ, Woodcock HV, Chambers RC,

Hinck AP, McSorley HJ, Maizels RM. 2017. A structurally distinct TGF-β mimic from an intestinal

helminth parasite potently induces regulatory T cells. *Nature Communications* 8:1741. DOI:

10.1038/s41467-017-01886-6.

Jung H, Jeon M-S, Hodgett M, Waterhouse P, Eyun S. 2020. Comparative Evaluation of Genome Assemblers

from Long-Read Sequencing for Plants and Crops. *Journal of Agricultural and Food Chemistry* 68:7670–

7677. DOI: 10.1021/acs.jafc.0c01647.

Kajitani R, Toshimoto K, Noguchi H, Toyoda A, Ogura Y, Okuno M, Yabana M, Harada M, Nagayasu E, Maruyama

H, Kohara Y, Fujiyama A, Hayashi T, Itoh T. 2014. Efficient de novo assembly of highly heterozygous

genomes from whole-genome shotgun short reads. *Genome Research*:gr.170720.113. DOI:

10.1101/gr.170720.113.

Kamath RS, Martinez-Campos M, Zipperlen P, Fraser AG, Ahringer J. 2000. Effectiveness of specific RNA-

mediated interference through ingested double-stranded RNA in Caenorhabditis elegans. *Genome*

Biology 2:research0002.1. DOI: 10.1186/gb-2000-2-1-research0002.

Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG Tools for Functional

Characterization of Genome and Metagenome Sequences. *Journal of Molecular Biology* 428:726–731.

DOI: 10.1016/j.jmb.2015.11.006.

Khelik K, Lagesen K, Sandve GK, Rognes T, Nederbragt AJ. 2017. NucDiff: in-depth characterization and

annotation of differences between two sets of DNA sequences. *BMC bioinformatics* 18:338. DOI:

10.1186/s12859-017-1748-z.

Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using repeat graphs.

Nature Biotechnology 37:540–546. DOI: 10.1038/s41587-019-0072-8.

Koo H, Lee G-W, Ko S-R, Go S, Kwon S-Y, Kim Y-M, Shin A-Y. 2023. Two long read-based genome assembly and

annotation of polyploidy woody plants, Hibiscus syriacus L. using PacBio and Nanopore platforms.

Scientific Data 10:713. DOI: 10.1038/s41597-023-02631-z.

Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read

assembly via adaptive k-mer weighting and repeat separation. *Genome Research* 27:722–736. DOI:

10.1101/gr.215087.116.

Kreis HA. 1964. Ein neuer Nematode aus dem äusseren Gehörgang von Zeburindern in Ostafrika, Rhabditis

bovis n.sp. (Rhabditidoidea, Rhabditidae). *Schweizer Archiv fur Tierheilkunde* 106:371–378.

Krzywinski M, Schein J, Birol İ, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA. 2009. Circos: An

information aesthetic for comparative genomics. *Genome Research* 19:1639–1645. DOI:

10.1101/gr.092759.109.

Laetsch DR, Blaxter ML. 2017. BlobTools: Interrogation of genome assemblies. DOI:

10.12688/f1000research.12232.1.

Laing R, Bartley DJ, Morrison AA, Rezansoff A, Martinelli A, Laing ST, Gilleard JS. 2015. The cytochrome P450

family in the parasitic nematode Haemonchus contortus. *International Journal for Parasitology* 45:243–

251. DOI: 10.1016/j.ijpara.2014.12.001.

Laing R, Kikuchi T, Martinelli A, Tsai IJ, Beech RN, Redman E, Holroyd N, Bartley DJ, Beasley H, Britton C, Curran

D, Devaney E, Gilabert A, Hunt M, Jackson F, Johnston SL, Kryukov I, Li K, Morrison AA, Reid AJ, Sargison

N, Saunders GI, Wasmuth JD, Wolstenholme A, Berriman M, Gilleard JS, Cotton JA. 2013. The genome

and transcriptome of Haemonchus contortus, a key model parasite for drug and vaccine discovery.

Genome Biology 14:R88. DOI: 10.1186/gb-2013-14-8-r88.

Leinonen R, Sugawara H, Shumway M. 2011. The Sequence Read Archive. *Nucleic Acids Research* 39:D19–D21.

DOI: 10.1093/nar/gkq1019.

Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. 2014. UpSet: Visualization of Intersecting Sets. *IEEE*

transactions on visualization and computer graphics 20:1983–1992. DOI: 10.1109/TVCG.2014.2346248.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100. DOI:

10.1093/bioinformatics/bty191.

Li H. 2023. Protein-to-genome alignment with miniprot. *Bioinformatics* 39:btad014. DOI:

10.1093/bioinformatics/btad014.

Li Z, Chen Y, Mu D, Yuan J, Shi Y, Zhang H, Gan J, Li N, Hu X, Liu B, Yang B, Fan W. 2012. Comparison of the two

major classes of assembly algorithms: overlap–layout–consensus and de-bruijn-graph. *Briefings in*

Functional Genomics 11:25–37. DOI: 10.1093/bfgp/elr035.

Lin Y, Yuan J, Kolmogorov M, Shen MW, Chaisson M, Pevzner PA. 2016. Assembly of long error-prone reads

using de Bruijn graphs. *Proceedings of the National Academy of Sciences* 113:E8396–E8405. DOI:

10.1073/pnas.1604560113.

Liu H, Wu S, Li A, Ruan J, Wu S, Li A, Ruan J. 2021. SMARTdenovo: a de novo assembler using long noisy reads.

Gigabyte 2021:1–9. DOI: 10.46471/gigabyte.15.

Lomsadze A, Ter-Hovhannisyan V, Chernoff YO, Borodovsky M. 2005. Gene identification in novel eukaryotic

genomes by self-training algorithm. *Nucleic Acids Research* 33:6494–6506. DOI: 10.1093/nar/gki937.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. 2021. BUSCO Update: Novel and Streamlined

Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic,

Prokaryotic, and Viral Genomes. *Molecular Biology and Evolution* 38:4647–4654. DOI:

10.1093/molbev/msab199.

Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. 2018. MUMmer4: A fast and versatile

genome alignment system. *PLOS Computational Biology* 14:e1005944. DOI:

10.1371/journal.pcbi.1005944.

Mieszkowska N, Wilson WH, Jenkins HL, Mrowicki R, Cunliffe M, Adkins P, Lonsdale P, Harley J, Bishop JDD,

Warring S, Blaxter ML. 2022. Sequence locally, think globally: The Darwin Tree of Life Project.

Proceedings of the National Academy of Sciences 119. DOI: 10.1073/pnas.2115642118.

Mohamed AM, Chin-Sang ID. 2011. The C. elegans nck-1 gene encodes two isoforms and is required for

neuronal guidance. *Developmental Biology* 354:55–66. DOI: 10.1016/j.ydbio.2011.03.019.

Nurk S, Walenz BP, Rhie A, Vollger MR, Logsdon GA, Grothe R, Miga KH, Eichler EE, Phillippy AM, Koren S. 2020.

HiCanu: accurate assembly of segmental duplications, satellites, and allelic variants from high-fidelity

long reads. :2020.03.14.992248. DOI: 10.1101/2020.03.14.992248.

Osbourn M, Soares DC, Vacca F, Cohen ES, Scott IC, Gregory WF, Smyth DJ, Toivakka M, Kemter AM, le Bihan T,

Wear M, Hoving D, Filbey KJ, Hewitson JP, Henderson H, Gonzàlez-Cìscar A, Errington C, Vermeren S,

Astier AL, Wallace WA, Schwarze J, Ivens AC, Maizels RM, McSorley HJ. 2017. HpARI Protein Secreted

by a Helminth Parasite Suppresses Interleukin-33. *Immunity* 47:739-751.e5. DOI:

10.1016/j.immuni.2017.09.015.

Oxford Nanopore Technologies. 2018. Medaka: Sequence correction provided by ONT Research.

Packer JS, Zhu Q, Huynh C, Sivaramakrishnan P, Preston E, Dueck H, Stefanik D, Tan K, Trapnell C, Kim J,

Waterston RH, Murray JI. 2019. A lineage-resolved molecular atlas of *C. elegans* embryogenesis at

single-cell resolution. *Science* 365:eaax1971. DOI: 10.1126/science.aax1971.

Pollo SMJ, Leon-Coria A, Liu H, Cruces-Gonzalez D, Finney CAM, Wasmuth JD. 2023. Transcriptional patterns of

sexual dimorphism and in host developmental programs in the model parasitic nematode

Heligmosomoides bakeri. *Parasites & Vectors* 16:171. DOI: 10.1186/s13071-023-05785-2.

Pollo SMJ, Reiling SJ, Wit J, Workentine ML, Guy RA, Batoff GW, Yee J, Dixon BR, Wasmuth JD. 2020.

Benchmarking hybrid assemblies of Giardia and prediction of widespread intra-isolate structural

variation. *Parasites & Vectors* 13:108. DOI: 10.1186/s13071-020-3968-8.

Pundir S, Martin MJ, O'Donovan C. 2017. UniProt Protein Knowledgebase. In: Wu CH, Arighi CN, Ross KE eds.

Protein Bioinformatics: From Protein Modifications and Networks to Proteomics. Methods in Molecular

Biology. New York, NY: Springer, 41–55. DOI: 10.1007/978-1-4939-6783-4_2.

R Core Team. 2022. R: A Language and Environment for Statistical Computing.

Roubin R, Naert K, Popovici C, Vatcher G, Coulier F, Thierry-Mieg J, Pontarotti P, Birnbaum D, Baillie D, Thierry-

Mieg D. 1999. let-756, a C. elegans fgf essential for worm development. *Oncogene* 18:6741–6747. DOI:

10.1038/sj.onc.1203074.

Rousakis A, Vlassis A, Vlanti A, Patera S, Thireos G, Syntichaki P. 2013. The general control nonderepressible-2

kinase mediates stress response and longevity induced by target of rapamycin inactivation in

Caenorhabditis elegans. *Aging Cell* 12:742–751. DOI: 10.1111/acel.12101.

Rual J-F, Ceron J, Koreth J, Hao T, Nicot A-S, Hirozane-Kishikawa T, Vandenhaute J, Orkin SH, Hill DE, Heuvel S

van den, Vidal M. 2004. Toward Improving Caenorhabditis elegans Phenome Mapping With an

ORFeome-Based RNAi Library. *Genome Research* 14:2162–2168. DOI: 10.1101/gr.2505604.

Ruan J, Li H. 2020. Fast and accurate long-read assembly with wtdbg2. *Nature Methods* 17:155–158. DOI:

10.1038/s41592-019-0669-3.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing genome

assembly and annotation completeness with single-copy orthologs. *Bioinformatics (Oxford, England)*

31:3210–3212. DOI: 10.1093/bioinformatics/btv351.

Slater GSC, Birney E. 2005. Automated generation of heuristics for biological sequence comparison. *BMC*

Bioinformatics 6:31. DOI: 10.1186/1471-2105-6-31.

Smyth DJ, Harcus Y, White MPJ, Gregory WF, Nahler J, Stephens I, Toke-Bjolgerud E, Hewitson JP, Ivens A,

McSorley HJ, Maizels RM. 2018. TGF-β mimic proteins form an extended gene family in the murine

parasite *Heligmosomoides polygyrus*. *International Journal for Parasitology* 48:379–385. DOI:

10.1016/j.ijpara.2017.12.004.

Stevens L, Martínez-Ugalde I, King E, Wagah M, Absolon D, Bancroft R, Gonzalez de la Rosa P, Hall JL, Kieninger

M, Kloch A, Pelan S, Robertson E, Pedersen AB, Abreu-Goodger C, Buck AH, Blaxter M. 2023. Ancient

diversity in host-parasite interaction genes in a model parasitic nematode. *Nature Communications*

14:7776. DOI: 10.1038/s41467-023-43556-w.

Stevens L, Rooke S, Falzon LC, Machuka EM, Momanyi K, Murungi MK, Njoroge SM, Odinga CO, Ogendo A,

Ogola J, Fèvre EM, Blaxter M. 2020. The Genome of Caenorhabditis bovis. *Current biology: CB* 30:1023-

1031.e4. DOI: 10.1016/j.cub.2020.01.074.

Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. 2021. The Dfam community resource of transposable element

families, sequence models, and genome annotations. *Mobile DNA* 12:2. DOI: 10.1186/s13100-020-

00230-y.

Stroehlein AJ, Young ND, Gasser RB. 2018. Improved strategy for the curation and classification of kinases, with

broad applicability to other eukaryotic protein groups. *Scientific Reports* 8:6808. DOI: 10.1038/s41598-

018-25020-8.

Sun J, Li R, Chen C, Sigwart JD, Kocot KM. 2021. Benchmarking Oxford Nanopore read assemblers for high-

quality molluscan genomes. *Philosophical Transactions of the Royal Society B: Biological Sciences*

376:20200160. DOI: 10.1098/rstb.2020.0160.

Sutton JM, Millwood JD, McCormack AC, Fierst JL. 2021. Optimizing experimental design for genome

sequencing and assembly with Oxford Nanopore Technologies. *GigaByte* 2021. DOI:

10.46471/gigabyte.27.

Tarailo-Graovac M, Chen N. 2009. Using RepeatMasker to identify repetitive elements in genomic sequences.

Current Protocols in Bioinformatics Chapter 4:4.10.1-4.10.14. DOI: 10.1002/0471250953.bi0410s25.

Vaser R, Sović I, Nagarajan N, Šikić M. 2017. Fast and accurate de novo genome assembly from long

uncorrected reads. *Genome Research* 27:737–746. DOI: 10.1101/gr.214270.116.

Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK,

Earl AM. 2014. Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection and Genome

Assembly Improvement. *PLOS ONE* 9:e112963. DOI: 10.1371/journal.pone.0112963.

Wasmuth J, Schmid R, Hedley A, Blaxter M. 2008. On the Extent and Origins of Genic Novelty in the Phylum

Nematoda. *PLOS Neglected Tropical Diseases* 2:e258. DOI: 10.1371/journal.pntd.0000258.

Weisman CM, Murray AW, Eddy SR. 2022. Mixing genome annotation methods in a comparative analysis

inflates the apparent number of lineage-specific genes. *Current biology: CB* 32:2632-2639.e2. DOI:

10.1016/j.cub.2022.04.085.

Wickham H. 2016. Data Analysis. In: Wickham H ed. *ggplot2: Elegant Graphics for Data Analysis*. Use R! Cham:

Springer International Publishing, 189–201. DOI: 10.1007/978-3-319-24277-4_9.