Prenatal Exposure to Environmentally Relevant Low Dosage Dibutyl Phthalate Reduces

Placental Efficiency in CD-1 Mice

Tasha Pontifex[1], Allison Yang[1], Ayna Tracy[1], Kimberlie Burns[1], Zelieann Craig[1], Chi Zhou[1,2,*]

[1]School of Animal and Comparative Biomedical Sciences, The University of Arizona, Tucson,

AZ, United States

[2]Department of Obstetrics and Gynecology, The University of Arizona, Tucson, AZ, United

States

Short title: Prenatal DBP Exposure Reduces Placental Efficiency

*Corresponding Authors:**

Chi Zhou, Ph.D., School of Animal and Comparative Biomedical Sciences, University of

Arizona, 4101 N Campbell Ave., Tucson, AZ 85719, USA.

Phone : (520) 621-2457. Fax : (520) 626-1283. E-mail: chizhou@arizona.edu

Abstract word count: 249

Abstract

Introduction: Dibutyl phthalate (DBP), a phthalate congener, is widely utilized in consumer products and medication coatings. Women of reproductive age have a significant burden of DBP exposure through consumer products, occupational exposure, and medication. Prenatal DBP exposure is associated with adverse pregnancy/fetal outcomes and cardiovascular diseases in the offspring. However, the mechanisms underlying DBP exposure-associated adverse pregnancy outcomes are unclear. We **hypothesize** that prenatal exposure to environmentally relevant low dosage DBP adversely affects fetal-placental vascular function and development during pregnancy. **Methods:** Adult female CD-1 mice (8-10wks) were orally treated with vehicle (control) or with environmentally relevant DBP dosages (100, 10, and 0.1 µg/kg/day; DBP100, DBP10, DBP0.1) daily from 30 days before pregnancy through gestational day (GD) 18.5. Dam body mass composition (fat/lean mass) was measured non-invasively using the echo-magnetic resonance imaging system. Placental lipid disposition was examined using Oil Red O staining. **Results:** DBP100 and 10 dose dependently increased weight of non-pregnant female mice, while DBP0.1 did not affect their weight. DBP0.1 exposure also did not affect the maternal weight gain pattern and body fat % during pregnancy. However, DBP0.1 exposure significantly increased the placental weight and decreased the fetal/placental weight ratio (indicative of decreased placental efficiency) at GD18.5. We further observed that while DBP0.1 did not affect lipid disposition in maternal decidual, it significantly decreased lipid disposition in fetal labyrinth of female, but not male placentas. **Conclusions:** Prenatal DBP exposure to environmentally relevant low-dosage adversely impacts the fetal-placental efficiency and lipid disposition in a fetal sex specific manner.

Keywords: Dibutyl phthalate; Prenatal Exposure; Fetal sex; Placenta; Lipid Disposition.

Introduction

Phthalates are a group of environmental endocrine disruptors that widely exist in the environment and adversely impact cardiovascular and metabolic function in humans[1-5]. Although general usage of phthalates has gradually reduced in the past 5-10 years, dibutyl phthalate (DBP), a phthalate congener, is still widely utilized in consumer products and medication coatings[3, 6-9]. Despite decreasing trends in legacy phthalate use in response to enhanced awareness and regulation, a strong need for phthalate research remains as reductions in phthalate burden differ by age, socio-economic, and geographical groups for specific phthalate metabolites[10-13]. Women of reproductive age have a significant burden of DBP exposure[3, 6, 7] through consumer products, occupational exposure, and medication (0.1-233μg/kg/day)[8, 9]. Higher urinary DBP metabolite levels (indicative of higher exposure)[14] were observed in Hispanic and non-Hispanic black women than in other ethnic groups, strongly suggesting disparities among women of different backgrounds.

Multiple biomonitoring studies reported DBP exposure in most mothers and babies during pregnancy[15-17]. Prenatal DBP exposure is associated with adverse pregnancy/fetal outcomes (e.g., fetal growth restriction)[15-17], childhood obesity[17, 18], and cardiovascular diseases in the offspring[5, 19-22]. Studies in rats have shown that prenatal DBP exposure at 500mg/kg/day leads to reduced placental weight[23, 24]. In humans, maternal DBP exposure in 1st trimester is reported to be associated with increased risks of fetal intrauterine growth restriction (which is associated with placental dysfunction) in pregnancies with male but not with female fetuses[5, 25]. However, mechanisms underlying these adverse effects of prenatal DBP exposure are unclear.

Fetal sex has significant impacts on the effects of prenatal stress exposure-associated adverse perinatal and pregnancy outcomes[26-29]. Pregnancies with male fetuses are reported to be at a higher risk of preterm birth, while female fetuses are at a higher risk of fetal growth restriction[27]. Although prenatal exposure to other phthalate congeners (DMP and DEHP) is

reported to be associated with fetal sex-specific fetal growth restriction[25] in humans, the role of

fetal sex in DBP exposure-associated adverse pregnancy/fetal outcomes is unclear.

To date, most DBP studies in the literature evaluated dosages that are much higher than

human environmental exposure[5, 30]. Furthermore, the majority of the limited existing literature

investigating environmentally relevant dosages of DBP exposure focused on the gonad toxicity in

non-pregnant mice. Although oral ingestion of environmentally relevant DBP exposure results in

detection of its active metabolite in the ovary and disruption of various ovarian processes and

pathways (e.g., folliculogenesis, steroidogenesis, xenobiotic metabolism, DNA repair and

apoptosis), the effects of environmentally relevant low dosages DBP exposure on pregnancy and

fetal development are still not clear.

In this study, we tested the hypothesis that environmentally relevant low dosage DBP

exposure differentially dysregulates male and female fetal placental function using an established

DBP exposure mouse model[8, 31]. We examined the effect of environmentally relevant DBP

exposure on the weight gain and body composition of adult female mice. We further determined

the effect of prenatal environmentally relevant low dosage DBP exposure on fetal/placental

development as well as placental lipid disposition at gestational day (GD) 18.5.

Methods

Ethical Approval

All animal experiments followed guidance provided in the Guide for the Care and Use of

Laboratory Animals by National Research Council[32] and the Animal Research: Reporting of In

Vivo Experiments (ARRIVE) guidelines[33] with the approval of the University of Arizona

Institutional Animal Care and Use Committee (Protocol# 20-670).

Animals

Adult (8-10wks) female CD-1 mice were obtained from Charles River Laboratories (Charles

River, California) and housed in single-use, BPA-free cages at the University of Arizona Animal

Facility. Water and food were provided ad libitum, their light cycle maintained at 14 L:10 D, and

temperature at 22 ± 1°C. After an acclimation period of at least 24 h, mice were randomly assigned

to receive oral tocopherol-stripped corn oil (MP Biomedicals, OH; vehicle; control group) or an

environmentally relevant[3, 6-9] DBP (Sigma-Aldrich, MO) dosage (100, 10, and 0.1 µg/kg/day; refer

as DBP100, DBP10, and DBP0.1).

Oral DBP Dosing in Non-pregnant and Pregnant Mice

Determining Environmentally Relevant DBP Dosages That Does Not Affect Weight Gain in Non-

Pregnant Adult Female Mice

To determine an environmentally relevant DBP dosages that does not affect the weight gain

in non-pregnant female CD-1 mice, adult female CD-1 mice (8-10wks) were orally treated with

vehicle (control group; n=14) or with environmentally relevant DBP dosages (DBP100, DBP10, or

DBP0.1; n=8 per group) daily for 30 days. Mice were subjected to daily weighing, vaginal smear

collections, and oral dosing with vehicle or DBP treatments. Weight of each mouse before the first

day of treatment was used as baseline weight to calculate the percentage of weight gained during

treatment.

Environmentally Relevant Low-Dosage DBP Dosing in Adult Female Mice Pre-Pregnant through

Pregnancy

Based on our dose finding experiment in non-pregnant mice, DBP0.1 did not affect the weight

gain in adult female mice. Hence, DBP0.1 was used in subsequent experiments to determine

physiological changes that are independent of weight gain. Adult female CD-1 mice (8-10wks)

were orally treated with vehicle (control [CT] group; n=18) or DBP0.1 (n=18) daily for 30 days

before mating with male CD-1 mice without DBP exposure and throughout pregnancy until

gestational day (GD) 18.5. The weight of each mouse was measured daily. Weight of each mouse

before the first day of treatment was used as baseline weight to calculate the percentage of weight-gain during treatment. At GD18.5, mice were euthanized under pre-sedation with inhaled isoflurane followed by decapitation. Major maternal organs (heart, lung, liver, kidney, ovary) were collected and weighted at GD18.5. In addition, all fetus and placentas of each dam were collected and weighted at GD18.5.

Body Mass Composition Measurement

Control and DBP0.1-treated dams were subjected to body mass composition (fat and lean mass) measurement non-invasively while conscious using an echo-magnetic resonance imaging system (EchoMRI-900 body composition analyzer with A10 insert for mice)[34, 35] at day 0, 14, 30 of pre-pregnancy treatment as well as GD5.5, GD11.5, and GD18.5 during pregnancy.

Histology and Ovarian Corpora Lutea (CL) Counts

After the end of the experiment, ovaries from each dam were dissected followed by fat and oviductal tissue removal. Both ovaries were fixed for subsequent histological processing. Fixation was done in neutral buffered formalin overnight at 4°C with shaking. After fixing, ovaries were washed in 70% ethanol, processed, and embedded in paraffin. Embedded ovaries were serially sectioned (5μm thickness) throughout the tissue. Two sequential sections were kept and mounted every 100μm to generate various histological evaluation levels throughout the ovary. The resulting sections were processed for hematoxylin (Richard-Allan, 7211) and eosin (Richard-Allan, 7111) staining using a standard procedure. Stained ovary slides were imaged prior to analysis as described[8]. Briefly, each ovarian section was imaged at 10x magnification using a Leica DMIL LED microscope. Ovarian sections too large to fit in one view, were imaged and captured in tiles, and then merged using the stitching tool in Adobe Photoshop. Only fully developed, functional CLs were counted and compared between treatments.

Oil Red O Staining of Mouse Placenta

The neutral lipids in mice placenta tissues were stained using an Oil Red O Staining Kit (IHC World Cat# IW-3008) following the manufacturer's instruction. In brief, mice placenta tissues were embedded in OCT and stored in -80°C until sectioning. Frozen mice placenta sections (5µm thick, 2 per individual placenta) were fixed in 10% formalin for 5 mins. After fixing, the placental sections were incubated in Pre-Stain Solution for 5 mins at room temperature followed with 10 min Oil Red O solution incubation at 60°C. After rinsing, nuclei of sections were The neutral lipids in mice placenta tissues were stained counterstained with Mayer's Hematoxylin for 1 min. After rinsing, sections were mounted using Aqueous Mounting Medium, and imaged at 40x magnification using a brightfield microscope (Leica DM5500B). Lipid droplets area in mice placental tissue sections were quantified using National Institutes of Health (NIH) ImageJ software (http://rsbweb.nih.gov/ij/).

Statistical Analyses

SigmaPlot software (Systat Software., San Jose, CA) was used for statistical analyses. Data are represented as the medians ± standard deviation (SD). Data analyses were performed using the Mann-Whitney Rank Sum Test or Kruskal-Wallis test as appropriate. Differences were considered significant when P <0.05. Benjamini and Hochberg False Discovery Rate (FDR)-adjustment[36, 37] was used for multiple comparison correction as appropriate.

Results

Effect of environmentally relevant dosages of DBP exposure on the weight gain of non-pregnant female mice

To determine the DBP exposure dosage that does not affect non-pregnant adult female mice, animals were orally treated with vehicle (CT group), DBP100, DBP10, or DBP0.1 for 30 days. Compared with control group, 30-day exposure to DBP100 and DBP10 does-dependently

increased the weight of non-pregnant female CD-1 mice (Fig.1A). Specifically, female mice with DBP100 exposure significantly gained weight starting after 18 days of treatment (10.8%) and peaked at 20 days of treatment (15.3%). Mice with DBP10 exposure exhibit a trend of weight gain after 20 days of treatment and reach significant weight gain of 5.6%~6.9% by 23 to 25 days after treatment. Similar to the control (CT) group, 30 days of DBP0.1 exposure does not cause significant weight gain of female mice (Fig.1B).

Effect of DBP0.1 exposure on female fertility efficiency in CD-1 mice

DBP0.1 exposure decreased the pregnancy rate in the female mice from 94.44% to 83.33% (Fig.2A), but it did not affect number of live pups per litter (litter size Fig.2B) andmale to female fetus ratio compared to CT group (Fig.2B-C). Interestingly, DBP0.1 exposed mice exhibited significantly lower pup to CL ratio at GD18.5 (defined as number of pups divided by number of CLs, Fig.2D).

Effect of DBP0.1 exposure on weight gain patterns of pregnant mice

Compared to control group, DBP0.1 exposure does not affect the pre-pregnancy and post pregnancy weight gain patterns (Fig.3A). Our body composition data (EchoMRI™ LLC, TX; Fig.3B) showed that DBP0.1 exposure does not affect the body fat percentage of dams (pre-pregnancy through GD18.5). Further, DBP0.1 exposure did not affect the weight of the maternal heart, lung, kidney, and liver at GD18.5 (Fig.3C-F).

Effect of prenatal DBP0.1 exposure on placental efficiency

Compared to the CT group, DBP0.1 exposure did not affect fetal weight at GD18.5 (Fig.4A). There is no difference between the weight of male and female fetuses from both CT and DBP0.1 dams. In CT group, male placental weights are significantly higher than female placentas. However, this fetal sex-specific placental weight differences disappeared in DBP group (Fig.4B).

DBP0.1 exposure significantly increased the placental weight and decreased the fetal/placental weight ratio (indicative of decreased placental efficiency, Fig.4B-C). Specifically, the DBP exposure-increased placenta weight is more dramatic in female than in male placentas.

Effect of prenatal DBP0.1 exposure on placental lipid disposition pattern.

In placentas from control dams, lipid disposition (calculated as the percentage of tissue area) in the fetal labyrinth area of the placenta is significantly higher in female than in male placentas (Fig.5A). DBP0.1 exposure significantly reduced the lipid disposition in the fetal labyrinth area of female placentas but did not affect the lipid disposition in male placentas. The fetal sex-specific fetal labyrinth lipid disposition pattern observed in control placentas was diminished in placentas from the DBP group. DBP0.1 exposure does not affect the lipid disposition in the maternal decidual area in both female and male placentas (Fig.5B).

Discussion

In this study, we have demonstrated that daily dosing with a low, environmentally relevant exposure to DBP of 0.1 µg/kg/day does not affect the weight (both pre-pregnant and during pregnancies) of adult female CD-1 mice but increases the placental weight at GD18.5. Further, we have shown for the first time that prenatal DBP0.1 exposure caused fetal sex-specific reductions in the fetal labyrinth lipid disposition in the placenta while not affecting the lipid disposition in the maternal decidual area at GD18.5 in CD-1 mice. These data clearly indicate that environmentally relevant low-dosage DBP exposure at 0.1µg/kg/day causes fetal sex-specific placental dysfunction.

Consistent with previous reports[8], we observed that 30 days of environmentally relevant dosages of DBP exposure (DBP100 and DBP10) dose-dependently increased the weight of non-pregnant adult female mice. However, our data showed that 30 days of environmentally relevant DBP exposure at a much lower dosage at DBP0.1 causes no significant weight gain nor body fat

percentage changes in non-pregnant adult female mice compared to control. DBP is known to be

associated with with obesity[38-40], metabolic disorders[41], and cardiovascular diseases[25, 42] in

humans. Maternal obesity is a prevalent risk factor for pregnancy complications (e.g., preterm

birth, fetal growth restriction)[43-47]. In studies utilizing higher DBP dosage that causes significant

maternal weight gains, it would be very difficult to separate the adverse pregnancy outcomes

caused by DBP exposure and by the maternal weight gain caused by DBP exposure. In this

study, we specifically chose an environmentally relevant DBP dosage that does not significantly

affect the weight gain in female mice, which allows us to evaluate the effect of maternal exposure

to environmentally relevant low-dosage DBP during pregnancy without being affected by the

maternal weight gain that would have occurred with higher DBP dosages.

Although the reproductive toxicity of DBP on male fertility is well established[48, 49], the effect

of DBP exposure on female fertility is not fully understood. Existing literature on the effects of DBP

exposure on pregnancy outcomes often utilizes DBP dosages that are much higher than the

human environment[5, 30], the effect of environmentally relevant low-dosage DBP exposure during

pregnancy on fetal development and pregnancy outcomes remains elusive.

Our data have shown that although 30 days of DBP0.1 exposure did not affect the body

weight or body fat composition during pregnancy, it significantly reduced the reproductive

efficiency of the female mice. When mating with untreated CD-1 male mice, the DBP0.1 exposure

reduced the pregnancy rate in female CD-1 mice from 93.33% to 82.35%. In addition, although

DBP0.1 exposure did not significantly affect the litter size, fetal sex ratio, and ovarian CL count, it

significantly reduced the pup-to-CL ratio which is indicative of a disconnect between ovulation

and live pups. These data agree with previous reports that DBP is an endocrine disruptor that

adversely affects pregnancy outcomes[30, 40]. Specifically, human epidemiological studies report

associations between phthalate exposures and increased risk of recurrent pregnancy loss[50]; thus,

the disconnect between CL number and pups could be due to embryo loss but further studies will

be needed to address this possibility directly. Observations in this current study extended the DBP

exposure-associated adverse effects on female fertility to an environmentally relevant DBP exposure dosage to as low as 0.1μg/kg/day.

Our data have shown that DBP0.1 exposure allows for examination of the effect the environmentally relevant DBP exposure during pregnancy independent of the effect of maternal weight gain that would usually cause by higher dosages of DBP. Specifically, adult female mice with 30-days pre-pregnancy DBP0.1 exposure and continued daily DBP0.1 exposure throughout pregnancy did not have significant changes in weight gain pattern and the body fat composition when compared with control animals. Further, this DBP0.1 treatment did not change the weight of major maternal organs at GD18.5. We further observed that the prenatal DBP0.1 exposure did not affect fetal weight but significantly increased placental weight and reduced fetal/placental weight ratio at GD18.5 indicative of reduced placental efficiency in pups from DBP0.1 exposed dams. Further, our data showed that DBP0.1 causes more dramatical placental weight increase in female than male placentas, which indicates more dramatical reduction in placental efficiency. All these data show that although DBP exposure at an environmentally relevant low dosage (DBP0.1) seems not to affect maternal weight and body composition during pregnancy, it still fetal sex-specifically affects placental development and placental efficiency.

Lipids play a crucial role in placental function[51, 52]. Dysregulation in placental lipid dynamics, as observed in our study, indicates disrupted nutrient transportation and overall placental function[52-55]. We observed a notable sexual dimorphism in fetal labyrinth lipid disposition in placentas from control mice, with female placentas exhibiting significantly higher fetal labyrinth lipid accumulation compared to their male counterparts. Strikingly, DBP0.1 exposure led to a significant reduction in fetal labyrinth lipid disposition in female placentas but does not affect the fetal labyrinth lipid disposition in male placentas. Our observation that DBP0.1 exposure abolishes the fetal sex-specific difference in fetal labyrinth lipid disposition, highlighting the disruptive effects of DBP on normal placental development and function. This is further supported by the fetal sex-specific dysregulation of placental weight in DBP0.1 exposed animals that observed in the current

study.

Interestingly, although DBP0.1 exposure leads to fetal sex-specific dysregulation of fetal

labyrinth lipid disposition, it does not affect lipid disposition in maternal decidual areas of placental

tissues both female and male. This data indicates that the DBP0.1 exposure mainly dysregulates

lipid disposition in the fetal labyrinth area of the placentas. This observation aligns with the

established role of the labyrinth as the primary nutrient exchange structure in the placenta[56, 57],

where dynamic lipid trafficking and metabolism is essential to sustain fetal growth and

development. This data also agrees with previous reports showing that DBP and its metabolite

can cross the placenta and play important roles in lipid metabolism and transportation[58-61].

In conclusion, our data have demonstrated that environmentally relevant low dosage of DBP

exposure at 0.1μg/kg/day reduces the reproductive efficiency in female mice and fetal sex-

specifically dysregulates the placental efficiency and lipid disposition. Findings in this study

highlight the significance of including fetal sex as a biological variable when examining the impact

of environmental endocrine disruptor exposures on placental function and fetal development.

Perspectives

To date, mechanisms underlying adverse effects of prenatal DBP exposure are unclear.

Further, most studies in the literature evaluated DBP dosages that are much higher than the

human environmental exposure. Here we reported that DBP exposure as low as 0.1μg/kg/day

during pregnancy, though does not affect maternal weight gain, still causes fetal sex-specific

dysregulation of placental development and fetal labyrinth lipid disposition. These observations

highlight the significance of further studying the effect of environmental endocrine disruptor

exposure during pregnancy on fetal development at a human environment relevant dosage as

well as the necessity to include fetal sex as a biological variable in future investigations.

Acknowledgments

We would like to thank Dr. Xiaosong Liu for technical assistance.

Sources of Funding

This study is supported by the American Heart Association (AHA) Transformative Project

Award 23TPA1066252 (Zhou), and Pilot grant (Zhou, Craig) from the Southwest Environmental

Health Sciences Center (National Institutes of Health [NIH] P30 ES006694). The content is solely

the responsibility of the authors and does not necessarily represent the official views of the NIH.

This study is also supported by the Research, Innovation & Impact (RII) and the Technology

Research Initiative Fund/Improving Health initiative at the University of Arizona (Zhou).

Conflict of Interest/Disclosure Statement

The authors have no conflict of interest.

References

1. Jaimes R, 3rd, Swiercz A, Sherman M, Muselimyan N, Marvar PJ and Posnack NG. Plastics
and cardiovascular health: phthalates may disrupt heart rate variability and cardiovascular
reactivity. *Am J Physiol Heart Circ Physiol*. 2017;313:H1044-h1053.
2. Jaimes R, 3rd, McCullough D, Siegel B, Swift L, McInerney D, Hiebert J, Perez-Alday EA,
Trenor B, Sheng J, Saiz J, Tereshchenko LG and Posnack NG. Plasticizer Interaction With
the Heart: Chemicals Used in Plastic Medical Devices Can Interfere With Cardiac
Electrophysiology. *Circ Arrhythm Electrophysiol*. 2019;12:e007294.
3. Saravanabhavan G, Guay M, Langlois É, Giroux S, Murray J and Haines D. Biomonitoring of
phthalate metabolites in the Canadian population through the Canadian Health Measures
Survey (2007-2009). *International journal of hygiene and environmental health*.
2013;216:652-661.
4. Wang Y, Zhu H and Kannan K. A Review of Biomonitoring of Phthalate Exposures. *Toxics*.
2019;7:21. doi: 10.3390/toxics7020021.
5. Mariana M and Cairrao E. Phthalates Implications in the Cardiovascular System. *Journal of*
cardiovascular development and disease. 2020;7:26. doi: 10.3390/jcdd7030026.

6. Mitro SD, Chu MT, Dodson RE, Adamkiewicz G, Chie L, Brown FM and James-Todd TM. Phthalate metabolite exposures among immigrants living in the United States: findings from NHANES, 1999-2014. *Journal of exposure science & environmental epidemiology*. 2019;29:71-82.

7. Silva MJ, Barr DB, Reidy JA, Malek NA, Hodge CC, Caudill SP, Brock JW, Needham LL and Calafat AM. Urinary levels of seven phthalate metabolites in the U.S. population from the National Health and Nutrition Examination Survey (NHANES) 1999-2000. *Environmental health perspectives*. 2004;112:331-338.

8. Liu X and Craig ZR. Environmentally relevant exposure to dibutyl phthalate disrupts DNA damage repair gene expression in the mouse ovary†. *Biology of reproduction*. 2019;101:854-867.

9. Hernández-Díaz S, Su YC, Mitchell AA, Kelley KE, Calafat AM and Hauser R. Medications as a potential source of exposure to phthalates among women of childbearing age. *Reproductive toxicology (Elmsford, NY)*. 2013;37:1-5.

10. Zota AR, Calafat AM and Woodruff TJ. Temporal trends in phthalate exposures: findings from the National Health and Nutrition Examination Survey, 2001-2010. *Environ Health Perspect*. 2014;122:235-41.

11. Shin HM, Dhar U, Calafat AM, Nguyen V, Schmidt RJ and Hertz-Picciotto I. Temporal Trends of Exposure to Phthalates and Phthalate Alternatives in California Pregnant Women during 2007-2013: Comparison with Other Populations. *Environ Sci Technol*. 2020;54:13157-13166.

12. Wu H, Kupsco AJ, Deierlein AL, Just AC, Calafat AM, Oken E, Braun JM, Mercado-Garcia A, Cantoral A, Téllez-Rojo MM, Wright RO and Baccarelli AA. Trends and Patterns of Phthalates and Phthalate Alternatives Exposure in Pregnant Women from Mexico City during 2007-2010. *Environ Sci Technol*. 2020;54:1740-1749.

13. Domínguez-Romero E, Komprdová K, Kalina J, Bessems J, Karakitsios S, Sarigiannis DA and Scheringer M. Time-trends in human urinary concentrations of phthalates and substitutes DEHT and DINCH in Asian and North American countries (2009-2019). *J Expo Sci Environ Epidemiol*. 2023;33:244-254.

14. James-Todd TM, Meeker JD, Huang T, Hauser R, Seely EW, Ferguson KK, Rich-Edwards JW and McElrath TF. Racial and ethnic variations in phthalate metabolite concentration changes across full-term pregnancies. *Journal of exposure science & environmental epidemiology*. 2017;27:160-166.

15. Huang Y, Li J, Garcia JM, Lin H, Wang Y, Yan P, Wang L, Tan Y, Luo J, Qiu Z, Chen JA and Shu W. Phthalate levels in cord blood are associated with preterm delivery and fetal growth parameters in Chinese women. *PloS one*. 2014;9:e87430.

16. Zhang Y, Lin L, Cao Y, Chen B, Zheng L and Ge RS. Phthalate levels and low birth weight: a nested case-control study of Chinese newborns. *The Journal of pediatrics*. 2009;155:500-504.

17. Whyatt RM, Adibi JJ, Calafat AM, Camann DE, Rauh V, Bhat HK, Perera FP, Andrews H, Just AC, Hoepner L, Tang D and Hauser R. Prenatal di(2-ethylhexyl)phthalate exposure and length of gestation among an inner-city cohort. *Pediatrics*. 2009;124:1213.

18. Latini G, De Felice C, Presta G, Del Vecchio A, Paris I, Ruggieri F and Mazzeo P. In utero exposure to di-(2-ethylhexyl)phthalate and duration of human pregnancy. *Environmental health perspectives*. 2003;111:1783-1785.

19. Mariana M, Feiteiro J, Verde I and Cairrao E. The effects of phthalates in the cardiovascular and reproductive systems: A review. *Environment international*. 2016;94:758-776.

20. Valvi D, Casas M, Romaguera D, Monfort N, Ventura R, Martinez D, Sunyer J and Vrijheid M. Prenatal Phthalate Exposure and Childhood Growth and Blood Pressure: Evidence from the Spanish INMA-Sabadell Birth Cohort Study. *Environmental health perspectives*. 2015;123:1022-1029.

21. Amin MM, Ebrahimpour K, Parastar S, Shoshtari-Yeganeh B, Hashemi M, Mansourian M, Poursafa P, Fallah Z, Rafiei N and Kelishadi R. Association of urinary concentrations of phthalate metabolites with cardiometabolic risk factors and obesity in children and adolescents. *Chemosphere*. 2018;211:547-556.

22. Vafeiadi M, Myridakis A, Roumeliotaki T, Margetaki K, Chalkiadaki G, Dermitzaki E, Venihaki M, Sarri K, Vassilaki M, Leventakou V, Stephanou EG, Kogevinas M and Chatzi L. Association of Early Life Exposure to Phthalates With Obesity and Cardiometabolic Traits in Childhood: Sex Specific Associations. *Frontiers in public health*. 2018;6:327.

23. Mahaboob Basha P and Radha MJ. Gestational di-n-butyl phthalate exposure induced developmental and teratogenic anomalies in rats: a multigenerational assessment. *Environ Sci Pollut Res Int*. 2017;24:4537-4551.

24. Strakovsky RS and Schantz SL. Using Experimental Models to Assess Effects of Bisphenol A (BPA) and Phthalates on the Placenta: Challenges and Perspectives. *Toxicological sciences : an official journal of the Society of Toxicology*. 2018;166:250-268.

25. Zhao Y, Chen L, Li LX, Xie CM, Li D, Shi HJ and Zhang YH. Gender-specific relationship between prenatal exposure to phthalates and intrauterine growth restriction. *Pediatric research*. 2014;76:401-408.

26. Wainstock T, Shoham-Vardi I, Glasser S, Anteby E and Lerner-Geva L. Fetal sex modifies effects of prenatal stress exposure and adverse birth outcomes. *Stress*. 2015;18:49-56.

27. Melamed N, Yogev Y and Glezerman M. Fetal gender and pregnancy outcome. *J Matern Fetal Neonatal Med*. 2010;23:338-44.

28. Al-Qaraghouli M and Fang YMV. Effect of Fetal Sex on Maternal and Obstetric Outcomes. *Front Pediatr*. 2017;5:144.

29. Yang J, Armson BA, Attenborough R, Carson GD, da Silva O, Heaman M, Janssen P, Murphy PA, Pasquier JC, Sauve R, Von Dadelszen P, Walker M and Lee SK. Survey of Mode of Delivery and Maternal and Perinatal Outcomes in Canada. *J Obstet Gynaecol Can*. 2022;44:960-971.

30. Heidari T, Batavani RA, Malekinejad H and Hobbenaghi R. Evaluation of di-n-butyl phthalate reproductive toxicity in pregnant rats and their offspring and assessment of vitamin E administration in reducing toxicity. *Vet Res Forum*. 2022;13:201-208.

31. Craig ZR, Hannon PR, Wang W, Ziv-Gal A and Flaws JA. Di-n-butyl phthalate disrupts the expression of genes involved in cell cycle and apoptotic pathways in mouse ovarian antral follicles. *Biol Reprod*. 2013;88:23.

32. National Research Council Committee for the Update of the Guide for the C and Use of Laboratory A. The National Academies Collection: Reports funded by National Institutes of Health *Guide for the Care and Use of Laboratory Animals* Washington (DC): National Academies Press (US) Copyright © 2011, National Academy of Sciences.; 2011.

33. Percie du Sert N, Hurst V, Ahluwalia A, Alam S, Avey MT, Baker M, Browne WJ, Clark A, Cuthill IC, Dirnagl U, Emerson M, Garner P, Holgate ST, Howells DW, Karp NA, Lazic SE, Lidster K, MacCallum CJ, Macleod M, Pearl EJ, Petersen OH, Rawle F, Reynolds P, Rooney K, Sena ES, Silberberg SD, Steckler T and Würbel H. The ARRIVE guidelines 2.0: Updated guidelines for reporting animal research. *PLoS Biol*. 2020;18:e3000410.

34. Dranse HJ, Waise TMZ, Hamr SC, Bauer PV, Abraham MA, Rasmussen BA and Lam TKT. Physiological and therapeutic regulation of glucose homeostasis by upper small intestinal PepT1-mediated protein sensing. *Nat Commun*. 2018;9:1118.

35. Geisler CE, Ghimire S, Bruggink SM, Miller KE, Weninger SN, Kronenfeld JM, Yoshino J, Klein S, Duca FA and Renquist BJ. A critical role of hepatic GABA in the metabolic dysfunction and hyperphagia of obesity. *Cell Rep*. 2021;35:109301.

36. Zhou C, Zou QY, Li H, Wang RF, Liu AX, Magness RR and Zheng J. Preeclampsia Downregulates MicroRNAs in Fetal Endothelial Cells: Roles of miR-29a/c-3p in Endothelial Function. *The Journal of clinical endocrinology and metabolism*. 2017;102:3470-3479.

37. Zhou C, Yan Q, Zou QY, Zhong XQ, Tyler CT, Magness RR, Bird IM and Zheng J. Sexual Dimorphisms of Preeclampsia-Dysregulated Transcriptomic Profiles and Cell Function in Fetal Endothelial Cells. *Hypertension (Dallas, Tex: 1979)*. 2019;74:154-163.

38. Xia B, Zhu Q, Zhao Y, Ge W, Zhao Y, Song Q, Zhou Y, Shi H and Zhang Y. Phthalate exposure and childhood overweight and obesity: Urinary metabolomic evidence. *Environment international*. 2018;121:159-168.

39. Harley KG, Berger K, Rauch S, Kogut K, Claus Henn B, Calafat AM, Huen K, Eskenazi B and Holland N. Association of prenatal urinary phthalate metabolite concentrations and childhood BMI and obesity. *Pediatric research*. 2017;82:405-415.

40. Qian Y, Shao H, Ying X, Huang W and Hua Y. The Endocrine Disruption of Prenatal Phthalate Exposure in Mother and Offspring. *Frontiers in public health*. 2020;8:366.

41. Radke EG, Galizia A, Thayer KA and Cooper GS. Phthalate exposure and metabolic effects: a systematic review of the human epidemiological evidence. *Environ Int*. 2019;132:104768.

42. Trasande L, Sathyanarayana S, Spanier AJ, Trachtman H, Attina TM and Urbina EM. Urinary phthalates are associated with higher blood pressure in childhood. *The Journal of pediatrics*. 2013;163:747-53.e1.

43. Jeyabalan A. Epidemiology of preeclampsia: impact of obesity. *Nutrition reviews*. 2013;71 Suppl 1:18.

44. Bodnar LM, Ness RB, Markovic N and Roberts JM. The risk of preeclampsia rises with increasing prepregnancy body mass index. *Annals of Epidemiology*. 2005;15:475-482.

45. Bodnar LM, Catov JM, Klebanoff MA, Ness RB and Roberts JM. Prepregnancy body mass index and the occurrence of severe hypertensive disorders of pregnancy. *Epidemiology (Cambridge, Mass)*. 2007;18:234-239.

46. Catov JM, Ness RB, Kip KE and Olsen J. Risk of early or severe pre-eclampsia related to pre-existing conditions. *International journal of epidemiology*. 2007;36:412-419.

47. Sanchez-Aranguren LC, Prada CE, Riano-Medina CE and Lopez M. Endothelial dysfunction and preeclampsia: role of oxidative stress. *Frontiers in physiology*. 2014;5:372.

48. Gray LE, Jr., Ostby J, Furr J, Price M, Veeramachaneni DN and Parks L. Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol Sci*. 2000;58:350-65.

49. Boberg J, Christiansen S, Axelstad M, Kledal TS, Vinggaard AM, Dalgaard M, Nellemann C and Hass U. Reproductive and behavioral effects of diisononyl phthalate (DINP) in perinatally exposed rats. *Reprod Toxicol*. 2011;31:200-9.

50. Liao KW, Kuo PL, Huang HB, Chang JW, Chiang HC and Huang PC. Increased risk of phthalates exposure for recurrent pregnancy loss in reproductive-aged women. *Environ Pollut*. 2018;241:969-977.

51. Biezenski JJ. Role of placenta in fetal lipid metabolism. I. Injection of phospholipids double labeled with C14-glycerol and P32 into pregnant rabbits. *Am J Obstet Gynecol*. 1969;104:1177-89.

52. Coleman RA. Placental metabolism and transport of lipid. *Fed Proc*. 1986;45:2519-23.

53. Lorentzen B and Henriksen T. Plasma lipids and vascular dysfunction in preeclampsia. *Semin Reprod Endocrinol*. 1998;16:33-9.

54. Lewis RM, Wadsack C and Desoye G. Placental fatty acid transfer. *Curr Opin Clin Nutr Metab Care*. 2018;21:78-82.

55. Jarvie E, Hauguel-de-Mouzon S, Nelson SM, Sattar N, Catalano PM and Freeman DJ. Lipotoxicity in obese pregnancy and its potential role in adverse pregnancy outcome and obesity in the offspring. *Clin Sci (Lond)*. 2010;119:123-9.

56. Simmons DG, Natale DR, Begay V, Hughes M, Leutz A and Cross JC. Early patterning of the chorion leads to the trilaminar trophoblast cell structure in the placental labyrinth. *Development*. 2008;135:2083-91.

57. Cross JC, Simmons DG and Watson ED. Chorioallantoic morphogenesis and formation of the placental villous tree. *Ann N Y Acad Sci*. 2003;995:84-93.

58. Saillenfait AM, Payan JP, Fabry JP, Beydon D, Langonne I, Gallissot F and Sabate JP. Assessment of the developmental toxicity, metabolism, and placental transfer of Di-n-butyl phthalate administered to pregnant rats. *Toxicol Sci*. 1998;45:212-24.

59. Clewell RA, Kremer JJ, Williams CC, Campbell JL, Jr., Andersen ME and Borghoff SJ. Tissue exposures to free and glucuronidated monobutylyphthalate in the pregnant and fetal rat following exposure to di-n-butylphthalate: evaluation with a PBPK model. *Toxicol Sci*. 2008;103:241-59.

60. Fennell TR, Krol WL, Sumner SC and Snyder RW. Pharmacokinetics of dibutylphthalate in pregnant rats. *Toxicol Sci*. 2004;82:407-18.

61. Thompson CJ, Ross SM and Gaido KW. Di(n-butyl) phthalate impairs cholesterol transport and steroidogenesis in the fetal rat testis through a rapid and reversible mechanism. *Endocrinology*. 2004;145:1227-37.

Figures and Figure Legends

Figure.1 Adult female CD-1 mice weight gain compared to control over 30 days of DBP exposure

at (A) 100, 10, and (B) 0.1 µg/kg/day. DBP100: 100µg/kg/day (n=8); DBP10: 10µg/kg/day (n=8);

DBP0.1: 0.1µg/kg/day (n=8); Control: Vehicle only (n=14). *: Differ from Control (P<0.05). Data

expressed as median ± standard error of mean. ᐃ: Differ between DBP100 and DBP10 (P<0.05).

Figure.2 Effect of DBP0.1 exposure on female fertility parameters ([A] pregnancy rate, [B] litter

size, [C] male/female fetus ratio per litter, and [D] pups to CLs ratio) in CD-1 mice. Data expressed

as median ± standard deviation. *: Differ between control (CT) and DBP0.1 groups (P<0.05). Each

data point represents the data from an individual dam. Sample size: (A) n=18 for each group; (B-

C) CT: n=12, DBP0.1: n=14; (D) CT: n=9; DBP0.1: n=13.

Figure.3 (A) Weight gain and (B) body fat % (fat mass/weight) of adult female CD-1 mice over 30

days of DBP0.1 exposure. CT: Control. (n= 6 dams per group). Data expressed as median ±

standard deviation. No statistical differences observed between CT and DBP dams. Each data

point represents the data from an individual dam. Sample size: (A) CT: n=15; DBP0.1: n=14; (B)

n=6 for each group; (C-F) CT: n=12; DBP n=14.

Figure.4 DBP0.1 exposure did not alter (A) fetal weight, while it significantly reduced (B) placental

weight and (C) fetal/placental weight ratio in female and male fetuses in CD-1 mice by GD18.5.

Data expressed as median ± standard deviation. *: P <0.05 (Two-way ANOVA). Median value of

all female or male fetuses from the same dam was shown as a single data point in figures. Sample

size: CT (n=12); DBP0.1 (n=14).

Figure.5 Lipid disposition in (A) fetal labyrinth and (B) maternal decidual of mice placental tissues

from CT and DBP groups. *: Differ between CT and DBP within the same fetal sex. Data

expressed as median ± standard deviation. $^{\Delta}$: Differ between female and male fetuses within the

same treatment group (P <0.05 Two-way ANOVA). The median value of all images from the same

placenta was shown as a single data point. Sample Size: CT.F (n=15); DBP.F (n=10); CT.M (n=

8); DBP.M (n=11).