Fes-deficient macrophages enhance CD8+ T cell priming and tumour control through increased proinflammatory cytokine production and presentation

Brian J. Laight[1,2], Natasha Dmytryk[1,2], Danielle Harper[1,2], Connie S. Zhang[1,2] Andrew Garven[1,2], Amber L. Simpson[3], David M. Berman[1,2,] Jacob Kment[1,3] Faizah Alotaibi[1,2,4], Changnian Shi[1], Yan Gao[1], Caitlyn Vlasschaert[5], David LeBrun[1,2], Kathrin Tyryshkin[2,6], Andrew W. Craig[1,3], Sameh Basta[1,3], Madhuri Koti[1,3], Peter A. Greer[1,2,3].

[1]Division of Cancer Biology and Genetics, Queen's Cancer Research Institute
[2]Department of Pathology and Molecular Medicine, Queen's University, Kingston, ON, Canada
[3]Department of Biomedical and Molecular Sciences, Queen's University, Kingston, ON, Canada
[4]King Saud bin Abdulaziz University for Heath Science, Riyadh, Saudia Arabia
[5]Department of Medicine, Queen's University, Kingston, ON, Canada
[6]School of Computing, Queen's University, Kingston, ON, Canada

ABSTRACT:

Activating the immune system is crucial for successful cancer immunotherapies, various proteins, such as the Fes non-receptor tyrosine kinase exist to limit activation and maintain homeostasis. However, in cancer settings, this serves as a barrier to the desired effects of immune activation following immunomodulatory treatment. Here, we demonstrate the role of Fes, a protein abundantly expressed in macrophages, as a novel innate intracellular immune checkpoint. Fes inactivity is associated with delayed tumour onset in a dose-dependent manner, and its deletion delays tumour growth, improves survival, enhances response to doxorubicin treatment, and sensitizes resistant tumours to PD-1 immune checkpoint inhibition. These effects are associated with an increase in Toll-like receptor signaling in antigen presenting cells, leading to an increase in proinflammatory cytokine production and cytotoxic T cell effector functions. Furthermore, we demonstrate a novel role for Fes in regulating the presentation of IL-12 on macrophage cell surfaces to enhance T-cell activation. Our results highlight Fes as a novel innate immune checkpoint with potential to serve as predictive biomarker to effective immune checkpoint blockade, and a potential novel therapeutic target for improved response to immunotherapy.

INTRODUCTION:

Cancer immunotherapy employs the patient's own immune system to help reduce tumour burden, and its use in the clinic has revolutionized patient care, highlighted by the use immune checkpoint inhibitors (ICI) (1, 2). The use and success of ICIs has primarily been focussed on enhancing T cell activity through inhibiting CTLA4 and PD-1 on the surface of T cells. Unfortunately, these inhibitors have not been one size fits all, and the heterogeneity of tumours and variety of infiltrating immune cell types (some immunosuppressive) provides both barriers to successful cancer immunotherapy(3), but also opportunities for improvement. Strategies to overcome the immune resistance exhibited by many tumour types include searching for novel immune checkpoints(4) or targeting different cell types, such as the cancer cells themselves, or other immune cells within the tumour microenvironment (TME)(5, 6).

One important class of immune cells involved in anti-cancer responses are antigen presenting cells (APCs), which play important roles including engulfing dying cancer cells and recruiting and priming cytotoxic T lymphocytes (CTLs). Many commonly used therapies are capable of inducing these anti-cancer immune responses, through a mechanism known as immunogenic cell death (ICD), such as anthracycline-based therapies (e.g., doxorubicin [Dox]) or radiotherapy(7). Upon induction of ICD, tumour associated antigens and damage associated molecular patterns (DAMPs) are released from dying cancer cells, activating an innate immune response in APCs which allows them to prime an anti-cancer adaptive immune response. For instance, the DNA-binding protein high mobility group box 1 (HMGB1) protein released from dying cancer cells during ICD binds to Toll-like receptor 4 (TLR) on APCs and activates a signaling cascade required for the production of pro-inflammatory cytokines, colloquially known as signal 3 (8-11). These pro-inflammatory cytokines play a crucial role in CTL priming, as their absence is associated with deficient CTL effector functions and a lack of memory formation(12, 13).

Because of the importance of APCs in initiating and sustaining adaptive anti-cancer immune responses, their inhibition or polarization to an immunosuppressive phenotype makes them a potential target for modulation. While pro-inflammatory cytokine production is one of the most important functions in priming anti-cancer immune responses, uncontrolled release of these cytokines can have detrimental and even fatal consequences, such as septic shock(14), auto-immune disorder(15), or cytokine storm(16). To combat this, the human body has evolved various mechanisms to attenuate immune responses to prevent these negative consequences. One such mechanism is through the non-receptor tyrosine kinase Fes, which is abundantly expressed in APCs(17)Fes limits TLR4 signaling(18), in part through facilitating TLR4 internalization and downregulation to prevent sustained stimulation(19). Fes has also been shown to associate with MHCI and suppress inflammatory signaling in macrophages through activation of the SH2 containing protein tyrosine phosphatase-2 (SHP2)(20). While mechanisms to suppress immune cell activation, such as PD-1, are important to maintain homeostatic conditions(21), in the context of cancer these mechanisms may enable tumour growth and limit the efficacy of treatments designed to activate anti-cancer immune responses(22).

Following previous observations implicating Fes in regulation of TLR internalization(19), involvement in vesicle trafficking(23), and signalling suppression through the MHC-I axis(20), we sought to explore its involvement in regulating anti-cancer immune responses. In this study, we demonstrate that Fes serves as an innate intracellular immune checkpoint. We show that Fes-deficiency is associated with delayed tumour initiation in a genetically engineered mouse model of breast cancer, which correlates with increased immune activity within the premalignant niche. In syngeneic engraftment models of triple negative breast cancer and melanoma, Fes-deficiency was associated with delayed tumour growth and improved survival, which was further enhanced by Dox treatment and immune checkpoint blockade (ICB) therapy. Tumours in Fes-deficient mice displayed increases in activated CTLs, which correlates with sensitization of tumours to ICB therapy. This increase in CTL activation can be attributed in part to an increased T cell priming capability of $fes^{-/-}$ bone-marrow derived macrophages (BMDMs) compared to their

wildtype counterparts, due in part to not only an increase in signal 3 cytokines production, namely interleukin-12 (IL-12), but also to the method of IL-12 presentation to CTLs through IL-12 membrane retention on the APCs. Taken together, these results demonstrate a novel role for Fes as an immune checkpoint, and a potential predictive biomarker for successful ICB therapy.

RESULTS:

Fes-deficiency delays tumour growth in a gene dose-dependent manner and is associated with increased immune infiltration in a transgenic mouse model of Her2+ breast cancer.

We previously showed that Fes-deficiency is associated with delayed tumour growth in an immunodeficient mouse engraftment model which lacks B and T cells(24). While that study implicated Fes-expressing macrophages in the modulation of tumourigenesis, we could not assess potential roles in the regulation of adaptive immunity. Here we have explored this question using immune competent models, beginning with a *MMTV-Neu* transgenic mouse model of HER2+ mammary tumourigenesis(25) crossed with transgenic mice targeted with a kinase-inactivating K588R knock-in mutation in *fes* (*fes^{KR}*). While the average tumour onset time in *fes$^{+/+}$* mice was 274 days, that was delayed to 316 days in *fes$^{KR/+}$* and 389 days in *fes$^{KR/KR}$* mice (Fig. 1A). These results demonstrated that a functional inhibition of 50% was sufficient to delay tumour onset, and that was even further delayed in mice lacking any Fes kinase activity. Interesting, once palpable tumours were detected, their growth rates were indistinguishable, suggesting tumour initiation had overcome some sort of suppression (Supplemental Figure 1). This observation correlates with a small magnitude, but significant effect of a single nucleotide polymorphism in human *FES* associated with differential Fes protein levels (rs17514846)(26), which predicts that higher Fes levels are correlated with higher cancer incidence rates (HR 1.03 [1.01 – 1.05], p=0.01) and risk of death (HR 1.06 [1.01 – 1.11], p=0.03) for those without cancer at baseline in the United Kingdom Biobank cohort of over 400,000 individuals.

To directly explore a potential role of the immune system on this delay in tumour onset, we isolated pre-malignant mammary glands (before palpable tumours could be detected) and performed immunohistochemical staining to assess the presence of myeloid and T cells. We found that there was an increase in both CD11b+ (Fig 1B,D) and CD3+ cells in *fes$^{KR/KR}$* glands (Fig.1C,E). Immunoblotting analysis of mammary gland lysates revealed an increase in phosphorylated NFκB in *fes$^{KR/KR}$* mice (Fig 1F,G); an important transcription factor for inflammatory signaling, including the expression of Signal 3 cytokines. Finally, an increase in immune cell infiltration was also apparent after tumours formed, where there was an increase in immune cell content relative to tumour volume (Fig 1H,I).

Fes-deficient mice have improved survival and delayed tumour growth in the presence or absence of doxorubicin treatment.

In the *MMTV-Neu* transgenic tumour-onset model, we saw that a reduction in functional Fes was associated with delayed tumour onset and an increase in immune cell infiltration and NFκB phosphorylation. This suggested a role for Fes in the regulation of anti-tumour immunity. To

further explore this, we employed the E0771 mammary carcinoma and B16F10 melanoma syngeneic orthotopic engraftment models using C57BL/6 mice targeted with a *fes* null mutation (*fes*$^{-/-}$) or wild type (*fes*$^{+/+}$) mice. Dox treatment was used in these studies to induce ICD and promote innate and adaptive anti-tumour immunity (Fig 2A,D). Tumour growth was significantly slower in *fes*$^{-/-}$ mice with or without Dox treatment, and Dox was more effective in *fes*$^{-/-}$ than *fes*$^{+/+}$ mice in both models (Fig 2B,E). Without Dox treatment, survival was significantly longer in *fes*$^{-/-}$ compared to *fes*$^{+/+}$ mice in both models (Fig 2C,F). Dox treatment prolonged survival of *fes*$^{-/-}$ mice in both models, but in *fes*$^{+/+}$ mice Dox treatment only improved survival in the B16F10 model. These observations are consistent with a Fes effect driven by the tumour immune microenvironment. Furthermore, neither E0771 nor B16F10 cells express detectable levels of Fes (Data not shown).

Fes-deficiency is correlated with an increased anti-tumoural immune microenvironment, enhanced by doxorubicin therapy.

Fes deficiency in cancer cell engraftment recipients was associated with slower tumour growth without treatment, and an additional delay in tumour growth and increased survival when treated with Dox (Fig. 2). These observations, coupled with delayed tumour onset and increased myeloid and T cell infiltration and NFκB activation in the pre-malignant niche of a Fes-deficient transgenic model of breast cancer (Fig.1), implicate Fes in regulating an anti-cancer immune response. This led us to next explore the tumour immune profiles of the E0771 and B16F10 engraftment models in *fes*$^{+/+}$ or *fes*$^{-/-}$ mice with or without Dox treatment (Fig 3A,H). There were no *fes* genome associated differences in the frequency of tumour associated CD8$^+$ T cells (CTLs) in either engraftment model (Fig 3B,I), and no differences in tumour associated CTL activation (CD69$^+$) between untreated and Dox treated *fes*$^{+/+}$ mice (Fig 3C,J). However, there was an increase in tumour associated CTL activation in untreated *fes*$^{-/-}$ mice compared to *fes*$^{+/+}$ mice in both models, which was further enhanced by Dox treatment (Fig 3C,J). In the spleens, *fes*$^{-/-}$ mice had higher levels of CTL activation (especially in the melanoma model), but this was not elevated by Dox treatment with either genotype or model (Supplemental Fig 2). Increased tumour associated CTL PD-1 positivity (indicative of more extensive and prolonged activation), was seen in *fes*$^{-/-}$ mice with Dox treatment in both models (Fig 3D,K). PD-1 positivity was inducible by Dox treatment in EO777 tumours only in *fes*$^{-/-}$ mice (Fig 3E); but was already higher in the B16F10 model in *fes*$^{-/-}$ mice without Dox treatment and was not further induced by Dox treatment (Fig 3K). In the spleen, there was a Dox-induced increase in PD-1 positivity only in *fes*$^{+/+}$ treated animals with the E0771 model (Supplemental Fig 2).

We next looked at the frequency of tumour associated macrophages and saw no *fes* genotype associated differences in either model (Fig 3E,L). However, we observed a *fes* genotype associated shift from immunosuppressive (CD206$^+$) macrophages (Fig 3F,M) to proinflammatory (CD206$^-$/CD80$^+$ and/or MHCII$^+$) macrophages (Fig 3G,N) in *fes*$^{-/-}$ mice in both engraftment models. This shift to more pro-inflammatory and less immunosuppressive macrophages was also seen in the spleen (Supplemental Fig 2). In both E0771 and B16F10 tumors, we saw no *fes* genotype associated changes in dendritic cell (DC) numbers (Fig 3N,P). However, in the E0771 model, we observed an increase in DC activation in *fes*$^{-/-}$ tumours (Fig

3O), but this was not observed in the B16F10 model (Fig 3Q). These results demonstrate that Fes deficiency is associated with increased CTL activation and PD-1 positivity, as well as an increase in pro-inflammatory macrophages and decrease in immunosuppressive macrophages in the tumor.

Fes deficient macrophages have increased TLR signaling, pro-inflammatory cytokine production and enhanced T cell priming functionality.

Fes displays tissue-specific expression, with highest levels observed in immune cells, primarily myeloid cells including antigen presenting macrophages and DCs, as well as NK and B cells; but Fes is not expressed in T cells. Since we see *fes* genome associated changes in the phenotype of tumour associated macrophages in both tumour models, we decided to explore the role of Fes in antigen presenting cells; specifically, macrophages since they made up most of the APCs within the tumour.

We have previously shown that Fes plays a role in downregulating inflammatory signaling in macrophages, in part through promoting LPS-induced endocytosis and downregulation of the TLR4 receptor, which suppresses TNFα expression(18). Additionally, it has been shown that Fes participates in suppressing inflammatory signaling in macrophages through association with MHC-I and phosphorylation of SHP2, which leads to suppression of TBK1 and NFκB pathways responsible for inflammatory cytokine expression(20). Here we explored the effects of Fes deficiency on the LPS-TLR4 signaling pathway in macrophages. To do this, we isolated BMDMs from $fes^{+/+}$ and $fes^{-/-}$ mice, treated them with 1µg/mL LPS and looked at downstream signaling. The $fes^{-/-}$ BMDMs displayed earlier and more robust activating phosphorylation of NFκB (Fig 4A,B) and TBK1 (Fig 4A,C) compared to $fes^{+/+}$ BMDMs. LPS-inducible activating phosphorylation of SHP-2 was apparent in the $fes^{+/+}$ BMDMs, while in $fes^{-/-}$ BMDMs, SHP2 displayed a baseline phosphorylation state that was not further induced by LPS (Fig 4A; Supplemental Fig3). We also observed increased NFκB and TBK1 phosphorylation in $fes^{-/-}$ bone marrow derived DCs compared with $fes^{+/+}$ DCs after LPS stimulation (Supplemental Fig 3). The $fes^{-/-}$ BMDMs were also more sensitive to LPS stimulation than $fes^{+/+}$, with increased NFκB and TBK1 phosphorylation even at concentration as low as 300pg/mL (Supplemental Fig 3E-G).

Seeing that Fes deficiency is associated with enhanced LPS-induced phosphorylation of NF-κB and TBK1, which play key roles in transcriptional upregulation of inflammatory cytokine genes, we next performed bulk RNAseq on $fes^{+/+}$ or $fes^{-/-}$ BMDMs before or after stimulation with LPS for 4 hours and performed a GO analysis on differentially expressed genes that met the following criteria: (1) were LPS inducible; (2) had at least a 1.25 fold increase in gene expression in LPS stimulated $fes^{-/-}$ compared to $fes^{+/+}$ BMDMs; and (3) the fold change associated with LPS stimulation was greater in $fes^{-/-}$ compared to $fes^{+/+}$BMDMs. The differentially expressed genes emerging from this analysis were significantly associated with increased inflammatory pathway activation and cytokine production, most specifically IL-12 and T cell activation pathways (Fig 4D), which was confirmed by looking at the fold change of several key genes following LPS stimulation (Fig 4E). When analyzed at the protein level by flow cytometry after LPS stimulation, cell-associated IL-12 was detected on more $fes^{-/-}$ than fes$^{+/+}$ BMDMs (Fig 4F); and

interestingly, this difference was even more apparent if the flow cytometry was restricted to assessing IL-12 on the external surface of nonpermeabilized cells (Fig 4G), suggesting that Fes plays a role in the trafficking of IL-12. We explored this further by immunofluorescence microscopy analysis of BMDMs in co-cultures with OT-I CD8+ T cells and observed co-localization of IL-12 and the IL-12 receptor on the surface of nonpermeabilized BMDMs (Fig 3H). Quantitation revealed higher IL-12 levels on the external surface of $fes^{-/-}$ compared to $fes^{+/+}$ BMDMs (Fig 3I).

These increases in TLR4 signaling, pro-inflammatory cytokine gene expression, and IL-12 on the surface of $fes^{-/-}$ BMDMs suggest that Fes deficient macrophages might be more effective at priming CTLs than their wildtype counterparts. To explore this, we pulsed $fes^{+/+}$ or $fes^{-/-}$ BMDMs with 10^{-10}M SIINFEKL peptide, and stimulated them with either the TLR4 agonist LPS (1µg/mL) or the TLR1/2 agonist Pam3Cys (0.1µg/mL) before co-culturing with splenic OT-1 CD8+ T cells in a 1:1 ratio. The $fes^{-/-}$ BMDMs activated a greater percentage of CD8$^+$ T cells both with and without LPS (Fig 4J) or Pam3Cys stimulation (Fig 4L). The $fes^{-/-}$ BMDMs also primed these CD8$^+$ T cells to a greater extent than $fes^{+/+}$ BMDMs in the presence of LPS (Fig 3K) and Pam3Cys (Fig 3M), as determined by the higher levels of IFNγ. This more efficient T cell priming by $fes^{-/-}$ BMDMs was preserved across a range of SIINFEKL concentrations (Supplemental Fig 3). However, there was no apparent fes genome associated difference in the ability of BMDMs to present the SIINFEKL peptide on MHCI (Supplemental Fig 3)

Fes-deficiency in BMDMs enabled them to prime CTLs more efficiently and to a greater extent, suggesting Fes-deficiency in APCs may allow them to promote more effective adaptive immunity in both tumour models. To directly show this, we performed adoptive macrophage transfer experiments with either $fes^{+/+}$ or $fes^{-/-}$ IFNγ-treated BMDMs injected directly into B16F10 engraftment tumours in $fes^{+/+}$ mice. Both $fes^{+/+}$ and $fes^{-/-}$ BMDMs reduced B16F10 tumour weights compared to mock injections with PBS, but $fes^{-/-}$ BMDMs were more effective at reducing tumour weights than $fes^{+/+}$ BMDMs (Fig 4N).

Tumours in Fes deficient mice are more sensitive to PD-1 therapy.

In light of our observation of increased PD-1 positivity on tumour associated CTLs in $fes^{-/-}$ mice with both engraftment models (Fig 3D,K), we decided to explore the effect of PD-1 therapy. Since CTLs in the E0771 model only show an increase in PD-1 positivity following Dox therapy, we treated all mice with Dox, and either an isotype antibody control (IgG) or anti-PD-1 antibody (Figure 5A, D). In both the E0771 and B16F10 models, Dox, and anti-PD-1 treated $fes^{+/+}$ mice show a decrease in tumour growth (Figure 5B, E), but no improvement in survival (Figure 5C, F) compared to Dox therapy alone, consistent with literature describing these tumour types as anti-PD-1 therapy resistant(27-32). Impressively, $fes^{-/-}$ mice treated only with Dox performed just as well (Fig 5B), or better (Fig 5E) than $fes^{+/+}$ treated with both Dox and anti-PD-1 therapy on delaying tumour growth. While anti-PD-1 therapy in $fes^{+/+}$ mice provided only a minimal tumour delay and no survival advantage, anti-PD-1 treated $fes^{-/-}$ mice demonstrated greater tumour delay and a greater survival advantage compared to all other treatment groups (Fig 5B,C,E,F). These observations demonstrated that Fes deficiency sensitizes these anti-PD-1 therapy resistant tumour types to this ICB treatment.

Discussion:

Here we provide novel evidence that the Fes non-receptor tyrosine kinase plays an intracellular innate immune checkpoint role in APCs that modulates anti-cancer adaptive immunity in part by regulating expression and presentation of the pro-inflammatory signal 3 cytokine IL-12 to CTLs. We show in a transgenic mouse breast cancer model that germline inactivating mutation in *fes* is associated with a delay in tumour onset. These *fes*[-/-] mice also displayed delayed tumour growth, enhanced Dox efficacy and improved response to anti-PD-1 ICB treatment in engraftment models of triple negative breast cancer and melanoma; and this correlated with increased T cell activation within the tumour and with increased pro-inflammatory signaling and IL-12 production in *fes*[-/-] macrophages.

In the transgenic tumour onset model, we see increased immune infiltration of CD3 and CD11b positive cells and inflammatory signaling (pNF-kB) correlating with delay in tumour onset. Inflammation can be a double-edged sword, where in some contexts it can induce tumour initiation and growth (chronic inflammation), and in others it can suppress tumour initiation and growth (acute inflammation) (33-35). It is well established that the immune system can suppress tumourigenesis until the cancer overcomes this through induction of immunosuppression(36). We first saw evidence that Fes might participate in immunosuppression in the transgenic tumour onset model, where delayed tumour onset correlated with increased density of CD3 and CD11b immune cells in the mammary glands prior to detection of palpable tumours in mice with targeted kinase-inactivating mutations in *fes*. However, once tumours were detected, their growth rates were the same across mouse *fes* genotypes, potentially demonstrating an adaptation of malignant cells to the tumour suppression. This could potentially explain the reduction in tumour incidence of people harbouring the rs17514846 SNP, which reduces Fes expression by ~50%(26).

Using BMDMs we demonstrated that *fes*[-/-] macrophages had increased inflammatory signaling as seen by the increase in phosphorylated (activated) NFκB and TBK1 downstream of TLR4 engagement by LPS. TBK1 goes on to phosphorylate and activate Interferon response factor 3 (IRF3). Activated IRF3 and NFκB are crucial for signal 3 cytokine production (e.g., type I IFNs and IL-12)(47). This was confirmed by RNAseq analysis showing that important signal 3 cytokines were transcriptionally upregulated following LPS stimulation to a higher degree in *fes*[-/-] compared to *fes*[+/+] BMDMs. Important signal 2 co-stimulatory molecules such as CD80, TL1A(48-50), and CD40(51, 52), were also expressed to higher levels in *fes*[-/-] BMDMs both in the presence and absence of LPS. These results may explain increased T cell activation by *fes*[-/-] BMDMs in the co-culture assay in the absence of LPS. The presence of increased co-stimulatory proteins and the enhanced sensitivity of *fes*[-/-] macrophages to respond to very low levels of LPS may potentially explain the already present increase in T cell activation in both E0771 and B16F10 models, and the ability for suboptimal induction of ICD to increase T cell activation in both models. During uninhibited tumour growth, there is still a degree of cell death that occurs(53, 54), which may produce sufficient DAMPs to activate innate immunity in *fes*[-/-] mice, but not *fes*[+/+] mice. This is supported not only by the increased sensitivity of *fes*[-/-] BMDMs to low concentrations of LPS, but also the enhanced CTL priming functionality at low antigen

levels. This may also explain how tumour initiation was delayed in the MMTV-Neu model, as the mice harbouring *fes* kinase-inactivating mutations not only had more CD11b myeloid cells within the premalignant glands, but macrophages or DCs among these cells may have had increased anti-cancer APC functionality. Additionally, the apparent increased functionality of *fes*$^{-/-}$ BMDMs over *fes*$^{+/+}$ was shown in the adoptive macrophage transfer. Cells from both genotypes were polarized to a proinflammatory phenotype by IFNγ stimulation; and while both genotypes reduced tumour volumes, the effect was greater in the mice which received *fes*$^{-/-}$ BMDMs. Given the results of the *in vitro* BMDM/T cell co-culture priming experiments, it is likely this improved tumour growth control was due to increased *in vivo* inflammatory cytokine presentation and T cell priming capabilities.

In BMDMs, IL-12-expressing *fes*$^{-/-}$ BMDMs were more abundant compared to *fes*$^{+/+}$, but what was most striking was the LPS-induced increase in membrane retained IL-12 in *fes*$^{-/-}$ BMDMs, which was not observed in *fes*$^{+/+}$ BMDMs. Not only were there more *fes*$^{-/-}$ cells expressing IL-12 on their surface, but the amount of IL-12 expressed on the surface, as determined by IF in the T cell co-culture assays, was also increased. This increase in cell-surface IL-12 may be partially responsible for improved T cell priming by *fes*$^{-/-}$ BMDMs. One group demonstrated that increased membrane-associated IL-12 improved DC-mediated priming of OT-II T cells, despite an apparent reduction in secreted IL-12(55). This suggests that membrane-associated IL-12 may be more important than secreted IL-12 in stimulating T cells. This provides a potential novel mechanism by which Fes might regulate T cell priming; by not only suppressing the expression of IL-12 following TLR stimulation, but also by promoting IL-12 secretion through a role in vesicular trafficking. Fes was previously reported to colocalize in the Golgi, as well as within cytoplasmic vesicles involved in both the endocytic and exocytic pathways(23). This is significant as IL-12 has also been shown to associated with Rab7 in late endosomes, and the deletion of the SNARE VAMP7, which is involved in exocytosis, caused a decrease in secreted IL-12(56). Future studies will be needed to explore potential trafficking interactions between Fes and IL-12 within APCs. Interestingly, the Fes-related kinase Fer has been shown to regulate ligand induced internalization of the EGFR(57), while Fes was implicated in regulating LPS-induced internalization of TLR4(19). These observations provide president for a role for this class of non-receptor tyrosine kinases in trafficking of signaling proteins.

Finally, an important finding for potential clinical applications is the effect of Fes deficiency on sensitivity to anti-PD-1 therapy. Immune checkpoint therapy has become one of the most widely used therapies in cancer, with over 1500 clinical trials using anti-PD-1 antibodies, and their approval as first-line treatment in certain cases (such as metastatic melanoma), and being used in TNBC in combination with chemotherapy in certain contexts (58). Metastatic melanoma has a response rate of ~32.6 – 45%, which represents one of the biggest successes of ICI therapy (59-61). Unfortunately, the response rate in TNBC remains low, from 5% in previously treated patients(62), up to 21% for first line therapy(63, 64). The combination of PD-1 therapy with chemotherapy shows improvements ranging from 25.8%(65), up to 64.8%(66) (albeit addition of PD-1 therapy improved outcomes by only 13.6% compared to chemotherapy alone). These results demonstrate that there is still significant room for improvement using combined chemoimmunotherapy. Another group showed that ICD induction with Dox, followed by

Nivolumab treatment results in a response rate of 35%, but most interestingly showed an increase in gene expression of immune-related genes involved in PD-1-PD-L1 and T cell cytotoxicity pathways(67). This reprogramming of the tumour microenvironment to one that is more inflamed and anti-tumoural, was recapitulated in our study, which showed increased T cell activation and PD-1 expression in $fes^{-/-}$, but not $fes^{+/+}$ mice. Furthermore, E0771 and B16F10 models are commonly reported in the literature as resistant to anti-PD-1 therapy(27-32, 68). This was seen in our studies where even in combination with Dox, anti-PD-1 elicited only a marginal improvement on tumour growth, and no improvement on survival in $fes^{+/+}$ mice in either model. However, the combination of Dox and anti-PD-1 therapy in $fes^{-/-}$ mice conferred a greater delay in tumour growth and a significant increase in survival compared to Dox alone. This demonstrated that host Fes-deficiency sensitizes these tumours to anti-PD-1 therapy, most likely because of increased T cell activation and PD-1 positivity seen in $fes^{-/-}$ tumours at baseline and following Dox therapy. This is reminiscent of the aforementioned combined chemotherapy plus PD-1 study of metastatic triple negative breast cancer which was reported in the TONIC clinical trial(67). These results highlight Fes as both a potential therapeutic target as well as a predictive biomarker for successful ICB.

Future work aims to explore the molecular mechanisms through which Fes regulates IL-12 expression, trafficking, secretion or presentation on the APC surface; correlating myeloid-specific Fes expression with patient outcomes through publicly available outcome data; and developing a novel and specific inhibitor for Fes with limited off-target effects. Currently, the ALK inhibitor Lorlatinib, developed for use in non small cell lung cancer has been shown to have off-target effects of inhibiting Fes and several other kinases(69). Due to this, an approach to selectively target Fes will be required.

In our study, we have shown that the non-receptor tyrosine kinase Fes, plays an important role in mediating successful anti-cancer immunotherapy. We believe that Fes falls into a novel group of immune checkpoints, we have coined as an innate intracellular immune checkpoint. Its deletion or inactivation leads to an increase in anti-tumour immune cell infiltration and activation within the tumour. This change in immune cell composition of the tumour correlates with increased tumour control and survival in both triple negative breast cancer and melanoma models, as well as sensitizing both models to additional anti-PD-1 ICB therapy, showing the role of Fes as a tumour agnostic checkpoint and identifying it as both a predictive biomarker for ICB therapy success, and an attractive therapeutic target.

Figure Captions:

Fig 1: Delay in tumour onset associated with *fes* inactivation and increased inflammation and immune infiltration into pre-malignant niche. (**A**) MMTV-Neu transgenic mice were crossed with either $fes^{+/+}$ mice (n=46), or with mice harbouring one ($fes^{KR/+}$) (n=47) or two ($fes^{KR/KR}$) (n=49) kinase-dead alleles, and tumour onset was assessed through manual palpation. Representative immunohistochemistry images and quantification of (**B,D**) CD11b staining in $fes^{+/+}$ (n=6) or $fes^{KR/KR}$ (n=6) mice or (**C,E**) CD3 staining in $fes^{+/+}$ (n=3) or 3 $fes^{KR/KR}$ (n=3) mice. (**F**) Pre-malignant glands were isolated from $fes^{+/+}$ (n=5) or $fes^{KR/KR}$ (n=5) mice and probed for pNFκB through western blot analysis and (**G**) densitometry assessed using imageJ. Mammary

glands containing similarly sized tumours were harvested from either $fes^{+/+}$ (n=11) or $fes^{KR/KR}$ (n=24) mice, **(H)** stained with H&E, and **(I)** were assessed by a blinded pathologist to assess stromal immune grading. Time-to-event (A) statistics were assessed through Log-rank (Mantel-Cox) test, and IHC and western blot statistical analysis was performed through an unpaired one-tailed t-test; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

Fig 2: Fes deficiency delays tumour growth, enhances survival, and improves doxorubicin effectiveness in a tumour agnostic manner. Schematic of the tumour growth and survival experiment for **(A)** E0771 and **(B)** B16F10 models. **(C,E)** E0771 cells or **(D,F)** B16F10 were orthotopically engrafted into either $fes^{+/+}$ or $fes^{-/-}$ mice (female for E0771 and male for B16F10), and treated with either 5% DMSO, or 5mg/kg Dox in the E0771 model, or 2.5mg/kg in the B16F10 model and were monitored using calipers to an endpoint of $1.1cm^3$ or physiological endpoint (e.g., ulceration) **(C,D)** and survival plotted **(E,F)**. Data represented as tumour volumes ± SEM of two independent experiments of both E0771 and B16F10 models with the following numbers of animals per group, respectively: $fes^{+/+}$ + DMSO (E0771 n=9, B16F10 n=10), $fes^{+/+}$ + Dox (E0771 n=10, B16F10 n=13), $fes^{-/-}$ + DMSO (E0771 n=9, B16F10 n=11), $fes^{-/-}$ + Dox (E0771 n=11, B16F10 n=12). Differences between tumour growth results were analyzed by two-way ANOVA with Tukey's multiple comparisons test and survival data was analyzed using Log-rank (Mantel-Cox) test; * p<0.05, ** p<0.01, *** p<0.001, ****p<0.0001.

Fig 3: Fes deficiency promotes a shift to an anti-tumour microenvironment and sensitizes tumours to doxorubicin-induced T cell activation. Schematic representation of immune profiling of **(A)** E0771 or **(B)** B16F10 tumours. Tumours were dissociated and stained for **(C,** E0771 n=14**; F,** B16F10 n=5) CD8+ T cells (CD45+, CD3+, CD8+; CTLs), **(D,** E0771 n=14**; G,** B16F10 n=5) CD69+ CTLs, **(E,** E0771 n=14**; H,** B16F10 n=5) PD-1+ CTLs, **(I,** E0771 n=18**; L,** B16F10 n=5) macrophages (CD45+, CD11b+, F4/80+), **(K,** E0771 n=18**; N,** B16F10 n=5) anti-inflammatory macrophages (CD206+), **(J,** E0771 n=18**; M,** B16F10 n=5) pro-inflammatory macrophages (CD206-, MHCII+ and/or CD80+), Results were analyzed by one-way ANOVA with Tukey's multiple comparisons test; * p<0.05, ** p<0.01, *** p<0.001, p<0.0001.

Fig 4: Fes deficiency is associated with a pro-inflammatory phenotype in bone marrow derived macrophages that correlates with enhanced T cell priming and tumour control through improved IL-12 signaling. (A-C) Bone marrow derived macrophages (BMDMs) were isolated from either $fes^{+/+}$ (n=5) or $fes^{-/-}$ (n=5) mice and stimulated with 1µg/mL LPS for the indicated time points before lysates were probed for (A) Fes, pNFκB, pTBK1, pSHP2 and GAPDH. Densitometry was assessed using ImageJ, and results were normalized to the band with the greatest density for either (B) pNFκB or (C) pTBK1 of the $fes^{+/+}$ BMDMs. **(D,E)** RNA was isolated from either $fes^{+/+}$ (n=3) or $fes^{-/-}$ (n=3) BMDMs that were either untreated or treated with 1µg/mL of LPS for 4 hours for bulk RNA sequencing. **(F,G)** BMDMs from either genotype were stimulated with 1µg/mL LPS for either 8 or 20hrs and stained either (F) intracellularly following permeabilization (n=3), or (G) extracellularly without permeabilization (n=3) for IL-12. **(H)** Representative images of BMDMs of either genotype were loaded with 10^{-10}M SIINFEKL and stimulated with 1µg/mL LPS for 4 hours prior to the addition of OT-I CTLs. After 30 mins of co-culture, cell conjugates were fixed and stained for CD3 (red), IL-12p35 (magenta), IL-12Rβ2

(green) and nuclear staining (blue). **(I)** IL-12 signal density per macrophage nuclei was measured using ImageJ analysis software from five different fields of view which included 47 $fes^{+/+}$ and 34 $fes^{-/-}$ macrophages. **(J-M)** BMDMs were pulsed with 10^{-10}M SIINFEKL peptide for 2 hours, washed, stimulated with the indicated (J,K) LPS or (L,M) Pam3Cys concentrations for 1 hour. OT-1 T cells were added to the macrophage culture in a 1:1 ratio for a total of 6 hours, with 5µg/mL of Brefeldin A added after 2 hours. Cells were then analyzed for (J,L) intracellular IFNγ positivity and (K,L) median fluorescence intensity (MFI). **(N)** B16F10 cells were orthotopically engrafted in $fes^{+/+}$ mice and when tumours reached 100mm^3 they were injected intratumourally with either PBS, or 3 million $fes^{+/+}$ or $fes^{-/-}$ BMDMs which had been stimulated for 16 hours with IFNγ. Nine days later, mice were euthanized, and tumours weights assessed, and normalized to the PBS control group. Figures B, C, F, G, J, K, L, M, N were analyzed by one-way ANOVA with Tukey's multiple comparisons test. Figure I was assessed using the Mann-Whitney test; * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

Fig 5: Fes deficiency enhances anti-PD-1 therapy and sensitizes previously resistant tumours. Schematic of tumour growth and survival using **(A)** E0771 and **(B)** B16F10 models. **(C,E)** E0771 cells or **(D,F)** B16F10 were orthotopically engrafted into either $fes^{+/+}$ or $fes^{-/-}$ mice (female for E0771 and male for B16F10), and all mice were treated with either 5mg/kg Dox (E0771 model), or 2.5mg/kg Dox (B16F10 model), and treated with either 200µg IgG antibody control, or 200µg anti-PD-1 antibody. Tumour growth (C,D) was monitored using calipers to an endpoint of 1.1cm^3 or humane endpoint (e.g., ulceration) and survival plotted (E,F). Data were represented as tumours volumes ± SEM of two (B16F10) or three (E0771) independent experiments with the following numbers of animals per group, respectively: $fes^{+/+}$ + Dox + αIgG (E0771 n=15, B16F10 n=12), $fes^{+/+}$ + Dox + αPD-1 (E0771 n=15, B16F10 n=12), $fes^{-/-}$ + Dox + αIgG (E0771 n=16, B16F10 n=13), $fes^{-/-}$ + Dox + αPD-1 (E0771 n=16, B16F10 n=14). Tumour growth results were analyzed by two-way ANOVA with Tukey's multiple comparisons test and survival data was analyzed using Gehan-Breslow-Wilcoxon Test for survival; * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

Supplemental Figure Captions

Supp Fig 1: MMTV induced tumours show no difference in growth rates. Tumours from either $fes^{+/+}$, $fes^{KR/+}$, or $fes^{KR/KR}$ mice were assessed by calipers until an endpoint of 1.1cm^3 or humane endpoint. Results were analyzed by unpaired one-tailed t-test, * p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

Supp Fig 2: Fes-deficiency and doxorubicin treatment induce tumoural and splenic changes in immune populations. Spleens were dissociated and stained for **(A,K)** CD8+ T cells (CD45+, CD3+, CD8+; CTLs), **(B,L)** CD69+ CTLs, **(C,M)** PD-1+ CTLs, **(D,N)** macrophages (CD45+, CD11b+, F4/80+), **(E,O)** M2 macrophages (CD206+), **(F,P)** M1 macrophages (CD206-, MHCII+ and/or CD80+), **(G,Q)** dendritic cells (F480-, MHCII+, CD11c+), **(H,R)** activated dendritic cells (CD80+). Tumours were also dissociated and stained for **(I,S)** dendritic cells, and **(J,T)** activated dendritic cells. Results were analyzed by one-way ANOVA with Tukey's multiple comparisons test; * p<0.05, ** p<0.01, *** p<0.001, p<0.0001.

Supp Fig 3: Fes-deficiency enhances inflammatory signaling in macrophages and dendritic cells. **(A)** Bone marrow derived macrophages (BMDMs) were isolated from $fes^{+/+}$ or $fes^{-/-}$ mice and treated with 1µg/mL LPS for the indicated time, assessed for pSHP2 by immunoblot analysis and densitometry assessed with ImageJ. **(B-D**) Bone marrow derived dendritic cells were isolated from $fes^{+/+}$ or $fes^{-/-}$ mice and treated with 1µg/mL LPS for the indicated time, assessed for pNFκB and pTBK1 by immunoblot analysis and densitometry assessed with ImageJ. **(E-G)** BMDMs were stimulated with the indicated LPS concentrations for 15 minutes before lysis and assessment for pNFκB and pTBK1 by immunoblot analysis and densitometry assessed with ImageJ. **(H)** BMDMs were pulsed with 10^{-10}M SIINFEKL for 2 hours, before stimulation with 1µg/mL LPS or PBS for 8 hours. BMDMs were then stained for MHCI-SIINFEKL protein and assessed by flow cytometry. BMDMs were pulsed with **(I)** 10^{-9}M, **(J)** 10^{-11}M, or **(K)** 10^{-12}M SIINFEKL peptide for 2 hours, washed, stimulated with the indicated LPS concentration for 1 hour. OT-1 T cells were added to the macrophage culture in a 1:1 ratio for a total of 6 hours, with 5µg/mL of Brefeldin A added after 2 hours. Cells were then analyzed for intracellular IFNγ positivity. Results were analyzed by one-way ANOVA with Tukey's multiple comparisons test; * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

Methods and Materials

Cells and cell culture conditions:

E0771 (CRL-3461), and 293T (CRL-3216) were purchased from the American Type Culture Collection (ATCC, Manassas, VA). B16F10 (CRL-6475) were a gift from Dr. Charles Graham (Kingston, Ontario, Canada). E0771, B16F10 and 293T cells were cultured in Dulbecco's Modified Eagle Medium (DMEM; Gibco cat#: 12100-061) supplemented with 10% fetal bovine serum (FBS; Corning Cat#:35-077-CV) 2mM L-glutamine (Corning; cat#:25-005-CI), and 1% antibiotic and antimycotics (A/A. Gibco; cat#:15240-062). Cell lines were cultured in an incubator held at 5% CO_2 and 37□. Cells used in all experiments were determined to be mycoplasma negative by the mycoplasma PCR detection kit.

Generation of bone marrow derived macrophages and dendritic cells:

Bone marrow derived macrophages (BMDMs) were isolated similarly to described previously(70, 71)In brief, on day 1, $fes^{+/+}$ or $fes^{-/-}$ (18)C57BL/6 mice (8 – 12 weeks old) were sacrificed and bone marrow flushed from both tibias and femurs in 5mL of Roswell Park Memorial Institute (RPMI) 1640 (Sigma-Aldrich; cat#:R8758-500mL) supplemented with 10% FBS and 1% A/A. Isolated bone marrow was resuspend by passing through a 23-gauge needle. Cells were passed through a 40µm cell strainer and pelleted at 1000rpm for 5 minutes. Cells were then resuspended in BMDM media (RPMI 1640 supplemented with 15% FBS, 1% A/A, 50µM α-monothioglycerate (α-MTG; Sigma; cat#:M-1753), and 20ng/mL macrophage colony-stimulating factor (M-CSF; a generous gift from Richard Stanley)(72)) and plated in 100mm cell culture plates. On day 2, non-adherent cells were collected, and pelleted at 1000rpm for 5 minutes. Cells were resuspended in 6mL of BMDM media and plated on a 60mm cell culture plate. On Day 4, non-adherent cells were collected, pelleted, resuspended in 10mL of BMDM media, and plated for onto untreated tissue culture plates. After 72 hours, cells were trypsonized,

counted, and plated at a concentration of 5e4 cells per well of a 96-well round bottom plate, or 1e6 cells/mL per well of a 6-well for all other experiments Bone marrow derived DCs were isolated as previously described (73) and cultured in DC media (15% FBS, 1% A/A, 50µM α-MTG, 20ng/mL granulocyte-macrophage colony stimulating factor (GM-CSF; Peprotech cat#: 315-03)

Immunoblotting analysis

Cells were washed twice with ice-cold phosphate buffered saline (PBS; 170mM NaCl, 3.5mM KCl, 10mM Na_2HPO_4, 2mM KH_2PO_4, pH 7.2) supplemented with 2mM sodium orthovanadate (ThermoFisher Scientific cat#: 205330500) and lysed in RIPA buffer (10 mM Tris pH7.2, 158 mM NaCl, 1 mM EDTA, 0.1% SDS, 1% Na Deoxycholate, 1% Triton-100) supplemented with Halt^{TM} Protease Inhibitor Cocktail (ThermoFisher Scientific cat#: 87785) and sodium orthovanadate. Protein quantification was performed using the Pierce^{TM} BCA Protein Assay Kit (ThermoFisher Scientific cat#: 23225) and immunoblotting was performed as previously described using the following antibodies and concentrations: Fes (1:1000; Cell Signaling Technologies cat#: 85704S), phospho-NF-κB (1:1000; Cell Signaling Technologies cat#: 3033S), , phospho-TBK1 (1:1000; Cell Signaling Technologies cat#: 5483), phospho-SHP2 (1:1000; Cell Signaling Technologies cat#:5431s), γ-tubulin (Sigma-Millipore, cat#: T5326), and GAPDH (1:50000; Cell Signaling Technologies cat#: 5174S).

BMDMs from fes+/+ and fes-/- mice were plated in 6-well flat bottom TC-treated plates (Corning, USA) at 2,000,000 cells per well in 2mL of complete macrophage media. The BMDMs were either left untreated, or they were stimulated with 1µg/mL LPS (Sigma-Aldrich, USA) for 4 hours. Cells were then lysed with 350µL of cold TRI Reagent® (Sigma-Aldrich, USA) for five minutes. RNA purification was performed according to Invitrogen's TrizolTM Regent protocol. Purified RNA sample concentrations were determined with a NanoDrop ND-1000 spectrophotometer for downstream cDNA synthesis. cDNA synthesis was carried out using the Maxima First Strand cDNA synthesis kit (Thermo Fisher Scientific, USA) according to manufacturer's instructions, with 372ng of RNA as the starting template. Samples were mixed, centrifuged, and placed in an Applied Biosystems thermocycler (Life Technologies, USA) for one cycle for 10 minutes at 25°C, 15 minutes at 50°C and finally 5 minutes at 85°C. RNA quality and quantity were confirmed via NanoDrop measurement. All samples possessed an A260/280 purity ratio >1.7 and were sent to Genome Quebec (Montreal, Canada) for poly-A enriched RNA-sequencing. Genome Quebec conducted an alignment of the sequencing data to the GRCm38 (Ensembl 102 release) murine genome to determine raw expression gene counts and transcripts per million (TPM) counts. The data was then pre-processed by removing genes with zero counts across all samples and filtering out genes with expression levels below the 75th percentile threshold (roughly 12 counts). The resulting pre-processed TPM counts for each gene were plotted for each genotype condition.

Macrophage and T cell coculture:

BMDMs were harvested as described above and were seeded in a 96-well plate. BMDMs were then treated for 2 hours with varying concentrations of OVA peptide (SIINFEKL: CPC Scientific

cat#: MISC-012). BMDMs were then washed three times with serum free RPMI, before treatment with1µg/mL of Lipopolysaccharides (LPS; *E. coli*, serotype 055:B5, Sigma Aldrich cat#: L2637), for 1 hour.cat#:. BMDMs were then washed three times again with serum free RPMI. Spleens from OT-1 mice were isolated, manually dissociated, filtered through a 40µm filter, and subjected to a CD8$^+$ T cell negative selection with the EasySep^{TM} Mouse T cell isolation kit (STEMCELL Technologies cat#: 19853), according to the manufacturer's protocol. Isolated CD8$^+$ T cells were seeded onto treated BMDMs in a ratio of 1:1 resuspended in T cell media (RPMI 1640, 10% FBS, 1% A/A, 1% gentamicin (ThermoFisher Scientific cat#: 15710064) and incubated for 2 hours after which cells were treated with 1x Brefeldin A (BioLegend; cat#: 420601) and incubated for a final 4 hours before analysis by flow cytometry for T cell activation.

Animal Studies:

All experiments involving the use of animals were performed in accordance with the Canadian Council on Animal Care guidelines and with the approval of Queen's University Animal Care Committee. All mouse strains (*fes*$^{KR/KR}$(74), *fes*$^{-/-}$ (18)and OT-1 (C57BL/6-Tg(TcraTcrb)1100Mjb/J, https://www.jax.org/strain/003831) were on a C57BL/6 background. Mice were bred in-house and 8–12-week-old males or females were used in engraftment studies for B16F10 or E0771 models, respectively with drug and antibody doses normalized to body weights. Animals were randomized at treatment start time, with any mice lacking tumour growth excluded from the study. Tumour length (*L*) and width (*W*) were measured by caliper, and tumour volume calculated with the following formula: (L x W^2)/2. Tumours were measured until an endpoint of 1.1cm^3 or human endpoint (e.g., ulceration) at which point mice were euthanized and survival analyzed.

Doxorubicin therapy model:

C57BL/6 mice (8-12 weeks old) were orthotopically engrafted with 1x10^5 E0771 cells injected into the fourth mammary fat pad with a Hamilton syringe, or with 6x10^4 B16F10 cells in a 1:1 ratio of Matrigel (Corning cat#: 354234) to PBS cell suspension injected subcutaneously with a 26-gauge needle into the left flank. Once tumours reached ~100mm^3 (E0771 model: *fes*$^{+/+}$ ~12 days; *fes*$^{-/-}$ ~14 days)(B16F10 model: *fes*$^{+/+}$ ~8 days; *fes*$^{-/-}$ ~10 days), E0771 tumour-bearing mice were treated with either 5mg/kg of Dox (Selleckchem cat#: S120813) or vehicle control in a 200µL 5% DMSO/PBS solution injected intraperitoneally (IP), repeated 5 and 10 days later; or for the B16F10 model, 2.5mg/kg Dox or vehicle control in a 200µL 5% DMSO/PBS solution injected IP every 2 days to a final cumulative dose of 20mg/kg. Tumour growth and survival were analyzed as described above.

Anti-PD-1 immune checkpoint blockade therapy model:

The anti-PD-1 treatment model followed a similar protocol to that described above; however, every mouse was treated with its tumour-specific Dox dose in combination with either 10mg/kg anti-IgG control antibody (BioXCell; cat#: BE0089) or 10mg/kg anti-PD-1 antibody (BioXCell; cat#:BE0146) IP, for a total of three doses, either with every Dox treatment in the E0771 model or every two Dox treatments starting on the third Dox treatment in the B16F10 model.

Harvesting tumours and spleens for tumour immune profiling

Mice were engrafted with cancer cells and treated as described above. However, three days following the second dose of Dox treatment in the E0771 model, or one day after the fourth dose in the B16F10 model, mice were euthanized, and tumours and spleens harvested. Tumours were cut into ~1mm^3 chunks with a straight razor, and spleens were pressed through a sieve with a syringe plunger. Tumour samples were suspended in RPMI 1640 supplemented with 10% FBS, 1mg/mL collagenase B (Sigma-Aldrich cat#: 11088815001) and 25µg/mL of DNAse I (Sigma-Aldrich Cat#: 10104159001), while spleens were resuspended in RPMI containing only 25µg/mL of DNAse I. Tumour chunks were then transferred into C Tubes (Miltenyi Biotec cat#: 130-096-334) and dissociated using the gentleMACS Dissociator following the manufacturer's protocol for tough (E0771) or soft (B16F10) tumours. Tumour and spleen homogenates were then filtered through a 70µM cell strainer followed by a 40µM cell strainer and then treated with Ammonia Chloride Potassium (ACK) lysis buffer to lyse red blood cells within the preparation. Cells were then washed and resuspended in fluorescence-activated cell sorting (FACS) buffer (PBS containing 0.75% bovine serum albumin (BSA; BioShop cat#: ALB001), 1mM EDTA in PBS) in preparation for flow staining.

Antibody staining and flow cytometry

Once cells of interested were harvested, 1x10^5 cells/well were plated into 96-well round bottom plates and centrifuged at 1000rpm for 5 minutes. Supernatants were carefully decanted, and cell pellets resuspended in 100µL FACS containing 0.2µg TruStain FcX™ (anti-mouse CD16/32) antibody or 100µL of PBS containing 0.2µg TruStain FcX™ and 0.04µL Zombie AquaTM Fixable Viability Kit (BioLegend cat#: 423102) and incubated at room temperature (RT) in the dark for 30 minutes. Next, 200µL of FACS buffer was added to each well, gently mixed, and pelleted at 1000rpm for 5 minutes, and carefully decanted. Next, cells were stained by resuspending in 100µL of FACS buffer containing appropriate antibody panels and incubated on ice in the dark for 30 minutes. 200µL of FACS was added to each well, gently mixed, and pelleted at 1000rpm for 5 minutes, and carefully decanted, followed by an additional wash in 200µL FACS buffer. If the cells were not being subjected to intracellular staining, cells were resuspended in 100µL of 1% paraformaldehyde in PBS and incubated at room temperature for 30 minutes. An additional 200µL of FACS buffer was added, cells pelleted at 1500rpm for 5 minutes, decanted, and resuspended in 200µL FACS buffer to be analyzed on a CytoFlex-S Flow Cytometer (Beckman Coulter). For intracellular staining, cells were stained following the ThermoFisher "BestProtocols: Staining Intracellular Antigens for Flow Cytometry – Protocol A" using the eBioscience™ Permeabilization Buffer (ThermoFisher Scientific cat#: 00-8333-56) and eBioscience™ IC Fixation Buffer (ThermoFisher Scientific cat#: 00-8222-49) before analysis on the CytoFlex-S Flow Cytometer (Beckman Coulter). Data analysis was performed using FlowJo® v10 software (FlowJo, LLC, Ashland, OR).

Macrophage adoptive transfer:

C57BL/6 *fes*$^{+/+}$ mice (8-12 weeks old) were engrafted with B16F10 cells as described above. BMDMs for adoptive transfer were isolated from 8-12 week old *fes*$^{+/+}$ or *fes*$^{-/-}$ mice as described above, except the day 4 adherent BMDMs were trypsonized and re-plated onto non-tissue culture grade plates for 48 hrs, then treated overnight with 20ng/mL interferon gamma (IFN-γ;

Peprotech cat#: 315-05) to polarize them into an M1-like state. These were harvested and 3×10^6 *fes*$^{+/+}$ or *fes*$^{-/-}$ M1-polarized BMDMs in 50µL of PBS, or PBS alone were injected intratumourally (IT) with a 26-gauge needle into 100mm^3 B16F10 tumours. Nine days later, mice were euthanized, and tumour weights determined.

References:

1. Pardoll DM. The blockade of immune checkpoints in cancer immunotherapy. Nature Reviews Cancer. 2012;12(4):252-64.

2. Postow MA, Callahan MK, Wolchok JD. Immune Checkpoint Blockade in Cancer Therapy. J Clin Oncol. 2015;33(17):1974-82.

3. Zhao X, Subramanian S. Intrinsic Resistance of Solid Tumors to Immune Checkpoint Blockade Therapy. Cancer Research. 2017;77(4):817-22.

4. Andrews LP, Marciscano AE, Drake CG, Vignali DAA. LAG3 (CD223) as a cancer immunotherapy target. Immunological Reviews. 2017;276(1):80-96.

5. Liu J, Chen Z, Li Y, Zhao W, Wu J, Zhang Z. PD-1/PD-L1 Checkpoint Inhibitors in Tumor Immunotherapy. Frontiers in Pharmacology. 2021;12.

6. Khan M, Arooj S, Wang H. NK Cell-Based Immune Checkpoint Inhibition. Front Immunol. 2020;11:167.

7. Zhou J, Wang G, Chen Y, Wang H, Hua Y, Cai Z. Immunogenic cell death in cancer therapy: Present and emerging inducers. Journal of Cellular and Molecular Medicine. 2019;23(8):4854-65.

8. Ahmed A, Tait SWG. Targeting immunogenic cell death in cancer. Molecular Oncology. 2020;14(12):2994-3006.

9. Mirlekar B, Pylayeva-Gupta Y. IL-12 Family Cytokines in Cancer and Immunotherapy. Cancers. 2021;13(2):167.

10. Boukhaled GM, Harding S, Brooks DG. Opposing Roles of Type I Interferons in Cancer Immunity. Annu Rev Pathol. 2021;16:167-98.

11. Lu C, Klement JD, Ibrahim ML, Xiao W, Redd PS, Nayak-Kapoor A, et al. Type I interferon suppresses tumor growth through activating the STAT3-granzyme B pathway in tumor-infiltrating cytotoxic T lymphocytes. Journal for ImmunoTherapy of Cancer. 2019;7(1):157.

12. Curtsinger JM, Lins DC, Mescher MF. Signal 3 Determines Tolerance versus Full Activation of Naive CD8 T Cells. 2003;197(9):1141-51.

13. Zagorulya M, Yim L, Morgan DM, Edwards A, Torres-Mejia E, Momin N, et al. Tissue-specific abundance of interferon-gamma drives regulatory T cells to restrain DC1-mediated priming of cytotoxic T cells against lung cancer. Immunity. 2023;56(2):386-405.e10.

14. Laight BJ, Jawa NA, Tyryshkin K, Maslove DM, Boyd JG, Greer PA. Establishing the role of the FES tyrosine kinase in the pathogenesis, pathophysiology, and severity of sepsis and its outcomes. Frontiers in Immunology. 2023;14.

15. Salvi V, Gianello V, Tiberio L, Sozzani S, Bosisio D. Cytokine Targeting by miRNAs in Autoimmune Diseases. Frontiers in Immunology. 2019;10.

16. Fajgenbaum DC, June CH. Cytokine Storm. New England Journal of Medicine. 2020;383(23):2255-73.

17. Haigh J, McVeigh J, Greer P. The fps/fes tyrosine kinase is expressed in myeloid, vascular endothelial, epithelial, and neuronal cells and is localized in the trans-golgi network. Cell Growth Differ. 1996;7(7):931-44.

18. Zirngibl RA, Senis Y, Greer PA. Enhanced Endotoxin Sensitivity in Fps/Fes-Null Mice with Minimal Defects in Hematopoietic Homeostasis. Molecular and Cellular Biology. 2002;22(8):2472-86.

19.   Parsons SA, Greer PA. The Fps/Fes kinase regulates the inflammatory response to endotoxin through down-regulation of TLR4, NF-κB activation, and TNF-α secretion in macrophages. Journal of Leukocyte Biology. 2006;80(6):1522-8.

20.   Xu S, Liu X, Bao Y, Zhu X, Han C, Zhang P, et al. Constitutive MHC class I molecules negatively regulate TLR-triggered inflammatory responses via the Fps–SHP-2 pathway. Nature Immunology. 2012;13(6):551-9.

21.   Chen RY, Zhu Y, Shen YY, Xu QY, Tang HY, Cui NX, et al. The role of PD-1 signaling in health and immune-related diseases. Front Immunol. 2023;14:1163633.

22.   Hanahan D, Weinberg Robert A. Hallmarks of Cancer: The Next Generation. Cell. 2011;144(5):646-74.

23.   Zirngibl R, Schulze D, Mirski SEL, Cole SPC, Greer PA. Subcellular Localization Analysis of the Closely Related Fps/Fes and Fer Protein-Tyrosine Kinases Suggests a Distinct Role for Fps/Fes in Vesicular Trafficking. Experimental Cell Research. 2001;266(1):87-94.

24.   Zhang S, Chitu V, Stanley ER, Elliott BE, Greer PA. Fes Tyrosine Kinase Expression in the Tumor Niche Correlates with Enhanced Tumor Growth, Angiogenesis, Circulating Tumor Cells, Metastasis, and Infiltrating Macrophages. Cancer Research. 2011;71(4):1465-73.

25.   Muller WJ, Sinn E, Pattengale PK, Wallace R, Leder P. Single-step induction of mammary adenocarcinoma in transgenic mice bearing the activated c-neu oncogene. Cell. 1988;54(1):105-15.

26.   Karamanavi E, McVey DG, van der Laan SW, Stanczyk PJ, Morris GE, Wang Y, et al. The FES Gene at the 15q26 Coronary-Artery-Disease Locus Inhibits Atherosclerosis. Circ Res. 2022;131(12):1004-17.

27.   Kleffel S, Posch C, Barthel SR, Mueller H, Schlapbach C, Guenova E, et al. Melanoma Cell-Intrinsic PD-1 Receptor Functions Promote Tumor Growth. Cell. 2015;162(6):1242-56.

28.   Tomita M, Yasui H, Higashikawa K, Nakajima K, Takakura H, Shiga T, et al. Anti PD-1 treatment increases [18F]FDG uptake by cancer cells in a mouse B16F10 melanoma model. EJNMMI Research. 2018;8(1).

29.   Waaler J, Mygland L, Tveita A, Strand MF, Solberg NT, Olsen PA, et al. Tankyrase inhibition sensitizes melanoma to PD-1 immune checkpoint blockade in syngeneic mouse models. Communications Biology. 2020;3(1).

30.   Nakamura T, Sato T, Endo R, Sasaki S, Takahashi N, Sato Y, et al. STING agonist loaded lipid nanoparticles overcome anti-PD-1 resistance in melanoma lung metastasis via NK cell activation. Journal for ImmunoTherapy of Cancer. 2021;9(7):e002852.

31.   Borchmann S, Selenz C, Lohmann M, Ludwig H, Gassa A, Brägelmann J, et al. Tripartite antigen-agnostic combination immunotherapy cures established poorly immunogenic tumors. Journal for ImmunoTherapy of Cancer. 2022;10(10):e004781.

32.   Wculek SK, Amores-Iniesta J, Conde-Garrosa R, Khouili SC, Melero I, Sancho D. Effective cancer immunotherapy by natural mouse conventional type-1 dendritic cells bearing dead tumor antigen. Journal for ImmunoTherapy of Cancer. 2019;7(1).

33.   Coussens LM, Werb Z. Inflammation and cancer. Nature. 2002;420(6917):860-7.

34.   Greten FR, Grivennikov SI. Inflammation and Cancer: Triggers, Mechanisms, and Consequences. Immunity. 2019;51(1):27-41.

35.   Zhao H, Wu L, Yan G, Chen Y, Zhou M, Wu Y, Li Y. Inflammation and tumor progression: signaling pathways and targeted intervention. Signal Transduction and Targeted Therapy. 2021;6(1):263.

36.   Gonzalez H, Hagerling C, Werb Z. Roles of the immune system in cancer: from tumor initiation to metastatic progression. Genes Dev. 2018;32(19-20):1267-84.

37.   Paridaens R, Biganzoli L, Bruning P, Klijn JGM, Gamucci T, Houston S, et al. Paclitaxel Versus Doxorubicin as First-Line Single-Agent Chemotherapy for Metastatic Breast Cancer: A European Organization for Research and Treatment of Cancer Randomized Study With Cross-Over. Journal of Clinical Oncology. 2000;18(4):724-.

38.   Ugurel M, Schadendorf D, Fink W, Zimpfer-Rechner C, Thoelke A, Figl R, Kaatz M. Clinical Phase II Study of Pegylated Liposomal Doxorubicin as Second-Line Treatment in Disseminated Melanoma. Onkologie. 2004;27(6):540-4.

39.   Smylie MG, Wong R, Mihalcioiu C, Lee C, Pouliot J-F. A phase II, open label, monotherapy study of liposomal doxorubicin in patients with metastatic malignant melanoma. Investigational New Drugs. 2007;25(2):155-9.

40.   Casares N, Pequignot MO, Tesniere A, Ghiringhelli Fo, Roux Sp, Chaput N, et al. Caspase-dependent immunogenicity of doxorubicin-induced tumor cell death. Journal of Experimental Medicine. 2005;202(12):1691-701.

41.   Kroemer G, Galluzzi L, Kepp O, Zitvogel L. Immunogenic Cell Death in Cancer Therapy. Annual Review of Immunology. 2013;31(1):51-72.

42.   Galluzzi L, Buqué A, Kepp O, Zitvogel L, Kroemer G. Immunogenic cell death in cancer and infectious disease. Nature Reviews Immunology. 2017;17(2):97-111.

43.   Ansar SM, Mudalige T. Characterization of doxorubicin liposomal formulations for size-based distribution of drug and excipients using asymmetric-flow field-flow fractionation (AF4) and liquid chromatography-mass spectrometry (LC-MS). International Journal of Pharmaceutics. 2020;574:118906.

44.   Franco Y, Vaidya T, Ait-Oudhia S. Anticancer and cardio-protective effects of liposomal doxorubicin in the treatment of breast cancer. Breast Cancer: Targets and Therapy. 2018;Volume 10:131-41.

45.   Merino M, Lozano T, Casares N, Lana H, Troconiz IF, ten Hagen TLM, et al. Dual activity of PD-L1 targeted Doxorubicin immunoliposomes promoted an enhanced efficacy of the antitumor immune response in melanoma murine model. Journal of Nanobiotechnology. 2021;19(1):102.

46.   Monteran L, Ershaid N, Doron H, Zait Y, Scharff Ye, Ben-Yosef S, et al. Chemotherapy-induced complement signaling modulates immunosuppression and metastatic relapse in breast cancer. Nature Communications. 2022;13(1):5797.

47.   Liu T, Zhang L, Joo D, Sun SC. NF-κB signaling in inflammation. Signal Transduct Target Ther. 2017;2:17023-.

48.   Migone TS, Zhang J, Luo X, Zhuang L, Chen C, Hu B, et al. TL1A is a TNF-like ligand for DR3 and TR6/DcR3 and functions as a T cell costimulator. Immunity. 2002;16(3):479-92.

49.   Papadakis KA, Prehn JL, Landers C, Han Q, Luo X, Cha SC, et al. TL1A synergizes with IL-12 and IL-18 to enhance IFN-gamma production in human T cells and NK cells. J Immunol. 2004;172(11):7002-7.

50.   Twohig JP, Marsden M, Cuff SM, Ferdinand JR, Gallimore AM, Perks WV, et al. The death receptor 3/TL1A pathway is essential for efficient development of antiviral CD4+ and CD8+ T-cell immunity. Faseb j. 2012;26(8):3575-86.

51.   Vonderheide RH. CD40 Agonist Antibodies in Cancer Immunotherapy. Annual Review of Medicine. 2020;71(1):47-58.

52.   Liu P-S, Chen Y-T, Li X, Hsueh P-C, Tzeng S-F, Chen H, et al. CD40 signal rewires fatty acid and glutamine metabolism for stimulating macrophage anti-tumorigenic functions. Nature Immunology. 2023;24(3):452-62.

53.   Labi V, Erlacher M. How cell death shapes cancer. Cell Death & Disease. 2015;6(3):e1675-e.

54.   Nössing C, Ryan KM. 50 years on and still very much alive: 'Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics'. British Journal of Cancer. 2023;128(3):426-31.

55.   Akhter N, Takeda Y, Nara H, Araki A, Ishii N, Asao N, Asao H. Apurinic/Apyrimidinic Endonuclease 1/Redox Factor-1 (Ape1/Ref-1) Modulates Antigen Presenting Cell-mediated T Helper Cell Type 1 Responses. The Journal of biological chemistry. 2016;291(45):23672-80.

56.   Chiaruttini G, Piperno GM, Jouve M, De Nardi F, Larghi P, Peden AA, et al. The SNARE VAMP7 Regulates Exocytic Trafficking of Interleukin-12 in Dendritic Cells. Cell Rep. 2016;14(11):2624-36.

57.   Sangrar W, Shi C, Mullins G, LeBrun D, Ingalls B, Greer PA. Amplified Ras-MAPK signal states correlate with accelerated EGFR internalization, cytostasis and delayed HER2 tumor onset in Fer-deficient model systems. Oncogene. 2015;34(31):4109-17.

58.   Gaynor N, Crown J, Collins DM. Immune checkpoint inhibitors: Key trials and an emerging role in breast cancer. Seminars in Cancer Biology. 2022;79:44-57.

59.   Wolchok JD, Chiarion-Sileni V, Gonzalez R, Grob J-J, Rutkowski P, Lao CD, et al. Long-Term Outcomes With Nivolumab Plus Ipilimumab or Nivolumab Alone Versus Ipilimumab in Patients With Advanced Melanoma. Journal of Clinical Oncology. 2022;40(2):127-37.

60.   Long GV, Stephen Hodi F, Lipson EJ, Schadendorf D, Ascierto PA, Matamala L, et al. Overall Survival and Response with Nivolumab and Relatlimab in Advanced Melanoma. NEJM Evidence. 2023;2(4).

61.   Larkin J, Chiarion-Sileni V, Gonzalez R, Grob J-J, Rutkowski P, Lao CD, et al. Five-Year Survival with Combined Nivolumab and Ipilimumab in Advanced Melanoma. New England Journal of Medicine. 2019;381(16):1535-46.

62.   Adams S, Schmid P, Rugo HS, Winer EP, Loirat D, Awada A, et al. Pembrolizumab monotherapy for previously treated metastatic triple-negative breast cancer: cohort A of the phase II KEYNOTE-086 study. Ann Oncol. 2019;30(3):397-404.

63.   Adams S, Loi S, Toppmeyer D, Cescon DW, De Laurentiis M, Nanda R, et al. Pembrolizumab monotherapy for previously untreated, PD-L1-positive, metastatic triple-negative breast cancer: cohort B of the phase II KEYNOTE-086 study. Annals of Oncology. 2019;30(3):405-11.

64.   Nanda R, Chow LQ, Dees EC, Berger R, Gupta S, Geva R, et al. Pembrolizumab in Patients With Advanced Triple-Negative Breast Cancer: Phase Ib KEYNOTE-012 Study. J Clin Oncol. 2016;34(21):2460-7.

65.   Tolaney SM, Kalinsky K, Kaklamani VG, D'Adamo DR, Aktan G, Tsai ML, et al. Eribulin Plus Pembrolizumab in Patients with Metastatic Triple-Negative Breast Cancer (ENHANCE 1): A Phase Ib/II Study. Clinical cancer research : an official journal of the American Association for Cancer Research. 2021;27(11):3061-8.

66.   Schmid P, Cortes J, Pusztai L, McArthur H, Kümmel S, Bergh J, et al. Pembrolizumab for Early Triple-Negative Breast Cancer. New England Journal of Medicine. 2020;382(9):810-21.

67.   Voorwerk L, Slagter M, Horlings HM, Sikorska K, van de Vijver KK, de Maaker M, et al. Immune induction strategies in metastatic triple-negative breast cancer to enhance the sensitivity to PD-1 blockade: the TONIC trial. Nature Medicine. 2019;25(6):920-8.

68.   Crosby EJ, Wei J, Yang XY, Lei G, Wang T, Liu CX, et al. Complimentary mechanisms of dual checkpoint blockade expand unique T-cell repertoires and activate adaptive anti-tumor immunity in triple-negative breast tumors. Oncoimmunology. 2018;7(5):e1421891.

69.   Zou HY, Li Q, Engstrom LD, West M, Appleman V, Wong KA, et al. PF-06463922 is a potent and selective next-generation ROS1/ALK inhibitor capable of blocking crizotinib-resistant ROS1 mutations. Proc Natl Acad Sci U S A. 2015;112(11):3493-8.

70.   Stanley ER. Murine Bone Marrow-Derived Macrophages. In: Pollard JW, Walker JM, editors. Basic Cell Culture Protocols. Totowa, NJ: Humana Press; 1997. p. 301-4.

71.   Weischenfeldt J, Porse B. Bone Marrow-Derived Macrophages (BMM): Isolation and Applications. Cold Spring Harbor Protocols. 2008;2008(12):pdb.prot5080.

72.   Ladner MB, Martin GA, Noble JA, Wittman VP, Warren MK, McGrogan M, Stanley ER. cDNA cloning and expression of murine macrophage colony-stimulating factor from L929 cells. Proc Natl Acad Sci U S A. 1988;85(18):6706-10.

73.   Oliveira MMS, D'Aulerio R, Yong T, He M, Baptista MAP, Nylén S, Westerberg LS. Increased cross-presentation by dendritic cells and enhanced anti-tumour therapy using the Arp2/3 inhibitor CK666. British Journal of Cancer. 2023;128(6):982-91.

74.	Senis Y, Zirngibl R, McVeigh J, Haman A, Hoang T, Greer PA. Targeted disruption of the murine fps/fes proto-oncogene reveals that Fps/Fes kinase activity is dispensable for hematopoiesis. Mol Cell Biol. 1999;19(11):7436-46.

Figure 3.

Figure 5.