The effect of molnupiravir and nirmatrelvir on SARS-CoV-2

genome diversity in severe models of COVID-19.

Rebekah Penrice-Randal[1*†], Eleanor G. Bentley[1*], Parul Sharma[1], Adam Kirby[1], I'ah

Donovan-Banfield[1,2], Anja Kipar[1,3], Daniele F. Mega[1] Chloe Bramwell[1,4], Joanne

Sharp[4,5], Andrew Owen[4,5], Julian A. Hiscox[1,2,6] and James P. Stewart[1,5].

[1]Department of Infection Biology and Microbiomes, University of Liverpool, Liverpool,

UK.

[2]NIHR Health Protection Research Unit in Emerging and Zoonotic Infections,

Liverpool, UK.

[3]Laboratory for Animal Model Pathology, Institute of Veterinary Pathology, Vetsuisse

Faculty, University of Zurich, Switzerland.

[4]Department of Pharmacology and Therapeutics, University of Liverpool, UK.

[5]Centre of Excellence in Long-acting Therapeutics (CELT), University of Liverpool,

UK.

[6]A*STAR Infectious Diseases Laboratories (A*STAR ID Labs), Agency for Science,

Technology and Research (A*STAR), Singapore.

*These authors contributed equally.

[†] Corresponding author: rebee@liverpool.ac.uk

Running title: Antiviral and SARS-CoV-2 genome diversity

Synopsis

Objectives. Immunocompromised individuals are susceptible to severe COVID-19 and potentially contribute to the emergence of variants with altered pathogenicity due to persistent infection. This study investigated the impact of immunosuppression on SARS-CoV-2 infection in k18-hACE2 mice and the effectiveness of antiviral treatments in this context during the first 7 days of infection.

Methods Mice were immunosuppressed using cyclophosphamide and infected with a B daughter lineage of SARS-CoV-2. Molnupiravir and nirmatrelvir, alone and in combination, were administered and viral load and viral sequence diversity was assessed.

Results Treatment of infected but immune compromised mice with both compounds either singly or in combination resulted in decreased viral loads and pathological changes compared to untreated animals. Treatment also abrogated infection of neuronal tissue. However, no consistent changes in the viral consensus sequence were observed, except for the emergence of the S:H655Y mutation. Molnupiravir, but not nirmatrelvir or immunosuppression alone, increased the transition/transversion (Ts/Tv) ratio, representative of G>A and C>U mutations and this increase was not altered by the co-administration of nirmatrelvir with molnupiravir.

Notably, immunosuppression itself did not appear to promote the emergence of mutational characteristic of variants of concern (VOCs).

Conclusions Further investigations are warranted to fully understand the role of immunocompromised individuals in VOC development, especially by taking persistence into consideration, and to inform optimised public health strategies. It is more likely that immunodeficiency promotes viral persistence but does not necessarily lead to substantial consensus-level changes in the absence of antiviral selection pressure. Consistent with mechanisms of action, molnupiravir showed a stronger mutagenic effect than nirmatrelvir in this model.

Keywords

SARS-CoV-2, COVID-19, immunocompromised, intra-host evolution, Molnupiravir,

Nirmatrelvir, Paxlovid.

Introduction

Unsurprisingly, since the start of the Severe Acute Respiratory Syndrome 2 (SARS-

CoV-2) pandemic and the first deposited genome sequences, and like other

coronaviruses, SARS-CoV-2 has diverged through single nucleotide polymorphism,

and homologous and heterologous recombination applications resulting in insertions

and deletions [1,2]. Over the course of the pandemic changes that have dominated

have resulted in increased transmissibility such as the P323L/D614G changes in

early 2020 [3-5], immune-evasion [6] and altered pathogenicity [7].

Founder effects, population bottlenecks, selection pressures and behaviour have

contributed to the diversification of the SARS-CoV-2 genome but also to the

apparent waves of different variants. Several Variants of Concern (VOCs) have

arisen that have a transmission advantage and/or potential immune evasion. Some

reports have suggested that such variants may have arisen in hosts with

compromised immunity and/or persistent infections, where infection leads to the

generation of more diverse variants through longer viral evolution within an individual

[8]. This includes a changing landscape of dominant viral genome sequence and

minor genomic variants in immune compromised individuals e.g. in a patient with

cancer [9]. Changes within the individual mapped to several different regions on the

SARS-CoV-2 genome including the spike glycoprotein and orf8.

Complicating the picture of potential rapid and dramatic genomic change in immune

compromised hosts is that similar changes can be observed in immune competent

patients. This can be either as part of the dominant genomic sequence [10] or minor

variant genomes [1]. Indeed, genomic variants with deletions can be identified in the

minor genomic variant population of Middle East respiratory syndrome coronavirus

(MERS-CoV) from patients [11] and as part of the dominant genomic sequence in

camels [12,13].

Parallels with other animal coronaviruses can be found where persistent infections
are established, and this might be associated in pathogenicity; an example are feline
coronavirus (FCoV) infections and feline infectious peritonitis (FIP) [14-17]. Thus, one
concern with long term persistence of SARS-CoV-2 in immune compromised
patients is that new transmissible variants could emerge [8].

Three small molecule direct acting anti-virals (DAAs) have received early use
authorisation for the treatment of COVID-19: remdesivir, molnupiravir (both
nucleoside analogues which target viral nucleic acid synthesis) and nirmatrelvir
(which targets the main viral protease). Unlike remdesivir, molnupiravir and
nirmatrelvir are orally administered and thus more readily deployed for treatment in
the community. Nirmatrelvir is packaged with ritonavir (as Paxlovid), this later
molecule acting as a pharmacokinetic boosting agent to inhibit P450 (CYP) 3A4.
However, adequate nirmatrelvir plasma concentrations can be achieved in mice
without the need for ritonavir boosting. In cell culture single or combination treatment
can result in decreased viral replication [18,19] and a natural extension is that such anti-
virals may be deployed as combination therapy to reduce the emergence of resistant
genotypes [20]. Resistant genotypes/phenotypes have been identified in vitro for
remdesivir [21]. Molnupiravir has previously been shown to enhance viral
transition/transversion mutations in a phase II clinical trial [22] and a molnupiravir
associated signature has been identified in circulating SARS-CoV-2 lineages since
the introduction of molnupiravir in 2022 [23].

Immunocompromised patients with a SARS-CoV-2 infection are treated as a priority
with anti-virals, including those compounds that generically target virus replication by
causing hyper-mutation or specifically preventing the function of a viral protein critical
to the life cycle of the virus. Such anti-virals may be deployed as combination
therapy to reduce the emergence of resistant genotypes [20] and may be particularly
relevant for patients with compromised immunity [24]. However, in the latter patients,
anti-virals may decrease viral loads but enhance genomic plasticity. To investigate
this, the genomic variation of SARS-CoV-2 was evaluated in an immune
compromised host over the first 7 days of infection, in the absence and presence of
medical countermeasures. We have developed animal models of COVID-19 to be
able to assess pathogenicity of new variants and develop interventions [25-27]. An

immune suppressed K18-hACE2 transgenic mouse model was used to simulate

patients with severe COVID-19 [28,29]. Two anti-virals, molnupiravir and nirmatrelvir,

were   evaluated   either   singly   or   in   combination.

Methods

Animal infection and treatment

A UK variant of SARS-CoV-2 (hCoV-2/human/Liverpool/REMRQ0001/2020), was

used as described previously [30,31]. Mutations belonging to the B daughter lineage

virus are outlined in table 1.

Table 1: Input virus used in this study

Input virus		
Nucleotide change	**Gene**	**Amino Acid Change**
A6948C	Nsp3	N1410T or N2228T
G11083T	Nsp6	L37F or L3606F
C21005T	Nsp16	A116V or A2513V
C25452T	Orf3a	I20 no change
C28253T	Orf8	F120 no change

Animal work was approved by the local University of Liverpool Animal Welfare and

Ethical Review Body and performed under UK Home Office Project Licence

PP4715265. Transgenic mice carrying the human ACE2 gene under the control of

the keratin 18 promoter (K18-hACE2; formally B6.Cg-Tg(K18-ACE2)2Prlmn/J) were

purchased from Jackson Laboratories (France) at 8 – 10 weeks of age. Mice were

maintained under SPF barrier conditions in individually ventilated cages and

underwent a week of acclimatisation in these conditions prior to experimental use.

Experimental design is shown in Fig. 1 and treatment groups detailed in Table 2.

Animals were randomly assigned into multiple cohorts of four animals using a

random number generator. For operational reasons at high containment the

treatment groups were not blinded during the experiment. Sample size was

determined using prior experience of similar experiments with SARS-CoV-2. For

SARS-CoV-2 infection, mice were anaesthetized lightly with isoflurane and

inoculated intra-nasally with 50 µl containing 10^4 PFU SARS-CoV-2 in PBS as

described previously [26]. Some cohorts of mice were immunosuppressed by

treatment with cyclophosphamide (100 mg/kg) intra-peritoneally (IP) at day -4 and -1

pre-infection. Molnupiravir was made up in 10% PEG400 and 2.5% cremophor in

water and used at 100 mg/kg. Nirmatrelvir was dissolved in 2% Tween 80 in 98%

(v/v) of 0.5% methyl cellulose and used at 500 mg/kg. These does were chosen

based on the known therapeutic range for these drugs in mice [32-35]. Both drugs were

administered via the oral route one hour prior to infection and then twice daily up to 4

days post-infection via the oral (PO) route. Groups of animals were kept in the same

cages during the experiment and were always weighed and treated in the same

order. Mice were sacrificed at day 6 (vehicle and cyclophosphamide treated group)

or 7 (all others) after infection by an overdose of pentobarbitone. Weights were

recorded daily, and tissues were removed immediately for downstream processing.

The right lung and nasal turbinates were frozen at -80 °C until further processing.

The left lung and heads were fixed in 10% neutral buffered formalin for 24-48 h and

then stored in 70%. No data were excluded from the analyses.

Histology, immunohistology and morphometric analysis

The fixed left lung was routinely paraffin wax embedded. Heads were sawn

longitudinally in the midline using a diamond saw (Exakt 300; Exakt) and the brain

left in the skull. Heads were gently decalcified in RDF (Biosystems) for twice 5 days,

at room temperature and on a shaker, then both halves paraffin wax embedded.

Consecutive sections (3-5 µm) were either stained with hematoxylin and eosin (HE)

or used for immunohistology (IH). IH was performed to detect viral antigen

expression using the horseradish peroxidase method and a rabbit anti-SARS-CoV

nucleocapsid protein (Rockland, 200-402-A50) as primary antibody, as previously

described [26,36,37].

For morphometric analysis, the immunostained sections were scanned

(NanoZoomer-XR C12000; Hamamatsu, Hamamatsu City, Japan) and analysed

using the software program Visiopharm (Visiopharm 2020.08.1.8403; Visiopharm,

Hoersholm, Denmark) to quantify the area of viral antigen expression in relation to

the total area (area occupied by lung parenchyma) in the sections. This was used to

compare the extent of viral antigen expression in the lungs between the different

treatment groups. A first app was applied that outlined the entire lung tissue as ROI

(total area). For this a Decision Forest method was used and the software was

trained to detect the lung tissue section (total area). Once the lung section was

outlined as ROI the lumen of large bronchi and vessels was manually excluded from

the ROI. Subsequently, a second app with Decision Forest method was trained to

detect viral antigen expression (as brown DAB precipitate) within the ROI.

RNA extraction

To inactivate virus in throat swabs, 260 µL of swab buffer was inactivated in a Class

II Biosafety cabinet using 750 µL of TRIzol LS reagent (ThermoFisher, Runcorn,

UK), and transferred into 2 mL screw-cap vials and mixed. Samples were stored at

−80 °C until further analysis. RNA samples were normalised to 20ng/µl before qPCR

and sequencing.

qRT-PCR for viral load

Viral loads were quantified using the GoTaq® Probe 1-Step RT-qPCR System

(Promega). For quantification of SARS-COV-2 the nCOV_N1 primer/probe mix from

the SARS-CoV-2 (2019-nCoV) CDC qPCR Probe Assay (IDT) were utilised and

murine 18S primers as described previously [25,26].

Sequencing of SARS-CoV-2

RNA samples were placed into plates based on very high viral load (Ct<18), high

viral load (Ct 20-24), medium viral load (Ct 25-28) or low viral load (Ct >28) to assist

with pooling strategy. Library preparation consisted of converting RNA to cDNA

using LunaScript™ (Thermofisher), then amplified by reverse complement (RC)-PCR

amplification (EasySeq™ SARS-CoV-2 Whole Genome Sequencing kit, Nimagen,

Netherlands). This kit barcodes and ligates Illumina adapters in a single PCR

reaction, with two separate pools of primers (pools 1 and 2). After amplification, each

amplicon library was pooled 1:1 before being cleaned with AmpliClean™ beads and

quantification. The two pools were then added together and denatured 2µl, 4µl, 8µl

and 16µl of each pool for very high, high, medium and low viral load was taken

respectively. Finally, the denatured amplicon library was loaded into the NovaSeq

cartridge (2 x 150 bp run).

Bioinformatics

Supplementary Fig. S1 provides an overview of the workflow used in this study. In

short, raw paired end fastq files were inputted into the EasySeq pipeline to generate

alignment files, vcf's and consensus sequences using the NC_045512.2 SARS-CoV-

2 reference[38].Consensus sequences were inputted into Nextclade for lineage

assignment and bam files were inputted into DiversiTools

(https://github.com/josephhughes/DiversiTools) to assess global minor variation.

Sequencing data was analysed as previously described and statistical analysis and

visualisation was performed in R [22]. In brief, the entropy outputs from DiversiTools

was imported to R and a coverage of at least 100 at each position was required, the

average quality scores, derived from phred scoring, for each position was less than

$2x10^{-6}$ indicative of low basecalling error. Raw fastq files are available under SRA

Project Accession: PRJNA886870. Code for analysis and figure generation is

available at https://github.com/Hiscox-lab/viral-genomics-immunosupression-and-

countermeasures.

Statistics

Graphs were prepared and statistics performed using Prism 10 (Graphpad Inc). *P*

values were set at 95% confidence interval. A repeated-measures two-way ANOVA

(Bonferroni post-test) was used for time-courses of weight loss; log-rank (Mantel-

Cox) test was used for survival curve and Mann-Whitney *U* test for side-by-side

comparisons. All differences not specifically stated to be significant were not

significant ($p > 0.05$). For all figures, *$p < 0.05$.

Results and Discussion

Since the emergence of the Alpha VOC there has been discussion on the involvement of the immunocompromised host and the generation of variants [8,39-43]. There are many case studies in the literature that follow SARS-CoV-2 evolution in immunocompromised hosts, however, the generation of VOCs is likely due to persistent infection as opposed to immunocompromised immune systems itself, little has been explored experimentally. In this study, mice were chemically immunocompromised with cyclophosphamide which is known to efficiently remove adaptive immunity in the form of B and T cells [44]. Additionally, therapeutic agents, molnupiravir and nirmatrelvir, were used independently and in combination to determine the effectiveness of these compounds in an immunocompromised model, and the impact of these compounds on viral sequence diversity during the first 7 days of infection.

Modelling an immunocompromised state in animal models in the context of SARS-CoV-2 is important for the consideration of countermeasures that may be utilised for humans who are considered vulnerable. Cyclophosphamide has been used previously to study the impact of immunosuppression in a hamster model [45-47], where intranasally infected hamsters with cyclophosphamide treatment before infection had prolonged weight loss and an inadequate neutralising antibody response to SARS-CoV-2. Distinct transcriptional profiles were identified between immunocompetent and immunosuppressed animals; however, the impact of antivirals or viral genome diversity was not investigated.

To investigate the frequency of genomic changes that occur in SARS-CoV-2 in the immune compromised or competent host in the presence or absence of antiviral drugs, K18-hACE2 transgenic mice were used as a model for severe SARS-CoV-2 infection in humans [48]. We have found that the pathological changes in the lungs in this model in many aspects resemble those in humans who have died of severe COVID-19 [26,28,29,36,37]. To mimic a host with compromised immunity, an experimental protocol was developed in which mice were exposed to cyclophosphamide [44] (Fig. 1, Table 2). Several anti-viral regimes in humans were simulated in the mouse model by giving a human equivalent dose of either molnupiravir (100 mg/kg), nirmatrelvir

273 (500 mg/kg) or both in combination. This included prophylactic followed by
274 therapeutic treatment. Mice were infected with 10^4 PFU of SARS-CoV-2.

Figure 1. *Schematic diagram of the experimental design for infection of immune compromised K18-hACE2 mice with SARS-CoV-2 and evaluation of two antiviral drugs given at a human equivalent dose; molnupiravir, a broad acting compound causing error catastrophe, or nirmatrelvir which specifically targets the viral 3C-like protease. Cyclophosphamide was used at 100 mg/kg via the intraperitoneal route to immunosuppress mice. Molnupiravir was used at 100 mg/kg and nirmatrelvir at 500 mg/kg both via the oral route. Effects of infection and treatment were evaluated by measuring the weight of the mice daily, determining viral loads in sequential oral/throat swabs and at day 7 post-infection, and examining nose, brain and lung at day 7 post infection for any histological changes and the expression of SARS-CoV-2 nucleoprotein.*

276 **Table 2. Treatment groups for in vivo analysis**

Group	Treatment
1	Control (vehicle)
2	Cyclophosphamide
3	Molnupiravir
4	Cyclophosphamide + molnupiravir
5	Cyclophosphamide + nirmatrelvir
6	Cyclophosphamide + molnupiravir + nirmatrelvir

Treatment with Molnupiravir or Nirmatrelvir either individually or in combination provides recovery in immune compromised mice infected with SARS-CoV-2.

Cyclophosphamide treatment prior to SARS-CoV-2 infection of hACE2 mice led to a more pronounced early weight loss in comparison to immunocompetent mice, a phenomenon previously reported in hamsters [47]. This was not associated with earlier mortality than in vehicle treated immunocompetent mice, although in human, a delayed adaptive immune response has been shown to be associated with fatality in COVID-19 patients, which may have been observed over longer timeframes [49]. Daily weighing of the animals indicated that all groups lost body weight after day 1 (Fig. 2). We attribute this to aversion to eating as all therapies were applied by gavage. However, starting at day 3 all groups, except for mice exposed to cyclophosphamide, or mice exposed to cyclophosphamide and treated with molnupiravir, started to gain, or stabilise weight. By days 5 and 6 a clear pattern had emerged where all groups treated with molnupiravir or nirmatrelvir either individually or in combination had regained their starting weight. The exception to this were mice exposed to vehicle only (controls) or cyclophosphamide; these reached a humane end point on day 6 (Fig. 2). Comparison of survival curves again indicated that immune compromised animals treated either singly or in combination with each therapeutic went on to survive (Fig. 3).

Figure 2: Treatment of SARS-CoV-2-infected mice leads to decreased weight loss. K18-hACE2 mice were challenged intranasally with 10^4 PFU SARS-CoV-2 Mice were monitored for weight at indicated time-points. (n = 4). Data represent the mean residual weight ± SEM. Comparisons were made using a repeated-measures two-way ANOVA (Bonferroni post-test). * on the represents $P < 0.05$. Asterisks below the curves represent * $P < 0.05$ and ** $P < 0.01$ between the cylophophamide and vehicle groups. Brackets and asterisk at the side represents $P < 0.05$ for the Vehicle/cycophosphamide groups and the drug treated groups.

Figure 3: Treatment of SARS-CoV-2-infected mice leads to enhanced survival.
K18-hACE2 mice were challenged intranasally with 10^4 PFU SARS-CoV-2. Survival
was assessed at indicated time points and significance determined using log rank
(Mantel-Cox) test (n = 4).

Viral load decreases in immune compromised mice treated with Molnupiravir

or Nirmatrelvir either individually or in combination.

Viral load in terms of copy numbers of the SARS-CoV-2 genome were calculated for

throat swabs during infection and compared to nasal tissue and lung tissue at the

end of the experiment. The data indicated that for throat swabs on days 1 and 3

post-infection there was a significant decrease in viral load in animals treated with

molnupiravir or nirmatrelvir either individually or in combination compared to

untreated controls (Figure 4A). At day 3 there was a significant difference between

both compounds used in combination and molnupiravir only (Figure 4A). No

significant differences were observed between vehicle control and

cyclophosphamide only groups.

Comparison of viral loads and titres in nasal and lung tissue respectively (Figure 4B

and 4C, respectively) at day 7 post-infection reflected that there was a significantly

lower viral load in animals treated with molnupiravir or nirmatrelvir either individually

or in combination compared to untreated mice. However, nirmatrelvir treatment

resulted in a greater decrease in viral load compared to molnupiravir. The

molnupiravir/nirmatrelvir combination was also more effective at decreasing viral

load than either drug alone, but this was only statistically significant in the case of

molnupiravir vs the drug combination.

Figure 4. Viral loads in swabs and tissues. *K18-hACE2 mice were challenged intranasally with 10^4 PFU SARS-CoV-2 and treated as indicated (n = 4 per group). RNA extracted from oral/throat swabs and nasal tissue was analysed for virus RNA load using qRT-PCR and primers specific for the SARS-CoV-2 N gene. Assays were normalised relative to levels of 18S RNA. Lung tissue was analysed for live virus by plaque assay. Data for individual animals are shown with the median value represented by a black line. (A) Throat swabs; (B) nasal tissue; (C) lung tissue. Comparisons were made using two-way ANOVA (Bonferroni post-test) in panel A and Mann-Whitney U test (Panels B and C). * Represents p < 0.05.*

Treatment with molnupiravir or nirmatrelvir or both in combination results in marked reduction of pulmonary infection and inhibits viral spread to the brain.

The lung, nose and brain of all animals were examined for any histopathological changes and the expression of viral antigen by immunohistology, to determine whether treatment of the animals with molnupiravir and/or nirmatrelvir influenced the outcome of infection. The lungs of vehicle treated, immunocompetent animals showed the typical changes previously reported in K18-hACE2 mice infected with this virus strain [26], i.e. multifocal areas with pneumocyte degeneration, type II pneumocyte activation, mild neutrophil infiltration, and mild vasculitis, with a diffuse increase in interstitial cellularity and widespread SARS-CoV-2 antigen expression in alveolar epithelial cells (Fig. 5A). In mice that had received cyclophosphamide alone, the changes were very similar, but slightly less widespread, with some unaltered parenchyma and less extensive viral antigen expression (Fig. 5B). With molnupiravir treatment, both inflammatory processes and viral antigen expression were markedly decreased; indeed, SARS-CoV-2 antigen was only found in disseminated patches of alveoli with positive pneumocytes (Fig. 5C). With cyclophosphamide and molnupiravir treatment, the lung parenchyma was widely unaltered, and there were only small patches of inflammation and alveoli with viral antigen expression, respectively (Fig. 5D). These were further reduced in number and size in animals that had received cyclophosphamide and nirmatrelvir (Fig. 5E). Treatment with all three compounds, cyclophosphamide, molnupiravir and nirmatrelvir, resulted in widely unaltered lung parenchyma with no or minimal viral antigen expression (Fig. 5F).

Figure 5: *K18-hACE2 mice were challenged intranasally with 10^4 PFU SARS-CoV-2 and treated as indicated below (n = 4 per group). Immunohistology for the detection of viral antigen in the lung at day 6 or 7 post infection. Sections from the formalin-fixed, paraffin embedded left lung lobe were stained using anti-SARS-CoV nucleoprotein and counterstained with hematoxylin. Representative images from the individual treatment groups are shown as follows: A. vehicle; B. cyclophosphamide; C. molnupiravir; D. cyclophosphamide and molnupiravir; E. cyclophosphamide and nirmatrelvir; F. cyclophosphamide, molnupiravir and nirmatrelvir. Viral antigen expression is restricted to pneumocytes in a few individual alveoli (higher magnifications in insets). Bars represent 2.5 mm (A-E), 1 mm (F) and 20 µm (F, insets).*

Examination of the heads using longitudinal sections (midline) revealed consistent
and widespread infection of the brain in animals treated with the vehicle or with
cyclophosphamide alone (Fig. 6A, B); this was associated with mild perivascular
mononuclear infiltration in particular in the brain stem, as described before in K18-
hACE2 mice infected with this virus strain [37]. In both groups of animals,
immunohistology confirmed viral antigen expression in the respiratory and/or
olfactory epithelium, in the latter with evidence of infection in olfactory sensory
neurons (Fig. 6A, B). In the other groups, there was no evidence of viral infection of
the brain (Fig. 6C-F), and viral antigen expression in the nasal mucosa was not seen
or restricted to scattered individual epithelial cells. In vehicle control and
cyclophosphamide mice, the nasal mucosa harboured viral antigen at this stage, in
the respiratory epithelium and in the olfactory epithelium; in the latter it also
appeared to be present in sensory neurons. Consequently, the virus had reached
and spread widely in the brain where it was detected in neurons; the infection was
associated with mild inflammatory response in particular in the brain stem, as
described before in K18-hACE2 mice infected with this virus strain [26,37]. After
treatment with all three compounds, cyclophosphamide, molnupiravir and
nirmatrelvir, the lung parenchyma was basically unaltered, with no or minimal viral
antigen expression. In all groups of mice, viral antigen expression in the nasal
mucosa was not seen or restricted to scattered individual epithelial cells and there
was no evidence of viral infection of the brain, suggesting that the antiviral treatment
blocked infection of the brain. Whether the latter is purely a consequence of reduced
viral replication in the upper respiratory tract cannot be assessed in the present
study; it does, however, appear likely.

Figure 6*: K18-hACE2 mice were challenged intranasally with 10^4 PFU SARS-CoV-2 and*
treated as indicated below (n = 4 per group). Immunohistology for the detection of viral
antigen in the brain and nose at day 6 or 7 post infection. Sections from formalin-fixed,
decalcified and paraffin embedded heads after longitudinal sawing in the midline were
stained using anti-SARS-CoV nucleoprotein, and counterstained with hematoxylin. Only
small fragments of nasal mucosa were available for the examination, as the nasal turbinates
had been sampled for PCR. Representative images from the individual treatment groups are
shown as follows: ***A****. Vehicle. There is widespread infection of the brain. The insets show*
infection of individual cells with the morphology of olfactory sensory neurons and epithelial
cells in the olfactory epithelial layer (left inset) and individual respiratory epithelial cells in the
nasal mucosa (arrowhead; right inset); ***B****. Cyclophosphamide. There is widespread infection*
of the brain. The inset shows a group of positive epithelial cells/sensory neurons in the
olfactory epithelial layer (arrowhead); ***C****. Molnupiravir. There is no evidence of brain*
infection. ***D****. Cyclophosphamide and molnupiravir. There is no evidence of brain infection.* ***E.***
Cyclophosphamide and nirmatrelvir. There is no evidence of brain infection. ***F.***
Cyclophosphamide, molnupiravir and nirmatrelvir. There is no evidence of brain infection.
Bars represent 2.5 mm (A-F) and 20 µm (A, B insets). FC – frontal cortex, HC –
hippocampus, MO – medulla oblongata, OB – olfactory bulb, OSN - olfactory sensory
neurons.

Evaluation of dominant and minor variants in SARS-CoV-2

To determine the impact of immunosuppression on viral diversity, 116 RNA samples from swabs and tissue were sequenced and analysed using the EasySeq WGS protocol by Nimagen. alignment files and associated index files were inputted into DiversiTools to provide mutation data and outputs were analysed in R. Samples with less than 90% breadth of coverage were discarded for mutational analysis (n=12), as well as samples that returned bad or mediocre quality scores in nextclade (n=13). The samples that were excluded were associated with higher Ct values and later time points belonging in the nirmatrelvir treatment groups. Sequencing data from 89 samples were taken forward in the analysis (swab, n=50, tissue n=39, Supplementary Table S1).

The input virus contained 5 substitutions and 3 amino acid substitutions in comparison to the reference sequence (NC_045512.2) and were thus not considered as changes during the analysis (Table 1). The S: H655Y mutation was present in 76% of the genomes that passed QC at the dominant level and observed as a minor variant across all samples (Supplementary Fig. S2). This mutation has been reported previously as a spike adaptation to other species such as cats, hamsters, and mink [50-52] and of course has independently arisen in human lineages such as Omicron [53]. As this mutation was clearly associated with a species adaptation, it was disregarded for the evaluation of treatment and immune status driven mutations. The other mutations appear to be novel at the time of writing; however, no distinct group was associated with driving these mutations, and can be overall interpreted as a rare event (Supplementary Table S2). The sequences showing the highest number of mutations were sequences derived from tissue samples. The species-specific adaptation S:H655Y was observed more frequently in the dataset, where there was little evidence of adaptations specific to immunocompromised and antiviral environments, putting the evolutionary pressures into perspective. Adaptations associated with immunosuppression and antiviral treatment may emerge in the context of persistent replication of the virus which would need to be investigated.

Molnupiravir increases the Ts/Tv ratio at the minor variant level in genomes derived from swabs

To further assess the impact of immunocompromising mice by cyclophosphamide,

and the therapeutic agents molnupiravir and the nirmatrelvir, a minor variant analysis

was conducted on samples derived from throat swabs as performed previously[22].

Only samples with a >90X coverage with a 100X depth were taken forward into this

analysis and the average depth was over 1400 for each sample (Fig S3). No

relationship was observed between ct value and the calculated average

transition/transversion (Ts/Tv) (Fig S4). The average Ts/Tv ratio for SARS-CoV-2

genomes from each mouse and the mean of each group was compared across

cohorts at day 1, day 3 and day 5 of infection in line with analysis performed

previously in a phase II clinical trial [22]. On day 1, an increase in Ts/Tv ration was

observed in the molnupiravir cohort and the cyclophosphamide and molnupiravir

cohort and had a p value < 0.05 when compared to the vehicle control and

cyclophosphamide only groups (Fig. 7). The number of samples analysed for

cyclophosphamide and nirmatrelvir only was too small for statistical analysis,

however, the trend resembles that of vehicle and cyclophosphamide only with little

change in the Ts/Tv ratio. Likewise, the combined cyclophosphamide and

molnupiravir and nirmatrelvir cohort is represented by one genome derived from one

mouse, due to low sequencing coverage obtained from the other mice within this

group, however, the trend resembles that of other genomes with exposure to

molnupiravir with an increased Ts/Tv ratio. The same is observed at day 3 of

sampling, however, there is a significant difference between the mean Ts/Tv ratio

between the molnupiravir only and cyclophosphamide and molnupiravir groups.

Importantly, the Ts/Tv ratios between the vehicle control and cyclophosphamide only

groups resemble each other demonstrating that immunosuppression itself does not

drive diversification of the viral genome over this time course. Curiously, when

looking at base changes independently of the Ts/Tv ratios, there are significant

changes between C to G, C to A and A to U in cyclophosphamide groups on day 1

and day 3 (Fig S3). This could be evidence of RNA editing through ROS.

Cyclophosphamide has been shown to activate oxidative stress pathways previously

and may be a consequence of this treatment[54,55]. This highlights a research gap in

events that can influence RNA editing in RNA viruses.The proportion of base

changes in the C to U and G to A transitions were significantly different in the

molnupiravir only group as previously seen in a phase II clinical trial [22,23] (Figure 8).

Further investigations are warranted to understand completely the role of
immunocompromised individuals in the development of SARS-CoV-2 variants. It is
more likely that immunodeficiency promotes viral persistence providing the virus
more opportunity to replicate and introduce mutations. Molnupiravir, compared to
nirmatrelvir, shows a stronger mutagenic effect in this model at the minor variant
level, however, data is insufficient to make conclusions regards consensus level
changes over the timeframes used in this study. When these therapies are used
individually or in combination, there is successful depletion in viral load and animals
recover from infection, whilst preventing infiltration into brain tissue. Given the
concern of molnupiravir associated lineages in circulation [23], combination therapy
may reduce this through more effective clearance of the virus [20], although this would
need to be evaluated over time in a real-world setting as the mutational signatures
were observed in the combined therapy group. The AGILE clinical trial is currently
ongoing to answer this question (ISRCTN: ISRCTN27106947). It is important to
note, that the mutational spectrum reported in this study is obtained by amplicon
sequencing data, and there is potential for RT-PCR errors within the data.

498
499
500 **Figure 7:** *The mean Ts/Tv ratio per genome plotted as boxplots. The plot is facetted by day*
501 *post infection. Less genomes were recovered for cyclophosphamide and* nirmatrelvir *and*
502 *cyclophosphamide,* nirmatrelvir *and molnupiravir, therefore statistical analysis returns the*
503 *differences as non-significant. Trends can be concluded with caution. * Represents a P*
504 *value <0.05 (Mann Whitney U test).*
505
506

507
508

Day of swab sample

Acknowledgements

The authors are grateful to the technical staff at the Histology Laboratory, Institute of

Veterinary Pathology, Vetsuisse Faculty, University of Zurich, for excellent technical

support.   JAH   is   a   member   of   the   ISARIC4C   consortium

(https://isaric4c.net/about/authors/), and we thank them for the use of the SARS-

CoV-2 isolate used in this study.

Funding

This work was funded by the MRC (MR/W005611/1) 'G2P-UK: A national virology

consortium to address phenotypic consequences of SARS-CoV-2 genomic variation'

and MR/Y004205/1 'The G2P2 virology consortium: keeping pace with SARS-CoV-2

variants, providing evidence to vaccine policy, and building agility for the next

pandemic' (co-Is JPS and JAH) and funded in part by U.S. Food and Drug

Administration Medical Countermeasures Initiative contract (75F40120C00085) to

JAH. The article reflects the views of the authors and does not represent the views

or policies of the FDA. Additionally, DNDi under the support by the Wellcome Trust

(Grant ref: 222489/Z/21/Z to AO and JPS) through the 'COVID-19 Therapeutics

Accelerator'. A.O. acknowledges funding by Wellcome Trust (222489/Z/21/Z),

EPSRC (EP/R024804/1; EP/S012265/1) and National Institute of Health (NIH)

(R01AI134091; R24AI118397). J.P.S. also acknowledges funding from the Medical

Research Council (MRC) (MR/R010145/1, MR/W021641/1, PA6162_G2P2-2023),

BBSRC (BB/R00904X/1; BB/R018863/1; BB/N022505/1) and Innovate UK

(TS/W022648/1). A.K. received support from the Swiss National Science

Foundation (SNSF; IZSEZ0 213289).

Transparency Declaration

A.O. is a director of Tandem Nano Ltd and co-inventor of patents relating to drug

delivery. A.O. has been co-investigator on funding received by the University of

Liverpool from ViiV Healthcare and Gilead Sciences in the past 3 years unrelated to

COVID-19. A.O. has received personal fees from Gilead and Assembly Biosciences

in the past 3 years, also unrelated to COVID-19. JPS has received funding from ENA

respiratory Pty Ltd, Bicycle Tx Ltd, and Infex Therapeutics Ltd unrelated to this

study. R.P.R. is an employee at TopMD Precision Medicine Ltd. No other conflicts

are declared by the authors.

References

1   Moore, S. C., Penrice-Randal, R., Alruwaili, M., Randle, N., Armstrong, S., Hartley, C., Haldenby, S., Dong, X., Alrezaihi, A., Almsaud, M., Bentley, E., Clark, J., Garcia-Dorival, I., Gilmore, P., Han, X., Jones, B., Luu, L., Sharma, P., Shawli, G., Sun, Y., Zhao, Q., Pullan, S. T., Carter, D. P., Bewley, K., Dunning, J., Zhou, E. M., Solomon, T., Beadsworth, M., Cruise, J., Crook, D. W., Matthews, D. A., Davidson, A. D., Mahmood, Z., Aljabr, W., Druce, J., Vipond, R., Ng, L., Renia, L., Openshaw, P. J. M., Baillie, J. K., Carroll, M. W., Stewart, J., Darby, A., Semple, M., Turtle, L. & Hiscox, J. A. Amplicon-Based Detection and Sequencing of SARS-CoV-2 in Nasopharyngeal Swabs from Patients With COVID-19 and Identification of Deletions in the Viral Genome That Encode Proteins Involved in Interferon Antagonism. *Viruses* **12** (2020). https://doi.org/10.3390/v12101164

2   McCarthy, K. R., Rennick, L. J., Nambulli, S., Robinson-McCarthy, L. R., Bain, W. G., Haidar, G. & Duprex, W. P. Natural deletions in the SARS-CoV-2 spike glycoprotein drive antibody escape. *bioRxiv*, 2020.2011.2019.389916 (2020). https://doi.org/10.1101/2020.11.19.389916

3   Korber, B., Fischer, W. M., Gnanakaran, S., Yoon, H., Theiler, J., Abfalterer, W., Hengartner, N., Giorgi, E. E., Bhattacharya, T., Foley, B., Hastie, K. M., Parker, M. D., Partridge, D. G., Evans, C. M., Freeman, T. M., de Silva, T. I., Sheffield, C.-G. G., McDanal, C., Perez, L. G., Tang, H., Moon-Walker, A., Whelan, S. P., LaBranche, C. C., Saphire, E. O. & Montefiori, D. C. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell* **182**, 812-827 e819 (2020). https://doi.org/10.1016/j.cell.2020.06.043

4   Alruwaili, M., Armstrong, S., Prince, T., Erdmann, M., Matthews, D. A., Luu, L., Davidson, A., Aljabr, W. & Hiscox, J. A. SARS-CoV-2 NSP12 associates with TRiC and the P323L substitution acts as a host adaption. *Journal of virology* **97**, e0042423 (2023). https://doi.org/10.1128/jvi.00424-23

5   Goldswain, H., Dong, X., Penrice-Randal, R., Alruwaili, M., Shawli, G. T., Prince, T., Williamson, M. K., Raghwani, J., Randle, N., Jones, B., Donovan-Banfield, I. a., Salguero, F. J., Tree, J. A., Hall, Y., Hartley, C., Erdmann, M., Bazire, J., Jearanaiwitayakul, T., Semple, M. G., Openshaw, P. J. M., Baillie, J. K., Baillie, J. K., Semple, M. G., Openshaw, P. J. M., Carson, G., Alex, B., Andrikopoulos, P., Bach, B., Barclay, W. S., Bogaert, D., Chand, M., Chechi, K., Cooke, G. S., da Silva Filipe, A., de Silva, T., Docherty, A. B., dos Santos Correia, G., Dumas, M.-E., Dunning, J., Fletcher, T., Green, C. A., Greenhalf, W., Griffin, J. L., Gupta, R. K., Harrison, E. M., Hiscox, J. A., Ho, A. Y. W., Horby, P. W., Ijaz, S., Khoo, S., Klenerman, P., Law, A., Lewis, M. R., Liggi, S., Lim, W. S., Maslen, L., Mentzer, A. J., Merson, L., Meynert, A. M., Moore, S. C., Noursadeghi, M., Olanipekun, M., Osagie, A., Palmarini, M., Palmieri, C., Paxton, W. A., Pollakis, G., Price, N., Rambaut, A., Robertson, D. L., Russell, C. D., Sancho-Shimizu, V., Sands, C. J., Scott, J. T., Sigfrid, L., Solomon, T., Sriskandan, S., Stuart, D., Summers, C., Swann, O. V., Takats, Z., Takis, P., Tedder, R. S., Thompson, A. A. R., Thomson, E. C., Thwaites, R. S., Turtle, L. C. W., Zambon, M., Hardwick, H., Donohue, C., Griffiths, F., Oosthuyzen, W., Donegan, C., Spencer, R. G., Norman, L., Pius, R., Drake, T. M., Fairfield, C. J., Knight, S. R., McLean, K. A., Murphy, D., Shaw, C. A., Dalton, J., Girvan, M., Saviciute, E., Roberts, S., Harrison, J., Marsh, L., Connor, M., Halpin, S., Jackson, C., Gamble, C., Plotkin, D., Lee, J., Leeming, G., Law, A., Wham, M., Clohisey, S., Hendry, R., Scott-Brown, J., Shaw, V., McDonald, S. E., Keating, S.,

Ahmed, K. A., Armstrong, J. A., Ashworth, M., Asiimwe, I. G., Bakshi, S., Barlow, S. L.,
Booth, L., Brennan, B., Bullock, K., Catterall, B. W. A., Clark, J. J., Clarke, E. A., Cole, S.,
Cooper, L., Cox, H., Davis, C., Dincarslan, O., Dunn, C., Dyer, P., Elliott, A., Evans, A.,
Finch, L., Fisher, L. W. S., Foster, T., Garcia-Dorival, I., Gunning, P., Jensen, R. L.,
Jones, C. B., Jones, T. R., Khandaker, S., King, K., Kiy, R. T., Koukorava, C., Lake, A.,
Lant, S., Latawiec, D., Lavelle-Langham, L., Lefteri, D., Lett, L., Livoti, L. A., Mancini,
M., McDonald, S., McEvoy, L., McLauchlan, J., Metelmann, S., Miah, N. S., Middleton,
J., Mitchell, J., Moore, S. C., Murphy, E. G., Pilgrim, J., Reynolds, W., Ridley, P. M.,
Sales, D., Shaw, V. E., Shears, R. K., Small, B., Subramaniam, K. S., Szemiel, A.,
Taggart, A., Tanianis-Hughes, J., Thomas, J., Trochu, E., van Tonder, L., Wilcock, E.,
Zhang, J. E., Flaherty, L., Maziere, N., Cass, E., Carracedo, A. D., Carlucci, N., Holmes,
A., Massey, H., Murphy, L., McCafferty, S., Clark, R., Fawkes, A., Morrice, K., Maclean,
A., Wrobel, N., Donnelly, L., Coutts, A., Hafezi, K., MacGillivray, L., Gilchrist, T.,
Adeniji, K., Agranoff, D., Agwuh, K., Ail, D., Aldera, E. L., Alegria, A., Allen, S., Angus,
B., Ashish, A., Atkinson, D., Bari, S., Barlow, G., Barnass, S., Barrett, N., Bassford, C.,
Basude, S., Baxter, D., Beadsworth, M., Bernatoniene, J., Berridge, J., Berry, C., Best,
N., Bothma, P., Chadwick, D., Brittain-Long, R., Bulteel, N., Burden, T., Burtenshaw,
A., Caruth, V., Chadwick, D., Chambler, D., Chee, N., Child, J., Chukkambotla, S., Clark,
T., Collini, P., Cosgrove, C., Cupitt, J., Cutino-Moguel, M.-T., Dark, P., Dawson, C.,
Dervisevic, S., Donnison, P., Douthwaite, S., Drummond, A., DuRand, I., Dushianthan,
A., Dyer, T., Evans, C., Eziefula, C., Fegan, C., Finn, A., Fullerton, D., Garg, S., Garg, S.,
Garg, A., Gkrania-Klotsas, E., Godden, J., Goldsmith, A., Graham, C., Hardy, E.,
Hartshorn, S., Harvey, D., Havalda, P., Hawcutt, D. B., Hobrok, M., Hodgson, L.,
Hormis, A., Jacobs, M., Jain, S., Jennings, P., Kaliappan, A., Kasipandian, V., Kegg, S.,
Kelsey, M., Kendall, J., Kerrison, C., Kerslake, I., Koch, O., Koduri, G., Koshy, G., Laha,
S., Laird, S., Larkin, S., Leiner, T., Lillie, P., Limb, J., Linnett, V., Little, J., Lyttle, M.,
MacMahon, M. & Investigators, I. C. The P323L substitution in the SARS-CoV-2
polymerase (NSP12) confers a selective advantage during infection. *Genome Biology*
24, 47 (2023). https://doi.org/10.1186/s13059-023-02881-5

6   Wang, B., Goh, Y. S., Prince, T., Ngoh, E. Z. X., Salleh, S. N. M., Hor, P. X., Loh, C. Y.,
Fong, S. W., Hartley, C., Tan, S. Y., Young, B. E., Leo, Y. S., Lye, D. C., Maurer-Stroh, S.,
Ng, L. F. P., Hiscox, J. A., Renia, L. & Wang, C. I. Resistance of SARS-CoV-2 variants to
neutralization by convalescent plasma from early COVID-19 outbreak in Singapore.
NPJ Vaccines **6**, 125 (2021). https://doi.org/10.1038/s41541-021-00389-2

7   Suzuki, R., Yamasoba, D., Kimura, I., Wang, L., Kishimoto, M., Ito, J., Morioka, Y., Nao,
N., Nasser, H., Uriu, K., Kosugi, Y., Tsuda, M., Orba, Y., Sasaki, M., Shimizu, R.,
Kawabata, R., Yoshimatsu, K., Asakura, H., Nagashima, M., Sadamasu, K., Yoshimura,
K., Genotype to Phenotype Japan, C., Sawa, H., Ikeda, T., Irie, T., Matsuno, K.,
Tanaka, S., Fukuhara, T. & Sato, K. Attenuated fusogenicity and pathogenicity of
SARS-CoV-2 Omicron variant. *Nature* **603**, 700-705 (2022).
https://doi.org/10.1038/s41586-022-04462-1

8   Corey, L., Beyrer, C., Cohen, M. S., Michael, N. L., Bedford, T. & Rolland, M. SARS-
CoV-2 Variants in Patients with Immunosuppression. *N Engl J Med* **385**, 562-566
(2021). https://doi.org/10.1056/NEJMsb2104756

9   Avanzato, V. A., Matson, M. J., Seifert, S. N., Pryce, R., Williamson, B. N., Anzick, S. L.,
Barbian, K., Judson, S. D., Fischer, E. R., Martens, C., Bowden, T. A., de Wit, E., Riedo,
F. X. & Munster, V. J. Case Study: Prolonged Infectious SARS-CoV-2 Shedding from an

637  Asymptomatic Immunocompromised Individual with Cancer. *Cell* **183**, 1901-1912
638  e1909 (2020). https://doi.org/10.1016/j.cell.2020.10.049

639 10 Young, B. E., Fong, S. W., Chan, Y. H., Mak, T. M., Ang, L. W., Anderson, D. E., Lee, C.
640  Y., Amrun, S. N., Lee, B., Goh, Y. S., Su, Y. C. F., Wei, W. E., Kalimuddin, S., Chai, L. Y.
641  A., Pada, S., Tan, S. Y., Sun, L., Parthasarathy, P., Chen, Y. Y. C., Barkham, T., Lin, R. T.
642  P., Maurer-Stroh, S., Leo, Y. S., Wang, L. F., Renia, L., Lee, V. J., Smith, G. J. D., Lye, D.
643  C. & Ng, L. F. P. Effects of a major deletion in the SARS-CoV-2 genome on the severity
644  of infection and the inflammatory response: an observational cohort study. *Lancet*
645  **396**, 603-611 (2020). https://doi.org/10.1016/S0140-6736(20)31757-8

646 11 Aljabr, W., Alruwaili, M., Penrice-Randal, R., Alrezaihi, A., Harrison, A. J., Ryan, Y.,
647  Bentley, E., Jones, B., Alhatlani, B. Y., AlShahrani, D., Mahmood, Z., Rickett, N. Y.,
648  Alosaimi, B., Naeem, A., Alamri, S., Alsran, H., Hamed, M. E., Dong, X., Assiri, A. M.,
649  Alrasheed, A. R., Hamza, M., Carroll, M. W., Gemmell, M., Darby, A., Donovan-
650  Banfield, I., Stewart, J. P., Matthews, D. A., Davidson, A. D. & Hiscox, J. A. Amplicon
651  and Metagenomic Analysis of Middle East Respiratory Syndrome (MERS) Coronavirus
652  and the Microbiome in Patients with Severe MERS. *mSphere* **6**, e0021921 (2021).
653  https://doi.org/10.1128/mSphere.00219-21

654 12 Chu, D. K. W., Hui, K. P. Y., Perera, R., Miguel, E., Niemeyer, D., Zhao, J.,
655  Channappanavar, R., Dudas, G., Oladipo, J. O., Traore, A., Fassi-Fihri, O., Ali, A.,
656  Demissie, G. F., Muth, D., Chan, M. C. W., Nicholls, J. M., Meyerholz, D. K., Kuranga,
657  S. A., Mamo, G., Zhou, Z., So, R. T. Y., Hemida, M. G., Webby, R. J., Roger, F.,
658  Rambaut, A., Poon, L. L. M., Perlman, S., Drosten, C., Chevalier, V. & Peiris, M. MERS
659  coronaviruses from camels in Africa exhibit region-dependent genetic diversity. *Proc*
660  *Natl Acad Sci U S A* **115**, 3144-3149 (2018).
661  https://doi.org/10.1073/pnas.1718769115

662 13 El-Kafrawy, S. A., Corman, V. M., Tolah, A. M., Al Masaudi, S. B., Hassan, A. M.,
663  Muller, M. A., Bleicker, T., Harakeh, S. M., Alzahrani, A. A., Alsaaidi, G. A., Alagili, A.
664  N., Hashem, A. M., Zumla, A., Drosten, C. & Azhar, E. I. Enzootic patterns of Middle
665  East respiratory syndrome coronavirus in imported African and local Arabian
666  dromedary camels: a prospective genomic study. *Lancet Planet Health* **3**, e521-e528
667  (2019). https://doi.org/10.1016/S2542-5196(19)30243-8

668 14 Rottier, P. J., Nakamura, K., Schellen, P., Volders, H. & Haijema, B. J. Acquisition of
669  macrophage tropism during the pathogenesis of feline infectious peritonitis is
670  determined by mutations in the feline coronavirus spike protein. *Journal of virology*
671  **79**, 14122-14130 (2005). https://doi.org/10.1128/jvi.79.22.14122-14130.2005

672 15 Licitra, B. N., Millet, J. K., Regan, A. D., Hamilton, B. S., Rinaldi, V. D., Duhamel, G. E.
673  & Whittaker, G. R. Mutation in spike protein cleavage site and pathogenesis of feline
674  coronavirus. *Emerg Infect Dis* **19**, 1066-1073 (2013).
675  https://doi.org/10.3201/eid1907.121094

676 16 Vennema, H., Poland, A., Foley, J. & Pedersen, N. C. Feline infectious peritonitis
677  viruses arise by mutation from endemic feline enteric coronaviruses. *Virology* **243**,
678  150-157 (1998). https://doi.org/10.1006/viro.1998.9045

679 17 Kipar, A., Meli, M. L., Baptiste, K. E., Bowker, L. J. & Lutz, H. Sites of feline
680  coronavirus persistence in healthy cats. *The Journal of general virology* **91**, 1698-
681  1707 (2010). https://doi.org/10.1099/vir.0.020214-0

682 18 Li, P., Wang, Y., Lavrijsen, M., Lamers, M. M., de Vries, A. C., Rottier, R. J., Bruno, M.
683  J., Peppelenbosch, M. P., Haagmans, B. L. & Pan, Q. SARS-CoV-2 Omicron variant is

highly sensitive to molnupiravir, nirmatrelvir, and the combination. *Cell Res* **32**, 322-
324 (2022). https://doi.org/10.1038/s41422-022-00618-w

19 Gidari, A., Sabbatini, S., Schiaroli, E., Bastianelli, S., Pierucci, S., Busti, C., Comez, L.,
Libera, V., Macchiarulo, A., Paciaroni, A., Vicenti, I., Zazzi, M. & Francisci, D. The
Combination of Molnupiravir with Nirmatrelvir or GC376 Has a Synergic Role in the
Inhibition of SARS-CoV-2 Replication In Vitro. *Microorganisms* **10** (2022).
https://doi.org/10.3390/microorganisms10071475

20 Hiscox, J. A., Khoo, S. H., Stewart, J. P. & Owen, A. Shutting the gate before the horse
has bolted: is it time for a conversation about SARS-CoV-2 and antiviral drug
resistance? *J Antimicrob Chemother* **76**, 2230-2233 (2021).
https://doi.org/10.1093/jac/dkab189

21 Szemiel, A. M., Merits, A., Orton, R. J., MacLean, O. A., Pinto, R. M., Wickenhagen, A.,
Lieber, G., Turnbull, M. L., Wang, S., Furnon, W., Suarez, N. M., Mair, D., da Silva
Filipe, A., Willett, B. J., Wilson, S. J., Patel, A. H., Thomson, E. C., Palmarini, M., Kohl,
A. & Stewart, M. E. In vitro selection of Remdesivir resistance suggests evolutionary
predictability of SARS-CoV-2. *PLoS Pathog* **17**, e1009929 (2021).
https://doi.org/10.1371/journal.ppat.1009929

22 Donovan-Banfield, I. a., Penrice-Randal, R., Goldswain, H., Rzeszutek, A. M., Pilgrim,
J., Bullock, K., Saunders, G., Northey, J., Dong, X. & Ryan, Y. Characterisation of SARS-
CoV-2 genomic variation in response to molnupiravir treatment in the AGILE Phase
IIa clinical trial. *Nature Communications* **13**, 7284-7284 (2022).

23 Sanderson, T., Hisner, R., Donovan-Banfield, I. a., Hartman, H., Løchen, A., Peacock,
T. P. & Ruis, C. A molnupiravir-associated mutational signature in global SARS-CoV-2
genomes. *Nature* **623**, 594-600 (2023). https://doi.org/10.1038/s41586-023-06649-6

24 Li, P., de Vries, A. C., Kamar, N., Peppelenbosch, M. P. & Pan, Q. Monitoring and
managing SARS-CoV-2 evolution in immunocompromised populations. *Lancet
Microbe* **3**, e325-e326 (2022). https://doi.org/10.1016/S2666-5247(22)00061-1

25 Bentley, E. G., Kirby, A., Sharma, P., Kipar, A., Mega, D. F., Bramwell, C., Penrice-
Randal, R., Prince, T., Brown, J. C., Zhou, J., Screaton, G. R., Barclay, W. S., Owen, A.,
Hiscox, J. A. & Stewart, J. P. SARS-CoV-2 Omicron-B.1.1.529 Variant leads to less
severe disease than Pango B and Delta variants strains in a mouse model of severe
COVID-19. *bioRxiv*, 2021.2012.2026.474085 (2021).
https://doi.org/10.1101/2021.12.26.474085

26 Clark, J. J., Penrice-Randal, R., Sharma, P., Kipar, A., Dong, X., Pennington, S. H.,
Marriott, A. E., Colombo, S., Davidson, A., Williamson, M. K., Matthews, D. A., Turtle,
L., Prince, T., Hughes, G. L., Patterson, E. I., Shawli, G., Mega, D. F., Subramaniam, K.,
Sharp, J., McLaughlin, L., Zhou, E.-M., Turner, J. D., Biagini, G., Owen, A., Hiscox, J. A.
& Stewart, J. P. Sequential infection with influenza A virus followed by severe acute
respiratory syndrome coronavirus 2 (SARS-CoV-2) leads to more severe disease and
encephalitis in a mouse model of COVID-19. *bioRxiv*, 2020.2010.2013.334532 (2023).
https://doi.org/10.1101/2020.10.13.334532

27 Gaynor, K. U., Vaysburd, M., Harman, M. A. J., Albecka, A., Jeffrey, P., Beswick, P.,
Papa, G., Chen, L., Mallery, D., McGuinness, B., Van Rietschoten, K., Stanway, S.,
Brear, P., Lulla, A., Ciazynska, K., Chang, V. T., Sharp, J., Neary, M., Box, H., Herriott,
J., Kijak, E., Tatham, L., Bentley, E. G., Sharma, P., Kirby, A., Han, X., Stewart, J. P.,
Owen, A., Briggs, J. A. G., Hyvönen, M., Skynner, M. J. & James, L. C. Multivalent
bicyclic peptides are an effective antiviral modality that can potently inhibit SARS-

CoV-2. *Nature Communications* **14**, 3583 (2023). https://doi.org/10.1038/s41467-
023-39158-1

28 Russell, C. D., Valanciute, A., Gachanja, N. N., Stephen, J., Penrice-Randal, R.,
Armstrong, S. D., Clohisey, S., Wang, B., Al Qsous, W., Wallace, W. A., Oniscu, G. C.,
Stevens, J., Harrison, D. J., Dhaliwal, K., Hiscox, J. A., Baillie, J. K., Akram, A. R.,
Dorward, D. A. & Lucas, C. D. Tissue Proteomic Analysis Identifies Mechanisms and
Stages of Immunopathology in Fatal COVID-19. *Am J Respir Cell Mol Biol* (2021).
https://doi.org/10.1165/rcmb.2021-0358OC

29 Dorward, D. A., Russell, C. D., Um, I. H., Elshani, M., Armstrong, S. D., Penrice-Randal,
R., Millar, T., Lerpiniere, C. E. B., Tagliavini, G., Hartley, C. S., Randle, N. P., Gachanja,
N. N., Potey, P. M. D., Dong, X., Anderson, A. M., Campbell, V. L., Duguid, A. J., Al
Qsous, W., BouHaidar, R., Baillie, J. K., Dhaliwal, K., Wallace, W. A., Bellamy, C. O. C.,
Prost, S., Smith, C., Hiscox, J. A., Harrison, D. J. & Lucas, C. D. Tissue-Specific
Immunopathology in Fatal COVID-19. *Am J Respir Crit Care Med* **203**, 192-201 (2021).
https://doi.org/10.1164/rccm.202008-3265OC

30 Patterson, E. I., Prince, T., Anderson, E. R., Casas-Sanchez, A., Smith, S. L., Cansado-
Utrilla, C., Solomon, T., Griffiths, M. J., Acosta-Serrano, A., Turtle, L. & Hughes, G. L.
Methods of Inactivation of SARS-CoV-2 for Downstream Biological Assays. *J Infect Dis*
222, 1462-1467 (2020). https://doi.org/10.1093/infdis/jiaa507

31 Davidson, A. D., Williamson, M. K., Lewis, S., Shoemark, D., Carroll, M. W., Heesom,
K. J., Zambon, M., Ellis, J., Lewis, P. A., Hiscox, J. A. & Matthews, D. A.
Characterisation of the transcriptome and proteome of SARS-CoV-2 reveals a cell
passage induced in-frame deletion of the furin-like cleavage site from the spike
glycoprotein. *Genome Med* **12**, 68 (2020).

32 Abdelnabi, R., Foo, C. S., Kaptein, S. J. F., Boudewijns, R., Vangeel, L., Jonghe, S. D.,
Jochmans, D., Weynand, B. & Neyts, J. A SCID Mouse Model To Evaluate the Efficacy
of Antivirals against SARS-CoV-2 Infection. *Journal of virology* **96**, e00758-00722
(2022). https://doi.org/doi:10.1128/jvi.00758-22

33 Jeong, J. H., Chokkakula, S., Min, S. C., Kim, B. K., Choi, W.-S., Oh, S., Yun, Y. S., Kang,
D. H., Lee, O.-J., Kim, E.-G., Choi, J.-H., Lee, J.-Y., Choi, Y. K., Baek, Y. H. & Song, M.-S.
Combination therapy with nirmatrelvir and molnupiravir improves the survival of
SARS-CoV-2 infected mice. *Antiviral Research* **208**, 105430 (2022).
https://doi.org/https://doi.org/10.1016/j.antiviral.2022.105430

34 Wahl, A., Gralinski, L. E., Johnson, C. E., Yao, W., Kovarova, M., Dinnon, K. H., Liu, H.,
Madden, V. J., Krzystek, H. M., De, C., White, K. K., Gully, K., Schäfer, A., Zaman, T.,
Leist, S. R., Grant, P. O., Bluemling, G. R., Kolykhalov, A. A., Natchus, M. G., Askin, F.
B., Painter, G., Browne, E. P., Jones, C. D., Pickles, R. J., Baric, R. S. & Garcia, J. V.
SARS-CoV-2 infection is effectively treated and prevented by EIDD-2801. *Nature* **591**,
451-457 (2021). https://doi.org/10.1038/s41586-021-03312-w

35 Stegmann, K. M., Dickmanns, A., Heinen, N., Blaurock, C., Karrasch, T., Breithaupt, A.,
Klopfleisch, R., Uhlig, N., Eberlein, V., Issmail, L., Herrmann, S. T., Schreieck, A.,
Peelen, E., Kohlhof, H., Sadeghi, B., Riek, A., Speakman, J. R., Groß, U., Görlich, D.,
Vitt, D., Müller, T., Grunwald, T., Pfaender, S., Balkema-Buschmann, A. &
Dobbelstein, M. Inhibitors of dihydroorotate dehydrogenase cooperate with
molnupiravir and N4-hydroxycytidine to suppress SARS-CoV-2 replication. *iScience*
25, 104293 (2022). https://doi.org/https://doi.org/10.1016/j.isci.2022.104293

36   De Neck, S., Penrice-Randal, R., Clark, J. J., Sharma, P., Bentley, E. G., Kirby, A., Mega,
D. F., Han, X., Owen, A., Hiscox, J. A., Stewart, J. P. & Kipar, A. The Stereotypic
Response of the Pulmonary Vasculature to Respiratory Viral Infections: Findings in
Mouse Models of SARS-CoV-2, Influenza A and Gammaherpesvirus Infections.
Viruses **15** (2023).

37   Seehusen, F., Clark, J. J., Sharma, P., Bentley, E. G., Kirby, A., Subramaniam, K.,
Wunderlin-Giuliani, S., Hughes, G. L., Patterson, E. I., Michael, B. D., Owen, A.,
Hiscox, J. A., Stewart, J. P. & Kipar, A. Neuroinvasion and Neurotropism by SARS-CoV-
2 Variants in the K18-hACE2 Mouse. *Viruses* **14** (2022).
https://doi.org/10.3390/v14051020

38   Coolen, J. P. M., Wolters, F., Tostmann, A., van Groningen, L. F. J., Bleeker-Rovers, C.
P., Tan, E. C. T. H., van der Geest-Blankert, N., Hautvast, J. L. A., Hopman, J.,
Wertheim, H. F. L., Rahamat-Langendoen, J. C., Storch, M. & Melchers, W. J. G. SARS-
CoV-2 whole-genome sequencing using reverse complement PCR: For easy, fast and
accurate outbreak and variant analysis. *Journal of Clinical Virology* **144**, 104993
(2021). https://doi.org/https://doi.org/10.1016/j.jcv.2021.104993

39   Avanzato, V. A., Matson, M. J., Seifert, S. N., Pryce, R., Williamson, B. N., Anzick, S. L.,
Barbian, K., Judson, S. D., Fischer, E. R., Martens, C., Bowden, T. A., de Wit, E., Riedo,
F. X. & Munster, V. J. Case Study: Prolonged Infectious SARS-CoV-2 Shedding from an
Asymptomatic Immunocompromised Individual with Cancer. *Cell* **183**, 1901-
1912.e1909 (2020). https://doi.org/10.1016/j.cell.2020.10.049

40   Burki, T. The origin of SARS-CoV-2 variants of concern. *Lancet Infect Dis* **22**, 174-175
(2022). https://doi.org/10.1016/s1473-3099(22)00015-9

41   Caccuri, F., Messali, S., Bortolotti, D., Di Silvestre, D., De Palma, A., Cattaneo, C.,
Bertelli, A., Zani, A., Milanesi, M., Giovanetti, M., Campisi, G., Gentili, V., Bugatti, A.,
Filippini, F., Scaltriti, E., Pongolini, S., Tucci, A., Fiorentini, S., d'Ursi, P., Ciccozzi, M.,
Mauri, P., Rizzo, R. & Caruso, A. Competition for dominance within replicating
quasispecies during prolonged SARS-CoV-2 infection in an immunocompromised
host. *Virus Evol* **8**, veac042 (2022). https://doi.org/10.1093/ve/veac042

42   Chen, L., Zody, M. C., Di Germanio, C., Martinelli, R., Mediavilla, J. R., Cunningham,
M. H., Composto, K., Chow, K. F., Kordalewska, M., Corvelo, A., Oschwald, D. M.,
Fennessey, S., Zetkulic, M., Dar, S., Kramer, Y., Mathema, B., Germer, S., Stone, M.,
Simmons, G., Busch, M. P., Maniatis, T., Perlin, D. S. & Kreiswirth, B. N. Emergence of
Multiple SARS-CoV-2 Antibody Escape Variants in an Immunocompromised Host
Undergoing Convalescent Plasma Treatment. *mSphere* **6**, e0048021 (2021).
https://doi.org/10.1128/mSphere.00480-21

43   Choi, B., Choudhary, M. C., Regan, J., Sparks, J. A., Padera, R. F., Qiu, X., Solomon, I.
H., Kuo, H.-H., Boucau, J., Bowman, K., Adhikari, U. D., Winkler, M. L., Mueller, A. A.,
Hsu, T. Y. T., Desjardins, M., Baden, L. R., Chan, B. T., Walker, B. D., Lichterfeld, M.,
Brigl, M., Kwon, D. S., Kanjilal, S., Richardson, E. T., Jonsson, A. H., Alter, G., Barczak,
A. K., Hanage, W. P., Yu, X. G., Gaiha, G. D., Seaman, M. S., Cernadas, M. & Li, J. Z.
Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host. *New*
England Journal of Medicine **383**, 2291-2293 (2020).
https://doi.org/10.1056/NEJMc2031364

44   Peng, K. W., Myers, R., Greenslade, A., Mader, E., Greiner, S., Federspiel, M. J.,
Dispenzieri, A. & Russell, S. J. Using clinically approved cyclophosphamide regimens

to control the humoral immune response to oncolytic viruses. *Gene Therapy* **20**, 255-
261 (2013). https://doi.org/10.1038/gt.2012.31

45 Schaecher, S. R., Stabenow, J., Oberle, C., Schriewer, J., Buller, R. M., Sagartz, J. E. &
Pekosz, A. An immunosuppressed Syrian golden hamster model for SARS-CoV
infection. *Virology* **380**, 312-321 (2008). https://doi.org/10.1016/j.virol.2008.07.026

46 Ramasamy, S., Kolloli, A., Kumar, R., Husain, S., Soteropoulos, P., Chang, T. L. &
Subbian, S. Comprehensive Analysis of Disease Pathology in Immunocompetent and
Immunocompromised Hosts following Pulmonary SARS-CoV-2 Infection.
Biomedicines **10**, 1343 (2022).

47 Brocato, R. L., Principe, L. M., Kim, R. K., Zeng, X., Williams, J. A., Liu, Y., Li, R., Smith,
J. M., Golden, J. W., Gangemi, D., Youssef, S., Wang, Z., Glanville, J. & Hooper, J. W.
Disruption of Adaptive Immunity Enhances Disease in SARS-CoV-2-Infected Syrian
Hamsters. *J Virol* **94** (2020). https://doi.org/10.1128/jvi.01683-20

48 Salzer, R., Clark, J. J., Vaysburd, M., Chang, V. T., Albecka, A., Kiss, L., Sharma, P.,
Gonzalez Llamazares, A., Kipar, A., Hiscox, J. A., Owen, A., Aricescu, A. R., Stewart, J.
P., James, L. C. & Lowe, J. Single-dose immunisation with a multimerised SARS-CoV-2
receptor binding domain (RBD) induces an enhanced and protective response in
mice. *FEBS Lett* **595**, 2323-2340 (2021). https://doi.org/10.1002/1873-3468.14171

49 Legebeke, J., Lord, J., Penrice-Randal, R., Vallejo, A. F., Poole, S., Brendish, N. J.,
Dong, X., Hartley, C., Holloway, J. W., Lucas, J. S., Williams, A. P., Wheway, G.,
Strazzeri, F., Gardner, A., Schofield, J. P. R., Skipp, P. J., Hiscox, J. A., Polak, M. E.,
Clark, T. W. & Baralle, D. Evaluating the Immune Response in Treatment-Naive
Hospitalised Patients With Influenza and COVID-19. *Front Immunol* **13**, 853265
(2022). https://doi.org/10.3389/fimmu.2022.853265

50 Escalera, A., Gonzalez-Reiche, A. S., Aslam, S., Mena, I., Laporte, M., Pearl, R. L.,
Fossati, A., Rathnasinghe, R., Alshammary, H., van de Guchte, A., Farrugia, K., Qin, Y.,
Bouhaddou, M., Kehrer, T., Zuliani-Alvarez, L., Meekins, D. A., Balaraman, V.,
McDowell, C., Richt, J. A., Bajic, G., Sordillo, E. M., Dejosez, M., Zwaka, T. P., Krogan,
N. J., Simon, V., Albrecht, R. A., van Bakel, H., García-Sastre, A. & Aydillo, T.
Mutations in SARS-CoV-2 variants of concern link to increased spike cleavage and
virus transmission. *Cell Host Microbe* **30**, 373-387.e377 (2022).
https://doi.org/10.1016/j.chom.2022.01.006

51 Braun, K. M., Moreno, G. K., Halfmann, P. J., Hodcroft, E. B., Baker, D. A., Boehm, E.
C., Weiler, A. M., Haj, A. K., Hatta, M., Chiba, S., Maemura, T., Kawaoka, Y., Koelle, K.,
O'Connor, D. H. & Friedrich, T. C. Transmission of SARS-CoV-2 in domestic cats
imposes a narrow bottleneck. *PLOS Pathogens* **17**, e1009373 (2021).
https://doi.org/10.1371/journal.ppat.1009373

52 Mizuki, Y., Keiko, T., Youko, H., Jun-ichiro, I., Yasushi, K. & Jin, G. SARS-CoV-2
Omicron spike H655Y mutation is responsible for enhancement of the endosomal
entry pathway and reduction of cell surface entry pathways. *bioRxiv*,
2022.2003.2021.485084 (2022). https://doi.org/10.1101/2022.03.21.485084

53 Ou, J., Lan, W., Wu, X., Zhao, T., Duan, B., Yang, P., Ren, Y., Quan, L., Zhao, W., Seto,
D., Chodosh, J., Luo, Z., Wu, J. & Zhang, Q. Tracking SARS-CoV-2 Omicron diverse
spike gene mutations identifies multiple inter-variant recombination events. *Signal
Transduction and Targeted Therapy* **7**, 138 (2022). https://doi.org/10.1038/s41392-
022-00992-2

54   Jeelani, R., Khan, S. N., Shaeib, F., Kohan-Ghadr, H. R., Aldhaheri, S. R., Najafi, T.,
Thakur, M., Morris, R. & Abu-Soud, H. M. Cyclophosphamide and acrolein induced
oxidative stress leading to deterioration of metaphase II mouse oocyte quality. *Free*
Radic Biol Med **110**, 11-18 (2017).
https://doi.org/10.1016/j.freeradbiomed.2017.05.006
55   Pimenta, G. F., Awata, W. M. C., Orlandin, G. G., Silva-Neto, J. A., Assis, V. O., da
Costa, R. M., Bruder-Nascimento, T., Tostes, R. C. & Tirapelli, C. R. Melatonin
prevents overproduction of reactive oxygen species and vascular dysfunction
induced by cyclophosphamide. *Life Sciences* **338**, 122361 (2024).
https://doi.org/https://doi.org/10.1016/j.lfs.2023.122361
879
880
881
882

SUPPLEMENTARY TABLES

Table S1: The number of samples per cohort, sample type and DPI that were included in the sequencing analysis. Samples with less than 90% coverage and poor or mediocre quality control results determined by Nextclade CLI, were excluded from the analysis. *Multiple end point time-points due to humane end-point variation.

Cohort	Sample type	DPI[*]	n
Cyclophosphamide and Molnupiravir	Swab	1	4
		3	4
		5	3
	Lung Tissue	7	4
	Nasal Tissue	7	4
Cyclophosphamide and Nirmatrelvir	Swab	1	1
		3	2
		5	1
	Lung Tissue	4	1
		7	2
	Nasal Tissue	4	1
		7	2
Cyclophosphamide only	Swab	1	4
		3	4
		4	1
		5	2
	Lung Tissue	4	2
		6	2
	Nasal Tissue	4	1
		6	2
Cyclophosphamide, Nirmatrelvir and Molnupiravir	Swab	1	1
		3	2

Cohort	Sample type	DPI[*]	n
	Lung Tissue	3	1
		7	2
	Nasal Tissue	3	1
		7	3
Molnupiravir	Swab	1	4
		3	4
		5	1
	Lung Tissue	7	3
	Nasal Tissue	7	1
Vehicle	Swab	1	4
		3	4
		5	4
	Lung Tissue	6	4
	Nasal Tissue	6	3

894 **Table S2:** *Unique phenotypic changes in SARS-CoV-2 within the dataset. Input virus*

895 *amino acids and S: H665Y were excluded.*

Amino acid change	Occurrence in dataset	Tissue type	Cohorts
S: D215H	2	Swab	Vehicle, Cyclophosphamide, Molnupiravir &

Amino acid change	Occurrence in dataset	Tissue type	Cohorts
			nirmatrelvir
ORF1a: G2581S	1	Swab	Cyclophosphamide & nirmatrelvir
ORF3a: P42L	1	Swab	Cyclophosphamide & Molnupiravir
ORF3a: Q57H	2	Nasal Tissue Swab	Cyclophosphamide, Molnupiravir & nirmatrelvir, Cyclophosphamide only
E: Y42H	1	Swab	Cyclophosphamide & nirmatrelvir
ORF1a: G82D S: V47I	1	Lung Tissue	Cyclophosphamide, nirmatrelvir and Molnupiravir Cyclophosphamide, nirmatrelvir and Molnupiravir
ORF1a: T568I ORF1a: C2160Y ORF1b: F312L ORF1b: V2515I ORF8: F108L S: A27T	1	Nasal Tissue	Cyclophosphamide, nirmatrelvir and Molnupiravir
ORF1a: F741L ORF1a: A872T ORF7b: S5P S: A890V S: D985N	1	Nasal Tissue	Cyclophosphamide & nirmatrelvir

Amino acid change	Occurrence in dataset	Tissue type	Cohorts
ORF1a: F190L	1	Nasal Tissue	Molnupiravir
ORF1a: G209S			
ORF1a: V2130I			
ORF1b: R1729H			
ORF3a:M1I			
ORF3a: G224S			
S: S247R			
S: V483I			

Supplementary Figures

899

900 **Figure S1**

901

902 Fig S1: Bioinformatic workflow used for sequencing analysis.

Fig S2: The proportion of amino acid H and Y at position 655 in the Spike protein over time or in lung or nasal tissues. The S: H655Y mutation was observed at the minor variant level if not observed in the consensus level. Each point represents one animal within the cohort.

903
904

905
906 Fig S3: The coverage (A) and average depth (B) of the swabs used in the minor
907 variant analysis.
908

Relationship Between Ct Value and Mean Ts/Tv Ratio

915
916
Fig S4: The scatter plot shows the relationship between Ct values and the mean
transition/transversion (Ts/Tv) ratios calculated with the DiversiTools output. Each
blue dot represents a swab sample used in the minor variant analysis. The red line
indicates the linear regression fit, with the shaded region showing the 95%
confidence interval. Statistical analysis revealed no significant correlation between Ct
values and Ts/Tv ratios (R^2 = 0.012 , p = 0.45), supporting the conclusion that
there is no association between the two variables.
924
925
926

935
936 *Fig S5: Minor variation for all base changes between day 1 and day 3 post infection.*
937 ** represents a P value <0.05 (Mann Whitney U test).*