Blood metabolomic profiling reveals new targets in the management of psychological symptoms associated with alcohol use disorder

Sophie Leclercq [a,1], Hany Ahmed[b,1], Camille Amadieu[c], Géraldine Petit[d], Ville Koistinen[b,e],

Quentin Leyrolle[g,c], Marie Poncin[d], Peter Stärkel[h], Eloise Kok[i,j], Pekka J. Karhunen[j], Philippe

De Timary[d], Sophie Layé[g], Audrey M. Neyrinck[c], Olli Kärkkäinen[f], Kati Hanhineva[b, e,2],

Nathalie M. Delzenne[c,2]

[1]Co-first authors

[2]Co-last authors

Affiliations:

[a]Laboratory of Nutritional Psychiatry, Institute of Neuroscience, UCLouvain, Université

catholique de Louvain, Brussels, Belgium

[b]Food Sciences Unit, Department of Life Technologies, University of Turku, Turku, Finland

[c]Metabolism and Nutrition Research Group, Louvain Drug Research Institute, UCLouvain,

Université catholique de Louvain, Brussels, Belgium

[d]Department of Adult Psychiatry, Cliniques Universitaires Saint-Luc and Institute of

Neuroscience, UCLouvain, Université catholique de Louvain, Brussels, Belgium

[e]School of Medicine, Institute of Public Health and Clinical Nutrition, University of Eastern

Finland, Kuopio, Finland

[f]School of Pharmacy, University of Eastern Finland, Kuopio, Finland

[g]Université de Bordeaux, INRAE, Bordeaux INP, NutriNeurO, UMR 1286, F-33000 Bordeaux,

France

[h]Department of gastro-enterology, Cliniques Universitaires Saint Luc, Brussels, Belgium

[i]Department of Pathology, University of Helsinki, Helsinki, Finland

[j]Faculty of Medicine and Health Technology, Tampere University and Fimlab Laboratories,

Tampere, Finland

Running title: Blood metabolomics and alcohol use disorder symptoms

Keywords: metabolomics; alcohol use disorder; depression; anxiety; craving; microbiota

Corresponding author: Nathalie M. Delzenne, Metabolism and Nutrition Research Group,

Louvain Drug Research Institute, UCLouvain, Université catholique de Louvain, avenue E.

Mounier   box   B1.73.11,   B-1200   Brussels,   Belgium;   E-mail   address:

nathalie.delzenne@uclouvain.be; Phone: +32 2 764 73 69

ABSTRACT

Alcohol use disorder (AUD) is a global health problem with limited therapeutic options. The biochemical mechanisms that lead to alcohol addiction are not yet fully understood, and in this respect, metabolomics represents a promising approach to decipher metabolic events related to AUD. The plasma metabolome contains a plethora of bioactive molecules that reflects the functional changes in host metabolism but also the impact of the gut microbiome and nutritional habits. In this study, we investigated the impact of chronic alcohol abuse, and of a three-week period of alcohol abstinence, on the blood metabolome (non-targeted LC-MS metabolomics analysis) in 96 patients diagnosed with severe alcohol use disorder (AUD). We found that the plasma levels of different lipids ((lyso)phosphatidylcholines, long-chain fatty acids), short-chain fatty acids (i.e. 3-hydroxyvaleric acid) and bile acids were altered in AUD patients. In addition, several microbial metabolites, including indole-3-propionic acid, p-cresol sulfate, hippuric acid, pyrocatechol sulfate, and metabolites belonging to xanthine class (paraxanthine, theobromine and theophylline) were sensitive to alcohol abuse and alcohol withdrawal. 3-Hydroxyvaleric acid, caffeine metabolites (theobromine, paraxanthine and theophylline) and microbial metabolites (hippuric acid and pyrocatechol sulfate) were correlated with anxiety, depression and alcohol craving. Metabolomics analysis in post-mortem samples of frontal cortex and cerebrospinal fluid of those consuming a high level of alcohol revealed that those metabolites can be found also in brain tissue. Our data allow to for the identification of neuroactive metabolites, from interactions between food components and microbiota, which may represent new targets in the management of neuropsychiatric diseases such as AUD.

The study was registered at clinicaltrial.gov under the identification number NCT03803709.

Introduction

Alcohol use disorder (AUD) is a global health problem accounting for substantial difficulties for the individuals who consume, their related persons, and for the society. The biochemical mechanisms that lead to alcohol addiction are not yet fully understood, and in this respect, metabolomics represents an interesting approach to decipher metabolic events related to AUD. Improving our understanding of the pathology could lead to discovering potential novel targets for therapies (1).

Alcohol consumption clearly leads to alterations of the circulating metabolome (1–4). For instance, changes in the levels of lipids (fatty acids, phosphatidylcholine, steroids) and amino acids (glutamine, tyrosine, alanine, serotonin, asparagine) are commonly observed and, interestingly, some changes in the blood metabolite profile precedes the emergence of alcohol use related diseases, such as lower levels of serotonin and asparagine (5). Metabolomics studies conducted in rodent models of alcohol exposure have mostly targeted the liver tissue or urine (6–8). In human, plasma or serum metabolomics studies have tested the effects of low/moderate/excessive alcohol intake (9) but are rarely performed in clinical populations of AUD patients, including analysis of central nervous system (CNS) tissues(10). Furthermore, the methodology could differ across studies, some using nuclear magnetic resonance (NMR) while others using more sensitive mass spectrometry (MS) coupled with liquid or gas chromatography (LC or GC), leading to the detection of different metabolites.

The human blood metabolome consists of 1) small molecules that directly represent the functional changes in host metabolism, 2) metabolites produced by intestinal micro-organisms and 3) metabolites originating from nutrition of other exogenous sources like drugs (1,11).

The aims of this study are multiple. First, we investigated the impact of severe AUD on the blood metabolome by non-targeted LC-MS metabolomics analysis. Second, we investigated the impact of a short-term alcohol abstinence on the blood metabolome followed by assessing the

correlations between the blood metabolome and psychological symptoms developed in AUD

patients. Last, we hypothesized that metabolites significantly correlated with depression,

anxiety or alcohol craving could potentially have neuroactive properties, and therefore the

presence of those neuroactive metabolites was confirmed in the central nervous system using

post-mortem analysis of frontal cortex and cerebrospinal fluid of persons with a history of heavy

alcohol use.

Our data bring new insights on xenobiotics- or microbial-derived neuroactive metabolites,

which can represent an interesting strategy to prevent or treat psychiatric disorders such as

AUD.

RESULTS

Clinical characteristics of the study participants

Two cohorts of AUD patients (ALCOHOLBIS and GUT2BRAIN) were used in this study.

Both groups of patients were similar in terms of age, gender, smoking and drinking habits and

presented with high scores of depression, anxiety and alcohol craving at T1 (Table 1). The only

differences between the cohorts resided in the number of previous withdrawal cures and the

score of craving (compulsion sub-score) which were higher in the GUT2BRAIN cohort

compared to ALCOHOLBIS cohort. Nevertheless, in both cohorts, the score of craving was

high, as usually seen in other cohorts of AUD patients (12–14). Therefore, for the analysis of

metabolomics data, both cohorts of AUD patients were combined, and compared to a group of

healthy controls.

Alterations in the plasma metabolome of AUD patients

The metabolomics analysis allowed for sorting out a total of 11,651 features from the four

analytical modes of the plasma samples. An unsupervised PCA model of the plasma

metabolomic profiles between healthy controls and AUD patients at the beginning of the

withdrawal (T1) is shown in Fig. 1a. In addition, the scores plot and the performance of a

supervised sPLS-DA model are shown in Figs. S1 and S2, respectively. Between healthy

controls and AUD T1, the annotated differential (Welch *t*-test $q<0.05$ and sPLS-DA VIP score

> 2.0) metabolites included compounds from several metabolite classes as indicated in Figs.

1b, 1c and Table S3. Compared to healthy controls, the metabolic profiles of AUD patients

were characterized by an increase in long-chain fatty acids, such as 16:1 (palmitoleic acid), 18:1

(octadecenoic acid) and 22:4 (docosatetraenoic acid) fatty acids and phospholipids holding

these fatty acids. In addition, several drugs (like diazepam, trazodone) and metabolites with

steroid backbone such glycinated bile acids (glycohyodeoxycholic acid and
glycochenodeoxycholic acid), steroid hormones and acylcarnitines were increased in the AUD
group. We also observed a significant increase in vitamin B6 metabolite, 4-pyridoxic acid,
nicotine metabolite cotinine, a hydroxy fatty acid 3-hydroxyvaleric acid and stress hormone
cortisol. However, LPCs holding a saturated odd-chain (e.g. LPC 15:0 and LPC 17:0),
polyunsaturated 18-carbon fatty acid or an ether bond (O-) containing lipid were consistently
decreased in AUD compared to controls (Fig. 1c). Further, we also showed a decrease in furan
fatty acids 3-carboxy-4-methyl-5-pentyl-2-furanpropanoic acid (3-CMPFP) and 3-carboxy-4-
methyl-5-propyl-2-furanpropionic acid (CMPF), in a carotenoid compound and in several
metabolites belonging to the family of xanthines. In addition, several amino-acid derived
bacterial metabolites such as pipecolic acid, 3-indole propionic acid, p-cresol sulfate and
hippuric acid were significantly decreased in AUD patients compared to controls.

We then conducted a correlational analysis between blood metabolites and alcohol consumption
reported by the patients. Alcohol intake was positively correlated with annotated bile acids,
steroids and drugs while xanthines (paraxanthine, theobromine and theophylline), odd-chain or
ether-bond LPCs and bacterial metabolite hippuric acid, p-cresol sulfate, pyrocatechol sulfate,
and 3-indole propionic acid showed negative correlations (Fig. 1d).

Alcohol withdrawal shapes the plasma metabolome

The score plot of the sPLS-DA in Fig. 2a shows a clear discrimination in plasma metabolomic
profiles in the course of withdrawal in AUD patients. The sPLS-DA model performance and
the unsupervised PCA model scores plot are shown in supplementary Figs. S3 and S4,
respectively. Annotated metabolites discriminating AUD groups (paired *t*-test $q<0.05$, sPLS-
DA VIP score > 2.0) before (T1) and after (T2) the 3-week withdrawal period included
metabolites from a range of chemical classes (Figs. 2b, 2c, Table S3). Apart from the

metabolites belonging to the class of bilirubins, the levels of identified metabolites that were significantly changed upon alcohol withdrawal (the major ones being indoles, acylcarnitines, glycerophospholipids and xanthines) came back towards to the levels observed in controls (Table S3). In the course of alcohol abstinence, we noted a significant decrease in 16-chain acylcarnitines, LPCs with 16- or 18-chain fatty acid tails excluding LPCs with ether bonds, retinol, tryptophyl-phenylalanine dipeptide and 3-hydroxyvaleric acid (Fig. 2c). On the contrary, LPCs with odd-chain fatty acid tails or ether bonds show a significant increase along with tryptophan derivatives 3-indoleacetic acid and 3-indolepropanoic acid and metabolites of the xanthine family such as theophylline, paraxanthine and trigonelline during alcohol abstinence. Moreover, Fig. 2c demonstrates that alcohol withdrawal shifted the AUD plasma metabolic fingerprint towards healthy controls.

Correlations between blood metabolites and psychological symptoms

Correlation analysis shows that, at T1, 96 annotated features were significantly ($p<0.05$) correlated with psychological scores of anxiety, depression and alcohol craving (with sub-scores of obsession and compulsion) (Fig 3). Annotated bile acids, drugs, lysophosphoethanolamines (LPE), fatty acids, LPCs apart from LPCs with an ether-bond or 15:0 or 17:0 fatty acid tail were consistently positively correlated with psychological symptoms, and more particularly with the compulsive component of alcohol craving. Members of the xanthine family, pyrocatechol sulfate, a pentose sugar (mannose/fructose), hippuric acid, 1-methyl-pyridone-carboxamide, acylcarnitines with maximum of 10 carbons, creatinine and kynurenine were negatively correlated with psychological symptoms. Within the acylcarnitine metabolite class, an interesting pattern was observed, as the shorter chain-length acylcarnitines were consistently negatively correlated with the psychological parameters, and the long-chain ones demonstrated positive correlation.

Alterations in the brain metabolome of AUD patients

Based on the annotated significantly altered plasma metabolites, we conducted a targeted search in a metabolomics dataset consisting of CSF and frontal cortex samples collected from deceased individuals with a history of heavy alcohol use and control individuals. 79 and 74 of the annotated plasma metabolites were identified from the CSF and frontal cortex, respectively (Fig. 4a, Table S3). We looked specifically at metabolites significantly correlated with at least one psychological symptom. 3-Hydroxyvaleric acid, cotinine, theobromine and paraxanthine were indeed present in the CNS and significantly (Welch t-test $p<0.05$) different between heavy alcohol use and control groups in both frontal cortex (Fig 4b) and CSF (Fig 4c). Additional significantly altered metabolites found only in the frontal cortex were LPE 20:3, nordiazepam, PC 16:0_18:1 and urea (Fig. 4b). In the CSF, the independent differential metabolites were FA 16:1 (palmitoleic acid), hippuric acid, LPCs 16:0 and 18:1, LPEs 16:0, 18:1 and 20:4 and pyrocatechol sulfate (Fig. 4c).

DISCUSSION

The circulating metabolome reflects the crosstalk between nutrition, microbiome and host metabolism (15), with diet and microbiome being the strongest determinants of the human blood metabolome (11,16). In this study, we showed the impact of chronic alcohol abuse, and the impact of a short-term abstinence, on the blood metabolome. We analyzed the correlations between blood metabolites and psychological symptoms, as well as the presence of identified metabolites in the CNS of individuals considered as heavy alcohol drinkers.

Impact of chronic alcohol abuse on the blood metabolome

In 2019, a review summarized the results obtained from 23 studies that have used a metabolomics approach for measuring changes in metabolite profiles in relation to alcohol use (1). Changes in lipids have been highlighted as the most consistent changes across studies. Lipids are an integral part of cell membranes and signaling molecules in the body. PCs and LPCs have been suggested to form a new class of biomarkers for alcohol consumption (2). For instance, in our study palmitoleic acid (FA 16:1) was largely increased in AUD patients compared to controls, and in other studies, this metabolite has likewise been significantly associated with alcohol consumption (17,18). Another clear observation in our study was the lower level of odd-chain lipids in AUD patients. Since the lipids containing FA 15:0 and FA 17:0 have been suggested to be products from bacterial metabolism (19,20), the existence of gut dysbiosis in AUD patients could explain the lower abundance of LPC 17:0 and LPC 15:0 (14).

We found that some bile acids, sulphated steroids and 3-hydroxyvaleric acid were positively associated with the amount of alcohol consumed. Sulphated steroids and hydroxyvalerate have previously been associated with alcohol intake (21,22). Metabolites belonging to the xanthine

family (theobromine, theophylline, paraxanthine) and microbial metabolites (hippuric acid, indole-3-propionic acid, p-cresol sulfate, pyrocatechol sulfate) were negatively correlated with alcohol consumption. Altogether, these results suggest that these metabolites are sensitive to alcohol exposure. Interestingly, these metabolites were also correlated with the severity of the psychological symptoms suggesting that they could play a role in the symptomatology of alcohol addiction.

Effect of short-term alcohol abstinence on the blood metabolome

Since alcohol consumption is known to influence lipid metabolism, it was expected that a short-term alcohol abstinence could reverse or ameliorate lipidomic alterations. Indeed, we found that some phospholipids that were increased in AUD patients at baseline, such as PC 16:0_18:1, PC18:1_14:0 and 16:0_16:1 as well as LPC 16:1, were downregulated during alcohol withdrawal to reach the levels of healthy controls after detox. On the other hand, LPC 15:0 and LPC 17:0 that were decreased in AUD patients at baseline, increased during alcohol withdrawal, but did not reach the levels of controls at the end of detox.

The metabolite that contributed the most to the differences observed with alcohol detoxification was bilirubin. While bilirubin was not statistically higher in AUD patients at baseline vs controls, we observed a significant reduction of this metabolite after a 3-week alcohol withdrawal. In a previous study, serum bilirubin was found to be associated with alcohol consumption, cigarette smoking and coffee consumption (23). Interestingly, the caffeine metabolites belonging to the xanthine family such as paraxanthine, theophylline and theobromine that were decreased at baseline in AUD patients compared to controls, increased significantly during alcohol withdrawal to reach the levels of healthy controls. Changes in dietary intake of coffee, tea and chocolate during alcohol withdrawal could explain these results (data not shown). Also, the bacterial metabolites indole derivatives such as 3-indolepropionic

acid and 3-indoleacetic acid increased during alcohol withdrawal to reach the levels of healthy

controls. Intriguingly, 3-hydroxyvaleric acid significantly decreased during alcohol withdrawal

and was found to be lower than healthy controls at the end of detoxification period.

Metabolites that remained significantly higher in AUD patients at the end of detoxification

compared to controls are stress hormone cortisol, palmitoleic acid (FA 16:1), some bile acids,

some drugs (diazepam, trazodone), vitB6 metabolite (4-pyridoxic acid, which is likely due to

the fact that patients received vitamin B supplement during their hospital stay) and cotinine

(nicotine metabolite that reveals the higher proportion of smokers in AUD patients compared

to controls).

Identification of blood metabolites with potential neuroactive properties

The metabolites belonging to the xanthine family (theobromine, paraxanthine and theophylline)

are metabolites of caffeine produced upon cytochrome P450 dependent oxidation in the liver.

They were all decreased in the blood of AUD patients at baseline and were negatively correlated

with alcohol intake, alcohol craving, depression and anxiety. The decrease in caffeine

metabolites has previously been described in the urine of AUD patients, that is linked to the

increasing severity of alcoholic liver disease (24).

Theobromine is the principle alkaloid found in cocoa beans and is responsible for the bitter taste

of chocolate. It is known for its mood improving effect (25). Like caffeine, theobromine is an

inhibitor of brain adenosine receptors and phosphodiesterase. A study in rats showed that the

antagonist of A2a adenosine receptor produced a reduction of ethanol reinforcement (26),

suggesting adenosine receptor as a potential target for the treatment of alcohol abuse. In a

randomized, double-blind, placebo-controlled trial, the phosphodiesterase inhibitor

pentoxifylline associated with escitalopram showed greater reduction of depression scores

compared to patients receiving escitalopram alone (27). In another study, Apremilast which is

also a phosphodiesterase inhibitor, suppressed excessive alcohol drinking in AUD patients (28).

Paraxanthine has a psychostimulant effect and can modulate dopamine release in the striatum

(29). Interestingly, in 2017 a systematic review indicated that consumption of coffee, tea and

cocoa could have protective effects against depression (30).

Lipids, and mostly LPCs (except ether LPC derivatives) and LPEs were significantly and

positively correlated with the compulsive component of alcohol craving. LPCs are secreted by

the liver and are actively transported via the blood-brain barrier (BBB) and have been associated

with pro-inflammatory events (31). LPCs are also precursors of brain lysophosphatidic acid

(LPA), which regulates glutamatergic transmission and cortical excitability within the CNS.

Recently, LPA has been shown to induce hyperphagia following food restriction and this effect

was dependent on hypothalamic agouti-related peptide (AgRP) neurons (32). AgRP neurons

have also been implicated in circuitry controlling non-feeding behavior, including those

associated with reward, anxiety and compulsive disorders, more particularly in anorexia

nervosa (33). Therefore, we hypothesize that the positive correlation between peripheral LPC

and compulsion for alcohol drinking found in AUD patients who have just been deprived of

alcohol could be mediated by the effect of LPA on AgRP neurons. Consistent with that,

postmortem brain tissues from patients consuming a high intake of alcohol showed increased

levels of many LPCs (10).

Circulating bile acids can reach the brain by crossing the BBB, either by simple diffusion or

active transport. Some bile acids show neuroprotective effects (34) while others are rather

neurotoxic (35). In Alzheimer disease patients, the levels of glycochenodeoxycholic acid was

associated with worse cognition (36). In our study, both primary (glycochenodeoxycholic acid

and glycocholic acid) and secondary (glycohyodeoxycholic acid, tauroursodeoxycholic acid)

bile acids were positively correlated with depression and anxiety in AUD patients.

3-Hydroxyvaleric acid, also called β-hydroxypentanoate, was significantly and positively

correlated with anxiety and alcohol craving. This metabolite is formed from odd carbon fatty

acids in the liver and can reach the brain. 3-hydroxyvaleric acid is a C5-ketone body and is a

precursor of propionyl-CoA that refills intermediates of citric acid cycle and is useful for

alternative energy fuel in the brain (37,38).

Other co-metabolites, i.e. produced by the gut microbiota and then processed by the liver, were

negatively correlated with all psychological symptoms. Pyrocatechol sulfate is a phenolic

compound derived from the gut microbiota, present in the CSF of mice, and implicated in

synapse formation and fear extinction learning (39). In Parkinson disease patients, the plasma

level of pyrocatechol sulfate is decreased compared to controls (40). In our study, blood

pyrocatechol sulfate was significantly and negatively correlated with all psychological

symptoms of AUD patients (i.e. anxiety, depression and craving) suggesting a neuroprotective

role of this metabolite. Interestingly, 4-ethylphenylsulfate, another gut-derived metabolite

linked with neurodevelopment abnormalities, autism and anxiety behavior in mice (41–43)

showed negative correlations with depression and the compulsive component of alcohol

craving.

Hippuric acid, the glycine conjugate of benzoic acid has long been associated with the microbial

degradation of specific dietary components, including polyphenolic compounds (like

chlorogenic acid and catechin) found in fruits, vegetables, coffee and tea (44). Hippuric acid is

indeed a host-microbe cometabolite (45). It is synthesized in the liver and in the renal cortex

from the microbial metabolite benzoate. The plasma concentration of hippuric acid has been

shown to be 17-fold higher in conventional mice compared with their germ-free counterparts

suggesting a substantial contribution of the gut microbiota in its production (46). We showed

that blood hippurate levels, that correlated negatively with anxiety, depression and craving,

were decreased in AUD patients, as shown in ethanol-treated mice (6) and humans

characterized by high alcohol intake and those with major depression (47,48). Urinary hippurate

excretion is also decreased in depression, schizophrenia and autism spectrum disorders patients

(44). A recent Mendelian randomization study including > 13,000 individuals from five

European cohorts characterized for depression suggested that low hippuric acid levels in the

circulation is part of the causal pathway leading to depression (49), which was consistent with

a significant decrease of the dietary sources of hippuric acid including fresh fruits and

vegetables in depressed patients (49).

Another way to support the neuroactive effects of the blood metabolites that are correlated with

one or several psychological factors is to demonstrate their presence in the brain. We therefore

conducted a targeted search in a database of post-mortem frontal cortex and CSF metabolomics

analysis (10) and found that 3-hydrovaleric acid, caffeine metabolites (theobromine,

paraxanthine and theophylline) and microbial metabolites (hippuric acid and pyrocatechol

sulfate) that were correlated with anxiety, depression and alcohol craving in our plasma cohort

were also present in the brain and in CSF, and the direction of their changes in the plasma

(increase or decrease) mimicked changes in the central nervous system.

Advantages and limitations of the study

Most of the studies assessing the impact of alcohol consumption on the blood metabolome were

cross-sectional, and only included male participants (47). Here, we reported longitudinal data

to assess the impact of a short-term alcohol abstinence on the blood metabolome, both in male

and female AUD patients. In the study of Zhu et al (9), the AUD patients recruited were alcohol

abstinent, but for various periods of abstinence. To avoid the bias of abstinence duration, our

patients were enrolled in a rigorous and standardized manner, within 24 hours after the last

drink. Furthermore, alcohol consumption was carefully evaluated with the time line follow back

method, which allows precise calculation of the amount (and type) of alcohol consumed (50).

This study also presents some limitations. First, the metabolomics analysis was conducted with LC-MS while some important molecules, like lipoproteins, could have been measured with NMR-based methods. Combining NMR and MS-based methods could have covered a wider spectrum of metabolites. However, the non-targeted metabolic profiling with two different chromatographic methods and ionization polarities covers a wide range of metabolites ideal for our discovery-based approach. Large studies are usually required in metabolomics to observe small and medium size changes. Here, we included only 96 AUD patients, but they were all well characterized and received standardized therapies (for instance, vitB supplementation) during alcohol withdrawal.

The selection of the control group is always challenging in alcohol research. Here, the healthy subjects were matched for sex, age and BMI but not for smoking status or nutritional intake. Alcohol addiction is a major cause of malnutrition in developed countries and tobacco smoking is more prevalent in alcohol users compared to healthy subjects. These two main confounding factors, although being an integral part of the alcoholic pathology, are known to influence the blood metabolome (51–53).

Conclusion

LC-MS metabolomics plasma analysis allowed for the identification of metabolites that were clearly linked to alcohol consumption, and reflected changes in metabolism, alterations of nutritional status, and gut microbial dysbiosis associated with alcohol intake. In particular the changes in lipid class involving odd-chain fatty acids and ether-bond lipids as well as compounds produced by gut microbiota seem to be the most prominent indicators of metabolic malfunction related to alcohol use disorder, and thus warrant further studies and targeted intervention. Also, the discovery of metabolites associated with behavioral and psychiatric traits

related to AUD were of importance, and could be considered potential new therapeutic targets

in the management of AUD, namely as adjuvants in the period of alcohol abstinence. The

novelty of our work was to characterize the impact of chronic alcohol abuse on the blood

metabolome, and the impact of a short-term alcohol abstinence in the same individuals, within

a cohort that included both male and female patients. Intervention studies are needed in order

to bring the proof of concept that nutritional interventions – namely the addition of specific

lipids, or of nutrients modulating the gut microbiome - for example prebiotic dietary fibers -

may be essential and so far underestimated components of alcohol withdrawal efficacy.

SUBJECTS AND METHODS

Study design and participants

A total of 96 AUD patients hospitalized for a 3-week detoxification program in the alcohol

withdrawal unit at Cliniques Universitaires Saint-Luc, Brussels, Belgium were recruited. These

patients belong to two different cohorts, namely ALCOHOLBIS (patients recruited in 2015 and

2019) and GUT2BRAIN (patients recruited in 2018-2019) (Table 1). The severity of AUD was

evaluated by a psychiatrist using the Diagnostic and Statistical Manual of Mental Disorders

(DSM) criteria, fourth edition (DSM-IV) (ALCOHOLBIS cohort) or fifth edition (DSM-5)

(GUT2BRAIN cohort). Patients were eligible if they had been drinking until the day of

admission to the detoxification unit or the day before, and if they also did not suffer from

inflammatory bowel disease, other chronic inflammatory diseases (such as rheumatoid arthritis)

or cancer, nor from metabolic disorders such as obesity (BMI > 30 kg/m2), diabetes and

bariatric surgery, or severe cognitive impairment (MMSE < 24). We also excluded subjects who

had taken antibiotics, probiotics, or prebiotics in the 2 months prior to enrolment and those who

were taking non-steroidal anti-inflammatory drugs or glucocorticoids within 1 month of

inclusion. Patients with known cirrhosis or significant liver fibrosis (≥F2) detected by Fibroscan

(>7.6 kPa) on the day of admission were also excluded from the study. No other psychiatric

diseases, and no other addiction (except tobacco) have been diagnosed in these patients.

AUD patients were tested twice, on the day following their admission (T1) and on days 18–19

(T2) corresponding to the last days of the detoxification program. The patients of the

GUT2BRAIN cohort were initially enrolled in a randomized, double-blind, placebo-controlled

study assessing the impact of prebiotic fiber supplementation on the gut-liver-brain axis (54,55).

For this reason, only biological and psychological data obtained at admission (T1), and before

the beginning of the prebiotic/placebo treatment, were considered. The patients of the

ALCOHOLBIS cohort did not take part in any other clinical study during the 3-week

hospitalization stay. Therefore, for this cohort, data obtained at both times of alcohol

withdrawal (T1 and T2) were considered. Thirty-two healthy controls (13 from the

GUT2BRAIN cohort and 19 from the ALCOHOLBIS cohort) matched for age, gender and BMI

with no AUD (Alcohol use disorders test [AUDIT] score <8 in males and <7 in females) were

also recruited using flyers posted in Brussel's public setting (Table S1). The inclusion/exclusion

criteria were the same as for AUD patients except for the alcohol related items. Healthy controls

and AUD patients were not matched for smoking status.

The study was approved by the "Comité d'éthique Hospitalo-facultaire Saint-Luc UCLouvain"

(2017/04JUL/354 and 2014/31dec/614, identification number NCT03803709 at

ClinicalTrials.gov). The study has been carried out in accordance with The Code of Ethics of

the World Medical Association and followed the ethical guidelines set out in the Declaration of

Helsinki. All participants provided written informed consent in compliance with the European

law 2001/20/CE guidelines.

For investigating presence of the potentially neuroactive metabolites in the CNS, we used

metabolomics data from frontal cortex (Broadman area 9) and CSF samples from the Tampere

Sudden Death Study (TSDS) cohort, which have been described in detail elsewhere (10). TSDS

was collected from forensic autopsies done in the area of the Pirkanmaa Hospital District during

2010-2015, a total of 700 subjects. Out of these we identified 97 heavy alcohol users based on

autopsy reports and medical records (diagnosis of alcohol-related diseases: ICD-10 codes

F10.X, G31.2, G62.1, G72.1, I42.6, K70.0-K70.4, K70.9, and K86.0, or signs of heavy alcohol

use in the clinical or laboratory findings, e.g., increased levels of gamma-glutamyl transferase,

mean corpuscular volume, carbohydrate-deficient transferrin). Lack of these findings was

inclusion criteria for the control group (n = 100), most of whom had died due to cardiovascular

diseases (Table S2). Samples were stored at -80 °C until use.

Assessment of psychological symptoms

AUD patients were tested for depression, anxiety and alcohol craving with self-reported

questionnaires: the Beck Depression Inventory [BDI] (56), the State-Trait Anxiety Inventory

(STAI form YA)(57), and the Obsessive-Compulsive Drinking Scale [OCDS](58) (see

supplemental material for details)

Biological sampling

To avoid variation due to fasting state and circadian rhythm, blood samples were collected in

all participants in the morning between 8:00 and 8:30 am after an overnight fasting, at T1 and

T2. Blood was drawn in tubes containing EDTA as an anticoagulant. The samples were centrifuged

at 1000 g for 15 min at 4°C and the plasma was frozen at −80°C until analysis.

Nontargeted metabolomics analysis

Plasma sample preparation and LC-MS measurement were performed as previously detailed

in Klåvus et al (59) (see supplemental material for details).

Statistical analysis

R software version 4.0.3. was used for statistical analyses. Multivariate analyses, namely PCA for dimension reduction and sPLS-DA for group discrimination, were conducted by 'mixOmics' R package v. 6.14.1 (60). For the sPLS-DA model we used a cross-validation (CV) procedure of 10-fold CV repeated 50 times. Univariate analyses were conducted by 'notame' R package v. 0.2.1(59). Significant features were shortlisted using Welch's and paired t-tests. All p-values were corrected using the Benjamini-Hochberg false discovery rate (FDR) to calculate the q-value. For all tests, p and q values < 0.05 were considered statistically significant. Visualizations were created by the previously mentioned R packages and GraphPad Prism v. 8.4.2. Correlation analyses were performed at T1 using R software version 3.6.1. Spearman coefficient was calculated and p-value < 0.05 was considered statistically significant.

REFERENCES

1. Voutilainen T, Kärkkäinen O (2019): Changes in the Human Metabolome Associated With

Alcohol Use: A Review. *Alcohol and Alcoholism* 54: 225–234.

2. Jaremek M, Yu Z, Mangino M, Mittelstrass K, Prehn C, Singmann P, *et al.* (2013):

Alcohol-induced metabolomic differences in humans. *Transl Psychiatry* 3: e276.

3. Würtz P, Cook S, Wang Q, Tiainen M, Tynkkynen T, Kangas AJ, *et al.* (2016): Metabolic

profiling of alcohol consumption in 9778 young adults. *Int J Epidemiol* 45: 1493–

1506.

4. Lehikoinen AI, Kärkkäinen OK, Lehtonen MAS, Auriola SOK, Hanhineva KJ, Heinonen

ST (2018): Alcohol and substance use are associated with altered metabolome in the

first trimester serum samples of pregnant mothers. *Eur J Obstet Gynecol Reprod Biol*

223: 79–84.

5. Kärkkäinen O, Klåvus A, Voutilainen A, Virtanen J, Lehtonen M, Auriola S, *et al.* (2020):

Changes in Circulating Metabolome Precede Alcohol-Related Diseases in Middle-

Aged Men: A Prospective Population-Based Study With a 30-Year Follow-Up.

Alcoholism: Clinical and Experimental Research 44: 2457–2467.

6. Gao X, Zhao A, Zhou M, Lin J, Qiu Y, Su M, Jia W (2011): GC/MS-based urinary

metabolomics reveals systematic differences in metabolism and ethanol response

between Sprague–Dawley and Wistar rats. *Metabolomics* 7: 363–374.

7. Ma T, Li Y, Zhu Y, Jiang S, Cheng C, Peng Z, Xu L (2020): Differential Metabolic

Pathways and Metabolites in a C57BL/6J Mouse Model of Alcoholic Liver Disease.

Med Sci Monit 26: e924602-1-e924602-10.

8. Fernando H, Kondraganti S, Bhopale KK, Volk DE, Neerathilingam M, Kaphalia BS, *et al.*

(2010): ^{1}H and ^{31}P NMR lipidome of ethanol-induced fatty liver. *Alcohol Clin Exp Res*

34: 1937–1947.

9. Zhu X, Huang J, Huang S, Wen Y, Lan X, Wang X, *et al.* (2021): Combining

Metabolomics and Interpretable Machine Learning to Reveal Plasma Metabolic

Profiling and Biological Correlates of Alcohol-Dependent Inpatients: What About

Tryptophan Metabolism Regulation? *Front Mol Biosci* 8: 760669.

10. Kärkkäinen O, Kokla M, Lehtonen M, Auriola S, Martiskainen M, Tiihonen J, *et al.*

(2021): Changes in the metabolic profile of human male postmortem frontal cortex

and cerebrospinal fluid samples associated with heavy alcohol use. *Addiction Biology*

26: e13035.

11. Bar N, Korem T, Weissbrod O, Zeevi D, Rothschild D, Leviatan S, *et al.* (2020): A

reference map of potential determinants for the human serum metabolome. *Nature*

588: 135–140.

12. Leclercq S, Cani PD, Neyrinck AM, Stärkel P, Jamar F, Mikolajczak M, *et al.* (2012):

Role of intestinal permeability and inflammation in the biological and behavioral

control of alcohol-dependent subjects [no. 6]. *Brain Behav Immun* 26: 911–918.

13. Leclercq S, De Saeger C, Delzenne N, de Timary P, Stärkel P (2014): Role of

inflammatory pathways, blood mononuclear cells, and gut-derived bacterial products

in alcohol dependence [no. 9]. *Biol Psychiatry* 76: 725–733.

14. Leclercq S, Matamoros S, Cani PD, Neyrinck AM, Jamar F, Stärkel P, *et al.* (2014):

Intestinal permeability, gut-bacterial dysbiosis, and behavioral markers of alcohol-

dependence severity [no. 42]. *Proc Natl Acad Sci U S A* 111: E4485–E4493.

15. Ahmed H, Leyrolle Q, Koistinen V, Kärkkäinen O, Layé S, Delzenne N, Hanhineva K

(2022): Microbiota-derived metabolites as drivers of gut–brain communication. *Gut*

Microbes 14: 2102878.

16. Chen L, Zhernakova DV, Kurilshikov A, Andreu-Sánchez S, Wang D, Augustijn HE, *et*

al. (2022): Influence of the microbiome, diet and genetics on inter-individual variation

in the human plasma metabolome. *Nat Med* 28: 2333–2343.

17. Guertin KA, Moore SC, Sampson JN, Huang W-Y, Xiao Q, Stolzenberg-Solomon RZ, *et*

al. (2014): Metabolomics in nutritional epidemiology: identifying metabolites

associated with diet and quantifying their potential to uncover diet-disease relations in

populations. *Am J Clin Nutr* 100: 208–217.

18. Zheng Y, Yu B, Alexander D, Steffen LM, Nettleton JA, Boerwinkle E (2014):

Metabolomic patterns and alcohol consumption in African Americans in the

Atherosclerosis Risk in Communities Study. *Am J Clin Nutr* 99: 1470–1478.

19. Hopkins MJ, Sharp R, Macfarlane GT (2001): Age and disease related changes in

intestinal bacterial populations assessed by cell culture, 16S rRNA abundance, and

community cellular fatty acid profiles. *Gut* 48: 198–205.

20. Taormina VM, Unger AL, Schiksnis MR, Torres-Gonzalez M, Kraft J (2020): Branched-

Chain Fatty Acids-An Underexplored Class of Dairy-Derived Fatty Acids. *Nutrients*

12: 2875.

21. Langenau J, Oluwagbemigun K, Brachem C, Lieb W, Giuseppe R di, Artati A, *et al.*

(2020): Blood Metabolomic Profiling Confirms and Identifies Biomarkers of Food

Intake [no. 11]. *Metabolites* 10: 468.

22. Wang Y, Gapstur SM, Carter BD, Hartman TJ, Stevens VL, Gaudet MM, McCullough

ML (2018): Untargeted Metabolomics Identifies Novel Potential Biomarkers of

Habitual Food Intake in a Cross-Sectional Study of Postmenopausal Women. *The*

Journal of Nutrition 148: 932–943.

23. Tanaka M, Budhathoki S, Hirata A, Morita M, Kono S, Adachi M, *et al.* (2013):

Behavioral and clinical correlates of serum bilirubin concentrations in Japanese men

and women. *BMC Endocr Disord* 13: 39.

24. Xu R, He L, Vatsalya V, Ma X, Kim S, Mueller EG, *et al.* (2023): Metabolomics analysis

of urine from patients with alcohol-associated liver disease reveals dysregulated

caffeine metabolism. *American Journal of Physiology-Gastrointestinal and Liver*

Physiology 324: G142–G154.

25. Franco R, Martínez-Pinilla E (2023): One-way or two-way sweet link between

theobromine and depression? *BMC Psychiatry* 23: 411.

26. Thorsell A, Johnson J, Heilig M (2007): Effect of the adenosine A2a receptor antagonist

3,7-dimethyl-propargylxanthine on anxiety-like and depression-like behavior and

alcohol consumption in Wistar Rats. *Alcohol Clin Exp Res* 31: 1302–1307.

27. El-Haggar SM, Eissa MA, Mostafa TM, El-Attar KS, Abdallah MS (2018): The

Phosphodiesterase Inhibitor Pentoxifylline as a Novel Adjunct to Antidepressants in

Major Depressive Disorder Patients: A Proof-of-Concept, Randomized, Double-Blind,

Placebo-Controlled Trial. *Psychother Psychosom* 87: 331–339.

28. Grigsby KB, Mangieri RA, Roberts AJ, Lopez MF, Firsick EJ, Townsley KG, *et al.*

(2023): Preclinical and clinical evidence for suppression of alcohol intake by

apremilast. *J Clin Invest* 133: e159103.

29. Orrú M, Guitart X, Karcz-Kubicha M, Solinas M, Justinova Z, Barodia SK, *et al.* (2013):

Psychostimulant pharmacological profile of paraxanthine, the main metabolite of

caffeine in humans. *Neuropharmacology* 67C: 476–484.

30. García-Blanco T, Dávalos A, Visioli F (2017): Tea, cocoa, coffee, and affective disorders:

vicious or virtuous cycle? *Journal of Affective Disorders* 224: 61–68.

31. Loppi S, Kolosowska N, Kärkkäinen O, Korhonen P, Huuskonen M, Grubman A, *et al.*

(2018): HX600, a synthetic agonist for RXR-Nurr1 heterodimer complex, prevents

ischemia-induced neuronal damage. *Brain Behav Immun* 73: 670–681.

32. Endle H, Horta G, Stutz B, Muthuraman M, Tegeder I, Schreiber Y, *et al.* (2022): AgRP

neurons control feeding behaviour at cortical synapses via peripherally derived

lysophospholipids [no. 6]. *Nat Metab* 4: 683–692.

33. Miletta MC, Iyilikci O, Shanabrough M, Šestan-Peša M, Cammisa A, Zeiss CJ, *et al.*

(2020): AgRP neurons control compulsive exercise and survival in an activity-based

anorexia model [no. 11]. *Nat Metab* 2: 1204–1211.

34. Palmela I, Correia L, Silva RFM, Sasaki H, Kim KS, Brites D, Brito MA (2015):

Hydrophilic bile acids protect human blood-brain barrier endothelial cells from

disruption by unconjugated bilirubin: an in vitro study. *Frontiers in Neuroscience* 9.

Retrieved July 20, 2023, from

https://www.frontiersin.org/articles/10.3389/fnins.2015.00080

35. Quinn M, McMillin M, Galindo C, Frampton G, Pae HY, DeMorrow S (2014): Bile acids

permeabilize the blood brain barrier after bile duct ligation in rats via Rac1-dependent

mechanisms. *Digestive and Liver Disease* 46: 527–534.

36. MahmoudianDehkordi S, Arnold M, Nho K, Ahmad S, Jia W, Xie G, *et al.* (2019):

Altered bile acid profile associates with cognitive impairment in Alzheimer's

disease—An emerging role for gut microbiome. *Alzheimer's & Dementia* 15: 76–92.

37. Brunengraber H, Roe CR (2006): Anaplerotic molecules: Current and future. *Journal of*

Inherited Metabolic Disease 29: 327–331.

38. Mochel F, DeLonlay P, Touati G, Brunengraber H, Kinman RP, Rabier D, *et al.* (2005):

Pyruvate carboxylase deficiency: clinical and biochemical response to anaplerotic diet

therapy. *Mol Genet Metab* 84: 305–312.

39. Chu C, Murdock MH, Jing D, Won TH, Chung H, Kressel AM, *et al.* (2019): The

microbiota regulate neuronal function and fear extinction learning [no. 7779]. *Nature*

574: 543–548.

40. Chen S-J, Lin C-H (2022): Gut microenvironmental changes as a potential trigger in

Parkinson's disease through the gut–brain axis [no. 1]. *J Biomed Sci* 29: 1–18.

41. Needham BD, Funabashi M, Adame MD, Wang Z, Boktor JC, Haney J, *et al.* (2022): A

gut-derived metabolite alters brain activity and anxiety behaviour in mice. *Nature* 602:

647–653.

42. Hsiao EY, McBride SW, Hsien S, Sharon G, Hyde ER, McCue T, *et al.* (2013):

Microbiota modulate behavioral and physiological abnormalities associated with

neurodevelopmental disorders [no. 7]. *Cell* 155: 1451–1463.

43. Needham BD, Adame MD, Serena G, Rose DR, Preston GM, Conrad MC, *et al.* (2021):

Plasma and Fecal Metabolite Profiles in Autism Spectrum Disorder. *Biol Psychiatry*

89: 451–462.

44. Lees HJ, Swann JR, Wilson ID, Nicholson JK, Holmes E (2013): Hippurate: The Natural

History of a Mammalian–Microbial Cometabolite. *J Proteome Res* 12: 1527–1546.

45. Pruss KM, Chen H, Liu Y, Van Treuren W, Higginbottom SK, Jarman JB, *et al.* (2023):

Host-microbe co-metabolism via MCAD generates circulating metabolites including

hippuric acid [no. 1]. *Nat Commun* 14: 512.

46. Wikoff WR, Anfora AT, Liu J, Schultz PG, Lesley SA, Peters EC, Siuzdak G (2009):

Metabolomics analysis reveals large effects of gut microflora on mammalian blood

metabolites. *ProcNatlAcadSciUSA* 106: 3698–3703.

47. Harada S, Takebayashi T, Kurihara A, Akiyama M, Suzuki A, Hatakeyama Y, *et al.*

(2016): Metabolomic profiling reveals novel biomarkers of alcohol intake and alcohol-

induced liver injury in community-dwelling men. *Environ Health Prev Med* 21: 18–
26.

48. Kärkkäinen O, Tolmunen T, Kivimäki P, Kurkinen K, Ali-Sisto T, Mäntyselkä P, *et al.*
(2024): Alcohol use associated alterations in the circulating metabolite profile in the
general population and in individuals with major depressive disorder. *Alcohol*.
https://doi.org/10.1016/j.alcohol.2024.01.005

49. van der Spek A, Stewart ID, Kühnel B, Pietzner M, Alshehri T, Gauß F, *et al.* (2023):
Circulating metabolites modulated by diet are associated with depression. *Mol*
Psychiatry 1–14.

50. Sobell LC, Sobell MB (1992): Timeline Follow-Back. In: Litten RZ, Allen JP, editors.
Measuring Alcohol Consumption: Psychosocial and Biochemical Methods. Totowa,
NJ: Humana Press, pp 41–72.

51. Hsu P-C, Lan RS, Brasky TM, Marian C, Cheema AK, Ressom HW, *et al.* (2017):
Metabolomic profiles of current cigarette smokers. *Mol Carcinog* 56: 594–606.

52. Barve S, Chen S-Y, Kirpich I, Watson WH, McClain C (2017): Development, Prevention,
and Treatment of Alcohol-Induced Organ Injury: The Role of Nutrition. *Alcohol Res*
38: 289–302.

53. Harrison ELR, Desai RA, McKee SA (2008): Nondaily smoking and alcohol use,
hazardous drinking, and alcohol diagnoses among young adults: findings from the
NESARC. *Alcohol Clin Exp Res* 32: 2081–2087.

54. Amadieu C, Coste V, Neyrinck AM, Thijssen V, Leyrolle Q, Bindels LB, *et al.* (2022):
Restoring an adequate dietary fiber intake by inulin supplementation: a pilot study
showing an impact on gut microbiota and sociability in alcohol use disorder patients.
Gut Microbes 14: 2007042.

55. Amadieu C, Maccioni L, Leclercq S, Neyrinck AM, Delzenne NM, de Timary P, Stärkel

P (2022): Liver alterations are not improved by inulin supplementation in alcohol use

disorder patients during alcohol withdrawal: A pilot randomized, double-blind,

placebo-controlled study. *EBioMedicine* 80: 104033.

56. Beck AT, Steer RA, Brown BK (1996): *Beck Depression Inventory (2nd Ed.)*, vol.

Corporation TP. San Antonio.

57. Spielberger CD, Gorsuch RL, Lushene R, Vagg PR, Jacobs GA (1983): *Manual for the*

State-Trait Anxiety Inventory, vol. Consulting Psychologist Press I. Palo Alto.

58. Anton RF, Moak DH, Latham P (1995): The Obsessive Compulsive Drinking Scale: a

self-rated instrument for the quantification of thoughts about alcohol and drinking

behavior. *Alcohol ClinExpRes* 19: 92–99.

59. Klåvus A, Kokla M, Noerman S, Koistinen VM, Tuomainen M, Zarei I, *et al.* (2020):

"Notame": Workflow for Non-Targeted LC–MS Metabolic Profiling [no. 4].

Metabolites 10: 135.

60. Rohart F, Gautier B, Singh A, Cao K-AL (2017): mixOmics: An R package for 'omics

feature selection and multiple data integration [no. 11]. *PLOS Computational Biology*

13: e1005752.

FUNDING

Gut2Behave project was initiated from ERA-NET NEURON network (Joint Transnational

Call 2019) and was financed by Academy of Finland, French National Research Agency

(ANR-19-NEUR-0003-03) and the Fonds de la Recherche Scientifique (FRS-FNRS) [PINT-

MULTI R.8013.19, Belgium). Metabolomics analysis of the TSDS samples was supported by

grant from the Finnish Foundation

ACKNOWLEDGMENTS AND DISCLOSURES

SL is a Research Associate of the Fonds de la Recherche Scientifique – FNRS. Collection of TSDS samples was supported by European Union 7th Framework Program (grant number 201668 for AtheroRemo Project, State Researh Funding for Tampere University Hospital, Finnish Foundation for Cardiovascular Research). PDT received funding from Fondation Saint Luc. NMD is a recipient of grants from the Fonds de la Recherche Scientifique (FRS-FNRS) [PDR T.0068.19], and from the Fédération Wallonie-Bruxelles (Action de Recherche Concertée ARC18- 23/092).

OK and KH are founders of Afekta Technologies Ltd. The other authors report no financial interests or potential conflicts of interest.

AUTHOR CONTRIBUTIONS

Conceptualized and designed the study: SL, CA, PS, PdT, NMD

Performed participant visits: SL, CA, GP, MP

Collected and provided the TSDS post-mortem samples: EK, PJK

Performed data analysis, interpreted the data and drafted the manuscript: SL, HA, CA, QL, PdT, SLa, AMN, OK, KH, NMD

Performed the analysis of the metabolomics data: HA, VK, OK, KH

Contributed to ethics application; SL, PS, PdT, NMD

All authors approved the final version of the manuscript.

664 **Table 1**. Clinical features of the study participants

	ALCOHOLBIS cohort	GUT2BRAIN cohort	All	p-value
Number of subjects	48	48	96	
Age	46 ± 10	48 ± 9	47 ± 10	0.24
Gender				
Men, n (%)	34 (71%)	30 (62.5%)	64 (67%)	0.39[$]
Women, n (%)	14 (29%)	18 (37.5%)	32 (33%)	
Smoking status				
Active smoker (%)	77	79	78	0.83[$]
Non-smoker (%)	23	21	22	
Alcohol history				
Alcohol consumption (g/day)	151 ± 112	139 ± 73	145 ± 94	0.54
Duration of drinking habits (years)	17 ± 10	16 ± 11	16 ± 11	0.76
Number of withdrawal cures	1 ± 2	2 ± 2	1.8 ± 2.3	0.015*
AUDIT score	31 ± 7			
DSM5 score		8 ± 2		
Depression				
Score at T1	23 ± 11	26 ± 12	25 ± 12	0.26
Anxiety				
Score at T1	44 ± 11	46 ± 15	45 ± 13	0.27
Alcohol craving				
Total score at T1	20 ± 7	25 ± 6	22 ± 7	0.00*
Obsession score at T1	9 ± 5	11 ± 4	10 ± 4	0.26
Compulsion score at T1	11 ± 3	14 ± 3	13 ± 3	0.00*

665 Results are means $\pm$ standard deviations. Independent t-tests to compare ALCOHOLBIS *versus*

666 GUT2BRAIN cohorts. [$]chi-square test for categorical variables; *p < 0.05

Figure legends

Figure 1. Chronic alcohol intake alters the plasma metabolome compared to healthy controls. **a** Principal component analysis score plot of the plasma metabolomic features between healthy controls and persons with AUD at T1. QC samples are colored in orange. **b** Top 50 annotated metabolites ranked on their variable importance in projections (VIP) scores in the sPLS-DA model. **c** Volcano plot depicting the effect size (Cohen's D) and -Log$_{10}$ transformed q values derived from Welch's t-test analysis of the metabolomic features different between healthy controls and persons with AUD at T1. **d** Annotated metabolites having significant association (Spearman $p<0.05$) with alcohol intake (g/day) in persons with AUD at T1. Circle size refers to the level of significance, blue gradient color to the strength of negative while red to the strength of positive correlation coefficients. 3-CMPFP 3-carboxy-4-methyl-5-propyl-2-furanpropionic acid; AC acylcarnitine; CLA conjugated linoleic acid; FA fatty acid; LPC lysophosphatidylcholine; LPE lysophosphatidylethanolamine; PC phosphatidylcholine; PE phosphatidylethanolamine; SM sphingomyelin.

Figure 2. 3-week alcohol withdrawal shapes the plasma metabolome. **a** Scores plot of the sparse partial least square discriminant analysis (sPLS-DA) of the plasma metabolomes at the start (T1) and end (T2) of the alcohol withdrawal in persons with AUD. **b** Top 50 annotated metabolites having a variable importance in projection (VIP) score of > 1.5 in the sPLS-DA model. **c** Lollipop plot of the effect size (Cohen's D) and -Log$_{10}$ transformed q values of the altered annotated metabolites between AUD T1 and healthy controls as well as AUD T1 and AUD T2. Circle size refers to the level of significance, grey color to the comparison between controls and AUD T1, orange color to relative increase while violet to relative decrease towards the end of alcohol withdrawal (T2). AC acylcarnitine; FA fatty acid; LPC lysophosphatidylcholine; LPE lysophosphatidylethanolamine; PC phosphatidylcholine.

Figure 3. Plasma metabolites associated with psychological symptom scores. Heatmap of the annotated metabolites having significant ($p<0.05$) Spearman correlation with one or more psychological symptom score of obsession, compulsion, alcohol craving, anxiety state or depression. Circle size refers to the level of significance, blue gradient color to the strength of negative while red to the strength of positive correlation coefficients. AC acylcarnitine; FA fatty acid; GM gut microbiota; LPC lysophosphatidylcholine; LPE lysophosphatidylethanolamine; PC phosphatidylcholine.

Figure 4. Overlapping metabolites within the plasma and brain metabolome. **a** Venn diagram of the annotated plasma metabolites that are also present in the frontal cortex and cerebrospinal fluid metabolome. **b** Significantly altered ($p<0.05$) frontal cortex metabolites with a corresponding plasma metabolite showing an association with one or more psychological symptom score. **c** Significantly altered ($p<0.05$) cerebrospinal fluid metabolite with a corresponding plasma metabolite showing an association with one or more psychological symptom score. Data expressed as mean ± SD with individual values shown. Statistical values derived from Welch's t-test comparing metabolomic features between control and alcohol groups.

Plasma annotated metabolites significantly (p<0.05) correlated with psychological symptom scores

Figure showing heatmap of Blood metabolic features vs Psychological symptom scores.

Row labels (top to bottom):
Creatinine, Epsilon–caprolactam, Kynurenine isomer 1, Kynurenine isomer 2, Tryptophyl–Phenylalanine

Acylcarnitines: AC 05:0, AC 08:1, AC 09:0, AC 10:3, AC 12:0, AC 14:0, Acetylcarnitine

Bile acids: 7a–hydroxy–3–oxo–4–cholestenoic acid, Glycochenodeoxycholic acid isomer 1, Glycochenodeoxycholic acid isomer 2, Glycocholic acid, Glycohyodeoxycholic acid, Tauroursodeoxycholic acid, Bile acid C26H43NO7S, Bile acid C26H43NO8S isomer 1, Bile acid C26H43NO8S isomer 3

Biliverdin isomer 1, 1–methyl–pyridone–carboxamide, 4–Pyridoxic acid, C11H18O4, C15H12O2, C8H15NO2, C9H15NO3, Cotinine

Drugs: Citalopram/Escitalopram, Diazepam, Nordiazepam, Oxaprozin, Pantoprazole, Temazepam, Trazodone, Venlafaxine

Fatty acids: FA 16:0, FA 16:1, FA 16:1;O, FA 18:0, FA 18:0;O, FA 18:3, 3–Hydroxyvaleric acid, Fatty acid_C27H50O2

GM-related: 3–Indoleacetic acid, 4–Ethylphenylsulfate, Hippuric acid, Phenylsulfate_C21H34O8S2, Pyrocatechol sulfate

LPCs: LPC 0:0/14:0, LPC 0:0/16:0, LPC 0:0/16:1, LPC 0:0/17:1, LPC 0:0/18:1, LPC 0:0/18:2, LPC 0:0/20:3, LPC 14:0/0:0, LPC 15:0_0:0, LPC 16:0/0:0, LPC 16:1/0:0, LPC 17:1/0:0, LPC 18:1/0:0, LPC 18:2/0:0, LPC 18:3_0:0, LPC 20:1_0:0, LPC 20:2_0:0, LPC 20:3/0:0, LPC 22:4_0:0, LPC 22:5, LPC O–18:0, LPC P–16:0/0:0, Lysophosphatidylcholine_C24H50NO8P

LPEs: LPE 0:0/18:2, LPE 0:0/22:6, LPE 16:0, LPE 18:1, LPE 18:2/0:0, LPE 18:3, LPE 20:3, LPE 20:4, LPE 20:5, LPE 22:5

PCs: PC 16:0/16:0, PC 16:0_18:1, PC 16:0_18:3, PC 16:1_18:2, PC 18:0_18:1, PC 18:1_14:0 and 16:0_16:1, PC 18:2_15:0, PC 30:1, PC 35:2 isomer 1, PC 35:2 isomer 2, PC O–38:6

Androsterone sulfate isomer 1, Dehydroandrosterone, Dehydroepiandrosterone sulfate, Urea

Xanthines: Hypoxanthine, Paraxanthine, Theobromine, Theophylline

Mannose/Fructose

Column labels (x-axis, Psychological symptom scores): Obsession (OCDS), Compulsion (OCDS), Craving (OCDS–Total Score), Anxiety state (STAI), Depression (BDI)

Y-axis: Blood metabolic features

−log10(p-value): 2, 3, 4

Correlation: 0.25, 0.00, −0.25

a

b

c