Near telomere-to-telomere genome assemblies of two *Chlorella* species unveil the composition and evolution of centromeres in green algae

Bo Wang[1,2†], Yanyan Jia[1,2†], Ningxin Dang[1,3†], Jie Yu[4], Stephen J. Bush[1], Shenghan Gao[1], Wenxi He[5], Sirui Wang[5], Hongtao Guo[6], Xiaofei Yang[6], Weimin Ma[4*], Kai Ye[1,2,3,5,7*]

[1]School of Automation Science and Engineering, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, China.

[2]MOE Key Lab for Intelligent Networks & Networks Security, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, China.

[3]Genome Institute, the First Affiliated Hospital of Xi'an Jiaotong University, Xi'an, China.

[4]College of Life Sciences, Shanghai Normal University, Shanghai, China

[5]School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, China.

[6]School of Computer Science and Technology, Faculty of Electronic and Information Engineering, Xi'an Jiaotong University, Xi'an, China.

[7]Faculty of Science, Leiden University, Leiden, The Netherlands

[†] These authors contributed equally.

* Correspondence author: kaiye@xjtu.edu.cn (K. Ye), wma@shnu.edu.cn (W. Ma)

Acknowledgments

This study was supported by the National Key R&D Program of China (2022YFC3400300) and the National Natural Science Foundation of China (32200510; 32070663; 32170257; 32125009; 62172325).

Conflict of interest

The authors declare that they have no conflicts of interest.

Abstract

Centromeres play a crucial and conserved role in cell division, although their composition and evolutionary history in green algae, the evolutionary ancestors of land plants, remains largely unknown. In this study, we constructed near telomere-to-telomere (T2T) assemblies for two Trebouxiophyceae species, *Chlorella sorokiniana* NS4-2 and *C. pyrenoidosa* DBH, with chromosome numbers of 12 and 13, and genome sizes of 58.11 Mb and 53.41 Mb, respectively. We identified and validated centromere sequences using CENH3 ChIP-seq and found that, similar to humans and higher plants, the centromere CENH3 signals of green algae display a pattern of hypomethylation. Interestingly, the centromeres of both species largely comprised transposable elements (TEs), although their composition showed significant differences. In *Chlorella* species, centromeres primarily consist of multiple elements, such as LINE/L1, LTR/Copia, LINE/RTEX, and Unknown families. However, in early-evolved species such as *Chamydomonas reinhardtii*, *Coccomyxa subellipsoidea*, and *Chromochloris zofingiensis*, centromere composition has evolved towards optimization with a single-element composition. Through phylogenomic and centromeric compositions analysis, We found that the centromeres of the most recent common ancestor of green algae are composed of two transposable element families, LINE/L1 and LTR/Copia. Moreover, we observed significant differences in the composition and structure of centromeres among chromosomes with strong collinearity within the *Chlorella* genus, suggesting that centromeres in green algae evolve more rapidly than non-centromeric regions. This study not only provides high-quality genome data for comparative genomics of green algae but gives insight into the composition and evolutionary history of centromeres in early plants, laying an important foundation for further research on their evolution.

Keywords

Genome hybrid assembly; Telomere-to-telomere; Centromere assembly; Transposable elements; Centromere evolution.

Introduction

Centromeres play a critical role in ensuring the accurate segregation of chromosomes during both mitosis and meiosis (Talbert and Henikoff, 2022). They are specialized, constricted regions on chromosomes responsible for chromatid cohesion and correct separation during cell division. Centromeres serve as the assembly and attachment points for the kinetochore, a large protein complex that facilitates the binding of spindle microtubules to chromatids, enabling their movement towards the poles (McAinsh and Marston, 2022). The 'centromere paradox' refers to the apparent contradiction between their essential role in chromosome segregation, which is conserved across all eukaryotes, and the significant variability in their organization and DNA composition, even within closely related species (Henikoff et al. 2001; Melters et al. 2013). This variability highlights the rapid evolution of centromeres.

The high prevalence of satellite repeats within centromere sequences also presents a considerable challenge in genome assembly for this region, frequently leading to the substantial loss of sequence information (Miga, 2020). This obstacle poses a significant impediment to the comprehensive investigation of centromere functionality. However, with advances in sequencing technology, in particular the PacBio high-fidelity (HiFi) and Oxford Nanopore Technologies (ONT) ultra-long methods, in conjunction with advances in assembly algorithms (Cheng et al. 2021; Rautiainen et al. 2023), significant progress has been made towards the complete assembly of centromeres in many model species, including but not limited to human (Nurk et al. 2022), *Arabidopsis thaliana* (Naish et al. 2021; Wang et al. 2022), rice (Song et al. 2021), and maize (Chen et al. 2023). Notably, both human and higher plant centromeres primarily comprise numerous satellite sequences, organized into higher-order repeat arrays (Miga, 2020; Gao et al. 2023). Interestingly, the centromeres of the oldest domesticated wheat (*Triticum monococcum*) were found to predominantly feature transposable elements (TEs) (Ahmed et al. 2023). In fact, a few years ago, scholars proposed that the plant centromeric satellites might result from the sequences of TEs specifically targeting the centromere (Presting 2018; Huang et al.

2021). Through high-quality genome assembly and putative centromere sequence analysis of the moss *Physcomitrium patens* and the model green alga *Chlamydomonas reinhardtii*, it was discovered that their centromeres are indeed composed of TEs (Craig et al. 2021; Bi et al. 2024). Whether centromeres in other green algae, more distantly related to *C. reinhardtii*, are also composed of TEs, and if so, whether the constituent elements are similar, as well as the evolutionary transitions between them, remains unclear.

In this study, we sequenced and assembled two *Chlorella* species to the T2T level, characterising and experimentally validating their centromeres. We found that the centromeres of the green algae species selected in this study are all composed of transposable elements (TEs), albeit with significant differences in composition. *Chlamydomonas reinhardtii* and *Coccomyxa subellipsoidea* predominantly feature elements from the LINE/L1 family, while *Chromochloris zofingiensis* is primarily composed of elements from the LTR/Copia family. In contrast, species within the *Chlorella* genus exhibit a more diverse centromere composition, with major constituents including families such as LTR/Copia, LINE/L1, and LINE/RTEX. This suggests that these species are in the process of optimizing one element to become the predominant constituent during evolution. Overall, our results give insight into the centromere evolution of green algae, which may be generalised to early plants, as well as providing high-quality genomic resources to facilitate future comparative analyses.

Results

Telomere-to-telomere assemblies of two *Chlorella* species

To investigate centromere evolution within the context of green algae, we assembled the genomes of two *Chlorella* species, *C. sorokiniana* NS4-2 and *C. pyrenoidosa* DBH (basionym: *Auxenochlorella pyrenoidosa*), both of which belong to the Trebouxiophyceae class, relatively distant from the Chlorophyceae class of *Chlamydomonas reinhardtii* in terms of phylogenetic distance (divergence time: 827 MYA, http://timetree.org/) (**Figure S1**) (Blanc et al. 2010). Their morphology is similar, but they vary in their chromosome numbers, with NS4-2 having 12

chromosomes and DBH having 13 (**Figure 1A, Figure 1B, Figure S2A and Figure S2B**). The genomes of NS4-2 and DBH were sequenced using HiFi reads with high coverage, namely 438× and 308×, respectively (**Table S1**). We assembled the HiFi reads (**Figure S3**) to produce a set of 16 contigs. These contigs had a minimum length of 141,360 bp and were selected based on the criterion of length exceeding 100 Kb. Following this, we generated a final set of 13 contigs by excluding the genome of the putative alga-benefitting commensal bacterium *Arthrobacter* sp. (Windler et al. 2022) as well as the fragments of rDNA and plastid DNA (21 Kb~107 Kb). Among these contigs, 11 contained the plant-type telomeric repeat unit (CCCTAAA/TTTAGGG) at both ends, signifying their completeness. Additionally, two contigs were found to have rDNA units at one end, as observed through collinearity analysis in comparison with the closely related species *Chlorella vulgaris* which was scaffold by optical mapping (Cecchin et al. 2019) (**Figure S4A**). To consolidate the rDNA ends originating from the same chromosome, we merged the two contigs and inserted 100 Ns as spacers. As a result of this process, the NS4-2 genome now represents a T2T assembly, comprised of 12 chromosomes. This conclusion is further supported by karyotyping results (**Figure 1B**) and Hi-C contact maps (**Figure 1C**). In the case of DBH, we initially assembled the HiFi reads using hifiasm (Cheng et al. 2021) (**Figure S3B**), which generated 38 contigs, each exceeding 100 Kb in length. However, upon closer examination, we discovered that only three of these contigs contained the telomeric repeat unit at both ends. This finding suggests that achieving a T2T-level assembly for DBH is more challenging compared to NS4-2 (**Figure S3A**). To overcome this challenge and obtain a T2T-level assembly for DBH, we generated ONT ultra-long reads (with a read length N50 of 59 Kb) at a high coverage of 368× (**Table S1**). We used *NextDenovo* (Hu et al. 2023) to assemble the ONT long reads, which initially yielded a set of 14 contigs. Out of the total contigs obtained, 10 of them exhibited the presence of telomeric repeat units at both ends, indicating complete resolution of the telomere regions. However, two contigs only had rDNA units at one end, suggesting unresolved telomere regions at the other. Additionally, the remaining two contigs contained the telomeric repeat unit on one end, indicating

that the telomere regions of the other end were not fully resolved. To address this issue and achieve a more comprehensive assembly, we utilized the HiFi contigs to replace the non-resolved ONT contigs (**Figure S3B**). By incorporating these updated HiFi contigs, we successfully achieved a gap-free assembly of 14 contigs. This updated assembly provides a more complete representation of the genome, with improved resolution and continuity. In a similar vein, for DBH, the collinearity analysis also revealed two contigs with rDNA units potentially originating from the same chromosome (**Figure S4B**). Following a similar approach to NS4-2, the rDNA ends were merged, and 100 Ns were introduced as spacers. This led to the generation of a T2T assembly with 13 chromosomes, again supported by karyotyping results (**Figure S2B**) and Hi-C contact maps (**Figure S2C**). The evaluation of genome completeness using Benchmarking Universal Single-Copy Orthologs (BUSCO) revealed that both genomes exhibited high genome completeness, with NS4-2 reaching 98.3% and DBH achieving 99.0%. The assembly of these two genomes represents a significant improvement compared to the existing genomes from the same species (**Table 1**). NS4-2 and DBH were predicted to have 12,691 and 11,327 protein coding genes, respectively. NS-2 and DBH exhibit a significant collinearity relationship, accompanied by chromosomal rearrangements (**Figure 1D**). All HiFi reads greater than 15 Kb were selected and used to perform mitochondrial and chloroplast genome assembly in Unicycler v. 0.5.0 (software optimized for circular genome assemblies) with default parameters (Wick et al. 2017). The chloroplast genome sizes of NS4-2 and DBH are 109,738 bp and 107,436 bp, respectively; while the mitochondrial genome sizes are 53,629 bp and 69,538 bp, respectively (GenBank accession: OP311589-OP311592).

Putative centromere identification and CENH3 ChIP-seq validation

Recent advancements in sequencing technologies have shed light on the crucial role of TEs in influencing the characteristics of centromeric sequences among green algal species (Craig et al. 2021; Payne et al. 2023). Notably, studies on species such as *Chlamydomonas reinhardtii* CC-4532 (v. 6.1) (Craig et al. 2023) and *Coccomyxa subellipsoidea* C-169 (v. 3.0) (Blanc et al. 2012) have revealed the prevalence of

LINE/L1 elements within their genomes, as well as LTR/Copia elements in *Chromochloris zofingiensis* (Roth et al. 2017), further emphasizing the impact of TEs in shaping these organism's centromeric regions. Upon examining the centromere regions of these species, a striking abundance of TEs becomes evident. These TEs form a unique and discernible structure when performing dot plots for self-alignment analysis (**Figure S5**). Notably, within this structure, specific clusters stand out, such as the *Zepp*-like cluster in *C. reinhardtii* and the *Zepp* cluster in *C. subellipsoidea* (**Figure S5**). The presence of these distinct TE clusters further underscores the significance of TEs in shaping the centromeric regions of these species. Leveraging this distinctive characteristic, we conducted dot plot analyses on a per-chromosome basis to pinpoint the potential centromeric regions within the genomes of NS4-2 (**Figure S6**) and DBH (**Figure S7**). In the case of NS4-2, we successfully validated putative centromeres for all 12 chromosomes using VerityMap (Mikheenko et al. 2020) (**Figure S8**), with lengths ranging from 28,465 bp to 104,090 bp (average length: 60 Kb) (**Table S2**). Subsequently, we employed centromeric Histone H3 chromatin immunoprecipitation followed by sequencing technology (CENH3 ChIP-seq) to validate the predicted potential centromere regions and found them to be consistent with experimental verification (**Figure 2 and Figure S9**). We then proceeded to use the dot plot method (**Figure S10**) for potential centromere prediction in DBH species. For DBH, we validated putative centromeres for all 13 chromosomes using VerityMap (**Figure S11**), with lengths ranging from 19,117 bp to 65,960 bp (average length: 39 Kb) (**Table S3**). Insights from circular plots indicate that the two species exhibit a strong degree of collinearity and the putative centromeres of NS4-2 and DBH are composed of TE elements (**Figure 1D**). This was especially apparent for NS4-2, in which the centromere regions were significantly enriched with LTR/Copia elements (**Figure 2 and Figure 1D**).

Centromere methylation analysis in two *Chlorella* species

In higher plants, such as *Arabidopsis thaliana* and *Zea mays*, satellite repeats that are associated with chromatin containing CENH3 exhibit lower levels of methylation when compared to the non-CENH3 centromeric regions (Zhang et al. 2008; Wang et

al. 2022). These findings imply that DNA methylation is a significant factor in the epigenetic demarcation of centromeric chromatin. It is currently unclear whether centromeres in green algae, primarily composed of transposable elements, exhibit similar methylation patterns. We utilized ccsmeth (Ni et al. 2023), a deep learning-based circular consensus sequencing (CCS) sequencing methylation calling software, to analyze the centromere methylation in the NS4-2 genome (**Figure 3**). Interestingly, we found that the centromeric region of chromosome 1 in NS4-2 contains hypomethylated intervals (**Figure 3A**). Subsequently, we further zoomed in on the regions 50 Kb ups- and downstream of the centromere and observed a negative correlation between hypomethylated regions and ChIP signals (**Figure 3A and Figure S12**). We divided the genome into four categories: centromeric regions, non-centromeric regions, ChIP peak regions in centromeric regions, and non-ChIP peak regions in centromeric regions, and conducted a statistical analysis of methylation frequencies. Our analysis revealed that centromeric regions exhibited significantly higher methylation levels compared to non-centromeric regions, primarily attributable to the low methylation status in ChIP peak regions (**Figure 3B**). For DBH, we employed HiFi and ONT sequencing to call methylation, and similarly identified regions of low methylation within the centromeric regions (**Figure S13**). We found that the methylation resolution of HiFi data calling is higher than that of ONT in the TE-rich centromeres (**Figure S13**). We observed the same phenomenon of centromeric low methylation as in the NS4-2 species (**Figure 3C**), indicating that despite the absence of highly tandemly repeated satellite sequences in green algal centromeres, the low methylation in ChIP signal regions resembles that of higher plants. These results suggest that despite rapid evolution, centromeric sequences remain evolutionarily conserved in terms of epigenetic inheritance.

Centromere size and composition in green algae

We selected six high-quality assembled green algal genomes for the identification and compositional analysis of potential centromeres, including *Chlamydomonas reinhardtii* CC-4532 (v. 6.1) (Craig et al. 2023), *Chromochloris zofingiensis* (Rotha et al. 2017), *Coccomyxa subellipsoidea* C-169 (v. 3.0) (Blanc et al. 2012), *Chlorella* sp.

SLA-04 (Goemann et al. 2023), and the two *Chlorella* species assembled in this study

(**Figure 4A**). Additionally, we used the red alga *Cyanidioschyzon merolae* (short low-

GC centromeric regions) (Nozaki et al. 2007; Kanesaki et al. 2015) as an outgroup for

the phylogenetic tree of the lineage (**Figure 4A**). We identified the potential

centromere positions of the above-green species, including *C. reinhardtii, C.*

zofingiensis, C. subellipsoidea and SLA-04, using dot plot analysis and references

from the literature (**Table S4−S7**). We utilized RepeatMasker to annotate the

centromere composition of the selected species based on TE-algae library. The

potential centromeric regions of *C. reinhardtii, C. zofingiensis, C. subellipsoidea*, and

SLA-04 span approximately from 96 Kb to 320 Kb, 10Kb to 45 Kb, 12 Kb to 36 Kb,

and 14 Kb to 118 Kb in length, respectively (**Table S4−S7**), with average sizes of

about 200 Kb, 27 Kb, 23 Kb, and 52 Kb. The model green algae has the largest

genome, measuring approximately 112 Mb, and also possesses the longest average

centromere length. We found a positive correlation between centromere size and

genome size in green algae, with a Pearson correlation coefficient of 0.87 (**Figure 4B**).

This fills a gap in the literature by providing data on green algae, which was

previously lacking (Plačková et al. 2021). We provided a detailed annotation of the

centromere composition in the selected green algal species (**Table S8-S13**), sorted

based on the top five of element's proportion within centromeric regions (**Figure 4C−**

H). In the centromere sequences of model green algae (**Figure 4C and Table S14**)

and *C. subellipsoidea* (**Figure 4E and Table S15**), we observed a predominant

presence of ZeppL-1_cRei (LINE/L1) elements in *C. reinhardtii* and Zepp (LINE/L1)

elements in *C. subellipsoidea*, consistent with previous reports (Craig et al. 2021).

Nearly all of these LINE/L1 elements are distributed in the centromere region,

indicating they are centromere-specific elements (**Table S8 and Table S10**). It is

noteworthy that although the species *C. zofingiensis* demonstrates a sister relationship

with *C. reinhardtii* in phylogenetic analysis, the centromeric composition of *C.*

zofingiensis is primarily comprised of the LTR/Copia family (**Figure 4D and Table**

S16), in stark contrast to *C. reinhardtii* (**Figure 4C**). In species SLA-04, the

centromeric component predominantly consists of Copia-2_Chlorella_SLA04

(LTR/Copia), followed by L1-5_Chlorella_SLA04 (LINE/L1) and RTE-X-

1_Chlorella_SLA04 (LINE/RTE-X), accounting for 37%, 22%, and 11% respectively

(**Table S11 and Table S17**). In species NS4-2, the centromeric component is

primarily composed of Copia-1_Chlorella_NS (LTR/Copia) and Unknown-

1_Chlorella_NS (Unknown), accounting for 50% and 21% respectively (**Table S12**

and Table S18). In the species DBH, the predominant centromeric component is

Unknown-2_Chlorella_DBH (Unknown), followed by RTE-X-1_Chlorella_DBH

(LINE/RTE-X), RTE-X-2_Chlorella_DBH (LINE/RTE-X), and Copia-

1_Chlorella_DBH (LTR/Copia), accounting for 25%, 20%, 19%, and 17%

respectively (**Table S13 and Table S19**). The elements mentioned above in species of

the *Chlorella* genus are all centromere-specific components (**Figure 4F-H**).

Centromere evolution in green algae

We systematically correlate the primary constituents of centromeres across diverse

species onto the phylogenomic tree framework (**Figure 5A**). This strategic approach

enables us to scrutinize the evolutionary trajectories of centromeric transposon

sequences among distinct taxa. In our investigation, we have identified the presence

of LINE/L1 and LTR/Copia elements within species belonging to both Chlorophyceae

and Trebouxiophyceae, two major classes within the green algae lineage (**Figure 5A**).

This finding suggests that these elements are encompassed within the centromeric

sequences of the most recent common ancestor (MCRA) of green algae. An intriguing

observation is the exclusive presence of LINE/RTEX and Unknown family elements

within the centromeric composition of species belonging solely to the Chlorella genus.

This suggests a unique formation of the most recent common ancestor (MCRA)

specific to Chlorella, underscoring distinct evolutionary dynamics within this genus

(**Figure 5A**). Through an extensive evolutionary timeline, the Chlorophyceae-

Chlorella divergence is estimated to have occurred approximately 827 million years

ago (MYA), while the Coccomyxa-Chlorella divergence is estimated to have occurred

around 544 MYA (source: http://www.timetree.org/). During this evolutionary

journey, species such as *C. reinhardtii*, *C. zofingiensis*, and *C. subellipsoidea* have

independently evolved centromere-specific transposable elements (TEs) (**Figure 5A**).

However, within the *Chlorella* genus, multiple TE elements still persist within

centromeric sequences (**Figure 5A**), suggesting an ongoing process of optimization,

potentially towards establishing a predominant TE element as the primary constituent

of their centromeres. For instance, elements like NS4-2 and SLA-04 may be refining

LTR/Copia elements into a singular composite component, while DBH could be

optimizing LINE/RTEX into a singular centromeric component (**Figure 5A**). We are

conducting comparative genomic analysis of *Chlorella*, refining the analysis to

examine the centromeres between each chromosome, aiming to investigate whether

there is high collinearity or if they originate from the same ancestor, and to

understand the composition and structure of their centromeres. We conduct

collinearity analysis on the three species assembled at T2T-level, and then painting

the collinearity relationships and the putative centromeres (**Figure 5B**). The

chromosome 3 of SLA-04, the chromosome 1 of NS4-2, and the chromosome 1 of

DBH exhibit strong collinearity, indicating they originate from the same ancestor

(**Figure 5B**). We further extracted the centromere sequences of these three

chromosomes for dot plot analysis and found significant structural differences among

them (**Figure 5C**). The composition of TE (transposable element) families among the

three chromosomes also exhibits significant differences (**Figure 5C**). Taking

LTR/Copia as an example, in SLA-04, CEN3 comprises 44% of the selected TE

family within the centromere. In NS4-2, CEN1 comprises 88% of the selected TE

family within the centromere. Meanwhile, in DBH, CEN1 comprises 27% of the

selected TE family within the centromere. These results indicate that the evolution of

centromere sequences is significantly faster than that of non-centromeric regions.

Discussion

Recent advances in third-generation sequencing technologies have enabled the

telomere-to-telomere (T2T) assembly of a number of genomes including the model

plant *Arabidopsis thaliana* (Wang et al. 2022) and the model green alga

Chlamydomonas reinhardtii (Payne et al. 2023). To complement these, we employed

HiFi reads, ONT ultra-long reads, and stringent assembly methods to generate T2T

assemblies for *C. sorokiniana* strain NS4-2 and *C. pyrenoidosa* strain DBH. These

T2T assemblies represent a significant improvement over previous genome

assemblies, such as *C. sorokiniana* UTEX1230 (Hovde et al. 2018) and *C.*

pyrenoidosa FACHB-9 (Fan et al. 2015), and in that respect they may serve as

valuable resources for future green algae research, in particular towards understanding

more complex regions of the genome, including the centromere.

In human centromeres, there are specific hypomethylated regions containing a

higher concentration of the centromeric histone CENP-A. This enrichment of CENP-

A suggests that these regions serves as functional kinetochore-binding sites (Logsdon

et al. 2021; Altemose et al. 2022). It's worth noting that this particular characteristic is

conserved in higher plants (Zhang et al. 2008; Wang et al. 2022). Interestingly, this

study revealed that the same epigenetic phenomenon is also present in green algae,

suggesting that this is a highly conserved epigenetic trait among eukaryotes in general.

Centromeric satellite sequences have a monomer size of 100-200 base pairs, typically

falling within the range of 150 to 180 bp, i.e. a length sufficient to wrap a single

nucleosome (Melters et al. 2013). However, the origin of these monomers remains

unclear. Analysis of ancestral crop species, such as *Gossypium raimondii* (Han et al.

2016), an ancestor species of tetraploid cotton, and *Triticum monococcum* (Ahmed et

al. 2023), the first domesticated wheat species, found that their centromeres were

comprised largely of retrotransposons rather than long arrays of centromeric satellites,

supporting the hypothesis that centromere repeats originated from non-random

retrotransposon insertions (Huang et al. 2021). We found that each of the green algae

in this study had TE-rich centromeres, further supporting the hypothesis that

centromeric monomers in higher plants originated from TEs in ancestral plants.

Nevertheless, different taxa of green algae exhibited significant differences in their

centromeric TE composition. For instance, *Chlamydomonas reinhardtii* and

Coccomyxa subellipsoidea both consist of LINE/L1 elements, but they are composed

of different subfamilies. Interestingly, the sister species of *C. reinhardtii*, *C.*

zofingiensis, primarily consists of LTR/Copia sequences in its centromeric sequences

(Roth et al, 2017), indicating that each species continually optimizes its specific

centromeric TE sequences during evolution. Despite being species of the same genus,

Chlorella, the composition and structure of centromeric sequences between

chromosomes derived from the common ancestor exhibit significant differences. It

can be said that the centromere is the fastest-evolving part of the chromosome

(Bensasson et al. 2008; Suzuki et al. 2020).

In summary, we assembled two *Chlorella* species (NS4-2 and DBH) to T2T level,

enabling a comprehensive analysis of their centromeric sequence composition and

evolution. These findings contribute to our understanding of centromere evolution in

green algae and, by extension, to their descendants.

Materials and methods

Culture conditions

Chlorella pyrenoidosa strain DBH (obtained from the Freshwater Algae Culture

Collection at the Institute of Hydrobiology) and *C. sorokiniana* strain NS4-2 (isolated

and purified from a water sample taken from Xueshi Lake at Shanghai Normal

University) were grown mixotrophically in a tris-acetate-phosphate (TAP) medium.

Both species were cultivated in Erlenmeyer flasks under continuous growth light

conditions (30-40 μmol photons m^{-2} s^{-1}) at a temperature of 25 °C, with agitation at

100 rpm, for maintenance purposes.

Karyotyping for the two algae species

The culture sample was harvested by centrifugation at 12,000 rpm for 2 minutes and

then washed sequentially with sterilized BG-11 medium and distilled water. The

resulting algal pellet was suspended in a 2 mM 8-hydroxyquinoline solution and kept

in darkness at 4 °C for 4 hours. Subsequently, the algae pellet was fixed with a

solution of 3 parts ethanol to 1 part acetic acid (v/v) for 24 hours to ensure fixation of

the algae cells in the metaphase of mitosis. After fixation, the cells were washed three

times with distilled water and then subjected to enzymatic digestion. The enzymatic

solution contained 2.5% (w/w) cellulose Onozuka R 10 (Yakult Pharmaceutical,

Tokyo, Japan) and 2.5% (w/w) pectinase Y23 (Yakult Pharmaceutical, Tokyo, Japan),

and the digestion process lasted for 18 hours at 37 °C. Following digestion, the cells

were homogenized at 15,000 rpm for 2 minutes and then centrifuged. The resulting

pellet was resuspended in 100 μL of acetic acid and vortexed for 20 seconds. The cell

suspension was dropped onto a sterile slide and allowed to air dry. Finally, the cells

were stained with an improved carbol fuchsin solution and examined under a

microscope to capture images of the metaphase stage.

388 Genomic DNA extraction

Samples were collected, and high molecular weight genomic DNA was prepared by

the CTAB method and followed by purification with QIAGEN® Genomic kit

(Cat#13343, QIAGEN) for regular sequencing, according to the standard operating

procedure provided by the manufacturer. The DNA degradation and contamination of

the extracted DNA was monitored on 1% agarose gels. DNA purity was then detected

using a NanoDrop One UV-VIS spectrophotometer (Thermo Fisher Scientific, USA),

of which OD 260/280 ranged from 1.8 to 2.0 and OD 260/230 from 2.0 to 2.2. DNA

concentration was subsequently measured using a Qubit 4.0 fluorometer (Invitrogen,

USA).

398 HiFi sequencing and genome assembly

SMRTbell libraries were prepared following the standard protocol of PacBio, using 15

Kb preparation solutions from Pacific Biosciences, USA. The library preparation

steps were conducted according to the method described in Wang et al. (2022).

Sequencing was performed on a PacBio Sequel II instrument with Sequencing Primer

V2 and Sequel II Binding Kit 2.0 at Xi'an Haorui Genomics Technology Co., Ltd.

(Xi'an, China). For the NS4-2 strain, a total of 25.87 Gb of HiFi reads were generated,

providing a coverage of 438×. The N50 of the reads was 15,694 bp. Regarding the

DBH strain, a total of 23.74 Gb of HiFi reads were generated, resulting in a coverage

of 308×. The N50 of the reads was 13,338 bp. The HiFi reads were assembled using

hifiasm v. 0.16.1-r375 (Cheng et al. 2021) with default parameters. gfatools

(https://github.com/lh3/gfatools) was employed to convert the sequence graphs in the

GFA format to FASTA format.

411 ONT sequencing and genome assembly

For the ultra-long Oxford Nanopore sequencing library, approximately 8–10 μg of

genomic DNA was selected (>50 kb) with the SageHLS HMW library system (Sage

Science, Beverly, MA), and then processed using the ligation sequencing 1D kit

(SQK-LSK109, Oxford Nanopore Technologies, Oxford, UK) according to the

manufacturer's instructions. DNA libraries (approximately 800 ng) were constructed

and sequenced using a PromethION (Oxford Nanopore Technologies, Oxford, UK) at

the Genome Center of Grandomics (Wuhan, China). A total of 22.12 Gb ONT long

reads with 368× coverage (read N50: 59,282 bp) were generated. To assemble these

into contigs, the long-read assembler NextDenovo v. 2.0 (Hu et al. 2023) was used

with parameters 'read_cutoff = 5k' and 'seed_cutoff = 132,382'. Contigs were then

polished using Nextpolish v. 1.3.1 (Hu et al. 2020) with parameters 'hifi_options = -

min_read_len 10k -max_read_len 45k' and 'hifi_minimap2_options = -x asm20'.

Hi-C sequencing and scaffolding

A Hi-C library was generated from cross-linked chromatins of algal cells using a

standard Hi-C protocol, and then sequenced using the NovaSeq 6000 platform. The

Hi-C sequencing data, as outlined in Table S1, were utilized to construct a Hi-C

contact map. The construction of the contact map was performed using Juicer v. 1.5

(Durand et al. 2016a) with parameter '-s DpnII'. The resulting contact map was

manually inspected and verified using Juicebox v. 1.11.08 (Durand et al. 2016b).

Evaluation of the putative centromeres

To assess the assembly structure of the predicted centromeric regions, we employed

VerityMap (Mikheenko et al. 2020). First, the PacBio HiFi reads were aligned to the

assembly using minimap2 v. 2.20-r1061 (Li, 2018) with '-x map-hifi' parameter. The

reads aligning to the centromeric regions were extracted using SAMtools v. 1.9 Li et

al. 2009). These extracted HiFi reads were then input into VerityMap using the

following parameters: '--careful --reads HiFi reads -d hifi '.

Genome comparisons

The synteny relationships among the chromosomes were estimated using MCScanX

(Wang et al. 2012) with parameter '-s 10' and visualized using Circos v. 0.69-8

(Krzywinski et al. 2009). Genomic alignment dot plot of chromosomes was generated

using Gepard v. 2.1 (Krumsiek et al. 2007). To assess genome completeness, we used

BUSCO v. 3.0.2 with the dataset 'chlorophyta_odb10 v. 4' (Simão et al. 2015).

Chromosome painting among three species of the genus Chlorella using the IAGS

software (Gao et al. 2022).

Genome annotation

For the NS4-2 and DBH genome annotations, we initially predicted the *de novo* gene

structure using GeneMark-ES with the parameters '--ES' (Borodovsky and Lomsadze,

2011). Subsequently, we employed the 'chlorella gene model' to train Augustus v. 3.3

(Mario and Burkhard, 2005). To identify protein-coding genes, we utilized the

MAKER2 pipeline (Cantarel et al. 2008), which integrated the gene models predicted

from GeneMark-ES and Augustus, as well as the protein sequences from closely

related organisms such as *Chlamydomonas reinhardtii* (GenBank accession:

GCF_000002595.2) (Merchant et al. 2007), *Chlorella variabilis* (GCF_000147415.1)

(Blanc et al. 2010), *Chlorella sorokiniana* (GCA_002245835.2) (Arriola et al. 2018),

and *Micractinium conductrix* (GCA_002245815.2) (Arriola et al. 2018), which are

known to be close relatives of NS4-2 and DBH. For the identification of repeat

sequences, we employed RepeatMasker v. 4.0.7 (http://www.repeatmasker.org) and

RepeatModeler v. 2.0.1 (Flynn et al. 2020). The TE-algae library used to annotate the

selected species consists of three parts: the *de novo* library of three Chlorella species

from RepeatModeler, a library from a previous study (Craig et al. 2021), and the

Chlorophyta Repbase library (https://www.repeatmasker.org/) (**Data S1**). The TE-

algae library underwent redundancy reduction using cd-hit (Li and Godzik 2006). The

karyoplots shown in this study were generated using KaryoploteR package in R (Gel

and Serra, 2017).

Phylogenetic tree construction

The phylogenetic trees depicted in Figure 1 and Figure 4 were generated utilizing

OrthoFinder v. 2.2.7 with the default parameters (Emms and Kelly, 2019). Figure

S1's phylogenetic tree was constructed using MEGA v. 7 softwareand employed the

Kimura 2-parameter model, accompanied by 1000 bootstrap replicates (Kumar et al.

2016). For Figure S13, the phylogenetic tree was crafted using RAxML-NG v. 1.1

(Kozlov et al. 2019) with the specified parameters '--all --bs-trees 1000', and the

HKY+G4 model was chosen through ModelTest-NG v. 0.1.7 (Darriba et al. 2020).

PacBio DNA methylation analysis

Methylation for NS4-2 and DBH was inferred with ccsmeth v.0.4.1 (Ni et al. 2023),

using kinetics features from PacBio CCS reads. Methylation of CCS reads was

predicted using 'ccsmeth call_mods' with the model

'model_ccsmeth_5mCpG_call_mods_attbigru2s_b21.v2.ckpt'. Then, we used

'ccsmeth align_hifi' to align the reads to their respective genome. The methylation

frequencies were calculated by using 'ccsmeth call_freqb' with the aggregate model

'model_ccsmeth_5mCpG_aggregate_attbigru_b11.v2p.ckpt'.

ONT DNA methylation analysis

Nanopolish v. 0.13.2 with the parameters 'call-methylation --methylation cpg' was

used to measure the frequency of CpG methylation. The ONT reads were aligned to

whole-genome assemblies via minimap2 (Li, 2018). The script

'calculate_methylation_frequency.py', provided in the methplotlib package (De

Coster et al. 2020), was then used to generate the methylation frequency. Methylation

density is defined as the number of occurrences with a methylation frequency greater

than 0.8 within genomic sliding windows (100-base pair window advanced at 50-base

pair increments).

CENH3 ChIP-seq

We first used BLASTP to identify the amino acid sequence of CENH3 in NS4-2, with

the reference sequence HTR12 from *Arabidopsis thaliana* (Genbank accession:

NP_001322867.1), and the '-evalue' parameter set to 1e-30. We selected the coding

gene NS|00958 as the unique matching result for the CENH3 sequence of the NS4-2

species for subsequent antibody preparation. An antigen with the peptide sequence

'MARTKQTKPSPSKRGA' corresponding to the N terminus of NS4-2_CENH3 was

used to produce the antibody (Beijing QiWei YiCheng Tech Co., Ltd.).

Shanghai Jiayin Biotechnology Co., Ltd conducted ChIP assays following a

modified version of the standard crosslinking ChIP protocol. In summary, the culture

sample was harvested by centrifugation at 12,000 rpm for 2 minutes and sequentially washed with sterilized BG-11 and distilled water. Subsequently, approximately 4g of algae pellet was obtained and ground in liquid nitrogen. The ground sample was fixed with 1% formaldehyde at room temperature for 15 minutes, followed by a 5-minute treatment with 0.125 M glycine. After fixation, the sample was washed, resuspended in a lysis buffer, and sonicated to generate DNA fragments. Following sonication, immunoprecipitation was carried out using the custom antibody NS4-2_CENH3. The immunoprecipitated complex was washed, and DNA was extracted. DNA purification was performed using the Universal DNA Purification Kit (#DP214). A ChIP-seq library was prepared by employing the ChIP-Seq DNA sample preparation kit (NEBNext® UltraTMII DNA), adhering to the manufacturer's instructions. The extracted DNA was ligated to specific adaptors and subsequently subjected to deep sequencing on the Illumina Novaseq 6000 platform in 150bp paired-end mode.

The ChIP/input paired-end reads were mapped to the NS4-2 genome using the 'mem' algorithm of BWA (Li and Durbin 2009). We determined the ChIP/input coverage ratio by employing the deeptools function 'bamCompare', with a threshold of MAPQ $\geq$ 30. The output format was in BigWig. We carried out peak calling using MACS2 (Zhang et al. 2008) with the parameters '-t ChIP.bam -c input.bam -f BAM --outdir chipseq -n chipseq -B --nomodel --keep-dup 1 -g 58000000'.

Data availability

The whole genome sequence data reported in this paper have been deposited in the Genome Warehouse in National Genomics Data Center, Beijing Institute of Genomics, Chinese Academy of Sciences / China National Center for Bioinformation (GWH: GWHBKBA00000000 for NS4-2; GWH: GWHBKBB00000000 for DBH) that is publicly accessible at https://ngdc.cncb.ac.cn/gwh. The genome annotations of NS4-2 and DBH has been deposited in figshare (10.6084/m9.figshare.24503881). The raw data for the PacBio HiFi reads, ONT long-reads, Hi-C Illumina short reads and ChIP-seq data have been deposited in the Genome Sequence Archive at the National Genomics Data Center, Beijing Institute of Genomics, Chinese Academy of Sciences / China National Center for Bioinformation (GSA: CRA007945), and are publicly

531   accessible at https://ngdc.cncb.ac.cn/gsa.

Author contributions

KY and WM conceived and designed the project. KY and BW designed the experiments. BW, YJ, SG, and HG analyzed the results. BW, YJ, ND, JY, SB, SG, WH, SW, and HG performed all the experiments and analyzed the data. BW, YJ and ND created the figures. BW, YJ and SB wrote the manuscript. SB, XY, WM, and KY reviewed the manuscript.

Table 1. Comparison genomic features of *Chlorella sorokiniana* NS4-2 and *C. pyrenoidosa* DBH assemblies.

Genome features	NS4-2	UTEX1230[a]	DBH	FACHB-9[b]
Nuclear genome size (Mb)	58.11	58.36	53.41	56.77
No. of chromosomes (count)	12	NA	13	NA
No. of contigs (count)	13	18	14	8193
GC content (%)	63.95	63.95	66.94	66.71
Contigs N50 (bp)	4,352,225	3,818,101	4,302,578	9757
Scaffolds N50 (bp)	4,714,146	4,091,730	4,383,467	1,392,758
Protein coding genes (count)	12,691	12,871	11,327	10,283
Repeat content (%)	8.84	8.89	8.91	5.46

[a] UTEX1230 is a strain of *C. sorokiniana* (Hovde et al. 2018).
[b] FACHB-9 is a strain of *C. pyrenoidosa* (Fan et al. 2015).

Figure legends

Figure 1. T2T genome assemblies and circular plot of *Chlorella sorokiniana* NS4-2 and *C. pyrenoidosa* DBH. (A) Micrograph of *C. sorokiniana* NS4-2, bar = 20 μm; **(B)** The karyotype diagram indicates that NS4-2 has 12 chromosomes; **(C)** Hi-C contact map of the NS4-2 T2T genome assembly (chromosome 1-12); **(D)** Circular plot shows the T2T genome features of NS4-2 and DBH. Track a shows the karyotype distribution of the assembled chromosomes; track b, GC density; track c, gene density; track d, LINE element density; track e, LTR element density; track f, putative centromeres. Densities was calculated in 50-Kb windows. The innermost of the circular plots represents the synteny relationship between NS4-2 and DBH.

Figure 2. CENH3 binding defines centromeres in *Chlorella sorokiniana* NS4-2.
Schematic overview of the chromosomes of NS4-2 showing the normalized CENH3
ChIP-seq read coverage (RPGC normalization in 1-Kb bins with 3-Kb smoothing,
plot generated using kpPlotBigWig function in karyoploteR), predicted centromere
regions, gene density, and LTR/Copia density. Densities was calculated in 5-Kb
windows.

Figure 3. Centromeric methylation features of *Chlorella sorokiniana* NS4-2. (A)
Methylation density distribution chart of chrosomome 1 in NS4-2. The red area
represents the centromere region. Zoomed-in overview of the chromosome 1 of NS4-
2 showing the methylation density distribution, CENH3 ChIP-seq signals (plot
generated using kpPlotBAMCoverage function in karyoploteR), as well as the regions
of centromere (green band) and ChIP-seq peaks called by MACS2 (light green); **(B)**
The box- and violin plots display a comparison of methylation density in NS4-2
among centromere (CEN), non-centromere (non-CEN), peaks within the centromere
(CEN_peaks), and non-peaks within the centromere (CEN_nonPeaks); **(C)** The box-
and violin plots display a comparison of methylation density in DBH among CEN and
nonCEN.

Figure 4. Centromeric size and compositions in green algae. (A) Phylogenomic
tree of six green algae species with the red algae *Cyanidioschyzon merolae* utilized as
an outgroup species. The figure on the right illustrates a comparison of genome size
and centromere length among these selected species; **(B)** Relationship between the
genome size of each chromosome (log-transformed) and each centromere size
(logtransformed). The figure was generated using basic plot functions in R; **(C-H)**
The distribution of different repeat subfamilies in centromeres (CEN; dark blue) and
across the genome (non-CEN; light gray) of *Chlamydomonas reinhardtii* (CR) **(C)**,
Chromochloris zofingiensis (CZ) **(D)**, *Coccomyxa subellipsoidea* (CS) **(E)**, *Chlorella
sorokiniana* (SLA-04) **(F)**, *Chlorella pyrenoidosa* (DBH) **(G)** and *Chlorella
sorokiniana* (NS4-2) **(H)**.

Figure 5. Centromeric TEs evolutionary in green algae and *Chlorella* species. (A) The representation of the main centromeric TE families of the selected species on the phylogenetic tree. MCRA stands for most recent common ancestor; **(B)** Synteny analyses of the 13 chromosomes of SLA-04, 12 chromosomes of NS4-2 and 13 chromosomes of DBH. The corresponding syntenic regions are marked using the same color. Approximate locations of putative centromeres are indicated by ellipses. The red circles represent the collinearity of the whole chromosomes; **(C)** The dotplot display of the centromeres showing the collinearity of entire chromosomes from three *Chlorella* species, along with the composition of TEs. Referring to the legend of (A), different colors represent different TE families.

Supporting information

Figure S1. Maximum likelihood phylogenetic tree based on the analysis of 18S gene sequences. GenBank accession numbers in brackets.

Figure S2. T2T genome assemblies and centromere identification of *Chlorella pyrenoidosa* strain DBH. **(A)** Micrograph of DBH,bar = 20 μm; **(B)** The karyotype diagram indicates that DBH has 13 chromosomes; **(C)** Hi-C contact map of the DBH T2T genome assembly (chromosome 1-13); **(D)** Schematic overview of the chromosomes of DBH showing the predicted centromere regions (green band).

Figure S3. High-quality genome assembly pipelines. **(A)** *Chlorella sorokiniana* NS4-2 genome assembly; **(B)** *C. pyrenoidosa* DBH genome assembly.

Figure S4. Demonstration of collinearity between chromosomes. **(A)** Between two contigs of *Chlorella sorokiniana* NS4-2 and chromosome 2 of *C. vulgaris*; **(B)** Between two contigs of *C. pyrenoidosa* DBH and chromosome 2 of *C. vulgaris.*

Figure S5. Self-Dot plot showing potential centromeric features and regions of two previous assemblies, *Chlamydomonas reinhardtii* and *Coccomyxa subellipsoidae.*

Figure S6. Self-dot plot of the 12 chromosomes of *Chlorella sorokiniana* NS4-2, with green boxes indicating potential centromeric regions.

Figure S7. Self-dot plot of the 13 chromosomes of *Chlorella pyrenoidosa* DBH, with

green boxes indicating potential centromeric regions.

Figure S8. Evaluation of the 12 putative centromere assemblies of NS4-2 using

VerityMap. The xaxis is centromere position, the y-axis is percent of deviated reads.

Only a low percentage (12.5%) of deviated reads were detected in CEN11, indicating

no potential large structural errors in the assembly.

FigureS9. ChIP/Input ratio of the 12 chromosomes of *Chlorella sorokiniana* NS4-2.

Schematic overview showing the ratio (plot generated using the kpPlotBigWig

function in karyoploteR) and the putative centromere regions (red band).

FigureS10. Using chromosome 10 of *Chlorella sorokiniana* NS4-2 as an example to

demonstrate the dot plot method for identifying potential centromere positions in the

genome.

Figure S11. Evaluation of the 13 putative centromere assemblies of *Chlorella*

pyrenoidosa DBH using VerityMap. The xaxis is centromere position, the y-axis is

percent of deviated reads.

Figure S12. Centromeric methylation features of *Chlorella sorokiniana* NS4-2.

Schematic overview depicting each centromere along with its 50 Kb upstream and

downstream regions of NS4-2. The illustration shows the distribution of methylation

density (determined by HiFi data), CENH3 ChIP-seq signals (plot generated using

kpPlotBAMCoverage function in karyoploteR), and the regions of centromeres (green

bands).

Figure S13. Centromeric methylation features of *Chlorella pyrenoidosa* DBH.

Schematic overview depicting each centromere along with its 50 Kb upstream and

downstream regions of DBH. The illustration shows the distribution of methylation

density determined by HiFi data, methylation density determined by ONT data, and

the regions of centromere (green band).

Table S1. Details of the sequencing data for *Chlorella sorokiniana* NS4-2 and *C.*

pyrenoidosa DBH.

Table S2. *Chlorella sorokiniana* NS4-2 putative centromeric coordinates.

Table S3. *Chlorella pyrenoidosa* DBH putative centromeric coordinates.

Table S4. *Chlamydomonas reinhardtii* CC4532 putative centromeric coordinates.

Table S5. *Chromochloris zofingiensis* putative centromeric coordinates.

Table S6. *Coccomyxa subellipsoidea* C169 putative centromeric coordinates.

Table S7. *Chlorella sp.* SLA-04 putative centromeric repeat annotations.

Table S8. Quantitative analysis of the number, length, and content of various transposable element families at the centromere of *Chlamydomonas reinhardtii*.

Table S9. Quantitative analysis of the number, length, and content of various transposable element families at the centromere of *Chromochloris zofingiensis*.

Table S10. Quantitative analysis of the number, length, and content of various transposable element families at the centromere of *Coccomyxa subellipsoidea* C-169.

Table S11. Quantitative analysis of the number, length, and content of various transposable element families at the centromere of *Chlorella* sp. SLA-04.

Table S12. Quantitative analysis of the number, length, and content of various transposable element families at the centromere of *Chlorella sorokiniana* NS4-2.

Table S13. Quantitative analysis of the number, length, and content of various transposable element families at the centromere of *Chlorella sorokiniana* DBH.

Table S14. *Chlamydomonas reinhardtii* putative centromeric repeat annotations.

Table S15. *Coccomyxa subellipsoidea* putative centromeric repeat annotations.

Table S16. *Chromochloris zofingiensis* putative centromeric repeat annotations.

Table S17. *Chlorella sp.* SLA-04 putative centromeric repeat annotations.

Table S18. *Chlorella sorokiniana* NS4-2 putative centromeric repeat annotations.

Table S19. *Chlorella pyrenoidosa* DBH putative centromeric repeat annotations.

Data S1. TE library used in this study.

References

Ahmed, H.I., Heuberger, M., Schoen, A., Koo, D.H., Quiroz-Chavez, J., Adhikari, L., Raupp, J., Cauet, S., Rodde, N., Cravero, C., Callot, C., Lazo, G.R., Kathiresan, N., Sharma, P.K., Moot, I., Yadav, I.S., Singh, L., Saripalli, G., Rawat, N., Datla, R., Athiyannan, N., Ramirez-Gonzalez, R.H., Uauy, C., Wicker, T., Tiwari, V.K., Abrouk, M., Poland, J., Krattinger, S. G. (2023). Einkorn genomics sheds light on history of the oldest domesticated wheat. Nature **620**, 830-838.

Altemose, N., Logsdon, G. A., Bzikadze, A. V., Sidhwani, P., Langley, S. A.,

Caldas, G. V., Hoyt, S. J., Uralsky, L., Ryabov, F. D., Shew, C. J., Sauria, M. E. G., Borchers, M., Gershman, A., Mikheenko, A., Shepelev, V. A., Dvorkina, T., Kunyavskaya, O., Vollger, M. R., Rhie, A., McCartney, A. M., Asri, M., Lorig-Roach, R., Shafin, K., Lucas, J.K., Aganezov, S., Olson, D., de Lima, L.G., Potapova, T., Hartley, G.A., Haukness, M., Kerpedjiev, P., Gusev, F., Tigyi, K., Brooks, S., Young, A., Nurk, S., Koren, S., Salama, S.R., Paten, B., Rogaev, E.I., Streets, A., Karpen, G.H, Dernburg, A.F., Sullivan, B.A., Straight, A.F., Wheeler, T.J., Gerton, J.L., Eichler, E.E., Phillippy, A.M., Timp, W., Dennis, M.Y., O'Neill, R.J., Zook, J.M., Schatz, M.C., Pevzner, P.A., Diekhans, M., Langley, C.H., Alexandrov, I.A., and Miga, K.H. (2022). Complete genomic and epigenetic maps of human centromeres. Science **376**, eabl4178.

Arriola, M.B., Velmurugan, N., Zhang, Y., Plunkett, M.H., Hondzo, H., and Barney, B.M. (2018). Genome sequences of *Chlorella sorokiniana* UTEX 1602 and *Micractinium conductrix* SAG 241.80: implications to maltose excretion by a green alga. Plant J **93**, 566-586.

Bensasson, D., Zarowiecki, M., Burt, A., and Koufopanou, V. (2008). Rapid evolution of yeast centromeres in the absence of drive. Genetics **178**, 2161-2167.

Blanc, G., Agarkova, I., Grimwood, J., Kuo, A., Brueggeman, A., Dunigan, D.D., Gurnon, J., Ladunga, I., Lindquist, E., Lucas, S., Pangilinan, J., Pröschold, T., Salamov, A., Schmutz, J., Weeks, D., Yamada, T., Lomsadze, A., Borodovsky, M., Claverie, J.M., Grigoriev, I.V., and Van Etten, J.L. (2012). The genome of the polar eukaryotic microalga *Coccomyxa subellipsoidea* reveals traits of cold adaptation. Genome Biol **13**, R39.

Blanc, G., Duncan, G., Agarkova, I., Borodovsky, M., Gurnon, J., Kuo, A., Lindquist, E., Lucas, S., Pangilinan, J., Polle, J., Salamov, A., Terry, A., Yamada, T., Dunigan, D.D., Grigoriev, I.V., Claverie, J.M., and Van Etten, J.L. (2010). The *Chlorella variabilis* NC64A genome reveals adaptation to photosymbiosis, coevolution with viruses, and cryptic sex. Plant Cell **22**, 2943-2955.

Bi, G., Zhao, S., Yao, J., Wang, H., Zhao, M., Sun, Y., Hou, X., Haas, F. B., Varshney, D., Prigge, M., Rensing, S. A., Jiao, Y., Ma, Y., Yan, J., and Dai, J. (2024). Near telomere-to-telomere genome of the model plant Physcomitrium patens. Nat Plants 10,327-343.

Borodovsky, M., and Lomsadze, A. (2011). Eukaryotic gene prediction using GeneMark.hmm-E and GeneMark-ES. Curr Protoc Bioinformatics **4**, 4.6.1-4.6.10.

Cantarel, B.L., Korf, I., Robb, S.M., Parra, G., Ross, E., Moore, B., Holt, C., Sánchez Alvarado, A., and Yandell, M. (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. Genome Res **18**, 188-196.

Cecchin, M., Marcolungo, L., Rossato, M., Girolomoni, L., Cosentino, E., Cuine, S., Li-Beisson, Y., Delledonne, M., and Ballottari, M. (2019). *Chlorella vulgaris* genome assembly and annotation reveals the molecular basis for

metabolic acclimation to high light conditions. Plant J **100**, 1289-1305.

Chen, J., Wang, Z., Tan, K., Huang, W., Shi, J., Li, T., Hu, J., Wang, K., Wang, C., Xin, B., Zhao, H., Song, W., Hufford, M.B., Schnable, J.C., Jin, W., and Lai, J. (2023). A complete telomere-to-telomere assembly of the maize genome. **Nat Genet** 55, 1221-1231.

Cheng, H., Concepcion, G.T., Feng, X., Zhang, H., and Li, H. (2021). Haplotype-resolved *de novo* assembly using phased assembly graphs with hifiasm. Nat Methods **18**, 170-175.

Craig, R.J., Hasan, A.R., Ness, R.W., and Keightley, P.D. (2021). Comparative genomics of *Chlamydomonas*. Plant Cell **33**, 1016-1041.

Craig, R.J., Gallaher, S.D., Shu, S., Salomé, P.A., Jenkins, J.W., Blaby-Haas, C.E., Purvine, S.O., O'Donnell, S., Barry, K., Grimwood, J., Strenkert, D., Kropat, J., Daum, C., Yoshinaga, Y., Goodstein, D.M., Vallon, O., Schmutz, J., and Merchant, S.S. (2023). The *Chlamydomonas* Genome Project, version 6: Reference assemblies for mating-type plus and minus strains reveal extensive structural mutation in the laboratory. Plant Cell **35**, 644-672.

Darriba, D., Posada, D., Kozlov, A. M., Stamatakis, A., Morel, B., and Flouri, T. (2020). ModelTest-NG: A new and scalable tool for the selection of DNA and protein evolutionary models. Mol Biol Evol **37**, 291-294.

Dias, G.B., Svartman, M., Delprat, A., Ruiz, A., and Kuhn, G.C. (2014). Tetris is a foldback transposon that provided the building blocks for an emerging satellite DNA of *Drosophila virilis*. Genome Biol Evol **6**, 1302-1313.

Durand, N.C., Robinson, J.T., Shamim, M.S., Machol, I., Mesirov, J.P., Lander, E.S., and Aiden, E.L. (2016a). Juicebox provides a visualization system for Hi-C contact maps with unlimited zoom. Cell Syst **3**, 99-101.

Durand, N.C., Shamim, M.S., Machol, I., Rao, S.S., Huntley, M.H., Lander, E.S., and Aiden, E.L. (2016b). Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. Cell Syst **3**, 95-98.

Emms, D. M., and Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative genomics. Genome Biol **20**, 238.

Fan, J., Ning, K., Zeng, X., Luo, Y., Wang, D., Hu, J., Li, J., Xu, H., Huang, J., Wan, M., Wang, W., Zhang, D., Shen, G., Run, C., Liao, J., Fang, L., Huang, S., Jing, X., Su, X., Wang, A., Bai, L., Hu, Z. X, J., and Li, Y. (2015). Genomic foundation of starch-to-lipid switch in oleaginous *Chlorella* spp. Plant Physiol **169**, 2444-2461.

Flynn, J.M., Hubley, R., Goubert, C., Rosen, J., Clark, A.G., Feschotte, C., and Smit, A.F. (2020). RepeatModeler2 for automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A **117**, 9451-9457.

Gao, S., Yang, X., Guo, H., Zhao, X., Wang, B., and Ye, K. (2023). HiCAT: A tool for automatic annotation of centromere structure. Genome Biol **24**, 58.

Gao, S., Yang, X., Sun, J., Zhao, X., Wang, B., and Ye, K. (2022). IAGS: Inferring ancestor genome structure under a wide range of evolutionary scenarios. Mol Biol Evol **39**, msac041.

Gel, B., and Serra, E. (2017). karyoploteR: an R/Bioconductor package to plot

customizable genomes displaying arbitrary data. Bioinformatics **33**, 3088-3090.

Goemann, C. L., Wilkinson, R., Henriques, W., Bui, H., Goemann, H. M.,
Carlson, R. P., Viamajala, S., Gerlach, Robin., and Wiedenheft, B. (2023).
Genome sequence, phylogenetic analysis, and structure-based annotation reveal
metabolic potential of *Chlorella* sp. SLA-04. Algal Res **69**, 102943.

Han, J., Masonbrink, R.E., Shan, W., Song, F., Zhang, J., Yu, W., Wang, K., Wu,
Y., Tang, H., Wendel, J.F., and Wang, K. (2016). Rapid proliferation and
nucleolar organizer targeting centromeric retrotransposons in cotton. Plant J **88**,
992-1005.

Henikoff, S., Ahmad, K., and Malik, H.S. (2001). The centromere paradox: stable
inheritance with rapidly evolving DNA. Science **293**, 1098-1102.

Hovde, B.T., Hanschen, E.R., Tyler, C.R. S., Lo, C.C., Kunde, Y., Davenport, K.,
Daligault, H., Msanne, J., Canny, S., Eyun, S.I., Riethoven, J.M., Polle, J.,
Starkenburg, S.R. (2018). Genomic characterization reveals significant
divergence within *Chlorella sorokiniana* (Chlorellales, Trebouxiophyceae). Algal
Res **35**, 449-461.

Hu, J., Fan, J., Sun, Z., and Liu, S. (2020). NextPolish: a fast and efficient genome
polishing tool for long-read assembly. Bioinformatics **36**, 2253-2255.

Hu, J., Wang, Z., Sun, Z., Hu, B., Ayoola, A.O., Liang, F., Li, J., Sandoval, J.R.,
Cooper, D.N., Ye, K., Ruan, J., Xiao, C., Wang, D., Wu, D., and Wang, S.
(2023). An efficient error correction and accurate assembly tool for noisy long
reads. bioRxiv, 2023.03.09.531669.

Huang, Y., Ding, W., Zhang, M., Han, J., Jing, Y., Yao, W., Hasterok, R., Wang,
Z., and Wang, K. (2021). The formation and evolution of centromeric satellite
repeats in *Saccharum* species. Plant J **106**, 616-629.

Kanesaki, Y., Imamura, S., Matsuzaki, M., and Tanaka, K. (2015). Identification
of centromere regions in chromosomes of a unicellular red alga,
Cyanidioschyzon merolae. FEBS Lett **589**, 1219-1224.

Kozlov, A.M., Darriba, D., Flouri, T., Morel, B., and Stamatakis, A. (2019).
RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood
phylogenetic inference. Bioinformatics **35**, 4453-4455.

Krumsiek, J., Arnold, R., and Rattei, T. (2007). Gepard: a rapid and sensitive tool
for creating dot plots on a genome scale. Bioinformatics **23**, 1026-1028.

Krzywinski, M., Schein, J., Birol, I., Connors, J., Gascoyne, R., Horsman, D.,
Jones, S.J., and Marra, M.A. (2009). Circos: an information aesthetic for
comparative genomics. Genome Res **19**, 1639-1645.

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: Molecular Evolutionary
Genetics Analysis Version 7.0 for Bigger Datasets. Mol Biol Evol **33**, 1870-1874.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences.
Bioinformatics, **34**, 3094-3100.

Li, H., Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
Wheeler transform. Bioinformatics **25**, 1754-1760.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing

Subgroup (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics **25**, 2078-2079.

Li, W., and Godzik, A. (2006). Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. Bioinformatics **22**, 1658-1659.

Logsdon, G.A., Vollger, M.R., Hsieh, P., Mao, Y., Liskovykh, M.A., Koren, S., Nurk, S., Mercuri, L., Dishuck, P.C., Rhie, A., de Lima, L.G., Dvorkina, T., Porubsky, D., Harvey, W.T., Mikheenko, A., Bzikadze, A.V., Kremitzki, M., Graves-Lindsay, T.A., Jain, C., Hoekzema, K., Murali, S.C., Munson, K.M., Baker, C., Sorensen, M., Lewis, A.M., Surti, U., Gerton, J.L., Larionov, V., Ventura, M., Miga, K.H., Phillippy, A.M., Eichler, E.E. (2021). The structure, function and evolution of a complete human chromosome 8. **Nature** 593, 101-107.

Stanke, M., and Morgenstern, B. (2005). AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. Nucleic Acids Res **33**, W465-W467.

Melters, D.P., Bradnam, K.R., Young, H.A., Telis, N., May, M.R., Ruby, J.G., Sebra, R., Peluso, P., Eid, J., Rank, D., Garcia, J.F., DeRisi, J.L., Smith, T., Tobias, C., Ross-Ibarra, J., Korf, I., and Chan, S.W. (2013). Comparative analysis of tandem repeats from hundreds of species reveals unique insights into centromere evolution. Genome Bio **14**, 1-20.

Merchant, S.S., Prochnik, S.E., Vallon, O., Harris, E.H., Karpowicz, S.J., Witman, G. B., Terry, A., Salamov, A., Fritz-Laylin, L.K., Maréchal-Drouard, L., Marshall, W.F., Qu, L.H., Nelson, D.R., Sanderfoot, A.A., Spalding, M.H., Kapitonov, V.V., Ren, Q., Ferris, P., Lindquist, E., Shapiro, H., ⋯ Grossman, A.R. (2007). The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. Science **318**, 245-250.

McAinsh, A.D., and Marston, A.L. (2022). The four causes: the functional architecture of centromeres and kinetochores. Annu Rev Genet **56**, 279-314.

Michael, T.P. (2014). Plant genome size variation: bloating and purging DNA. Brief Funct Genomics **13**, 308-317.

Miga, K.H. (2020). Centromere studies in the era of 'telomere-to-telomere' genomics. Exp Cell Res **394**, 112127.

Mikheenko, A., Bzikadze, A.V., Gurevich, A., Miga, K.H., and Pevzner, P.A. (2020). TandemTools: mapping long reads and assessing/improving assembly quality in extra-long tandem repeats. Bioinformatics **36**, i75-83.

Naish, M., Alonge, M., Wlodzimierz, P., Tock, A.J., Abramson, B.W., Schmücker, A., Mandáková, T., Jamge, B., Lambing, C., Kuo, P., Yelina, N., Hartwick, N., Colt, K., Smith, L. M., Ton, J., Kakutani, T., Martienssen, R.A., Schneeberger, K., Lysak, M.A., Berger, F., ⋯ Henderson, I.R. (2021). The genetic and epigenetic landscape of the Arabidopsis centromeres. Science **374**, eabi7489.

Ni, P., Nie, F., Zhong, Z., Xu, J., Huang, N., Zhang, J., Zhao, H., Zou, Y., Huang, Y., Li, J., Xiao, C.L., Luo, F., and Wang, J. (2023). DNA 5-methylcytosine detection and methylation phasing using PacBio circular consensus sequencing.

Nat Commun **14**, 4054.

Nozaki, H., Takano, H., Misumi, O., Terasawa, K., Matsuzaki, M., Maruyama, S.,
Nishida, K., Yagisawa, F., Yoshida, Y., Fujiwara, T., Takio, S., Tamura, K.,
Chung, S.J., Nakamura, S., Kuroiwa, H., Tanaka, K., Sato, N., and Kuroiwa,
T. (2007). A 100%-complete sequence reveals unusually simple genomic
features in the hot-spring red alga *Cyanidioschyzon merolae*. BMC Biol **5**, 1-8.

Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., ...
& Phillippy, A.M. (2022). The complete sequence of a human genome. Science
376, 44-53.

Payne, Z.L., Penny, G.M., Turner, T.N., and Dutcher, S.K. (2023). A gap-free
genome assembly of *Chlamydomonas reinhardtii* and detection of translocations
induced by CRISPR-mediated mutagenesis. Plant Commun **4**, 100493.

Plačková, K., Bureš, P., and Zedek, F. (2021). Centromere size scales with genome
size across Eukaryotes. Sci Rep **11**, 19811.

Presting, G.G. (2018). Centromeric retrotransposons and centromere function. Curr
Opin Genet Dev **49**, 79-84.

Rautiainen, M., Nurk, S., Walenz, B.P., Logsdon, G.A., Porubsky, D., Rhie, A.,
Eichler, E.E., Phillippy, A.M., and Koren, S. (2023). Telomere-to-telomere
assembly of diploid chromosomes with Verkko. Nat Biotechnol **41**, 1474-1482.

Roth, M. S., Cokus, S. J., Gallaher, S. D., Walter, A., Lopez, D., Erickson, E.,
Endelman, B., Westcott, D., Larabell, C. A., Merchant, S. S., Pellegrini, M.,
and Niyogi, K. K. (2017). Chromosome-level genome assembly and
transcriptome of the green alga *Chromochloris zofingiensis* illuminates
astaxanthin production. Proc Natl Acad Sci U S A **114**, E4296-E4305.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., and Zdobnov,
E.M. (2015). BUSCO: assessing genome assembly and annotation completeness
with single-copy orthologs. Bioinformatics **31**, 3210-3212.

Song, J.M., Xie, W. Z., Wang, S., Guo, Y.X., Koo, D.H., Kudrna, D., Gong, C.,
Huang, Y., Feng, J.W., Zhang, W., Zhou, Y., Zuccolo, A., Long, E., Lee, S.,
Talag, J., Zhou, R., Zhu, X.T., Yuan, D., Udall, J., Xie, W., ⋯ Chen, L.L.
(2021). Two gap-free reference genomes and a global view of the centromere
architecture in rice. Mol Plant **14**, 1757-1767.

Suzuki, Y., Myers, E. W., and Morishita, S. (2020). Rapid and ongoing evolution of
repetitive sequence structures in human centromeres. Sci Adv **6**, eabd9230.

Talbert, P.B., and Henikoff, S. (2010) Centromeres convert but don't cross. PLoS
Biol **8**, e1000326.

Talbert, P. B., and Henikoff, S. (2020). What makes a centromere?. Exp Cell Res
389, 111895.

Wang, B., Yang, X., Jia, Y., Xu, Y., Jia, P., Dang, N., Wang, S., Xu, T., Zhao, X.,
Gao, S., Dong, Q., and Ye, K. (2022). High-quality Arabidopsis thaliana
genome assembly with nanopore and HiFi long reads. Genom Proteom Bioinf **20**,
4-13.

Wang, Y., Tang, H., Debarry, J.D., Tan, X., Li, J., Wang, X., Lee, T.H., Jin, H.,
Marler, B., Guo, H., Kissinger, J.C., and Paterson, A.H. (2012). MCScanX: a

toolkit for detection and evolutionary analysis of gene synteny and collinearity.
Nucleic Acids Res **40**, e49.

Wick, R.R., Judd, L.M., Gorrie, C.L., and Holt, K.E. (2007). Unicycler: Resolving
bacterial genome assemblies from short and long sequencing reads. PLoS
Comput Biol **13**, e1005595.

Windler, M., Stuart, R., Deutzmann, J.S., Mayali, X., Navid, A., D'haeseleer, P.,
Marcu, O.E., Lipton, M., Nicora, C., and Spormann, A.M. (2022). Bacterial
exometabolites influence *Chlamydomonas* cell cycle and double algal
productivity. FEMS Microbiol Ecol **98**, fiac091.

Zhang, W., Lee, H. R., Koo, D. H., and Jiang, J. (2008). Epigenetic modification of
centromeric chromatin: hypomethylation of DNA sequences in the CENH3-
associated chromatin in Arabidopsis thaliana and maize. Plant Cell **20**, 25-34.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,
Nusbaum, C., Myers, R.M., Brown, M., Li, W., and Liu, X.S. (2008). Model-
based analysis of ChIP-Seq (MACS). Genome Biol **9**, R137.

A

Trebouxiophyceae

Chlorophyceae

0.2

Chlorella sorokiniana NS4-2
Chlorella pyrenoidosa DBH
Chlorella sp. SLA-04
Coccomyxz subellipsoidea
Chromochloris zofingiensis
Chlamydomonas reinhardtii
Cyanidioschyzon merolae

0.0   0.2   0.4   0.6   0.8   1.0

LINE/L1   LINE/RTEX   LINE/Dualen   Unknown
DNA/EnSpm   LTR/Gypsy   LTR/Copia   MCRA

B

Chlorella sp. SLA-04   *Chlorella sorokiniana NS4-2*   *Chlorella pyrenoidosa DBH*

C

Chlorella sp. SLA-04 (CEN3)   *Chlorella sorokiniana NS4-2* (CEN1)   *Chlorella pyrenoidosa DBH* (CEN1)