**Establishment of an Inhalation Administration Non-invasive Murine Model for**

**Rapidly Testing Drug Activity against *Mycobacterium tuberculosis***

Xirong Tian [1,2,3,4*], Yamin Gao [1,2,3,4*], Wanli Ma [1,2,3,4], Jingran Zhang [1,2,3,5], Yanan Ju

[1,2,3,5], Jie Ding [1,2,3, 6], Sanshan Zeng [1,2,3,4], H.M. Adnan Hameed [1,2,3,4], Nanshan Zhong

[7], Gregory M. Cook [4,8,9,10], Jinxing Hu [7, 11#] and Tianyu Zhang [1,2,3,4#]

[1] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine

and Health (GIBH), Chinese Academy of Sciences (CAS), Guangzhou 510530,

China

[2] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious

Diseases, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese

Academy of Sciences (CAS), Guangzhou 510530, China

[3] University of Chinese Academy of Sciences (UCAS), Beijing 100049, China

[4] China-New Zealand Joint Laboratory on Biomedicine and Health, Guangzhou

Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou

510530, China

[5] School of Life Sciences, University of Science and Technology of China, Hefei

230026, China

[6] Institute of Physical Science and Information Technology, Anhui University, Hefei,

230601, China

[7] Guangzhou National Laboratory, Guangzhou 510005, China

[8] Department of Microbiology and Immunology, School of Biomedical Sciences,

University of Otago, Dunedin, 9054, New Zealand

[9] Maurice Wilkins Centre for Molecular Biodiscovery, University of Auckland,

Private Bag 92019, Auckland, 1042, New Zealand

[10] Translational Research Institute, Queensland University of Technology, Brisbane,

Queensland, 4102, Australia

[11] State Key Laboratory of Respiratory Disease, Guangzhou Chest Hospital,

Guangzhou, China

[*] These authors contributed equally to this work.

[#] Corresponding author: Jinxing Hu (hujinxing2000@163.com) and Tianyu Zhang

(zhang_tianyu@gibh.ac.cn)

**ABSTRACT**

The efficacy of many compounds against *Mycobacterium tuberculosis* is often

limited when administered via conventional oral or injection routes due to

suboptimal pharmacokinetic characteristics. Inhalation delivery methods have been

investigated to achieve high local therapeutic doses in the lungs. However, previous

models, typically employing wild-type *M. tuberculosis* strains, were intricate, time-

consuming, labor-intensive, and with poor repeatability. In this study, we developed

an autoluminescence-based inhalation administration model to evaluate drug activity

by quantifying relative light units (RLUs) emitted from live mice infected with

autoluminescent *M. tuberculosis*. This novel approach has several improvements: it

eliminates the need for anesthesia in mice during administration and simplifies the

instrument manipulation; it is cost-effective by utilizing mice instead of larger

animals; it shortens time from several months to 16 or 17 days for obtaining result; it

is non-invasive by measuring the live RLUs of mice; up to six mice can be

administrated daily and simultaneously, even 2-3 times/day; results are relatively

objective and repeatable minimizing human factors. Proof-of-concept experiments

demonstrated that inhalable rifampicin, isoniazid, and ethambutol showed anti-*M.*

*tuberculosis* activity at concentrations as low as 0.5, 0.5, and 0.625 mg/mL,

respectively, as evidenced by comparing the live RLUs of mice. Furthermore,

consistency between RLUs and colony-forming units of the lungs reaffirms the

reliability of RLUs as an indicator of drug efficacy, highlighting the potential of this

approach for accurately assessing anti-*M. tuberculosis* activity *in vivo*. This

autoluminescence-based and non-invasive inhalation model offers a substantial

reduction in the time, effort, and cost required for evaluating the efficacy of

screening new drugs and repurposing old drugs *in vivo* via inhalation administration.

**KEYWORDS**

Inhalation administration, Autoluminescence, Tuberculosis, Murine model,

Chemotherapy

**INTRODUCTION**

*Mycobacterium tuberculosis*, primarily transmitted through respiratory droplets, is the causative agent of human tuberculosis (TB) and is responsible for the highest mortality rate attributed to a single infectious agent before the 2019 coronavirus pandemic (McQuaid et al., 2021, Shah et al., 2022). The emergence of drug-resistant *M. tuberculosis* strains presents substantial public health challenges and economic burdens (Dousa et al., 2020, Hameed et al., 2018, McGrath et al., 2014, Shoen et al., 2018). Therefore, there is an urgent need for innovative *in vivo* drugs screening and evaluation tools to identify efficacious antimicrobial agents for TB control.

Conventional *in vivo* efficacy evaluation experiments for anti-TB drugs often employ oral or various injection routes of administration (Namvarpour et al., 2019, Yu et al., 2020, Yu et al., 2021). However, these methods fail to specifically target the lungs and may induce significant toxicity in other organs during absorption, transport, and metabolism (Xu et al., 2019, Yoon et al., 2017, Zhao et al., 2023). Over several decades, inhalation therapy, delivering high local therapeutic doses to the lungs, has emerged as a preferred treatment method for respiratory diseases (Barnes et al., 2015, Garcia-Contreras et al., 2007, Gonzalez-Juarrero et al., 2012, Hickey et al., 2016, Sanchis et al., 2013). However, previous attempts to develop inhalable anti-TB drug administration models have encountered several challenges, including the time-consuming process of counting colony-forming units (CFUs), the complexities associated with anesthesia during administration, the difficulty in instrument operation, and the poor repeatability of experimental results (Garcia-

Contreras et al., 2010, Verma et al., 2013).

Meanwhile, most existing drugs evaluation models mentioned above are based on

wild-type *M. tuberculosis* and require several months for obtaining results, as it takes

3-6 weeks for the *M. tuberculosis* to form visible colonies on agar plates. Recently,

some researchers have introduced autoluminescent *M. tuberculosis* into the process

of drugs or vaccine evaluation (Yang et al., 2015, Zhang et al., 2012). The *luxAB*

genes encode the enzymes that catalyze the reaction resulting in the emission of

blue-green light, and *luxCDE* genes are responsible for the synthesis and recycling of

the aldehyde substrate (Hakkila et al., 2002). Light production by the

autoluminescent *M. tuberculosis* carrying the *luxCDABE* gene cluster has been

utilized as a surrogate of CFUs or visible bacterial growth for high-throughput

evaluation of antimycobacterial compounds *in vitro* and in *vivo* (Li et al., 2023,

Yang et al., 2015, Zhang et al., 2012). The combination of improved inhalation

delivery method and autoluminescent *M. tuberculosis* may offers a promising

approach for assessing drug activities.

Here, we have developed an autoluminescence-based, inhalable, time-efficient, non-

invasive, and cost-effective antimicrobial assessing model that provides a valuable

tool for evaluating the *in vivo* efficacy of lead compounds, frontline anti-TB

medications, and therapeutic regimens, thereby offering insights crucial for clinical

practice.

**MATERIALS AND METHODS**

**Mycobacterial strains and culture conditions**

The genetically engineered autoluminescent *M. tuberculosis* H37Rv (AlRv) strain,

which harbors the *luxCDABE* gene cluster within its genome, was preserved at -80□

(Yang et al., 2015). AlRv displays a consistent emission of blue-green light

independently, a characteristic attributed to the enzymatic activity encoded by the

*luxAB* genes. *luxCDE* genes are responsible for the continuous production of the

substrate aldehyde necessary for bioluminescence. Remarkably, AlRv demonstrates

drug susceptibility and growth rate comparable to that of the wild-type *M.*

*tuberculosis* strain. The traditional quantification method of CFUs was replaced by

the assessment of relative light units (RLUs), maintaining an approximate ratio of

10:1.

The AlRv was subcultured in Middlebrook 7H9 broth (Difco, Detroit, MI, USA),

supplemented with 10% oleic-acid-albumin-dextrose-catalase enrichment medium

(BBL, Sparks, MD, USA), 0.2% glycerol, and 0.05% Tween80 at 37□. The AlRv

culture was grown until the optical density, measured at 600 nm, reached 0.6-0.8,

with a corresponding RLUs level of $5 \times 10^6$/mL. The optical density was measured

using a V-1000 spectrophotometer (AOE, Shanghai, China), and the RLUs were

quantified using the GLOMAX 20/20 luminometer (Promega, Madison, USA).

**Antimicrobials**

All pharmaceutical drugs, including rifampicin (RIF), isoniazid (INH), and

ethambutol (EMB) were purchased from Meilun (Dalian, China) with a minimum

purity threshold of ≥ 95%. RIF and INH were solubilized in sterilized water, serving

as the solvent, to attain stock concentrations of 2 mg/mL, which were subsequently

diluted to 0.5 and 0.25 mg/mL, respectively. EMB was similarly dissolved into

sterilized water at its original concentration of 10 mg/mL, then further diluted to 2.5

and 0.625 mg/mL. The prepared solutions of the drugs were refrigerated at 4□ until

administration.

**Aerosol infection**

The female BALB/c mice (Gempharmatech, Foshan, China), aged six to eight weeks,

underwent an acclimatization period of five to seven days before the initiation of

experimental procedures. Subsequently, all mice were exposed to 10 mL broth

culture of AlRv using an inhalation exposure system (Glas-Col, Terre Haute, IN)

(Yu et al., 2020, Yu et al., 2021).

**Determination of the initial bacterial number**

After infection, the mice were randomly divided into treatment and control groups,

each containing six mice. On the initial day of treatment administration, anesthesia

was administered individually to six mice by placing them in a container with 0.5

mL of isoflurane. It took around ten seconds for the loss of consciousness.

Subsequently, the live RLUs of mice were measured by aligning their chests with the

detection position of the GLOMAX 20/20 luminometer. After detection of the live

RLUs, six mice were euthanized via cervical dislocation and positioned on a foam

board with large head pins to secure them, ensuring their abdomens were oriented

upwards. The mice were sterilized by spraying 75% ethanol and dissected to obtain

the lungs. The organs were then placed into 2 mL sterilized phosphate-buffered

saline, washed once with phosphate-buffered saline, and homogenized using glass

tissue grinders. The lung suspensions were diluted using a 10-fold serial dilution

method. Subsequently, the RLUs of 0.5 mL of undiluted and 10-fold diluted lung

suspensions were measured using the GLOMAX 20/20 luminometer. Based on the

detected RLUs and the ratio of CFUs to RLUs, the CFUs of the lung suspension

were briefly speculated upon. The 0.5 mL properly diluted lung suspension was

plated on 7H10 plates containing trimethoprim, actidione, carbenicillin, and

polymyxin B to prevent contamination. All plates were then incubated in a 37□

thermostatic incubator for four weeks, following which the CFUs were determined.

**Chemotherapy**

Each cohort of six mice underwent daily administration of either 4 mL of solvent or

drug solutions utilizing the Tow Systems Nose-Only Exposure Units (Tow-Int,

Shanghai, China), as previously described (Fan et al., 2022, Yang et al., 2021, Zhao

et al., 2023). The device of administration is illustrated in Figure S1. Mice were

subsequently secured in a holder and connected to the exposure cabinet. The drug or

solvent solution was injected into the atomizer, which uses high-frequency

oscillation technology through microporous screens to generate nebulized aerosols

with diameters ranging from 1 to 5 micrometers for inhalation. The drug exposure

duration for each group was approximately 20-25 minutes until 4 mL of the

solutions had been administered.

The study consists of two main experiments, delineated as Experiment 1 and

Experiment 2. The experimental protocols remained largely consistent, with the

primary variations involving alterations in pharmacological dosages and the timing

of medication administration subsequent to infection. Detailed protocols for

Experiments 1 and 2 were provided in Tables 1 and 2, respectively. The live RLUs

of the solvent group were continuously monitored from the day 11 for Experiment 1

(or day 13 for Experiment 2) post-infection. Measurements of RLUs for all groups

were performed once the live RLUs of the solvent-treated mice exceeded 400. Based

on our experiences, live RLUs exceeding 400 indicated a higher level of lung RLUs

and CFUs, suitable for evaluating drug efficacy. Treatment of mice was discontinued

upon reaching live RLUs exceeding 400. The total duration of administration for

Experiments 1 and 2 was 14 and 15, respectively. Subsequently, all mice were

euthanized the following day to determine lung RLUs and CFUs following the

method mentioned above. The plates were then incubated at 37□ for four weeks, and

then CFUs were determined.

**Statistical analysis**

Log-transformed data using GraphPad Prism version 8.3.0 was used to determine

drug efficacy through a two-way analysis of variance, with $P < 0.05$ as statistical

significance.

**RESULTS**

**The results of experiment 1.**

In experiment 1, the initial CFUs of the infected bacteria were approximately 4000.

The live RLUs of all treatment groups were significantly lower than those of the

solvent group, with all differences exceeding 200 (Figure 1, $P < 0.0001$).

Additionally, these live RLUs were comparable to those of the before-treatment (BT)

group (Figure 1, $P > 0.05$). Furthermore, the lung RLUs of all drug-treated groups

were substantially lower than those of the solvent group, with all $\log_{10}$RLUs/lung

differences exceeding 1.5 (Figure 1, $P < 0.0001$). Specifically, the lung RLUs of the

RIF or EMB-treated groups were nearly identical to those of the BT group (Figure 1,

$P > 0.05$), while the INH-treated group exhibited slightly lower lung RLUs than the

BT group (Figure 1, $P < 0.05$). Moreover, the lung CFUs of all treatment groups

were markedly lower than that of the solvent group, as all differences were over 1.5

(Figure 1, $P < 0.0001$). The RIF group had higher lung CFUs, whereas the INH

group had lower lung CFUs compared to the BT group (Figure 1, $P < 0.01$ and $P <$

0.0001, respectively). The lung CFUs of the EMB-treated group were almost similar

to those of the BT group (Figure 1, $P > 0.05$). The data from experiment 1 indicated

that the autoluminescence-based inhalation delivery method can be utilized to

establish a non-invasive model for testing drugs activities *in vivo*, using the same

batch of live mice within 18 days after infection. Furthermore, the effectiveness of

drugs can be further evaluated as either bacteriostatic or bactericidal by comparing

lung CFUs of the different treatment groups rather than RLUs, as lung RLUs of the

BT group were close to the background values.

**The results of experiment 2.**

Based on the results of Experiment 1, two additional dosages of each drug were

added in Experiment 2. The initial CFUs of the infected bacteria in Experiment 2

were found to be roughly equivalent to those recorded in Experiment 1. The live

RLUs of the groups treated with RIF 2, INH 2, INH 0.5, EMB 10, EMB 2.5, and

EMB 0.625 mg/mL were significantly lower compared to the solvent group (Figure

2a, $P < 0.0001$, $P < 0.0001$, $P < 0.01$, $P < 0.001$, $P < 0.001$, and $P < 0.05$,

respectively). Conversely, the live RLUs of RIF 0.5, RIF 0.125 and INH 0.125

mg/mL groups did not show significant differences from those of the solvent group

(Figure 2a, $P > 0.05$). These findings were also largely supported by both the lung

RLUs and CFUs (Figure 2b and 2c), except for the lung CFUs of the RIF at 0.5

mg/mL group, which were slightly significantly lower than that of the solvent group

(Figure 2c, $P < 0.05$). In summary, these findings suggest a negative correlation

between live RLUs and administered dosages, indicating that live RLUs could serve

as an initial criterion for assessing the therapeutic efficacy.

The lung RLUs of the RIF 2 mg/mL group demonstrated a significant reduction

compared to those in the solvent group, with a $\log_{10}$RLUs/lung difference exceeding

1.5 and even similar to that of the BT group (Figure 2b, $P < 0.0001$ and $P > 0.05$,

respectively). Conversely, the lung RLUs in the RIF 0.5 and 0.125 mg/mL groups

were comparable to that in the solvent group, indicating a lack of evident anti-*M.*

*tuberculosis* activity at these dosages (Figure 2b, $P > 0.05$). The lung RLUs of the

INH 2 and 0.5 mg/mL groups were significantly lower than that of the solvent group

(Figure 2b, $P < 0.0001$), as all differences in $\log_{10}$RLUs/lung exceeded 1.5 and were

comparable to that of the BT group (Figure 2b, $P > 0.05$). The lung RLUs of the INH

0.125 mg/mL group were similar to those of the solvent group (Figure 2b, $P > 0.05$).

In comparison, the lung RLUs of the EMB 10 and 2.5 mg/mL groups exhibited a

pronounced decrease compared to those in the solvent group, with $\log_{10}$RLUs/lung

differences exceeding 1.5 and 1, respectively (Figure 2b, $P < 0.0001$). Meanwhile,

the lung RLUs of the EMB 10 mg/mL group were equivalent to that of the BT group

(Figure 2b, $P > 0.05$), whereas the lung RLUs in the EMB 0.625 mg/mL group

showed less significant differences from the solvent group (Figure 2b, $P < 0.05$). In

conclusion, the lung RLUs results are consistent with those obtained from the live

RLUs, providing further support for the utility of detecting live RLUs in determining

drug activity.

The lung CFUs in the RIF 2 mg/mL group exhibited a significant reduction

compared to those in the solvent group with a difference in $\log_{10}$CFUs/lung

exceeding 1.5 (Figure 2c, $P < 0.0001$). The lung CFUs in the RIF 0.5 and 0.25

mg/mL groups were approximately comparable to the solvent group (Figure 2c, $P <$

and close to 0.05 and $P > 0.05$, respectively). Notably, no colonies were observed on

the plates of the INH 2 mg/mL group, indicating possibly eradication of AlRv from

the mice lungs, considering that only a fraction of the lung suspension from each

mouse was plated. The CFUs of the INH 0.5 mg/mL group were also significantly

lower than that of the solvent group, with a difference of $\log_{10}$CFUs/lung exceeding

1.5 and even lower than the BT group (Figure 2c, $P < 0.0001$). Moreover, all EMB

dosages resulted in significantly reduced lung CFUs compared to the solvent control

(Figure 2c, $P < 0.0001$, $P < 0.0001$, and $P < 0.001$, respectively). However, only the

$\log_{10}$CFUs/lung difference between the EMB 10 mg/mL and solvent group exceeded

1.5, with the lung CFUs in the EMB 10 group being nearly identical to those in the

BT group (Figure 2c, $P > 0.05$). These CFU results were consistent with the findings

derived from live or lung RLUs data. Overall, based on the comprehensive

experimental data presented, we have successfully established an autoluminescence-

based inhalation administration murine model for evaluating drugs activities against

*M. tuberculosis in vivo*.

**DISCUSSION**

*M. tuberculosis* imposes a substantial burden, particularly in developing countries

(Chong and Lim, 2009, Grüber, 2020). Despite ongoing efforts, the development of

effective anti-*M. tuberculosis* drugs remains confronted with numerous challenges,

including the complexity of evaluation process and the emergence of drug-resistant

strains (McGrath et al., 2014, Olaru et al., 2015). Over the past several decades,

inhalable therapy has emerged as a promising approach for treating pulmonary

infections, offering improved cure rates and bacterial eradication (Riveiro et al.,

2023). However, the evaluation of inhalable drugs still presents challenges such as

complex manipulation and time-consuming results completion. Recent researches

have revealed that a recombinant autoluminescent strain of *M. tuberculosis* carrying

*luxCDABE* gene cluster facilitates rapid, non-invasive, real-time monitoring of the

RLUs in live mice during experimental chemotherapy, providing a valuable tool for

assessing treatment efficacy (Yang et al., 2015, Zhang et al., 2012). Hence, we have

developed a novel murine model by combining the improved inhalation method and

autoluminescent *M. tuberculosis*.

As of the present, four primary research reports, encompassing three inhalation

administration models, have explored the detection of anti-TB drug efficacy through

airway administration (Garcia-Contreras et al., 2007, Garcia-Contreras et al., 2010,

Gonzalez-Juarrero et al., 2012, Verma et al., 2013). Among these studies, only two

have employed murine models for their investigations. Table S1 presents a

comprehensive comparison of the advantages and disadvantages of each report

relative to our study. Compared to these existing inhalation administration model,

our innovative method achieves several improvements. Firstly, it is cost-effective

and user-friendly, employing mice instead of larger animals. Secondly, it eliminates

the need for anesthesia in mice during administration, simplifying the instrument

operation. We also attempted to employ published administration method in mice

infected with AlRa via tail vein injection, but encountered significant challenges in

administering drugs daily due to the mice developing swollen throats (Gonzalez-

Juarrero et al., 2012). However, we did observe that mice treated with INH exhibited

reduced RLUs compared to both the untreated control and the day before

administration (unpublished data). Thirdly, the results are relatively objective and

repeatable, minimizing human factors. Fourthly, the same treatment group,

consisting of six mice, is administrated daily and simultaneously. In fact, we indeed

succeeded in administrating the same six mice 2-3 times/day and even with different

drugs. Additionally, our method is non-invasive, measuring the live RLUs of mice as

a surrogate marker for CFUs for drug *in vivo* activity testing. Lastly, it reduces the

time required for obtaining results from several months to 16 or 17 days.

We have evaluated the *in vivo* efficacy of three first-line TB drugs using our model,

including RIF, INH, and EMB. Our model can accelerate assessment of drug activity,

with results obtainable within 17 or 18 days or even shorter by detecting either live

RLUs of mice or RLUs of lung suspensions. Our findings demonstrate that RIF, INH,

and EMB, at the dosage of 0.5, 0.5, and 0.625 mg/mL, exhibited significant anti-*M.*

*tuberculosis* activity *in vivo*. Notably, INH exhibits potential for complete

eradication of lung bacteria at increased dosage up to 2 mg/mL in the current model.

Moreover, the CFUs results corroborate the findings inferred from live RLUs,

affirming the utility of live RLUs as a surrogate marker for CFUs in mice. This

innovative model holds promise for enhancing the assessment of anti-*M.*

*tuberculosis* drug efficacy *in vivo*, thereby reducing the likelihood of overlooking

potentially effective treatments and minimizing the risk of adverse effects.

The drug's activity can be determined by comparing the lung suspension RLUs of

different groups within a shorter duration than the current 17-day period (from

infection to get the results), since the lung RLUs become significantly higher than

the background value a few days earlier than the live RLUs do. Furthermore, drug

efficacy could potentially be observed within a considerably shorter time frame in

live mice if mice infected with high dose AlRv via tail vein injection route, similar to

the previous studies showed that the live RLUs could be detected at the day after

infection, although this may result in a higher standard deviation (Liu et al., 2019,

Olaru et al., 2015, Zhang et al., 2012). Further investigation is warranted to

determine whether other anti-TB drugs or potent compounds demonstrate efficacy

using our model. Additionally, exploring alternative formulations or solvents may

further enhance the drug's effectiveness of drugs beyond sterilized water.

In conclusion, our study establishes an autoluminescence-based, inhaled, and non-invasive administration murine model for evaluating anti-*M. tuberculosis* drug *in vivo* efficacy. This model holds promise for expediting the development of innovative anti-*M. tuberculosis* drugs and treatment regimens, addressing the urgent need for more effective TB therapies. Meanwhile, this improved inhalation model can also be used for developing drugs targeting other diseases and assessing the toxicity of compounds.

**DATA AVAILABILITY STATEMENT**

The original contributions presented in the study are included in the article, further inquiries can be directed to the corresponding authors.

**ETHICS STATEMENT**

The animal care and experimental protocols were approved by the Laboratory Animal Ethics Committee of the Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences.

**AUTHOR CONTRIBUTIONS**

Conceived the project: Tianyu Zhang, Jinxing Hu, Xirong Tian, Yamin Gao; Designed the research: Tianyu Zhang, Jinxing Hu, Xirong Tian, Yamin Gao; Performed the studies: Xirong Tian, Yamin Gao, Jingran Zhang, Yanan Ju, Jie Ding, Sanshan Zeng, Wanli Ma; Interpreted the results: Tianyu Zhang, Xirong Tian, Yamin Gao; Drafted the manuscript: Tianyu Zhang, Jinxing Hu, Xirong Tian, Yamin Gao, H.M. Adnan Hameed, Nanshan Zhong, Gregory M. Cook; Final approval of the manuscript: all authors.

**FUNDING**

This work was supported by the National Key R&D Program of China

(2021YFA1300900; 2023YFFO713605), the National Natural Science Foundation

of China (81973372, 21920102003, 82061138019, 32300152), Guangdong

Provincial Basic and Applied Basic Research Fund (2022A1515110505),

Guangzhou Science and Technology (2024A04J4273), Chinese Academy of

Sciences (CAS)-President's International Fellowship Initiative (2023VBC0015),

China Postdoctoral Science Foundation (2022M723164), the Chinese Academy of

Sciences Grants (YJKYYQ20210026), State Key Laboratory of Respiratory Disease

(SKLRD-Z-202301, SKLRD-Z-202414), and Guangzhou Science and Technology

Planning Project (2023A03J0992).

**REFERENCES**

Barnes PJ, Burney PGJ, Silverman EK, Celli BR, Vestbo J, Wedzicha JA, et al.

(2015). Chronic obstructive pulmonary disease. *Nat. Rev. Dis. Primers*. 1, 15076.

Chong VH, Lim KS. (2009). Gastrointestinal tuberculosis. *Singapore. Med. J.* 50(6),

638-645.

Dousa KM, Kurz SG, Bark CM, Bonomo RA, Furin JJ. (2020). Drug-resistant

tuberculosis: A glance at progress and global challenges. *Infect. Dis. Clin. North. Am*.

34(4), 863-886.

Fan Z, Zhou B, Liu Y, Sun W, Fang Y, Lu H, et al. (2022). Optimization and

application of an efficient and stable inhalation exposure system for rodents. *AAPS.*

*PharmSciTech*. 23(1), 50.

Garcia-Contreras L, Fiegel J, Telko MJ, Elbert K, Hawi A, Thomas M, et al.(2007).

Inhaled large porous particles of capreomycin for treatment of tuberculosis in a

guinea pig model. *Antimicrob. Agents. Chemother.* 51(8), 2830-2836.

Garcia-Contreras L, Sung JC, Muttil P, Padilla D, Telko M, Verberkmoes JL, et al.

(2010). Dry powder PA-824 aerosols for treatment of tuberculosis in guinea pigs.

*Antimicrob. Agents. Chemother.* 54(4), 1436-1442.

Gonzalez-Juarrero M, Woolhiser LK, Brooks E, DeGroote MA, Lenaerts AJ. (2012).

Mouse model for efficacy testing of antituberculosis agents via intrapulmonary

delivery. *Antimicrob. Agents. Chemother.* 56(7), 3957-3959.

Grüber G.(2020). Introduction: Novel insights into TB research and drug discovery.

*Prog. Biophys. Mol. Biol.* 152, 2-5.

Hakkila K, Maksimow M, Karp M, Virta M. (2002). Reporter genes *lucFF*,

*luxCDABE*, *gfp*, and *dsred* have different characteristics in whole-cell bacterial

sensors. *Anal. Biochem.* 301(2), 235-242.

Hameed HMA, Islam MM, Chhotaray C, Wang C, Liu Y, Tan Y, et al. (2018).

Molecular targets related drug resistance mechanisms in MDR-, XDR-, and TDR-

*Mycobacterium tuberculosis* strains. *Front. Cell. Infect. Microbiol.* 8, 114.

Hickey AJ, Durham PG, Dharmadhikari A, Nardell EA. (2016). Inhaled drug

treatment for tuberculosis: Past progress and future prospects. *J. Control. Release.*

240, 127-134.

Li C, Tian X, Huang Z, Gou X, Yusuf B, Li C, et al. (2023). Structure-activity

relationship of novel pyrimidine derivatives with potent inhibitory activities against

*Mycobacterium tuberculosis*. *J. Med. Chem.* 66(4), 2699-2716.

Liu Y, Gao Y, Liu J, Tan Y, Liu Z, Chhotaray C, et al. (2019). The compound TB47 is

highly bactericidal against *Mycobacterium ulcerans* in a Buruli ulcer mouse model.

*Nat. Commun.* 10(1), 524.

McGrath M, Gey van Pittius NC, van Helden PD, Warren RM, Warner DF. (2014).

Mutation rate and the emergence of drug resistance in *Mycobacterium tuberculosis*. *J.*

*Antimicrob. Chemother.* 69(2), 292-302.

McQuaid CF, Vassall A, Cohen T, Fiekert K, White RG. (2021). The impact of

COVID-19 on TB: A review of the data. *Int. J. Tuberc. Lung. Dis.* 25(6), 436-446.

Namvarpour M, Tebianian M, Mansouri R, Ebrahimi SM, Kashkooli S. (2019).

Comparison of different immunization routes on the immune responses induced by

*Mycobacterium tuberculosis* ESAT-6/CFP-10 recombinant protein. *Biologicals.* 59,

6-11.

Olaru ID, von Groote-Bidlingmaier F, Heyckendorf J, Yew WW, Lange C, Chang

KC. (2015). Novel drugs against tuberculosis: a clinician's perspective. *Eur. Respir. J.*

45(4), 1119-1131.

Riveiro V, Casal A, Abelleira R, Valdés L. (2023). Advances in inhaled antibiotics

for management of respiratory tract infections. *Curr. Opin. Pulm. Med.* 29(3), 160-

167.

Sanchis J, Corrigan C, Levy ML, Viejo JL. (2013). Inhaler devices - from theory to

practice. *Respir. Med.* 107(4), 495-502.

Shah T, Shah Z, Yasmeen N, Baloch Z, Xia X. (2022). Pathogenesis of SARS-CoV-2

and *Mycobacterium tuberculosis* coinfection. *Front. Immunol*. 13, 909011.

Shoen C, DeStefano M, Hafkin B, Cynamon M. (2018). *In vitro* and *in vivo* activities

of contezolid (MRX-I) against *Mycobacterium tuberculosis*. *Antimicrob. Agents.*

*Chemother*. 62(8), e00493-18.

Verma RK, Germishuizen WA, Motheo MP, Agrawal AK, Singh AK, Mohan M, et al.

(2013). Inhaled microparticles containing clofazimine are efficacious in treatment of

experimental tuberculosis in mice. *Antimicrob. Agents. Chemother*. 57(2), 1050-

1052.

Xu J, Wang B, Fu L, Zhu H, Guo S, Huang H, et al. (2019) *In vitro* and *in vivo*

activities of the riminophenazine TBI-166 against *Mycobacterium tuberculosis*.

*Antimicrob. Agents. Chemother*. 63(5), e02155-18.

Yang F, Njire MM, Liu J, Wu T, Wang B, Liu T, et al. (2015). Engineering more

stable, selectable marker-free autoluminescent mycobacteria by one step. *PLoS. One*.

10(3), e0119341.

Yang X, Liang J, Wang Z, Su Y, Zhan Y, Wu Z, et al. (2021). Sesamolin protects

mice from ovariectomized bone loss by inhibiting osteoclastogenesis and RANKL-

mediated NF-κB and MAPK signaling pathways. *Front. Pharmacol*. 12, 664697.

Yoon HK, Park Y-B, Rhee CK, Lee JH, Oh Y-M. (2017). Summary of the chronic

obstructive pulmonary disease clinical practice guideline revised in 2014 by the

Korean academy of tuberculosis and respiratory disease. *Tuberc. Respir. Dis. (Seoul).*

80(3), 230-240.

Yu W, Chiwala G, Gao Y, Liu Z, Sapkota S, Lu Z, et al. (2020). TB47 and

clofazimine form a highly synergistic sterilizing block in a second-line regimen for

tuberculosis in mice. *Biomed. Pharmacother*. 131, 110782.

Yu W, Yusuf B, Wang S, Tian X, Hameed HMA, Lu Z, et al. (2021). Sterilizing

effects of novel regimens containing TB47, clofazimine, and linezolid in a murine

model of tuberculosis. *Antimicrob. Agents. Chemother*. 65(10), e0070621.

Zhang T, Li S-Y, Nuermberger EL. (2012). Autoluminescent *Mycobacterium*

*tuberculosis* for rapid, real-time, non-invasive assessment of drug and vaccine

efficacy. *PLoS. One*. 7(1), e29774.

Zhao G, Ho W, Chu J, Xiong X, Hu B, Boakye-Yiadom KO, et al. (2023). Inhalable

siRNA nanoparticles for enhanced tumor-targeting treatment of KRAS-mutant non-

small-cell lung cancer. *ACS. Appl. Mater. Interfaces*. 15(26), 31273-31284.

**Table 1. Scheme of experiment 1.**

| Drug/Dosage[1] (mg/mL) | Number of mice sacrificed at the given time points | | | | Total |
|---|---|---|---|---|---|
| | D0[2] | D3[3] | D11-17 | D18[4] | |
| Solvent | | 6 | Detecting chest | 6 | 12 |
| RIF 2 | | Treatments | RLUs of live mice | 6 | 6 |
| INH 2 | | initiation | in solvent group | 6 | 6 |
| EMB 10 | Infection | | daily from day 11 | 6 | 6 |
| | | | until live RLUs | | |
| | | | exceed 400 | | |
| Total | 30 | 6 | | 24 | 30 |

[1] RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the

number after the drug represents the dosage.

[2] D0, a total of 30 mice were infected with AlRv by aerosol.

[3] D3, the mice were randomly divided into groups and treated daily from the day 3 to

day 16. The live RLUs, the lung suspension RLUs and CFUs of six mice were

detected before the treatment initiation.

[4] D18, the mice were sacrificed and the lungs were removed for detecting RLUs and

CFUs.

All CFUs were counted 4 weeks after incubation at 37□.

**Figure 1. Anti-TB drug activities by inhalation administration in Experiment 1.**

BT, before treatment; RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the number after the drug represents the

dosage; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; ns, $P > 0.05$, no significance. The dotted lines represent the background

values.

**Table 2. Scheme of experiment 2.**

| Drug/Dosage[1] (mg/mL) | Number of mice sacrificed at the given time points | | | | Total |
| --- | --- | --- | --- | --- | --- |
| | D0[2] | D1[3] | D13-16 | D17[4] | |
| Solvent | | 6 | | 6 | 12 |
| RIF 2 | | | | 6 | 6 |
| RIF 0.5 | | | Detecting chest | 6 | 6 |
| RIF 0.125 | | | RLUs of live mice | 6 | 6 |
| INH 2 | | | in solvent group | 6 | 6 |
| INH 0.5 | Infection | Treatments initiation | daily from day 13 | 6 | 6 |
| INH 0.125 | | | until live RLUs | 6 | 6 |
| EMB 10 | | | exceed 400 | 6 | 6 |
| EMB 2.5 | | | | 6 | 6 |
| EMB 0.625 | | | | 6 | 6 |
| Total | 66 | 6 | | 60 | 66 |

[1] RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the number after the drug represents the dosage.

[2] D0, a total of 66 mice were infected with AlRv by aerosol.

[3] D1, the mice were randomly divided into groups and treated daily from day 1 to day 15. The live RLUs, the lung suspension RLUs and CFUs of six mice were detected before the treatment initiation.

[4] D17, the mice were sacrificed, and the lungs were removed for detecting RLUs and CFUs.

All CFUs were counted 4 weeks after incubation at 37□.

**Figure 2. Anti-TB drug activities by inhalation administration in Experiment 2.**

(a) Light intensities of live mice. (b) The suspension light intensities. (c) The lung *M. tuberculosis* burden.

BT, before treatment; RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the number after the drug represents the dosage; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; ns, $P > 0.05$, no significance. The dotted lines represent the background values.