Establishment of an Inhalation Administration Noninvasive Murine Model for

Rapidly Testing Drug Activity against *Mycobacterium tuberculosis*

Xirong Tian [1,2,3,4,*], Yamin Gao [1,2,3,4,*], Wanli Ma [1,2,3,4], Jingran Zhang [1,2,3,5], Yanan Ju

[1,2,3,5], Jie Ding [1,2,3, 6], Sanshan Zeng [1,2,3,4], H.M. Adnan Hameed [1,2,3,4], Nanshan Zhong

[7], Gregory M. Cook [4,8,9,10], Jinxing Hu [7, 11, #] and Tianyu Zhang [1,2,3,4,7,#]

[1] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine

and Health (GIBH), Chinese Academy of Sciences (CAS), Guangzhou 510530,

China, [2] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious

Diseases, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese

Academy of Sciences (CAS), Guangzhou 510530, China, [3] University of Chinese

Academy of Sciences (UCAS), Beijing 100049, China, [4] China-New Zealand Joint

Laboratory on Biomedicine and Health, Guangzhou Institutes of Biomedicine and

Health, Chinese Academy of Sciences, Guangzhou 510530, China, [5] School of Life

Sciences, University of Science and Technology of China, Hefei 230026, China, [6]

Institute of Physical Science and Information Technology, Anhui University, Hefei,

230601, China, [7] Guangzhou National Laboratory, Guangzhou 510005, China, [8]

Department of Microbiology and Immunology, School of Biomedical Sciences,

University of Otago, Dunedin, 9054, New Zealand, [9] Maurice Wilkins Centre for

Molecular Biodiscovery, University of Auckland, Private Bag 92019, Auckland,

1042, New Zealand, [10] Translational Research Institute, Queensland University of

Technology, Brisbane, Queensland, 4102, Australia, [11] State Key Laboratory of

Respiratory Disease, Guangzhou Chest Hospital, Guangzhou, China

* These authors contributed equally to this work.

Corresponding author: Jinxing Hu (hujinxing2000@163.com) and Tianyu Zhang

(zhang_tianyu@gibh.ac.cn)

ABSTRACT

Mycobacterium tuberculosis, as one of the most serious infectious pathogens, results

in millions of deaths and a tremendous economic burden for both individuals and

governments. Many potential drugs for the treatment of tuberculosis show excellent

activity against *M. tuberculosis in vitro*, but lack significant activity *in vivo* when

administered traditionally by either oral or injection delivery routes. Even though

previous researches revealed that inhaled administration method can enhance the

efficacy of drugs against mycobacteria, while this evaluating process was

complicated and time-consuming by using wild type *M. tuberculosis*. Conversely,

our research uncovered that the inhalation administration model using strains of

autoluminescent *M. tuberculosis* may provide a more rapid, non-invasive, lung-

targeted mid-throughput method, may avoid significant toxic effects on other organs,

for evaluating the anti-tuberculosis activity of compounds in mouse models of

infection. The live relative light units, quantifying the light emitted by

autoluminescent *M. tuberculosis* $H_{37}Rv$, can be used to preliminarily evaluate the

therapeutic effect of different drug doses, which can significantly shorten the

evaluation time. This model offers the ability to assess the efficacy of new lead

compounds and TB drugs that can be administered by inhalation.

KEYWORDS

Inhalation administration, Autoioluminescence, Tuberculosis, Murine model,

Chemotherapy

INTRODUCTION

Mycobacterium tuberculosis (Mtb) primarily transmitted via respiratory droplets, stands as the causative agent of human tuberculosis (TB) and accounted for the highest mortality attributed to a single infectious agent prior to the 2019 coronavirus pandemic (McQuaid et al., 2021, Shah et al., 2022). Approximately one quarter of the global population is latently infected with Mtb, with an estimated 5-10% progressing to active TB over their lifetimes (Ahmad et al., 2022). The rise in prevalence of drug-resistant Mtb strains is largely attributable to antibiotic misuse and patient non-compliance, posing substantial public health risks and imposing considerable economic burdens (Dousa et al., 2020, Hameed et al., 2018, McGrath et al., 2014, Shoen et al., 2018). Consequently, there is a pressing demand for novel anti-TB medications characterized by enhanced efficacy, minimal toxicity, and a lack of cross-resistance with existing therapies to address these challenges (Grüber, 2020, Liu Pingxian et al., 2019).

In conventional *in vivo* efficacy evaluation experiments for anti-tuberculosis drugs, oral or various injection routes of administration are frequently employed (Namvarpour et al., 2019, Yu et al., 2020, Yu et al., 2021). However, these traditional administration methods fail to specifically target the lungs during absorption, transport, and metabolism, while also inducing significant toxicity in other organs (Xu et al., 2019, Yoon et al., 2017, Zhao et al., 2023). Over recent decades, inhalation therapy has become a preferred treatment method for respiratory diseases (Barnes et al., 2015, Sanchis et al., 2013). Despite previous attempts to

develop models for assessing inhaled drug administration, these models encounter

various challenges necessitating further improvement.

The lung-targeted inhalation drug delivery method devised in this study effectively

overcomes the aforementioned limitations. Drug activities administered via

inhalation can be assessed in live mice infected with autoluminescent Mtb H37Rv

(AlRv) in a time-efficient, non-invasive, and cost-effective mid-throughput manner.

This model represents a valuable tool for evaluating the efficacy of inhalable lead

compounds, frontline anti-TB medications, and therapeutic regimens, thereby

providing clinicians with invaluable insights.

MATERIALS AND METHODS

Mycobacterial strains and culture conditions

The AlRv strain, genetically engineered to contain the *luxCDABE* gene cluster

within its genome, was preserved at -80 □ (Zhang et al., 2012). AlRv displays a

consistent emission of blue-green light independently, attributed to the enzymatic

activity encoded by the *luxCDE* genes, which continuously produces the necessary

substrate aldehyde (Zhang et al., 2012). Remarkably, AlRv demonstrates a

susceptibility to drugs comparable to that of the wild type Mtb H37Rv strain (Zhang

et al., 2012). The quantification of colony forming units (CFUs) was replaced with

the assessment of relative light units (RLUs), maintaining a ratio of approximately

10:1 (Zhang et al., 2012).

The AlRv was subcultured in Middlebrook 7H9 broth (Difco, Detroit, MI, USA),

supplemented with 10% oleic-acid-albumin-dextrose-catalase enrichment medium

(BBL, Sparks, MD, USA), 0.2% glycerol, and 0.05% Tween 80 at 37 □. The AlRv

was cultured until the optical density, measured at 600 nm, reached 0.6-0.8, with a

corresponding RLUs level of 5×10^6/mL. The optical density was measured using a

V-1000 spectrophotometer (AOE, Shanghai, China), and the RLUs were quantified

using the GLOMAX 20/20 luminometer (Promega, Madison, USA).

Antimicrobials

All pharmaceutical compounds, including rifampicin (RIF), isoniazid (INH) and

ethambutol (EMB) were purchased from Meilun (Dalian, China) with a minimum

purity threshold of $\geq$ 95%. RIF and INH were solubilized in sterilized water, serving

as the solvent, to attain stock concentrations of 2 mg/mL, which were subsequently

diluted to 0.5 and 0.25 mg/mL, respectively. EMB were similarly dissolved into

sterilized water at its original concentration of 10 mg/mL, then further diluted to 2.5

and 0.625 mg/mL. The prepared solutions of the drugs were refrigerated at 4°C until

administration.

Aerosol infection determination of infection bacterial burden

The female BALB/c mice (Gempharmatech, Foshan, China), aged six to eight weeks,

were subjected to an acclimatization period lasting five to seven days before the

initiation of experimental procedures. Subsequently, all mice were exposed to

homogenous AlRv culture of 10 mL using an inhalation exposure system (Glas-Col,

Terre Haute, IN) (Yu et al., 2020, Yu et al., 2021). The total exposure duration was

two hours, comprising preheating (15 minutes), atomization (60 minutes), defogging

(15 minutes), and disinfection (25 minutes) procedures.

Determination of the initial bacterial number

After infection, the mice were randomly divided into treatment and solvent groups, each consisting of six individuals. On the initial day of treatment administration, anesthesia was administered individually to six mice via exposure to a container containing 0.5 mL of isoflurane. Loss of consciousness typically occurred within approximately 10 seconds. Subsequently, the live RLUs of mice were measured by aligning their chests with the detection position of the GLOMAX 20/20 luminometer. Upon detection of the live RLUs, six mice were euthanized via cervical dislocation and positioned on a foam board with large head pins to secure them, ensuring their abdomens were oriented upwards. The mice were sterilized by spraying 75% ethanol and dissected to obtain the lungs. The organs were then placed into 2 mL sterilized phosphate-buffered saline, washed once with phosphate-buffered saline, and homogenized using glass tissue grinders. The lung suspensions were diluted using a 10-fold serial dilution method. Subsequently, the RLUs of 0.5 mL of undiluted and 10-fold diluted lung suspensions were measured using the GLOMAX 20/20 luminometer. Based on the detected RLUs and the ratio of CFUs to RLUs, the CFUs of the lung suspension were briefly speculated upon. The o.5 mL properly diluted lung suspension was plated on 7H10 plates containing trimethoprim, actidione, carbenicillin and polymyxin B to prevent contamination. All plates were then incubated in a 37□ thermostatic incubator for four weeks, following which the CFUs were determined.

Chemotherapy

Each group of six mice received daily administration of either 4 mL of solvent or

drug solutions using the Tow Systems Nose-Only Exposure Units (Tow-Int,

Shanghai, China), as previously described (Fan et al., 2022, Yang et al., 2021, Zhao

et al., 2023). The device of administration is illustrated in Figure 1. Subsequently,

each mouse was secured using the holder and connected to the exposure cabinet. The

drug solution or solvent were injected into the atomizer, which utilizes high

frequency oscillation technology through microporous screens to generate nebulized

aerosols with diameters ranging from 1 to 5 micrometers for inhalation. The drug

exposure duration for each group spanned approximately 20-25 minutes, until 4 mL

of the solutions had been administered.

The study consists of two main experiments, labeled as Experiment 1 and

Experiment 2. The experimental protocols remained largely consistent, with the

primary variations involving adjustments in drug dosages and the timing of drug

administration post-infection. Detailed protocols for Experiment 1 and 2 were

provided in Tables 1 and 2, respectively. The live RLUs of the solvent group were

continuously monitored from the day 11 for Experiment 1 (or day 13 for Experiment

2) post-infection. Once the live RLUs of the solvent -treated mice exceeded 400, the

RLUs of all groups were measured. Based on our experiences, live RLUs exceeding

400 indicated a higher level of lung RLUs and CFUs, suitable for evaluating drug

efficacy. Treatment of mice was discontinued upon reaching live RLUs exceeding

400. The total duration of administration for Experiment 1 and 2 were 14 and 15,

respectively. Subsequently, all mice were euthanized the following day to determine

lung RLUs and CFUs following the method mentioned above. All plates were then

incubated in a 37□ thermostatic incubator for four weeks, after which CFUs were

determined.

Statistical analysis

Log-transformed data using GraphPad Prism version 8.3.0 was used to determine

drug efficacy through a two-way analysis of variance with $P < 0.05$ as statistical

significance.

RESULTS

The results of experiment 1.

In experiment 1, the initial CFUs of the infected bacteria approached approximately

4000. The live RLUs of all treatment groups were significantly lower than that of the

solvent group, with all differences exceeding 200 (Figure 1, $P < 0.0001$).

Additionally, these live RLUs were comparable to those of the before-treatment (BT)

group (Figure 1, $P > 0.05$). Furthermore, the lung RLUs of all drug-treated groups

were substantially lower than those of the solvent group, with all $log_{10}RLUs/lung$

differences exceeding 1.5 (Figure 1, $P < 0.0001$). Specifically, the lung RLUs of the

RIF or EMB-treated groups were nearly identical to those of the BT group (Figure 1,

$P > 0.05$), while the INH-treated group exhibited slightly lower lung RLUs than the

BT group (Figure 1, $P < 0.05$). Moreover, the lung CFUs of all treatment groups

were markedly lower than that of the solvent group, as all differences were over 1.5

(Figure 1, $P < 0.0001$). The RIF group had higher lung CFUs, whereas the INH

group had lower lung CFUs compared to the BT group (Figure 1, $P < 0.01$ and $P <$

0.0001, respectively). The lung CFUs of EMB-treated group were almost similar to

those of the BT group (Figure 1, $P > 0.05$). The data from experiment 1 indicated

that the autoluminescence-based inhalation delivery method can be utilized to

establish a noninvasive model for testing drugs activities *in vivo*, using the same

batch of live mice within 18 days after infection. Furthermore, the effectiveness of

drugs can be further evaluated as either bacteriostatic or bactericidal by comparing

lung CFUs of the different treatment groups rather than RLUs, as lung RLUs of the

BT group were close to the background values.

The results of experiment 2.

Based on the findings of Experiment 1, two additional dosages of each drug were

added in Experiment 2. The initial CFUs of the infected bacteria, in the Experiment

2, were found to be approximately equivalent to those recorded in Experiment 1. The

live RLUs of the groups treated with RIF 2, INH 2, INH 0.5, EMB 10, EMB 2.5 and

EMB 0.625 mg/mL were significantly lower compared to the solvent group (Figure

2a, $P < 0.0001$, $P < 0.0001$, $P < 0.01$, $P < 0.001$, $P < 0.001$ and $P < 0.05$,

respectively). Conversely, the live RLUs of RIF 0.5, RIF 0.125 and INH 0.125

mg/mL groups did not exhibit significant differences from those of the solvent group

(Figure 2a, $P > 0.05$). These findings were also largely supported by both the lung

RLUs and CFUs (Figure 2b and 2c), except for the lung CFUs of the RIF at 0.5

mg/mL group, which were slightly significantly lower than that of the solvent group

(Figure 2c, $P < 0.05$). Collectively, these findings indicate a negative correlation

between live RLUs and administered dosages, suggesting that live RLUs could serve

as an initial criterion for assessing the therapeutic efficacy.

The lung RLUs of the RIF 2 mg/mL group exhibited a significant reduction

compared to those in the solvent group, with a $\log_{10}$RLUs/lung difference exceeding

1.5 and even similar to that of the BT group (Figure 2b, $P < 0.0001$ and $P > 0.05$,

respectively). Conversely, the lung RLUs in the RIF 0.5 and 0.125 mg/mL groups

were similar to that in the solvent group, indicating a lack of evident anti-Mtb

activity at these dosages (Figure 2b, $P > 0.05$). The lung RLUs of the INH 2 and 0.5

mg/mL groups were markedly lower than that of the solvent group (Figure 2b, $P <$

0.0001), as all differences in $\log_{10}$RLUs/lung exceeded 1.5 and comparable to that of

the BT group (Figure 2b, $P > 0.05$). The lung RLUs of INH 0.125 mg/mL group

were similar to that of the solvent group (Figure 2b, $P > 0.05$). In comparison, the

lung RLUs of the EMB 10 and 2.5 mg/mL groups exhibited a pronounced decrease

compared to those in the solvent group, with $\log_{10}$RLUs/lung differences exceeding

1.5 and 1, respectively (Figure 2b, $P < 0.0001$). Meanwhile, the lung RLUs of the

EMB 10 mg/mL group were basically equivalent to that of the BT group (Figure 2b,

$P > 0.05$), whereas the lung RLUs in the EMB 0.625 mg/mL group showed less

significant differences from the solvent group (Figure 2b, $P < 0.05$). In conclusion,

the lung RLUs results are consistent with those obtained from the live RLUs,

providing further support for the utility of detecting live RLUs in determining drug

activity.

The lung CFUs in the RIF 2 mg/mL group exhibited a significant reduction

compared to those in the solvent group, given that the difference of $\log_{10}$CFUs/lung

exceeded 1.5 (Figure 2c, $P < 0.0001$). The lung CFUs in the RIF 0.5 and 0.25

mg/mL groups were approximately comparable to that of the solvent group (Figure

2c, $P <$ and close to 0.05 and $P > 0.05$, respectively). Notably, no colonies were

observed on the INH 2 mg/mL group's plates, indicating potential complete

clearance of AlRv from the mice lungs, given that only a fraction of the lung

suspension from each mouse was plated. The CFUs of the INH 0.5 mg/mL group

were also considerably lower than that of the solvent group, with a difference of

$\log_{10}$CFUs/lung exceeding 1.5 and even lower than BT group (Figure 2c, $P <$

0.0001). Furthermore, all EMB dosages resulted in significantly reduced lung CFUs

compared to the solvent control (Figure 2c, $P < 0.0001$, $P < 0.0001$ and $P < 0.001$,

respectively). While only the $\log_{10}$CFUs/lung difference between the EMB 10

mg/mL and solvent group exceeded 1.5, with the lung CFUs in the EMB 10 group

nearly identical to those in the BT group (Figure 2c, $P > 0.05$). These CFU results

were consistent with the findings derived from live or lung RLUs data. Overall,

based on the comprehensive experimental data presented, we have successfully

established an autoluminescence-based inhalation administration murine model for

evaluating drugs activity against Mtb *in vivo*.

DISCUSSION

Mtb poses a substantial burden of morbidity and mortality, primarily in developing

countries (Chong and Lim, 2009, Grüber, 2020). Despite concerted efforts, the

development of effective anti-Mtb drugs remains confronted with numerous

challenges and uncertainties, including the complexity of the drug activity evaluation

process and the continued emergence of drug-resistant strains, particularly

multidrug-resistant and extensively drug-resistant Mtb (McGrath et al., 2014, Olaru

et al., 2015). Over several decades of continuous research, only bedaquiline,

pretomanid and delamanid have received approved for clinical use (Pontali et al.,

2013). Recently, inhaled administration has emerged as a promising approach for the

treatment of plumonary infections, offering improved cure rates and bacterial

eradication (Riveiro et al., 2023). However, the evaluation of inhalable drugs still

presents challenges such as complex manipulation and time-consuming processes.

Addressing these challenges is crucial to optimizing the efficacy and accessibility of

inhaled drug delivery in clinical settings. Therefore, the development of a rapid,

accurate and efficient activity evaluation model with impressive clinical applicability

performance has the potential to significantly accelerate drug development progress.

In this study, we implemented an autoluminesence-based inhalation administration

murine model to assess the *in vivo* anti-Mtb activity of three first-line TB drugs. The

assessment process can be significantly shortened by detecting either the live RLUs

of mice or the RLUs of lung suspensions, thereby yielding prompt results. Our

findings demonstrate the successful establishment of an autoluminescence-based

murine model for assessing respiratory administration efficacy against Mtb, offering

substantial reductions in time, cost, and labor. The autoluminesence-based inhalation

administration murine model established holds promise for improving the

assessment of anti-Mtb drug efficacy *in vivo*, thus preventing the omission of

potentially effective anti-Mtb drugs and minimizing the risk of serious side effects.

The drug's activity can be determined by comparing the lung suspension RLUs of

different groups within a shorter duration than the current 17-day period (from

infection to get the results), since the lung RLUs become significantly higher than

the background value a few days earlier than the live RLUs do. Additionally, drug

efficacy could potentially be observed within a considerably shorter timeframe in

live mice if higher infection dosages were administered, such as via injection,

although this may result in a higher standard deviation (Liu Yang et al., 2019, Olaru

et al., 2015). Further investigation is warranted to determine whether other anti-

tuberculosis drugs or potent compounds demonstrate efficacy using this model.

Exploring alternative formulations or solvents could also be of interest to enhance

the drug's effectiveness beyond sterilized water. In conclusion, the autoluminesence-

based inhalation administration murine model for evaluating anti-Mtb drug efficacy

in vivo has the potential to expedite the development of innovative anti-Mtb drugs

and treatment regimens.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article, further

inquiries can be directed to the corresponding authors.

ETHICS STATEMENT

The animal care and experimental protocols were approved by the Laboratory

Animal Ethics Committee of the Guangzhou Institutes of Biomedicine and Health,

Chinese Academy of Sciences.

AUTHOR CONTRIBUTIONS

Conceived the project: Tianyu Zhang, Jinxing Hu, Xirong Tian, Yamin Gao;

Designed the research: Tianyu Zhang, Jinxing Hu, Xirong Tian, Yamin Gao;

Performed the studies: Xirong Tian, Yamin Gao, Jingran Zhang, Yanan Ju, Jie Ding,

Sanshan Zeng, Wanli Ma; Interpreted the results: Tianyu Zhang, Xirong Tian,

Yamin Gao; Drafted the manuscript: Tianyu Zhang, Jinxing Hu, Xirong Tian, Yamin

Gao, H.M. Adnan Hameed, Nanshan Zhong, Gregory M. Cook; Final approval of

the manuscript: all authors.

FUNDING

This work was supported by the National Key R&D Program of China

(2021YFA1300900; 2023YFFO713605), the National Natural Science Foundation

of China (81973372, 21920102003, 82061138019, 32300152), Guangdong

Provincial Basic and Applied Basic Research Fund (2022A1515110505), China

Postdoctoral Science Foundation (2022M723164), the Chinese Academy of Sciences

Grants (YJKYYQ20210026), State Key Laboratory of Respiratory Disease

(SKLRD-Z-202301, SKLRD-Z-202414), and Guangzhou Science and Technology

Planning Project (2023A03J0992).

REFERENCES

Ahmad F, Rani A, Alam A, Zarin S, Pandey S, Singh H, et al. (2022). Macrophage: A

Cell With Many Faces and Functions in Tuberculosis. *Front Immunol*, 13, 747799.

Barnes PJ, Burney PGJ, Silverman EK, Celli BR, Vestbo J, Wedzicha JA, et al.

(2015). Chronic obstructive pulmonary disease. *Nat Rev Dis Primers*, 1, 15076.

Chong VH, Lim KS. (2009). Gastrointestinal tuberculosis. *Singapore Med J*, 50(6),

638-645.

Dousa KM, Kurz SG, Bark CM, Bonomo RA, Furin JJ. (2020). Drug-Resistant

Tuberculosis: A Glance at Progress and Global Challenges. *Infect Dis Clin North Am*,

34(4), 863-886.

Fan Z, Zhou B, Liu Y, Sun W, Fang Y, Lu H, et al. (2022). Optimization and

Application of an Efficient and Stable Inhalation Exposure System for Rodents.

AAPS PharmSciTech. 23(1), 50.

Grüber G. (2020). Introduction: Novel insights into TB research and drug discovery.

Prog Biophys Mol Biol. 152, 2-5.

Hameed HMA, Islam MM, Chhotaray C, Wang C, Liu Y, Tan Y, et al. (2018).

Molecular Targets Related Drug Resistance Mechanisms in MDR-, XDR-, and TDR-

Mycobacterium tuberculosis Strains. *Front Cell Infect Microbiol*. 8, 114.

Liu P, Yang Y, Tang Y, Yang T, Sang Z, Liu Z, et al. (2019). Design and synthesis of

novel pyrimidine derivatives as potent antitubercular agents. *Eur J Med Chem*. 163,

169-182.

Liu Y, Gao Y, Liu J, Tan Y, Liu Z, Chhotaray C, et al. (2019). The compound TB47 is

highly bactericidal against *Mycobacterium ulcerans* in a Buruli ulcer mouse model.

Nat Commun. 10(1), 524.

McGrath M, Gey van Pittius NC, van Helden PD, Warren RM, Warner DF. (2014).

Mutation rate and the emergence of drug resistance in *Mycobacterium tuberculosis*. *J*

Antimicrob Chemother. 69(2), 292-302.

McQuaid CF, Vassall A, Cohen T, Fiekert K, White RG. (2021). The impact of

COVID-19 on TB: a review of the data. *Int J Tuberc Lung Dis*. 25(6), 436-446.

Namvarpour M, Tebianian M, Mansouri R, Ebrahimi SM, Kashkooli S. (2019).

Comparison of different immunization routes on the immune responses induced by

Mycobacterium tuberculosis ESAT-6/CFP-10 recombinant protein. *Biologicals*. 59,

6-11.

Olaru ID, von Groote-Bidlingmaier F, Heyckendorf J, Yew WW, Lange C, Chang

KC. (2015). Novel drugs against tuberculosis: a clinician's perspective. *Eur Respir J*.

45(4), 1119-1131.

Pontali E, Matteelli A, Migliori GB. (2013). Drug-resistant tuberculosis. *Curr Opin*

Pulm Med. 19(3), 266-272.

Riveiro V, Casal A, Abelleira R, Valdés L. (2023). Advances in inhaled antibiotics

for management of respiratory tract infections. *Curr Opin Pulm Med*. 29(3), 160-167.

Sanchis J, Corrigan C, Levy ML, Viejo JL. (2013) Inhaler devices - from theory to

practice. *Respir Med*. 107(4), 495-502.

Shah T, Shah Z, Yasmeen N, Baloch Z, Xia X. (2022). Pathogenesis of SARS-CoV-2

and *Mycobacterium tuberculosis* Coinfection. *Front Immunol*. 13, 909011.

Shoen C, DeStefano M, Hafkin B, Cynamon M. (2018). *In Vitro* and *In Vivo*

Activities of Contezolid (MRX-I) against *Mycobacterium tuberculosis*. *Antimicrob*

Agents Chemother. 62(8), e00493-18.

Xu J, Wang B, Fu L, Zhu H, Guo S, Huang H, et al. (2019) *In Vitro* and *In Vivo*

Activities of the Riminophenazine TBI-166 against *Mycobacterium tuberculosis*.

Antimicrob Agents Chemother. 63(5), e02155-18.

Yang X, Liang J, Wang Z, Su Y, Zhan Y, Wu Z, et al. (2021). Sesamolin Protects

Mice From Ovariectomized Bone Loss by Inhibiting Osteoclastogenesis and

RANKL-Mediated NF-κB and MAPK Signaling Pathways. *Front Pharmacol*. 12,

664697.

Yoon HK, Park Y-B, Rhee CK, Lee JH, Oh Y-M. (2017). Summary of the Chronic

Obstructive Pulmonary Disease Clinical Practice Guideline Revised in 2014 by the

Korean Academy of Tuberculosis and Respiratory Disease. *Tuberc Respir Dis*

(Seoul). 80(3), 230-240.

Yu W, Chiwala G, Gao Y, Liu Z, Sapkota S, Lu Z, et al. (2020). TB47 and

clofazimine form a highly synergistic sterilizing block in a second-line regimen for

tuberculosis in mice. *Biomed Pharmacother*. 131, 110782.

Yu W, Yusuf B, Wang S, Tian X, Hameed HMA, Lu Z, et al. (2021). Sterilizing

Effects of Novel Regimens Containing TB47, Clofazimine, and Linezolid in a

Murine Model of Tuberculosis. *Antimicrob Agents Chemother*. 65(10), e0070621.

Zhang T, Li S-Y, Nuermberger EL. (2012). Autoluminescent *Mycobacterium*

tuberculosis for rapid, real-time, non-invasive assessment of drug and vaccine

efficacy. *PLoS One.* 7(1), e29774.

Zhao G, Ho W, Chu J, Xiong X, Hu B, Boakye-Yiadom KO, et al. (2023). Inhalable

siRNA Nanoparticles for Enhanced Tumor-Targeting Treatment of KRAS-Mutant

Non-Small-Cell Lung Cancer. *ACS Appl Mater Interfaces*. 15(26), 31273-31284.

Figure 1. The operating mechanism of the Tow Systems Nose-Only Exposure

Units.

381 **Table 1. Scheme of experiment 1.**

Drug/Dosage[1] (mg/mL)	Number of mice sacrificed at the given time points				Total
	D0[2]	D3[3]	D11-17	D18[4]	
Solvent		6	Detecting chest	6	12
RIF 2		Treatments	RLUs of live mice	6	6
INH 2		initiation	in solvent group	6	6
EMB 10	Infection		daily from day 11	6	6
			until live RLUs		
			exceed 400		
Total	30	6		24	30

[1] RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the

number after the drug represents the dosage.

[2] D0, a total of 30 mice were infected with AlRv by aerosol.

[3] D3, the mice were randomly divided into groups and treated daily from the day 3 to

day 16. The live RLUs, the lung suspension RLUs and CFUs of six mice were

detected before treatment initiated.

[4] D18, the mice were sacrificed and the lungs were removed for detecting RLUs and

CFUs.

All CFUs were read 4 weeks after incubation at 37 □.

Figure 2. Anti-TB drug activities by inhalation administration in Experiment 1.

BT, before treatment; RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the number after the drug represents the

dosage; *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$; ns, $P > 0.05$, no significance. The dotted lines represent the background

values.

Table 2. Scheme of experiment 2.

Drug/Dosage[1] (mg/mL)	Number of mice sacrificed at the given time points				Total
	D0[2]	D1[3]	D13-16	D17[4]	
Solvent		6		6	12
RIF 2				6	6
RIF 0.5			Detecting chest	6	6
RIF 0.125			RLUs of live mice	6	6
INH 2			in solvent group	6	6
INH 0.5	Infection	Treatments initiation	daily from day 13	6	6
INH 0.125			until live RLUs	6	6
EMB 10			exceed 400	6	6
EMB 2.5				6	6
EMB 0.625				6	6
Total	66	6		60	66

[1] RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent, sterilized water; the number after the drug represents the dosage.

[2] D0, a total of 66 mice were infected with AlRv by aerosol.

[3] D1, the mice were randomly divided into groups and treated daily from day1 to day 15. The live RLUs, the lung suspension RLUs and CFUs of six mice were detected before treatment initiated.

[4] D17, the mice were sacrificed and the lungs were removed for detecting RLUs and CFUs.

All CFUs were read 4 weeks after incubation at 37 □.

Figure 3. Anti-TB drug activities by inhalation administration in Experiment 2.

(a) Light intensities of live mice. (b) The suspension light intensities. (c) The lung

Mtb burden.

BT, before treatment; RIF, rifampicin; INH, isoniazid; EMB, ethambutol; Solvent,

sterilized water; the number after the drug represents the dosage; *, $P < 0.05$; **, $P <$

0.01; ***, $P < 0.001$; ****, $P < 0.0001$; ns, $P > 0.05$, no significance. The dotted

lines represent the background values.

drug solutions / sterilized water

atomizer

RATE: 0.2 mL/min
TIME: 20-25 min

controller

exposure cabinet

holder chamber