Hydroxysteroid 17-beta dehydrogenase 13 (*Hsd17b13*) knockdown attenuates liver steatosis in high-fat diet obese mice

Shehroz Mahmood[1,2], Nicola Morrice[1,2,4], Dawn Thompson[1,2], Sara Milanizadeh[1,2], Sophie Wilson[1,2], Philip D. Whitfield[5], George D. Mcilroy[1,3], Justin J. Rochford[1,3], Nimesh Mody[1,2]

1. Aberdeen Cardiovascular & Diabetes Centre,

2. Institute of Medical Sciences,

3. The Rowett Institute,

1,2,3. School of Medicine, Medical Sciences & Nutrition, University of Aberdeen, Aberdeen, AB25 2ZD, Scotland, UK

4. Centre for Genome Enabled Biology & Medicine, University of Aberdeen, Aberdeen, AB24 3UU, Scotland, UK

5. Glasgow Polyomics, University of Glasgow, Garscube Campus, Glasgow, G61 1QH, UK.

Correspondence: Nimesh Mody, n.mody@abdn.ac.uk

RESEARCH ARTICLE

RUNNING HEAD:

Hsd17b13 RNAi knockdown attenuates liver steatosis in mice

ABSTRACT

Hydroxysteroid 17-beta dehydrogenase 13 (*HSD17B13*) loss-of-function gene variants are associated with decreased risk of 'metabolic dysfunction-associated steatotic liver disease' (MASLD). Our RNA-seq analysis of steatotic liver from obese mice -/+ Fenretinide treatment identified major beneficial effects of Fenretinide on hepatic gene expression including *Hsd17b13*. We sought to determine the relationship between Hsd17b13 expression and MASLD and to validate it as a therapeutic target by liver-specific knockdown. *Hsd17b13* expression, which is unique to hepatocytes and associated with the lipid-droplet, was elevated in multiple models of MASLD and normalised with prevention of obesity and steatotic liver. Direct, liver-specific, shRNA-mediated knockdown of *Hsd17b13* (shHsd17b13) in high-fat diet (HFD)-obese mice, markedly improved hepatic steatosis with no effect on body weight, adiposity or glycaemia. shHsd17b13 decreased elevated serum ALT, serum FGF21 levels and markers of liver fibrosis e.g. *Timp2*. shHsd17b13 knockdown in HFD-obese mice and Hsd17b13 overexpression in cells reciprocally regulated expression of lipid metabolism genes e.g. *Cd36*. Global lipidomic analysis of liver tissue revealed a major decrease in diacylglycerols (e.g. DAG 34:3) with shHsd17b13 and an increase in phosphatidylcholines containing polyunsaturated fatty acids (PUFA) e.g. PC 34:3 and PC 42:10. Expression of key genes involved in phospholipid and PUFA metabolism e.g. *Cept1*, were also reciprocally regulated suggesting a potential mechanism of Hsd17b13 biological function and role in MASLD. In conclusion, *Hsd17b13* knockdown in HFD-obese adult mice was able to alleviate MASLD via regulation of fatty acid and phospholipid metabolism, thereby confirming HSD17B13 as a genuine therapeutic target for MASLD and development of liver fibrosis.

NEW & NOTEWORTHY

RNA-seq analysis of steatotic liver identified beneficial effects of Fenretinide on hepatic gene expression including Hsd17b13. Liver-specific shRNA knockdown of Hsd17b13 (shHsd17b13) in obese mice markedly improved hepatic steatosis and markers of liver health e.g. Fgf21. Hsd17b13 influenced expression of lipid/phospholipid metabolism genes e.g. Cd36 and Cept1 and phosphatidylcholines PC 34:3 and PC 42:10. Our study suggests a mechanism of HSD17B13's biological function and that the translational potential of targeting HSD17B13 for MASLD/MASH is strong.

Keywords: Fenretinide, fibrosis, non-alcoholic fatty liver disease (NAFLD), Hydroxysteroid 17-Beta Dehydrogenase 13 (Hsd17b13), phospholipids

INTRODUCTION

Metabolic dysfunction-associated steatotic liver disease (MASLD), previously termed non-alcoholic fatty liver disease (NAFLD) is a common complication associated with obesity and diabetes. MASLD can progress to the more severe metabolic dysfunction-associated steatohepatitis (MASH), previously termed non-alcoholic steatohepatitis (NASH) and therefore requires urgent therapeutic attention (1). MASLD is characterised by an accumulation of triglycerides in the liver and is estimated to affect 25-30% of the global population as well as up to 90% of patients classed as morbidly obese (1). Currently, there are no approved therapeutics available for MASLD other than dietary and lifestyle interventions, however human trials suggest medicines with known benefits against obesity and diabetes may be useful to also treat MASLD (2). In addition, several novel pharmacologic therapies are in various phases of clinical development for the specific treatment of MASLD/MASH (3).

Several studies including human genome-wide association and multi-omics data analysis have identified genes that are key drivers of MASLD (4). For example, 17-beta hydroxysteroid dehydrogenase 13 (HSD17B13) is a novel lipid-droplet protein that has been identified from human steatotic liver biopsies (5). Three independent gene variants in *HSD17B13* which generate loss-of-function proteins are reported to associate with protection from liver injury (4, 6, 7). HSD17B13 gene expression and protein levels are upregulated in patients and mouse models of MASLD and overexpression of HSD17B13 in mice promotes rapid lipid accumulation in the liver of mice (5). However, the traditional gene knockout of *Hsd17b13* in mice had no beneficial effect on the development of liver steatosis with various dietary interventions (8). This surprising result suggests that the normal biological role of HSD17B13 and its metabolite pathway maybe important for normal liver function from birth, but abnormally elevated levels of HSD17B13 in adulthood is a pathological driver of steatotic liver.

In the present study, our analysis of unbiased, whole-genome RNA-sequencing in liver tissue from mice with high-fat diet (HFD) induced obesity and type-2 diabetes and mice treated with the synthetic retinoid Fenretinide to inhibit adiposity, insulin resistance and liver steatosis

(9, 10) revealed genes that are key drivers of MASLD including Hsd17b13. Thus, we

hypothesised that direct inhibition of elevated levels of liver HSD17B13 using RNAi-mediated

knockdown in vivo may reveal a beneficial phenotype and lead to novel insights into the

biological function of this uncharacterised lipid-droplet protein.

MATERIALS AND METHODS
Animal studies

All animal procedures were performed under a project license approved by the U.K. Home

Office under the Animals (Scientific Procedures) Act 1986 (PPL P1ECEB2B6) and the

University of Aberdeen Ethics Review Board. The Fenretinide HFD and *db/db* studies were

previously described (9, 10). For the Hsd17b13 knockdown study, 3-4 week male C57BL/6J mice

were purchased from Charles River. Mice were maintained at 22–24°C on 12-h light/dark cycle

with free access to food/water. At 11 weeks age, 18 mice were placed on to a 45% kcal HFD

(Research diets, D12451) whilst 8 remained on regular Chow. Following 21 weeks on HFD,

mice were randomised into two groups for intraperitoneal (i.p.) injection of AAV8-shscrmbl or

AAV8-shHsd17b13 (Vectorbuilder) at a virus titer of $1x10^{11}$ virus particles. Chow controls

received a saline injection of the same volume administered by i.p. injection. Body weight was

measured weekly and physiological tests performed. Two weeks (14 days) after virus-shRNA

injections, mice were killed (following 5 hours fasting) by CO2-induced anaesthesia and cervical

dislocation. Trunk blood following decapitation was collected and serum stored at -80°C. All

tissues were rapidly dissected, frozen in liquid nitrogen, and stored at -80°C. Tissues for

histology were fixed in formalin for 48h at 4°C, then stored at 4°C in phosphate-buffered saline

(PBS) until further processing.

Physiological tests

Body composition was measured before virus-shRNA injection to confirm HFD-induced obesity

and aid with randomization into treatment/control groups. The measurement was repeated 7 days

after shRNA treatment. Each mouse was analysed using an Echo MRI-3-in-1 scanner (Echo

Medical Systems, Houston, TX, USA). Glucose tolerance test (GTT) was performed at 10 days

post shRNA treatment. Briefly, mice were fasted for 5hrs, baseline glucose levels were sampled

from tail blood using glucose meters (AlphaTRAK, Abbott Laboratories, Abbot Park, IL, USA).

Subsequently mice were injected i.p. with 20% glucose (w/v) and blood glucose measured at 15,
30, 60 and 90 mins post-injection.

Stable expression of HSD17B13 in cells

HEK293 cells and HepG2, were maintained in Dulbecco's Modified Eagle Medium (DMEM)
GlutaMAX with 10% foetal bovine serum (FBS) at 37 ^{o}C and 5% CO_2. Human *HSD17B13*
(NCBI Accession number: NM_178135.5) with HA-tag on the C-terminus was cloned into
pcDNA3.1 (Genscript). HEK293 cells and HepG2 cells were transfected with HSD17B13 or
empty control vector using Lipofectamine 2000, cultured for 2-3 weeks in G418 for neomycin
resistance and positive colonies selected. Stable cell lines were treated with retinoic acid (RA, 1
μm, 2 h) or DMSO (vehicle control). Fatty acid (1mM 2:1 ratio of oleic and palmitic acid, 24 h)
or 1% methanol (vehicle control) was prepared with 10% bovine serum albumin (fatty acid free)
and 1% FBS in DMEM+G418.

Immunoblotting

Frozen liver tissues were homogenised in 400μl of ice-cold Radioimmunoprecipitation assay
buffer (10mM Tris-HCl pH 7.4, 150mM NaCl, 5mM EDTA pH 8.0, 1mM NaF, 0.1% SDS, 1%
Triton X-100, 1% Sodium Deoxycholate with freshly added 1mM $NaVO_4$ and protease
inhibitors) using a PowerGen 125 homogeniser and lysates normalised to 1μg/μl. Proteins were
separated on a 4-12% Bis-Tris gel (Fisher Scientific) by gel electrophoresis and transferred onto
a nitrocellulose membrane. Membranes were probed with the following primary antibodies:
HSD17B13 (1/500, ab122036, Abcam), GFP (1/1000, A11122, Invitrogen), GAPDH (1/1000,
14C10) and Vinculin (1/1000, 13901) both Cell Signaling Technology. Immunoblotting for RBP
(Dako) was from 1μl serum (11).

RNA extraction and RT-qPCR

Frozen tissues were lysed in TRIzol reagent (Sigma) and RNA isolated using phenol/chloroform
extraction according to manufacturer's instructions. RNA was then synthesized into cDNA
Tetrokit (Bioline, UK) and subjected to qPCR analysis using SYBR green and LightCycler 480
(Roche). Gene expression was determined relative to the reference gene *Nono* for mouse liver
tissue and *Hprt* for human cell lines (HEK and HepG2) and analysis was performed using the
Pfaffl method (12). Primer sequences can be found in Supplemental Table 1.

Liver Histology

Liver tissues were embedded in paraffin, sectioned and stained with haematoxylin and eosin (H&E). Images were taken using a light microscope, EVOS XL (Thermofisher Scientfic, UK) at 20x and 40x magnification. Frozen liver tissue was used for fluorescent staining. Briefly, processed slides were washed PBS for 10-12 min. Subsequently, slides were incubated with Blocking (5% goat serum in PBS) and then with HCS Lipidtox^{TM} deep red neutral lipid stain (H34477) (1:200 in PBS) in RT each for 1 h. Furthermore, they were mounted with fluoroshield mounting medium with DAPI (ab104139).

Liver and serum assays

50-100mg of frozen liver tissues were homogenised in 1ml of PBS and frozen in liquid nitrogen to enable further cell lysis. Samples were left to thaw at room temperature and centrifuged for 15s at 7500rpm to pellet debris. The remaining fat cake was resuspended in PBS, the resulting supernatant used to determine total liver triglycerides according to manufacturer's instructions (Sigma, cat: MAK266). Serum was analysed for alanine aminotransferase (ALT) activity (Abcam, ab105134), FGF21 (R&D Systems, #MF2100) and total triglycerides (Sigma, cat: MAK266) following manufacturer's guidelines.

Lipidomics Analysis

Global lipidomics analysis was performed by high resolution liquid chromatography-mass spectrometry (LC-MS) using an Exactive Orbitrap mass spectrometer (Thermo Scientific, Hemel Hempsted, UK) interfaced to a Thermo UltiMate 3000 RSLC system. Samples (10 μl) were injected onto a Thermo Hypersil Gold C18 column (1.9 μm; 2.1 mm x 100 mm) maintained at 50°C. Mobile phase A consisted of water containing 10 mM ammonium formate and 0.1% (v/v) formic acid. Mobile phase B consisted of a 90:10 mixture of isopropanol-acetonitrile containing 10 mM ammonium formate and 0.1% (v/v) formic acid. The initial conditions for analysis were 65%A-35%B, increasing to 65%B over 4 min and then to 100%B over 15 min, held for 2 min prior to re-equilibration to the starting conditions over 6 min. The flow rate was 400 μl/min. Analyses were undertaken in positive and negative ion modes at a resolution of 100,000 over the mass-to-charge ratio (m/z) range of 250 to 2,000. Progenesis QI v3.0 (Nonlinear Dynamics,

171 Newcastle upon Tyne, UK) was used to process the data sets and lipids were identified through
172 interrogation of HMDB (http://www.hmdb.ca/), LIPID MAPS (www.lipidmaps.org/).
173 Multivariate statistical analysis was performed using SIMCA-P v13.0 (Umetrics, Umea,
174 Sweden) and the heatmap was generated with the online available platform MetaboAnalyst 5.0
175 (https://www.metaboanalyst.ca/) (13).

RNA sequencing

As previously described (10). Total RNA was extracted from frozen liver using TriZOL reagent
(Sigma Aldrich, Irvine, UK) to manufacturer's instructions. RNA was purified using RNeasy
miniprep kit (Qiagen, Manchester, UK) and quantified on a Bioanalyzer 2000 (Agilent
Technologies, Edinburgh, UK). Ribosomal RNA was removed from samples using the Ribo-
Zero rRNA removal kit (Illumina, San Diego, CA, USA). Sequencing libraries were prepared
using the TruSeq RNA Library Preparation Kit v2 (Illumina) following the low sample protocol,
using 1 μg RNA. Libraries were sequenced on the NextSeq-500 Desktop sequencer platform
(Illumina) with a 2×75 paired-end read length giving 719 million raw reads (average depth 45
million reads per sample). Reads were filtered using the FASTQ Toolkit v2.0.0 in the Illumina
BaseSpace cloud computing environment to remove adapter sequences and reads with phred
score <20 or length <20 bp. Filtered reads were checked using FastQC before aligning with
STAR 2.0 to the *Mus musculus* UCSC mm10 genome and differential expression measured using
DESeq2 with default parameters. RNA-seq data generated in this study has been deposited in the
NCBI Gene Expression Omnibus (GEO) database accession number GSE220684.

Statistical Analysis.

192 All values are expressed as mean + S.E.M, unless otherwise stated. Statistical analyses were
193 performed by using one- or two-way ANOVA (where stated) followed by multiple-comparison
194 tests to compare the means of three or more groups. In all figures, $*/^{\#}p \leq 0.05$, $**/^{\#\#}p \leq 0.01$,
195 $***/^{\#\#\#}p \leq 0.001$. All analyses were performed using GraphPad Prism (GraphPad Software).

RESULTS
Hepatic RNA-seq reveals signature gene expression alterations in HFD-induced obese steatotic liver and rescue by beneficial effects of Fenretinide

We have previously reported the effects of Fenretinide to inhibit adiposity, insulin resistance and the accumulation of liver triglycerides. To characterise the molecular mechanism of beneficial effects of Fenretinide in the liver, we performed RNA-seq (GEO number GSE220684) on liver tissue from HFD-induced obese C57BL/6 mice +/- Fenretinide for 20 weeks, compared to lean control mice (Figure 1A) (9, 10). HFD-induced obesity resulted in upregulation of 941 liver genes and downregulation of 637 genes (Figure 1B). Many of the genes upregulated in HFD steatotic liver were involved in triglyceride synthesis and fatty acid metabolism and well-established targets of PPAR-alpha signalling and/or GWAS identified genes associated with MASLD (e.g Hsd17b13, Figure 1C) (14). Fenretinide treatment prevented the increase in expression of many of these genes (Figure 1C).

To further characterise the effects on gene expression changes linked to steatotic liver, we compared the effects of Fenretinide on hepatic gene expression in HFD-induced and genetically obese *db/db* mouse model (10). Both HFD and *db/db* obese mice had increased expression of hepatic lipid metabolism genes (e.g. *Mogat* and *Crat*) (Figure 1D and 1E). Fenretinide treatment prevented this increase in HFD mice but had no effect in *db/db* mice (Figure 1D and 1E). Similarly, HFD repressed other genes (e.g. *Avpr1a* and *Hsd17b2*) and Fenretinide treatment prevented this decrease in HFD mice but had no effect in *db/db* mice (Figure 1F and 1G). These findings suggest the beneficial effects of Fenretinide treatment on hepatic gene expression are associated with inhibition of HFD-induced adiposity and associated insulin resistance and MASLD but not with genetically-obese *db/db* where adiposity and MASLD were not inhibited by Fenretinide (9, 10).

Hsd17b13 expression is upregulated in HFD-induced and genetic mouse models of obesity, hyperglycaemia and MASLD.

17β-Hydroxysteroid dehydrogenases (17β-HSDs) are a family of 15 members, which play a key role in sex hormone metabolism by catalyzing steps of steroid biosynthesis (e.g. Hsd17b1 and b2) (15, 16). Some may also play a role in cholesterol and fatty acid metabolism. Of the hydroxysteroid 17-beta dehydrogenase family of genes the most closely related to *Hsd17b13* is *Hsd17b11* although it is not specific to hepatocytes and also expressed in adipocytes (17). HFD-induced obesity also upregulated *Hsd17b11 and Hsd17b7* hepatic gene expression in C57BL/6 mice and Fenretinide treatment decreased their gene expression compared to HFD mice (new Figure 2A RNA-seq mini-table Hsd17b family of genes). HFD also upregulated *Hsd17b4* and *Hsd17b12* hepatic gene expression in C57BL/6 mice, however, Fenretinide treatment had no effect to decrease these genes. *Hsd17b13* has been linked to MASLD and was one of the top 10% most upregulated genes in C57BL/6 HFD mice, increased 1.5-fold (Figure 2A and 2B).

Hsd17b13 hepatic protein expression was also induced by a similar amount in HFD mice compared to lean controls (Figure 2C). Fenretinide treatment was able to prevent this induction, with both gene and protein expression levels close to lean C57BL/6 control mice (Figure 2A, 2B and 2C). Fenretinide treatment also repressed Hsd17b13 expression in parallel with prevention of liver steatosis in LDLR$^{-/-}$ mice fed-HFD plus high-cholesterol diet model of dyslipidemia (Thompson et al., 2023 and Figure 2D) (18). *Hsd17b13* gene expression levels were induced in two more genetical models of metabolic disease and steatotic liver, in leptin-deficient obese *ob/ob* mice (Figure 2E) and in lipodystrophic Seipin/*Bscl2*$^{-/-}$ mice (Figure 2F) (19). FEN treatment could not inhibit the increase in *Hsd17b13* obese *ob/ob* mice but gene therapy to restore WAT in *Bscl2*$^{-/-}$ mice inhibited MASLD and *Hsd17b13* gene expression (Figure 2F) (20). Taken together these data suggest a key role of 17β-Hydroxysteroid dehydrogenases and in particular Hsd17b13 in MASLD.

RNAi-mediated Hsd17b13 knockdown in HFD-induced obesity and MASLD does not affect body weight, adiposity or glucose homeostasis

To evaluate the role of Hsd17b13 in driving steatotic liver, we proceeded to directly knockdown the elevated levels of this protein in adult mice with HFD-induced MASLD thereby

circumventing the lack of a protective effect of Hsd17b13 whole-body, total genetic knockout

from birth in mice. Candidate short hairpin RNA (shRNA) were tested for knockdown efficiency

in HEK293 cells overexpressing mouse *Hsd17b13*, and the most potent (hereafter renamed

shHsd17b13) was selected for treatment *in vivo*.

HFD caused increased weight gain and obesity over time compared to chow-fed

C57BL/6 mice, as expected (Figure 3A). At 21 weeks on HFD, obese mice were randomised for

administration of adeno-associated virus (AAV)-8 constructs encoding either shHsd17b13 or

scrambled control (shScrmbl) with the U6 promoter for high expression of shRNAs. shHsd17b13

treatment had no effect on body weight or adiposity compared to HFD-shscrmbl control in the

following 2 weeks (Figure 3B and C). Hepatic GFP co-expression confirmed successful AAV8

delivery in HFD mice (Figure 3D). shHsd17b13 treatment suppressed the elevated *Hsd17b13*

expression compared to shScrmbl treatment, normalising gene and protein expression to a level

similar to that in chow-fed lean mice (Figure 3E and F). HFD-induced obesity caused glucose

intolerance and shHsd17b13 knockdown had no effect to alter this compared to shScrmbl control

mice (Figure 3G, H and I).

RNAi-mediated Hsd17b13 knockdown prevents HFD-induced MASLD and markers of

fibrotic injury.

HFD-induced obesity and glucose intolerance led to steatotic liver characterised by

increased number and size of lipid droplets as determined by H&E staining of liver tissues and

fluorescent staining of lipid droplets (Figure 4A and B). Strikingly, *Hsd17b13* knockdown

substantially decreased the number and size of hepatic lipid droplets in HFD mice. In addition,

shHsd17b13 decreased liver triglycerides by 45% from the elevated level in HFD-shscrmbl and

back towards a normal level in CHOW mice (Figure 4C *first panel*). HFD-induced liver steatosis

was associated with elevated levels of serum ALT and serum Fgf21 compared to lean mice

(Figure 4C *second and third panels*). shHsd17b13 decreased both serum ALT levels and serum

Fgf21 indicating biomarkers of improved liver health compared to HFD-shscrmbl mice.

Persistent excess lipid accumulation and the progression to MASH, is associated with

activation of hepatic stellate cells by TGF-β which leads to the development of fibrosis by

collagens and matrix remodeling by tissue inhibitors of matrix metalloproteinases (21). HFD

increased the expression of *Col1a1*, *Timp1*, *Timp2*, *Tgf-β* (Figure 4D). Hsd17b13 knockdown

significantly decreased expression of *Timp2* and trended to decrease expression of *Col1a1*,

Col4a1 and *Timp1, Tgf-β* in HFD mice. Similar changes in the pro-inflammatory macrophage

marker *Cd68* were determined. Thus, *Hsd17b13* knockdown and attenuation of excess hepatic

triglycerides storage, appeared to inhibit the development of fibrosis and MASH, via suppressing

gene expression alterations associated with hepatic stellate cell activation and a pro-

inflammatory environment.

Increasing or decreasing Hsd17b13 levels does not affect retinoid transcriptional regulator

genes or serum RBP4.

Hsd17b13 has been reported to have *in vitro* dehydrogenase activity towards retinol and thus

modulate retinoic acid levels in cells transfected with *Hsd17b13* (6). Thus, we determined

whether targets of retinoic acid signalling were altered with Hsd17b13 knockdown. HFD

increased hepatic *Lrat* expression which is associated with altered retinyl ester dynamics in

stellate cells and MASH (Figure 5A) (22, 23). Hsd17b13 knockdown trended to decrease

expression of *Lrat.* shHsd17b13 had no effect to alter the expression of nuclear receptors *Rara,*

Rxra, Rxrb or *Rarb,* a classic retinoic acid target gene (Figure 5A). HFD increased serum RBP4

levels in shScrmbl control mice compared to chow control and shHsd17b13 treatment had no

effect (Figure 5B). Our results suggest that Hsd17b13 does not modulate retinoic acid levels or

retinol homeostasis *in vivo*. To extend these studies further, we examined the effect of Hsd17b13

over-expression and retinoic acid in cells. Although retinoic acid could increase the expression of

Cyp26a1 and *Rarb*, two classic retinoic acid inducible genes, Hsd17b13 over-expression had no

effect alone or in combination with retinoic acid to alter their gene expression (Figure 5C).

Hsd17b13 over-expression did increase *Raldh1* expression by >2-fold (p<0.001 by t-test) as did

retinoic acid treatment however there was no additive or synergistic interaction (by two-way

ANOVA). Our results suggest that Hsd17b13 does not play a significant role in modulating

retinoic acid levels or retinol homeostasis.

Changing Hsd17b13 levels alters hepatic lipid and phospholipid metabolism.

Dis-regulation of a network of hepatic genes involved in fatty acid metabolism in

response to over-nutrition contribute to excess hepatic lipid accumulation and thus MASLD. We

hypothesised that *Hsd17b13* knockdown mediated decrease in steatotic liver may be associated

with a rescue of this impairment. HFD increased hepatic expression of PPARα target genes

Cd36, Vldlr, Crat and *Mogat1* involved in fatty acid transport and metabolism*,* and shHsd17b13

treatment inhibited this induction (Figure 6A). To determine if these gene expression changes

were due to acute changes in HSD17B13 protein or linked secondarily to decreased hepatic lipid

accumulation next, we examined the effect of increased Hsd17b13 expression in cells. Increased

Hsd17b13 expression increased triglyceride storage in cells (Figure 6B) and increased *Cd36*

expression without additional fatty acids (Figure 6C). Fatty acid treatment further increased

triglyceride storage in cells with Hsd17b13 but did not change *Cd36* expression.

Next, we performed lipidomic analysis of liver tissues to understand the mechanism of

HSD17B13 biological function and role in MASFD. shHsd17b13 treatment decreased

diacylglycerols and increased phospholipids, in particular phosphatidylcholines containing

polyunsaturated fatty acids (PUFA) e.g. PC 34:3 and PC 42:10 (Figure 6D). We studied the

pathway which is responsible for the production of phospholipids and PUFAs. HFD increased

the expression of hepatic *Fads1* and *Lpcat3/Mboat5* (p=0.04 by t-test) and shHsd17b13

treatment did not significantly alter these (Figure 6E). HFD markedly increased hepatic *Cept1*

expression and shHsd17b13 treatment normalised *Cept1* expression in mice (Figure 6F). CEPT1

or Choline-enthanolamine phosphotransferase 1 catalyzes the terminal step of the Kennedy

pathway and is responsible for phosphatidylcholine (PC) and phosphatidylethanolamine (PE)

production (24). It is also reported to produce the endogenous ligand for PPAR-alpha, a

transcription factor that is a key regulator of hepatic lipid metabolism genes (25). Increased

Hsd17b13 expression in cells and increased *Cept1* expression -/+ additional fatty acids (Figure

6G). Thus, Hsd17b13 is a key driver of increased hepatic triglyceride storage and phospholipid

remodelling, likely via increased *Cept1* and *Cd36* expression, and may explain at least part of the

mechanism of it biological function and role in MASFD.

DISCUSSION

The multiple overlapping secondary pathologies in response to diet-induced obesity such as the development of type 2 diabetes, MASLD, atherosclerosis and CVD are a global medical burden. Currently there is no treatment for MASLD or MASH and thus therapies are urgently needed. We have previously reported the beneficial effects of Fenretinide to prevent the accumulation of hepatic triglycerides via decreased whole-body adiposity and associated insulin resistance. Here our RNA-seq study of liver tissue from HFD obese mice -/+ Fenretinide identified major beneficial effects of Fenretinide on hepatic gene expression including key drivers of MASLD e.g. HSD17B13. Thus, we sought to directly knockdown the elevated levels of Hsd17b13 to evaluate its role in driving MASLD and regulation of lipid/phospholipid metabolism.

HSD17B13 gene/protein expression are upregulated in MASLD and transgenic overexpression of HSD17B13 promotes rapid lipid accumulation in the liver of mice (5). However, the traditional gene knockout of HSD17B13 challenged with several diets to induce steatotic liver and/or fibrosis did not lead to the hypothesised protective phenotype (8). Our RNAi therapeutic approach, to suppress Hsd17b13 levels in adult mice with existing HFD-induced liver steatosis to decrease hepatic triglyceride storage is a translational approach to investigate the role of Hsd17b13 in MASLD. This RNAi knockdown strategy has been taken by pharma/biotech in on-going clinical trials (26). Possible explanations for the discrepancy between the phenotype of mouse genetic knockout and the RNAi knockdown approach, are if HSD17B13 affects other proteins in a mechanism independent of its enzymatic activity or total loss of Hsd17b13 activity is compensated for by another protein/pathway.

Su, Guan and co-workers (27) reported that Hsd17b13 interacts with ATGL on lipid droplets to regulate lipolysis in response to cAMP-PKA signalling, a mechanism that resembles a role played by Plin5 (28). Thereby, both Hsd17b13 and Plin5 have been shown to be phosphorylated leading to an increase in ATGL-mediated lipolysis. While this may be beneficial for MASLD, increased hepatic cAMP-PKA signalling is also associated with excess endogenous glucose production via gluconeogenesis in obesity and type 2 diabetes (29). Thus, it is not clear if this signalling network can be delineated sufficiently to facilitate targeting for the treatment of metabolic diseases. Recently, the X-ray crystal structure of Hsd17b13 has provided insights into a mechanism for association with the lipid droplet and interaction with ATGL or other potential

binding partners (30). The findings suggest that HSD17B13 may have an important scaffold

function at the lipid droplet.

Distinct changes occur in the liver lipidome with MASLD (31-33). Moreover, many genes

involved in hepatic PUFA and phospholipid metabolism, are important in the formation of

biologically important lipids and are linked to metabolic disease by human GWAS and mouse

knockout studies e.g. FADS1, ELOVL5, LPCAT3, MBOAT7, CEPT1, *PLA2G4A*/cPLA2α,

PLA2G6/iPLA2β, PTGDS and PTGES (34, 35) .The expression of these genes is also altered by

dysregulation of a network of transcription factors such as PPARa, LXR, SREBP. Our

lipidomics data suggests that HSD17B13 plays a key role in hepatic triglyceride and

phospholipid remodelling, whereby knockdown promoted an increase in phosphatidyl-cholines

(PC) containing PUFAs eg. PC 34:3 and PC 42:10. Studies of the human genetic variants of

Hsd17b13 (protective against severe liver disease) have reported an increase in phospholipids

including PC 34:3 (36). shHsd17b13 knockdown normalised liver *Cept1* expression in mice and

increased Hsd17b13 expression in cells led to increased *Cept1* expression. These results suggest

that Hsd17b13 may regulate phospholipid metabolism and may explain at least part of the

mechanism of it biological function and role in MASLD.

Several human genome-wide association studies have identified gene variants in

HSD17B13 which generate loss-of-function proteins that associate primarily with protection

from severity of MASLD and fibrosis i.e. progression to MASH (6, 7, 36-38). In agreement with

our *in vivo* data, mouse models have shown that inhibition/knockdown of Hsd17b13 protects

against both hepatic triglyceride accumulation and fibrosis (27, 39, 40) . We determined that

shHsd17b13 knockdown repressed hepatic fatty acid transporter Cd36 expression in mice and

reciprocally, increased Hsd17b13 expression in cells led to increased Cd36 expression.

Hepatocyte CD36 is a key factor in driving *de novo* lipogenesis and in the development of

MASLD (41-43). Thus, Hsd17b13 may play a role in the regulation of key transcription factors

(such as PPARa, LXR and SREBP) which coordinate the expression of hepatic lipid homeostasis

genes in response to physiological signals.

These data provide strong evidence for an important role of Hsd17b13 in driving

MASLD via regulation of hepatic triglyceride storage and phospholipid metabolism.

Importantly, this beneficial effect is present without an effect on body weight, adiposity or

glucose homeostasis suggesting that directly targeting Hsd17b13 is superior to the use of FEN,

which has diverse mechanisms of action. In addition, treating MASLD may prevent the

development of fibrosis/MASH and create an opportunity to clinically treat these without the

need to treat co-morbidities of obesity, type-2 diabetes and cardiovascular diseases. Thus, the

translation potential of targeting Hsd17b13 for MASLD/MASH is strong (26, 38).

DATA AVAILABILITY

Data will be made available upon reasonable request. Hepatic RNA-seq data generated in this study (HFD+/-Fenretinide, 20 weeks in C57BL/6J mice) has been deposited in the NCBI Gene Expression Omnibus (GEO) database accession number GSE220684.

SUPPLEMENTAL MATERIAL

Supplementary Table 1: qPCR Primers

ACKNOWLEDGMENTS

The authors thank the University of Aberdeen (UoA) animal research facility, qPCR core facility (Institute of Medical Sciences, UoA) and Linda Davidson (NHS Grampian) for their technical contributions regarding animal studies, qPCR and histology respectively. RNA-sequencing was carried out at the UoA Centre for Genome Enabled Biology and Medicine (CGEBM); thanks to Diane Stewart, Sophie Shaw and Elaina Collie-Duguid for support with sample prep, bioinformatics reporting and project supervision. Thanks to Lars Grøntved (University of Southern Denmark) for invaluable discussions about the RNA-seq bioinformatic analysis. Lipidomics was performed at Glasgow Polyomics (University of Glasgow). The authors also wish to thank Mirela Delibegovic (UoA) for invaluable discussions throughout the study and review of the manuscript and members of the Delibegovic-Mody joint lab for their technical contributions.

GRANTS

This study was supported by a James Mearns Trust PhD studentship (to S. Mahmood); British Heart Foundation (PG16/90/32518) project grant and Wellcome Trust ISSF small project grant (to N. Mody) and Wellcome Trust ISSF international partnership award (to N. Mody and Lars Grøntved, University of Southern Denmark); UoA CGEBM PhD studentship (to N. Morrice).

DISCLOSURES

None

DISCLAIMERS

None

AUTHOR CONTRIBUTIONS

GM and N.Mody conceived and designed research; S.Mahmood, N. Morrice, DT, S. Milanizadeh, SW and PW performed experiments and analyzed data; JR and N.Mody interpreted results of experiments; S.Mahmood, S. Milanizadeh, SW, DT and PW prepared figures; S.Mahmood drafted the manuscript and N.Mody edited and revised the manuscript, S.Mahmood, N. Morrice, GM and N.Mody approved final version of manuscript.

REFERENCES

1. Pafili K, Roden M: Nonalcoholic fatty liver disease (NAFLD) from pathogenesis to treatment concepts in humans. *Mol Metab.* 50:101122, 2021

2. Luo Q, Wei R, Cai Y, Zhao Q, Liu Y, Liu WJ: Efficacy of off-label therapy for non-alcoholic fatty liver disease in improving non-invasive and invasive biomarkers: A systematic review and network meta-analysis of randomized controlled trials. *Front Med (Lausanne).* 9:793203, 2022

3. Esler WP, Bence KK: Metabolic targets in nonalcoholic fatty liver disease. *Cell Mol Gastroenterol Hepatol.* 8:247-267, 2019

4. Chella Krishnan K, Kurt Z, Barrere-Cain R, Sabir S, Das A, Floyd R, Vergnes L, Zhao Y, Che N, Charugundla S, Qi H, Zhou Z, Meng Y, Pan C, Seldin MM, Norheim F, Hui S, Reue K, Lusis AJ, Yang X: Integration of multi-omics data from mouse diversity panel highlights mitochondrial dysfunction in non-alcoholic fatty liver disease. *Cell Syst.* 6:103-115.e7, 2018

5. Su W, Wang Y, Jia X, Wu W, Li L, Tian X, Li S, Wang C, Xu H, Cao J, Han Q, Xu S, Chen Y, Zhong Y, Zhang X, Liu P, Gustafsson JA, Guan Y: Comparative proteomic study reveals 17beta-HSD13 as a pathogenic protein in nonalcoholic fatty liver disease. *Proc Natl Acad Sci U S A.* 111:11437-11442, 2014

6. Ma Y, Belyaeva OV, Brown PM, Fujita K, Valles K, Karki S, de Boer YS, Koh C, Chen Y, Du X, Handelman SK, Chen V, Speliotes EK, Nestlerode C, Thomas E, Kleiner DE, Zmuda JM,

Sanyal AJ, (for the Nonalcoholic Steatohepatitis Clinical Research Network), Kedishvili NY, Liang TJ, Rotman Y: 17-beta hydroxysteroid dehydrogenase 13 is a hepatic retinol dehydrogenase associated with histological features of nonalcoholic fatty liver disease. *Hepatology.* 69:1504-1519, 2019

7. Abul-Husn NS, Cheng X, Li AH, Xin Y, Schurmann C, Stevis P, Liu Y, Kozlitina J, Stender S, Wood GC, Stepanchick AN, Still MD, McCarthy S, O'Dushlaine C, Packer JS, Balasubramanian S, Gosalia N, Esopi D, Kim SY, Mukherjee S, Lopez AE, Fuller ED, Penn J, Chu X, Luo JZ, Mirshahi UL, Carey DJ, Still CD, Feldman MD, Small A, Damrauer SM, Rader DJ, Zambrowicz B, Olson W, Murphy AJ, Borecki IB, Shuldiner AR, Reid JG, Overton JD, Yancopoulos GD, Hobbs HH, Cohen JC, Gottesman O, Teslovich TM, Baras A, Mirshahi T, Gromada J, Dewey FE: A protein-truncating HSD17B13 variant and protection from chronic liver disease. *N Engl J Med.* 378:1096-1106, 2018

8. Ma Y, Brown PM, Lin DD, Ma J, Feng D, Belyaeva OV, Podszun MC, Roszik J, Allen JN, Umarova R, Kleiner DE, Kedishvili NY, Gavrilova O, Gao B, Rotman Y: 17-beta hydroxysteroid dehydrogenase 13 deficiency does not protect mice from obesogenic diet injury. *Hepatology.* 73:1701-1716, 2021

9. Mcilroy GD, Delibegovic M, Owen C, Stoney PN, Shearer KD, McCaffery PJ, Mody N: Fenretinide treatment prevents diet-induced obesity in association with major alterations in retinoid homeostatic gene expression in adipose, liver, and hypothalamus. *Diabetes.* 62:825-836, 2013

10. Morrice N, Mcilroy GD, Tammireddy SR, Reekie J, Shearer KD, Doherty MK, Delibegovic M, Whitfield PD, Mody N: Elevated fibroblast growth factor 21 (FGF21) in obese, insulin resistant states is normalised by the synthetic retinoid fenretinide in mice. *Sci Rep.* 7:43782, 2017

11. Yang Q, Graham TE, Mody N, Preitner F, Peroni OD, Zabolotny JM, Kotani K, Quadro L, Kahn BB: Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes. *Nature.* 436:356-62, 2005

12. Pfaffl MW: A new mathematical model for relative quantification in real-time RT-PCR.

Nucleic Acids Res. 29:e45, 2001

13. Pang Z, Chong J, Zhou G, de Lima Morais DA, Chang L, Barrette M, Gauthier C, Jacques P,

Li S, Xia J: MetaboAnalyst 5.0: Narrowing the gap between raw spectra and functional insights.

Nucleic Acids Res. 49:W388-W396, 2021

14. Kersten S: Integrated physiology and systems biology of PPARalpha. *Mol Metab.* 3:354-371,

2014

15. Poutanen M, Penning TM: Biology and clinical relevance of hydroxysteroid (17beta)

dehydrogenase enzymes. *Mol Cell Endocrinol.* 489:1-2, 2019

16. Marchais-Oberwinkler S, Henn C, Moller G, Klein T, Negri M, Oster A, Spadaro A, Werth

R, Wetzel M, Xu K, Frotscher M, Hartmann RW, Adamski J: 17beta-hydroxysteroid

dehydrogenases (17beta-HSDs) as therapeutic targets: Protein structures, functions, and recent

progress in inhibitor development. *J Steroid Biochem Mol Biol.* 125:66-82, 2011

17. Horiguchi Y, Araki M, Motojima K: 17beta-hydroxysteroid dehydrogenase type 13 is a liver-

specific lipid droplet-associated protein. *Biochem Biophys Res Commun.* 370:235-238, 2008

18. Thompson D, Mahmood S, Morrice N, Kamli-Salino S, Dekeryte R, Hoffmann PA, Doherty

MK, Whitfield PD, Delibegovic M, Mody N: Fenretinide inhibits obesity and fatty liver disease

but induces Smpd3 to increase serum ceramides and worsen atherosclerosis in LDLR(-/-) mice.

Sci Rep. 13:3937-w, 2023

19. Mcilroy GD, Suchacki K, Roelofs AJ, Yang W, Fu Y, Bai B, Wallace RJ, De Bari C,

Cawthorn WP, Han W, Delibegovic M, Rochford JJ: Adipose specific disruption of seipin causes

early-onset generalised lipodystrophy and altered fuel utilisation without severe metabolic

disease. *Mol Metab.* 10:55-65, 2018

20. Sommer N, Roumane A, Han W, Delibegovic M, Rochford JJ, Mcilroy GD: Gene therapy

restores adipose tissue and metabolic health in a pre-clinical mouse model of lipodystrophy. *Mol*

Ther Methods Clin Dev. 27:206-216, 2022

21. Friedman SL, Neuschwander-Tetri BA, Rinella M, Sanyal AJ: Mechanisms of NAFLD
development and therapeutic strategies. *Nat Med.* 24:908-922, 2018

22. Molenaar MR, Penning LC, Helms JB: Playing jekyll and hyde-the dual role of lipids in fatty
liver disease. *Cells.* 9:2244. doi: 10.3390/cells9102244, 2020

23. Saeed A, Dullaart RPF, Schreuder TCMA, Blokzijl H, Faber KN: Disturbed vitamin A
metabolism in non-alcoholic fatty liver disease (NAFLD). *Nutrients.* 10:29. doi:
10.3390/nu10010029, 2017

24. Gibellini F, Smith TK: The kennedy pathway--de novo synthesis of
phosphatidylethanolamine and phosphatidylcholine. *IUBMB Life.* 62:414-428, 2010

25. Chakravarthy MV, Lodhi IJ, Yin L, Malapaka RRV, Xu HE, Turk J, Semenkovich CF:
Identification of a physiologically relevant endogenous ligand for PPARalpha in liver. *Cell.*
138:476-488, 2009

26. Amangurbanova M, Huang DQ, Loomba R: Review article: The role of HSD17B13 on
global epidemiology, natural history, pathogenesis and treatment of NAFLD. *Aliment Pharmacol*
Ther. 57:37-51, 2023

27. Su W, Wu S, Yang Y, Guo Y, Zhang H, Su J, Chen L, Mao Z, Lan R, Cao R, Wang C, Xu H,
Zhang C, Li S, Gao M, Chen X, Zheng Z, Wang B, Liu Y, Liu Z, Wang Z, Liu B, Fan X, Zhang
X, Guan Y: Phosphorylation of 17beta-hydroxysteroid dehydrogenase 13 at serine 33 attenuates
nonalcoholic fatty liver disease in mice. *Nat Commun.* 13:6577-1, 2022

28. Keenan SN, De Nardo W, Lou J, Schittenhelm RB, Montgomery MK, Granneman JG, Hinde
E, Watt MJ: Perilipin 5 S155 phosphorylation by PKA is required for the control of hepatic lipid
metabolism and glycemic control. *J Lipid Res.* 62:100016, 2021

29. Yang H, Yang L: Targeting cAMP/PKA pathway for glycemic control and type 2 diabetes
therapy. *J Mol Endocrinol.* 57:R93-R108, 2016

30. Liu S, Sommese RF, Nedoma NL, Stevens LM, Dutra JK, Zhang L, Edmonds DJ, Wang Y,

Garnsey M, Clasquin MF: Structural basis of lipid-droplet localization of 17-beta-hydroxysteroid

dehydrogenase 13. *Nat Commun.* 14:5158-0, 2023

31. McGlinchey AJ, Govaere O, Geng D, Ratziu V, Allison M, Bousier J, Petta S, de Oliviera C,

Bugianesi E, Schattenberg JM, Daly AK, Hyotylainen T, Anstee QM, Oresic M: Metabolic

signatures across the full spectrum of non-alcoholic fatty liver disease. *JHEP Rep.* 4:100477,

2022

32. Ooi GJ, Meikle PJ, Huynh K, Earnest A, Roberts SK, Kemp W, Parker BL, Brown W,

Burton P, Watt MJ: Hepatic lipidomic remodeling in severe obesity manifests with steatosis and

does not evolve with non-alcoholic steatohepatitis. *J Hepatol.* 75:524-535, 2021

33. Velenosi TJ, Ben-Yakov G, Podszun MC, Hercun J, Etzion O, Yang S, Nadal C, Haynes-

Williams V, Huang WA, Gonzalez-Hodar L, Brychta RJ, Takahashi S, Akkaraju V, Krausz KW,

Walter M, Cai H, Walter PJ, Muniyappa R, Chen KY, Gonzalez FJ, Rotman Y: Postprandial

plasma lipidomics reveal specific alteration of hepatic-derived diacylglycerols in nonalcoholic

fatty liver disease. *Gastroenterology.* 162:1990-2003, 2022

34. Zhang JY, Kothapalli KSD, Brenna JT: Desaturase and elongase-limiting endogenous long-

chain polyunsaturated fatty acid biosynthesis. *Curr Opin Clin Nutr Metab Care.* 19:103-110,

2016

35. Jalil A, Bourgeois T, Menegaut L, Lagrost L, Thomas C, Masson D: Revisiting the role of

LXRs in PUFA metabolism and phospholipid homeostasis. *Int J Mol Sci.* 20:3787. doi:

10.3390/ijms20153787, 2019

36. Luukkonen PK, Tukiainen T, Juuti A, Sammalkorpi H, Haridas PAN, Niemela O, Arola J,

Orho-Melander M, Hakkarainen A, Kovanen PT, Dwivedi O, Groop L, Hodson L, Gastaldelli A,

Hyotylainen T, Oresic M, Yki-Jarvinen H: Hydroxysteroid 17-beta dehydrogenase 13 variant

increases phospholipids and protects against fibrosis in nonalcoholic fatty liver disease. *JCI*

Insight. 5:e132158. doi: 10.1172/jci.insight.132158, 2020

37. Anstee QM, Darlay R, Cockell S, Meroni M, Govaere O, Tiniakos D, Burt AD, Bedossa P,

Palmer J, Liu Y, Aithal GP, Allison M, Yki-Jarvinen H, Vacca M, Dufour J, Invernizzi P, Prati

D, Ekstedt M, Kechagias S, Francque S, Petta S, Bugianesi E, Clement K, Ratziu V,

Schattenberg JM, Valenti L, Day CP, Cordell HJ, Daly AK, EPoS Consortium Investigators:

Genome-wide association study of non-alcoholic fatty liver and steatohepatitis in a histologically

characterised cohort(☆). *J Hepatol.* 73:505-515, 2020

38. Zhang H, Su W, Xu H, Zhang X, Guan Y: HSD17B13: A potential therapeutic target for

NAFLD. *Front Mol Biosci.* 8:824776, 2022

39. Wang M, Li J, Li H, Dong B, Jiang J, Liu N, Tan J, Wang X, Lei L, Li H, Sun H, Tang M,

Wang H, Yan H, Li Y, Jiang J, Peng Z: Down-regulating the high level of 17-beta-

hydroxysteroid dehydrogenase 13 plays a therapeutic role for non-alcoholic fatty liver disease.

Int J Mol Sci. 23:5544. doi: 10.3390/ijms23105544, 2022

40. Luukkonen PK, Sakuma I, Gaspar RC, Mooring M, Nasiri A, Kahn M, Zhang X, Zhang D,

Sammalkorpi H, Penttila AK, Orho-Melander M, Arola J, Juuti A, Zhang X, Yimlamai D, Yki-

Jarvinen H, Petersen KF, Shulman GI: Inhibition of HSD17B13 protects against liver fibrosis by

inhibition of pyrimidine catabolism in nonalcoholic steatohepatitis. *Proc Natl Acad Sci U S A.*

120:e2217543120, 2023

41. Koonen DPY, Jacobs RL, Febbraio M, Young ME, Soltys CM, Ong H, Vance DE, Dyck

JRB: Increased hepatic CD36 expression contributes to dyslipidemia associated with diet-

induced obesity. *Diabetes.* 56:2863-2871, 2007

42. Wilson CG, Tran JL, Erion DM, Vera NB, Febbraio M, Weiss EJ: Hepatocyte-specific

disruption of CD36 attenuates fatty liver and improves insulin sensitivity in HFD-fed mice.

Endocrinology. 157:570-585, 2016

43. Rada P, Gonzalez-Rodriguez A, Garcia-Monzon C, Valverde AM: Understanding

lipotoxicity in NAFLD pathogenesis: Is CD36 a key driver? *Cell Death Dis.* 11:802-w, 2020

 FIGURE LEGENDS

Figure 1: <u>**Hepatic gene expression in high-fat diet-induced obese and genetically-obese mice**</u>
<u>**with liver steatosis, and effects of Fenretinide treatment.**</u>

A. Graphical illustration of mouse liver study described.

B. Volcano plot of hepatic RNA-seq data of HFD (obese-steatotic liver) most highly induced
genes by fold change and significance, compared to lean control mice (10).

C. Shortlist of differential gene expression from RNA-seq data, reciprocally regulated genes in
HFD (obese-steatotic liver) +/-FEN, involved in triglyceride synthesis and fatty acid metabolism
and genes associated with MASLD. HFD induced genes vs lean control mice and FEN-HFD
repressed genes vs HFD, significance, adjusted p-value *** $<10^{-15}$, ** $<10^{-8}$, * $<10^{-3}$, ns not
significant.

D. and **F**. Hepatic gene expression in HFD (obese-steatotic liver) mice +/-FEN.

E. and **G**. Hepatic gene expression in *db/db* (genetically-obese steatotic liver) mice +/- FEN.

Data (n=7-8 per group) are presented as mean + SEM and analysed by one-way ANOVA
followed by Bonferroni multiple comparison tests where *p≤0.05, **p≤0.01 and ***p≤0.001
(HFD vs CHOW and *db/db* vs lean) or # p≤0.05 and ## p≤0.01 and ### p≤0.001 (FEN-HFD vs
HFD and FEN-*db/db* vs *db/db*).

Figure 2: <u>**Increased Hsd17b13 expression in a HFD-induced and genetic mouse models of**</u>

<u>**obesity, hyperglycaemia and MASLD.**</u>

A. Differential gene expression from RNA-seq data. Hydroxysteroid 17-beta dehydrogenase family of genes reciprocally regulated in HFD (obese-steatotic liver) +/-FEN. HFD induced genes vs lean control mice and FEN-HFD repressed genes vs HFD, significance, adjusted p-value *** $<10^{-15}$, ** $<10^{-8}$, * $<10^{-3}$, ns not significant.

B. *Hsd17b13* gene expression in HFD (obese-steatotic liver) mice +/-FEN and chow fed lean controls, normalised to *Nono*, (n=7-8 per group).

C. Western blot of Hsd17b13 protein expression in HFD (obese-steatotic liver) mice +/-FEN and chow fed lean controls, and quantification normalised to Gapdh, (n=4 per group).

D. Western blot and quantification of HSD17B13 protein expression in *Ldlr*[-/-] HFD (obese-steatotic liver) mice +/-FEN and control (10% kcal) fat diet lean controls, and quantification normalised to vinculin, (n=5-6 per group).

E. *Hsd17b13* gene expression in *ob/ob* (genetically-obese steatotic liver) mice +/-FEN and control (10% kcal) fat diet lean controls, normalised to *Nono*, (n=6-7 per group).

F. *Hsd17b13* gene expression in *Bscl2*[-/-] (lipodystrophic steatotic liver) mice +/- human *Bscl2*[-/-] rescue and wildtype control, normalised to *Nono*, (n=8-9 per *Bscl2*[-/-] groups, n=11 wildtype control).

Data are presented as mean + SEM and analysed by one-way ANOVA followed by Bonferroni multiple comparison tests where *p≤0.05, **p≤0.01 and ***p≤0.001 (respective MASLD model vs control) or ## p≤0.01 (FEN-HFD vs HFD).

Figure 3: <u>Effect of RNAi-mediated Hsd17b13 knockdown on HFD-induced obesity and glucose homeostasis</u>

A. Body weights of C57BL/6 mice on high-fat diet (HFD, 45% kCal fat) or control CHOW diet for 23 weeks. At 21 weeks on HFD, obese mice were randomised for administration of AAV8-GFP-shHsd17b13 or scrambled control shRNA (shScrmbl), n=8-9 per group.

B. Change in body weight from 21 to 23 weeks diet and

C. Adiposity (% body fat) at week 22 HFD in response to administration of AAV8-GFP-shRNA.

At 23 weeks, following administration of AAV8-GFP-shHsd17b13 or shScrmbl,

D. Western blot of GFP in representative mouse liver tissues

E. *Hsd17b13* gene expression in mouse livers, n=8 per group

F. Western blot of Hsd17b13 protein expression and vinculin (loading control) in mouse livers and quantification, n=4-5 per group.

G. Fasting (5 hours) glycaemia (basal glycemia from GTT). $*p \leq 0.05$ (shHsd17b13 HFD vs Chow)

H. Glucose tolerance test (GTT) performed in week 2 post-treatment with shHsd17b13 or shScrmbl, n=8-9 per group,

I. Area under curve (AUC) from GTT for each individual mouse.

All data presented as mean ± SEM and analysed by one-way ANOVA followed by Bonferroni multiple comparison tests where

$*p \leq 0.05$, $**p \leq 0.01$ and $***p \leq 0.001$ (HFD groups vs Chow)

$\#p \leq 0.05$ and $\#\#\# \, p \leq 0.001$ (HFD- shHsd17b13 vs HFD-shscrmbl)

In addition, in **F**. Western blot quantification, One-way ANOVA $\#p \leq 0.05$ (HFD- shHsd17b13 vs HFD-shscrmbl) but required t-test for $*p \leq 0.05$, (HFD-shscrmbl vs Chow).

in **H**. GTT, t-tests were performed at individual time-points and where $\#\#\# \, p \leq 0.001$ (HFD-shHsd17b13 vs Chow).

Figure 4: <u>**Effect of RNAi-mediated Hsd17b13 knockdown on HFD-induced MASLD**</u>.

A. H&E staining of representative liver tissues at 20x and 40x magnification and

B. fluorescent staining of lipid droplets

in C57BL/6 control CHOW diet mice or HFD obese mice treated with shHsd17b13 or scrambled control (shScrmbl).

Scale bars are 200 μm (20x) and 100 μm (40x) in H&E images and 20 μm in fluorescent images respectively.

C. Liver triglyceride levels, serum ALT activity and serum FGF21 levels.

D. Gene expression of fibrosis markers in liver tissue, Collagen type I alpha 1 (*Col1a1*), Collagen type IV alpha 1 (*Col4a1*), Tissue inhibitor of metalloproteinases (*Timp*)-*1* and *Timp2*, Transforming growth factor beta (*Tgf-β*)-1 and Cluster of Differentiation (*Cd*)-*68*.

Data in **C**) and **D**) are presented as mean ± SEM (n=8 per group) and analysed by one-way ANOVA followed by Bonferroni multiple comparison tests where * $p \leq 0.05$, ** $p \leq 0.01$ and *** $p \leq 0.001$ (comparing HFD groups to control CHOW diet) or # $p \leq 0.05$ and ## $p \leq 0.01$ (comparing HFD-shHsd17b13 to HFD-shscrmbl).

Figure 5: <u>**Effect of altered Hsd17b13 levels on retinoid genes and serum RBP4.**</u>

A. Gene expression of hepatic lecithin-retinol acyltransferase (*Lrat*) and *Retinoic Acid* and

Retinoid X receptors (*Rar* and *Rxr*) transcription factors (n=8 per group)

in C57BL/6 control CHOW diet mice or HFD obese mice treated with shHsd17b13 or scrambled

control (shScrmbl).

B. Western blot and quantification of serum RBP4 levels (n=5-6 per group).

Data is presented as mean ± SEM with analysed by One-way ANOVA (*$p \leq 0.05$) followed by

Bonferroni multiple comparison tests where * $p \leq 0.05$ and ** $p \leq 0.01$ (comparing HFD groups to

control CHOW diet).

C. HEK293 cells were transfected with HA-tagged Hsd17b13 or pcDNA3.1 control plasmid to

stably express Hsd17b13 and treated with retinoic acid (RA, 1 μm, 2 h) or DMSO (vehicle

control).

Gene expression of *Hsd17b13* and retinoic acid metabolism genes *Cyp26a1*, *Raldh1* and *Rarb.*

Data is presented as mean ± SEM from n=3 biological replicates and analysed by two-way

ANOVA where *** $p \leq 0.001$ (Hsd17b13 cells vs pcDNA control cells), # $p \leq 0.05$ and ## $p \leq 0.01$

and ### $p \leq 0.001$ (RA vs DMSO control). Hsd17b13 over-expression did increase *Raldh1*

expression by >2-fold ($p < 0.001$ by t-test)

Figure 6: <u>Effect of altered Hsd17b13 levels on lipid and phospholipid metabolism</u>

A. Gene expression of hepatic lipid metabolism genes (n=8 per group)

in C57BL/6 control CHOW diet mice or HFD obese mice treated with shHsd17b13 or scrambled shRNA (shScrmbl).

B. HepG2 cells stably expressing HA-tagged Hsd17b13 or pcDNA3.1 control plasmid were treated with 1mM fatty acid (2:1 ratio of oleic and palmitic acid) or methanol (vehicle control). Cells were stained with oil red O. Images are representative of 24 hour treatment. Scale bar = 100μM.

C. Fatty acid translocase *Cd36* gene expression HEK293 cells expressing Hsd17b13 or pcDNA3.1 control and treated with 1mM fatty acid (2:1 ratio of oleic and palmitic acid) or methanol (vehicle control).

D. Lipidomic analysis heatmap of most significant (p<0.05) changes in lipids, enriched in diacylglycerides and phospholipid species in liver tissue from C57BL/6 HFD obese mice treated with shHsd17b13 or scrambled shRNA (shScrmbl), n=3 for each.

E. and **F**. Gene expression of key regulators of phospholipid metabolism (n=8 per group)

G. *Cept1* (Choline-enthanolamine phosphotransferase 1) gene expression HEK293 cells expressing Hsd17b13 or pcDNA3.1 control and treated with 1mM fatty acid (2:1 ratio of oleic and palmitic acid) or methanol (vehicle control).

All data is presented as mean ± SEM.

A., **E**. and **F**. were analysed by one-way ANOVA followed by Bonferroni multiple comparison tests where * $p \leq 0.05$, ** $p \leq 0.01$ and *** $p \leq 0.001$ (HFD groups vs CHOW diet) or # $p \leq 0.05$, ## $p \leq 0.01$ and ### $p \leq 0.001$ (shHsd17b13 vs shscrmbl). Fads1 and Lpcat3 were increased in HFD, $p \leq 0.05$ by t-test.

C. and **G**. were analysed by two-way ANOVA

where * $p \leq 0.05$ and *** $p \leq 0.001$ (Hsd17b13 cells vs pcDNA control cells), # $p \leq 0.05$ and ## $p \leq 0.01$ and ### $p \leq 0.001$ (fatty acid vs methanol control).

FIGURE 1

A HFD / FEN HFD → RNA-seq, 20 weeks (obese)

B HFD-obese, fatty liver

Volcano plot: Significance, adj. p-value (-log10) vs Induced, fold change. Labeled genes: *Vnn1*, *Crat*, *Abca1b*, *Agpat9*, *Pex11a*, *Rdh16*, *Acot4*, *Mogat*, *Fabp2*, *Gstm2*, *Cyp46a1*, *Cyp2b13*, *Slc39a5*

C

Gene name	Induced (in HFD-obese)	Repressed (in FEN-HFD)
Mogat	14.0***	2.7*
Crat	5.1***	2.4***
Vldlr	3.6**	2.4*
Cyp46a1	5.1***	3.0*
Rdh16	3.2***	2.5**
Gstm2	3.6***	2.1*
Hsd17b13	1.5*	2.3*
Fads1	2.8*	2.2**
Pnpla3	2.4*	1.7(ns)
Inhbe	1.6*	1.3(ns)

Fold change

D Relative expression bar graphs (*Mogat*, *Crat*, *Vldlr*, *Cyp46a1*, *Rdh16*, *Gstm2*) — CHOW, HFD, FEN-HFD

E Relative expression bar graphs (*Mogat*, *Crat*, *Vldlr*, *Cyp46a1*, *Rdh16*, *Gstm2*) — Lean, db/db, FEN-db/db

F Relative expression bar graphs (*Avpr1a*, *Hsd17b2*) — CHOW, HFD, FEN-HFD

G Relative expression bar graphs (*Avpr1a*, *Hsd17b2*) — Lean, db/db, FEN-db/db

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 6