Khdc3 Regulates Metabolism Across Generations in a DNA-Independent Manner

Liana Senaldi[1,2], Nora Hassan[1], Sean Cullen[1,2], Uthra Balaji[2], Natalie Trigg[3], Jinghua Gu[2],

Hailey Finkelstein[1], Kathryn Phillips[1], Colin Conine[3], Matthew Smith-Raska[1,2]

1. Division of Neonatology, Department of Pediatrics, Weill Cornell Medicine, New York-
   Presbyterian Hospital, New York, NY, USA.

2. Drukier Institute for Children's Health, Weill Cornell Medicine, New York, NY, USA.
   mrs7001@med.cornell.edu.

3. Departments of Genetics and Paediatrics, University of Pennsylvania Perelman School of
   Medicine and Children's Hospital of Philadelphia, Philadelphia, PA, USA.

Abstract

Genetic variants can alter the profile of heritable molecules such as small RNAs in sperm and oocytes, and in this manner ancestral genetic variants can have a significant effect on offspring phenotypes even if they are not themselves inherited. Here we show that wild type female mice descended from ancestors with a mutation in the mammalian germ cell gene *Khdc3* have hepatic metabolic defects that persist over multiple generations. We find that genetically wild type females descended from *Khdc3* mutants have transcriptional dysregulation of critical hepatic metabolic genes, which persist over multiple generations and pass through both female and male lineages. This was associated with dysregulation of hepatically-metabolized molecules in the blood of these wild type mice with mutational ancestry. The oocytes of *Khdc3*-null females, as well as their wild type descendants, had dysregulation of multiple small RNAs, suggesting that these epigenetic changes in the gametes transmit the phenotype between generations. Our results demonstrate that ancestral mutation in *Khdc3* can produce transgenerational inherited phenotypes, potentially indefinitely.

Introduction

It is becoming increasingly apparent that non-DNA molecules inherited from germ cells contribute significantly to the transmission of traits and diseases across generations. This has been demonstrated in response to a variety of exposures such as diet or stress in model organisms (1, 2) and has been confirmed in human epidemiological studies (3, 4). The mechanisms driving this process remain poorly defined and are complicated by the erasure of acquired epigenetic molecules such as DNA methylation early in embryonic development. Expression of germ cell small RNAs such as microRNA (miRNA) and tRNA fragments (tsRNA) are responsive to exposures and

inherited at fertilization, making them potential candidates for passage of phenotypic information

across generations (5). Microinjection of these small RNAs from the germ cells of an exposed

mouse into an unexposed embryo is sufficient to recapitulate the descendant's phenotypes,

demonstrating causality (1). However, the absence of an RNA-dependent RNA polymerase in

mammals implies that small RNAs by themselves cannot propagate phenotypes across multiple

generations.

Non-inherited ancestral genetic variants can also affect descendants' risk of disease, as has been

demonstrated in cancer (6, 7), thyroid hormone metabolism (8), body weight (9), anxiety (10), type

I diabetes (11), and folate metabolism (12). This phenomenon is often referred to as "genetic

nurture" based on the effect of non-inherited DNA variants nurturing phenotypes in the next

generation (13). In all described cases, the observed phenotypes disappear after a few generations.

The molecular mechanisms driving these cross-generational traits, in the absence of inheritance of

the causal variant, remains largely undescribed, with the exception of paramutation effects that are

driven by inherited RNA molecules (14, 15). Some of the genes associated with this phenomenon

are thought to directly alter the molecular profile of the gamete, for example with testicular germ

cell tumors and RNA-modifying genes that are expressed in sperm (6). Other reported examples

involve genes that do not have any known function in the germ cells, and likely alter descendants'

phenotypes indirectly, by changing underlying physiology in a manner that subsequently alters

germ cell heritable molecules, not dissimilar from the mechanisms of exposure-based changes to

heritable germ cell molecules.

Khdc3 is a mammalian gene expressed in the male and female germ cells that encodes a protein

containing an RNA-binding KH domain that localizes to a multi-protein complex at the oocyte

periphery known as the subcortical maternal complex (16, 17). *Khdc3*-null female mice have

decreased fertility caused by defects in maintaining euploidy during embryogenesis (18). Human

females with homozygous mutations in the ortholog *KHDC3L* are infertile, which has been

associated with abnormal DNA methylation in oocytes (16). *KHDC3L* (and *Khdc3)* does not

localize to the nucleus and does not contain a DNA methyltransferase domain, suggesting that the

DNA methylation defects are secondary outcome and not the main function of this gene.

We demonstrate that wild type female descendants of *Khdc3*-null mice have dysfunctional

expression of hepatic metabolic genes that persists over multiple generations, and that this effect

is transmitted from both male and female mutant ancestors. This corresponds with abnormal levels

of hepatic metabolites in the serum of these wild type mice with mutant ancestors. The persistence

of these abnormalities in genetically wild type mice suggests that altered epigenetic information

in the germ cells is transmitting the inherited metabolic phenotypes. Accordingly, we observed

that the oocytes of *Khdc3*-null females and their wild type descendants have multiple dysregulated

miRNAs and tsRNAs, suggesting a mechanism of inheritance.

Results

Wild type females descended from Khdc3-null ancestors display hepatic transcriptional

dysregulation.

Global transcriptome analysis of *Khdc3*-null oocytes revealed significant dysregulation of genes

important in metabolic processes regulated by the liver, especially the metabolism of lipids

(atherosclerosis, PPAR signaling, and pantothenate and CoA metabolism) and of carbohydrates

(glycolysis/gluconeogenesis, and fructose and mannose metabolism) (Fig. 1A) (19-22). Despite

the abnormal expression of metabolic genes in the *Khdc3*-null oocyte, this gene is only expressed

in female reproductive tissue, with no detectable expression in the liver and minimal expression in

other metabolic tissues such as adipose or pancreas (Fig. 1B).

Expression of *Cyp17a1*, a gene central to lipid metabolism (23, 24), was increased in the livers of

Khdc3-null (knockout, or "KO") females, despite a lack of expression or any known function of

Khdc3 in wild type livers. Elevated expression of *Cyp17a1* was observed in KO females generated

from KO parents (referred to as "KO[KO]") as well as KO females generated by mating *Khdc3*

heterozygote parents (referred to as "KO*") (Fig. 1C). *Cyp17a1* expression was also significantly

increased, to the same extent as KO[KO] and KO* females, in genetically wild type females

descended from heterozygous parents (referred to as "WT*" to indicate genetically wild type mice

that have descended from *Khdc3* heterozygous mutant parents and a combination of wild type and

Khdc3-null grandparents). Thus, having an ancestor that carried a *Khdc3* mutation was associated

with elevated *Cyp17a1* expression, and the *Khdc3* genotype had no effect on *Cyp17a1* expression,

revealing a DNA-independent form of inheritance (Fig. 1C). Increased *Cyp17a1* mRNA

expression occurred in WT* females generated from all possible combinations of male and female

KO grandparents (Fig. 1D). In addition to conventional genotyping, the wild type genotype of

these mice was confirmed with RNA-Seq of ovaries (Supplemental Figure 1).

RNA-Seq of female WT* livers revealed a pattern of global transcriptional dysregulation that

overlapped significantly with KO* and KO[KO] females (Fig. 1E-F), demonstrating that shared

ancestry rather than genotype accounts for the transcriptional abnormalities in WT* mice. This is

supported by the observation that most significantly dysregulated genes in WT* livers were

similarly dysregulated in KO* livers (Fig. 1G). Gene ontology of the common dysregulated genes

in livers of WT* and KO* mice revealed enrichment in metabolic processes in which the liver

plays a central role, especially lipid and glucose metabolism (22-25) (Fig. 1H).

WT defects persist over multiple generations.*

When WT* male and female mice were mated with each other to generate the next generation

(referred to as WT**, signifying the 2^{nd} generation of genetically wild type mice), hepatic

transcriptional dysregulation of metabolic genes persisted. There was a similar pattern of

dysregulation in WT** females derived from either male or female *Khdc3*-null ancestors (denoted

WT**(P) and WT**(M), respectively, denoting Paternal or Maternal ancestral mutant history)

(Fig. 2A). Comparison of dysregulated genes between WT**(P) and WT**(M) females showed

significant overlap (Fig. 2C) revealing that the same pattern of abnormal gene expression is

observed in wild type females descended from both male and female mutant ancestors.

WT(P) male and female mice were mated with each other in order to examine passage these defects

over successive generations. Descendants of male mutants were examined to avoid potential

confounding from non-germ cell mechanisms of inheritance such as mitochondrial inheritance,

fetal-maternal communication across the placenta, or transmission of molecules via breastmilk.

Expression of *Cyp17a1* and *2610507I01Rik*, two genes dysregulated in WT*, KO*, KO^{KO}, and

WT**(P) female mice, demonstrated persistent and worsening dysregulated expression in the 2^{nd}

through 6^{th} generation of females (Fig. 2D). In a minority of mice, expression reverted back to

WT levels (Fig. 2D). Importantly, those mice that reverted to WT levels of *Cyp17a1* maintained

elevated expression of *2610507I01Rik*, and vice-versa, suggesting that the abnormal expression of

these genes is driven by independent units of inheritance.

Khdc3-null females were outcrossed with a true wild type male without any ancestral history of

Khdc3 mutation, in order to examine the persistence of the observed defects through only the

maternal line without any contribution from males that descended from mutants. 100% of the F2

outcrossed females, descended from a maternal grandmother mutant, had persistent transcriptional

dysregulation of metabolism-related genes in the liver (Fig. 2E). Many of these dysregulated genes

overlapped with WT** females descended from both male (WT**(P)) and female (WT**(M))

homozygous-null ancestors (Fig 2F). In this mating scheme, persistence of hepatic transcriptional

dysregulation in all 7 examined females is not consistent with mechanisms of inheritance involving

in-cis transmission, including an unaccounted DNA polymorphism/mutation, DNA methylation,

or histone protein modification, in which any chromosome from the *Khdc3*-null grandmother

would have been inherited by ~50% of the F2 outcross generation.

WT* females have abnormal levels of serum hepatic metabolites.

Global serum metabolomic analysis was performed to examine the metabolic consequences of the

observed transcriptional dysregulation. Both WT**(P) and WT****(P) female mice had abnormal

levels of multiple bile acids, which are synthesized in the liver (26-29)(Fig. 3A). The magnitude

of dysregulation was similar to that observed in KO^{KO} females (Fig. 3A), suggesting that

mutational ancestry was more important than the organism's genotype. These females also had

abnormal concentrations of various xenobiotics (Fig. 3B) and metabolic cofactors (Fig. 3C), both

of which are associated with hepatic metabolism (30-33).

WT****(P) mice were metabolically challenged with 8 weeks of a high fat diet (HFD). When

compared with true wild type mice exposed to a HFD, WT****(P) females exposed to a HFD

showed significant increase in multiple lipid molecules that were not dysregulated in WT****(P)

females that consumed a conventional diet, or in WT mice consuming a HFD (Fig. 3D). Thus,

there were latent metabolic abnormalities that could not be detected unless stressed with a high fat

diet. The HFD WT****(P) serum also had decreased levels of multiple other hepatic metabolites

(34-36)(Fig. 3E), some of which were also decreased in WT HFD females although at a much

lesser magnitude (Fig. 3F). These findings demonstrate that the hepatic transcriptional

dysregulation observed in these mice has significant effects on multiple metabolites found in the

blood.

Small RNA dysregulation.

Livers of WT**(P) females had dysfunctional expression of multiple small RNA-processing

genes, especially genes that regulate tRNA (*Trmt9b (37)*, *Ang (38)*, *Nsun6 (39)*, *Elac1 (40)*) (Fig.

3A) and miRNA processing (*Mettl1 (41, 42)*, *Ago2 (43)*, *Exosc10*, *Syncrip (44)*) (Fig. 4A). Based

on this observation, in combination with the fact that gamete small RNAs can transmit non-

genetically inherited phenotypes in mice, we hypothesized that the observed phenotypes were

driven by the defective inheritance of small RNAs from the germ cells of *Khdc3* mutant ancestors.

Small RNA-Seq of KO^{KO} oocytes revealed dysregulation of multiple miRNAs and tRNA

fragments, with minimal piRNA dysregulation (Fig. 4B). In WT**(P) oocytes, there was also

abnormal expression of tRNA fragments and miRNA, most of which were different small RNAs

than the ones dysregulated in KO^{KO} oocytes (Fig. 4C). Of note, the WT**(P) oocytes had

downregulation of the tRNA fragment Gly-GCC, for which expression in sperm has been

associated with inherited metabolic and hepatic gluconeogenesis phenotypes in offspring (1, 45,

46). There were 3 tRNA fragments and 18 miRNAs that were commonly dysregulated in both

KO^{KO} and WT**(P) oocytes (Fig. 3D), suggesting that their abnormal expression is established in

KO^{KO} oocytes and is not normalized with reintroduction of a wild type *Khdc3* allele. The function

of most of these commonly dysregulated tsRNAs and miRNAs remains undescribed, however

miR-107, which was upregulated in both KO^{KO} and WT**(P) oocytes, is involved in hepatic lipid

metabolism (47).

Discussion

The presence of abnormal phenotypes in wild type organisms descended from mutant-carrying

ancestors has been described in mice and humans, however all previously observed defects

disappear after one or two generations. This can occur from mutations that affect epigenetic

phenomenon in the germ cell (6, 7, 48), but in other cases the mutation affects a process that has

no clear connection with heritable molecules in the germ cell, such as with thyroid metabolism or

type I diabetes (8, 11).

We find that loss-of-function mutation in *Khdc3* alters the hepatic metabolism of female wild type

descendants over at least 6 generations in a manner that cannot be rescued with re-introduction of

the *Khdc3* allele. This is paralleled by abnormal levels of multiple metabolites in the serum of

these mice. Because there is no detectable *Khdc3* expression in the liver, we suspect that the

defects in WT* mice are caused by inherited molecules that affect hepatic metabolism independent

of the inheritance of a functional *Khdc3* gene. We expect that other metabolic tissues such as the

pancreas and adipose will also demonstrate evidence of metabolic dysregulation.

We have demonstrated that *Khdc3*-null oocytes have abnormal expression of multiple miRNAs

and tsRNAs, some of which persist over multiple generations in WT* mice, providing a potential

mechanism of inheritance. Microinjection experiments using sperm RNA have demonstrated that

small RNAs are sufficient to drive metabolic derangements in offspring (1, 49), however the

maintenance of defects to the successive generation remains unexplained. The amount of RNA

needed for these experiments currently prevents using oocyte RNA in such an experimental

paradigm, because of the low total RNA yield obtained from the oocytes of a single mouse.

We propose that the metabolic phenotype is at least partially driven by abnormal oocyte small

RNAs, however the persistence of a phenotype over multiple generations requires maintenance of

small RNAs via a RNA-dependent RNA polymerase, which mammals do not possess. It is

possible that small RNAs inherited from the oocyte drive abnormal metabolic profiles in offspring,

and oocyte small RNA defects are re-established in the germ cells of the following generation in

response to the abnormal metabolism. A recent study demonstrated that transfer of serum from a

stress-exposed mouse into a non-exposed mouse can recapitulate the effects observed in the

exposed's offspring, suggesting serum factors can modify the composition of heritable molecules

in the germ cells and affect offspring traits (50).

We utilized an outcrossing scheme to demonstrate that the observed defects persisted when passed

only through the maternal line. In this mating scheme, 100% of the observed F2 outcrossed female

mice had evidence of hepatic metabolic dysregulation. This pattern of inheritance is not consistent

with in-cis mechanisms such as an unaccounted DNA variant, DNA methylation, or histone protein

modification, because in these scenarios the alleles in the *Khdc3*-null maternal grandmother would

be present in ~50% of the F2 outcrossed females.

We chose to examine females because of the reported fertility phenotypes in *KHDC3* human (16,

17) and *Khdc3* mouse female mutants (18). Human *KHDC3*-null females are infertile, while

mouse *Khdc3*-null females are subfertile. The fertility defects are associated with oocyte DNA

methylation abnormalities that we suspect are sequelae of a more primary defect in mutant

organisms. There are no reports of abnormal phenotypes in *Khdc3*-null males, however our

detected metabolic abnormalities could affect males as well, which will be an important future

investigation. Furthermore, it will be important to examine how *Khdc3*-null mice transmit

exposure-based information across generations to affect phenotypes in non-exposed offspring, as

has been observed in numerous studies, most thoroughly with a high fat diet or stress.

One important conclusion from studies on genetic nurture is that wild type mice descended from

mutant ancestors should be used thoughtfully and with caution, because phenotypes can be

obscured that would be apparent with the use of wild type mice without any ancestral history of

mutation. In sum, the effect of non-inherited genetic variants on offspring phenotypes remains a

relatively unexplored and potentially significant contributor to traits and disease risk.

Materials and Methods

Mice

FVB/N WT (Jax 001800) mice were purchased from The Jackson Laboratory and bred in-house. Frozen *Khdc3-null* sperm was obtained from the Mutant Mouse Resource and Research Center (MMRRC, North Carolina) and the *Khdc*3 KO mouse was rederived. Female mice were used at 8 weeks of age. In individual experiments, all animals were age-matched. All mice were maintained under specific pathogen-free (SPF) conditions on a 12-hour light/dark cycle, and provided food and water ad libitum. All mouse experiments were approved by, and performed in accordance with, the Institutional Animal Care and Use Committee guidelines at Weill Cornell Medicine.

Mouse Genotyping

As described in this study, WT mice were generated from WT parents. WT* and KO* mice were generated from the mating of *Khdc3* heterozygote parents that were themselves generated from mating of WT and KO^{KO} mice. KO^{KO} mice were generated from *Khdc3* KO parents. DNA was extracted from tail biopsies by incubating the tails biopsies in an alkaline lysis buffer (24mM NaOH and 0.2mM disodium EDTA) for 30 minutes at 95°C and then on ice for 10 minutes. A neutralization buffer (40 mM Tris-HCl) was added to the samples and diluted 1:10 with water. PCR to detect *Khdc3* deletion was performed with 50–200 ng of DNA using the Phire Green Hot Start II DNA Polymerase Master Mix (Thermo Scientific) in the Bio-Rad T100 Thermal Cycler. PCR for *Khdc3* deletion was performed using a forward primer for the wild type allele F1, a separate forward primer that incorporates the deletion allele F2, and a common reverse primer R1. The primers were as follows: P1: 5'-TGCCTGGGCAGGTTATTTAG-3', P2: 5'-

CGAGCGTCTGAAACCTCTTC-3' and P3: 5'-AGCTAGCTTGGCTGGACGTA-3'. P1 and P2

amplified the wild type allele and P1 and P3 amplified the *Khdc3* mutant allele (KO). The PCR

products were separated on a 1% agarose gel with SYBR safe DNA gel stain. Because of the

different sizes of the PCR products, the genotypes can be easily determined from the band patterns

on DNA gels.

Liver RNA Extraction

Total RNA extraction was performed on the liver of 8 week old mice using the Qiagen RNeasy

Lipid Tissue Mini Kit (Qiagen) according to the manufacturer's instructions. Samples were eluted

in 30μl nuclease-free water. Nucleic acid concentration, A260/280 and A260/230 ratios was

determined via NanodropD-1000 (Thermo Scientific). Extracted samples were aliquoted and

stored at -80°C.

RT-PCR and quantitative RT-PCR

2000ng RNA per sample was used to generate cDNA using SuperScript III First-Strand Synthesis

System (Invitrogen) following the manufacturer's instructions. cDNA was diluted to 1:4 in water

before performing qPCR using SYBR Green PCR Master Mix (Applied Biosystems). qPCR

reactions were performed on a QuantStudio 6 Flex Real Time PCR Instrument. Cycling conditions

were as follows: Initial denaturation 95°C for 3 minutes, 40 cycles of denaturation at 95°C for 15

seconds followed by annealing/extension at 60°C for 60 seconds. Relative expression was

calculated using the ΔΔCt method, using YWHAZ as the reference gene. Primers used are listed

in Supplemental Table 1.

Oocyte RNA Isolation

The oocytes were dissected following a published protocol with some modifications (51). Briefly, the ovaries were first dissected from unstimulated 8-week-old WT and *Khdc3*-null mice and placed in PBS. The ovary was dissected from surrounding para-ovarian fat and subsequently placed in a 35-mm culture dish with 2mL PBS and 20μl collagenase and 20μl DNase in a 37°C incubator for 20 minutes. Using a dissecting microscope, the oocytes were separated from granulosa cells, theca cells, and stromal cells, and transferred to a petri dish with PBS using a mouth-controlled micropipette. RNA was extracted from the oocytes using the PicoPure RNA Isolation kit (Thermo Fisher) following the manufacturer's instructions and eluted in 20 µL of the provided elution buffer.

Liver RNA-seq Library Preparation

Sequencing libraries were prepared using the Illumina TruSeq Stranded Total RNA kit according to the manufacturer's protocol and sequenced to a depth of 40 million reads per sample. The paired-end (PE) libraries were sequenced on Novaseq platform.

Oocyte RNA-seq Library Preparation

Sequencing libraries were prepared using the SMART-Seq v4 Ultra Low Input RNA plus Nextera XT DNA Sample Preparation according to the manufacturer's protocol and sequenced to a depth of 40 million reads per sample. The paired-end (PE) libraries were sequenced on Novaseq platform.

RNA-seq Bioinformatics

320 All sequenced reads were assessed for quality using FastQC (52). Adapter trimming and filtering

321 of low-quality bases (<20) were performed using cutadapt (53). The trimmed reads were then

322 aligned with Hisat2 (54) against the mouse reference (GRCm38) genome. Per gene counts were

323 computed using htseq-count (55) (genocode M15). DESeq2 (56) bioconductor package was used

324 for normalization and differential gene expression analysis.

326 **Oocyte small RNA isolation**

327 Eight week old female mice were sacrificed and 30 oocytes were dissected under stereomicroscopy

328 from the ovaries of each mouse, as described above. Four separate WT, *Khdc3*-null, and WT(P)**

329 RNA samples were generated from separate mice. Total RNA was isolated using a magnet-based

330 method (ChargeSwitch Total RNA Cell Kit).

332 **Oocyte small RNA-Seq Library Preparation**

333 Small RNA quality and concentration was assessed using an Agilent Bioanalyzer and small RNA

334 Chip to ensure a clear small RNA population of more than 1ng. Subsequently, small RNA libraries

335 were generated and sequenced to a depth of 10 million reads per sample. Small RNA libraries were

336 generated using the NEXTFLEX Small RNA-seq Kit, which were sequenced on the Novaseq

337 platform.

339 **Small RNA-seq Bioinformatics**

340 Read quality was assessed using FastQC (v0.11.8), and adapter sequences were trimmed using

341 Trimmomatic (v0.39). After adapter trimming, reads were mapped sequentially to rRNA mapping

342 reads, miRbase, murine tRNAs, pachytene piRNA clusters (57), repeatmasker and Refseq using

Bowtie 2 alignment algorithm (v2.3.5) and totaled using Feature Counts on the Via Foundry

(v1.6.4) platform (58). To assess for differentially expressed small RNAs , data was loaded into R

Statistical Software and analyzed using the DESeq2 package (56). Differentially abundant small

RNAs were determined as those with a log fold change > 0.58 and P-value < 0.05.

Serum preparation

Blood was collected from the submandibular (facial) vein of mice. Approximately 400-500μl of

whole blood was collected from each mouse and placed in an SST amber microtainer blood

collection tube (BD). Blood collection tubes were placed in a 37°C incubator for 30 minutes,

followed by 10 minutes at 4°C. Samples were then centrifuged at 5500 RPM for 10 minutes at

4°C. The serum was collected from the top of the tube and stored at -80°C until needed.

Metabolomic profiling

Metabolomics profiling was conducted using ultra-high-performance liquid chromatography-

tandem mass-spectrometry by the metabolomics provider Metabolon Inc. (Morrisville, USA) on

mouse serum samples from WT, WT**(P), WT****(P) and KO^{KO} mice, as well as these mice

after consuming a high fat diet. The metabolomic dataset measured by Metabolon includes known

metabolites containing the following broad categories – amino-acids, peptides, carbohydrates,

energy intermediates, lipids, nucleotides, cofactors and vitamins, and xenobiotics. Statistical

differences were determined using unpaired t-test.

High fat Diet

Female WT and WT****(P) mice were divided into two diet groups, one group receiving a high-

fat diet (HFD, D12492i; Research Diets Inc., New Brunswick, NJ) and the other group received a

normal diet with 10% fat (D12450Ji; Research Diets Inc., New Brunswick, NJ). All mice were

given access to food and water *ad libitum* and were maintained on a 12:12-h light-dark artificial

lighting cycle. After 8 weeks of each diet, serum was collected and stored in the -80°C.

Statistical Analysis and data visualization

Gene ontology performed with WebGestalt (https://www.webgestalt.org/) using KEGG analysis.

Volcano plots generated with ggVolcanoR (https://ggvolcanor.erc.monash.edu). Venn diagrams

generated with InteractiVenn (http://www.interactivenn.net/). Scatter plots were generated with

Prism 9.2.0. The manuscript was written using Microsoft Word v16.66.1. Figures were

assembled on Microsoft PowerPoint v16.66.1

Data Availability: All RNA-Seq and small RNA-Seq data are available in the NCBI GEO

database.

Acknowledgements

This work was supported by the Young Investigator Award (LS) at Weill Cornell Medicine,

Pediatric Scientist Development Program under the National Institute of Child Health under award

number K12HD000850 (SC), Friedman Family Clinical Scholar Award (MSR), and National

Institute of Child Health and Human Development (NICHD) of the National Institutes of Health

under award number P50HD104454 (MSR).

Author Contributions

L.S. and M.S.R. conceived and planned the experiments. L.S., S.C., N.H., H.F., K.P., and M.S.R

carried out the experiments. U.B., N.T., J.G., and C.C. performed bioinformatic analysis of the

RNA-Seq and small RNA-Seq data. L.S. and M.S.R. prepared the manuscript. All authors

provided critical feedback and contributed to shaping the research, analysis, and manuscript.

References

1.   Q. Chen *et al.*, Sperm tsRNAs contribute to intergenerational inheritance of an acquired
metabolic disorder. *Science* **351**, 397-400 (2016).
2.   U. Sharma, Paternal Contributions to Offspring Health: Role of Sperm Small RNAs in
Intergenerational Transmission of Epigenetic Information. *Front Cell Dev Biol* **7**, 215
(2019).
3.   L. O. Bygren, G. Kaati, S. Edvinsson, Longevity determined by paternal ancestors'
nutrition during their slow growth period. *Acta Biotheor* **49**, 53-59 (2001).
4.   G. Kaati, L. O. Bygren, S. Edvinsson, Cardiovascular and diabetes mortality determined
by nutrition during parents' and grandparents' slow growth period. *Eur J Hum Genet* **10**,
682-688 (2002).
5.   C. C. Conine, F. Sun, L. Song, J. A. Rivera-Pérez, O. J. Rando, Small RNAs Gained during
Epididymal Transit of Sperm Are Essential for Embryonic Development in Mice. *Dev Cell*
46, 470-480.e473 (2018).
6.   D. Carouge *et al.*, Parent-of-origin effects of A1CF and AGO2 on testicular germ-cell
tumors, testicular abnormalities, and fertilization bias. *Proc Natl Acad Sci U S A* **113**,
E5425-5433 (2016).
7.   M. Y. Lam, J. D. Heaney, K. K. Youngren, J. H. Kawasoe, J. H. Nadeau, Trans-
generational epistasis between Dnd1Ter and other modifier genes controls susceptibility to
testicular germ cell tumors. *Hum Mol Genet* **16**, 2233-2240 (2007).
8.   J. Anselmo, N. H. Scherberg, A. M. Dumitrescu, S. Refetoff, Reduced Sensitivity to
Thyroid Hormone as a Transgenerational Epigenetic Marker Transmitted Along the
Human Male Line. *Thyroid* **29**, 778-782 (2019).
9.   S. N. Yazbek, S. H. Spiezio, J. H. Nadeau, D. A. Buchner, Ancestral paternal genotype
controls body weight and food intake for multiple generations. *Hum Mol Genet* **19**, 4134-
4144 (2010).
10.   V. R. Nelson, S. H. Spiezio, J. H. Nadeau, Transgenerational genetic effects of the paternal
Y chromosome on daughters' phenotypes. *Epigenomics* **2**, 513-521 (2010).
11.   S. T. Bennett *et al.*, Insulin VNTR allele-specific effect in type 1 diabetes depends on
identity of untransmitted paternal allele. The IMDIAB Group. *Nat Genet* **17**, 350-352
(1997).
12.   N. Padmanabhan *et al.*, Mutation in folate metabolism causes epigenetic instability and
transgenerational effects on development. *Cell* **155**, 81-93 (2013).

13. A. Kong *et al.*, The nature of nurture: Effects of parental genotypes. *Science* **359**, 424-428
(2018).
14. I. R. Adams, R. R. Meehan, From paramutation to paradigm. *PLoS Genet* **9**, e1003537
(2013).
15. S. Yuan, D. Oliver, A. Schuster, H. Zheng, W. Yan, Breeding scheme and maternal small
RNAs affect the efficiency of transgenerational inheritance of a paramutation in mice. *Sci*
Rep **5**, 9266 (2015).
16. H. Demond *et al.*, A KHDC3L mutation resulting in recurrent hydatidiform mole causes
genome-wide DNA methylation loss in oocytes and persistent imprinting defects post-
fertilisation. *Genome Med* **11**, 84 (2019).
17. D. Bebbere, L. Masala, D. F. Albertini, S. Ledda, The subcortical maternal complex:
multiple functions for one biological structure? *J Assist Reprod Genet* **33**, 1431-1438
(2016).
18. P. Zheng, J. Dean, Role of Filia, a maternal effect gene, in maintaining euploidy during
cleavage-stage mouse embryogenesis. *Proc Natl Acad Sci U S A* **106**, 7473-7478 (2009).
19. Y. Wang, T. Nakajima, F. J. Gonzalez, N. Tanaka, PPARs as Metabolic Regulators in the
Liver: Lessons from Liver-Specific PPAR-Null Mice. *Int J Mol Sci* **21** (2020).
20. P. R. Nagareddy, S. K. Noothi, M. C. Flynn, A. J. Murphy, It's reticulated: the liver at the
heart of atherosclerosis. *J Endocrinol* **238**, R1-R11 (2018).
21. A. Czumaj *et al.*, The Pathophysiological Role of CoA. *Int J Mol Sci* **21** (2020).
22. H. S. Han, G. Kang, J. S. Kim, B. H. Choi, S. H. Koo, Regulation of glucose metabolism
from a liver-centric perspective. *Exp Mol Med* **48**, e218 (2016).
23. A. Milona *et al.*, Steroidogenic control of liver metabolism through a nuclear receptor-
network. *Mol Metab* **30**, 221-229 (2019).
24. T. T. Wu *et al.*, Mutated CYP17A1 promotes atherosclerosis and early-onset coronary
artery disease. *Cell Commun Signal* **21**, 155 (2023).
25. P. Nguyen *et al.*, Liver lipid metabolism. *J Anim Physiol Anim Nutr (Berl)* **92**, 272-283
(2008).
26. J. Y. Chiang, Bile acid metabolism and signaling. *Compr Physiol* **3**, 1191-1212 (2013).
27. S. A. Ahmed *et al.*, Xanthosine, a purine glycoside mediates hepatic glucose homeostasis
through inhibition of gluconeogenesis and activation of glycogenesis via regulating the
AMPK/ FoxO1/AKT/GSK3β signaling cascade. *Chem Biol Interact* **371**, 110347 (2023).
28. M. Kusaczuk, Tauroursodeoxycholate-Bile Acid with Chaperoning Activity: Molecular
and Cellular Effects and Therapeutic Perspectives. *Cells* **8** (2019).
29. F. Brial *et al.*, The Natural Metabolite 4-Cresol Improves Glucose Homeostasis and
Enhances β-Cell Function. *Cell Rep* **30**, 2306-2320.e2305 (2020).
30. J. E. Klaunig, X. Li, Z. Wang, Role of xenobiotics in the induction and progression of fatty
liver disease. *Toxicol Res (Camb)* **7**, 664-680 (2018).
31. I. Lahdou *et al.*, Increased serum levels of quinolinic acid indicate enhanced severity of
hepatic dysfunction in patients with liver cirrhosis. *Hum Immunol* **74**, 60-66 (2013).
32. A. Sambeat *et al.*, Endogenous nicotinamide riboside metabolism protects against diet-
induced liver damage. *Nat Commun* **10**, 4291 (2019).
33. G. Liang *et al.*, Nicotinamide N-methyltransferase and liver diseases. *Genes Dis* **10**, 1883-
1893 (2023).

34.   M. H. Kim *et al.*, Daidzein supplementation prevents non-alcoholic fatty liver disease through alternation of hepatic gene expression profiles and adipocyte metabolism. *Int J Obes (Lond)* **35**, 1019-1030 (2011).

35.   L. Seidemann, A. Krüger, V. Kegel-Hübner, D. Seehofer, G. Damm, Influence of Genistein on Hepatic Lipid Metabolism in an In Vitro Model of Hepatic Steatosis. *Molecules* **26** (2021).

36.   T. Kose, J. Moreno-Fernandez, M. Vera-Aviles, P. A. Sharp, G. O. Latunde-Dada, Ferulic acid protects HepG2 cells and mouse liver from iron-induced damage. *Biochem Biophys Rep* **35**, 101521 (2023).

37.   P. C. Dedon, T. J. Begley, Dysfunctional tRNA reprogramming and codon-biased translation in cancer. *Trends Mol Med* **28**, 964-978 (2022).

38.   P. Ivanov, M. M. Emara, J. Villen, S. P. Gygi, P. Anderson, Angiogenin-induced tRNA fragments inhibit translation initiation. *Mol Cell* **43**, 613-623 (2011).

39.   S. Haag *et al.*, NSUN6 is a human RNA methyltransferase that catalyzes formation of m5C72 in specific tRNAs. *RNA* **21**, 1532-1543 (2015).

40.   M. C. J. Yip, S. Savickas, S. P. Gygi, S. Shao, ELAC1 Repairs tRNAs Cleaved during Ribosome-Associated Quality Control. *Cell Rep* **30**, 2106-2114.e2105 (2020).

41.   R. García-Vílchez *et al.*, METTL1 promotes tumorigenesis through tRNA-derived fragment biogenesis in prostate cancer. *Mol Cancer* **22**, 119 (2023).

42.   L. Pandolfini *et al.*, METTL1 Promotes let-7 MicroRNA Processing via m7G Methylation. *Mol Cell* **74**, 1278-1290.e1279 (2019).

43.   J. O'Brien, H. Hayder, Y. Zayed, C. Peng, Overview of MicroRNA Biogenesis, Mechanisms of Actions, and Circulation. *Front Endocrinol (Lausanne)* **9**, 402 (2018).

44.   L. Santangelo *et al.*, The RNA-Binding Protein SYNCRIP Is a Component of the Hepatocyte Exosomal Machinery Controlling MicroRNA Sorting. *Cell Rep* **17**, 799-808 (2016).

45.   B. Wang *et al.*, Paternal High-Fat Diet Altered Sperm 5'tsRNA-Gly-GCC Is Associated With Enhanced Gluconeogenesis in the Offspring. *Front Mol Biosci* **9**, 857875 (2022).

46.   U. Sharma *et al.*, Biogenesis and function of tRNA fragments during sperm maturation and fertilization in mammals. *Science* **351**, 391-396 (2016).

47.   H. Bhatia, G. Verma, M. Datta, miR-107 orchestrates ER stress induction and lipid accumulation by post-transcriptional regulation of fatty acid synthase in hepatocytes. *Biochim Biophys Acta* **1839**, 334-343 (2014).

48.   B. J. Lesch *et al.*, Intergenerational epigenetic inheritance of cancer susceptibility in mammals. *Elife* **8** (2019).

49.   Q. Chen, W. Yan, E. Duan, Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications. *Nat Rev Genet* **17**, 733-743 (2016).

50.   G. van Steenwyk *et al.*, Involvement of circulating factors in the transmission of paternal experiences through the germline. *EMBO J* **39**, e104579 (2020).

51.   S. El-Hayek, Q. Yang, L. Abbassi, G. FitzHarris, H. J. Clarke, Mammalian Oocytes Locally Remodel Follicular Architecture to Provide the Foundation for Germline-Soma Communication. *Curr Biol* **28**, 1124-1131.e1123 (2018).

52.   S. Andrews (2010) *FastQC: a quality control tool for high throughput sequence data*. p *Available online at:* http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

53.   M. Martin, Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* (2011).

54.	D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907-915 (2019).
55.	S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166-169 (2015).
56.	M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
57.	X. Z. Li *et al.*, An ancient transcription factor initiates the burst of piRNA production
during early meiosis in mouse testes. *Mol Cell* **50**, 67-81 (2013).
58.	O. Yukselen, O. Turkyilmaz, A. R. Ozturk, M. Garber, A. Kucukural, DolphinNext: a
distributed data processing platform for high throughput genomics. *BMC Genomics* **21**,
310 (2020).

Figures

531 **Figure 1**

532
Figure 1. Female mice with an ancestral history of Khdc3 mutation manifest hepatic

transcriptional dysregulation that is independent of genotype.

(a) Gene ontology scatterplot for dysregulated genes in *Khdc3*-null oocytes, based on RNA-Seq.

The y-axis represents −log FDR (false discovery rate) and the x-axis represents log2 enrichment

ratio. Dysregulation of genes important in lipid and carbohydrate metabolism are highlighted. **(b)**

Murine expression of *Khdc3* in various tissues, detected by qPCR (N=3). **(c)** Relative mRNA

expression of *Cyp17a1* in the livers of WT, WT*, KO^KO, and KO* mice measured by qPCR. Each

dot represents an individual mouse. **(d)** Relative mRNA expression of *Cyp17a1* in the livers of

WT* female mice generated from various male and female *Khdc3*-null grandparents measured by

qPCR. **(e)** Volcano plots depicting differentially expressed genes identified by RNA-seq in the

livers of WT vs. WT*, KO* and KO^{KO} mice, respectively (N=4-6). Red dots represent upregulated

genes in WT*, KO* and KO^{KO} mice while blue dots represent genes that were downregulated in

WT*, KO* and KO^{KO} mice. The x-axis shows log$_2$ fold change values, and the y-axis denotes

−log$_{10}$ p-values. **(f)** Dot plots of common dysregulated liver genes amongst the WT*, KO*, and

KO^{KO} mice compared to WT mice; ***p-adjusted < 1 x 10^{-5}. **(g)** Dot plot of the significantly

dysregulated genes in KO* livers (x-axis) versus WT* livers (y-axis), based on adjusted p-

value. Red dots reveal significantly dysregulated genes in KO* mice, blue dots reveal significantly

dysregulated genes in WT* mice, and grey dots represent commonly significantly dysregulated

genes in both KO* and WT* mice. **(h)** Gene ontology of the most common dysregulated genes

identified in the livers of both WT* and KO* revealed abnormalities in pathways critical for lipid

and glucose metabolism.

Figure 2

555
Figure 2. WT* defects persist over multiple generations and can be passed through the

maternal and paternal ancestral lines. (a) Schematic of experimental mating to form WT**(P)

mice. *Khdc3*-null mice are represented by black filled shapes, WT mice represented by white filled

shapes, and *Khdc3*-heterozygote mice represented by half-filled black shapes. Volcano plot

displaying dysregulated genes in the livers of WT**(P) mice compared to WT mice (N=4). Red

dots represent upregulated genes and blue dots represent downregulated genes in the WT**(P)

mice. **(b)** Schematic of experimental mating to form WT**(M) mice. Volcano plot displaying

dysregulated genes in the livers of WT**(M) mice compared to WT mice (N=4). Red dots

represent upregulated genes and blue dots represent downregulated genes in the WT**(M) mice.

(c) Venn diagram representing the number of significantly dysregulated liver genes of WT**(P)

and WT**(M) female mice compared to WT mice. **(d)** Pedigree schematic representing mating of

WT*(P) male and female mice over successive generations to form WT******(P) mice. Dot plots

represent relative liver mRNA expression of *Cyp17a1* and *2610507I01Rik* of the 2^{nd} through 7^{th}

generation WT female mice descended from male *Khdc3*-null mice as measured by qPCR. **(e)**

Pedigree schematic representing creation of a F2 outcross WT female from the mating of a female

Khdc3-null female and WT male and subsequent F1 generation *Khdc3*-heterozygote female with

WT male. Venn diagram depicting the overlap of commonly dysregulated genes in the livers of

F2 outcross WT and WT**(M) female mice compared to WT mice (N=4-7). **(f)** Dot plots

representing liver gene dysregulation amongst WT, WT*(P), WT*(M), and F2 outcross WT

female mice. ***$p < 1 \times 10^{-5}$.

Figure 3

Figure 3. Wild type mice with ancestral history of *Khdc3* mutation have dysregulation of multiple hepatically-metabolized molecule in the serum. Dot plots revealing abnormal concentration of **(a)** bile acids, **(b)** xenobiotics, and **(c)** metabolic cofactors present in KO^KO, WT**(P), and WT****(P) female mice compared to WT mice. **(d)** WT****(P) female mice exposed to a HFD had elevated serum concentration of lipid molecules compared to WT**** mice fed a conventional diet and WT mice fed a HFD. **(e, f)** Female WT****(P) female mice exposed to a HFD had decreased levels of multiple other hepatic metabolites, some of which were decreased in WT HFD female mice to a lesser magnitude *p < 0.05, **p < 0.005, ***p < 0.0005.

     Figure 4

589   **Figure 4. Oocytes of KOKO and WT**(P) mice have dysregulated expression of multiple**

590   **miRNAs and tsRNAs. (a)** Dot plots representing dysfunctional expression of small RNA-

591   processing genes of the livers of WT**(P) mice as measured in RNA-seq**. (b)** Volcano plots

592   depicting differentially expressed miRNAs and tRNA fragments in KO^{KO} oocytes compared to

593   WT oocytes. Red dots represent upregulated small RNAs, and blue dots represent downregulated

594   small RNAs (N=4). **(c)** Volcano plots depicting differentially expressed miRNAs and tRNA

fragments in WT**(P) oocytes compared to WT oocytes. Red dots represent upregulated small

RNAs, and blue dots represent downregulated small RNAs (N=4). **(d)** Scatterplot of the most

significantly dysregulated small RNAs of WT**(P) oocytes (x-axis) versus KO^{KO} oocytes (y-

axis), based on p-value. Green dots reveal dysregulated miRNAs and tRNA fragments in WT**(P)

oocytes, brown dots reveal dysregulated miRNAs and tRNA fragments in KO^{KO} oocytes, and

black dots represent commonly dysregulated small RNAs in both WT**(P) and KO^{KO} oocytes.

Additional Information

Supplementary Information is available for this paper.

Supplemental Material

Supplemental Figure 1.

Khdc3 expression in ovaries of WT, KO[*], and WT* mice, from RNA-Seq.

Supplemental Table 1: Primer sequences used for qPCR

Gene	Primer Sequence
YWHAZ	Forward primer: 5' CAGAAGACGGAAGGTGCTGAGA 3' Reverse primer: 5' CTTTCTGGTTGCGAAGCATTGGG 3'
Cyp17a1	Forward primer: 5'ACTGCAGTGATTGTCGGTCA 3' Reverse primer: 5' CTAGAGTCACCATCTGGGGC 3'
2610507I01Rik	Forward primer: 5' GGATCTGATAGTCGCCCGTG 3' Reverse primer: 5' TCGCAAGAGTTCCCTGCTTT 3'

610